Low-input soil management increases yield and decreases CO₂-emissions
but aggravates risk of nitrate leaching and diseases in winter wheat
cropping systems under climate change

Running title: Time travelling with *Triticum*

Michel J[1,2,a,*], Leemans V[2,3,a], Weinmann M[4], Balanzategui-Guijarro I[5], Bin J[1], Biver S[1],
Blum A[1], Börger R[6], Cao D[7], Him SL[1], Kirbas G[1], Le Gouis J[8], Moya-Laraño J[5], Persyn M[9],
Pierreux J[10], Quenon A[1], Sanchez-Moreno S[11], Symanczik S[12], Vanden Brande F[3], Van Der
Straeten D[7], Wagner M[4], Waibel M[12], Xayphrarath A[9], Vanderschuren H[1,2,13], Thonar C[1,14,b],
Delaplace P[1,2,b]

1 Plant Genetics and Rhizosphere Processes Laboratory, TERRA teaching and research centre, Gembloux Agro-
Bio Tech, University of Liège, Gembloux, Belgium
2 Transdisciplinary Agroecosystem Platform for Integrated Research (TAPIR), TERRA teaching and research
centre, Gembloux Agro-Bio Tech, University of Liège, Gembloux, Belgium
3 Biosystems Dynamics and Exchanges (BIODYNE), TERRA teaching and research centre, Gembloux Agro-
Bio Tech, University of Liège, Gembloux, Belgium
4 Institute of Crop Science, Nutritional Crop Physiology (340h), University of Hohenheim, 70593 Stuttgart,
Germany
5 Functional and Evolutionary Ecology, EEZA – CSIC, Almería, Spain
6 Harris Manchester College, University of Oxford, Oxford, United Kingdom
7 Laboratory of Functional Plant Biology, Department of Biology, Faculty of Sciences, Ghent University, Ghent,
Belgium
8 UMR GDEC, INRAE, Université Clermont Auvergne, Clermont-Ferrand, France
9 Polytech Lille, Université de Lille, Lille, France
10 Experimental Farm, Gembloux Agro- Bio Tech, Liege University, Gembloux, Belgium
11 Department of the Environment and Agronomy, National Centre Institute for Agricultural and Food Research
and Technology, INIA-CSIC, Madrid, Spain
12 Department of Soil Sciences, Research Institute of Organic Agriculture (FiBL), 5070 Frick, Switzerland
13 Tropical Crop Improvement Lab, Department of Biosystems, KU Leuven, Heverlee, Belgium
14 Agroecology Lab, Faculty of Sciences, Université Libre de Bruxelles (ULB), Brussels, Belgium
* Correspondence: jennifer.michel@uliege.be, +32495337705
a,b Contributed equally

Abstract

Understanding how climate change will affect crop performance is critical to ensure global food security and sustainability. Empirical data is key to anticipate the impact of climate change on cropping systems, but multifactorial climate change experiments remain scarce. In this study, the growth of winter wheat was examined in two agricultural soil management systems: one with long-term low organic inputs and the other one with high organic inputs. The wheat was grown in these differentially managed soils in an Ecotron, where the plant-soil mesocosms were subjected to three different climatic conditions. These conditions represent a gradient of ongoing climate change, simulating the weather patterns of the years 2013, 2068, and 2085 respectively. This approach allows to study the combined effects of projected increases in temperature, atmospheric CO_2-concentrations, solar irradiation and altered precipitation patterns on the cropping system (wheat growth, grain yield, rhizosphere processes, greenhouse gases, disease dynamics). The low-input system outperformed the high-input system with higher yields and lower CO_2-emissions in the future climates. On the other hand, the risk for plant diseases and nitrate leaching was also increased in the low-input system. To reduce the environmental impact of high-yielding cropping systems in the future it is therefore essential to identify management practices which allow fertiliser application and nutrient buffering without necessarily increasing organic inputs, like fertigation or biological nitrification inhibition. Under both here studied soil management systems the wheat plants developed natural coping mechanisms such as enhanced root growth and increased levels of proline and silicon to mitigate the adverse effects of environmental and biotic stresses. Unravelling the molecular mechanisms that trigger such inherent plant defences is a further interesting target for breeding future crops. Adapting crop rotations and cover crops to the shorter wheat cycle in the future is also an opportunity to break disease cycles.

Keywords

climate change, cropping system, ecotron, greenhouse gas emissions, nitrate leaching, plant health, soil health, soil management, sustainable agriculture, Triticum aestivum, winter wheat

Introduction

Climate change is expected to significantly impact winter wheat production in Central Europe over the next 100 years, with a theoretical potential for increased yields, but also an increased risk of crop failure following increased heat and water stress (European Environment Agency, 2024). Wheat is currently the most cultivated cereal in Europe, representing 34% of global production, and is the third most cultivated cereal in the world after rice (*Oriza sativa* L.) and maize (*Zea mays* L.), with over 95% of wheat produced worldwide being bread wheat (*Triticum aestivum* L.) (FAO, 2021). While wheat is a staple food in Europe which provides carbohydrates, protein, essential minerals like iron and zinc, and also vitamins like thiamine and pantothenic acid, its large-scale production is critical in terms of its environmental impact, notably nitrate pollution and greenhouse gas emissions (Shewry, 2013; Tilman et al., 2011; Asseng et al., 2019; Cao et al., 2024). Farmers therewith face a triple challenge: increase yields to feed a growing world population, reduce the negative impact of cropping systems on the environment and adapt land management to more and more challenging climatic conditions.

The optimal growth temperature for common winter wheat is 21°C with a non-stressing range between 9 and 22°C, it is sensitive to water logging and drought (Dickin & Wright, 2008; Lamba et al., 2023; Kumar et al., 2023). Under the Representative Concentration Pathways scenario RCP 8.5 W m^{-2} the Western European climate is predicated to be warmer (+3°C), with more (+140 mm) but unequally distributed rain and elevated CO_2 concentrations (+375 ppm) by the end of this century (IPCC, 2021). While warmer temperatures might initially benefit yields, extreme heat events during critical growth phases (like flowering) can lead to heat stress and reduced yields (Lamba et al., 2023; Riedesel et al., 2023). Rising temperatures, especially combined with elevated atmospheric CO_2 concentrations are also going to affect growth cycles, potentially leading to earlier wheat maturation and shorter growth cycles (Rezaei et al., 2018). This bears particular risks for winter wheat in central Europe as crops may be exposed to unfavourable weather conditions during their sensitive growth stages, for example heat waves during grain filling. Rising atmospheric carbon dioxide levels as such can have both beneficial and detrimental effects on crops. While increased CO_2 levels can initially benefit wheat growth through the CO_2-fertlisation effect, this is only observed under optimal temperature conditions. Beyond these conditions, the benefits diminish, and elevated CO_2 can decrease yields and protein levels (Myers et al., 2014; Porter et al., 2014). Additionally, elevated CO_2 levels can promote rhizodeposition,

which involves an increased release of root exudates intensifying microbial decomposition of

soil organic carbon and potentially increasing CO_2-emissions from the soil (Jansson and

Hofmockel, 2020). Moreover, changes in rainfall, including both increased frequency of

heavy rainfall and prolonged droughts, can negatively impact crop growth and ultimately

yields, especially if wheats suffer from insufficient water supply in critical growth stages

and/or excessive rainfall increases disease pressure and hinders field operations. In general,

warmer temperatures and higher humidity levels can facilitate the spread of pests and

diseases such as powdery mildew, septoria or take-all, which may lead to poor crop growth or

even crop losses if not managed effectively. Similarly, changes in temperature and moisture

influence soil microbial communities and nutrient cycling, potentially reducing soil health

and fertility over time with negative consequences for the harvested crops.

A further key point to consider is disease propagation in winter wheat cropping systems under

climate change, which is influenced by various factors including temperature, humidity, and

pathogen biology (Juroszek & von Tiedemann, 2013; Chakraborty & Newton, 2011). Warmer

temperatures can accelerate pathogen life cycles and increase disease severity, especially

fungal diseases may become more prevalent. Altered precipitation patterns and higher

humidity can also promote the growth of diseases such as powdery mildew and take-all

(Ghini et al., 2011; Miedaner, 2018). With shifts to the timing of winter and spring, crop

phenology and pathogen life cycles may face new combinations with currently unknown

consequences for crop vulnerability towards pathogens. Integrated pest management

strategies will be crucial for adapting to these changes, but these strategies depend on

accurate information about the anticipated disease pressure.

To prevent devastating impacts of climate change on winter wheat cropping systems and

anticipate negative consequences it is crucial to have accurate empirical data on future crop

growth cycles, plant nutrient requirements and disease propagation. To date, most

experiments focus on individual climate change factors such as elevated CO_2, however one of

the cruxes of climate change is the multifactorial aspect of abiotic stresses combining

elevated CO_2, altered precipitation patterns and rising temperatures at varying intensity. Here,

an Ecotron facility was used to expose wheat plants growing in two soils with contrasting

organic matter management to three climate change scenarios representing the meteorological

conditions of the years 2013, 2068 and 2085 respectively. Above and belowground cropping

system performance was monitored during the full growth cycle and thus unique insights to

the agronomic performance and the environmental impact of the wheat of the future were

obtained. The experimental insights provide new information about phenological shifts,

disease propagation, plant adaptation strategies, plant nutrient uptake and yield, as well as

nitrate leaching and greenhouse gas emissions.

The experiment addresses two main research questions: (i) how do future meteorological

conditions impact winter wheat cropping systems and (ii) can low or high organic input soil

management strategies prevent some of the anticipated negative impacts of climate change on

crop system performance and yield?

Methods

132 **Methods**

Time travelling with Triticum: experimental set-up in the Ecotron

Time travelling with Triticum: experimental set-up in the Ecotron

The experiment was implemented in the TERRA-Ecotron facility at Gembloux Agro-Bio

Tech, University of Liège, Belgium. The Ecotron was built in 2018 and currently has six

controlled environment rooms (CER). The experiment was carried out in mesocosms, which

have a soil compartment of 125 L each (cubes of 50x50x50cm), which allows to place nine

mesocosms in each CER, resulting in a total of n=54 experimental mesocosms. The first

factor "climate" was implemented with three levels, i.e. wheat was grown under the

meteorological conditions of the three years 2013, 2068 and 2085 respectively. The second

factor crossed with "climate" was "soil management", which had two levels that is the low

organic input (S1) and high organic input (S2). This resulted in a total of six modalities

(2013.S1, 2013.S2, 2068.S1, 2068.S2, 2085.S1, 2085.S2). Each modality was implemented

with eight replicates, half of which were kept untouched until final harvest for realistic

estimation of yield components and the other half was sampled repeatedly during the

experiment for destructive measurements such as root growth or leaf elemental composition,

each time taking 3-5 plants. In addition, for each soil type in each climate an unplanted

control cube was kept to measure baseline soil processes. Having six CERs meant that each

climate (2013, 2068, 2085) was replicated in two CERs and the replicate cubes of each of the

two soil management type were randomly distributed amongst these CERs (Fig. 1).

Three climate scenarios

Three climate scenarios

Historical data of continuous climate observations from the Ernage meteorological station

(50°34'33"N, 4°43'1"E, Belgium, since 1980) was used as well as predicted future

meteorological conditions using the Alaro-0 model (Giot et al., 2016). The model ran for the

Representative Concentration Pathway (RCP) scenario 8.5 W m^{-2} (IPCC, 2014) and the two

time periods 2040-2070 and 2070-2100 (Fig. 2). The three years selected from these

predictions (2013, 2068, 2085) align on a continuous gradient of increasing temperature,

precipitation, hydrothermal index (HI) and atmospheric CO_2 concentrations (Table 1). HI is a

measure of the relationship between precipitation and temperature, calculated as the ratio of

total precipitation to one-tenth of the sum of mean temperatures (Meshcherskaya &

Blezhevich 1997). A higher HI indicates wetter and/or cooler conditions, suggesting more

favourable moisture availability relative to temperature, which can benefit crops and

vegetation. Conversely, a lower HI signifies drier and/or warmer conditions, which may

indicate drought stress or reduced water availability. The Ecotron reproduces the simulated

weather conditions at a very high temporal resolution, where the key environmental

parameters such as sun light intensity and temperature are adjusted every five minutes. By

these means, diurnal and seasonal variabilities are accurately reproduced, and plant behaviour

can be studied under realistic climate scenarios. For example, the irradiation includes natural

sun rise and sun set patterns. The main climatic components that were manipulated for each

year in this experiment were atmospheric CO_2 concentration, temperature and precipitation.

CO_2 concentrations were along a gradient for the three years, with 420 ppm in 2013, 550 ppm

in 2068 and 775 ppm in 2085. The historic reference year 2013 was characterised by a mean

temperature of 7.59°C during the wheat growth cycle, with a mean precipitation of 2.12 mm

d^{-1} (HI=3.99). Interestingly, this year is the most extreme in terms of maximal and minimal

temperature (26°C, -7.6°C), and also the year with the longest periods outside the optimum

range for wheat in terms of cold and hot days (n=94 days below 4°C and n=9 days above

22°C). The year had a long cold winter from approximately 51-176 days after sowing, during

which the soil was frozen and therewith the water present in the soil was not readily available

to plants. The year 2013 also had the largest number of rain-free days (n=135), but the lowest

number of rainy days (n=57 days with precipitation >=4mm). The 2068 climate was

characterised by mean temperature higher than 2013 with 10.17°C but still had a significant

winter period with n=40 days where temperatures dropped below 4°C. Mean precipitation

was slightly higher than in 2013 with 2.4 mm d^{-1} and only n=4 days without rain, but more

rainy days (n=62). The climate of the year 2085 was the smoothest in regards of temperatures

around 12.1°C and very rarely below 4°C or above 22°C. The winter period was short with

only n=3 days below 4°C. In regards of precipitation, 2085 was the wettest scenario with

n=72 days of rain (>=4mm). Interestingly, all three years experience at least one day with

very high rain (>30mm), though these maximum rain events occurred at different moments

during the season: in March for 2068, in May for 2085 and in July for 2013.

Two soil types

The experiment tests two soils which are very closely related in terms of pedogenesis, both originating from the Walloon Brabant in Belgium (S1: 0°38'35.1474"N, 4°37'22.0123"E, S2: 50°39'12.8668"N, 4°38'10.7664"E). In this area limestone formations are prevalent and soils are often clay-rich, which helps retain moisture and nutrients making them very suitable for agriculture (Gentile et al., 2009). Both soils are classified as Aba(b)0 in the Belgian soil classification system and characterised as silty loam (Cigale, 2021). Both soils are from fields that are under agricultural management since generations and implement the regular local crop rotations with winter wheat and root vegetables mainly. In recent years, cover crops with plants such as phacelia, oats, radish and clover have always been grown between two main crops in S2. In both fields standard and reduced tillage was applied regularly, as well as commercial fertilisers, green and brown manure and occasionally herbicides (supplementary material SM1). The main differences between the two soils is that soil two (S2) has received significantly higher quantities of organic inputs than soil one (S1). While the soils have similar C:N of 10.5 and pH just above 8, these higher organic inputs left S2 with more than doubled humus, carbon (C), nitrogen (N), phosphorous (P), potassium (K), magnesium (Mg) and calcium (Ca) contents as compared to S1 (Table 2). Another difference is in the soil texture, which is sandier for the low organic input S1 soil. In each field, a total of n=27 cubes were sampled in November 2022 and moved to the Ecotron. Soils were sampled as undisturbed soil monoliths (50x50x50cm) with a surface of 0.25 m^2 and each weighing approximately 200 kg. Monoliths were taken to realistically represent field conditions in the Ecotron, avoid disturbance of sensitive soil organisms and keep the soil structure intact. One cube of each soil type in each CER was placed on a scale to monitor the weight of the cubes and improve estimates of evapotranspiration.

Crop management and monitoring

During an initial acclimatisation period of three weeks, the soil cubes were kept under respective climates in the CERs of the Ecotron. At the time of the monolithic sampling for the Ecotron trial the S2 field was pre-cropped with a radish mix, in grass cover with a dense layer of mulch in the sub-surface (wheat straw). There was no cover crop sown in the S1 field prior to sampling the cubes, but a straw layer had also been incorporated to the soil before the sampling. The soils in all cubes were weeded and surface tillage (raking to 15cm depth) was applied during the acclimatisation period. On 23/12/22, which corresponds to 01/11 Ecotron date, 54 cubes were planted with winter wheat (*Triticum aestivum* (L.) var. Asory) at a density

of 308 seeds m^{-2} (77 seeds per cube) as recommended for this region (Livre Blanc Céréales,

2022). Cubes were weeded manually when needed and no herbicides were applied. At the

beginning of the experiment, metaldehyde pellets were applied to control molluscs. The

pellets were placed in the reversed lids of 50 ml Falcon tubes which were then placed on the

soil to minimise impact on soil chemistry and other soil organisms (Birkhofer et al., 2008;

Iglesias et al., 2003). After germination, plant development was closely monitored and

several agronomic and environmental parameters were regularly recorded (Fig. 3,

supplementary material SM2). For the aboveground compartment, plant BBCH growth

stages, plant height and leaf area index (LAI) were measured to quantify plant growth and

maximum quantum efficiency of photosystem II (Fv/Fm) was measured as an indicator of

plant performance (Meier, 2001; Vlaović et al., 2020). At three time points (BBCH30/50/80)

foliar silicon and foliar proline were determined as important molecules in plant stress

response (Hayat et al., 2012; Wang et al., 2017). For the belowground compartment, soil

microbial biomass and total root length were measured at the same time points and root

infestation with the fungus *Gaeumannomyces tritici* (take-all disease) was quantified once at

230 DAS (Campbell et al., 2003; Seethepalli et al.,2021; BBA, 1999). Unless prevented by

drought conditions, interstitial soil pore water was extracted weekly to quantify freely

available nitrate in aqueous soil solution and glucose equivalents as indicators of root

exudation (Folegatti et al., 2005; Yemm & Williams, 1954). During the experiment, both soils

in all climates were fertilised with ammonium nitrate (N: 27%), which was applied in three

doses according to plant growth stage, namely at the end of tillering/stem-elongation, 2. node

and flag leaf in each climate respectively. The quantities of N-input varied between 150 and

205 kg N ha^{-1} per application depending on the pre-existing soil N content for the two soils,

with S1 receiving approximately 50 kg N ha^{-1} more than S2 (supplementary material SM1).

Plants were harvested when fully ripe (BBCH89) and for each cube aboveground biomass

was determined (straw/leaves/heads), as well as number of heads, grain fresh weight, grain

humidity, grain yield at 14% moisture, thousand grain weight and grain nitrogen content.

DNDC model

DNDC (i.e. DeNitrification-DeComposition) is a process-based computer simulation model

of carbon and nitrogen biogeochemistry in agro-ecosystems (Giltrap et al., 2010; Gilhespy et

al., 2014). DNDC predicts soil environmental factors, C sequestration, and emissions of C

and N gases primarily based on microbe-mediated biogeochemical processes, including

decomposition, nitrification, denitrification, fermentation, and methanogenesis (Li et al.,

1992a, 1992b, Li, 2000). DNDC simulates these processes based on the activity of different

functional groups of microbes under different environmental conditions including

temperature, moisture, pH, redox potential (Eh) and substrate concentration gradients in soils.

For example, nitrification is modelled as first-order process based on soil ammonium

concentration (NH_4^+) under aerobic conditions and nitrous oxide production (N_2O) is

modelled as a fraction of the overall nitrification rate. Soil Eh is calculated with the Nernst

equation at a daily time step following soil saturation and then used to determine anaerobic

microbial group activity under a given set of soil conditions. Anaerobic microbial group

activity is then modelled using standard Michaelis-Menten-type kinetics. The DNDC model

has been extensively evaluated against datasets of trace gases fluxes that were measured

worldwide (Gilhespy et al., 2014; Giltrap et al., 2010). To access the accuracy of the model

for this experiment, the model's predicted outputs were compared with measured Ecotron

variables and the coefficient of determination (R^2) was used as a measure of goodness of fit

(supplementary material SM3).

Statistical analysis

The effects of climate and soil type on the empirically measured parameters (supplementary

material SM2, lines 1-12) were assessed using linear mixed-effects models (LMM) with

climate and soil as fixed effects, and random intercepts for time and CER to account for the

within-subject correlation of repeated measures and replicate rooms. Post-hoc comparisons

using estimated marginal means (EMMs) were used where appropriate to elucidate

differences between the levels of each fixed effect, i.e. the interaction between climate and

soil (2013/2068/2085xS1/S2). For yield components (Table 3, Fig. 3Q) and take-all index

(Fig. 3L) analysis of variance (ANOVA) and post-hoc Tukey's HSD were used to identify

differences between climate and soil type. The parameters derived from the DNDC model,

that is upscaled budgets of CO_2, N_2O, N leaching and transpiration, have only one

observation per modality (Fig. 3R-T). They are descriptive but were indicatively compared

via Krustkal-Wallis test for climate and soil individually. To reduce the overall dimensionality

of the dataset to better understand what the core characteristics of each cropping system in

each climate are, probabilistic principal component analysis (pPCA) was performed (Fig.4).

pPCA extends traditional PCA by incorporating a probabilistic framework, which allows the

estimation of principal components to identify underlying latent structures in the presence of

noise (Tipping & Bishop, 1999). Statistical analysis was carried out using R 4.4.1 (R Core

Team, 2024) with the additional packages car (Fox & Weisberg, 2019), emmeans (Lenth,

2024), ggplot2 (Wickham, 2016), lmerTest (Kuznetsova et al., 2017), missForest (Stekhoven
& Buehlmann, 2012) and multcompView (Graves et al., 2024).

Results

Phenological advance in future meteorological conditions

Warmer temperatures and more intense rain, but overall less stressful weather with higher
atmospheric CO_2 concentrations meant that crops grew faster and bigger the further in the
future the climate scenario was (Fig.3E-G). The phenological advance manifested from
February onwards, with plants maturing faster in 2068 and 2085 (BBCH p<0.0001). There
was a significant effect of soil type in interaction with climate for plant height (p<0.0001),
with taller plants the further in the future the climate scenario. Notably, plants in 2085 were
almost twice as high as in 2013 (Fig.3F) and plant height was one of the main characteristics
of the 2085 cropping systems on both soil types in the ordination (Fig. 4, supplementary
material SM4). In 2013, plants in S2 grew taller than plants in S1 (2013.S1: 19.3±0.445,
2013.S2: 20.8±0.476). The soil type effect was reversed for the two future climates where
plants grew taller in S1 compared to S2 and (2068.S1: 25.6±0.445, 2068.S2: 24.0±0.445,
2085.S1: 39.4±0.445, 2085.S2: 37.2±0.445). Small leaf area index (LAI) was one of the main
characteristics of the 2013 cropping systems, particularly S2 (Fig. 3G, Fig. 4, supplementary
material SM4) and considerably increased in the future climates (p<0.0001).

Early harvest and yield components

The phenological advance entailed earlier harvests in the future years, with approximately
two weeks between each of the climate scenarios. Plants in 2068 were harvested first in early
July, followed by 2085 in late July, while 2013 was harvested beginning of August (all dates
refer to Ecotron months). Visually distinguishable differences in plant developmental stages
between the two soil types were minor and plants from both soils were harvested
simultaneously once all grains were fully ripened for the respective year.

Grain yield was always higher in S1 as compared to S2 (Fig.3Q, Table 3), with an overall
significant soil type effect and minor differences between climate scenarios (p=0.04). Most
notably, yield increased proportional to the increasing hydrothermal index across the three
years for S1, meaning that for this soil type the further in the future the climate scenario, the
higher the yield. The globally highest aboveground biomass and also highest grain yield were
achieved for S1 in 2085, while the lowest grain yield was recorded for S2 in 2085. Thousand

grain weight was also significantly affected by soil and climate (p<0.0001) with lowest

values in 2013 and highest values for S2 in 2068 (Table 3). Grain nitrogen was not

significantly different between soils or climates, however trended to be higher in S1 as

compared to S2 and was highest in 2013 (p=0.13).

Plant health

The maximum quantum efficiency of photosystem II, represented as Fv/Fm (Fig.3I), is a

crucial parameter in plant physiology with values around 0.8 considered optimal and lower

values indicating stress leading to impaired PSII efficiency (Maxwell & Johnson, 2000;

Baker, 2008). Fv/Fm is measured at leaf level and requires significantly wide leaf area, which

is why the measurement was initially not possible in 2013. For the data available, Fv/Fm was

not significantly different between soil types, but significantly different between climates

(p<0.002), with on average higher Fv/Fm in 2085 than in 2068 (than in 2013 when

measured). A notable drop in Fv/Fm was recorded for 2068 compared to 2085 at the

beginning of the experiment (March Ecotron month) and overall lowest Fv/Fm was measured

for S2.2068 which distinguished this cropping system (Fig. 4, supplementary material SM4).

Leaf silicon and proline levels as key indicators of plant stress and adaptation mechanisms

were also characterised (Fig. 3J, 3K). There was notably elevated foliar silicon in S2.2085 at

BBCH50 and elevated proline in S1.2068 at BBCH80 (Fig. 4, supplementary material SM4).

Overall, leaf silicon levels were most distinguished 2085 (p=0.02) with the lowest levels for

S1.2085 and the highest foliar silicon for S2.2085 (Fig.3J). Foliar silicon tended to be

globally higher in S2 than in S1 (p=0.095). Foliar proline also stood out in 2085 (p=0.07),

with an opposite trend to silicon, having lowest values in S2.2085 and highest in S1.2085

together with S1.2068 (Fig.3K).

Take-all disease is a common struggle in winter wheat plantations (Palma-Guerrero et al. ,

2021) and was quantified at BBCH80 in this experiment (Fig.3L). Symptoms of root

infestation with the fungus *Gaeumannomyces tritici* (take-all index) were increased in 2085,

especially in S1, but overall no significant climate or soil effect was detected (p=0.06).

Rhizosphere processes and environmental impact

Decreased total root length (TRL) was amongst the main characteristics of the two future

cropping systems on S1 in the ordination (Fig. 3O, Fig. 4, supplementary material SM4).

TRL differed particularly at the early stages of plant development where it was higher in S2

compared to S1, but TRL was in frequentist approach not significantly different (p=0.08).

Microbial biomass in the root zone was quantified at three time points corresponding to

BBCH 30/50/80 (Fig.3E), where it was always higher in S2 compared to S1 (p=0.03).

Microbial biomass was one of the main characteristics of the S1.2068 cropping system,

together with glucose equivalent (Fig. 4, supplementary material SM4). Detectable glucose

levels averaged around 0.038 mg ml-1 in all modalities and were one of the main

characteristics of the S1.2013 cropping system (Fig.4, supplementary material SM4). There

was a periodical variation in glucose concentrations with a common low level during winter

and then varying peaks according to climate and soil type (Fig. 3N). The overall lowest levels

of glucose were detected in S1.2013. Later in the experiment, the largest spikes were detected

in the 2085 climate, with the all-time highest peak at the beginning of August in S2.2085

(Fig.3N). Freely available nitrate presented high temporal variation just as did glucose levels

(Fig.3M), yet nitrate in soil solution was overall significantly higher in the two future

climates (p=0.02) and globally not significantly different between soil types. Most notable

spikes in free nitrate were detected in S1.2068 and S1.2085 (Fig. 3M) and according to the

DNDC model the risk of nitrate leaching was always higher in S1 compared to S2 (Fig.3T),

but not statistically significant for soil nor climate. The DNDC model also predicted both

higher soil CO_2 emissions and higher soil N_2O emissions for S2 compared to S1 (both

p=0.05). Comparing predicted GHG emissions across the three climate scenarios revealed an

overall trend towards decreasing soil N_2O and increasing soil CO_2 emissions with increasing

hydrothermal index (Fig.3R,3S).

Discussion

Yield and soil organic C: Less is more

In all three climate scenarios, plants in the low input soil S1 yielded more grains than plants

in the high input soil S2 (p=0.007). The proportional increase in yield with the hydrothermal

index for S1 supports previous findings that optimal moisture and temperature conditions can

lead to substantial yield improvements in the future (Wilcox & Makowski, 2014). That this

effect was observed with S1 but not with S2 aligns with other research showing that soil

properties play a key role in wheat productivity, especially as climate conditions fluctuate

(Zhao et al., 2022). However, the better realisation of the yield potential under future climates

in S1 adds some important nuance as to whether increasing organic inputs and soil organic

carbon per se is an advisable farming practice. It may be that in soils with higher organic

carbon content and larger microbial communities, the competition between soil microbes and

plants is enhanced to a degree which prevents optimal plant nutrient uptake and growth

(Kuzyakov & Xu, 2013). For harnessing the yield potential in this study, a tipping point was

identified for S2 where yields increased for the near future, but decreased strongly for the far

future scenario (Fig.3Q, Tabel 3). The result of this study with a yield gap between the two

soil types aligns with a mesocosm experiment in California which also found that higher soil

organic carbon does not relate to higher wheat yields (Kelley et al., 2024) and with a survey

of European farms which found a poor association between soil organic matter and crop

yields (Vonk et al., 2020). Future studies could investigate under which conditions of nutrient

and water availability lower organic inputs can sustain similar to higher yields as high input

systems. In addition to higher yields, the cropping systems with S1 also emitted less CO_2 than

S2 systems, which is most likely explained by the enhanced microbial biomass in the S2 soils

(ESDAC, 2020). This underlines that definitions of "healthy" cropping systems cannot solely

rely on high humus/ soil C content and increased microbial biomass C contents, as these often

link to higher microbial activity and therewith accelerated soil CO_2 emissions, a potentially

negative feedback loop to climate change (Hamamoto et al., 2022; Moitinho et al., 2021).

Therefore, it is critical to consider the relationship between soil carbon content and nutrient

cycling, as these links are fundamental for understanding soil health and its role in

biogeochemical cycles (Schröder et al., 2016; Rocci et al., 2024). Overall, the here observed

divergence in yields as a function of soil type demonstrates the sensitivity of cropping

systems to climate change (Benton, 2020; Hamamoto et al., 2022). To avoid detrimental

losses of yield when agricultural systems cross tipping points and to ensure food security

worldwide even under climate change, it needs to be carefully evaluate which management

practices provide resilience to cropping systems in the long-term (Kornhuber et al. 2023).

Quality of harvested grains

To prevent malnutrition, it will also be important to determine the technological properties

and nutritious value of the harvested grains (Lowe, 2021). In this study, grain nitrogen

content was not statistically different between soil types or climates but tended to decrease

the further in the future the climate scenario (Table 3). Such a trend of decreasing nutrient

levels with increasing temperatures and CO_2 is often reported in the literature, but our results

provide an interesting example of how crops can compensate dilution effects from CO_2-

fertilisation, which merits further study (Liang et al., 2019; Govindasamy et al., 2023).

Interestingly, in this study thousand grain weight (TGW), which is an important component

of crop yield linked to potential flour yield (Bordes et al., 2008), increased in parallel to the

overall yield for the future climate scenarios, but had no consistent relation with soil type

(Table 3). This presents a particular challenge for breeding programs, as TGW is closely

linked to the stress resilience of wheat plants. Factors like drought and nutrient deficiency can

lead to premature ripening and reduced TGW, while also influencing seedling vigour in the

next generation (Shahwani et al., 2014). However, developing a soil management plan to

target these traits requires more empirical data to establish consistent trends between soil

management practices and nutrient redistribution during grain filling, taking the shorter

growth cycle in the future climates into account.

Naturally balancing the nitrogen cycling?

N_2O is a relevant greenhouse gas with much higher global warming potential than CO_2

(IPCC, 2014). Agricultural land is one of the main N_2O sources and as higher temperatures

stimulate mineralization and nitrification processes and therewith substrate availability for

denitrification, increases in N_2O emissions are expected in the future (Lamprea et al., 2021).

However, depending on local conditions such as soil moisture and N availability, reductions

in soil N_2O emissions are also possible (Muños et al., 2010; Reay et al., 2012). In this study,

the DNDC model predicted an overall decrease in soil N_2O emissions for future climates,

with higher N_2O-emissions in S2 compared to S1, while the difference between S1 and S2

regarding N_2O-emissions did not change between the years (Fig.3S). As the future years were

smoother in the rainfall distribution, the soils were overall drier (Fig.3D,H) and therewith the

risk of waterlogging and anaerobic conditions was reduced, limiting denitrification and N_2O

production (Venterea, 2007). Moreover, there was no indication of limited plant nitrogen

uptake in the future climates (Table 3), which could indicate that the increased soil

mineralisation did not lead to nutrient losses in GHGs, but instead nitrogen was taken up by

the growing plants. Another mechanism which could have successfully prevented gaseous N-

losses could be associated with greater N immobilization belowground, with surplus

inorganic NO_3-N being incorporated into microbial biomass, supporting greater N recycling

and retention (Buckeridge et al., 2020; Cao et al., 2021; Pausch et al. 2024). The process of

enhanced nutrient immobilisation in microbial biomass could also explain the decreased risk

of nitrate leaching predicted for S2 as compared to S1, which has higher microbial biomass,

while the sandy texture of S1 likely contributed to its higher susceptibility to nutrient

leaching (Gaines & Gaines, 1994). This means that while low organic input may be a soil

management strategy to reduce GHG-emissions without compromising yield, soils with

higher drainage may experience greater nitrate loss through leaching due to the soils' inability

to retain nutrients effectively. To reduce the environmental impact of high-yielding cropping

systems in the future it is therefore essential to identify soil management practices which

allow nutrient buffering without necessarily increasing organic inputs, like fertigation or

biological nitrification inhibition.

Stimulating natural plant adaptation mechanisms to mitigate stress

Plants in the future climatic conditions were more prone to stress because of their changed

phenology and a higher potential for disease propagation. For example, leaf area index (LAI)

increased significantly under future climate scenarios, which not only indicates increased

rates of photosynthesis, but a larger leaf area also allows for greater stomatal conductance,

which facilitates more water vapor loss through transpiration and may thus stress plants and

increase the water requirements of the cropping system (Zhang et al., 2021). Therewith,

plants with larger LAI are more vulnerable to drought spells which can be detrimental for

yields (Farooq et al., 2009; Li et al., 2023). Similarly, taller growing plants as observed in the

future climates in this experiment are more vulnerable to physical damage from wind and

require better rooting systems and stronger cell walls to withstand high wind speeds (Zhao et

al., 2022; Jia et al., 2021; Gardiner et al., 2016). In addition, plant health is expected to

worsen under climate change when autumns and winters become milder and wetter which

favours water logging and the spread of fungal diseases which may ultimately pose severe

risks to food production systems (Chakraborty & Newton, 2011). In this study, the pathogenic

root fungus *Gaeumannomyces tritici* which causes take-all disease was enhanced in the two

future climates, most notably in the 2085 climate, and there particularly in S1, but the

pathogenic root infestation did eventually not threaten the yields (Fig. 3L, 3Q). This indicates

that the plants could develop successful stress mitigation strategies to combat the pathogen.

Two promising factors which could have contributed to the plants' successful defence are

proline and silicon. Proline functions as a potent antioxidant, scavenging reactive oxygen

species (ROS) and reducing oxidative damage, which can help protect cellular structures and

macromolecules during stress conditions (Hayat et al., 2012). Moreover, exogenous proline

application has been shown to improve photosynthetic parameters such as chlorophyll

content, stomatal conductance, and PSII efficiency in stressed plants (Sporman et al., 2023)

and proline accumulation is associated with increased resistance to various pathogens acting

as a signalling molecule triggering defence responses and the production of antimicrobial

compounds (Kaur & Asthir, 2015). Silicon on the other hand can contribute to plant health by

impregnating cell walls and thus forming a barrier that impedes pathogen penetration, while it

also physically strengthens plant stems which could benefit crops that grow taller and are

expose to higher wind speeds (Wang et al., 2017). Silicon has also been shown to improve

drought tolerance which could become even more important in the future as crops grow with

larger leaf area like in this study (Wang et al., 2021) and like proline has been related to

oxidative stress mitigation (Kim et al., 2017).

Implications of phenological advance on farming practices

Plants use various signals to respond to environmental changes, including solar signals which

determine the photoperiod, past seasonal experiences like winter chilling, and current

conditions such as temperature and moisture (Tang et al. 2016). The advancement in harvest

date of approximately two weeks between each climate scenario observed in this study

reflects this response and aligns with trends observed in other experiments which report that

warmer temperatures accelerate crop development and lead to earlier maturity dates and that

in spring photoperiod and winter chilling work together to determine plant growth (Tang et

al. 2016; Harkness et al., 2020). The here observed phenological advance in future

meteorological conditions has significant implications for agricultural management, for

example harvesting winter wheat up to four weeks earlier requires adapting the crop rotation

cycle including sowing dates and identifying wheat varieties with shorter maturation cycles.

Another observation in this experiment was that plants in 2085 reached heights nearly double

those recorded in 2013, which is in line with other studies (Quan et al., 2024). These taller

plants may require more physical protection against wind by means of hedges or trees and

also imply that harvesting equipment may need to be adapted to the higher positioning of the

grains (Miller et al., 2022). Overall, the projected advances in crop maturity and plant height

could have profound implications for agricultural practices and as climate conditions

continue to evolve, further research will be needed to refine models predicting phenological

responses and to develop adaptive strategies for sustainable agricultural practices in changing

environments.

Limitations of the Ecotron

510 While Ecotrons are unique tools to study agroecosystems under climate change, they are a

511 compromise operating at an intermediate scale between simplistic microcosm experiments

512 and real-world ecosystems, which still cannot fully replicate the complexity of natural

513 environments (Roy et al., 2021; Schmidt et al., 2021). It is therefore vital to cross-validate

514 observations from Ecotron experiments with data from field experiments and to replicate

experiments sufficiently. A further challenge to Ecotron experiments is the timescale, as exposing soil and miniature agroecosystems to future meteorological conditions without several generations of adaptive changes occurring implies a very abrupt change in climate conditions. In this experiment for example, soil monoliths sampled in November 2022 had about one month to adapt to the conditions of the simulated November 2085 rather than slowly evolving through 63 years of climate change. However, the experiment with its accelerated evolutionary approach remains realistic, especially for wheat as a plant with a long growth cycle in a crop rotation, because even in current meteorological conditions, year-to-year climatic variability can result in substantial shifts in meteorological conditions between two wheat seasons (Fig. 2). Future experiments could look at how the accumulation of several more extreme seasons affects cropping systems and how the occurrence of individual more extreme events within one growing season impacts crop performance.

Conclusion

This study provides new insights into the complex interplay between climate change, soil management practices, and winter wheat performance. The observed phenological advances, increased yields in low-input soils (S1) as compared to high-input soil (S2), and natural stress adaptation mechanisms such as proline and silicon accumulation highlight the capacity of crops to respond to future climate conditions. However, additional research would be needed to better understand why in some soil conditions (like S2 in this study) the CO_2-fertilization effect remains limited and yields decrease under future climates. This may be linked to higher nutrient immobilisation in high organic input systems compared to low input systems, which could also explain the lower risk of nitrate leaching in S2. To ensure sustainable performance of future cropping systems it would therefore be key to further develop management practices that allow fertiliser application and nutrient buffering without necessarily increasing organic inputs because of their enhanced CO_2-emissions and nutrient immobilisation. Fertigation or biological nitrification inhibition should be tested under future meteorological conditions to assess their potential to optimise nutrient stability and use efficiency in agroecosystems. This study further highlights the need to investigate links between altered soil processes and plant diseases under future climates, where shorter cropping cycles may provide an opportunity to break disease cycles with inter-cropping.

Acknowledgments

This research is part of the BIOFAIR project (www.biofair.uliege.be) funded through the 2019-2020 BiodivERsA joint call for research proposals, under the BiodivClim ERA-Net COFUND programme, and with the funding organisations Agence Nationale de la Recherche (ANR, France; ANR-20-EBI5-0002), Agencia Estatal de Investigación (AEI, Spain, PCI2020-120713-2), Deutsches Zentrum fuer Luft- und Raumfahrt Projektträger (DLR-PT, Germany), Fonds de la Recherche Scientifique (FNRS, Wallonia, Belgium; R.8001.20), Fonds voor Wetenschappelijk Onderzoek - Vlaanderen (FWO, Flanders, Belgium; FWO ERA-NET G0H7320N) & Schweizerischer Nationalfonds (SNF, Switzerland; 31BD30_193869).

References

Asseng S, Martre P, Maiorano A, et al. (2019) Climate change impact and adaptation for wheat protein. *Glob Change Biol*. 25: 155–173. https://doi.org/10.1111/gcb.14481.

Baker, N. R. (2008). Chlorophyll fluorescence: a probe of photosynthesis in vivo. Annual Review of Plant Biology, 59, 89-113. https://doi.org/10.1146/annurev.arplant.59.032607.092759.

BBA (1999). F. 01 Deutscher Vorschlag für eine EPPO-Richtlinie zur Prüfung der Wirksamkeit von Saatgutbehandlungsmitteln gegen luft- und bodenbürtige Krankheitserreger an Getreide. Biologischen Bundesanstalt für Land- und Forstwirtschaft, Braunschweig, Germany.

Benton, T.G. (2020). Running AMOC in the farming economy. Nat Food 1, 22–23 https://doi.org/10.1038/s43016-019-0017-x.

Birkhofer K, Fliessbach A, Gavín-Centol MP, Hedlund K, Ingimarsdóttir M, Jørgensen HB, Kozjek K, Meyer S, Montserrat M, Moreno SS, Laraño JM, Scheu S, Serrano-Carnero D, Truu J, Kundel D. (2021). Conventional agriculture and not drought alters relationships between soil biota and functions. Sci Rep.11(1):23975. doi: 10.1038/s41598-021-03276-x.

Buckeridge, K.M., Mason, K.E., McNamara, N.P. et al. (2020). Environmental and microbial controls on microbial necromass recycling, an important precursor for soil carbon stabilization. Commun Earth Environ 1, 36 https://doi.org/10.1038/s43247-020-00031-4.

Campbell, CD, Chapman, SJ, Cameron, CM, Davidson, MS, Potts JM (2003). A rapid microtiter plate method to measure carbon dioxide evolved from carbon substrate amendments so as to determine the physiological profiles of soil microbial communities by using whole soil. Appl. Environ. Microbiol., 69 (6): 3593-3599. doi: 10.1128/AEM.69.6.3593-3599.2003.

Cao, D., Heughebaert, L. Boffel, L., Stove, C. & Van Der Straeten, D. (2024). Simultaneous quantification of seven B vitamins from wheat grains using UHPLC-MS/MS. Food Chemistry 139667. https://doi.org/10.1016/j.foodchem.2024.139667.

Cao Y et al. (2021). Organic-C quality as a key driver of microbial nitrogen immobilization in
soil: A meta-analysis. Geoderma 383, 114784.
https://doi.org/10.1016/j.geoderma.2020.114784.

Chakraborty, S. · Newton, A.C. (2011). Climate change, plant diseases and food security: an
overview. Plant Pathol. 60:2-14

Cigale: Consultation de l'Information Géographique pour l'Agriculture, Les ressources
naturelles et l'Environnement en Wallonie V4.3.2 (api-4.8), Service public de Wallonie, Here
[Navteq], AGDP et IGN (2021) Carte des Sols de Wallonie.
http://geoapps.wallonie.be/Cigale/Public/#VIEWER=CNSW#BBOX=158223.8715386281,1
78729.12088246012,135715.83238417906,151881.90638299374. Accessed on 01/11/23.

Dickin E & Wright D (2008). The effects of winter waterlogging and summer drought on the
growth and yield of winter wheat (Triticum aestivum L.), European Journal of Agronomy
28(3): 234-244, https://doi.org/10.1016/j.eja.2007.07.010.

ESDAC (2020). Soil microbial biomass and respiration. European Soil Data Centre
(ESDAC), European Commission, Joint Research Centre.
https://esdac.jrc.ec.europa.eu/content/soil-microbial-biomass-and-respiration.

European Environment Agency (2024). European Climate Risk Assessment (EUCRA). EEA
Report No 1/2024. Luxembourg: Publications Office of the European Union.
doi:10.2800/20424. https://www.eea.europa.eu/publications/european-climate-risk-
assessment.

Folegatti, M.V., Blanco, F.F., Boaretto, R.M. and Boaretto, A.E. (2005) Calibration of cardy-
ion meters to measure nutrient concentrations in soil solution and in plant sap. Sci. Agric.
62(1): 8-11. https://doi.org/10.1590/S0103-90162005000100002.

Food and Agriculture Organization (FAO). (2021). FAOSTAT Statistical Database. Rome:
FAO. Retrieved from
https://www.fao.org/faostat/fr/#rankings/commodities_by_regions_imports 23/07/2023.

Farooq, M., Hussain, M., & Siddique, K. H. M. (2009). Drought stress in plants: An overview
on its consequences and management strategies. Understanding Drought Impact on Crop
Yield. DOI:10.1007/978-3-642-32653-0_1.

Fox J, Weisberg S (2019). An R Companion to Applied Regression, Third edition. Sage,
Thousand Oaks CA. https://www.john-fox.ca/Companion/.

Juroszek, P., & von Tiedemann, A. (2013). Climate change and potential future risks through
wheat diseases: a review. European Journal of Plant Pathology, 136(1), 21-33.
https://doi.org/10.1007/s10658-012-0144-9.

Chakraborty, S., & Newton, A. C. (2011). Climate change, plant diseases and food security:
an overview. Plant Pathology, 60(1), 2-14. https://doi.org/10.1111/j.1365-3059.2010.02411.x.

Gentile AR, Barceló-Cordón S, Van Liedekerke M (2009). Soil Country Analyses Belgium.
EUR 23959 EN/2. European Commission, Joint Research Centre, Institute for Environment
and Sustainability. https://publications.jrc.ec.europa.eu/repository/bitstream/JRC53315/
publication_scr_belgium_final.pdf.

Ghini, R., Bettiol, W., & Hamada, E. (2011). Diseases in tropical and plantation crops as
affected by climate changes: current knowledge and perspectives. Plant Pathology, 60(1),
122-132. https://doi.org/10.1111/j.1365-3059.2010.02403.x.

Gaines, T. P., & Gaines, S. T. (1994). Soil texture effect on nitrate leaching in soil percolates.
Communications in Soil Science and Plant Analysis, 25(13–14), 2561–2570.
https://doi.org/10.1080/00103629409369207.

Gardiner, B et al. (2016) Review: Wind impacts on plant growth, mechanics and damage.
Plant Science 245 : 94-118. https://doi.org/10.1016/j.plantsci.2016.01.006.

Gilhespy, S. L., Anthony, S., Cardenas, L., Chadwick, D., del Prado, A., Li, C., Misselbrook,
T., Rees, R. M., Salas, W., Sanz-Cobena, A., Smith, P., Tilston, E. L., Topp, C. F. E., Vetter,
S., & Yel-ripati, J. B. (2014). First 20 years of DNDC (DeNitrification DeComposition):
Model evolution. Ecological Modelling, 292, 51-62.

Giltrap, D. L., Li, C., & Saggar, S. (2010). DNDC: A process-based model of greenhouse gas
fluxes from agricultural soils. Agriculture, Ecosystems & Environment, 136(3-4), 292-300.
https://doi.org/10.1016/j.agee.2009.06.014.

Giot, O., Termonia, P., Degrauwe, D., De Troch, R., Caluwaerts, S., Smet, G., Berckmans, J.,
Deckmyn, A., De Cruz, L., De Meutter, P., Duerinckx, A., Gerard, L., Hamdi, R., Van den
Bergh, J., Van Ginderachter, M., & Van Schaeybroeck, B. (2016). Validation of the ALARO-0
model within the EURO-CORDEX framework. Geoscientific Model Development, 9(3),
1143-1152. https://doi.org/10.5194/gmd-9-1143-2016.

Govindasamy, P., Muthusamy, S. K., Bagavathiannan, M., Mowrer, J., Jagannadham, P. T. K.,
Maity, A., ... & Tiwari, G. (2023). Nitrogen use efficiency - a key to enhance crop
productivity under a changing climate. Frontiers in Plant Science, 14, 1121073.
https://doi.org/10.3389/fpls.2023.1121073.

Graves S, Piepho H, Dorai-Raj LS (2024). multcompView: Visualizations of Paired
Comparisons. R package version 0.1-10. https://CRAN.R-
project.org/package=multcompView.

Hamamoto, T., Nhamo, N., Chikoye, D. *et al.* (2022). Responses of CO_2 emissions and soil
microbial community structures to organic amendment in two contrasting soils in Zambia. *Sci
Rep* **12**, 6368 https://doi.org/10.1038/s41598-022-10368-9.

Harkness, C., Semenov, M. A., Areal, F., Senapati, N., Trnka, M., Balek, J., & Bishop, J.
(2020). Adverse weather conditions for UK wheat production under climate change.
Agricultural and Forest Meteorology, 282, 107862.
https://doi.org/10.1016/j.agrformet.2019.107862.

Hayat S, Hayat Q, Alyemeni MN, Wani AS, Pichtel J, Ahmad A. (2012) Role of proline under
changing environments: a review. Plant Signal Behav. (11):1456-66. doi:10.4161/psb.21949.

Iglesias, J., Castillejo, J., & Castro, R. (2003). The effects of repeated applications of the
molluscicide metaldehyde and the biocontrol nematode Phasmarhabditis hermaphrodita on
molluscs, earthworms, nematodes, acarids and collembolans: a two-year study in north-west
Spain. Pest Management Science: Formerly Pesticide Science, *59*(11), 1217-1224.

IPCC, Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. (2021) Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Jansson, J.K., Hofmockel, K.S. (2021) Soil microbiomes and climate change. Nat Rev Microbiol 18, 35–46 (2020). https://doi.org/10.1038/s41579-019-0265-7.Khan A, Ahmad M, Ahmed M, Iftikhar Hussain M. Rising Atmospheric Temperature Impact on Wheat and Thermotolerance Strategies. Plants 10(1):43. https://doi.org/10.3390/plants10010043.

Jia, K.; Yang, Y.Z.; Dong, G.L.; Zhang, C.; Lang, T.T. (2021) Variation and determining factor of winter wheat water requirements under climate change. Agric. Water Manag. 254, 106967. DOI: 10.1016/j.agwat.2021.106967.

Kaur G. & Asthir B. (2015) Proline: a key player in plant abiotic stress tolerance. Biologia plantarum 59 (4): 609-619. https://doi.org/10.1007/s10535-015-0549-3.

Kelley, LA et al. (2024) Changes in soil N2O emissions and nitrogen use efficiency following long-term soil carbon storage: Evidence from a mesocosm experiment. Agriculture, Ecosystems & Environment 370, 109054. https://doi.org/10.1016/j.agee.2024.109054.

Kim YH, Khan AL, Waqas M, Lee IJ. (2017) Silicon Regulates Antioxidant Activities of Crop Plants under Abiotic-Induced Oxidative Stress: A Review. Front Plant Sci. 8:510. doi: 10.3389/fpls.2017.00510.

Kornhuber, K., Lesk, C., Schleussner, C.F. et al. (2023). Risks of synchronized low yields are underestimated in climate and crop model projections. Nat Commun 14, 3528 https://doi.org/10.1038/s41467-023-38906-7.

Kumar H, Chugh V, Kumar M, Gupta V, Prasad S, Kumar S, Singh CM, Kumar R, Singh BK, Panwar G, Kumar M. (2023) Investigating the impact of terminal heat stress on contrasting wheat cultivars: a comprehensive analysis of phenological, physiological, and biochemical traits. Front Plant Sci.14:1189005. doi: 10.3389/fpls.2023.1189005.

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) "lmerTest Package: Tests in Linear Mixed Effects Models." Journal of Statistical Software 82(13): 1-26. doi:10.18637/jss.v082.i13.

Kuzyakov, Y. and Xu, X. (2013), Competition between roots and microorganisms for nitrogen: mechanisms and ecological relevance. New Phytol, 198: 656-669. https://doi.org/10.1111/nph.12235.

Lamba, K., Kumar, M., Singh, V. et al. (2023). Heat stress tolerance indices for identification of the heat tolerant wheat genotypes. Sci Rep 13, 10842 https://doi.org/10.1038/s41598-023-37634-8.

Lamprea Pineda, P. A., Bauters, M., Verbeeck, H., Baez, S., Barthel, M., Bodé, S., and Boeckx, P. (2021). Ideas and perspectives: patterns of soil CO_2, CH_4, and N_2O fluxes along an altitudinal gradient – a pilot study from an Ecuadorian neotropical montane forest, Biogeosciences, 18, 413–421, https://doi.org/10.5194/bg-18-413-2021.

Lenth R (2024) emmeans: Estimated Marginal Means, aka Least-Squares Means. R package
version 1.10.5. https://CRAN.R-project.org/package=emmeans.

Li, C., Frolking, S., & Harriss, R. (1992a). Modeling carbon biogeochemistry in agricultural
soils. Global Biogeochemical Cycles, 6(4), 237-254. https://doi.org/10.1029/92GB01402.

Li, C., Frolking, S., & Harriss, R. (1992b). Modeling N2O emissions from agricultural fields.
Global Biogeochemical Cycles, 6(4), 287-306. https://doi.org/10.1029/92GB02092.

Li, C.S. (2000). Modeling trace gas emissions from agricultural ecosystems. Nutrient Cycling
in Agroecosystems, 58(1-3), 259-276. https://doi.org/10.1023/A:1009859006242.

Li, W., Pacheco-Labrador, J., Migliavacca, M. *et al.* (2023). Widespread and complex
drought effects on vegetation physiology inferred from space. *Nat Commun* 14, 4640
https://doi.org/10.1038/s41467-023-40226-9.

Liang, S., Zhang, X., Sun, N. et al. (2019). Modeling crop yield and nitrogen use efficiency in
wheat and maize production systems under future climate change. Nutr Cycl Agroecosyst
115, 117–136 https://doi.org/10.1007/s10705-019-10013-4.

Livre Blanche Céréales (2022) Gembloux Agro-Bio Tech, Université de Liège, Belgium.
https://livre-blanc-cereales.be/wp-content/uploads/2022/04/Complet.pdf.

Lowe NM. (2021) The global challenge of hidden hunger: perspectives from the field.
Proceedings of the Nutrition Society 80(3):283-289. doi:10.1017/S0029665121000902.

Maxwell, K. & Johnson, GN (2000). Chlorophyll fluorescence—a practical guide, Journal of
Experimental Botany (51) 345, 659–668. https://doi.org/10.1093/jexbot/51.345.659.

Meier, U. (ed.). (2001). Growth Stages of Mono- and Dicotyledonous Plants. BBCH
Monograph. 2nd edition. Federal Biological Research Centre for Agriculture and Forestry,
Germany. DOI: 10.5073/20180906-074619.

Meshcherskaya A. V. & Blazhevich V. G. (1997) The Drought and Excessive Moisture
Indices in a Historical Perspective in the Principal Grain-Producing Regions of the Former
Soviet Union. Journal of Climate 10(10): 2670–2682. https://doi.org/10.1175/1520-
0442(1997)010<2670:TDAEMI>2.0.CO;2.

Miedaner, T. (2018). Breeding strategies for improving plant resistance to diseases. In
Advances in Plant Breeding Strategies: Agronomic, Abiotic and Biotic Stress Traits (pp. 561-
599). Springer, Cham. https://doi.org/10.1007/978-3-319-22518-0_15.

Moitinho, M.R., Teixeira, D., Bicalho, E. et al. (2021). Soil CO2 emission and soil attributes
associated with the microbiota of a sugarcane area in southern Brazil. Sci Rep 11, 8325
https://doi.org/10.1038/s41598-021-87479-2.

Muñoz, C., Paulino, L., Monreal, C., & Zagal, E. (2010). Greenhouse gas (CO2 and N2O)
emissions from soils: a review. Chilean journal of agricultural research, 70(3), 485-497.
http://www.bioline.org.br/pdf?cj10053.

Myers, S., Zanobetti, A., Kloog, I. et al. Increasing CO2 threatens human nutrition. Nature
510, 139–142 (2014). https://doi.org/10.1038/nature13179.

Pais IP, Moreira R, Semedo JN, Ramalho JC, Lidon FC, Coutinho J, Maçãs B, Scotti-Campos
P. (2022) Wheat Crop under Waterlogging: Potential Soil and Plant Effects. Plants
(Basel);12(1):149. https://doi:10.3390/plants12010149.

Palma-Guerrero, J et al. (2021) Take-All Disease: New Insights into an Important Wheat
Root Pathogen. Trends in Plant Science, Volume 26, Issue 8, 836 – 848.
https://doi.org/10.1016/j.tplants.2021.02.009.

Pausch, J., Holz, M., Zhu, B. & Cheng, W. (2024) Rhizosphere priming promotes plant
nitrogen acquisition by microbial necromass recycling. Plant, Cell & Environment, 47, 1987–
1996. https://doi.org/10.1111/pce.14858.

Porter JR, Xie L, Challinor AJ, Cochrane K, Howden SM, Iqbal MM, Lobell DB, Travasso
MI. (2014). Food security and food production systems. In: Field CB, Barros VR, Dokken
DJ, Mach KJ, Mastrandrea MD, Bilir TE, Chatterjee M, Ebi KL, Estrada YO, Genova RC,
Girma B, Kissel ES, Levy AN, MacCracken S, Mastrandrea PR, White LL, (Eds.). Climate
Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects.
Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental
Panel on Climate Change Cambridge University Press, Cambridge, United Kingdom and
New York, NY, USA, pp. 485-533.

Quan, Q., He, N., Zhang, R. et al. (2024). Plant height as an indicator for alpine carbon
sequestration and ecosystem response to warming. Nat. Plants 10, 890–900.
https://doi.org/10.1038/s41477-024-01705-z.

R Core Team (2024) R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Reay, D., Davidson, E., Smith, K. et al. (2012). Global agriculture and nitrous oxide
emissions. Nature Clim Change 2, 410–416. https://doi.org/10.1038/nclimate1458.

Rezaei, E. E., Siebert, S., Hüging, H., & Ewert, F. (2018). Climate change effect on wheat
phenology depends on cultivar change. Scientific Reports, 8, 4891.
https://doi.org/10.1038/s41598-018-23101-2.

Riedesel L, Möller M, Horney P, Golla B, Piepho HP, Kautz T, Feike T. (2023) Timing and
intensity of heat and drought stress determine wheat yield losses in Germany. PLoS
One;18(7):e0288202. doi: 10.1371/journal.pone.0288202.

Rocci et al. (2024) Aligning theoretical and empirical representations of soil carbon-to-
nitrogen stoichiometry with process-based terrestrial biogeochemistry models. Soil Biology
and Biochemistry 189: 109272. https://doi.org/10.1016/j.soilbio.2023.109272.

Roy J, Rineau F, De Boeck HJ, Nijs I, Pütz T, Abiven S, Arnone JA 3rd, Barton CVM,
Beenaerts N, Brüggemann N, Dainese M, Domisch T, Eisenhauer N, Garré S, Gebler A,
Ghirardo A, Jasoni RL, Kowalchuk G, Landais D, Larsen SH, Leemans V, Le Galliard JF,
Longdoz B, Massol F, Mikkelsen TN, Niedrist G, Piel C, Ravel O, Sauze J, Schmidt A,
Schnitzler JP, Teixeira LH, Tjoelker MG, Weisser WW, Winkler B, Milcu A. (2021) Ecotrons:
Powerful and versatile ecosystem analysers for ecology, agronomy and environmental
science. Glob Chang Biol.(7):1387-1407. doi: 10.1111/gcb.15471.

Schmidt A, Hines J, Türke M, Buscot F, Schädler M, Weigelt A, Gebler A, Klotz S, Liu T,
Reth S, Trogisch S, Roy J, Wirth C, Eisenhauer N. (2021) The iDiv Ecotron-A flexible
research platform for multitrophic biodiversity research. Ecol Evol.11(21):15174-15190. doi:
10.1002/ece3.8198.

Schröder, J.J., Schulte, R.P.O., Creamer, R.E., Delgado, A., van Leeuwen, J., Lehtinen, T.,
Rutgers, M., Spiegel, H., Staes, J., Tóth, G. and Wall, D.P. (2016), The elusive role of soil
quality in nutrient cycling: a review. Soil Use Manage, 32: 476-486.
https://doi.org/10.1111/sum.12288.

Seethepalli, A., Dhakal, K., Griffiths, M., Guo, H., Freschet, G. T., York, L. M. (2021).
RhizoVision Explorer: Open-source software for root image analysis and measurement
standardization. AoB PLANTS, plab056, https://doi.org/10.1093/aobpla/plab056.

Shahwani, A. R., Baloch, S. U., Baloch, S. K., Mengal, B., Bashir, W., Baloch, H. N., ... &
Mengal, A. (2014). Influence of seed size on germinability and grain yield of wheat (Triticum
aestivum L.) varieties. Journal of Natural Sciences Research, 4(23), 147-155.

Shewry, PR. (2013) Improving Grain Quality: Wheat, In: Seed Genomics, ed.PW. Becraft,
pp. 159-178. John Wiley & Sons, Inc. https://doi.org/10.1002/9781118525524.ch9.

Spormann S, Nadais P, Sousa F, Pinto M, Martins M, Sousa B, Fidalgo F, Soares C. (2023)
Accumulation of Proline in Plants under Contaminated Soils-Are We on the Same Page?
Antioxidants (Basel). 12(3):666. doi: 10.3390/antiox12030666.

Stekhoven D. J., & Buehlmann, P. (2012). MissForest - non-parametric missing value
imputation for mixed-type data. Bioinformatics 28(1), 112-118. DOI:
10.1093/bioinformatics/btr597.

Tang, J., C. Körner, H. Muraoka, S. Piao, M. Shen, S. J. Thackeray, and X. Yang. (2016).
Emerging opportunities and challenges in phenology: a review. Ecosphere 7(8):e01436.
10.1002/ecs2.1436.

Tilman, D., Balzer, C., Hill, J., & Befort, B. L. (2011). Global food demand and the
sustainable intensification of agriculture. *Proceedings of the National Academy of Sciences*,
108(50), 20260-20264. https://doi.org/10.1073/pnas.1116437108.

Tipping, M. E., & Bishop, C. M. (1999). Probabilistic Principal Component Analysis. Journal
of the Royal Statistical Society: Series B (Statistical Methodology), 61(3), 611-622.
https://doi.org/10.1111/1467-9868.00196.

Venterea, R.T. (2007). Nitrite-driven nitrous oxide production under aerobic soil conditions:
kinetics and biochemical controls. Global Change Biology 13: 1798-1809.
https://doi.org/10.1111/j.1365-2486.2007.01389.x

Vonk, W.J., van Ittersum, M.K., Reidsma, P. *et al.* (2020). European survey shows poor
association between soil organic matter and crop yields. *Nutr Cycl Agroecosyst* 118, 325–334.
https://doi.org/10.1007/s10705-020-10098-2.

Wang M, Gao L, Dong S, Sun Y, Shen Q, Guo S. (2017) Role of Silicon on Plant-Pathogen
Interactions. Front Plant Sci.;8:701. doi: 10.3389/fpls.2017.00701.

Wang, M., Wang, R., Mur, L.A.J. et al. (2021). Functions of silicon in plant drought stress
responses. Hortic Res 8, 254 https://doi.org/10.1038/s41438-021-00681-1.

Wilcox J & Makowski D (2014) A meta-analysis of the predicted effects of climate change on
wheat yields using simulation studies. Field Crops Research 156: 180-190.
https://doi.org/10.1016/j.fcr.2013.11.008.

Yemm EW, Williams AJ (1954). The estimation of carbohydrates in plant extracts by
anthrone. Biochem J. 57(3):508-14. https://doi.org/0.1042/bj0570508.

Zhang, G., Ming, B., Shen, D., Xie, R., Hou, P., Xue, J., Wang, K., & Li, S. (2021).
Optimizing Grain Yield and Water Use Efficiency Based on the Relationship between Leaf
Area Index and Evapotranspiration. Agriculture, 11(4), 313.
https://doi.org/10.3390/agriculture11040313.

Zhao, H., Zhang, L., Kirkham, M.B. *et al.* (2022). U.S. winter wheat yield loss attributed to
compound hot-dry-windy events. *Nat Commun* 13, 7233 https://doi.org/10.1038/s41467-022-
34947-6.

Display items for

Low-input soil management increases yield and decreases CO$_2$-emissions but aggravates risk of nitrate leaching and diseases in winter wheat cropping systems under climate change

Running title: Time travelling with *Triticum*

Graphical abstract

Main: 4 figures & 2 tables

1. Figure 1: Experimental set-up

2. Figure 2: Climate scenarios

3. Table 1: Physicochemical characterisation of the two soils

4. Figure 3: Environmental, soil and crop parameters

5. Table 2: Yield components

6. Figure 4: Ordination of parameters characterizing each cropping system (pPCA)

Graphical abstract

Yield (t ha⁻¹)

| 2013 | 2068 | 2085 |

S1: 3.79±0.51
S2: 3.50±0.34

S1: 4.76±0.37 (+26%)
S2: 3.71±0.32 (+6%)

S1: 5.09±0.50 (+34%)
S2: 3.05±0.59 (-13%)

Proline↑
(S1)

Silicon↑
(S2)

Total root
length ↑
(S2)

*Gaeumanno-
myces tritici* ↑
(S1)

Figure 1: Experimental set-up. Left: Schematic representation of the TERRA-Ecotron which consists of six controlled environment rooms (CERs), in each of which n=9 cubes are placed. Each cube measures 50x50x50cm and contains a soil monolith originating either from the field with low organic inputs ("S1") or the field with high organic inputs ("S2"). In each CER one soil cube is unplanted ("C_") while all other cubes are sown with *Triticum aestivum* var. Asory. In each of the CERs one meteorological condition is employed representing either one of the years 2013, 2068 or 2085, with each year being replicated in two CERs (2013: CER1, CER4, 2068: CER2, CER5, 2085: CER3, CER6). Darker shadows of the cubes indicate positioning of a scale underneath the cube. Right: View inside one of the CERs during the experiment: Each CER is equipped with a lightening system combing plasma, halogen and LED lamps which can reach maximum photon fluxes of 1200 µmol m^{-2} s^{-1}. Sensors for photosynthetic active radiation (PAR) and irradiance are located at canopy height.

2013

CER 1

S2	C_S2	S1
S2	S1	S1
S2	S1	S2

CER 4

C_S1	S2	X-S2
S1	S2	S2
S1	S2	S1

2068

CER 2

S1	S1	S2
S2	S2	S1
S1	C_S2	S2

CER 5

S1	S2	S2
S1	S1	S2
C_S1	S2	S1

2085

CER 3

S2	S2	S1
C_S1	S1	S2
S1	S1	S2

CER 6

S2	S1	S1
S1	S2	S2
S1	S2	C_S2

Figure 2: Climate scenarios. Left: Wheat agroecosystems were exposed to the meteorological conditions of three years representing the present climate (2013), the near future (2068) and a farther future (2085). The three years align on a continuous gradient of hydrothermal index with increasing temperature, precipitation and atmospheric CO_2-concentrations. The climate scenarios are based on continuous meteorological observations from the Ernage weather station (50°34'33"N, 4°43'1"E, Belgium, since 1980) and the predicted future climates were simulated using the Alaro-0 model. The historic observations (HO) cover the period 1981-2017 (blue) and the model ran for the Representative Concentration Pathway (RCP) scenario 8.5 W m^{-2} for the two time periods 2041-2070 (green) and 2070-2100 respectively (rosé). Each dot represents a year and the ellipses represent the 95% confidence interval of the three periods. Right: Table summarizing the key climatic parameter of the selected years.

Climatic parameters	2013	2068	2085
Mean CO_2 ppm	420	550	775
Mean radiation µmol m^{-2} s^{-1}	51.81	53.98	54.49
Mean temperature °C d^{-1}	7.6	10.2	12.1
Mean precipitation mm d^{-1}	2.1	2.4	3
Hydrothermal index (rel.)	4	4.5	4.7
Maximum temperature °C	26.5	23.4	22.2
Minimum temperature °C	-7.6	-5.8	3.1
n days temperature ≤ 4°C	94	40	3
n days temperature ≥ 22°C	9	7	2
Maximum precipitation mm	46.3	39.0	47.3
Minimum precipitation mm	0.0	0.0	0.0
n days precipitation ≤ 0mm	135	4	3
n days precipitation ≥ 4mm	57	62	72

Table 2: Physicochemical characterisation of the two soil types at the beginning of the experiment, one composite sample per soil type, sampling depth: 0-20cm.

	C (g kg⁻¹)	N (g kg⁻¹)	$\frac{C}{N}$	P (mg 100g⁻¹)	K (mg 100g⁻¹)	Mg (mg 100g⁻¹)	Ca (mg 100g⁻¹)	pH (H₂0)	Humus (%)	Clay (%)	Silt (%)	Sand (%)	Soil classification
Soil One (S1) "low-input"	9.92	0.94	10.5	13.60	31.20	8.37	218.31	8.04	1.98	12.15	67.13	20.72	Aba(b)0 Silt loam
Soil Two (S2) "high-input"	22.01	2.09	10.5	39.79	72.51	14.70	534.42	8.08	4.23	13.55	78.85	7.60	Aba(b)0 Silt loam

Figure 3: Environmental, crop and soil parameters of two management systems (S1: continuous line/fill and S2: dotted line/hatched area) in the meteorological conditions of the years 2013 (blue), 2068 (green) and 2085 (rosé). Details of each measurement method and replication for each parameter in supplementary material SM2. A: Photon flux µmol m^{-2} s^{-1}, B: Air temperature °C day^{-1},C: Soil temperature °C day^{-1}, D: Soil relative humidity % day^{-1}, E: BBCH scale plant growth stages, F: Plant height cm, G: Leaf area index (LAI), H: Precipitation mm day^{-1}, I: Maximum quantum efficiency of photosystem II (Fv/Fm), J: Leaf silicon mg kg^{-1}, K: Leaf proline µg g^{-1}, L: Take-all index (% *Gaeumannomyces tritici* infestation), M: Nitrate (NO$_3$) mg L^{-1}in aqueous soil solution, N: Glucose equivalent in aqueous soil solution mg ml^{-1}, O: Total root length (TRL) mm for 0-10cm soil depth, P: Microbial biomass CO$_2$-C mg g^{-1}, Q: Grain yield t ha^{-1} at 14% grain humidity, R: Annual soil CO$_2$-C budget kg ha^{-1} yr^{-1}, S: Annual soil N$_2$O-N budget kg ha^{-1} yr^{-1}, Annual soil N leaching kg ha^{-1} yr^{-1}, A-D, H: Daily measurements, E: BBCH growth stages evaluated for all cubes and only one value noted per modality, F-G, I-Q: Dots represent means ± SD of n=4 or n=8, R-T: Data modelled with DNDC and only one value per modality.

Table 3: Yield components for winter wheat (*Triticum aestivum* var. Asory) grown in two differentially managed soil types (low input S1 and high input S2) in the meteorological conditions of the years 2013, 2068 and 2085. Probability values following analysis of variance with asterisks indicating significance levels at <0.001***, ≤0.05*, >0.05 not significant (ns) and letters indicating grouping based on TukeyHSD test.

	2013		2068		2085		
	S1	**S2**	**S1**	**S2**	**S1**	**S2**	**p-value**
Grain yield t ha^{-1}	3.79±0.51 ab	3.50±0.34 ab	4.76±0.37 ab	3.71±0.32 ab	5.09±0.50 a	3.05±0.59 b	0.0375 *
Thousand grain weight	31.12±2.35 a	30.45±6.85 a	35.62±2.77 ab	47.13±8.48 c	45.18±1.95 bc	41.29±3.98 abc	0.0004 ***
Grain nitrogen (%)	2.26±0.20	2.25±0.18	2.00±0.11	1.87±0.11	2.11±0.39	1.97±0.22	0.13 (ns)
Number of heads cube^{-1}	108±20	108±19	111±11	88±9	118±20	101±27	0.36 (ns)
Total fresh weight straw g cube^{-1}	121.54±17.53 a	131.74±52.51 a	176.55±22.84 a	186.71±40.81 a	327.29±71.40 b	206.21±49.92 a	0.00009 ***

Figure 4: Ordination of agronomic and environmental parameter in probabilistic principle component analysis (pPCA) and clustering by simulated year and soil type. Main parameters which distinguish the groups: 2013.S1: glucose equivalent, take-all index; 2013.S2: leaf area index; 2068.S1: foliar proline, microbial biomass carbon, total root length; 2068.S2: basal CO_2, maximum quantum efficiency PSII; 2085.S1: plant height, aq. NO_3, total root length; 2085.S2: foliar silicon, plant height (please see supplementary material SM4 for further details on vector loadings).

Supplementary material for

Low-input soil management increases yield and decreases CO$_2$-emissions but aggravates risk of nitrate leaching and diseases in winter wheat cropping systems under climate change

Running title: Time travelling with *Triticum*

SM1: a: Soil management history, b: monolith sampling, c: fertilization during Ecotron experiment

SM2: Details on methods of empirical and modelled parameter to quantify agronomic performance and environmental impact

SM3: DNDC model evaluation

SM4: Loadings for the main vector of each modality (Year.Soil)

SM1a: Detailed soil management

Soil type 1 (S1) 50°38'35.1474"N, 4°37'22.0123"E

Date	Crop / intervention	Dose	Comments/Details
2015	Wheat		
2016	Chicories		
2017	Wheat		
07/08/17	1 pass of harrowing		
24/08/17	Icorporation of Haspargit® and scums		Haspargit®, low-chlorine potassium fertilizer, contains sulfur and calcium
05/03/18	350 kg/ha of Sulfonitrate N26+ 31 SO3(commercial name)	350 kg/h	
06/04/18	350kg/ha Nitrogen (N27 %)	350 kg/h	
15/04/18	CARYX®	1,25 L/ha	Growth Regulator + Fungicide
04/05/18	Tangus Gold	0,5 kg/ha	Fungicide
14/07/18	Canola harvest		
21/08/18	Harrowing		
03/10/18	Tillage		
03/10/18	Plantation escourgeon multi		
06/11/18	HEROLD®	0,5 L /ha	HEROLD®, contact and residual broad-spectrum herbicide based on the leading grass-weed active ingredient flufenacet, for pre- and post-crop emergence application
06/11/18	Chlortoluron	1,2 L/ha	Herbicide
06/11/18	Patriot protect	0,32 L/ha	Patriot® Selective Herbicide
26/02/19	Nitrogen N27 %	250 kg/ha	
28/03/19	KANTIK®	1 L/ha	Fungicide
28/03/19	MEDAX ® TOP	1,2 L/ha	MEDAX® TOP is a growth regulator for cereals
28/03/19	MAGNESIE	2,5 kg/ ha	
02/04/19	SULFAZOTE 22%N (28,6 U)	200 L/ha	
20/04/19	TERPAL ®	1,2L /ha	Terpal® shortens the internodes and strengthens the stem wall, treated cereal crops become more resistant to lodging
20/04/19	INPUT	0,6L /ha	
02/05/19	VELOGY ERA	0,7L / ha	FUNGICIDE
02/05/19	BRAVO	1,2L /ha	FUNGICIDE
02/05/19	MAGNESIE	2,5KG /ha	
02/05/19	Patriot protect	0,2L /ha	Patriot® Selective Herbicide
08/07/19	Harverst escourgeon multi		
24/08/19	Glyfall plus	6L/ha	Total herbicide
02/09/19	Harrowing		
09/09/19	Harrowing		
11/09/19	Sowing seeds (radish, clover, phacelia)	12kg/ha	
2021	Sugar beet/wheat culture		Variety BTS 4860 + KWS Tessilia + Caprianna kws + Bts 3480 (6 boxes of each)

Date	Operation	Rate	Description
03/04/20	Sugar beet sowing		
21/08/20	Fertilizer 0-5-16+15 CaO+2 MgO+12 EB	1113 kg/ha	
02/09/20	Harrowing		
07/09/20	Harrowing		
07/09/20	Liquid nitrogen	120 l/ha	
07/09/20	Green manure seeding 15kg/ha (base 5kg + early 10kg) brand Lemken		
31/10/20	Fast plow		
06/12/20	Tillage		
23/03/21	7 So3	800 kg/ha	Fertilizer
23/03/21	Nitrogen N27	150 kg/ha	
23/03/21	7 So3	800 kg/ha	Fertilizer
26/03/21	Liquide nitrogen	350 L/ha	
02/04/21	Soil preparation		
04/04/21	Beetix	1,75 L/kg	Herbicide
04/04/21	Centium®	0,05 L/kg	Herbicide
01/05/21	Dianal	0,9 L/ha	
01/05/21	Ethomat	0.3	Herbicide to control annual dicot weeds and annual grass weeds in sugar beet and fodder beet.
01/05/21	Safari®	15 gr/ha	Safari® herbicide for post-emergence control of broadleaf weeds in sugar beets, red beets, chicory, and endive
01/05/21	Goltix® Queen	1 L/ha	Goltix® Queen is a combination of the active ingredient metamitron, and an addition of the active ingredient quinmerac, long-lasting control on regrowth, effectively controlling a broad spectrum of weeds
01/05/21	Vegetop	0,5 L/ha	Additive to enhance herbicide performance.
01/05/21	Magnesia	2,3kg /ha	
07/05/21	Dianal	0,9 L/ha	
07/05/21	Ethomat	0.3	Herbicide to control annual dicot weeds and annual grass weeds in sugar beet and fodder beet.
07/05/21	Goltix® Queen	1 L/ha	Goltix® Queen is a combination of the active ingredient metamitron, and an addition of the active ingredient quinmerac, long-lasting control on regrowth, effectively controlling a broad spectrum of weeds
07/05/21	Vegetop	0,5 L/ha	Additive to enhance herbicide performance.
07/05/21	Magnesia	2,3kg /ha	
11/05/21	Hoeing		
29/05/21	Dianal	0,9 L/ha	
29/05/21	Ethomat	0.3	The product is intended to be used as herbicide to control annual dicot weeds and annual grass weeds in sugar beet and fodder beet.
29/05/21	Beetix	0,6 L/ha L/kg	Herbicide
29/05/21	Tamaris	0,6 L/ha	

29/05/21	Lenazar	150 gr/ha	Selective herbicide against annual dicotyledonous weeds
07/05/21	Vegetop	0,5 L/ha	Additive to enhance herbicide performance.
15/06/21	Centium®	0,05 L/kg	Herbicide
15/06/21	Frontier® Optima	0,5 L/ha	Frontier Optima is a soil herbicide primarily absorbed through the coleoptile of grasses as the seedling penetrates the upper layer of the soil.
15/06/21	Boron	3 L/ha	Fertilizier
15/06/21	Magnesia	2,3kg /ha	Fertilizier
05/08/21	Agora	0,35l /ha	Fungicide
05/08/21	Magnesia	5,9kg /ha	
01/09/21	Bicanta	1 L/ha	Bicanta: product to combat diseases in sugar beet cultivation, combination of two active ingredients 125 g/l of difenoconazole and 125 g/l of azoxystrobin, effective control against all major foliar diseases, including Cercospora, Stemphylium, downy mildew, Ramularia, and rust
01/09/21	Magnesia	5 kg/ha	
16/10/21	Harvest		
15/11/21	Harvest		
2022	Wheat		
18/10/21	Sowing wheat	153 kg/ha	Variety Campesino PMG:44 gr
06/11/21	Thin sowing wheat		
01-03-22	18 SO3	260 kg/ha	
15-03-22	Sulfazote 22%	200 l/ha	
23-03-22	Osmose	0,2 l/ha	
23-03-22	SAVVY®	20 gr/ha	Herbicide, 200g/kg of Metsulfuron-methyl
23-03-22	Sigma® Maxx	1,1 L/ha	Selective post-emergence herbicide in winter wheat, spring wheat, rye, triticale, and spelt against annual grasses and dicotyledons
23-03-22	Magnesia	2 kg/ha	
05-04-22	Nitrogene N27 %	194 kg/ha	Fertilizer
18-04-22	FERTILEADER® Trio		Improves plant resistance to stress, stimulates metabolism, improves plant photosynthesis, contains patented Seactiv® complex (trademark of Timak Agro)
18-04-22	Magnesia	2,3 kg/ha	
18-04-22	Osmose	0,2 l/ha	
18-04-22	Chlormequat	1 L/ha	Growth regulator, can control the vegetative growth of plants (i.e., root and leaf growth), promote the reproductive growth of plants (i.e., flower and fruit growth), and enhance the plant's fruit-setting rate.
27-04-22	Tebecur	2,7 kg/ha	All-round fungicide with efficacy over a period of several weeks, controlling numerous pathogens in various crops, effective on i.a. Fusarium, Alternaria, Septoria nodorum, rusts, Sclerotinia, powdery mildew, leaf spots
27-04-22	Osmose	0,2 l/ha	

27-04-22	Magnesia	2,7 kg/ha	
15-05-22	Nitrogen N38	31 L/ha	Foliar fertilizer
18-05-22	Osmose	0,2 l/ha	
18-05-22	Magnesia	3,6 kg/ha	
18-05-22	REVYTREX®	1,1 L/ha	New cereal fungicide based on Revysol®
18-05-22	COMET ®	0,4 L/ha	Preventive-action fungicide designed to combat foliar diseases in various cereals (eg fight dwarf rust)
29/07/22	Harverst wheat		
05/08/22	Harrowing		
05-08-22	KALCIPHOS P-K (S)	(0-5-15) 1000 kg/ha	
19/08/22	Cattle manure	15t/ha	
24/09/22	Green fertilizer	11,6 kg/ha	
24/09/22	Fast plowing		

Soil type 2 (S2) 50°39'12.8668"N, 4°38'10.7664"E

Date	Crop / intervention	Dose	Comments/Details
2017	Beet culture		
2018	Wheat culture		
2018	Cover crop after harvest (oats, mustard, radish, clover, sunflower, peas).		
2019	Potato culture		
2019-2020	wheat		
20/10/19	Mixed vegetation seeding	145 kg/ha	
02/03/20	Nitrogen N39	175 L/kg	
01/02/20	TMS	80 kg/ha	TMS, mineral CE-fertilizer, regulates microbial flora and organic matter evolution, improves chemical, physical and biological soil fertility.
16/03/20	Sigma® Flex	140 g/ha	Herbicide (not sure about yhe unit looks like g/ha)
16/03/20	Brodway	150 g/ha	Herbicide (not sure about yhe unit looks like g/ha)
16/03/20	Oil	1l/ha	
06/04/20	Cycocel		Cycocel is a plant growth regulator
18/04/20	Nitrogen N39	170 L/ha	
18/04/20	Humic and fulvic acids	1 L/ha	Fertilizer
16/05/20	Tebuzifo	04 L/ha	
16/05/20	TMF	1L/ha	Fertilizer
16/05/20	MgSo4	1,8 kg/ha	Fertilizer
20/05/20	Nitrogen N 27	200 kg/ha	Fertilizer
02/06/20	MgSo4	1,8 kg/ha	Fertilizer
02/06/20	TMF	1L/ha	Fertilizer
02/06/20	Opustear	0,35 l/ha	

Date	Product	Rate	Type
02/06/20	Sportitck	0,36 l/ha	
02/06/20	Kestrel	0,36 l/ha	
20/08/20	Cow manure	12000L	Fertilizer
	Cover crop seeding (phacelia, oats, radish, clover)		
	Grazing of cover crops, CRAW Janart trials		
08/03/21	Glypho	1L/ha	Herbicide
23/02/21	TMS	80 kg/ha	Fertilizer
03/03/21	Unreadable	300 kg	
18/03/21	Nitrogen	100L/ha	Fertilizer
18/03/21	Humactiv	3,5 L/ha	Root stimulant
11/05/21	Flax		
11/05/21	Kaei		
11/05/21	TMS	1 L/ha	
11/05/21	MgSo4	3kg/ha	
11/06/21	Rudis®	1 L/ha	Fungicide
11/06/21	TMF	0,4 L/ha	Fertilizer
11/06/21	éthéplum	4 L/ha	
15/06/21	TOPREX	0,3 L/ha	TOPREX, co-formulation of difenoconazole and paclobutrazol, use as plant growth regulator, also as a fungicide against phoma and light leaf spot in winter oilseed rape
15/06/21	Flax		
16/10/21	Wheat Sowing		
05/11/21	Liberator + Flufenacet	0,58 L/ha	Herbicides
05/11/21	Mateno® Duo	0,35 L/ha	Herbicides
26/02/22	TMS	80 kg/ha	Fertilizer
14/03/22	Nitrogen N39	200 L/ha	Fertilizer
05/04/22	Nitrogen N39	175 L/ha	Fertilizer
13/04/22	MgSO4	3,5 kg/ha	Fertilizer
13/04/22	TMF	1 L/ha	Fertilizer
13/04/22	Cycocel	1L /ha	Cycocel is a plant growth regulator
25/04/22	Nitrogen N39	125 L/ha	Fertilizer
22/05/22	Kestrel	1 L/ha	
22/05/22	MGSO4	3kg/ha	Fertilizer
22/05/22	TMF	1L/ha	Fertilizer
01/06/22	Tebuzip ?? Maybe Tebuconazole	0,7 L/ha	??
01/06/22	TMF	1L/ha	Fertilizer
01/06/22	Prinus	0,1 ??	
	Wheat	108 Qx/ha	
15/08/22	Horse manure	15 T/ha	Fertilizer
17/09/22	Mixture Scam Selfie		Cover crop

SM1b: Soil monolith sampling

SM1c: Fertilisation in BIOFAIR ecotron trial

The total required amount of nitrogen fertiliser was split into three doses, each of which was applied according to plant growth stage in respective climate. The recommendation was made for each soil type in each CER and based on the guidelines of the Wallon Centre of Agronomic Research (CRA-W) as described in Le Livre Blanc: La fertilisation azotée. The recommendation consists in calculating a forecast N balance for the crop, taking into account the soil nitrogen supplies and the estimated needs of the intended crop. It makes it possible to assess the adequate quantity of fertilizer to bring. Soil nitrogen supplies are determined based on soil characteristics (humus content, remaining mineral nitrogen at the end of winter, mineralization of organic nitrogen in the soil), the phytotechnical history of the plot (crop residues, previous crop) and mineral and previous organic nitrogen fertilizer inputs by the farmer.

Simulated year	CER	Soil	End of winter kg N-NO_3 ha^{-1} 0-25 + 25-50cm		Total amount of fertiliser needed kg N ha^{-1}	1. dose: start stem-elongation kg N ha^{-1} (ecotron date of application)	2. dose: 2. node kg N ha^{-1} (ecotron date of application)	3. dose: flag leaf kg N ha^{-1} (ecotron date of application)
2085	3	1	11	10	200	65 (10/02/2085)	65 (04/03/2085)	70 (22/04/2085)
2085	3	2	19	10	150	50 (10/02/2085)	50 (04/03/2085)	50 (22/04/2085)
2085	6	1	11	11	200	65 (10/02/2085)	65 (04/03/2085)	70 (22/04/2085)
2085	6	2	23	18	155	54 (10/02/2085)	50 (04/03/2085)	50 (22/04/2085)
2068	2	1	2	5	205	70 (17/04/2068)	65 (12/05/2068)	70 (23/05/2068)
2068	2	2	5	10	155	55 (17/04/2068)	50 (12/05/2068)	50 (23/05/2068)
2068	5	1	3	10	200	70 (17/04/2068)	65 (12/05/2068)	65 (23/05/2068)
2068	5	2	3	9	160	55 (17/04/2068)	50 (12/05/2068)	55 (23/05/2068)
2013	1	1	3	5	205	70 (12/05/2013)	65 (23/05/2013)	70 (12/06/2013)
2013	1	2	3	<2	155	55 (12/05/2013)	50 (23/05/2013)	50 (12/06/2013)
2013	4	1	2	4	200	70 (12/05/2013)	65 (23/05/2013)	65 (12/06/2013)
2013	4	2	2	<2	160	55 (12/05/2013)	50 (23/05/2013)	55 (12/06/2013)

SM2: Empirical and modelled parameter to quantify agronomic performance and environmental impact

Parameter	Unit and scale of inference	Measurement
BBCH plant developmental stage	Unitless, visual accessment of all plants within CERs, bi-weekly	Meier, U. (ed.). (2001). Growth Stages of Mono- and Dicotyledonous Plants. BBCH Monograph. 2nd edition. Federal Biological Research Centre for Agriculture and Forestry, Germany. DOI: 10.5073/20180906-074619.
Plant height	cm, 3 plants per cube, 8 cubes per soil type x climate, monthly	Metric
Leaf area index	Unitless, 3 plants per cube, 4 cubes per soil type x climate, monthly June-August	Scan and image analysis (pixel)
Total root length	mm, 2 plants per cube, 2 cubes per soil type x climate, BBCH 30, 50, 80	Wash, scan, image analysis Seethepalli, A., Dhakal, K., Griffiths, M., Guo, H., Freschet, G. T., York, L. M. (2021). RhizoVision Explorer: Open-source software for root image analysis and measurement standardization. AoB PLANTS, plab056, https://doi.org/10.1093/aobpla/plab056.
Maximum quantum efficiency PSII	Fv/Fm, 3 leaves per cube, 8 cubes per soil type x climate, weekly for sufficiently large leaf surface area	HandyP+, Hansatech Instruments, UK Vlaović J., Balen J., Grgić K., Žagar D., Galić V. & Šimić D., 2020. An Overview of Chlorophyll Fluorescence Measurement Process, Meters and Methods. In: 2020 International Conference on Smart Systems and Technologies (SST). Presented at the 2020 International Conference on Smart Systems and Technologies (SST), 245–250. doi: 10.1109/SST49455.2020.9264091.
Foliar silicon	mg kg-1, 3 leaves per cube, 4 cubes per soil type x climate, BBCH 30, 50, 80	Inductively coupled plasma optical emission spectroscopy (ICP-OES), Core Facility Hohenheim (CFH), University of Stuttgart, Germany
Foliar proline	µg g-1, 3 leaves per cube, 4 cubes per soil	Acid-ninhydrin quantification. Bates, L.S., Waldren, R.P. & Teare, I.D. Rapid determination of free proline for

	type x climate, BBCH 30, 50, 80	water-stress studies. Plant Soil 39, 205–207 (1973). https://doi.org/10.1007/BF00018060.
Take-all index	%, visual accessment, 3 plants per cube, 4 cubes per soil type x climate, BBCH 40-50	Cook, RJ (2003) Take-all of wheat. Physiological and Molecular Plant Pathology 62(2):73-86. https://doi.org/10.1016/S0885-5765(03)00042-0. BBA (1999). F. 01 Deutscher Vorschlag für eine EPPO-Richtlinie zur Prüfung der Wirksamkeit von Saatgutbehandlungsmitteln gegen luft- und bodenbürtige Krankheitserreger an Getreide. Biologischen Bundesanstalt für Land- und Forstwirtschaft, Braunschweig, Germany
NO_3 aq.	mg L-1, interstitial soil pore water extracts using rhizons at 8-12cm depth, 0-8 cubes per soil type x climate if water extraction possible, weekly	LAQUAtwin NO3-11C, Horiba Ltd., Kyoto, Japan Folegatti, M.V., Blanco, F.F., Boaretto, R.M. and Boaretto, A.E. (2005) Calibration of cardy-ion meters to measure nutrient concentrations in soil solution and in plant sap. Sci. Agric. 62(1): 8-11. https://doi.org/10.1590/S0103-90162005000100002
Glucose equivalent	mg ml-1, interstitial soil pore water extracts using rhizons at 8-12cm depth, 0-8 cubes per soil type x climate if water extraction possible, weekly	Yemm EW, Williams AJ (1954). The estimation of carbohydrates in plant extracts by anthrone. Biochem J. 57(3):508-14. https://doi.org/0.1042/bj0570508.
Microbial biomass carbon	CO_2-C mg g-1, BBCH 30, 50, 80	MicroResp, James Hutton Institute, UK Campbell, CD, Chapman, SJ, Cameron, CM, Davidson, MS, Potts JM (2003) A rapid microtiter plate method to measure carbon dioxide evolved from carbon substrate amendments so as to determine the physiological profiles of soil microbial communities by using whole soil. Appl. Environ. Microbiol., 69 (6): 3593-3599.
Grain yield	t ha-1 at 14% grain moisture, harvest, 4 undisturbed cubes per soil type x climate,	Weight and water content

Soil N leaching	kg N ha-1 yr-1, cummulative	DNDC (DeNitrification-DeComposition) model, Institute for the Study of Earth, Oceans, and Space, University of New Hampshire Durham, USA https://www.dndc.sr.unh.edu/
Soil CO_2-emission	CO_2-C kg ha-1 yr-1, cummulative	DNDC (DeNitrification-DeComposition) model, Institute for the Study of Earth, Oceans, and Space, University of New Hampshire Durham, USA https://www.dndc.sr.unh.edu/
Soil N_2O-emission	N_2O-N kg ha-1 yr-1, cummulative	DNDC (DeNitrification-DeComposition) model, Institute for the Study of Earth, Oceans, and Space, University of New Hampshire Durham, USA https://www.dndc.sr.unh.edu/

SM3: DNDC model evaluation against measured data

To evaluate the accuracy of the modelled parameters from the DNDC model, the results were compared against measured values using the coefficient of determination (R^2). The correlation was excellent for the DNDC estimate of CO_2-emission and the basal soil microbial respiration accessed with the MicroResp kit (R^2=0.98). While no empirical data directly on denitrification was gathered during the Ecotron experiment, the DNDC estimate of soil N_2O-emission correlated well with the metabolic quotient (qCO_2), probably reflecting that microbes that actively denitrify (under anaerobic conditions) also engage in respiration, leading to CO_2 emissions (R^2=0.6). The correlation between the measured nitrate in aqueous solution and the predicted soil N-leaching was good (R^2=0.71) if 2013.S1 was not taken into account. Similarly, a moderate correlation (R^2=0.53) was observed for the root exudation predicted by the DNDC model and the measured values of glucose equivalent, if 2013.S1 was excluded (R^2=0.53). The data distribution suggests that the NO_3 in aqueous solution in 2013.S1 was actually higher than what was measured, which could be related to the fast turnover time of NO_3 in soil where it is quickly taken up by plants and microbes. For root exudation, in DNDC root exudates are represented as a portion of the root-derived carbon where the model considers both root growth and root turnover, while the empirical measurement performed here only gives an estimate of low-molecular weight sugars measured as glucose equivalents which may explain discrepancies. The data distribution suggests that both the DNDC model and the measurements underestimated root exudation in 2013.S1, which similarly to NO_3 could be due to the high reactivity of these molecules in the soil. Overall, the modelled and measured parameter were well aligned for gaseous emissions of CO_2 and N_2O (R^2=0.98, R^2=0.6), and moderately well aligned for nitrate and glucose equivalent (R^2=0.71, R^2=0.53) with 2013.S1 not fitting the distribution, possibly due to the technical limitations of the empirical measurements.

SM4: Loadings for the main vector of each modality (Year.Soil)

	2013.S1	2013.S2	2068.S1	2068.S2	2085.S1	2085.S2
Basal.CO2.02.05				-0.8399		1.55331
Basal.CO2.30.05	-0.7269	-0.8814		1.23257		
Basal.CO2.03.07				-0.8972	1.25774	
Basal.CO2.07.08			-0.0394			
Foliar.proline_BBCH80			0.14672			
Foliar.silicon_BBCH50						1.36197
Glucose.equivalent.27.02	0.82706		-0.1687			
Glucose.equivalent.14.03	-0.6393					
Glucose.equivalent.02.05	0.67934			-0.7745		
Glucose.equivalent.21.05		-0.896				
Glucose.equivalent.03.07			-0.0938			
Glucose.equivalent.19.07	-0.7882	-1.0001				
Glucose.equivalent.25.08			-0.1407			
Leaf.area.index.March	-0.864	-0.8922				1.31855
Leaf.area.index.April		-1.0344	0.09927			1.53179
Leaf.area.index.May		-1.141				
Leaf.area.index.June		-1.1419				
Maximum.quantum.efficiency.PSII.23.03				-0.7742	1.17837	1.12584
Maximum.quantum.efficiency.PSII.07.04				0.82972		-1.2996
Maximum.quantum.efficiency.PSII.19.04			-0.1273			
Maximum.quantum.efficiency.PSII.10.05				-0.9076		
Maximum.quantum.efficiency.PSII.08.06		-0.9777				
Maximum.quantum.efficiency.PSII.21.06	-0.8253	-1.2397				
Maximum.quantum.efficiency.PSII.05.07				0.89958		
Microbial.biomass.carbon_BBCH30			-0.1362			
Microbial.biomass.carbon_BBCH50			0.03226	0.72059		
NO3.aq.14.03					-1.3856	
NO3.aq.15.05	0.75471			-0.8281		
NO3.aq.12.06					1.53893	
NO3.aq.03.07		-0.9794				
NO3.aq.19.07	-0.7774					
NO3.aq.07.08					1.16468	
Plant.height.March					1.4301	1.14948
Plant.height.April					1.43003	1.14928
Plant.height.May					1.43006	1.14936
Plant.height.June					1.43012	1.14976
Take.all.index	-0.722					
Total.root.length_BBCH30			-0.0929		1.13926	