Short title: Byers & Jacobs – Quantitative insect GC-EAD for plant interaction analysis

Title: Quantitative analysis of gas chromatography-coupled electroantennographic detection (GC-EAD)
of plant volatiles by insects

Kelsey J.R.P. Byers[1] & Robert N. Jacobs[2]

[1] Department of Cell and Developmental Biology, John Innes Centre, Norwich Research Park, Norwich,
NR4 7UH, United Kingdom. To whom correspondence may be addressed: Kelsey.Byers@jic.ac.uk, ORCID:
0000-0002-0923-8264

[2] 6 Greenways, Norwich, NR4 6HE, United Kingdom. Correspondence: robert.n.jacobs@gmail.com

Abstract

Premise: Plant-interacting insects receive plant volatile signals through their antennae, and voltage
changes across an antenna exposed to volatile stimuli can be measured to determine if insects can
perceive them, often by coupling gas chromatography to electroantennographic detection (GC-EAD).
Current methods for analysing GC-EAD data rely on *qualitative* observation, rather than using
quantitative analysis, and are thus prone to bias.

Methods and Results: We developed a novel *quantitative* methodology for analysis of GC-EAD data
using a signal processing technique on EAD data to compare responses of hawkmoths and bumble bees
to a library of common floral volatiles. Responses varied between species and sexes and these responses
were in some cases affected by compound type or modification.

Conclusions: Our novel GC-EAD analysis technique is robust to baseline drift and low signal:noise ratios
commonly found in GC-EAD data, and provides a way forward for quantitative studies of plant volatile-
mediated plant-insect interactions.

INTRODUCTION

Plants produce a wide array of secondary metabolites – i.e. those that are not immediately tied to growth and reproduction – with over 200,000 isolated and described across the plant kingdom (Kessler and Kalske, 2018). Many of these compounds play direct roles in the interactions of plants with the abiotic and biotic environment, and have been exploited for this purpose by humanity, including the agricultural (e.g. geraniol, a terpenoid volatile, which repels eggplant-infesting pest moths (Ghosh et al., 2023; Byers, 2023)) and flavour and fragrance industries (e.g. vanillin, a volatile product of the phenylpropanoid pathway, which is the dominant compound in natural vanilla extract (Gallage et al., 2014)). Although these compounds can be exploited in such ways, and although chemical structures of numerous plant metabolites have been described, their biological functions remain unknown in many cases (Dixon and Dickinson, 2024). Many of these compounds are volatiles, which are especially well known for influencing plants' interactions with insects by attracting both beneficial and antagonistic species (e.g. pollinators and herbivores (Raguso, 2008)), with significant implications for plant ecology and evolution (Byers, 2021) as well as agriculture (Pickett and Khan, 2016).

Identifying which volatile cues emitted by plants are perceived by plant-interacting insects is key to our understanding of these cross-kingdom interactions. In particular, identifying insect-relevant volatiles can facilitate work in plant secondary metabolism, ecology, and evolution. For example, identifying pest-resistant crop varieties by screening for known resistance volatiles is cheaper and easier than performing field trials (e.g. Ghosh et al., 2023), while determining which volatiles are key to the ecology and evolution of plant-pollinator interactions can facilitate genetic mapping of only those volatiles rather than the entire plant headspace (e.g. Byers et al., 2014a, 2014b). In short, identifying insect-relevant volatiles – often only a fraction of a plant's emitted volatiles and individually not always the most abundant volatiles (Clavijo McCormick et al., 2014) – can dramatically reduce the experimental scope and increase the tractability of plant secondary metabolite study.

Electroantennography – the detection and measurement of electrical signals from whole insect

antennae – is often used to screen plant volatiles for insect perceptibility. The compounds to which the

insect antenna "responds" with some voltage change are taken to be physiologically active and may be

further tested via behavioural assays with single pure or mixed compounds or with plant lines known to

be high or low in the relevant compounds. Although these techniques have been in use for nearly

seventy years, being first described in 1957 (Schneider, 1957), the resulting data are always presented

qualitatively – "response/no response" (e.g. Peakall et al., 2010) – with no quantitative, repeatable

measure used to define antennal "response" versus background activity. Indeed, a comprehensive and

well-cited overview of GC-EAD includes no mention of how data analysis is performed or what criteria

are used to define an "active" peak (Schiestl and Marion-Poll, 2002). This has hampered the field by

limiting reproducibility and cross-study or cross-observer comparisons, introducing error, and reducing

ease of application to broader plant volatile studies. This may be a particular issue with many plant

volatiles, which – unlike insect pheromones – may not have the same degree of "dedicated hardware"

on the insect antenna (Akers and Getz, 1993; Schiestl and Marion-Poll, 2002) and thus may produce

weaker signals, with the exception of sexually deceptive insect pollination systems which use insect

pheromone compounds to attract pollinators.

In this Protocol Note, I present "best practice" statistical approaches incorporating techniques from

signal processing to analyse gas chromatography-coupled electroantennographic detection (GC-EAD)

data in a quantitative fashion. Methods for correcting for baseline drift (specifically, the use of the

antennal signal first forward difference), background antennal noise (specifically, comparison with

response to the solvent control), and antennal sensitivity loss (via use of windowing) are presented. A

sample data set from six *Manduca sexta* hawkmoths (three males and three females), four *Bombus*

terrestris audax bumble bees (all workers), and 28 common volatile stimuli of different volatile chemical

skeleton types (all present in a variety of plant species) are used to demonstrate the methods and their

application in detail and show that they work for a variety of volatile plant secondary metabolites and

two major model pollinating insect species. I show that quantitative analyses of electroantennogram

data are indeed possible and quite feasible, pointing the way forward for robust, statistically-based

analyses of antennal detection ability in volatile-driven plant-insect interactions.

METHODS AND RESULTS

Methods: Gas chromatography-coupled electroantennographic detection (GC-EAD) setup and

recording

Insect sourcing

Adult *Manduca sexta* (L.) (Lepidoptera: Sphingidae) hawkmoth males and females were sourced from a

maintained colony reared on artificial diet (Frontiers Scientific Services, Newark, Delaware, USA) at the

John Innes Centre Entomology platform facility; adults were not fed but females were allowed to

oviposit freely on live *Nicotiana benthamiana* plants and males and females were kept together prior to

collection. Worker *Bombus terrestris audax* (Harris) (Hymenoptera: Apidae) were sourced from a single

Bumblebee New Season Hive (Biobest, purchased from Agralan Ltd, Wiltshire, UK) which were allowed

to freely forage in a custom-built acrylic cage with 25% sucrose and ground pollen *ad libitum* (Sevenhills

Wholefoods Organic Raw Bee Pollen, Dinnington, UK).

EAD test mixture preparation and analysis

To assess antennal responses to multiple compounds delivered at approximately equal flame ionization

detector (FID) sensitivities, a mixture of 28 authentic reference standards was assembled, representing

29 GC-FID peaks (farnesol is sold as a mixture of isomers and both elute separately on the GC-FID).

Standards were obtained from a variety of established sources (see Table S1 for vendor addresses and

Table S2 for vendor and product numbers) and diluted to 10μg/μl in hexane (HPLC-grade, ca. 95%

purity). The final compound list was selected to include a variety of chemical skeletons (terpenoids,

aromatics, fatty acid derived compounds, and one nitrogenous compound) that would separate well on

a 10°C per minute GC oven ramp program. Due to differences in FID responsiveness to the individual

compounds (i.e. some compounds may produce different peak areas on the FID despite being at equal

injected amounts), different volumes of the 10μg/μl dilutions were added to the mixture (see Table S2

for final concentrations, which ranged from 93ng/μl for (R)-(+)-limonene to 2004ng/μl (2.004μg/μl) for

indole).

To determine retention times and mixture peak identities, 3ul of the test mixture was manually injected

into an Agilent 7890B GC (Agilent, Stockport, UK) fitted with an Agilent 5977A/B MS detector (GC-MS)

and a Phenomenex Zebron ZB-5HT column (30m x 0.25mm x 0.10μm; Phenomenex UK Ltd, Macclesfield,

United Kingdom). The GC run parameters were as follows: the column flow was 1.0603 ml/min of

helium carrier gas with an oven program of 50°C for 4 minutes followed by a temperature ramp of

10°C/minute to 230°C, which was then held for 4 minutes. The inlet was splitless and held at 250°C and

the MS source was held at 230°C. Compounds were identified using Agilent Unknowns software (v10.1,

Agilent, Stockport, UK) and via comparison to Kovats Retention Indices of the individual compounds run

separately using the same GC-MS system, calculated using an alkane ladder (49452-U, Sigma-Aldrich,

Watford, UK).

EAD antennal preparation and assembly

Manduca sexta male and female adults: single antennae were excised using microdissection scissors

from three individuals each of living adult male and female *Manduca sexta*. Due to biosecurity

requirements for this non-native potential plant pest species, excised antennae were placed in a 15ml

conical tube on wet tissue paper to maintain antennal humidity levels and carried to the

electrophysiology setup in a separate building. The fine tip of an antenna (representing ca. 7 segments,

cut where the tip abruptly broadens) was removed using a microscalpel under a dissection microscope

and the antenna mounted on an antenna electrode fork (Syntech PRG-2) using electrode gel (Signa Gel,

Parker Laboratories, Fairfield, New Jersey, USA), which was then plugged in to the electrode mount

(Syntech MP-15 with EAG Combi Probe) and moved into position in front of the GC eluent-carrying air

stream approximately 1cm from the air stream outlet.

Bombus terrestris audax workers: individuals were removed from the Entomology platform facility in

50ml conical tubes, then kept at 4°C for 30 minutes to enable safe antennal removal. Antennae were

then removed using microdissection scissors under red light (with the insect held using a set of forceps)

and the tip of the final flagellar segment removed using a microscalpel under a dissection microscope,

then treated as above.

GC-EAD setup and recording procedure

The GC-FID setup consisted of an Agilent 8890 GC fitted with a flame ionization detector (FID). The

column (a Phenomenex Zebron ZB-5HT column (35m x 0.25mm x 0.10μm), identical to the one used for

GC-MS analysis) was fitted to a splitless inlet kept at 250°C. The detector end of the column was fitted to

the bottom end of an inert glass Y-splitter (Agilent UltraInert Universal PressFit Y-Splitter , part 5190-

6980). The two other arms of the splitter were fitted with deactivated column segments of equal 30cm

length (Agilent CP803210, 0.32mm inner diameter), with one end running to the FID (which was kept at

300°C) and the other end leaving the GC oven through a port on the side which was fitted with a

Syntech EC-03 heated transfer tube kept at 250°C. The GC used hydrogen as a carrier gas, which was

generated with a Parker Hydrogen Generator (H2PEM-100, Thames Restek UK Ltd, Saunderton, United

Kingdom). Nitrogen gas at a flow of 25ml/min was used as a make-up gas, and the FID was also fed with

compressed air at 400ml/min to ensure adequate flame gas supply. The GC run parameters were as

follows: the column flow was 4.7 ml/min with an oven program of 50°C for 4 minutes followed by a

temperature ramp of 10°C/minute to 230°C, which was then held for 4 minutes. Each manual sample

injection consisted of 3ul of the EAD test mixture, for a total of 0.279-6.012µg of compound injected.

GC-FID chromatogram waveforms were exported from Agilent OpenLab CDS v2.5 software as CSV files

consisting of an FID amplitude value (in pA) for each given sampled timepoint.

The EAD recording setup consisted of a Syntech IDAC-2 (Ockenfels Syntech, Buchenbach, Germany)

connected to the desktop computer controlling the GC-FID system. A constant air flow (setting 8 on the

CS-55) was delivered using a Syntech CS-55 Stimulus Controller throughout the experiment. Prior to and

shortly after each GC-FID run, a puff of the EAD test mixture (via a Pasteur pipette fitted with filter

paper, with 10µl of the EAD test mixture added onto the paper, also setting 8 on the CS-55) was

delivered using the CS-55 to ensure antennal viability had been maintained throughout the GC-FID run.

EAD signals were captured using Syntech GcEAD 2014 v1.2.5 software with default settings and

exported as CSV waveforms in GcEAD 2014 (measured as millivolts) prior to analysis.

Methods: Data processing and analysis

GC-FID peak identification

A custom Perl script (Perl, 2023; see Appendix for script) was developed to determine the "onset" and

"offset" timings (i.e. when a peak began and when it ended) of GC-FID peaks above a certain user-

defined threshold from the FID CSV data. In the presented experimental data, the threshold was set at

50pA to capture all significant peaks while avoiding inclusion of small peaks likely representing

contamination from the hexane solvent. Note that this resulted in the merger of the closely eluting

caryophyllene oxide and ethyl laurate peaks, which were removed from the analyses of compound type

and compound modification impact on GC-EAD responses as a result.

Data loading and initial processing

EAD and FID waveforms were loaded as CSV files into R (R Core Team, 2024; see Appendix for example

script). GC-FID onsets and offsets determined in the previous section were also loaded. As the EAD setup

samples at a rate of 6000 datapoints per second and does not export a timestamp to its CSV file, a new

time variable was defined corresponding to the sampling rate. The GC-FID injection time and a 20

minute endpoint (after all tested peaks had eluted) relative to the EAD time were also defined, as were

EAD timestamps corresponding to each GC-FID peak onset and offset. As injections were done manually

for the GC-FID, there was occasionally a slight disparity between GC-FID peak onsets and the onset of a

significant-seeming EAD pattern, and so a user-defined offset was determined by comparing the most

significant EAD pattern onset with the previously determined corresponding GC-FID peak onset.

First Forward Difference (FFD) processing and significant FFD value identification

Using the diff() function in base R (specifically as 'diff(EADamplitude,lag=1,differences=1)'), the First

Forward Difference of the antennal amplitude was calculated from the input antennal amplitude data.

The FFD vector was subsetted into five windows of four minutes each for a total GC-FID run of 20

minutes. For each window, the average and standard deviation of the FFD within that window was

calculated. A significance threshold of 3σ above or below the average was calculated, and data points

that exceeded the significance threshold for their respective windows were marked as significant.

Calculating the number of significant FFD hits per second for each identified GC-FID peak

For each GC-FID peak identified, the number of significant hits as defined above within its onset-offset

window was determined using the 'between' function in the R package dplyr (Wickham et al., 2023).

This value was divided by the length of the GC-FID peak in seconds (offset time minus onset time) to get

the significant hit frequency of hits/second. As EAD data can have significant background noise (Schiestl

and Marion-Poll, 2002), we defined the frequency of the initial hexane peak 1 as the threshold for

whether a peak attracted significant EAD activity: if the frequency for a given GC-FID peak exceeded the

frequency of the hexane peak, it was counted as significant, and otherwise considered insignificant.

Statistical analysis

A "significant response" across each species-sex/caste category (female *M. sexta*, male *M. sexta*, and

worker *B. terrestris*) was identified for compounds where >50% of individuals tested showed a

significant hit frequency greater than that of the initial large solvent (hexane) peak.

To investigate whether the results were repeatable between moths, we calculated the Pearson

correlation coefficient between the significant hit frequency for each pair of moths within a sex using

cor.test() in R. We repeated this test with a binary vector (1 = GC-FID peak had a significant hit

frequency; 0 = it did not). We then used lm() in R to calculate the slope and intercept of the relationship

between each pair of moths within a sex. Finally, we used lm() and emmeans() (Lenth, 2024) to

determine whether compound skeletal or modification type had an influence on significant hit

frequency within each species-sex/caste group (female *M. sexta*, male *M. sexta*, and worker *B.*

terrestris).

Results

The new method allows quantitative analysis and avoids baseline drift

Traditional GC-EAD analysis is done in a visual qualitative fashion by comparing GC-FID peak timings with

EAD readouts by eye (e.g. comparison of **Figure 1A** with **Figure 1B**, see also **Figures S1-S15**). It also

suffers from baseline drift and signal noise, two phenomena that make it difficult to detect whether an

EAD data point is indeed a "significant" response to a GC-provided stimulus or not (Schiestl and Marion-

Poll, 2002). As the antenna degrades over time and accommodates in the long term to compound

stimuli (see Byers et al., 2020), antennal responsiveness and the ability to detect significant responses by

eye also degrades and the noise level often increases, also impeding qualitative visual comparison of GC-

FID and EAD data.

Here we present a method that avoids the otherwise significant biases inherent in GC-EAD *qualitative*

analysis methodology by the use of a signal processing technique, the *first forward difference* (FFD), to

remove the influence of baseline drift, essentially flattening out the EAD signal and replacing its

waveform with a series of individual spikes (**Figure 1C**, see also **Figures S16-S30**). A significance

threshold may then be set for each window (horizontal lines in **Figure 1C** and **Figures S16-S30**, both

showing ± 3 standard deviations above/below the mean for that window), and the frequency of spikes

with an amplitude over that significance threshold during a given GC-FID peak may be easily calculated

and analysed using traditional statistical methods. This results in a *quantitative* rather than *qualitative*

assessment of whether a GC-EAD peak is physiologically active or not.

GC-FID peaks with significant EAD responses

Table 1 shows the list of compounds tested in the EAD test mixture along with their response rates for

female and male moths and worker bumble bees and how many individuals of each had a response rate

over that of the main hexane peak. In general, female moths were most responsive to fatty acid-derived

compounds (FADs, red in **Figure 2A**, see also **Figure S31**) and to some but not all aromatic volatiles (blue

in **Figure 2A**). Responses to terpenoids were more mixed (green in **Figure 2A**), with some compounds

(e.g. (1S)-(-)-verbenone) eliciting significant responses, while others (e.g. camphene) eliciting no

significant responses at all. The sole tested nitrogenous compound, indole (yellow in **Figure 2A**), did not

elicit significant responses in female moths. Overall, female moths responded significantly to 15 of the

31 tested compounds (excluding the hexane peaks). Male moths were far less responsive than females

to the majority of compounds, with overall significant responses to only four of the 31 tested

compounds (excluding the hexane peaks) and with no significant difference in response between

compound skeleton types (**Figure S31A,B**). Worker bumble bees responded significantly to 10 of the 31

compounds. Bees responded equally to all types of compound classes (**Figure S31C**), although there was

no response to indole, the sole nitrogenous compound tested, in any bee.

Compounds containing oxygenated modifications seemed to produce the highest level of response in

female moths, but not in males (**Figure S31D,E**). Females responded significantly to 6 of 10 presented

esters (analyzed without the co-eluting peaks ethyl laurate and caryophyllene oxide), 2/6 alcohols, 3/3

aldehydes, and 3/3 ketones, but not to any of the five presented unmodified monoterpenes ((1R)-α-

pinene, camphene, and R-(+)-limonene) or sesquiterpenes (caryophyllene, α-humulene), both groups of

which lack oxygen atoms. Male moths responded to 1 of 10 esters, 0/6 alcohols, 1/3 aldehydes, and 0/3

ketones but, similarly to females, only responded to 1/5 of the unmodified (mono/sesqui)terpenes. Both

females and males also responded significantly overall to the "Unknown 2" compound, whose identity

remains unclear despite GC-MS analysis but which likely also is oxygenated based on tentative NIST

library matches. For bees, there was no difference in response based on compound modification,

oxygenated or otherwise (**Figure S31F**). Bees responded to 3 of 10 esters, 3/6 alcohols, 1/3 aldehydes,

2/3 ketones, and 1/5 unmodified (mono/sesqui)terpenes.

Repeatability within moth sexes and comparisons between sexes

Responses to non-hexane compounds were highly repeatable between female moths (**Figure S32A**,

Pearson's correlation coefficient range 0.265 to 0.637, all $P < 3 \times 10^{-7}$), but not between male moths

(**Figure S32B**, Pearson's correlation coefficient range -0.238 to -0.054, all $P \geq 0.20$). In general, male

moths were substantially less responsive to non-hexane chemical stimuli in this experiment, as

represented by their much lower average significant hit frequency (**Table 1,** range 0-4.000 significant

hits/second) versus females (range 0-21.919 significant hits/second). Responses between males and

females were generally uncorrelated (Pearson's r = 0.073, *P* = 0.696, **Figure S33**). The only two

compounds that resulted in significant hit frequencies in ≥ 2/3 of moths of both sexes were

phenylacetaldehyde and "Unknown 1".

Repeatability and feasibility across species and comparison to existing literature

To ensure that our results were not simply due to some property of *Manduca sexta* antennae, we also

analysed the same stimulus mixture with four workers of the polylectic (wide dietary breadth, Rasmont

et al., 2008) bumble bee *Bombus terrestris audax* using the same data collection and analysis

methodologies. We found no compounds that were significant in >50% of tested individuals across all

three species-sex/caste combinations (female hawkmoths, male hawkmoths, worker bumble bees).

However, we did identify a short list of compounds that were significant in >50% of tested individuals for

both female *Manduca sexta* and worker *Bombus terrestris audax*, namely benzaldehyde, (*Z*)-3-hexenyl

acetate, acetophenone, (1S)-(-)-verbenone, and geraniol (**Table 1**). Results from individuals of all three

groups are presented in **Figure 3**. We also observed overall weaker responses in bumble bees compared

with hawkmoths, despite robust responses to single puffs of the EAD stimulus mixture. The signal to

noise threshold in the bumble bee antennae appears to be weaker than in the hawkmoths (as shown by

much lower first forward difference significance thresholds and a lower range of significant average hit

frequencies (range 0-1.752 hits/second compared with hawkmoth values of 0-21.919 and 0-4.000 for

females and males respectively). We separately conducted a somewhat haphazard literature review to

identify whether our tested compounds had previously been identified as electroantennographically

active in either *Manduca sexta* or *Bombus terrestris*, with the results shown in **Table S3**. In general,

many of our tested compounds had not previously been tested with GC-EAD or puff-based

electroantennography, and those that had been often produced results that differed between studies,

further suggesting the need for a quantitative analysis technique such as the one presented here.

CONCLUSIONS

In this Protocol Note, we have presented a novel signal processing-based methodology for use with gas

chromatography-coupled electroantennographic detection (GC-EAD) measurements of insect antennae

in response to plant volatiles. Female *Manduca sexta* hawkmoths were responsive to approximately half

of tested compounds in a highly repeatable fashion, while male hawkmoths of the same species

responded to only about one eighth of tested compounds and were far less repeatable between

individuals. In female moths, compound skeleton type (FAD, aromatic, terpenoid, or nitrogenous) and

modification (none, alcohol, aldehyde, ester, ketone, or oxide) played a significant role in response

patterns, while in male moths this was only true for modification type but not compound skeleton type.

Worker *Bombus terrestris audax* bumble bees showed an intermediate level of response, with

significant responses to one third of tested compounds, and with no difference in response rate relative

to compound type or modification.

Our haphazard literature review to identify compounds that had previously been identified as

electroantennographically active in either *Manduca sexta* or *Bombus terrestris* identified large gaps in

our knowledge, as well as discrepancies between studies in whether compounds were designated as

physiologically active or not in the antenna. Consistent with our findings, some compounds were

significant in some studies but not others, emphasizing the importance of testing stimuli on multiple

individuals of a given species rather than relying on one or two individuals, given that we found

variability even between individuals of the same commercial bumble bee hive and hawkmoth colony.

The weaker responses seen in bumble bees further necessitate the use of this new technique to ensure

unbiased assessment of electroantennographic detection studies in potentially less strongly responsive

insects.

By studying responses to common plant volatiles in two model pollinator species, we have shown that

responses are species- and sex-specific, and that it will be difficult to generalize results from one

pollinator group to another. Further study will be needed to determine whether results can be

generalized within a genus of pollinators, or between those that are generalists and specialists.

However, at least one species of pollinating insect was responsive to the majority of the plant

compounds tested here. Interestingly, there was no significant GC-EAD response in any species tested to

the monoterpene (R)-(+)-limonene. Limonene is the most widely distributed floral volatile in terms of

angiosperm families in which it is found (Schiestl, 2010), and is known to provoke a responsive in the

antennal lobe of *Bombus vosnesenskii* (Byers et al., 2014a). We also did not see any response to the

common nitrogenous volatile indole. The latter is utilized in many dung-mimicking brood site-deceptive

pollination systems (Jürgens et al., 2013), which does not include the insects tested here.

The methods introduced make use of existing typical GC-EAD recording setups and methods, differing

only in the downstream processing of collected GC-EAD data, and thus may be used with both existing

as well as novel datasets. In addition, they are simple to execute, requiring only minimal implementation

in R or another statistical processing language (example code is provided in the Appendix). This study

provides the first truly quantitative analysis of insect responses to common plant volatiles using GC-EAD,

and paves the way forward for future GC-EAD analyses to incorporate quantitative, unbiased data

processing methodologies to determine whether plant volatiles are physiologically active to interacting

insect species.

AUTHOR CONTRIBUTIONS

KJRPB conceived the project, designed and executed the data collection and data analysis methodology,

and wrote the manuscript. RNJ contributed significant signal processing methodology ideas and

expertise to the data analysis methodology development. Both authors approved the final version of the

manuscript.

ACKNOWLEDGMENTS

The authors thank the John Innes Centre (JIC) Entomology platform for maintaining the *Manduca sexta*

and *Bombus terrestris* colonies and the JIC Metabolomics platform for GC-MS maintenance. Funding was

provided by the UK Biotechnology and Biological Sciences Research Council Institute Strategic

Programmes (Harnessing Biosynthesis for Sustainable Food and Health (HBio) (grant no. BB/X01097X/1)

and Building Robustness in Crops (BRiC) (grant no. BB/X01102X/1)). We also thank anonymous

reviewers and the editor for their helpful suggestions.

DATA AVAILABILITY

341   All raw data (GC-FID CSV files, electroantennography CSV files, and relevant GC-FID injection

342   timestamps) are available on FigShare (doi: XXX). Scripts referenced in the main text and appendices are

343   also available on GitHub ().

REFERENCES

Akers, R. P., and W. M. Getz. 1993. Response of olfactory receptor neurons in honeybees to odorants
and their binary mixtures. *Journal of Comparative Physiology A* 173: 169-185.

Byers, K. J. R. P. 2021. "As if they discovered it by the scent": improving our understanding of the
chemical ecology, evolution, and genetics of floral scent and its role in pollination. *American Journal of*
Botany 108: 729-731.

Byers, K. J. R. P. 2023. Reducing eggs on eggplant: a common naturally emitted plant volatile could
replace insecticides in the 'king of vegetables'. *New Phytologist* 240: 915-917.

Byers, K. J. R. P., H. D. Bradshaw, Jr., and J. A. Riffell. 2014a. Three floral volatiles contribute to
differential pollinator attraction in monkeyflowers (*Mimulus*). *Journal of Experimental Biology* 217: 614-
623.

Byers, K. J. R. P., J. P. Vela, F. Peng, J. A. Riffell, and H. D. Bradshaw, Jr. 2014b. Floral volatile alleles can
contribute to pollinator-mediated reproductive isolation in monkeyflowers (*Mimulus*). *The Plant Journal*
80: 1031-1042.

Byers, K. J. R. P., K. Darragh, J. Musgrove, D. Abondano Almeida, S. Fernanda Garza, I. A. Warren, P. M.
Rastas, M. Kučka, Y. F. Chan, R. M. Merrill, S. Schulz, W. O. McMillan, and C. D. Jiggins. 2020. A major
locus controls a biologically active pheromone component in *Heliconius melpomene*. *Evolution* 74: 349-
364.

Clavijo McCormick, A., J. Gershenzon, and S. B. Unsicker. 2014. Little peaks with big effects: establishing
the role of minor plant volatiles in plant-insect interactions. *Plant, Cell & Environment* 37: 1836-1844.

Dixon, R. A., and A. J. Dickinson. 2024. A century of studying plant secondary metabolism - From "what"
to "where, how, and why?" *Plant Physiology* 195: 48-66.

Gallage, N. J., E. H. Hansen, R. Kannangara, C. E. Olsen, M. S. Motawia, K. Jørgensen, I. Holme, K.
Hebelstrup, M. Grisoni, and B. L. Møller. 2014. Vanillin formation from ferulic acid in *Vanilla planifolia* is
catalysed by a single enzyme. *Nature Communications* 5: 4037.

Ghosh, R., D. Metze, S. Sant, M. Shaikh, A. Deshpande, D. M. Firake, and S. Pandit. 2023. Chemical
ecology of Himalayan eggplant variety's antixenosis: identification of geraniol as an oviposition
deterrent against the eggplant shoot and fruit borer. *New Phytologist* 240: 1259-1274.

Jürgens, A., S.-L. Wee, A. Shuttleworth, and S. D. Johnson. 2013. Chemical mimicry of insect oviposition
sites: a global analysis of convergence in angiosperms. *Ecology Letters* 16: 1157-1167.

Kessler, A., and A. Kalske. 2018. Plant secondary metabolite diversity and species interactions. *Annual*
Review of Ecology, Evolution, and Systematics 49: 115-138.

Lenth, R. 2024. Version 1.10.0 (2024-04-06). emmeans: Estimated Marginal Means, aka Least-Squares
Means. Website https://CRAN.R-project.org/package=emmeans [accessed 1 December 2024].

Peakall, R., D. Ebert, J. Poldy, R. A. Barrow, W. Francke, C. C. Bower, and F. P. Schiestl. 2010. Pollinator
specificity, floral odour chemistry and the phylogeny of Australian sexually deceptive *Chiloglottis*
orchids: implications for pollinator-driven speciation. *New Phytologist* 188: 437-450.

Perl. 2023. Version 5.38.2 (2023-11-29). Perl. Website https://www.perl.org [accessed 1 December
2024].

Pickett, J. A., and Z. R. Khan. 2016. Plant volatile-mediated signalling and its application in agriculture:
successes and challenges. *New Phytologist* 212: 856-870.

R Foundation for Statistical Computing. 2024. Version 4.3.2 (2023-10-31). The R Foundation for
Statistical Computing, Vienna Austria. Website http://www.r-project.org/ [accessed 1 December 2024].

Raguso, R. A. 2008. Wake up and smell the roses: the ecology and evolution of floral scent. *Annual*
Review of Ecology, Evolution, and Systematics 39: 549-569.

Rasmont P., A. Coppee, D. Michez, and T. De Meulemeester. 2008. An overview of the *Bombus terrestris*
(L. 1758) subspecies (Hymenoptera: Apidae). *Annales de la Société entomologique de France (N.S.)* 44:
243-250.

Schiestl, F. P. 2010. The evolution of floral scent and insect chemical communication. *Ecology Letters* 13:
643-656.

Schiestl, F. P., and F. Marion-Poll. 2002. Detection of physiologically active flower volatiles using gas
chromatography coupled with electroantennography. In J. F. Jackson, H. F. Linskens [eds.], Analysis of
Taste and Aroma, Molecular Methods of Plant Analysis, volume 21, 173-198. Springer, Berlin,
Heidelberg, Germany.

Schneider, D. 1957. Electrophysiological investigation on the antennal receptors of the silk moth during
chemical and mechanical stimulation. *Experientia* 13: 89-91.

Wickham, H., R. François, L. Henry, K. Müller, and D. Vaughan. 2023. Version 1.1.4 (2023-11-17). dplyr: A
Grammar of Data Manipulation. Website https://CRAN.R-project.org/package=dplyr [accessed 1
December 2024].

APPENDIX

The methods used in this Protocol Note may be found in detail in the Methods section, but included
below are two computational code examples that users may find helpful:

1. The Perl script used to identify peak onsets and offsets in GC-FID CSV data;

2. Example R code to process GC-EAD data and identify GC-FID peaks with significant hit rates over the
hexane peak background level.

FIGURES

Figure 1: Comparison of raw EAD and first forward difference-processed EAD traces from a single female
Manduca sexta hawkmoth antenna between 4-8 minutes of GC elapsed time. (A) GC-FID trace annotated
with automatically detected peak starts and ends (colored vertical lines) and peak identities. (B) Raw
EAD amplitude annotated with GC-FID peak starts and ends. (C) First forward difference-processed EAD
amplitude annotated with GC-FID peak starts and ends. Note the "wandering baseline" in (B) that is
absent in (C).

Figure 2: First forward difference-processed EAD responses to 33 compounds from the test mixture,
colored by compound type (note that the Y-axis scaling differs for (C)). Note for caryophyllene oxide*:
this box includes responses to ethyl laurate, as the peaks were merged in the GC-FID annotation
thresholding step so are treated as a single peak by the analysis. (A) Data from three *Manduca sexta*
females. (B) Data from three *Manduca sexta* males. (C) Data from four *Bombus terrestris audax* workers.

Figure 3: Antennal responses of three female *Manduca sexta* ("MsextaF[1-3]"), three male *Manduca sexta* ("MsextaM[1-3]"), and four *Bombus terrestris audax* workers ("Bterr[1-4]") expressed as a ratio of the significant hit frequency of each compound to that of the significant hit frequency of the initial hexane solvent peak ("value"), with warmer colors and larger circles denoting higher ratios and blue and smaller circles denoting lower ratios. Absent points indicate the specific peak was not detected in the GC-FID run for a given insect.

TABLES

Table 1: Compounds used in the EAD test mixture with details of their retention time, compound
skeleton type and subtype or modification categorization, and average significant hit frequency (spikes
per second, bold text indicates higher than first hexane peak frequency for that species and sex/caste)
for female and male *Manduca sexta* hawkmoths and worker *Bombus terrestris audax* bumble bees. The
ratio after the significant hit frequency indicates the number of samples where the given compound
elicited a greater response than the main hexane peak versus the total number of samples with that
peak present (bold: >50% of samples reacted to that peak for a given species-sex/caste combination).
Unknown superscripts: 1[A]: an unclear oxygenated cyclic molecule from GC-MS results; 2[B]: likely an
unknown sesquiterpenoid oxide from GC-MS analysis results.

Retention time (GC-FID, minutes)	Compound identity	Skeleton	Modifications / subcategory	Average *Manduca sexta* female response (significant hit frequency, spikes/second)	Average *Manduca sexta* male response (significant hit frequency, spikes/second)	Average *Bombus terrestris audax* worker response (significant hit frequency, spikes/second)
1.973	Hexane, main peak	FAD	None	0.843 (NA)	0.896 (NA)	0.229 (NA)
2.687	Hexane, small peak	FAD	None	0.606 (1/3)	8.485 (1/3)	0 (0/1)
3.755	Ethyl L-lactate	FAD	Ester	0.700 (1/3)	**0.867 (2/3)**	**0.483 (4/4)**
4.291	(*Z*)-3-hexen-1-ol, shoulder peak	FAD	Alcohol	0.663 (1/3)	1.292 (1/3)	**0.413 (2/3)**
4.458	(*Z*)-3-hexen-1-ol, main peak	FAD	Alcohol	**4.889 (3/3)**	4.000 (1/3)	0.055 (0/4)
4.741	Isoamyl acetate	FAD	Ester	**7.566 (3/3)**	0.818 (1/3)	0.165 (1/4)
5.184	Ethyl valerate	FAD	Ester	**13.438 (3/3)**	2.519 (1/3)	0.594 (2/4)
5.716	(1R)-(+)-α-pinene	Terpenoid	Monoterpene	0.123 (0/3)	0.494 (1/3)	0.251 (1/4)
5.985	Camphene	Terpenoid	Monoterpene	0 (0/3)	0.160 (0/3)	0.276 (2/4)
6.292	Benzaldehyde	Aromatic	Aldehyde	**0.657 (2/3)**	0.845 (1/3)	**1.752 (4/4)**
7.200	(*Z*)-3-hexenyl acetate	FAD	Ester	**21.919 (3/3)**	0.657 (1/3)	**0.730 (3/4)**
7.505	(R)-(+)-limonene	Terpenoid	Monoterpene	0.741 (1/3)	0 (0/3)	0.348 (2/4)
7.847	Phenylacetaldehyde	Aromatic	Aldehyde	**5.374 (3/3)**	**1.455 (2/3)**	0.265 (2/4)
8.222	Acetophenone	Aromatic	Ketone	**2.871 (3/3)**	0.827 (1/3)	**1.152 (4/4)**
8.691	Methyl benzoate	Aromatic	Ester	**4.202 (3/3)**	0.448 (0/3)	0.210 (1/4)
9.355	(1R)-(+)-nopinone	Terpenoid	Monoterpenoid ketone	**1.137 (2/3)**	0.879 (1/3)	0.502 (2/4)
9.818	Benzyl acetate	Aromatic	Ester	**9.561 (3/3)**	0.155 (0/3)	0.190 (1/4)
10.279	1-phenylethyl	Aromatic	Ester	**2.304 (2/3)**	0.637 (1/3)	0.337 (1/4)

	acetate					
10.511	(1S)-(-)-verbenone	Terpenoid	Monoterpenoid ketone	**3.154 (3/3)**	0.599 (0/3)	**0.299 (3/4)**
10.609	Unknown 1[A]	Unclear	Unclear	**2.241 (3/3)**	**1.322 (2/3)**	0.282 (1/2)
11.240	Geraniol	Terpenoid	Monoterpenoid alcohol	**2.583 (2/3)**	0.778 (1/3)	**0.491 (3/4)**
11.459	(*E*)-cinnamaldehyde	Aromatic	Aldehyde	**1.645 (2/3)**	1.021 (1/3)	0.167 (1/4)
11.777	Indole	Nitrogenous	Aromatic	0.788 (1/3)	0.896 (1/3)	0 (0/4)
12.691	3-phenylpropyl acetate	Aromatic	Ester	1.189 (1/3)	0.207 (1/3)	**1.057 (4/4)**
13.254	β-caryophyllene	Terpenoid	Sesquiterpene	0.496 (1/3)	**0.780 (2/3)**	1.038 (4/4)
13.683	α -humulene	Terpenoid	Sesquiterpene	0.240 (0/3)	0.863 (1/3)	0.200 (2/4)
14.111	Phenethyl 2-methylbutyrate	Aromatic	Ester	0 (0/3)	0.159 (0/3)	0.235 (1/4)
14.285	Methyl isoeugenol	Aromatic	Alcohol	0.387 (0/3)	0.731 (1/3)	0.450 (2/4)
15.265	Caryophyllene oxide + ethyl laurate (peaks merged by FID peak identification script)	Terpenoid + FAD	Sesquiterpenoid oxide + ester	0.816 (1/3)	1.551 (1/3)	0.727 (2/4)
15.476	Unknown 2[B]	Terpenoid	Unknown sesquiterpenoid oxide	0 (0/3)	1.667 (1/3)	1.111 (1/2)
16.474	Farnesol, isomer 1	Terpenoid	Sesquiterpenoid alcohol	0.564 (1/3)	0.154 (0/3)	0.646 (2/4)
16.734	Farnesol, isomer 2	Terpenoid	Sesquiterpenoid alcohol	0.333 (0/3)	0.095 (0/3)	**1.284 (3/4)**
17.189	Benzyl benzoate	Aromatic	Ester	1.323 (1/3)	0.153 (0/3)	0.212 (2/4)

SUPPORTING INFORMATION: SUPPLEMENTAL FIGURE LEGENDS AND TABLE

Figures S1-S5: Raw EAD amplitude for three female hawkmoths. Top pane: GC-FID trace annotated with
peak starts and ends (colored vertical lines) and peak identities. Bottom three panes: individual female
hawkmoth EAD traces annotated with peak starts and ends (colored vertical lines). Figure S1: 0-4
minutes of GC-FID elapsed time. Figure S2: 4-8 minutes. Figure S3: 8-12 minutes. Figure S4: 12-16
minutes. Figure S5: 16-20 minutes. In Figure S5, the large peaks in the second two EAD traces after the
benzyl benzoate interval are mixture puffs to test antennal viability.

Figures S6-S10: Raw EAD amplitude for three male hawkmoths. Top pane: GC-FID trace annotated with
peak starts and ends (colored vertical lines) and peak identities. Bottom three panes: individual male
hawkmoth EAD traces annotated with peak starts and ends (colored vertical lines). Figure S1: 0-4
minutes of GC-FID elapsed time. Figure S2: 4-8 minutes. Figure S3: 8-12 minutes. Figure S4: 12-16
minutes. Figure S5: 16-20 minutes. In Figure S10, the large peaks in the EAD traces after the benzyl
benzoate interval are mixture puffs to test antennal viability.

Figures S11-S15: Raw EAD amplitude for four bumble bee workers. Top pane: GC-FID trace annotated
with peak starts and ends (colored vertical lines) and peak identities. Bottom four panes: individual
bumble bee worker EAD traces annotated with peak starts and ends (colored vertical lines). Figure S11:
0-4 minutes of GC-FID elapsed time. Figure S12: 4-8 minutes. Figure S13: 8-12 minutes. Figure S14: 12-16
minutes. Figure S15: 16-20 minutes.

Figures S16-S20: First forward difference-processed EAD amplitude for three female hawkmoths. Top
pane: GC-FID trace annotated with peak starts and ends (colored vertical lines) and peak identities.
Bottom three panes: individual female hawkmoth first forward difference-processed EAD traces
annotated with peak starts and ends (colored vertical lines) and significant peak hit frequency
(spikes/second), with red indicating frequency over the median hexane solvent threshold. Figure S16: 0-
4 minutes of GC-FID elapsed time. Figure S17: 4-8 minutes. Figure S18: 8-12 minutes. Figure S19: 12-16
minutes. Figure S20: 16-20 minutes. In Figure S20, the large peaks in the second two EAD traces after
the benzyl benzoate interval are mixture puffs to test antennal viability.

Figures S21-S25: First forward difference-processed EAD amplitude for three male hawkmoths. Top
pane: GC-FID trace annotated with peak starts and ends (colored vertical lines) and peak identities.
Bottom three panes: individual male hawkmoth first forward difference-processed EAD traces
annotated with peak starts and ends (colored vertical lines) and significant peak hit frequency
(spikes/second), with red indicating frequency over the median hexane solvent threshold. Figure S21: 0-
4 minutes of GC-FID elapsed time. Figure S22: 4-8 minutes. Figure S23: 8-12 minutes. Figure S24: 12-16
minutes. Figure S25: 16-20 minutes. In Figure S25, the large peaks in the EAD traces after the benzyl
benzoate interval are mixture puffs to test antennal viability.

Figures S26-S30: First forward difference-processed EAD amplitude for four bumble bee workers. Top
pane: GC-FID trace annotated with peak starts and ends (colored vertical lines) and peak identities.
Bottom three panes: individual bumble bee worker first forward difference-processed EAD traces
annotated with peak starts and ends (colored vertical lines) and significant peak hit frequency

(spikes/second), with red indicating frequency over the median hexane solvent threshold. Figure S16: 0-
4 minutes of GC-FID elapsed time. Figure S17: 4-8 minutes. Figure S18: 8-12 minutes. Figure S19: 12-16
minutes. Figure S20: 16-20 minutes.

Figure S31: First forward difference-processed EAD significant peak hit frequency (spikes/second) across
(A,B,C) four chemical skeleton classes (FADs, terpenoids, aromatics, and nitrogenous compounds) and
(D,E,F) six compound modification types tested (none, alcohols, aldehydes, esters, and ketones) for (A,D)
three *Manduca sexta* females, (B,E) three *Manduca sexta* males, and (C,F) four *Bombus terrestris audax*
workers.

Figure S32: Inter-sex EAD response (repeatability) comparisons between the average values of male and
female *Manduca sexta* hawkmoths for each GC-FID peak.

Figure S33: Inter-individual EAD response (repeatability) comparisons between *Manduca sexta*
hawkmoths of the same sex. (A) All pairwise comparisons between three females. (B) All pairwise
comparisons between three males.

Table S1: Names and addresses of vendors from whom authentic reference standards were sourced.

Table S2: Sources of authentic reference standards used in the EAD test mixture.

Table S3: Literature sources demonstrating previous electroantennogram activity or lack thereof in
Manduca sexta and *Bombus terrestris* to the compounds used in this study. Note the frequent
discrepancies in activity (especially for activity in *B. terrestris*), even for compounds with multiple
citations. Compounds with at least one active citation in either species are highlighted in bold.
Compounds shaded in yellow were EAD-active in more than 50% of *M. sexta* males or females and/or *B.*
terrestris worker individuals from this study.