# Canadian Journal of Fisheries and Aquatic Sciences

## Monitoring for fisheries or for fish? Declines in monitoring of salmon spawners continue despite a conservation crisis

| | |
|---|---|
| Journal: | *Canadian Journal of Fisheries and Aquatic Sciences* |
| Manuscript ID | cjfas-2024-0387.R1 |
| Manuscript Type: | Research Article |
| Date Submitted by the Author: | 09-Jun-2025 |
| Complete List of Authors: | Atkinson, Emma; University of Alberta, Biological Sciences Carturan, Bruno; Pacific Salmon Foundation, Salmon Watersheds Program Atkinson, Clare; Pacific Salmon Foundation, Salmon Watersheds Program Bateman, Andrew; Pacific Salmon Foundation; University of Toronto,  Ecology and Evolutionary Biology Connors, Katrina; Pacific Salmon Foundation, Salmon Watersheds Program Hertz, Eric; Pacific Salmon Foundation, Salmon Watersheds Program; Simon Fraser University,  Biological Sciences Peacock, Stephanie J.; Pacific Salmon Foundation, Salmon Watersheds Program |
| Is the manuscript for consideration in a Special Issue or Collection?: | Commemorating the work and ideas of Jeff Hutchings |
| Keyword: | Pacific salmon, Long-term monitoring, Wild Salmon Policy, MARINE FISHERIES < General, Marine conservation |
| | |

SCHOLARONE™
Manuscripts

# 1  Monitoring for fisheries or for fish?

# 2  Declines in monitoring of salmon

# 3  spawners continue despite a

# 4  conservation crisis

**Emma M Atkinson[1][*][+], Bruno S Carturan[2][+], Clare P Atkinson[2], Andrew W Bateman[2,3,4],**

**Katrina Connors[2], Eric Hertz[2,5], Stephanie J Peacock[2,3]**

[1] Department of Biological Sciences, University of Alberta, CCW 405, Biological Sciences
Building, Edmonton, AB T6G 2E9, Canada
[2] Salmon Watersheds Program, Pacific Salmon Foundation, #320, 1385 West 8th Ave,
Vancouver, BC V6H 3V9, Canada
[3] Salmon Coast Field Station, General Delivery, Simoom Sound, BC V0P 1S0
[4] Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks St,
Toronto, ON M5S 3B2, Canada
[5] Department of Biology, Simon Fraser University, B8255, 8888 University Drive, Burnaby, BC
V5A 1S6, Canada

*Corresponding author: emmamargaretatkinson@gmail.com
[+]These authors contributed equally to this work.

# Abstract

Monitoring of salmon in Pacific Canada has been declining for decades. Counts of spawning salmon are critically important, enabling researchers to quantify impacts of stressors, identify when and where management interventions are required to avoid extirpations, and evaluate the efficacy of recovery efforts. These data are more important now than ever, as uniquely adapted spawning populations underlie salmon resilience and their ability to adapt to climate change, and fine-scale data can inform sustainable fishing opportunities including the revitalization of terminal fisheries. We revisit the state of Pacific salmon spawner data from the Yukon to southern British Columbia. Almost two-thirds of salmon populations that were historically monitored have no reported estimates in 2014-2023 – the worst decade for data since broadscale spawner surveys began in the 1950s. We found a positive association between the number of populations monitored and landed value for three of the five Pacific salmon species, suggesting that monitoring is, to some extent, motivated by the information needs of commercial fisheries management. We recommend aligning monitoring objectives and strategic investments to improve monitoring outcomes for salmon, ecosystems, and the communities that depend on them. We emphasize data stewardship, as ensuring access to these baseline data is a cornerstone for rebuilding wild Pacific salmon.

# Introduction

Collecting baseline, fishery-independent data is a pillar of sustainable fisheries management, providing the foundation for evidence-based decisions about the conservation and management of aquatic species (Hilborn and Walters 1992; Punt et al. 2020; Punt 2023). While commercial fisheries data are important components of sustainable fisheries management frameworks, relying solely on fishery-dependent data is broadly acknowledged as insufficient to meet sustainable fisheries objectives (Walsh et al. 2018; Free et al. 2020; Ovando et al. 2021). In a recent example underlining this point, Canada's Pacific salmon fishery was forced to withdraw from the Marine Stewardship Council certification program in 2022, in large part due to a lack of sufficient data to assess salmon abundance trends (Raincoast Conservation Foundation 2019; Marine Stewardship Council 2022). The history of colonial fisheries and their management in Canada is punctuated with consequential failures resulting from insufficiently considering the biology of fished species and the harvest levels they can sustain. From the 1990s collapse of Atlantic cod (Hutchings and Rangeley 2011) to the mid-2000s decline of Fraser sockeye in the Pacific (Peterman and Dorner 2011; Castañeda et al. 2020), retrospective reviews emphasized the importance of decision-making grounded in robust, data-driven science. Maintaining up-to-date and quality-controlled datasets, required for rigorous science, is crucial to effectively managing fish populations. In the absence or decline of commercial fisheries, as in the case of wild salmon in Pacific Canada (Walters et al. 2019), baseline data remain instrumental to stewarding populations through stressors such as climate change and habitat degradation.

Multiple types of data inform efforts to study, monitor, and manage Pacific salmon. The federal
department of Fisheries and Oceans Canada (DFO) tracks how many fish are caught in
commercial fisheries and uses fin-clips or scale samples from captured fishes to provide
information on their genetic origins. Coded wire tags, PIT (Passive Integrated Transponder)
tags, and acoustic tags provide detailed information on ocean survival, movement, and
exploitation rates. These data, and more, help to manage commercial fisheries and are critical
to understand, track, and conserve the diversity and distribution of salmon. An evidence-based
approach to conserving wild salmon, however, also requires simply counting them. Fortunately
for us, and unlike other marine fishes, salmon show up in our proverbial backyards each year
for us to count as they return home to spawn (Fig. 1).
73
In Pacific Canada, the history of counting salmon – in one way or another – as they return to
spawn extends back thousands of years. For millennia, First Nations communities have
developed deep-rooted knowledge of salmon passed down through generations, informed by
diverse approaches to monitoring local populations (Atlas et al. 2021a; Reid et al. 2022). This
paper focuses on salmon counting programs that extend back to the 1930s, which is relatively
modern history but a period for which centrally compiled, broad-scale, and publicly accessible
data are available. The evolution of salmon counting coordinated by the federal government
over the last one hundred years feeds directly into the data available today. Beginning in the
1920s, the "Annual Reports of Salmon Stream and Spawning Grounds," commonly known as
the "BC16s", were released annually (Fig. 2). Initially, reports were sent as personal letters to
federal fisheries supervisors, describing runs as "good", "average", "well seeded", etc.
(McNairnay 1987). In the early 1930s, the reports were formalized to provide an estimated
range for the approximate number of spawners on the spawning grounds. This shift was not
intended to produce actual estimates of abundance but rather to allow considerations of
watershed size, given that "a small stream well seeded could probably have one hundred parent
salmon, but a larger stream equally well seeded would have perhaps five hundred parent
salmon…" (McNairnay 1987). By the 1980s, the estimates for most populations had transitioned
to "point estimates" of the number of individual spawning salmon (Fisheries and Oceans
Canada 2023a). In 1995, responsibility for salmon counting moved to DFO's Science branch,
and counting data were brought together in a centralized database in what is now known as the
New Salmon Escapement Database (NuSEDS), including details of estimation methods and
other descriptive information.
96
Today, NuSEDS houses the most spatially and temporally comprehensive spawner abundance
estimates available for Pacific salmon in Canada. The database is publicly available online and
is intended to be updated annually (Fisheries and Oceans Canada 2024a). The specific process
for feeding data from local salmon systems to NuSEDS varies by region and data processing
protocols are not publicly available. Generally, within a given region, spawner surveys are
conducted annually by DFO, local First Nations, consultants, contracted charter patrol officers,
and/or local community groups contracted by DFO. The methods for conducting spawner counts
vary but generally include stream walks, boat surveys, aerial surveys, snorkel surveys, and
fence counts. During the spawning season, regional DFO staff compile and process spawner
survey data (e.g., generate total abundance estimates from survey counts using a range of

methods). Post-season, DFO NuSEDS staff compile spawner data from across regions and
process them for integration into the public database (Fig. 3). In the current Canadian context,
NuSEDS is fundamental to assessing salmon status (Box 1), estimating population-level effects
of localised stressors (Box 2), and evaluating the efficacy of recovery actions (Box 3).
111
While rarely acknowledged directly, the importance of counting salmon is implicitly woven into
the policy history of Pacific salmon in Canada, most recently embodied in the Wild Salmon
Policy and its plan for implementation (Fisheries and Oceans Canada 2005, 2018). The Wild
Salmon Policy formalised a shift, from viewing salmon simply as "stocks" to be harvested,
toward conservation of the inherent diversity and distribution of wild Pacific salmon. This shift
required federal policy to consider what it means to define, monitor, restore, and maintain
diverse salmon populations. One step in this process included defining "Conservation Units"
(CUs) – genetically and ecologically distinct groups of salmon unlikely to recolonize within a
human lifetime if extirpated (Fisheries and Oceans Canada 2005). Delineating CUs at
biologically and socially relevant scales is not a trivial task and has been a focus of extensive
research (Holtby and Ciruna 2007; Weinstein 2007; Fisheries and Oceans Canada 2009). While
imperfect, CUs laid the foundation for implementing the Wild Salmon Policy and, once defined,
begged the questions: What data are required to monitor and manage CUs? In what state are
those data? Indeed, the first strategy of the Wild Salmon Policy is to "[standardize] monitoring of
wild salmon status" (Fisheries and Oceans Canada 2005). For over 20 years, researchers and
salmon advocates have noted that the number of populations monitored in various regions of
BC has steadily declined (Harvey and MacDuffee 2002; Price et al. 2008, 2017; English 2016;
Atkinson et al. 2020; Price and Moore *In review*). However, a coastwide assessment of the
trajectory of salmon counting as captured by NuSEDS has not been published. Such an
assessment is necessary to understand whether previous calls for increased monitoring have
led to change, especially in light of substantial recent investments by the federal and provincial
governments through the Pacific Salmon Strategy Initiative ($647 million) and the BC Salmon
Restoration and Innovation Fund ($143 million).
135
Inspired by the late Dr. Jeffrey Hutchings and his commitment to investigating the scientific and
policy connections between data and decision-making, we set out to produce an updated,
broad-scale assessment of the state of Pacific salmon spawner abundance data in Canada. We
focused on NuSEDS, the most comprehensive public database of spawner counts in Pacific
Canada and considered monitoring as a two-part process: (1) data collection and (2) data
integration into the public database. We assessed the number of populations and CUs
monitored annually from 1926-2023 and investigated the relationship between commercial
catch, price-per-kilogram, and monitoring effort. Finally, we provide recommendations for
optimizing monitoring to meet the conservation objectives laid out in Canada's Wild Salmon
Policy and for improving data management to ensure that timely and accurate information can
be shared publicly and appropriately in the coming decades.

# Boxes

## Box #1: Assessing salmon status

The primary metric used to assess biological status of CUs is current spawner abundance (Holt
et al. 2009; Pestal et al. 2023; Pacific Salmon Foundation 2024). Estimates of CU-level
abundance can be compared to benchmarks – generally set from historical abundances or
social-biologically relevant reference points (e.g. the number of fish that maximise fisheries
yield) – to determine the status of CUs and determine the potential for fisheries (Chum
Technical Committee 2023; Chinook Technical Committee 2024) or the need for conservation
intervention. This seemingly simple goal involves hidden complexity and assumptions required
to estimate CU-level abundance. In some cases, such as status assessments of geographically
constrained lake-type sockeye, spawner abundance can be obtained from a single survey or
indicator population that is representative of the aggregate stock or CU. However, in most
cases, assumptions must be made to "expand" counts from a limited number of spawner
surveys to estimate spawners for an aggregate unit, such as a CU, Stock Management Unit, or
region. The expanded estimates are also often required for calculating total returns and
exploitation rates, because harvest of salmon does not discriminate between fish that are
returning to monitored and unmonitored streams (with the exception of some mark-selective
fisheries). Thus, to ensure that catch and spawner data correspond to the same component
populations and can be added together, spawner abundance must be extrapolated to
unmonitored populations. Although common expansion approaches are fairly robust (Peacock
et al. 2020), a severe reduction in the number of monitored populations or decline in monitoring
effort over time can invalidate the assumptions of these expansions and create uncertainty
around the estimated aggregate spawner abundance. For example, in 2021, only one out of 61
spawning populations was monitored for the West Haida Gwaii chum CU, with a count of 500
spawners (Pacific Salmon Foundation 2023). Based on the historical proportional contribution of
the monitored stream to the CU, these 500 spawners were expanded to a CU-level abundance
of over 40,000 spawners. Of particular concern is the tendency to stop monitoring salmon
populations that have declined to very low abundances and focus on more abundant
populations, which may overestimate the expanded number of spawners and present
conservation risks to the aggregate on top of potential undocumented extirpations of individual
populations (Price et al. 2008).

## Box #2: Assessing impacts on salmon

The spatial and temporal breadth of spawner counts in Pacific Canada has enabled a wide body
of research into the effects of local stressors on the responses of Pacific salmon populations. By
comparing spawner abundances, or recruits-per-spawner for individual populations, across a
range of exposure to a given stressor, the impacts of that stressor on salmon can be estimated.
With this approach, research has leveraged salmon abundance data to investigate, for example,
the population level-effects of climate change (Hinch and Martins 2011; Malick et al. 2017),
parasite transmission from salmon farms (Connors et al. 2010; Krkošek and Hilborn 2011;

Peacock et al. 2013), marine mammal predation (Nelson et al. 2019), and forestry (Holtby 1988;
Bradford and Irvine 2000; Wilson et al. 2021). All these studies examined stressors that show a
range of effect sizes at fine spatial resolution in freshwater or near-shore marine environments
(e.g., watersheds or inlets), and many relied on salmon spawner abundance records assembled
and maintained by DFO in the New Salmon Escapement Database (NuSEDS) to quantify
salmon responses at an appropriate spatial resolution. More spatially constrained data, such as
survival estimates from coded wire tags of select indicator stocks, or data with coarser spatial
resolution, such as catch within Pacific Fisheries Management Areas, fail to represent the effect
sizes and resolution of many pressures. Thus, the decline of spawner abundance monitoring in
Pacific Canada limits the ability to quantify effects of localised freshwater and near-shore
stressors on salmon abundance and productivity.

## 197   Box #3: Assessing the efficacy of recovery actions

Monitoring spawner abundances before and after recovery actions is essential to track progress
towards recovery and facilitate adaptive management. In the past five years, the Canadian
government has renewed efforts to address the declining abundance of Pacific salmon by
establishing systematic planning processes such as rebuilding plans, prescribed in revisions to
the *Fisheries Act,* and Integrated Planning for Salmon Ecosystems, piloted under the Pacific
Salmon Strategy Initiative (Fisheries and Oceans Canada 2022, 2024b). For example, resource-
intensive efforts led by DFO are underway to create a rebuilding plan for West Coast Vancouver
Island Chinook salmon and pilot Integrated Plans for Salmon Ecosystems in the Thompson-
Shuswap, Nicola, and Yukon watersheds. While historical spawner abundance data may be
sufficient to inform planning and implementation, assessing the success of interventions may be
impossible if monitoring declines continue. The efficacy of recovery actions undertaken at finer
spatial scales, such as localized habitat restoration or water management interventions, can
only be understood by monitoring the affected populations rather than aggregate stocks.
Mismatches between recovery strategies and monitoring coverage, both temporally and
spatially, may result in instances where significant investments are made in recovery planning
and implementation, but the success of such efforts cannot be demonstrated and may be
unable to inform future actions.

## 215   Methods

The terminology used for different biological and management scales of salmon is a common
source of confusion in analyses such as ours, so we begin with some definitions. For the
purpose of our analysis, we define a "population" as the spawners returning to a unique
spawning site (e.g., stream, river, or lake) within a CU. These populations mostly reflect the
biological unit for which spawner abundances are reported in NuSEDS (Fisheries and Oceans
Canada 2024a), except that we consider pink salmon counted at the same site in even- and
odd-years as separate populations, consistent with the evidence for genetically distinct even-
and odd-year lines (Beacham et al. 2012). A relatively small number ($n = 87$) of NuSEDS
populations could not be reliably assigned to a CU; these populations were kept to count the

number of populations monitored and were excluded to determine the proportion of CUs
monitored (cf. Supplement A). We only included data in our analysis that were available through
NuSEDS, and thus define 'monitoring' as data collection *and* integration into NuSEDS. In many
regions, spawner surveys may be undertaken but the data are either not submitted to or not
included in NuSEDS. This potential reporting bias is addressed in recommendation #4 in the
Discussion.
231
We downloaded two datasets from NuSEDS on January 21st, 2025 (Fisheries and Oceans
Canada 2024a). The first dataset (*all_areas_nuseds*) contains the annual counts (e.g. number
of natural adult and jack spawners) for each population. The second dataset
(*conservation_unit_system_sites*) associates each population to its respective CU and site. We
merged the two datasets to associate each population's time series to its CU. There were
numerous errors and omissions between these two datasets including duplicate time series,
abundance estimates assigned to wrong populations or CUs, or missing CU information. We
undertook a systematic review of the data to identify and correct these cases (cf. Supplement A
for details and access to R code). The resulting cleaned spawner survey data are available
online (Carturan and Peacock 2024).
242
243  We assigned each population to one of nine broad-scale regions based on the site location
244  (latitude/longitude) in NuSEDS to assess broad spatial patterns in monitoring (Fig. 1). The final
245  dataset contained spawner abundance data from 1926 to 2023 for the five Pacific salmon
246  species (Chinook, chum, coho, pink, and sockeye), 392 CUs, 6,973 populations, and 2,345 sites
247  (Table 1).
248
From the cleaned NuSEDS data, we calculated the annual "number of populations monitored"
as the number of populations having a numeric estimate in any of the following fields: natural
adult spawners, natural jack spawners, total natural spawners, adult broodstock removals, jack
broodstock removals, total broodstock removals, other removals, or total return to river. There
were 3,449 yearly counts (2%) with a maximum value of zero across these fields. We found
abrupt increases in the frequency of zeroes starting in certain years and only for certain regions
and species (e.g., Fig. S6). Given the available information, it was not possible to determine
whether these zeros represented monitored populations with true zero counts. As such, we
opted to conduct the analysis under two different assumptions: in the main text we present
results assuming that counts of zero indicate no monitoring and in Supplement B, we present
results under the assumption that monitoring did occur when zero spawners are reported.
260
We divided the number of populations monitored each year by the total number of populations
monitored to obtain the annual "proportion of populations monitored". The annual "proportion of
CUs monitored" was the number of CUs with at least one population monitored divided by the
total number of CUs (which differed in even and odd years because pink salmon CUs are only
monitored every other year). We summarised declines in monitoring for each species and
region as the average change per year in the number of populations monitored. We estimated
the average change per year as the slope of a linear regression model fitted to the number of
populations monitored each year from the peak of monitoring in 1986 to 2023.

Spawner monitoring arises, in part, from the need to make informed commercial fishery
management decisions, such as defining Total Allowable Catch, estimating exploitation rates,
etc. If commercial fisheries are a significant driver for monitoring, reducing the size, value,
and/or spatial extent of these fisheries might be expected to result in corresponding declines in
monitoring given lower urgency or fewer requirements to generate quantitative estimates that
rely on abundance data. To investigate whether and for which species this relationship existed,
we quantified the association between the number of populations surveyed and commercial
catch (kilograms), price per kilogram, and species. We defined two models to compare the
effect of landed value (the product of catch and price) and catch alone. We fitted a negative
binomial generalised linear model (GLM) with the log-link function and compared the two
models using AIC and pseudo-$R^2$. Further details of model fitting and assessment can be found
in Supplement B. We downloaded catch data from the North Pacific Anadromous Fish
Commission website (NPAFC; 2024) in July 2024. We considered the Canadian, commercial
catches expressed in kilograms for each of the five Pacific salmon species. We imported the
price-per-kg for each species from DFO's Economic Analysis (Fisheries and Oceans Canada
2023).
285
We used R (version 4.5.0; R Core Team 2025) and the R packages `tidyr` 1.3.1 (Wickham et
al. 2024) and `dplyr` 1.1.4 (Wickham et al. 2023) to handle and analyse the datasets and
produce the figures. We used the R packages `sf` 1.0-20 (Pebesma and Bivand 2023) to group
sites in their respective region, `stats` 4.5.0 to fit the GLMs. The workflow is fully reproducible
by accessing the R scripts and datasets in the Zenodo repository at
https://zenodo.org/records/14248905.

# Findings

Monitoring and reporting of salmon spawner abundances throughout BC and the Yukon peaked
in the mid-1980s, with over 2,700 individual populations across ~1,300 unique sites being
monitored annually (Fig. 2). Since 1986, there has been a marked decline in the number of
populations monitored annually, with an average loss of 41 populations per year (1.5% per year)
in the NuSEDS data from 1986 to 2023. Across 6,973 unique populations (Table 1), only 37%
have spawner records in the most recent decade (2014-2023) compared to 69% in the period
1980-1989 (i.e., a decline of 31%).
300
Coho and chum populations saw the greatest decline in monitoring, particularly for Vancouver
Island & Mainland Inlets, the Central Coast, and Haida Gwaii (Fig. 4c,f,g, Fig. S1b,c). Pink
salmon also saw major reductions in monitoring on the Central Coast, Skeena, and Haida Gwaii
(Fig. 4c,f,e, Fig. S1d). Fraser Chinook and Fraser sockeye saw the only increases in monitoring
since the 1980s among all regions and species (Fig. 5), although the number of Fraser Chinook
populations monitored peaked later, in 2000, and has also since declined (Fig. 4h).
307
Trends in monitoring effort across species and regions through time tend to reflect trends in
commercial catches, particularly over the last 40 years (Fig. 2). For example, declines in coho

monitoring in the early 2000s coincided with the moratorium on coho fisheries following a period
of low marine survival that started in the 1990s (Decker et al. 2014). When there was no longer
such a strong need to inform coho fishery management, monitoring diminished. Additionally, this
reduction in coho monitoring may have inadvertently led to a parallel decline in chum
monitoring, most notably in the Vancouver Island & Mainland Inlets region (Fig 3g). Given their
overlapping spawn timing, chum may have been monitored opportunistically during surveys
targeting coho. Despite the increase of many coho populations in the late 2000s and 2010s,
with the reduction of harvest and more recent shift in ocean condition (Wilson et al. 2025),
monitoring has not returned to historical coverage levels. As the Interior Fraser coho
Management Unit recently exceeded historical average abundances for the first time in decades
(Connors et al. 2024), we anticipate the resumption of fisheries. Whether the diversity and
distribution of coho within the Fraser watershed have also recovered, and how vulnerable this
stock is to another collapse, are difficult to assess in the absence of broad-scale river-level
monitoring (Boxes #2-3). Pervasive declines of chum salmon over the past five years (North
Pacific Anadromous Fish Commission 2023; Connors et al. 2024) have added to the urgent
need to better understand the abundance and distribution of these two species in late-season
spawner surveys.
327
We found some empirical support for the hypothesis that monitoring effort reflects economically
important fisheries. There is a positive relationship between the number of monitored
populations and the annual landed commercial value for coho, chum and pink (Fig. 6). The
model with landed value as a predictor was better supported than the model with catch (Δ
AIC=30.2). Overall, there is a strong temporal component to this relationship, with more recent
years having low monitoring effort and landed value. Chinook and sockeye show the lowest
levels of association among species between monitoring effort and landed value, a pattern
reflecting the relatively low monitoring of these species in the early part of the time series when
catches were moderate, and relatively high levels of monitoring as catch declined in the 1990s
and 2000s. Several general and contextual factors may explain the discrepancy observed for
the two species. First, both species hold high social and cultural value independent of their
economic value and may also be more frequently subject to monitoring requirements through
treaty obligations, species-at-risk processes, etc. Second, the relationship for sockeye might be
influenced by the significant decline in Fraser Sockeye returns during the 1990s that prompted
investigations into potential causes and the formulation of recommendations (e.g., Pacific
Fisheries Resource Conservation Council 2002; Lapointe et al. 2003; Cohen 2012). This
notable event may explain why Fraser Sockeye is the only species-region combination where
monitoring effort has increased since the late 1990s (Fig. 4). In the other regions, however, the
total number of monitored sockeye populations have experienced declines and the overall
proportion of monitored sockeye CUs in Pacific Canada has declined (Fig. 7c). Third,
recreational fisheries have been the dominant source of Chinook harvest at least since 2000 – a
trend unique to this species (Fisheries and Oceans Canada 2023). Consequently, commercial
landed value might not represent the species' economic value accurately.
351
Evidence suggests that the value of commercial fisheries influences spawner monitoring effort,
but multiple factors likely contribute to the history of spawner monitoring over time. We cannot

account for every nuance that drives decisions regarding monitoring effort but, with limited
resources, high-profile populations tend to be prioritised. But this approach to prioritisation does
not implicitly guarantee optimal monitoring with respect to meeting conservation objectives such
as those outlined in the Wild Salmon Policy. In an example of this pattern, 14 of the 30 most
frequently monitored Chinook populations in the Vancouver Island & Mainland Inlets region are
"indicator stocks" reported on by the PSC's Chinook Technical Committee, with an average of
64 years of data per indicator population. By contrast, the 255 Chinook populations in the region
that are not indicator stocks have an average of 17 years of data, and 66% of them have not
been monitored at all in the most recent decade (2014-2023).
363
Even within streams, monitoring tends to increase in years with more abundant brood lines. As
a result, pink salmon are often predominantly monitored in either even or odd years (e.g., Fig.
4b-h). In the Vancouver Island & Mainland Inlets region, monitoring was historically predominant
in even years, following the dominant return year (Fig. S3e,f). "Cyclic" sockeye in the Fraser
River, which show marked patterns across four brood lines, tend to have higher monitoring
effort in the higher-abundance "dominant" years, a pattern that has intensified since the early
2000s (Fig. 4h). While this prioritisation may have practical justifications, it also makes it difficult
to detect shifts in dominance patterns – which have already been shown to occur in some pink
salmon populations (Krkošek et al. 2011; Irvine et al. 2014) – and declines in non-dominant
years.
374
A corollary of "monitoring for fisheries" is the dearth of data for less abundant populations.
These monitoring data are, however, essential for understanding patterns of salmon distribution
and diversity, which include both large and small runs. Monitoring only abundant populations
reduces the power to detect shifts in abundance, such as shifts in dominance for pink salmon
(Irvine et al. 2014), or declines in biodiversity, such as the erosion of sub-dominant and off-cycle
lines for Fraser sockeye (Ricker 1997), that may affect the sustainability of management
decisions and the resilience of salmon. Furthermore, ground-based salmon counting surveys
provide valuable information beyond abundance counts. They are an opportunity to monitor the
system more broadly, including documenting disease events, pre-spawn mortality, river
blockages, and high temperatures. Less abundant or declining populations have been termed
"ghost streams" in reference to their undetermined status and potential for extirpations to go
unrecorded (Price et al. 2008).
387
The proportion of CUs with at least one population monitored has also broadly declined since
the mid-1980s across all regions and species (Fig. 7). CUs were only defined after the
implementation of Canada's Wild Salmon Policy in 2005 (Fisheries and Oceans Canada 2005),
which emphasized the importance of monitoring these ecologically and genetically unique
groups of populations. Despite this recognition, monitoring of CUs has declined since 2005
(Price et al. 2017; Price and Moore *In review*), continuing even after 2018 when DFO renewed
its commitment to the policy with the release of the Wild Salmon Policy Implementation Plan
(Fisheries and Oceans Canada 2018).
396

In some regions, patterns in monitoring may reflect a reporting bias rather than an actual
reduction in effort, although it is difficult to disentangle the two if the data are not made
available. For example, no data have been included in NuSEDS for Yukon populations since
2008 (Fig. 4a), yet spawner surveys are reported in other publications (Yukon River Joint
Technical Committee 2025a). In other cases, the apparent disappearance of monitoring may
reflect a shift towards aggregate assessments at more downstream locations. This is the case
for Fraser River pink salmon, for which monitoring of individual spawning populations appears to
have stopped in the 1990s (Fig. 4h) when the Pacific Salmon Commission began recording pink
salmon abundance using a hydroacoustic station near the mouth of the Fraser (Glaser et al.
2025). Although this method provides rapid and accurate counts of the number of pink salmon
returning in-season to the Fraser, and is used to inform fisheries management, the population-
level survival of those fish to their spawning grounds and distribution throughout the Fraser
River basin is now unknown.
410
We define monitoring effort as the number of populations monitored, without accounting for
differences in data quality, which may be a limitation. Indeed, higher quality spawner surveys
(e.g., sonar and video weirs) are more expensive and could therefore be considered to
represent higher monitoring effort than simpler surveys (e.g., stream walks). Since these
surveys were more likely implemented in the later part of the assessed time period, the decline
in the number of populations monitored might reflect, to some extent, the trade-off between
quantity and quality (i.e., higher quality abundance estimates from fewer systems). Although
data quality is reported in NuSEDS, we excluded it from our analysis because consistent data
quality scores are only available from 1998 onward (Fig. S7). However, declines in the number
of populations monitored are evident across all survey quality categories over the available time
period (Fig. S9). This suggests that the reduction in monitored populations is unlikely to be the
result of a strategic shift toward higher-quality surveys focused on fewer populations, but rather
reflects a broader decline in overall monitoring investment.
424
Retaining observations with zero counts in the NuSEDS data yields similar results overall (Table
S1; Figs. S4, S8, S10, S22). The main differences pertain to sockeye populations in the Fraser,
which accounts for 56% of the zero counts, most recorded after 1998 (Fig. S6). This abrupt
reporting of zero counts dampens the magnitude of fluctuations in monitoring effort after 2000
(Fig. S4h) and increases the average annual change in monitored population since 1998 (Fig.
S5). While these differences do not undermine our conclusions, they underscore the extent to
which the interpretation of results hinges on the consistency and reliability of how zero counts
versus NAs are reported.

# Discussion

Spawner abundance monitoring and reporting has historically focused on informing commercial
fisheries management, resulting in monitoring efforts being concentrated on economically
important species and numerically abundant populations. Commercial catches of salmon in BC
and the Yukon have collapsed in recent years (Walters et al. 2019; Yukon River Joint Technical
Committee 2025), along with the effort towards collecting and sharing spawner data that was

previously used to inform management of those fisheries. For the commercial salmon fisheries
that persist, lack of spawner abundance data has led to loss of sustainability certifications
(Raincoast Conservation Foundation 2019; Marine Stewardship Council 2022) which impedes
access to diverse markets that increasingly favour sustainable seafood. While well-managed
salmon fisheries will continue to be sustainable into the future, the current magnitude of
commercial fisheries is very different than it once was, and monitoring priorities must shift to
reflect the changing seascape of salmon fisheries. As public interest and funding shifts towards
salmon conservation and recovery planning, the limitations of past, commercial fishery-oriented
monitoring objectives are becoming apparent: we currently lack the data to assess the status of
diverse Conservation Units or understand threats to their survival (Boxes #1-2). The absence of
this information impedes recovery planning and the assessment of recovery actions (Box #3).
450
We recognize that the patterns we observed may be due to a combination of declining
monitoring effort and potential reporting bias. It takes time and effort for data to be verified and
entered into NuSEDS, creating the possibility of lag that may be misinterpreted as a decline in
effort. However, the long time period over which we observed consistent declines in monitoring
– more than 40 years – suggests that such a lag is not the only factor at play. We observed
some regions and species where monitoring appears to have declined or completely stopped
based on the NuSEDS data even though we know that spawner surveys have continued and
are reported elsewhere. For example, in the Yukon and Transboundary regions, many spawner
surveys are reported on by the Technical Committees tasked with informing salmon
management under the Pacific Salmon Treaty (Transboundary Technical Committee 2022;
Yukon River Joint Technical Committee 2025b), but the very same surveys are inconsistently
updated in NuSEDS (Peacock et al. *In press*). Regardless of whether data continue to be
collected, their usefulness to research and broad-scale prioritization and planning is undermined
if they are not compiled and included in a single, centralized, and publicly accessible database –
NuSEDS.
466
As climate change requires salmon to adapt, tracking changes in their distribution and diversity
through spawner surveys has never been more critical. Monitoring programs that rely on a
handful of key indicator stocks may miss potential changes in diversity, which underpins the
resilience of salmon social-ecological systems (Schindler et al. 2010; Nesbitt and Moore 2016).
Numerically abundant indicator populations may persist as surrounding (unmonitored)
populations decline, and aggregate stocks may continue to have large returns even as the
diversity of their components is undermined (Price et al. 2021). This can lead to apparently
healthy populations, until a tipping point is reached or a catastrophic event threatens the viability
of the aggregate stock. For example, preceding the collapse of Atlantic cod fisheries in eastern
Canada, aggregate offshore surveys continued to suggest healthy cod stocks, while inshore
fishers observed local extirpations – proverbial canaries in the coal mine (Hutchinson 2008).
Climate changes are now exacerbating other pressures on salmon, yet insufficient data hinders
our ability to understand the adaptive capacity – or fragility – of salmon populations.
480

481   Here, we provide recommendations with the goal to encourage and enable strategic
482   investments in spawner abundance monitoring that can support informed, evidence-based
483   decisions about salmon conservation, management, and recovery.

# 484   Recommendations

We, the authors, are non-Indigenous researchers with extensive technical experience working
with large salmon datasets to serve the scientific objectives of a broad range of groups including
academic researchers, non-governmental organisations, federal and provincial governments,
First Nations governments, and other Indigenous organizations. We are best positioned to make
recommendations that would improve the technical utility of salmon abundance data generally
and NuSEDS in particular. We also recognize that data (and their handling) are intertwined with
the decisions they inform, and we draw on our experience and a growing body of literature on
Indigenous Data Sovereignty to offer suggestions intended to be respectful of Indigenous rights.
493
494   Solutions to the evident decline in monitoring will involve increased investments in boots on the
495   ground, but the burden of the costs can be mediated by strategic investments in collecting data,
496   employing new technologies, and making improvements to how data are processed and shared.
497
**Recommendation #1: Optimize spawner monitoring to serve the assessment of salmon**
**diversity, for the purpose of maintaining it.**
In the Wild Salmon Policy (Fisheries and Oceans Canada 2005, 2018), DFO identified
monitoring wild salmon status as the foundation of its strategy to accomplish a core objective:
"restore and maintain healthy and diverse salmon populations". However, our assessment of
current monitoring efforts suggests that the institutional shift required to apply this strategy has
yet to occur (also, see Price and Moore *In review*). We have shown that, historically,
investments in spawner surveys have mirrored commercial catches through time. As
commercial catches have declined to zero in some regions, accompanying declines in
monitoring efforts have undermined our ability to assess salmon status, understand threats, and
implement effective recovery plans. Clarifying monitoring objectives in-practice to include
assessing changes in salmon biodiversity would help inform conservation and recovery efforts
focused on maintaining that diversity, while promoting sustainable management of modern-day
fisheries – balancing local and regional objectives, First Nations rights, and both recreational
and commercial interests.
513
Annual spawner surveys of all ~2,345 locations listed in NuSEDS (Table 1) may not be feasible,
but strategic monitoring programs can be designed to meet clearly stated objectives with
minimal costs, making the most of limited resources. Spawner monitoring focused on assessing
the status of Conservation Units, as outlined in the Wild Salmon Policy, provides a starting point
for assessing and conserving salmon diversity that is necessary to support sustainable fisheries
and ensure salmon resilience in the face of environmental change. With almost 400 individual
CUs (Baxter and Hamilton 2024), many of which are remote and logistically challenging to
monitor, annual monitoring of each may not be possible. When prioritisation among CUs is
necessary, monitoring could aim to distribute effort across all salmon-bearing regions rather

than focus on the regions with the most high-profile populations or historically-renowned
fisheries.
525

Some CUs encompass hundreds of spawning populations and thus monitoring a single site is
likely insufficient. In such cases, the total number of populations of interest may exceed the
capacity for monitoring. Answering the simple question, 'how many populations should we
count?' is deceptively difficult because it depends on the monitoring objectives, the nuances of
particular regions, and the available resources. While it is beyond the scope of this paper to
provide specific numbers, an interested reader can refer to abundant resources on the design
and implementation of salmon monitoring programs (National Centre for Ecological Analysis
and Synthesis 2010). Extensive work has been published on regionally specific monitoring
designs (English 2016; Atlas et al. 2021b), quantifying consequences of missing spawner data
in run reconstructions, and optimal monitoring designs to assess conservation status (Holt et al.
2011; Peacock and Holt 2012). Techniques such as "rotating panel" monitoring designs
(Peacock and Holt 2012) can be employed to maximize the information obtained with limited
resources. These designs, where some indicator populations are assessed annually while other
populations are assessed on a rotating basis, allow one to efficiently detect contractions in the
spatial distributions of spawners.
541

**Recommendation #2: Provide sufficient and consistent resources to spawner monitoring**
**programs to meet stated objectives.**
The Wild Salmon Policy set out a forward-looking roadmap for the conservation of wild Pacific
salmon that included monitoring and assessment of Conservation Units, but the challenge of
following that roadmap has not been met with sufficient and consistently available resources.
Ironically, there have recently been historic investments in salmon ($647 million) through the
Pacific Salmon Strategy Initiative (Government of Canada 2021) and BC Salmon Restoration
and Innovation Fund (Government of Canada et al. 2019). We argue that these investments,
while unprecedented in their total amount, have not led to the stable long-term funding and
coordinated approach needed to rebuild salmon monitoring in Pacific Canada.
552

A major barrier to revitalizing spawner monitoring in Pacific Canada is the apparent lack of
integration and communication between different initiatives within DFO. Rebuilding spawner
monitoring requires long-term funding but it will also require a coordinated approach to identify
short- and long-term priorities, streamline data handling processes, and coordinate within and
between regions. There are likely multiple ways to meet this challenge, but we suggest two
specific targets for reinvestment at each level of the spawner monitoring data flow (Fig. 2). First,
we recommend investing to build capacity within the NuSEDS team to support their work
developing and maintaining a quality-controlled, well-documented database (see
Recommendation #4). Second, we recommend investing in well-staffed regional DFO teams
with capacity to steward spawner data from collection to integration within NuSEDS and support
the many non-DFO groups conducting spawner abundance surveys. Systems with
accountability for collecting, documenting, and making information accessible must be a core
responsibility for all DFO staff tasked with managing data.
566

Investment to reinvigorate spawner monitoring across BC and the Yukon is a common-sense
approach to salmon conservation and management. It is also a practical opportunity to create
local jobs in the many rural coastal and inland communities working to expand and diversify
labour opportunities. Stream-walking jobs to monitor spawning salmon are community based,
provide connection between people and places, and have the potential to be purposeful,
meaningful occupations for people in Pacific Canada. Reviving salmon counting will likely
involve reinvesting in DFO-led stock assessment programs, but in many salmon watersheds the
practical solution may involve supporting local communities and First Nations already
stewarding watersheds and salmon. For example, DFO reported working with 40 unique
Indigenous collaborators in 2023 and 2024 on salmon stock assessment activities (Fisheries
and Oceans Canada 2024c). Many locally based salmon counters are funded through DFO
contracts that have shrunk in value and scope year-over-year (e.g., Billy Proctor, *personal*
*communication*, 2022). Nonetheless, these groups often have ample expertise and enthusiasm,
and sufficient investment can enable them to count more populations.
581
**Recommendation #3: Invest in new technologies that can improve monitoring efficiency.**
The rapid advancement of computer analysis and remote sensing has significant potential to
improve the quality of data and reduce costs. Historically, new monitoring technologies have
focused on providing more in-depth knowledge about survival of particular stocks, rather than
improving broad-scale coverage of monitoring. For example, coded wire tags, PIT tags, and
acoustic tags provide detailed information on ocean survival and movements but are only
deployed on a limited number of stocks, requiring strong assumptions (e.g. about the
representativeness of coded wire tag stocks) to extrapolate findings to broader scales.
Emerging technologies, on the other hand, may make monitoring more efficient, thereby
allowing for increased coverage. The application of computer vision and deep learning to
process salmon sonar and video data, collected at in-river weirs, can reduce the need for
manual review of hundreds of hours of footage, with potential to reduce costs significantly (Atlas
et al. 2023). Uncrewed aerial vehicles (UAVs), commonly known as drones, have been shown –
in amenable systems – to provide reliable indices of salmon spawner abundance for relatively
low cost (Ponsioen et al. 2023). Environmental DNA, or eDNA, is another emerging technology
that may be suitable for detecting, and potentially estimating abundance of, salmonids (Levi et
al. 2019).
599
600    Although these technologies are in their infancy, and cost savings may not be immediate,
601    investing in their development and application in Canada could speed up broad-scale
602    application and its associated benefits. To ensure salmon abundance data maintain their value
603    as long time series, it will be critical that new technologies are applied together with existing
604    monitoring approaches to validate and calibrate between methods.
605
**Recommendation #4: Compile and maintain high-quality spawner data across**
**government regional divisions and organizations.**
In some cases, monitoring gaps may be due to a failure to integrate existing spawner
abundance data into NuSEDS, a process which is slow or inconsistent across regions. NuSEDS
represents the most comprehensive dataset on salmon abundance, and yet we found many

instances where data were not contributed to NuSEDS despite being published by DFO in other
reports or made available on request. We mentioned examples in the Yukon and
Transboundary regions above, but the monitoring gaps for Columbia sockeye and Chinook
salmon (Fig. 4i) can also be filled by looking at alternative sources (Hyatt and Stockwell 2019;
Matylewich et al. 2019). For Fraser chum salmon, the most recent data in NuSEDS are from
2020, four years behind collection, and the last year of data currently available for the Harrison
River (a major chum population of the Fraser watershed) is 2015, almost 10 years ago. This
lack of data centralisation limits the power and application of the data to improve our
understanding and timely management of Pacific salmon (e.g. Boxes #1-3). In the context of
this study, it also hinders our ability to document putative declines in monitoring effort. Following
the collapse of Atlantic cod and declines in Pacific salmon, several noted experts recommended
that all scientific information on stock abundances be made public as soon as it is presented to
DFO (Hutchings et al. 1997). Even 'interim' or 'provisional' data are valuable for timely detection
of conservation risks and should be made available. Any errors could be corrected in future
versions of the data. As an excellent example of this proactive approach, the Pacific Salmon
Commission Secretariat releases 'in-season' estimates of Fraser sockeye abundance that are
retroactively finalized and re-released pending estimates of pre-spawn mortality (Pacific Salmon
Commission 2024). Institutions responsible for publicly funded monitoring are ultimately
accountable to ensure that resulting data are open and accessible in a timely manner.
630
Our analyses revealed numerous errors in NuSEDS that needed correction, including duplicate
entries and mis-assigned population identifiers (undermining the efficacy of the slow-release
approach discussed in the last paragraph). Data anomalies also included inconsistent use of
zero and "NA" values, complicating efforts to discern whether monitoring occurred in these
cases. Inconsistent use of zeroes and NAs hampers even simple investigations of monitoring
efforts such as our analysis. Although many of these errors were easily corrected once
identified, it is unclear if effort is made (or if sufficient capacity exists) to improve quality across
the dataset. At a minimum, a transparent forum for documenting data errors (and ideally a
mechanism to correct those errors) is needed.
640
As monitoring becomes increasingly distributed across DFO, First Nations, and non-
governmental organisations (e.g., academic researchers, community groups, etc.), integrating
data into NuSEDS will require increased investment in capacity to review and clean data to
respect investment in monitoring (see Recommendation #2). Being more willing to share
"preliminary" data, embracing crowd-source error checking, instituting a streamlined mechanism
to correct identified errors, and implementing modern version control could help moderating the
investment required.
648
**Recommendation #5: Work to develop data sharing protocols that facilitate data use**
**while respecting Indigenous Data Sovereignty.**
FAIR data are Findable, Accessible, Interoperable, and Reusable - a set of principles that has
been lauded by the scientific community for almost a decade (Wilkinson et al. 2016). More
recently, the CARE principles for Indigenous Data Governance have also been developed to
address concerns around Indigenous Data Sovereignty. CARE stands for data that are used for

**collective** benefit, where Indigenous People maintain their **authority** to control their data, non-
Indigenous institutions and individuals share a **responsibility** to engage respectfully, and data
are applied following Indigenous Peoples' **ethics** (Carroll et al. 2021). These two sets of
principles can be complementary, and given that salmon are core to Indigenous livelihoods and
cultures and salmon data are often collected by First Nations staff or members, both FAIR and
CARE principles are applicable here (Carroll et al. 2021).
661
Facilitating access to salmon data requires understanding the hesitations and concerns of the
data owners and custodians and this must be approached contextually. Fear of misuse or
misinterpretation of data is an oft-cited explanation from DFO for limiting access to data, but
ensuring that data are well documented with appropriate caveats and limitations is a logical
antidote that is supported by existing government policy (Fisheries and Oceans Canada 2013).
The Pacific Salmon Commission Fraser River Panel's online tools for viewing and downloading
both in-season and annual data on Fraser pink and sockeye abundance and timing (Pacific
Salmon Commission 2024) demonstrate that timely, transparent, and appropriately caveated
sharing of data is entirely feasible given appropriate resourcing. Rather than causing
misunderstanding, publicly distributed data provide common baseline information that is
foundational for equitable discussions about salmon management (Walters et al. 2008).

# 673 Conclusions

Salmon are a foundational species in Pacific Canada, and counting them each year as they
return home to spawn is integral to assessing and tracking biological status, managing local and
regional fisheries, estimating the impact of myriad stressors, and evaluating recovery efforts.
With the ongoing decline of monitoring salmon populations, both realised and potential
knowledge gained from decades of counting salmon is eroding. Until now, conventional
monitoring and management by DFO has focused on collecting the information needed to
manage commercial harvest. By available public records, the proportion of salmon populations
counted over the past decade (at the scale of individual populations spawning in their respective
rivers/streams/lakes *or* the coarser scale of Conservation Units) is at the lowest it has been
since the growth of counting programs in the 1950s. Pervasive salmon declines and the demise
of the commercial salmon industry demand that we shift monitoring priorities to support the
admirable blueprint set forth by the Wild Salmon Policy. As salmon distributions appear to be
contracting, with pervasive declines at coarse scales (Connors et al. 2024), there is a need for
*increased* monitoring of at-risk populations to avoid the undocumented erosion of salmon
diversity. Revitalizing salmon spawner abundance monitoring – through redefined priorities,
focused investments in salmon communities, and robust data practices – has the potential to
advance shared goals and benefit both salmon themselves and the people and ecosystems that
depend on them.

# 692 References

Atkinson, E.M., Guinchard, C.E., Kamarainen, A.M., Peacock, S.J., and Bateman, A.W. 2020.
The status of Pacific salmon in the Broughton Archipelago, northeast Vancouver Island,

and mainland inlets. Salmon Coast Field Station, Gilford Island, BC.

Atlas, W.I., Ban, N.C., Moore, J.W., Tuohy, A.M., Greening, S., Reid, A.J., Morven, N., White,
E., Housty, W.G., Housty, J.A., Service, C.N., Greba, L., Harrison, S., Sharpe, C., Butts,
K.I.R., Shepert, W.M., Sweeney-Bergen, E., MacIntyre, D., Sloat, M.R., and Connors, K.
2021a. Indigenous Systems of Management for Culturally and Ecologically Resilient
Pacific Salmon (Oncorhynchus spp.) Fisheries. BioScience **71**(2): 186–204.
doi:10.1093/biosci/biaa144.

Atlas, W.I., Connors, K., Honka, L., Moody, J., Service, C.N., Brown, V., Reid, M., Slade, J.,
McGivney, K., Nelson, R., Hutchings, S., Greba, L., Douglas, I., Chapple, R., Whitney,
C., Hammer, H., Willis, C., and Davies, S. 2021b. A Salmon Monitoring & Stewardship
Framework for British Columbia's Central Coast. Pacific Salmon Foundation, Vancouver,
BC, Canada.

Atlas, W.I., Ma, S., Chou, Y.C., Connors, K., Scurfield, D., Nam, B., Ma, X., Cleveland, M.,
Doire, J., Moore, J.W., Shea, R., and Liu, J. 2023. Wild salmon enumeration and
monitoring using deep learning empowered detection and tracking. Frontiers in Marine
Science **10**. doi:https://doi.org/10.3389/fmars.2023.1200408.

Baxter, B., and Hamilton, S. 2024. Pacific Salmon Conservation Units, Sites & Status. Open
Data Canada. Available from https://open.canada.ca/data/en/dataset/1ac00a39-4770-
443d-8a6b-9656c06df6a3 [accessed 14 November 2024].

Beacham, T.D., McIntosh, B., MacConnachie, C., Spilsted, B., and White, B. 2012. Population
structure of pink salmon (Oncorhynchus gorbuscha) in British Columbia and
Washington, determined with microsatellites. Fisheries Bulletin **110**: 242–256.

Bradford, M.J., and Irvine, J.R. 2000. Land use, fishing, climate change, and the decline of
Thompson River, British Columbia, coho salmon. Can. J. Fish. Aquat. Sci. **57**(1): 13–16.
doi:10.1139/f99-283.

Carroll, S.R., Herczog, E., Hudson, M., Russell, K., and Stall, S. 2021. Operationalizing the
CARE and FAIR Principles for Indigenous data futures. Sci Data **8**(1): 108.
doi:10.1038/s41597-021-00892-0.

Carturan, B., and Peacock, S. 2024, November 20. Cleaned spawner survey data. Zenodo.
doi:10.5281/zenodo.14194638.

Castañeda, R.A., Burliuk, C.M.M., Casselman, J.M., Cooke, S.J., Dunmall, K.M., Forbes, L.S.,
Hasler, C.T., Howland, K.L., Hutchings, J.A., Klein, G.M., Nguyen, V.M., Price, M.H.H.,
Reid, A.J., Reist, J.D., Reynolds, J.D., Van Nynatten, A., and Mandrak, N.E. 2020. A
Brief History of Fisheries in Canada. Fisheries **45**(6): 303–318. doi:10.1002/fsh.10449.

Chinook Technical Committee. 2024. Annual Report of Catch and Escapement for 2023.
Available from https://www.psc.org/reports/tcchinook-24-01.

Chum Technical Committee. 2023. 2019 Post-Season Summary Report. Pacific Salmon
Commission Joint Chum Technical Committee **3–11**: viii + 60.

Cohen, B.I. 2012. Commission of Inquiry into the decline of Sockeye Salmon in the Fraser
River. Final Report, Canada. Privy Council., Ottawa, Ontario. Available from
https://publications.gc.ca/site/eng/432516/publication.html.

Connors, B.M., Krkošek, M., Ford, J., and Dill, L.M. 2010. Coho salmon productivity in relation
to salmon lice from infected prey and salmon farms: Salmon populations and parasites
from aquaculture. Journal of Applied Ecology **47**(6): 1372–1377. doi:10.1111/j.1365-
2664.2010.01889.x.

Connors, K., Jones, E., Peacock, S., and Belton, K. 2024. State of Salmon 2024 Report. Pacific
Salmon Foundation, Vancouver, BC. Available from https://doi.org/10.60740/3867-G827.

Decker, A.S., Hawkshaw, M.A., Patten, B.A., Sawada, J., and Jantz, A.L. 2014. Assessment of
the Interior Fraser Coho Salmon (Oncorhynchus kisutch) Management Unit Relative to
the 2006 Conservation Strategy Recovery Objectives. Research Document, Fisheries
and Oceans Canada. Available from https://waves-vagues.dfo-mpo.gc.ca/library-

746            bibliotheque/360009.pdf.
747    English, K.K. 2016. Review of Escapement Indicator Streams for the North and Central Coast
748            Salmon Monitoring Program Final Report. LGL Limited.
749    Fisheries and Oceans Canada. 2005. Canada's Policy for Conservation of Wild Pacific Salmon.
750            Fisheries and Oceans Canada, Vancouver, BC. Available from
751            https://salmonwatersheds.ca/document/lib_578/ [accessed 23 March 2022].
752    Fisheries and Oceans Canada. 2009. Framework for implementation of the Wild Salmon Policy:
753            Initial lists of Conservation Units for British Columbia. Science Advisory Report,
754            Fisheries and Oceans Canada.
755    Fisheries and Oceans Canada. 2013. Policy for Scientific Data. Integrated Science Data
756            Management Service, Ottawa, Canada. Available from https://www.dfo-
757            mpo.gc.ca/about-notre-sujet/publications/science/datapolicy-politiquedonnees/index-
758            eng.html.
759    Fisheries and Oceans Canada. 2018. Wild Salmon Policy 2018-2022 Implementation Plan.
760            Fisheries and Oceans Canada, Vancouver, BC.
761    Fisheries and Oceans Canada. 2022. Regulations Amending the Fishery (General) Regulations:
762            SOR/2022-73. Available from https://gazette.gc.ca/rp-pr/p2/2022/2022-04-13/html/sor-
763            dors73-eng.html [accessed 26 November 2024].
764    Fisheries and Oceans Canada. 2023a. NuSEDS-New Salmon Escapement Database System
765            Background Information. Available from
766            https://open.canada.ca/data/en/dataset/c48669a3-045b-400d-b730-
767            48aafe8c5ee6/resource/0f29fda6-2402-3766-9fe6-5dfd504d7add [accessed 11
768            November 2024].
769    Fisheries and Oceans Canada. 2023b. B.C. commercial salmon fisheries: A 70 year
770            perspective. Available from https://www.pac.dfo-mpo.gc.ca/analyses-econom-
771            analysis/analyses/econ-perspective-salmon-saumon-eng.html [accessed 17 April 2025].
772    Fisheries and Oceans Canada. 2024a. NuSEDS - New Salmon Escapement Database System.
773            Pacific Biological Station. Available from
774            https://open.canada.ca/data/en/dataset/c48669a3-045b-400d-b730-48aafe8c5ee6
775            [accessed 22 January 2024].
776    Fisheries and Oceans Canada. 2024b. Integrated Planning for Salmon Ecosystems. Available
777            from https://www.pac.dfo-mpo.gc.ca/pacific-smon-pacifique/ipse-pies-base-eng.html
778            [accessed 26 November 2024].
779    Fisheries and Oceans Canada. 2024c. Fieldnotes: Spotlight on salmon stock assessment.
780            Fisheries and Oceans Canada.
781    Free, C.M., Jensen, O.P., Anderson, S.C., Gutierrez, N.L., Kleisner, K.M., Longo, C., Minto, C.,
782            Osio, G.C., and Walsh, J.C. 2020. Blood from a stone: Performance of catch-only
783            methods in estimating stock biomass status. Fisheries Research **223**: 105452.
784            doi:10.1016/j.fishres.2019.105452.
785    Glaser, D.M., Connors, B.M., Dionne, K., and Huang, A.-M. 2025. Estimating precautionary
786            approach reference points and assessing consequences of harvest control rules for
787            Fraser River pink salmon (Oncorhynchus gorbuscha). Research Document, Fisheries
788            and Oceans Canada, Ottawa ON.
789    Government of Canada, F. and O.C. 2021, June 8. Pacific Salmon Strategy Initiative. Available
790            from https://www.dfo-mpo.gc.ca/campaign-campagne/pss-ssp/index-eng.html [accessed
791            28 October 2024].
792    Government of Canada, Fisheries and Oceans Canada, and Province of British Columbia.
793            2019, March 15. British Columbia Salmon Restoration and Innovation Fund. Available
794            from https://www.dfo-mpo.gc.ca/fisheries-peches/initiatives/fish-fund-bc-fonds-peche-
795            cb/index-eng.html [accessed 28 October 2024].
Harvey, B., and MacDuffee, M. 2002. Ghost Runs: The Future of Wild Salmon on the North and

Central Coasts of British Columbia. Raincoast Conservation Society, Victoria, BC.
Available from https://www.raincoast.org/2002/12/ghost-runs-the-future-of-wild-salmon-
on-the-north-and-central-coasts-of-british-columbia/ [accessed 17 May 2024].

Hilborn, R., and Walters, C.J. 1992. Stock and Recruitment. *In* Quantitative Fisheries Stock
Assessment: Choice, Dynamics and Uncertainty. Springer US, Boston, MA. pp. 241–
296. doi:10.1007/978-1-4615-3598-0_7.

Hinch, S.G., and Martins, E.G. 2011. A review of potential climate change effects on survival of
Fraser River sockeye salmon and an analysis of interannual trends in en-route loss and
pre-spawn mortality. Fisheries and Oceans Canada, Vancouver, BC.

Holt, C.A., Cass, A., Holtby, B., and Riddell, B. 2009. Indicators of Status and Benchmarks for
Conservation Units in Canada's Wild Salmon Policy. Research Document, Fisheries and
Oceans Canada.

Holt, K.R., Peterman, R.M., and Cox, S.P. 2011. Trade-offs between monitoring objectives and
monitoring effort when classifying regional conservation status of Pacific salmon
(Oncorhynchus spp.) populations. Can. J. Fish. Aquat. Sci. **68**(5): 880–897. NRC
Research Press. doi:10.1139/f2011-022.

Holtby, B.L., and Ciruna, K.A. 2007. Conservation Units for Pacific Salmon under the Wild
Salmon Policy. Canadian Science Advisory Secretariat Research Document **070**.
Available from https://salmonwatersheds.ca/document/lib_4/ [accessed 19 October
2023].

Holtby, L.B. 1988. Effects of Logging on Stream Temperatures in Carnation Creek British
Columbia, and Associated Impacts on the Coho Salmon (Oncorhynchus kisutch).
Canadian Journal of Fisheries & Aquatic Sciences **45**: 502–515.

Hutchings, J.A., and Rangeley, R.W. 2011. Correlates of recovery for Canadian Atlantic cod
(Gadus morhua). Can. J. Zool. **89**(5): 386–400. NRC Research Press. doi:10.1139/z11-
022.

Hutchings, J.A., Walters, C., and Haedrich, R.L. 1997. Is scientific inquiry incompatible with
government information control? Can. J. Fish. Aquat. Sci. **54**(5): 1198–1210.
doi:10.1139/f97-051.

Hutchinson, W.F. 2008. The dangers of ignoring stock complexity in fishery management: the
case of the North Sea cod. Biology Letters **4**(6): 693–695.

Hyatt, K.D., and Stockwell, M.M. 2019. Chasing an Illusion? Successful Restoration of
Okanagan River Sockeye Salmon Oncorhynchus nerka in a Sea of Uncertainty. *In* From
Catastrophe to Recovery: Stories of Fish Management Success. American Fisheries
Society. American Fisheries Society. pp. 65–100. Available from
https://salmonwatersheds.ca/library/lib_515/.

Irvine, J.R., Michielsens, C.J.G., O'Brien, M., White, B.A., and Folkes, M. 2014. Increasing
Dominance of Odd-Year Returning Pink Salmon. Transactions of the American Fisheries
Society **143**(4): 939–956. doi:10.1080/00028487.2014.889747.

JTC. 2024. Yukon River Salmon 2023 Season Summary and 2024 Season Outlook. Yukon
River Joint Technical Committee Report **YUKON JTC (24)-01**: vi + 181 p.

Krkošek, M., and Hilborn, R. 2011. Sea lice (Lepeophtheirus salmonis) infestations and the
productivity of pink salmon (Oncorhynchus gorbuscha) in the Broughton Archipelago,
British Columbia, Canada. Can. J. Fish. Aquat. Sci. **68**(1): 17–29. doi:10.1139/F10-137.

Krkošek, M., Hilborn, R., Peterman, R.M., and Quinn, T.P. 2011. Cycles, stochasticity and
density dependence in pink salmon population dynamics. Proc. R. Soc. B. **278**(1714):
2060–2068. doi:10.1098/rspb.2010.2335.

Lapointe, M., Cooke, S.J., Hinch, S.G., Farrell, A.P., Jones, S., MacDonald, S., Patterson, D.,
Healey, M.C., and Van Der Kraak, G. 2003. Late-run Sockeye Salmon in the Fraser
River, British Columbia, are Experiencing Early Upstream Migration and Unusually High
Rates of Mortality—What is Going On? Vancouver, BC, Canada. p. 14. Available from

https://watershedwatch.ca/wp-content/uploads/2011/04/Exh-72-NonRT.pdf.
Levi, T., Allen, J.M., Bell, D., Joyce, J., Russell, J.R., Tallmon, D.A., Vulstek, S.C., Yang, C.,
and Yu, D.W. 2019. Environmental DNA for the enumeration and management of Pacific
salmon. Molecular Ecology Resources **19**(3): 597–608. doi:10.1111/1755-0998.12987.
Malick, M.J., Cox, S.P., Mueter, F.J., Dorner, B., and Peterman, R.M. 2017. Effects of the North
Pacific Current on the productivity of 163 Pacific salmon stocks. Fish. Oceanogr. **26**(3):
268–281. doi:10.1111/fog.12190.
Marine Stewardship Council. 2022. British Columbia Salmon MSC Notice of Withdrawal. Lloyd's
Register. Available from https://fisheries.msc.org/en/fisheries/british-columbia-
salmon/@@assessments [accessed 26 November 2024].
Matylewich, M., Oatman, M., Parken, C., Riddell, B., Tweit, B., Wright, H., Baldwin, C., Garrison,
R., Lothrop, R., and McGrath, E. 2019. A Summary of Okanagan Chinook Information
Requested by the Pacific Salmon Commission. Pacific Salmon Commission Technical
Report No. 42. Available from https://salmonwatersheds.ca/library/lib_502/.
McNairnay, H.R. 1987. "BC 16" Annual report of salmon stream and spawning grounds: An
Historical Review. Fisheries and Oceans Canada. Available from https://waves-
vagues.dfo-mpo.gc.ca/library-bibliotheque/358188.pdf.
National Centre for Ecological Analysis and Synthesis. 2010. Salmon Monitoring Advisor.
Available from https://groups.nceas.ucsb.edu/monitoring-kb.html.
Nelson, B.W., Walters, C.J., Trites, A.W., and McAllister, M.K. 2019. Wild Chinook salmon
productivity is negatively related to seal density and not related to hatchery releases in
the Pacific Northwest. Can. J. Fish. Aquat. Sci. **76**(3): 447–462. doi:10.1139/cjfas-2017-
0481.
Nesbitt, H.K., and Moore, J.W. 2016. Species and population diversity in Pacific salmon
fisheries underpin indigenous food security. Journal of Applied Ecology **53**(5): 1489–
1499. doi:10.1111/1365-2664.12717.
North Pacific Anadromous Fish Commission. 2023. The Status and Trends of Pacific Salmon
and Steelhead Trout Stocks with Linkages to their Ecosystem. North Pacific
Anadromous Fish Commission, Vancouver, Canada. Available from
https://doi.org/10.23849/LOEX7610.
North Pacific Anadromous Fish Commission (NPAFC). 2024. NPAFC statistics: description of
Pacific salmonid catch and hatchery release data files (updated 28 June 2024). North
Pacific Anadromous Fish Commission, Vancouver, B.C. Available from
https://www.npafc.org.
Ovando, D., Free, C.M., Jensen, O.P., and Hilborn, R. 2021. A history and evaluation of catch-
only stock assessment models. Fish and Fisheries. doi:10.1111/faf.12637.
Pacific Fisheries Resource Conservation Council. 2002. Annual Report 2001–2002. Pacific
Fisheries Resource Conservation Council, Vancouver, BC, Canada. Available from
https://psf.ca/wp-content/uploads/2021/10/Download-PDF551-1.pdf.
Pacific Salmon Commission. 2024. PSC Annual Report Application. Release 2024.04 [Updated
2024 apr 03; accessed 2024 Apr 05]. Available from
https://www.psc1.shinyapps.io/PSC_Annual_Fraser/.
Pacific Salmon Foundation. 2023. Haida Gwaii Salmon Assessment Update. Vancouver, BC.
Available from https://salmonwatersheds.ca/document_library_files/lib_534.pdf
[accessed 25 November 2024].
Pacific Salmon Foundation. 2024. Methods for Assessing Status and Trends in Pacific Salmon
Conservation Units and their Freshwater Habitats. Pacific Salmon Foundation,
Vancouver, BC, Canada. Available from https://salmonwatersheds.ca/document/lib_475
[accessed 21 March 2024].
Peacock, S., Honka, L., Bryan, K., Porter, M., Hertz, E., Mather, V., and Connors, K. 2024. The
Status of Salmon and their Habitats in BC's Northern Transboundary Region: 2024

Snapshot Report. Pacific Salmon Foundation, Vancouver, BC. Available from
https://salmonwatersheds.ca/document/lib_600/.
Peacock, S.J., Hertz, E., Holt, C.A., Connors, B., Freshwater, C., and Connors, K. 2020.
Evaluating the consequences of common assumptions in run reconstructions on Pacific
salmon biological status assessments. Canadian Journal of Fisheries and Aquatic
Sciences **77**(12): 1904–1920. Canadian Science Publishing. doi:10.1139/CJFAS-2019-
0432/ASSET/IMAGES/CJFAS-2019-0432IEQ43.GIF.
Peacock, S.J., and Holt, C.A. 2012. Metrics and sampling designs for detecting trends in the
distribution of spawning Pacific salmon (Oncorhynchus spp.). Canadian Journal of
Fisheries and Aquatic Sciences **69**(4): 681–694. doi:10.1139/f2012-004.
Peacock, S.J., Krkošek, M., Proboszcz, S., Orr, C., and Lewis, M.A. 2013. Cessation of a
salmon decline with control of parasites. Ecological Applications **23**(3): 606–620.
doi:10.1890/12-0519.1.
Pebesma, E., and Bivand, R. 2023. Spatial Data Science: With Applications in R. *In* 1st edition.
Chapman and Hall/CRC, New York. doi:10.1201/9780429459016.
Pestal, G., MacDonald, B.L., Grant, S.C.H., and Holt, C.A. 2023. State of the salmon: rapid
status assessment approach for Pacific salmon under Canada's Wild Salmon Policy.
Canadian Technical Report of Fisheries and Aquatic Sciences **3570**: xiv + 200 p.
Peterman, R.M., and Dorner, B. 2011. Fraser River Sockeye Production Dynamics. Cohen
Commission, Vancouver, BC, Canada. Available from www.cohencommission.ca
[accessed 21 September 2021].
Ponsioen, L., Kapralova, K.H., Holm, F., and Hennig, B.D. 2023. Remote sensing of salmonid
spawning sites in freshwater ecosystems: The potential of low-cost UAV data. PLOS
ONE **18**(8): e0290736. Public Library of Science. doi:10.1371/journal.pone.0290736.
Price, M.H.H., Darimont, C.T., Temple, N.F., and MacDuffee, S.M. 2008. Ghost runs:
Management and status assessment of Pacific salmon (Oncorhynchus spp.) returning to
British Columbia's central and north coasts. Canadian Journal of Fisheries and Aquatic
Sciences **65**(12): 2712–2718. doi:10.1139/F08-174.
Price, M.H.H., English, K.K., Rosenberger, A.G., Macduffee, M., and Reynolds, J.D. 2017.
Canada's wild salmon policy: An assessment of conservation progress in British
Columbia. Canadian Journal of Fisheries and Aquatic Sciences **74**(10): 1507–1518.
Canadian Science Publishing. doi:10.1139/cjfas-2017-0127.
Price, M.H.H., and Moore, J.W. In review. From policy to practice: monitoring declines and
Pacific salmon conservation in Canada. Canadian Journal of Fisheries & Aquatic
Sciences.
Price, M.H.H., Moore, J.W., Connors, B.M., Wilson, K.L., and Reynolds, J.D. 2021. Portfolio
simplification arising from a century of change in salmon population diversity and
artificial production. Journal of Applied Ecology **58**(7): 1477–1486. doi:10.1111/1365-
2664.13835.
Punt, A.E. 2023. Those who fail to learn from history are condemned to repeat it: A perspective
on current stock assessment good practices and the consequences of not following
them. Fisheries Research **261**: 106642. doi:10.1016/j.fishres.2023.106642.
Punt, A.E., Dunn, A., Elvarsson, B.Þ., Hampton, J., Hoyle, S.D., Maunder, M.N., Methot, R.D.,
and Nielsen, A. 2020. Essential features of the next-generation integrated fisheries stock
assessment package: A perspective. Fisheries Research **229**: 105617.
doi:10.1016/j.fishres.2020.105617.
R Core Team. 2025. R: A Language and Environment for Statistical Computing. R, R
Foundation for Statistical Computing, Vienna, Austria. Available from https://www.R-
project.org/.
Raincoast Conservation Foundation. 2019. Backgrounder on Canada's Pacific salmon fishery
losing its Marine Stewardship Council certification. Available from

https://www.raincoast.org/2019/10/backgrounder-on-canadas-pacific-salmon-fishery-
losing-its-marine-stewardship-council-certification/ [accessed 26 November 2024].
Reid, A.J., Young, N., Hinch, S.G., and Cooke, S.J. 2022. Learning from Indigenous knowledge
holders on the state and future of wild Pacific salmon. FACETS **7**: 718–740.
doi:10.1139/facets-2021-0089.
Ricker, W.E. 1997. Cycles of abundance among Fraser River sockeye salmon (Oncorhynchus
nerka). Canadian Journal of Fisheries & Aquatic Sciences **54**.
Schindler, D.E., Hilborn, R., Chasco, B., Boatright, C.P., Quinn, T.P., Rogers, L.A., and
Webster, M.S. 2010. Population diversity and the portfolio effect in an exploited species.
Nature **465**(7298): 609–612. doi:10.1038/nature09060.
Transboundary Technical Committee. 2022. Final Estimates of Transboundary River Salmon
Production, Harvest and Escapement and a Review of Joint Enhancement Activities in
2020. Pacific Salmon Commission Transboundary Technical Committee Report **TCTR
(22)-01**: xii + 162 p.
Walsh, J.C., Minto, C., Jardim, E., Anderson, S.C., Jensen, O.P., Afflerbach, J., Dickey-Collas,
M., Kleisner, K.M., Longo, C., Osio, G.C., Selig, E.R., Thorson, J.T., Rudd, M.B.,
Papacostas, K.J., Kittinger, J.N., Rosenberg, A.A., and Cooper, A.B. 2018. Trade-offs for
data-limited fisheries when using harvest strategies based on catch-only models. Fish
Fish **19**(6): 1130–1146. doi:10.1111/faf.12316.
Walters, C., English, K., Korman, J., and Hilborn, R. 2019. The managed decline of British
Columbia's commercial salmon fishery. Marine Policy **101**(May 2018): 25–32. Elsevier
Ltd. doi:10.1016/j.marpol.2018.12.014.
Walters, C.J., Lichatowich, J.A., Peterman, R.M., and Reynolds, J.D. 2008. Report of the
Skeena Independent Science Review panel. A report to the Canadian Department of
Fisheries and Oceans and the British Columbia Ministry of the Environment.
Weinstein, M.S. 2007. Design possibilities for a regional resource management board for the
implementation of the Wild Salmon Policy within the MTTC First Nations' territories.
Watershed Watch Salmon Society, Coquitlam, BC.
Wickham, H., François, R., Henry, L., Müller, K., Vaughan, D., Software, P., and PBC. 2023.
dplyr: A Grammar of Data Manipulation. Available from https://cran.r-
project.org/web/packages/dplyr/index.html [accessed 23 August 2024].
Wickham, H., Vaughan, D., Girlich, M., Ushey, K., Software, P., and PBC. 2024. tidyr: Tidy
Messy Data. Available from https://cran.r-project.org/web/packages/tidyr/index.html
[accessed 23 August 2024].
Wilkinson, M.D., Dumontier, M., Aalbersberg, Ij.J., Appleton, G., Axton, M., Baak, A., Blomberg,
N., Boiten, J.-W., da Silva Santos, L.B., Bourne, P.E., Bouwman, J., Brookes, A.J.,
Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C.T., Finkers, R.,
Gonzalez-Beltran, A., Gray, A.J.G., Groth, P., Goble, C., Grethe, J.S., Heringa, J., 't
Hoen, P.A.C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S.J., Martone, M.E., Mons,
A., Packer, A.L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-
A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M.A., Thompson, M., van
der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft,
K., Zhao, J., and Mons, B. 2016. The FAIR Guiding Principles for scientific data
management and stewardship. Sci Data **3**(1): 160018. Nature Publishing Group.
doi:10.1038/sdata.2016.18.
Wilson, K.L., Atlas, W.I., Whitney, C.K., Reid, M., Service, C.N., Connors, B.M., and Adams,
M.S. 2025, March 3. Using fisheries risk assessment to inform precautionary and
collaborative management in a declining coho salmon fishery. bioRxiv.
doi:10.1101/2025.02.26.640431.
Wilson, K.L., Bailey, C.J., Davies, T.D., and Moore, J.W. 2021. Marine and freshwater regime
changes impact a community of migratory Pacific salmonids in decline. Glob Change

Biol: gcb.15895. doi:10.1111/gcb.15895.
Yukon River Joint Technical Committee. 2025a. Yukon River Salmon 2024 Season Summary
and 2025 Season Outlook. Yukon River Joint Technical Committee Report **YUKON JTC**
**(25)-01**: vi + 181 p.
Yukon River Joint Technical Committee. 2025b. Yukon River Salmon 2024 Season Summary
and 2025 Season Outlook. Yukon River Joint Technical Committee Report **YUKON JTC**
**(25)-01**: vii + 186 p.

# 1008 Acknowledgements

We thank the Pacific Salmon Foundation for ongoing support of the Salmon Watersheds
Program's work to democratise salmon data, and for supporting staff to pursue this analysis. We
thank Michael Price for reading an early draft of the manuscript and providing constructive
feedback. Finally, we acknowledge and thank Dr. Jeff Hutchings, who has inspired our work,
here and elsewhere, most clearly through his commitment to the integrity of the processes that
connect science to decision makers in the context of fisheries.

# 1015 Competing interests

1016 The authors declare no competing interests.

# 1017 Author contributions

Conceptualization - EA, AB, EH, CA
Data curation - BC, SP
Formal analysis - BC, SP
Funding acquisition - KC, EA
Investigation - EA, BC, EH, AB, SP
Methodology - BC, EA, SP, EH, AB
Project administration - EA
Resources - EA, KC
Software - BC, SP, EA
Supervision - SP, KC
Validation - SP, BC
Visualization - BC, EA, SP
Writing – original draft - EA, SP, EH, CA, BC
Writing – review & editing - EA, SP, BC, EH, AB, KC

# 1032 Funding statement

1033 The work was supported by a Vanier Canada Graduate Scholarship from the Natural Sciences
1034 and Engineering Research Council of Canada and a Mitacs Accelerate Scholarship to E.M.A.
1035 The Salmon Watershed Program at Pacific Salmon Foundation supported staff members

# Data availability statement

1038

1039    Cleaned spawner survey data are available at https://zenodo.org/records/14194638. Additional
1040    data and R code to reproduce all our analyses for this submission are archived at
1041    https://zenodo.org/records/14248905.

# Supplement A: Monitoring for fisheries or for fish? Declines in monitoring of salmon spawners continue despite a conservation crisis

**Emma M Atkinson[1][*][+], Bruno S Carturan[2][+], Clare P Atkinson[2], Andrew W Bateman[2,3,4], Katrina Connors[2], Eric Hertz[2,5], Stephanie J Peacock[2,3]**

To access Supplement A, please follow this link:
https://bookdown.org/salmonwatersheds/1_nuseds_collation_atkinson/1_nuseds_collation.html

# Supplement B: Monitoring for fisheries or for fish? Declines in monitoring of salmon spawners continue despite a conservation crisis

**Emma M Atkinson[1*+], Bruno S Carturan[2+], Clare P Atkinson[2], Andrew W Bateman[2,3,4], Katrina Connors[2], Eric Hertz[2,5], Stephanie J Peacock[2,3]**

To access Supplement B, please follow this link:
https://bookdown.org/salmonwatersheds/atkinson_et_al_supp_main/

**Table 1.** Number of Conservation Units (CUs), populations, and sites present in the NuSEDS data in each region after cleaning (zeros and NAs counts were removed). Sites are unique locations where spawners from one or more CUs may be counted. Populations are unique combinations of CUs and sites.

| Region | Number of CUs | Number of populations | Number of sites |
|---|---|---|---|
| Yukon | 15 | 60 | 54 |
| Transboundary | 22 | 65 | 37 |
| Haida Gwaii | 28 | 756 | 233 |
| Nass | 20 | 307 | 82 |
| Skeena | 54 | 777 | 265 |
| Central Coast | 111 | 1681 | 404 |
| Vancouver Island & Mainland Inlets | 83 | 2297 | 666 |
| Fraser | 57 | 1028 | 603 |
| Columbia | 2 | 2 | 1 |
| **Total** | **392** | **6,973** | **2,345** |

**Figure 1**. Map of Pacific Canada showing the nine broad-scale regions (colours) and 2,300 sites (points) where spawner monitoring is recorded in NuSEDS. Open circles reflect sites which were not monitored at least once within the 2014-2023 decade.

**Figure 2.** A timeline of the number of populations for which spawner counts appear in NuSEDS (blue) and total commercial salmon catch (brown) in Pacific Canada, highlighting notable events pertaining to abundance monitoring. A population is a grouping of salmon that can return to breed in the same spawning location, usually a stream or river (but sometimes a smaller geographic area). The blue line depicts the number of populations monitored over time (as inferred from the number of records in NuSED, see Methods for full details). The brown line depicts total commercial salmon fishery catch among five salmon species over time as reported by the North Pacific Anadromous Fish Commission (NPAFC 2024). This figure focuses specifically on contemporary salmon abundance monitoring as curated in NuSEDs but many of these populations have been monitored, in one way or another, for thousands of years.

**Figure 3.** High-level schematic illustrating the data flow and processing steps within the NuSEDS pipeline. General data flow adapted from information provided by the Fisheries & Assessment Data Section at DFO. Spawner abundance surveys are conducted by a range of groups, compiled and processed at a regional level, and subsequently consolidated and processed across regions by NuSEDS staff. Dashed lines highlight places along the data flow where spawner abundance data may be lost from inclusion in NuSEDs.

**Figure 4.** The annual number of salmon populations monitored in NuSEDS, broken down by region (panels a through i) and by species (Chinook, chum, coho, pink, and sockeye).

**Figure 5.** The average annual change in populations monitored since 1986 as estimated via linear regression for each species and region. Dots and horizontal bars represent the slopes and their 95% confidence intervals, respectively.

**Figure 6.** Prediction of the number of populations monitored as a function of landed value (in million of CAD), which is the product between commercial catch (in kg) and price (in CAD per kg). A generalized linear model with a negative binomial distribution and log-link function was fitted with catch, price and species as predictive variables (n = 345). Lines and polygons represent model fits and 95% confidence intervals, respectively; circles represent the data, which span from 1952 and 2020. Zero counts in NuSEDS were filtered out.

**Figure 7.** Annual proportion of conservation units (CUs) with at least one population monitored in NuSEDS.