# Repeated evolution of supergenes on an ancient social chromosome

Aparna Lajmi[1,2,*], Pnina Cohen[1,3], Chih-Chi Lee[1,4,5], Zeev Frenkel[1], Yoann Pellen[1], Eyal Privman[1,*]

[1] Department of Evolutionary and Environmental Biology, Institute of Evolution,
University of Haifa, Haifa, Israel

[2] Department of Life Sciences, Gandhi Institute of Technology and Management, Visakhapatnam, India

[3] Department of Archaeology, Zinman Institute, University of Haifa, Haifa 3498837, Israel.

[4] Agricultural Biotechnology Research Center, Academia Sinica, Taipei 115, Taiwan

[5] Biotechnology Center in Southern Taiwan, Academia Sinica, Tainan 711, Taiwan

* Co-corresponding authors

# Abstract

Supergenes are non-recombining chromosomal regions that code for complex polymorphic traits. With the advance of population genomics, studies in diverse organisms revealed supergenes associated with a myriad of complex traits, ranging from morphology to sexual behavior, from butterfly wing patterns to floral morphs. Two supergenes have been described on non-homologous "social chromosomes" in two ant lineages, *Solenopsis* and *Formica*, that diverged more than 90 million years ago. In both lineages, the supergene is associated with social polymorphism, where some colonies are headed by a single queen (monogyne) while others harbor multiple queens (polygyne). We discovered a new supergene associated with similar social polymorphism in the desert ant *Cataglyphis niger*. Although *Cataglyphis* is more closely related to *Formica* than to *Solenopsis*, the social chromosome in *C. niger* is homologous to the *Solenopsis* social chromosome, with conservation of synteny in gene content as well as gene order. Therefore, this is an ancient chromosome dating back to the common ancestor of *Solenopsis* and *Cataglyphis*, at least 90 million years ago. Previous work dated the *Solenopsis* supergene to be only one million years old, therefore, we propose that the two supergenes have evolved independently in the two diverged lineages on this ancient social chromosome. We surveyed hymenopteran genomes to discover a bee chromosome homologous to the ants' social chromosome. The conservation of the gene set on these homologous chromosomes in ants and bees suggests that this chromosome plays an important role in social behavior across social Hymenoptera. We hypothesize that the conserved gene set in the social chromosome was repeatedly used as a pre-adapted toolkit for the evolution of social traits in general and specifically in the evolution of polygyne social structure in ants.

**Keywords:** Supergene, social chromosome, polygyne, complex trait, Hymenoptera, social insects, population genomics

## 1. Introduction

The convergent evolution of traits is a common phenomenon across the tree of life[1]. Little is known about the genomic architectures that facilitate the repeated evolution of complex traits, but the recent accessibility of genomic approaches allows studying these processes across diverse lineages. Ants have repeatedly evolved the trait of multiple-queen colonies (polygyny) from an ancestral phenotype of a single-queen colony (monogyny)[2,3]. Polygyny is a complex trait involving a suite of life-history, behavioral, physiological, and morphological modifications. Moreover, it results in nestmates that have very low relatedness, which has consequences for the evolution of cooperation by means of kin selection[3,4]. Hughes *et al.*[2] reported that polygyne colonies were observed for 54 out of 130 (42%) ant species included in that study. Some species of ants are socially polymorphic, where both monogyne and polygyne colonies coexist in the same population. A genetic basis of this social polymorphism has been documented and examined in detail in two distantly related ant lineages, *Solenopsis* fire ants and *Formica* wood ants. In both lineages, a supergene was found to be associated with this polymorphism[5]. Supergenes are chromosomal regions with suppressed recombination, containing multiple tightly linked loci that code for a complex trait. The most well-known type of supergenes are sex chromosomes, but virtually any other complex polymorphic trait may be coded by a supergene[6]. The supergenes responsible for variation in queen number offer an opportunity to understand the genomic basis of the recurrent evolution of complex traits.

Independently evolved supergenes on non-homologous chromosomes underlie similar social polymorphism in both *Solenopsis* and *Formica*[7,8]. However, there are several important similarities and differences in how the supergene is maintained. The first social supergene was discovered by Wang and colleagues in the red fire ant *Solenopsis invicta*, which is native to tropical South America[7]. In monogyne colonies of this species, the queen and the workers are MM homozygotes at the supergene (the *Solenopsis* supergene haplotypes were originally named *SB* and *Sb*, but we are using M for monogyne-associated and P for polygyne-associated haplotypes across all species for

consistency). On the other hand, the polygyne queens are always MP heterozygous at the supergene, and produce workers of all three genotypes: MM, MP, and rarely PP, by mating with both M and P haploid males[9]. The absence of the PP genotype in queens led Wang and his colleagues to refer to this chromosome as a "Y-like social chromosome". This supergene was found across a few closely related, socially polymorphic *Solenopsis* species that speciated approximately one million years ago (MYA)[10]. The formation of the supergene was dated approximately to the same time based on the diverges of the M and P haplotype sequences[11,12]. In the Alpine wood ant *Formica selysi*, where the second social supergene was discovered, monogyne queens and workers are MM homozygotes at the supergene while the polygyne queens and workers are either PP homozygotes or MP heterozygotes. Unlike in *S. invicta*, MM genotypes in polygyne colonies are eliminated in this species, which was suggested to be due to a maternal effect killing[13]. Supergenes homologous to the *F. selysi* supergene were found in numerous *Formica* species and therefore evolved in their common ancestor at least 30 MYA[14]. *Formica* and *Solenopsis* belong to two different ant subfamilies (Formicinae and Myrmicinae) that diverged more than 90 MYA[15]. The young age of the two supergenes relative to the lineage divergence and the fact that they reside in non-homologous chromosomes implies that the two supergenes have independent origins.

The desert ant *Cataglyphis niger* is a socially polymorphic species, with populations consisting of both monogyne and polygyne colonies. These populations were initially thought to consist of multiple species based on divergent mitochondrial sequences (mitotypes) that are associated with the social form[16]. However, nuclear genomic sequencing revealed no genetic divergence between monogyne and polygyne colonies, demonstrating that *C. niger* is a socially polymorphic species[17]. The genera *Cataglyphis* and *Formica* belong to the same tribe Formicini within the Formicinae subfamily, which diverged approximately 40 MYA[15]. We examined the genomic basis of social polymorphism in *C. niger* by generating reduced representation genome sequencing data for 672 individuals from 30 nests representing both social forms from a single population. The social form of the nests was ascertained by estimating relatedness

among nestmates, as well as other lines of evidence, including aggressive behavior between colonies. We discovered a novel "Y-like" social chromosome in *C. niger* harboring a supergene that is maternally inherited. Surprisingly, we find that the *Cataglyphis* social chromosome is homologous to the *Solenopsis* social chromosome, even though the two lineages diverged more than 90 MYA. A wider survey across multiple hymenopteran genomes revealed conserved synteny of the social chromosome across bees and ants.

Extensive previous literature suggested that the evolution of sociality in diverse hymenopteran lineages involved reuse of a core set of ancestral genes, a so-called "molecular toolkit"[18–20]. Our study revealed the conservation of synteny of an ancestral gene set on an ancient chromosome that was reused for the evolution of supergenes coding for alternative social structures in diverged ant lineages. We suggest that this gene set represents such a "toolkit" for evolving sociobiological traits.

## 2. Results and discussion

### 2.1 Social polymorphism in *Cataglyphis niger* is associated with chromosome 2

We fully excavated 30 nests from a single population in Tel Baruch, Israel, where both monogyne and polygyne nests were known to co-exist. Genomic data was generated for 672 samples (28 queens and 644 workers) from the 30 nests (20-24 individuals each) using double-digest restriction-site associated DNA sequencing (RAD-seq), yielding genotype data for 30,955 high-confidence single nucleotide polymorphisms (SNPs). As queens were found only in 14 nests (7 had one queen and 7 had more than one queen), the social form of the sampled nests was assigned based on the relatedness within the nest.

A pairwise kinship matrix constructed based on the total length of shared identity-by-descent (IBD) haplotypes across 672 sampled individuals showed some nests with relatedness between all nestmates and somewhere most pairs were unrelated (Figure 1a), as expected for monogyne and polygyne nests, respectively. In monogyne nests,

we expected three clusters of relatedness scores— full siblings or super sisters sharing 75% of their genome, mother-daughter pairs with 50% relatedness, and half-sisters (from different fathers) with 25% relatedness. The nests with higher relatedness had broadly three clusters of relatedness scores consistent with our expectations in a monogyne nest (Figure 1b). However, we did find several worker pairs with 50% relatedness, which could result from the queen mating with brothers. As expected for polygyne nests, most pairs of workers had relatedness scores lower than half-sister relatedness (25%). Based on the distribution of kinship scores, 13 nests were classified as monogyne and 17 as polygyne for downstream analyses. Aggression assays also corroborated this classification of the nests: aggression scores were significantly higher in interactions between two monogyne nests and between a monogyne and a polygyne nest, compared to between two polygyne nests (Figure 1c; $t$-test $p$-value << 0.05 for all comparisons). The number of queens found during sampling was also in line with these results (full details provided in Supplementary Figures S1 & S2).

**Figure 1:** Identification of social form using kinship inference based on the total length of identity-by-descent (IBD) haplotypes shared between nestmates, **(a)** plotted as a kinship matrix demarketing monogyne nests in blue and polygyne nests in green. **(b)** Distributions of relatedness within nests. **(c)** Aggression scores from the dyadic assay showed significantly higher aggression between monogyne and monogyne-polygyne nests compared to between polygyne nests.

We examined $F_{ST}$ at each SNP between monogyne and polygyne samples (Figure 2a). The average weighted $F_{ST}$ across the genome was low (0.019) as expected due to gene flow and lack of genetic differentiation between the two social forms[17]. However, the average weighted $F_{ST}$ across chromosome 2 was much higher (0.12) than the rest of the genome. The high $F_{ST}$ (>0.2) region was a ~5.8Mb stretch extending from position 3,930,062 bp to 9,747,310 bp in the Cnig_gn3.1 genome assembly.

**Figure 2:** Genetic divergence and high linkage disequilibrium on chromosome 2 suggest the presence of a supergene. **(a)** Plot of $F_{ST}$ between social forms (defined based on kinship) showing high $F_{ST}$ values on chromosome 2 . **(b)** Heatmaps of linkage disequilibrium in monogyne (above the diagonal) and polygyne (below the diagonal) population samples (displaying only $r^2$>0.2) across all chromosomes. **(c)** Heterozygosity along chromosome 2 in 69 full sisters from a polygyne colony. **(d)** Comparison of the boundaries of the supergene region according to the different criteria.

## 2.2 Suppression of recombination in chromosome 2

Supergenes maintain complex polymorphic traits despite high gene flow between morphs by suppression of recombination that prevents the shuffling of the alleles associated with the trait at different loci. We tested for suppression of recombination

using two approaches— (1) testing for linkage disequilibrium (LD) in the population sample and (2) testing for linkage in progeny of a single polygyne queen. We conducted LD analyses separately on monogyne and polygyne samples to examine if the supergene region on chromosome 2 showed suppression of recombination. The analysis revealed high LD ($r^2 \geq 0.2$) in polygynes but not in monogynes across most SNP pairs in the supergene region (chromosome 2 positions 3,813,975 – 10,147,746 bp), aligning with the region of high $F_{ST}$ SNPs (Figure 2b). We also noted LD decaying rapidly with SNP distance across all chromosomes of monogyne and polygyne samples except in chromosome 2 in polygynes, where LD did not decay over long distances (Supplementary Figure S3). Adjacent to the large 5.7Mb LD block, a smaller 270 kb LD block can be observed in both monogyne and polygyne samples (Supplementary Figure S4). Similarly, four additional small LD blocks spanning over 249 kb were observed in other chromosomes of both the social forms (chr07: 298 kb; chr11: 249 kb; chr13: 434 kb; chr17: 297 kb) (Figure 2b).

Additionally, a linkage analysis was conducted based on RAD sequencing of 225 brood samples collected from a single polygyne queen. The mother was also sequenced. Six patrilines were identified in this data set and the largest patriline (69 full sisters) was used to build a linkage map. However, the genotype data did not allow for building a map of chromosome 2 because of an unusual pattern. All sisters had the same genotypes across the supergene region: 131 SNPs (70%) were always heterozygous and 44 SNPs (24%) always had the same homozygous genotype, out of 186 SNPs that were heterozygous in the mother (Supplementary Figure S5). Plotting the heterozygosity along chromosome 2 (Figure 2c) showed that SNPs in the supergene region were either 100% or 0% heterozygous. The extent of markers with heterozygosity of 100% spanned chromosome position 3,986,390 - 10,237,665bp. This pattern can be explained by a combination of two effects: (1) there were no recombination events within the supergene region, and (2) all the daughters received the same haplotype of the supergene from their mother. That is, recombination in the supergene was fully suppressed and one of the two maternal haplotypes is lethal. We

note that this pattern is reminiscent of maternal effect killing of homozygous MM daughters in polygyne *F. selysi* colonies.

## 2.3 Maternally inherited 'Y-like' supergene

We examined the genotypes of 164 high $F_{ST}$ (>0.2) SNPs on chromosome 2, by generating a heatmap (Figure 3a). The heatmap showed that polygyne samples were predominantly heterozygous (127/164 loci on average), whereas monogyne samples were largely homozygous for one allele (132/164 loci on average). This suggests that the polygyne social form is associated with a 'Y-like' supergene haplotype, which we denote as the P haplotype. That is, polygyne samples carry a heterozygous MP supergene genotype, while monogyne samples are MM homozygotes. This pattern holds for all queens and most of the workers. One conspicuous exception to this rule was nest M31, which was designated as a monogyne nest based on high relatedness, and yet, all samples in this nest had the MP genotype. The phenotype of high relatedness in this nest could have been observed if polygyne nests in *Cataglyphis* were also founded independently by queens and not, as commonly assumed, through nest budding. We speculate that some polygyne nests may be founded by a single MP queen, which is subsequently joined by additional MP queens. Indeed, this nest had a relatively small number of workers, as expected for a newly established nest. This is also in line with the geographic location of this nest among other polygyne nests (Figure 3c). The mitotype (discussed below) and aggression scores of this nest are also as expected for the polygyne form. The aggression scores of this nest against polygyne nests (Supplementary Figure S1) were relatively low (0.2 – 0.9) compared to its score against a monogyne nest (1.3). There were seven other exceptions to the broad pattern observed in the heat map, all of which were workers: five samples from polygyne nests had an MM genotype (four from P34 and one from P35) and two samples from monogyne nests had an MP genotype (M19-44 and M27-19). Nest details along with supergene genotypes and mitotypes are given in Supplementary Table S1.

**Figure 3: (a)** Heatmap showing the genotypes at SNPs with high $F_{ST}$ (>0.2) on chromosome 2. Generally, one allele is only found in heterozygous genotypes in polygyne colonies, whereas the other allele is homozygous in monogyne colonies. Exceptions to the rule are highlighted in nests M19, M27, M31, P34. **(b)** Mitochondrial phylogeny with a deep divergence between monogyne and polygyne samples indicating maternal inheritance of the social chromosome. **(c)** Geographic location of the nests where the social form was identified based on kinship and supergene genotype.

Previous work on this species suggested deep divergence between mitotypes associated with monogyne and polygyne social forms based on samples collected across Israel[16]. To ascertain if this pattern was consistent across the nests sampled in this study, we sequenced part of the mitochondrial gene Cytochrome *b* from two to three individuals from each nest. The mitochondrial phylogeny (Figure 3b) resulted in two deeply divergent well-supported monophyletic clades largely corresponding to monogyne and polygyne samples. Previously published mitotypes associated with monogyne nests (previously named the "savignyi" and "drusus" mitotypes) were nested within the monogyne clade and mitotypes associated with polygyne nests (the "niger" mitotype) were nested within the polygyne clade. This divergence in mitotypes

associated with social form indicates a strict, non-Mendelian, maternal inheritance of the P chromosome. Five samples were an exception to this rule. Both samples sequenced from nest M31 (a queen and a worker), which had the MP supergene genotype, also carried a polygyne-associated mitotype. Two worker samples from polygyne nests with MP supergene genotypes carried a monogyne-associated mitotype.

One conspicuous result from the heat map is the absence of both homozygous PP and MM genotypes from polygyne nests, where virtually all individuals have a heterozygous MP genotype. The absence of MM genotypes suggests negative selection, possibly a result of mechanisms such as meiotic drive, maternal effect killer, or green beard effect, which were suggested for the social supergenes in other species. Maternal effect killing was proposed to eliminate MM genotypes in polygyne colonies of *F. selysi*[13], whereby the presence of the P chromosome in the mother results in the death of progeny that did not inherit at least one P chromosome. Another possibility is a green beard effect, whereby workers carrying the P chromosome would kill brood that do not, similar to the killing of MM queens in polygyne colonies of *S. invicta*[21]. The absence of PP genotypes also indicates negative selection, potentially due to deleterious mutations, which may be similar to the low fitness of PP queens in *S. invicta*. Such deleterious mutations may also result in P being lethal in males (although this is not the case in *S. invicta*). If so, MP daughters would always inherit their P chromosome from their mother (and M from their father), in line with the mitochondrial phylogeny. Another possible explanation for the maternal inheritance of the P chromosome is that P males do exist, but monogyne queens do not mate with them.

## 2.4 The supergene is distributed across multiple populations in Israel

We used a previously published RAD-seq dataset of worker samples collected across Israel to verify that the supergene is distributed beyond our study population of Tel Baruch[17]. This study used different enzymes from our study to generate genomic libraries resulting in a different set of genomic loci. We used the mitotype of the samples as a proxy to identify their social form and calculated $F_{ST}$ between inferred monogyne and polygyne samples at each SNP across the genome. High $F_{ST}$ SNPs were detected

on chromosome 2 between positions 3,985,614-9,804,010, which overlapped the supergene region identified in the Tel Baruch population (Supplementary Figures S6a and S6b). Similar to the pattern reported above (Figure 3a), the 164 high $F_{ST}$ SNPs (>0.4) in this dataset were mostly homozygous MM (52/56) in the monogyne mitotype samples and heterozygous MP (13/14) in the polygyne mitotype samples (Supplementary Figure S6c). This sampling across Israel shows that most of the range consists of pure MM populations, whereas the P haplotypes are always found in polymorphic populations (MM and MP) limited to the central coastline region between Ashkelon and Netanya (Supplementary Figure S6d).

## 2.5 Repeated evolution of social supergenes on an ancient conserved chromosome

Non-homologous social chromosomes are known to be associated with social polymorphism in *Solenopsis* and *Formica*. We wanted to examine whether the *Cataglyphis* social chromosome is homologous to either of these chromosomes. *Cataglyphis* and *Formica* both belong to the subfamily Formicinae, whereas *Solenopsis* belongs to the subfamily Myrmicinae. The two subfamilies diverged around 93 MYA[15] (95% confidence interval 91-95 MYA). Synteny dot plots based on one-to-one orthologs showed that the social chromosomes of *Cataglyphis* and *Formica* were not homologous (Figure 4a). Surprisingly, despite belonging to different subfamilies, the social chromosomes of *Cataglyphis* and *Solenopsis* are homologous (Figure 4b). Synteny is not only conserved in terms of gene content (macrosynteny) but also gene order (microsynteny), especially in the supergene region (red points in Figure 4c). Interestingly, the largest region with conserved gene order is inside both supergenes, while the non-overlapping regions of the supergenes show conservation of gene order to a lesser extent.

**Figure 4:** Conserved synteny of the social chromosome across ants and bees. Dot plots of one-to-one orthologs **(a)** between *Cataglyphis* and *Solenopsis* and **(b)** between *Cataglyphis* and *Formica*. Orange arrows indicate the social chromosome in *Solenopsis* and *Formica*, and the orange box indicates the social chromosome in *Cataglyphis*. **(c)** Social chromosomes with the supergene regions of *Cataglyphis* in blue and *Solenopsis* in green. Supergene coordinates in *Cataglyphis* are defined based on the high $F_{ST}$ region (3,930,062 to 9,747,310bp), and in *Solenopsis* based on the inversion breakpoints (Yan *et al.*[12]). The overlapping supergene region highlighted in red shows remarkable conservation in gene order. The yellow triangles indicate odorant receptors (ORs) and red stars indicate odorant-binding proteins (OBPs). **(d)** *Cataglyphis* and *Apis* dot plot. *Cataglyphis* chromosome 2 shows notable conservation of gene content and homology to *Apis* chromosome 15.

(a)

(b)

**Figure 5:** Conservation of synteny of all *Cataglyphis* chromosomes. **(a)** Percent of one-to-one orthologous genes in each *Cataglyphis* chromosome that reside in one chromosome in other Hymenoptera species. **(b)** Synteny plot showing three representative species of ants, two Apoidae species, and one wasp outgroup. The figure highlights three broad regions of *Cataglyphis* chromosome 2, the supergene region (red), the region upstream (green), and the region downstream (blue) of the supergene. The blue box marks a cluster of genes that appears to have been convergently fused to the social chromosome in both *Cataglyphis* and *Solenopsis* lineages (see Supplementary Figure S7).

To investigate the extent of the conservation of synteny of the social chromosome, we conducted a comparative genomic analysis of all published annotated, highly contiguous ant assemblies. We also added representatives of bees and wasps to this analysis amounting to 22 genomes that allowed us to evaluate the conservation of synteny across Hymenoptera. As an estimate of macrosynteny, we calculated the percentage of *Cataglyphis* genes found on the homologous chromosome (the chromosome carrying the largest number of homologous genes) in each of these species (one-to-one orthologs only). Figure 5a shows that *Cataglyphis* chromosome 2 is one of the more highly conserved chromosomes across Hymenoptera, especially relative to other large chromosomes (chromosomes are numbered from largest to smallest). This chromosome is notably more conserved in distant lineages such as *Solenopsis* (75% of one-to-one orthologs on the same chromosome) and even in the honey bee *Apis mellifera* (41%).

Figure 5b shows the conservation of microsynteny of genes on the social chromosome in three ants, two Apoidea species (the honey bee and a digger wasp), and the jewel wasp *Nasonia* as an outgroup (Supplementary Figure S7 shows all 22 Hymenoptera). Microsynteny in the supergene and the region upstream to the supergene in *Cataglyphis* appear to be conserved to a certain extent across the available genomes from Myrmicinae and Formicinae. In the clonal raider ant *Ooceraea biroi,* however, this region is broken up and represented on five chromosomes. This is likely to have happened after the lineage leading to *O. biroi* diverged from the lineage leading to Myrmicinae and Formicinae. It remains to be determined whether this is common to other members of the subfamily Dorylinae or specific to the clonal raider ant. The downstream region of the *Cataglyphis* chromosome is broken in many formicine species (e.g. *Formica*) and myrmicine species (e.g. *Pogonomyrmex*), and also in bees. Either the downstream region was part of the ancestral chromosome and broke off in multiple mymricine and formicine lineages, or it was not part of the ancestral chromosome, and the same synteny group of genes was fused to the social chromosome in both *Cataglyphis* and *Solenopsis* lineages (marked in Figure 5b; see also Supplementary Figure S7).

Most genes in the upstream region of the *Cataglyphis* social chromosome and the supergene region (red and green lines in Figure 5b) are found on chromosome 15 of *Apis*, followed by chromosomes 1, 7, 8, and 14. This conservation of macrosynteny is evident across all the six representatives of the superfamily Apoidea, which includes the bees and the crabronid wasps (Supplementary Figure S7). These results indicate deep conservation of the social chromosome, dating back to the divergence of Apoidea and Formicidae (ants) around 175 million years ago[15]. We propose that conserved gene content in this region is an ancestral genetic toolkit that has been repeatedly used in the evolution of sociobiological traits in hymenopteran lineages.

The association of social polymorphism with a supergene on the social chromosome in both *Cataglyphis* and *Solenopsis* can be interpreted in two ways. We may propose that the supergene was present in the common ancestor and conserved in both lineages. However, based on the divergence between the two haplotypes of the *Solenopsis* supergene, it was estimated to be no more than one MYA[7,11]. While we do not yet have information on the age of the supergene in *Cataglyphis niger*, there were no polygyne colonies reported for the most closely related *Cataglyphis* species (bicolor group). Therefore, we propose that the two lineages evolved analogous supergenes on the same chromosome independently. While this may seem surprising, studies on sex chromosome turnover show that certain chromosomes repeatedly transition to being sex chromosomes, possibly because of the genes they carry[22,23]. Similarly, if the social chromosome contains an ancient genetic toolkit for sociobiological traits, this would explain the repeated evolution of supergenes on this chromosome. Examining the genomic basis of multiple socially polymorphic species in other ant lineages could help shed light on this question.

## 2.5 Gene content

We examined the gene content of the *Cataglyphis* and *Solenopsis* social chromosomes to find 786 and 1088 putative protein-coding genes, respectively. The two supergene regions contain 248 and 441 putative protein-coding genes in *Cataglyphis* and *Solenopsis*, respectively. Next, we identified orthologous genes between the two

chromosomes. In the *Solenopsis* supergene region, there are 207 functionally-annotated protein-coding genes with one-to-one orthologs in *Cataglyphis*, and 112 such genes in the *Cataglyphis* supergene, with 84 of these being found in both the supergene regions. All but one of the one-to-one orthologs of genes in the *Cataglyphis* supergene are in the *Solenopsis* social chromosome, and one gene is in an unassembled scaffold (Supplementary Table S2), demonstrating the exceptional degree of gene content conservation of this chromosomal region. Other genes have many-to-many orthology relationships between the two species (Supplementary Table S3; genes with functional annotation). The *Cataglyphis* social chromosome contains 492 such genes in total, of which 149 are in the supergene region. The *Solenopsis* social chromosome contains 665 such genes, 279 of which are in the supergene region.

The homologous supergene regions contain a large and diverse set of molecular functions. It is difficult to identify specific candidate genes without additional functional information. Nevertheless, we reviewed the list and looked for gene functions that may be implicated in the sociobiology of ants (Supplementary Tables S2 and S3 – one-to-one and many-to-many; Supplementary Figure S8 – dot plot with gene annotations). One notable example is a cluster of three cytochrome P450 4C1 genes in the *Cataglyphis* supergene, which are orthologous to nine genes in the *Solenopsis* supergene. Cytochrome P450 genes are a large and diverse family of enzymes, which was noted already when the first ant genomes were sequenced. The genome of the harvester ant *Pogonomyrmex barbatus* was found to contain 72 cytochrome P450 genes, which is a significant expansion relative to other insect species[24]. Enzymes of this gene family participate in a wide range of metabolic processes[25], including synthesis and degradation of hormones such as juvenile hormone, a major regulator of insect development that plays a key role in caste determination in social insects[26–29].

The total number of functionally annotated one-to-one orthologs shared between the *Cataglyphis* and *Solenopsis* social chromosomes and *Apis mellifera* (anywhere in the bee genome) is 305. Of these, 164 (54%) are located on *Apis* chromosome 15 (Supplementary Table S4), and 34 out of the 164 are located in the supergenes of both

*Solenopsis* and *Cataglyphis*. These 34 genes include dopamine receptor 2 (*Cataglypis* gene_1588 / *Solenopsis* XP_039314227.1 / *Apis* XP_026301043.1), odorant binding protein-9 (*Cataglyphis* gene_1588 / *Solenopsis* XP_039314227.1 / *Apis* XP_026301043.1), and neuronal acetylcholine receptor subunit alpha-10 (*Cataglyphis* gene_2000 / *Solenopsis* XP_025994007.1 / *Apis* XP_392070.3). Dopamine is implicated in modulating social and reproductive behavior in Hymenoptera[30–34].

We also took note of the downstream region of the *Cataglyphis* chromosome, which is homologous to the upstream region (short arm) of the *Solenopsis* chromosome, because this region may have been convergently fused to the chromosome in both lineages (Figure 5b; Supplementary Figure S7). This region contains 121 functionally annotated one-to-one orthologous genes, including a steroid hormone receptor (*Cataglypis* gene_1712 / *Solenopsis* XP_039315095.1), an ecdysone 20-monooxygenase (*Cataglyphis* gene_2123 / *Solenopsis* XP_011172207.1), and an ecdysteroid-regulated 16 kDa protein (*Cataglyphis* gene_2111 / *Solenopsis* XP_011170763.1). Genes implicated in such functions may have significance for the sociobiology of ants, as different levels of ecdysteroids and ecdysone receptors were found in the brains of foragers versus nurses in *Camponotus* and *Harpegnathos*[35,36].

We were particularly interested in genes implicated in olfactory functions, which were extensively studied in *Solenopsis* because of their potential role in chemical signaling and the regulation of social structure. Previous studies reported differences between monogyne and polygyne queens in cuticular hydrocarbons and other compounds that were suggested to function as queen pheromones involved in the regulation of social form and the green beard effect[21,37,38]. Other studies focused on odorant binding proteins (OBPs) and odorant receptors (ORs) in the *Solenopsis* supergene, which may be involved in olfactory response to such chemical signals[39–42]. The *Solenopsis* supergene contains a large array of 23 ORs and 3 more separate ORs[42]. We annotated ORs in the *Cataglyphis niger* genome and constructed a gene tree including ORs from *C. niger, Solenopsis invicta, Camponotus floridanus,* and *Linepithema humile* (Supplementary Figure S8). The 23 *Solenopsis* ORs and their orthologs in *Cataglyphis*

belong to a single subfamily of the OR gene superfamily. The gene tree shows that the 23 *Solenopsis* ORs have one-to-one or many-to-many orthology relationships with ORs in two regions of the *Cataglyphis* social chromosome, but outside of the supergene (marked by red stars on the dot plot in Figure 4c). Three more ORs in the *Solenopsis* supergene (belonging to a different OR subfamily) are orthologs of two *Cataglyphis* ORs, which also reside outside of the supergene. Given that the ORs are a fast-evolving family, there is no clear orthology relationship between ant and bee ORs.

The *Solenopsis* supergene also contains 9 out of a total of 23 OBPs in the genome[41]. One of the OBPs is *Gp-9*, the first gene discovered in the *Solenopsis* supergene in the 1990s and the one protein that shows the largest number of amino acid substitutions between the M and the P chromosomes, indicating positive selection[39–41]. Interestingly, the homologous chromosome in the honey bee, chromosome 15, also contains 9 out of 21 OBPs, but only one of these (AmelOBP9) is orthologous to one of the supergene OBPs (SiOBP9). The 9 OBPs in the *Solenopsis* supergene are orthologous to 3 OBPs in the *Cataglyphis* supergene (Fig. 4c). Cnig_gene_1607 is orthologous to 7 of the *Solenopsis* OBPs; Cnig_gene_1608 is the ortholog of the *Solenopsis Gp-9* gene (SiOBP3); and Cnig_gene_1586 is orthologous to SiOBP9 and AmelOBP9 (Fig. 6). We note that the OBP cluster containing *Gp-9* is in the largest block of conserved microsynteny between *Solenopsis* and *Cataglyphis*. Altogether, these gene families implicated in olfaction are promising candidates for the toolkit that may have been reused for social evolution in multiple hymenopteran lineages.

**Figure 6:** Gene tree of the odorant binding protein (OBP) gene family in the ants *Cataglyphis niger* and *Solenopsis invicta* and the honey bee *Apis mellifera*. OBPs marked by rectangles reside on the social chromosomes (*Cataglyphis* chromosome 2, *Solenopsis* chromosome 16, *Apis* chromosome 15).

## 3. Conclusion

The complex trait of multi-queen (polygyne) colonies has evolved repeatedly from single-queen (monogyne) colonies in ants[2,3]. Multiple ant species display polymorphism in social form, providing a unique opportunity to investigate the evolutionary genomic basis of this trait. We investigated the genomic architecture associated with social form in a polymorphic population of *Cataglyphis niger*. Our data show that social polymorphism is associated with a large supergene— a region of suppressed recombination across at least 5.8Mb of chromosome 2. This supergene is virtually always heterozygous (MP) in polygyne individuals and homozygous (MM) in monogynes. Surprisingly, this supergene is homologous to the supergene of the fire ant *Solenpsis invicta*, even though these two lineages diverged more than 90 million years ago. These species are representatives of the two largest ant subfamilies Formicinae and Myrmicinae. Most of *Cataglyphis* chromosome 2 is homologous to most of *Solenopsis* chromosome 16, with a substantial overlap of the supergene regions where the gene order is highly conserved. The specific structure of this set of genes suggests that this chromosome has conserved functions related to social colony structure and the sociobiology of ants, which is likely to be shared across many ant species. Conservation of synteny in gene content of the social chromosome is also observed in representatives of Apoidea, including the honey bee *Apis mellifera*. The observation of conserved synteny dating back to around 175 million years suggests that the genomic architecture of this chromosome has adaptive significance. The conserved gene set could have served as a toolkit for evolving social traits across bee and ant lineages. Specifically in ants, this genetic toolkit might have facilitated the repeated evolution of polygyne social structure. This discovery opens the way for both evolutionary and mechanistic studies of the genetic basis and genomic architecture of complex phenotypes related to insect sociality.

# 4. Methods

## 4.1 Sample collection

Thirty nests of *Cataglyphis niger* were collected from a single population in Tel Baruch beach, Israel, between November and December 2019. This location is known to harbor both monogyne and polygyne colonies based on Reiner Brodetzki *et al.*[17]. We targeted at least two monogyne and two polygyne nests in every sampling trip based on their geographical distribution in that previous study. This also allowed us to conduct aggression assays (see below) between monogyne and polygyne nests after every batch of sampling. A distance of at least 100m was maintained between two sampled nests as these ants are known to be polydomous. An attempt was made to collect the queens by excavating the entire nest. Nest samples were carried back to the lab (TBA14 - 44) to perform behavioral experiments and extract genomic DNA. An additional polygyne nest (TBA67) with several queens was collected in January 2021 and was used for building a linkage map. The queens from this nest were isolated and 225 samples from the brood of a single queen were collected for RAD sequencing.

## 4.2 Molecular work

Samples not used for the behavioral assay were stored at -80C immediately after sampling. 22-24 workers from each nest in addition to the queens amounting to 714 samples were used for the molecular work from sampling carried out in 2019, henceforth referred to as the SC (social chromosome) dataset. An additional 225 brood samples from a single polygyne mother were used to construct a linkage map (linkage map dataset). DNA was extracted from the workers' abdomen and the queen's thorax using a liquid handling robot (Tecan Freedom EVO 150 NGS) and DNeasy (QIAGEN) protocol. The extracts were cleaned using ethanol precipitation and stored at -20C. A partial fragment of 450bp of Cytochrome b (*cyt b*) was PCR amplified and Sanger sequenced for 2-3 workers and queens from each nest. Double-digest Restriction-site Associated DNA sequencing (RAD-seq) was used to generate genomic libraries following Brelsford *et al.*[43]. Briefly, the genomic DNA was cut using restriction enzymes EcoRI and BfaI. Adaptors with individual barcodes were ligated to each sample for multiplexing. The ligated products were PCR amplified using barcoded primers and

pooled. Size selection of DNA fragments between 330–1000bp was carried out using a Bluepippin instrument (Sage Science). A detailed protocol can be found in Lajmi *et al.*[44]. Libraries were sequenced on an Illumina HiSeq X machine with paired-end 150bp reads.

## 4.3 Molecular Analysis

### 4.3.1 Mitochondrial phylogeny

The Sanger sequences were aligned in MEGA X[45] using the Muscle algorithm[46]. To assign our samples to previously described mitotype clades, published *cyt b* sequences of monogyne and polygyne colonies were also included from Reiner Brodetzki *et al.*[17]. Mitotypes of *Cataglyphis israelensis* were used as an outgroup. A maximum likelihood tree was constructed using raxmlGUI 2.0[47] using a GTR+G model of sequence evolution and 1000 bootstraps.

### 4.3.2 RAD-seq pipeline

The bioinformatics pipeline was run separately for the SC data and linkage map, as well as the two datasets combined. The raw sequencing data were examined using FastQC[48] for quality. Adaptors and low-quality reads were then trimmed/filtered using Trimmomatic[49]. Two mismatches were allowed in identifying the adaptors, with eight being the minimum length for identifying an adaptor, a sliding window size of 4bp with an average required quality of 15 was used. FLASH[50] was used to combine overlapping paired reads. The data were then demultiplexed using the command process_radtags.pl from the Stacks2 package[51]. The reads were then mapped to the *Cataglyphis niger* genome (assembly version Cnig_gn3.1; NCBI accession number: xxxxxxxxx [52] using Bowtie2[53] with the --end-to-end and --very-sensitive options. SNP calling was conducted using Ref_map.pl from Stacks2 to obtain a VCF file. Samples with more than 50% missing data were dropped from the analyses. The SNP calling was done separately for the SC and linkage map datasets, as well as the combined dataset. A blacklist of loci with potential artifacts was created based on the heterozygous genotype calls in three or more samples out of 64 haploid males. These loci were excluded from the VCF. Such

artifacts may be the result, for example, of errors in the assembly of the reference genome, where repetitive sequences are collapsed into a single copy.

Along with removing the blacklisted loci, the following loci were removed from the VCF file for the SC dataset using VCFtools version 0.1.14[54]: i) genotype calls based on less than 12 reads; ii) loci with more than 30% missing data; iii) loci with a minor allele frequency of less than 0.4% or 3 allele copies; iv) loci with a mean depth greater than 175 reads per individual. The final VCF file for the SC dataset consisted of 672 individuals comprising 28 queens and 644 workers. This resulted in a catalog of 30,955 high-confidence SNPs that were used for most of our analyses. The linkage map data VCF file varied slightly in the filtration in that we required only 8 reads for a genotype call, a maximum mean depth of 150 reads per individual (because of lower average depth), and a minor allele frequency of 1% or 3 allele copies (because of the smaller number of samples). The final VCF file for the linkage map consisted of 225 offspring and their mother and amounted to 17,984 high-confidence SNPs.

## 4.4 Identification of the social form

*Cataglyphis niger* is a polydomous species, i.e., a colony may have multiple nests. This made finding the queens of each colony difficult. Therefore, we relied on kinship estimates among worker samples to identify the colony's social form. Monogyne nests were expected to show high kinship between nestmates as they are expected to be full or half-siblings, compared to polygyne nests where workers are not necessarily related. We also conducted behavioral assays, as additional supporting evidence for the colony's social structure.

### 4.4.1 Kinship

Kinship inference was carried out for the two genomic datasets separately. For the SC dataset, kinship was used to identify the social form of the nests. Whereas in the linkage map dataset, kinship was used to identify the patrilines. The RAD-seq SNP catalog was imputed and phased using Beagle version 5.0[55] and refined-IBD[56] was used to identify shared identity-by-descent (IBD) haplotypes between samples. The kinship between a pair of samples was estimated based on the combined lengths of their shared IBD

haplotypes, resulting in a kinship matrix. To obtain an estimate of the relatedness coefficient we divided the total IBD length by twice the total genome assembly length (because the samples are diploid). A total of 7,651,126 and 6,238,277 IBD haplotypes were identified in the SC dataset (among the 672 samples) and the linkage map dataset (among the 225 samples), respectively. Social form was assigned to each nest based on the distribution of kinship scores. Nests were classified as polygyne if more than 75% of pairs of nestmates had less than 20% relatedness, and as monogyne otherwise.

### 4.4.2 Behavioral experiment

Behavioral assays were conducted within a week of collection from the field. These assays were based on the assumption that workers from different colonies would show high aggression towards each other. Therefore, workers from two different monogyne nests were expected to show higher aggression than workers from two polygyne nests, as we expect all polygyne ants to be part of a supercolony[17]. Dyadic aggression assays were performed, where two ants from different nests (test) or the same nest (control) were introduced into a round 12 cm diameter arena and their behavior was video recorded for three minutes. Behavior aggression scores were coded as antenation=0, mandibular threat=1, short bite with jump=2, bite=3, spraying=4 and the total aggression score was calculated as

$$\sum_{i=1}^{n} \frac{A_i * t_i}{T}$$

where $A_i$ and $t_i$ are the aggression score and duration of each behavior over the total time $T$, respectively. Ten trials were conducted for each pair of nests that was tested and an average score was calculated. A total of 1029 assays were carried out to get 104 averaged scores between pairs of nests. A two-sample *t*-test was conducted to compare the aggression scores between nestmates, polygyne-polygyne nests, and monogyne/polygyne-monogyne nest. The test was performed in R using command t.test.

### 4.5 Identifying genomic regions associated with social form

Weir and Cockerham (1984)[57] $F_{ST}$ was calculated for every SNP using VCFtools version 0.1.14[54] between 282 monogyne samples and 388 polygyne samples, where the social

form was defined based on the distribution of kinship score within the nest. Linkage disequilibrium (LD; $r^2$) analysis was performed among SNPs in each chromosome for monogyne and polygyne samples separately, using Plink v1.9[58]. Only SNPs exhibiting a minor allele frequency surpassing 0.05 for each social form were included in this analysis.

### 4.5.1 Linkage analysis in the progeny of a polygyne queen

To test for recombination in the supergene in polygyne colonies, we isolated a single polygyne queen from nest TBA67 with workers (TBA67c). 269 samples consisting of larvae and pupae were collected in a span of 60 days after the queen was isolated. RAD-seq libraries were constructed for the queen and her offspring using the same protocol as above. Patrilines were identified using kinship analysis (as above) and a linkage map was constructed for the patriline with the largest number of samples (69 samples). A linkage map was constructed based on 3,149 bi-allelic loci where the minor allele was present in at least two individuals from the same patriline. However, this analysis failed in the supergene region of chromosome 2 because all 69 samples had the same genotype in every SNP. Therefore, it was not possible to estimate the frequency of recombination and test for linkage between neighboring SNPs. To illustrate this unusual genetic pattern we plotted the heterozygosity in the 69 progeny along the chromosome for all SNPs that were heterozygous in the mother queen.

### 4.6 Synteny analysis

Synteny analysis was carried out to evaluate the conservation of synteny in the chromosome carrying the supergene across 22 hymenopteran species with the oldest divergence dating to ~230 Million years ago. The analysis included the ant subfamilies Formicinae (*Cataglyphis niger*, *C. hispanica*, a *Formica aquilonia x polyctena* hybrid*, Camponotus floridanus,* and *Lasius platythorax*), Myrmicinae (*Monomorium pharaonis*, *Solenopsis invicta*, *Pogonomyrmex californicus,* and *Trachymyrmex septentrionalis*) and Dorylinae (*Ooceraea biroi*). Six species belonging to the Apoidea superfamily, the sister clade to ants, were also included in this analysis. This included bee families Apidae (*Apis mellifera*) and Colletidae (*Colletes gigas*), a solitary bee belonging to Megachilidae (*Osmia bicornis*), and solitary wasps belonging to Crabronidae (*Pemphredon lugubris*,

*Nysson spinosus*, and *Ectemnius continuus*). Six additional wasp lineages were included— families Vespidae (*Vespula vulgaris* and *ancistrocerus nigricornis*), Chalcidoidea (*Nasonia vitripennis*), Ichneumonidae (*Venturia canescens*), Ichneumonoidea (*Mocroplitis mediator*), and Pompilidae (*Anoplius nigerrimus)*. The genomes for this analysis were chosen based on their availability on NCBI, a scaffold N50 of more than 1MB, and gene annotation for the genome assembly. OMA v2.6.0[59] was used to infer orthogroups and transfer functional annotations from *Solenopsis invicta* orthologs to *Cataglyphis niger* genes.

## 4.7 OR and OBP phylogenetic analysis

Odorant receptors (ORs) were annotated using HAPpy-ABCENTH[60], based on OR gene models from 5 manually annotated ant species. The first part of the program locates putative genetic loci through homology, and the second part, ABCENTH, is a gene finder specifically built for multigene families like ORs, with high sequence divergence but a very conserved exon structure. OR protein sequences were compiled from *Cataglyphis niger, Solenopsis invicta*, *Linepithema humile,* and *Camponotus floridanus.* Odorant Binding Protein (OBP) sequences for *Apis mellifera*[61] and *Solenopsis invicta*[41] were downloaded from NCBI and *Cataglyphis* annotations were used from Inbar *et al.*[52] and analyzed separately. For each of these datasets, MAFFT[62] was used to build a multiple sequence alignment. OR and OBP gene trees were reconstructed using RAxML v8.2.12[63], with the PROTCALG model, the LG rate matrix, and 100 bootstraps repeats.

# 5. Acknowledgments

We thank Abraham Hefetz for introducing us to the polymorphic study population and for background knowledge on this species and its social forms. We are grateful to Jessica Purcell and Alan Brelsford for their advice on RAD sequencing and the study of supergenes. EP would like to thank Laurent Keller for inspiring and motivating the study of social supergenes. AL would like to thank Praveen Karanth for his support during the analysis and writing of this paper. AL was supported by the "Study in Israel" Fellowship provided by the Planning & Budgeting Committee (PBC) of the Council for Higher

Education (CHE). This study was funded by Israel Science Foundation grants 646/15 and 1845/22.

# 6. Citations

1. Blount, Z. D., Lenski, R. E. & Losos, J. B. Contingency and determinism in evolution: Replaying life's tape. *Science* **362**, eaam5979 (2018).
2. Hughes, W. O. H., Ratnieks, F. L. W. & Oldroyd, B. P. Multiple paternity or multiple queens: two routes to greater intracolonial genetic diversity in the eusocial Hymenoptera. *J of Evolutionary Biology* **21**, 1090–1095 (2008).
3. Hughes, W. O. H., Oldroyd, B. P., Beekman, M. & Ratnieks, F. L. W. Ancestral Monogamy Shows Kin Selection Is Key to the Evolution of Eusociality. *Science* **320**, 1213–1216 (2008).
4. Keller, L. Social life: the paradox of multiple-queen colonies. *Trends in Ecology & Evolution* **10**, 355–360 (1995).
5. Kay, T., Helleu, Q. & Keller, L. Iterative evolution of supergene-based social polymorphism in ants. *Philosophical Transactions of the Royal Society B: Biological Sciences* **377**, 20210196 (2022).
6. Schwander, T., Libbrecht, R. & Keller, L. Supergenes and Complex Phenotypes. *Current Biology* **24**, R288–R294 (2014).
7. Wang, J. *et al.* A Y-like social chromosome causes alternative colony organization in fire ants. *Nature* **493**, 664–668 (2013).
8. Purcell, J., Brelsford, A., Wurm, Y., Perrin, N. & Chapuisat, M. Convergent Genetic Architecture Underlies Social Organization in Ants. *Current Biology* **24**, 2728–2732 (2014).
9. Ross, K. G. Multilocus Evolution in Fire Ants: Effects of Selection, Gene Flow and Recombination. *Genetics* **145**, 961–974 (1997).
10. Cohen, P. & Privman, E. Speciation and hybridization in invasive fire ants. *BMC Evol Biol* **19**, 111 (2019).
11. Cohen, P. & Privman, E. The social supergene dates back to the speciation time of two Solenopsis fire ant species. *Sci Rep* **10**, 11538 (2020).
12. Yan, Z. *et al.* Evolution of a supergene that regulates a trans-species social polymorphism. *Nat Ecol Evol* **4**, 240–249 (2020).
13. Avril, A., Purcell, J., Béniguel, S. & Chapuisat, M. Maternal effect killing by a supergene controlling ant social organization. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 17130–17134 (2020).
14. Brelsford, A. *et al.* An Ancient and Eroded Social Supergene Is Widespread across Formica Ants. *Current Biology* **30**, 304-311.e4 (2020).

15. Moreau, C. S. & Bell, C. D. Testing the museum versus cradle tropical biological diversity hypothesis: Phylogeny, diversification, and ancestral biogeographic range evolution of the ants. *Evolution* **67**, 2240–2257 (2013).

16. Eyer, P. A., Seltzer, R., Reiner-Brodetzki, T. & Hefetz, A. An integrative approach to untangling species delimitation in the Cataglyphis bicolor desert ant complex in Israel. *Molecular Phylogenetics and Evolution* **115**, 128–139 (2017).

17. Reiner Brodetzki, T. *et al.* The Interplay between Incipient Species and Social Polymorphism in the Desert Ant Cataglyphis. *Sci Rep* **9**, 9495 (2019).

18. Toth, A. L. & Robinson, G. E. Evo-devo and the evolution of social behavior. *Trends in Genetics* **23**, 334–341 (2007).

19. Rittschof, C. C. & Robinson, G. E. Behavioral Genetic Toolkits. in *Current Topics in Developmental Biology* vol. 119 157–204 (Elsevier, 2016).

20. Rehan, S. M. & Toth, A. L. Climbing the social ladder: the molecular evolution of sociality. *Trends in Ecology & Evolution* **30**, 426–433 (2015).

21. Keller, L. & Ross, K. G. Selfish genes: a green beard in the red fire ant. *Nature* **394**, 573–575 (1998).

22. Marshall Graves, J. A. & Peichel, C. L. Are homologies in vertebrate sex determination due to shared ancestry or to limited options? *Genome Biol* **11**, 205 (2010).

23. Jeffries, D. L. *et al.* A rapid rate of sex-chromosome turnover and non-random transitions in true frogs. *Nat Commun* **9**, 4088 (2018).

24. Smith, C. R. *et al.* Draft genome of the red harvester ant *Pogonomyrmex barbatus*. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 5667–5672 (2011).

25. Feyereisen, R. INSECT P450 ENZYMES. *Annu. Rev. Entomol.* **44**, 507–533 (1999).

26. Robinson, G. E. & Vargo, E. L. Juvenile hormone in adult eusocial hymenoptera: Gonadotropin and behavioral pacemaker. *Arch. Insect Biochem. Physiol.* **35**, 559–583 (1997).

27. Hartfelder, K. Insect juvenile hormone: from 'status quo' to high society. *Braz J Med Biol Res* **33**, 157–177 (2000).

28. Lillico-Ouachour, A. & Abouheif, E. Regulation, development, and evolution of caste ratios in the hyperdiverse ant genus Pheidole. *Current Opinion in Insect Science* **19**, 43–51 (2017).

29. Nijhout, H. F. & Wheeler, D. E. Juvenile Hormone and the Physiological Basis of Insect Polymorphisms. *The Quarterly Review of Biology* **57**, 109–133 (1982).

30. Fletcher, D. J. C. & Blum, M. S. Pheromonal Control of Dealation and Oogenesis in Virgin Queen Fire Ants. *Science* **212**, 73–75 (1981).

31. Harris, J. W. & Woodring, J. Elevated brain dopamine levels associated with ovary development in queenless worker honey bees (Apis mellifera L.). *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology* **111**, 271–279 (1995).

32. Sasaki, K. & Nagao, T. Distribution and levels of dopamine and its metabolites in brains of reproductive workers in honeybees. *Journal of Insect Physiology* **47**, 1205–1216 (2001).

33. Boulay, R., Hooper-Bui, L. M. & Woodring, J. Oviposition and oogenesis in virgin fire ant females *Solenopsis invicta* are associated with a high level of dopamine in the brain. *Physiological Entomology* **26**, 294–299 (2001).

34. Beggs, K. T. *et al.* Queen pheromone modulates brain dopamine function in worker honey bees. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 2460–2464 (2007).

35. Nemoto, M. & Hara, K. Ecdysone receptor expression in developing and adult mushroom bodies of the ant Camponotus japonicus. *Dev Genes Evol* **217**, 619–627 (2007).

36. Penick, C. A., Liebig, J. & Brent, C. S. Reproduction, dominance, and caste: endocrine profiles of queens and workers of the ant Harpegnathos saltator. *J Comp Physiol A* **197**, 1063–1071 (2011).

37. Eliyahu, D., Ross, K. G., Haight, K. L., Keller, L. & Liebig, J. Venom Alkaloid and Cuticular Hydrocarbon Profiles Are Associated with Social Organization, Queen Fertility Status, and Queen Genotype in the Fire Ant Solenopsis invicta. *J Chem Ecol* **37**, 1242–1254 (2011).

38. Zeng, H., Millar, J. G., Chen, L., Keller, L. & Ross, K. G. Characterization of Queen Supergene Pheromone in the Red Imported Fire Ant Using Worker Discrimination Assays. *J Chem Ecol* **48**, 109–120 (2022).

39. Krieger, M. J. B. & Ross, K. G. Identification of a Major Gene Regulating Complex Social Behavior. *Science* **295**, 328–332 (2002).

40. Gotzek, D., Robertson, H. M., Wurm, Y. & Shoemaker, D. Odorant Binding Proteins of the Red Imported Fire Ant, Solenopsis invicta: An Example of the Problems Facing the Analysis of Widely Divergent Proteins. *PLoS ONE* **6**, e16289 (2011).

41. Pracana, R. *et al.* Fire ant social chromosomes: Differences in number, sequence and expression of odorant binding proteins. *Evolution Letters* **1**, 199–210 (2017).

42. Cohanim, A. B., Amsalem, E., Saad, R., Shoemaker, D. & Privman, E. Evolution of olfactory functions on the fire ant social chromosome. *Genome Biology and Evolution* (2018) doi:10.1093/gbe/evy204.

43. Brelsford, A., Dufresnes, C. & Perrin, N. High-density sex-specific linkage maps of a European tree frog (Hyla arborea) identify the sex chromosome without information on offspring sex. *Heredity* **116**, 177–181 (2016).

44. Lajmi, A., Glinka, F. & Privman, E. Optimizing ddRAD sequencing for population genomic studies with ddgRADer. *Molecular Ecology Resources* 1755–0998.13870 (2023) doi:10.1111/1755-0998.13870.

45. Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and Evolution* **35**, 1547–1549 (2018).

46. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research* **32**, 1792–1797 (2004).

47. Edler, D., Klein, J., Antonelli, A. & Silvestro, D. raxmlGUI 2.0: A graphical interface and toolkit for phylogenetic analyses using RAxML. *Methods Ecol Evol* **12**, 373–377 (2021).

48. Andrews, S. FastQC: a quality control tool for high throughput sequence data. (2010).

49. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

50. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* **27**, 2957–2963 (2011).

51. Rochette, N. C., Rivera-Colón, A. G. & Catchen, J. M. Stacks 2: Analytical methods for paired-end sequencing improve RADseq-based population genomics. *Mol Ecol* **28**, 4737–4754 (2019).

52. Inbar, S. *et al.* Genetic basis of cuticular hydrocarbons in the desert ant. *In prep*.

53. Langmead B & Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9**, 357–359 (2012).

54. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).

55. Browning, B. L., Zhou, Y. & Browning, S. R. A One-Penny Imputed Genome from Next-Generation Reference Panels. *The American Journal of Human Genetics* **103**, 338–348 (2018).

56. Browning, B. L. & Browning, S. R. Improving the Accuracy and Efficiency of Identity-by-Descent Detection in Population Data. *Genetics* **194**, 459–471 (2013).

57. Weir, B. S. & Cockerham, C. C. Estimating F-Statistics for the Analysis of Population Structure. *Evolution* **38**, 1358 (1984).

58. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *The American Journal of Human Genetics* **81**, 559–575 (2007).

59. Altenhoff, A. M. *et al.* OMA standalone: orthology inference among public and custom genomes and transcriptomes. *Genome Res.* **29**, 1152–1163 (2019).

60. HAPpy-ABCENTH https://github.com/biorover/HAPpy-ABCENTH.

61. Forêt, S. & Maleszka, R. Function and evolution of a gene family encoding odorant binding-like proteins in a social insect, the honey bee ( *Apis mellifera* ). *Genome Res.* **16**, 1404–1413 (2006).

62. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**, 772–780 (2013).

63. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

# 7. Supplementary material

Supplementary Figure S1: Matrix of average aggression scores among nests. The intensity of cell color is proportional to the aggression score. The first four rows provide nest IDs, kinship, genotype at the supergene, number of queens found, and mitotypes (monogyne/polygyne). The monogyne form is denoted in blue and polygyne in green.

Supplementary Figure S2: (a) Histogram of aggression scores. (b) Aggression scores across social forms assigned based on mitotype.

Supplementary Figure S3: LD decay plot for *Cataglyphis niger* chromosomes. Blue dots represent the LD at corresponding SNP distances in monogyne samples, while green dots represent polygyne samples. The blue dashed and solid green lines indicate the average LD at each SNP distance in monogyne and polygyne samples, respectively.

Supplementary Figure S4: A plot of linkage disequilibrium (LD) along chromosome 2 in (a) polygyne and (b) monogyne samples.

(a)

(b)

Supplementary Figure S5: Phased genotypes of 65 daughters of a single polygyne queen and a single father (full-sisters). Only sites that were polymorphic in these samples are shown. The supergene region in chromosome 2 stands out for having 100% heterozygosity.

Supplementary Figure S6: Genomic data of broad scale sampling of populations of *C. niger* across Israel from Reiner Brodetzki et al. (2019), showing the presence of the supergene on chromosome 2. Monogyne and polygyne samples were assigned based on the mitotype. (a) Genome-wide $F_{ST}$ analysis using broad sampling of *C. niger* also shows a large number of high $F_{ST}$ SNPs on chromosome 2. (b) The high $F_{ST}$ region on chromosome 2 is consistent with our data from a single population. (c) Genotypes at the high $F_{ST}$ SNPs show the association between mitotypes and supergene genotypes. Samples bearing the polygyne mitotype are mostly heterozygous MP and monogyne samples are homozygous MM at the supergene. (d) Distribution of supergene genotypes across Israel.

(a)

(b)

(c)

(d)

Supplementary Figure S7: Synteny plot with 22 hymenopteran species based on one-to-one orthologous genes on the social chromosome of *Cataglyphis niger*.

Supplementary Figure S8: Dotplot of *Solenopsis* and *Cataglyphis* social chromosomes with gene names for one-to-one orthologs. Red dots indicate genes in the supergene region in both species.

Supplementary Figure S9: Subtree of the Odorant Receptor (OR) gene tree including genes in the *Cataglyphis* and *Solenopsis* social chromosomes. CnigOR = *Cataglyphis niger* OR; CfOR = *Camponotus floridanus* OR; SiOR = *Solenopsis invicta* OR; LhOR = *Linepithema humile* OR.

**Supplementary Table 1**: Information on nests sampled in Tel Baruch beach, geographic location, number of queens found, supergene genotype, and mitotype.

| Date | Nest ID | lat | long | Number of queens | kinship | Supergene genotype | Mitotype |
|---|---|---|---|---|---|---|---|
| 13.11.19 | TBA14 | 32.124964 | 34.784754 | – | monogyne | monogyne | savigyni |
| 13.11.19 | TBA15 | 32.1242833 | 34.7847167 | – | monogyne | monogyne | savigyni |
| 13.11.19 | TBA17 | 32.1258333 | 34.7873 | 1 | monogyne | monogyne | savigyni |
| 20.11.19 | TBA18 | 32.1252167 | 34.7858 | – | monogyne | monogyne | savigyni |
| 20.11.19 | TBA19 | 32.1259 | 34.7856333 | – | monogyne | monogyne | niger/savignyi |
| 20.11.19 | TBA20 | 32.12675 | 34.786556 | 2 | polygyne | polygyne | niger/savignyi |
| 20.11.19 | TBA21 | 32.1287 | 34.7878833 | – | polygyne | polygyne | niger |
| 20.11.19 | TBA22 | 32.129534 | 34.78787 | – | polygyne | polygyne | niger |
| 20.11.19 | TBA23 | 32.129225 | 34.7881031 | 2 | polygyne | polygyne | niger |
| 20.11.19 | TBA24 | 32.1299 | 34.7888333 | – | polygyne | polygyne | niger |
| 27.11.19 | TBA25 | 32.1252136 | 34.7881939 | – | monogyne | monogyne | drusus |
| 27.11.19 | TBA26 | 32.1262297 | 34.7862736 | 1 | monogyne | monogyne | drusus |
| 27.11.19 | TBA27 | 32.1245667 | 34.7855167 | – | monogyne | monogyne | drusus |
| 27.11.19 | TBA28 | 32.1280333 | 34.7871833 | 2 | polygyne | polygyne | niger |
| 27.11.19 | TBA29 | 32.1283833 | 34.7871333 | – | polygyne | polygyne | niger |
| 27.11.19 | TBA30 | 32.1280667 | 34.7870167 | – | polygyne | polygyne | niger |
| 02.12.19 | TBA31 | 32.123017 | 34.786583 | 1 | Monogyne | Polygyne | niger |
| 12.02.19 | TBA32 | 32.123513 | 34.786879 | 1 | polygyne | polygyne | niger |
| 12.02.19 | TBA33 | 32.1246006 | 34.7881331 | 1 | monogyne | monogyne | drusus |
| 12.02.19 | TBA34 | 32.1218833 | 34.7874167 | – | polygyne | polygyne | niger |
| 12.02.19 | TBA35 | 32.120884 | 34.787851 | 2 | polygyne | polygyne | niger |
| 12.02.19 | TBA36 | 32.1302175 | 34.79003 | – | polygyne | polygyne | niger |
| 12.02.19 | TBA37 | 32.1308167 | 34.7887833 | 2 | polygyne | polygyne | niger |
| 12.02.19 | TBA38 | 32.129777 | 34.789511 | 1 | polygyne | polygyne | niger/ drusus |
| 15.12.19 | TBA39 | 32.1249781 | 34.7865431 | – | monogyne | monogyne | savigyni |
| 15.12.19 | TBA40 | 32.1241111 | 34.7876389 | – | monogyne | monogyne | savigyni |
| 15.12.19 | TBA41 | 32.1250333 | 34.78595 | – | monogyne | monogyne | savigyni |
| 15.12.19 | TBA42 | 32.1310719 | 34.7905367 | 6 | polygyne | polygyne | niger |
| 15.12.19 | TBA43 | 32.130685 | 34.789609 | 3 | polygyne | polygyne | niger |
| 15.12.19 | TBA44 | 32.1305167 | 34.7893167 | 1 | polygyne | polygyne | niger |

**Supplementary Table 2:** List of one-to-one orthologous genes with functional annotation found on *Solenopsis* chromosome 16 and Cataglyphis chromosome 2. The highlighted cells indicate the genes found in the supergene region of the respective species.

| HOG | Cnig ch | Cnig_id | Cnig_start | Cnig_end | Sinv ch | Sinv_id | Sinv_start | Sinv_end | Sinv_function |
|---|---|---|---|---|---|---|---|---|---|
| HOG2874 | chr02 | gene_1564_tr1 | 3963068 | 3963349 | chr16 | XP_025990922.1 | 25532553 | 25534584 | glutenin, high molecular weight subunit DX5 |
| HOG14365 | chr02 | gene_1567_tr2 | 3971998 | 3974172 | chr16 | XP_011156788.1 | 25520361 | 25523691 | stAR-related lipid transfer protein 7, mitochondrial isoform X1 |
| HOG14370 | chr02 | gene_1937_tr1 | 3975447 | 3977432 | chr16 | XP_011156786.2 | 25517499 | 25519872 | DNA-directed primase/polymerase protein |
| HOG14988 | chr02 | gene_1922_tr1 | 3230128 | 3231288 | chr16 | XP_011164983.1 | 9360203 | 9363557 | DDRGK domain-containing protein 1 isoform X1 |
| HOG3816 | chr02 | gene_1894_tr1 | 2558786 | 2559621 | chr16 | XP_011156749.1 | 24916104 | 24918107 | histone RNA hairpin-binding protein |
| HOG8963 | chr02 | gene_1537_tr1 | 3150294 | 3154642 | chr16 | XP_039314572.1 | 12972095 | 13055442 | coiled-coil domain-containing protein R3HCC1L isoform X1 |
| HOG10455 | chr02 | gene_1510_tr1 | 2549965 | 2551574 | chr16 | XP_039315055.1 | 24927324 | 24931095 | S-adenosylmethionine mitochondrial carrier protein |
| HOG13778 | chr02 | gene_1540_tr2 | 3237179 | 3238657 | chr16 | XP_039314532.1 | 12063131 | 12112267 | DNA excision repair protein ERCC-1 isoform X1 |
| HOG10884 | chr02 | gene_1938_tr1 | 3979161 | 3980572 | chr16 | XP_011156782.1 | 25514403 | 25516090 | cuticle protein 21.3 |
| HOG9890 | chr02 | gene_1852_tr1 | 1685469 | 1700678 | chr16 | XP_039314798.1 | 15844676 | 15869563 | ABC transporter G family member 23 isoform X1 |
| HOG15358 | chr02 | gene_1930_tr1 | 3427464 | 3429176 | chr16 | XP_011167880.1 | 14926676 | 14929867 | pre-mRNA-splicing factor Slu7 |
| HOG15304 | chr02 | gene_1568_tr1 | 3983445 | 3986702 | chr16 | XP_025990992.2 | 25505722 | 25510892 | molybdenum cofactor sulfurase 3 |
| HOG9148 | chr02 | gene_1858_tr1 | 2051611 | 2054176 | chr16 | XP_011170086.1 | 16968597 | 16976194 | phosphoglucomutase isoform X1 |
| HOG15068 | chr02 | gene_1569_tr1 | 3988663 | 3991066 | chr16 | XP_025990993.1 | 25495806 | 25500737 | ras-related protein Rab-27A |
| HOG15085 | chr02 | gene_1939_tr1 | 4056694 | 4063002 | chr16 | XP_025990986.1 | 25336872 | 25415644 | protein artichoke isoform X1 |
| HOG8690 | chr02 | gene_1940_tr1 | 4381399 | 4384558 | chr16 | XP_011156768.2 | 25042442 | 25044976 | alpha-glucosidase |
| HOG13962 | chr02 | gene_1843_tr1 | 1377153 | 1379481 | chr16 | XP_011166565.2 | 24474013 | 24478058 | nicalin |
| HOG8446 | chr02 | gene_1844_tr1 | 1518193 | 1532996 | chr16 | XP_011164628.1 | 15263526 | 15305312 | homeotic protein empty spiracles |
| HOG9085 | chr02 | gene_1447_tr2 | 771890 | 772300 | chr16 | XP_011165219.2 | 23177534 | 23178507 | mitochondrial import inner membrane translocase subunit Tim16 |
| HOG8671 | chr02 | gene_1570_tr1 | 4414120 | 4420669 | chr16 | XP_011156769.2 | 24995645 | 25088183 | protein artichoke |
| HOG7911 | chr02 | gene_1867_tr1 | 2175099 | 2181479 | chr16 | XP_011170059.1 | 16779865 | 16787958 | protein FAM160B1 isoform X1 |

| HOG | chr | gene | start | end | chr | protein | start | end | description |
|---|---|---|---|---|---|---|---|---|---|
| HOG15320 | chr02 | gene_1538_tr1 | 3187547 | 3188773 | chr16 | XP_025996242.1 | 10068516 | 10130643 | tubulin polymerization-promoting protein homolog isoform X1 |
| HOG10354 | chr02 | gene_1442_tr2 | 598774 | 600601 | chr16 | XP_011165314.2 | 22885865 | 22888640 | rRNA N6-adenosine-methyltransferase ZCCHC4 |
| HOG7789 | chr02 | gene_1942_tr1 | 4466584 | 4467522 | chr16 | XP_011156758.1 | 24940738 | 24942742 | ELMO domain-containing protein 2 |
| HOG7302 | chr02 | gene_1943_tr1 | 4481091 | 4484240 | chr16 | XP_039314108.1 | 24577632 | 24585030 | cell death specification protein 2 |
| HOG3972 | chr02 | gene_1573_tr1 | 4492366 | 4493178 | chr16 | XP_039314761.1 | 24558695 | 24560167 | nucleolin-like |
| HOG8813 | chr02 | gene_1905_tr2 | 2847105 | 2848201 | chr16 | XP_011175190.1 | 17717140 | 17718878 | charged multivesicular body protein 5 |
| HOG10946 | chr02 | gene_1501_tr2 | 2452175 | 2454290 | chr16 | XP_011170021.1 | 16312880 | 16315978 | TGF-beta-activated kinase 1 and MAP3K7-binding protein 1 |
| HOG8441 | chr02 | gene_1945_tr2 | 4495244 | 4499484 | chr16 | XP_011170055.1 | 16730385 | 16734694 | 60S ribosomal protein L4 |
| HOG15067 | chr02 | gene_1575_tr7 | 4520673 | 4523032 | chr16 | XP_025989498.1 | 16702011 | 16707132 | aldo-keto reductase family 1 member B1 |
| HOG15839 | chr02 | gene_1582_tr1 | 4561481 | 4562970 | chr16 | XP_011170040.1 | 16662165 | 16664578 | peptide deformylase, mitochondrial |
| HOG8215 | chr02 | gene_1504_tr1 | 2476545 | 2484677 | chr16 | XP_039315026.1 | 16336210 | 16350025 | LOW QUALITY PROTEIN: serine/threonine-protein phosphatase 1 regulatory subunit 10 |
| HOG15092 | chr02 | gene_1948_tr4 | 4566904 | 4596170 | chr16 | XP_025989494.1 | 16575605 | 16659139 | tachykinin-like peptides receptor 99D isoform X1 |
| HOG8214 | chr02 | gene_1870_tr2 | 2218820 | 2226623 | chr16 | XP_039314336.1 | 24529290 | 24544420 | transferrin |
| HOG13033 | chr02 | gene_1949_tr1 | 4650675 | 4652957 | chr16 | XP_011170039.1 | 16566292 | 16569457 | sphingosine-1-phosphate lyase |
| HOG15150 | chr02 | gene_1950_tr1 | 4657835 | 4677811 | chr16 | XP_011170038.2 | 16538591 | 16564671 | tubby-related protein 4 |
| HOG14128 | chr02 | gene_1584_tr1 | 4680005 | 4681037 | chr16 | XP_011156805.1 | 25882931 | 25886321 | RWD domain-containing protein 1 |
| HOG14108 | chr02 | gene_1951_tr4 | 4684226 | 4690590 | chr16 | XP_011156808.3 | 25887243 | 25903896 | NADP-dependent malic enzyme isoform X1 |
| HOG17235 | chr02 | gene_1586_tr1 | 5014932 | 5016002 | chr16 | XP_011173007.1 | 13472076 | 13475943 | general odorant-binding protein 83a |
| HOG8114 | chr02 | gene_1588_tr1 | 5024531 | 5025745 | chr16 | XP_039314227.1 | 13391052 | 13471488 | dopamine receptor 2 |
| HOG14573 | chr02 | gene_1859_tr1 | 2065162 | 2066044 | chr16 | XP_011170087.1 | 16952983 | 16954745 | ATP synthase subunit C lysine N-methyltransferase |
| HOG8827 | chr02 | gene_1589_tr1 | 5045800 | 5046548 | chr16 | XP_011161724.2 | 13267097 | 13271493 | probable E3 ubiquitin-protein ligase RHY1A |
| HOG8077 | chr02 | gene_1502_tr1 | 2465955 | 2468779 | chr16 | XP_011170022.1 | 16326307 | 16330617 | V-type proton ATPase subunit B |
| HOG8826 | chr02 | gene_1958_tr1 | 5052175 | 5054200 | chr16 | XP_039314418.1 | 13289459 | 13298659 | UPF0687 protein C20orf27 homolog |

| HOG15232 | chr02 | gene_1932_tr1 | 3463774 | 3465313 | chr16 | XP_025993452.2 | 14979589 | 14984897 | TBC1 domain family member 7 isoform X1 |
|---|---|---|---|---|---|---|---|---|---|
| HOG15520 | chr02 | gene_1462_tr3 | 1350205 | 1352805 | chr16 | XP_011169804.1 | 24221257 | 24293784 | tachykinins |
| HOG8584 | chr02 | gene_1592_tr1 | 5061102 | 5063487 | chr16 | XP_011155907.1 | 21850661 | 21853990 | insulin-like growth factor-binding protein-related protein 1 |
| HOG15356 | chr02 | gene_1887_tr7 | 2500497 | 2507595 | chr16 | XP_011156723.1 | 24664117 | 24678278 | signal transducer and activator of transcription 5B |
| HOG16061 | chr02 | gene_1959_tr1 | 5063881 | 5081427 | chr16 | XP_011155899.1 | 21745846 | 21850839 | prominin-2 isoform X1 |
| HOG17057 | chr02 | gene_1440_tr1 | 577319 | 581594 | chr16 | XP_011165379.1 | 22852621 | 22858466 | coiled-coil domain-containing protein 142 isoform X1 |
| HOG8794 | chr02 | gene_1904_tr2 | 2840928 | 2843949 | chr16 | XP_011175214.1 | 17709621 | 17713021 | nuclear envelope integral membrane protein 1 isoform X1 |
| HOG8492 | chr02 | gene_1497_tr1 | 2394239 | 2399165 | chr16 | XP_011164711.1 | 16158349 | 16165729 | lysyl oxidase homolog 3 |
| HOG8964 | chr02 | gene_1916_tr1 | 3156807 | 3159565 | chr16 | XP_011166899.1 | 13059468 | 13063879 | signal recognition particle subunit SRP72 |
| HOG8673 | chr02 | gene_1594_tr1 | 5111114 | 5144049 | chr16 | XP_025987255.1 | 21751143 | 21823735 | trace amine-associated receptor 9 |
| HOG15025 | chr02 | gene_1458_tr2 | 1247917 | 1274925 | chr16 | XP_039315079.1 | 23900432 | 24065116 | protein tincar isoform X1 |
| HOG8672 | chr02 | gene_1960_tr2 | 5155982 | 5192533 | chr16 | XP_011155897.1 | 21475608 | 21744759 | leishmanolysin-like peptidase isoform X1 |
| HOG10640 | chr02 | gene_1524_tr2 | 2991447 | 3001477 | chr16 | XP_011170093.1 | 16977851 | 16988129 | huntingtin isoform X1 |
| HOG10453 | chr02 | gene_1885_tr1 | 2458091 | 2464714 | chr16 | XP_011170017.1 | 16318164 | 16325215 | 4-hydroxybutyrate coenzyme A transferase isoform X1 |
| HOG8674 | chr02 | gene_1961_tr4 | 5394807 | 5397857 | chr16 | XP_011155894.2 | 21468490 | 21472522 | ferrochelatase, mitochondrial |
| HOG10670 | chr02 | gene_1436_tr4 | 555219 | 560423 | chr16 | XP_011174290.1 | 22795556 | 22799567 | casein kinase I isoform X1 |
| HOG8675 | chr02 | gene_1595_tr1 | 5397895 | 5399539 | chr16 | XP_011155895.1 | 21465338 | 21469039 | EEF1A lysine methyltransferase 2 |
| HOG13826 | chr02 | gene_1526_tr2 | 3021334 | 3025332 | chr16 | XP_011170099.1 | 17016110 | 17024394 | endoplasmic reticulum metallopeptidase 1 isoform X1 |
| HOG8120 | chr02 | gene_1964_tr2 | 5421067 | 5421712 | chr16 | XP_025987260.1 | 21422668 | 21424863 | high affinity copper uptake protein 1 isoform X1 |
| HOG8210 | chr02 | gene_1486_tr2 | 2212659 | 2217507 | chr16 | XP_011156707.1 | 24545232 | 24553641 | cytoplasmic dynein 2 intermediate chain 1 isoform X1 |
| HOG8118 | chr02 | gene_1596_tr1 | 5425828 | 5435317 | chr16 | XP_011155888.2 | 21405290 | 21422491 | endoplasmic reticulum aminopeptidase 1 isoform X1 |
| HOG8665 | chr02 | gene_1906_tr2 | 3009188 | 3013281 | chr16 | XP_011170095.1 | 16992073 | 17009406 | probable sodium/potassium/calcium exchanger CG1090 |

| HOG8187 | chr02 | gene_1597_tr1 | 5443206 | 5533951 | chr16 | XP_011155884.1 | 21285578 | 21397783 | segmentation protein cap'n'collar isoform X1 |
|---|---|---|---|---|---|---|---|---|---|
| HOG8312 | chr02 | gene_1599_tr1 | 5570470 | 5579495 | chr16 | XP_011158158.1 | 21155372 | 21199818 | RNA/RNP complex-1-interacting phosphatase |
| HOG8186 | chr02 | gene_1600_tr1 | 5695305 | 5708193 | chr16 | XP_011158115.1 | 21018836 | 21079706 | fibroblast growth factor 17 isoform X1 |
| HOG8266 | chr02 | gene_1929_tr2 | 3421778 | 3426438 | chr16 | XP_039314407.1 | 14909767 | 14922009 | activin receptor type-2A isoform X1 |
| HOG13656 | chr02 | gene_1463_tr1 | 1358112 | 1362142 | chr16 | XP_011166559.1 | 24437663 | 24446348 | trimeric intracellular cation channel type 1B.1 |
| HOG8329 | chr02 | gene_1967_tr1 | 5744411 | 5815901 | chr16 | XP_011158097.2 | 20910154 | 20987545 | transcription factor collier isoform X1 |
| HOG8817 | chr02 | gene_1601_tr1 | 5870774 | 5872329 | chr16 | XP_039314709.1 | 20850242 | 20864404 | glutamine amidotransferase-like class 1 domain-containing protein 3A, mitochondrial |
| HOG16969 | chr02 | gene_1603_tr3 | 6164963 | 6195172 | chr16 | XP_039314822.1 | 20496762 | 20533250 | Kv channel-interacting protein 4 isoform X1 |
| HOG8721 | chr02 | gene_1972_tr1 | 6201764 | 6203381 | chr16 | XP_011158029.1 | 20490961 | 20493299 | protein farnesyltransferase/geranylgeranyltransferase type-1 subunit alpha |
| HOG8084 | chr02 | gene_1973_tr5 | 6204904 | 6208979 | chr16 | XP_011158015.1 | 20485060 | 20490596 | heat shock 70 kDa protein cognate 5 |
| HOG8364 | chr02 | gene_1974_tr1 | 6210501 | 6218674 | chr16 | XP_025987284.1 | 20473483 | 20484396 | peroxidasin homolog pxn-2 |
| HOG8547 | chr02 | gene_1975_tr1 | 6219804 | 6222112 | chr16 | XP_025987285.1 | 20468781 | 20473347 | U4/U6 small nuclear ribonucleoprotein Prp3 isoform X1 |
| HOG8499 | chr02 | gene_1925_tr1 | 3255213 | 3256394 | chr16 | XP_025993462.1 | 14621720 | 14625536 | charged multivesicular body protein 3 |
| HOG8670 | chr02 | gene_1976_tr1 | 6241885 | 6280032 | chr16 | XP_011157936.1 | 20053347 | 20462470 | nucleolysin TIAR isoform X1 |
| HOG8376 | chr02 | gene_1873_tr1 | 2264436 | 2284940 | chr16 | XP_039314194.1 | 16006971 | 16033707 | MAP kinase-activating death domain protein isoform X1 |
| HOG6288 | chr02 | gene_1977_tr1 | 6679051 | 6679895 | chr16 | XP_011157914.1 | 20000044 | 20001599 | peritrophin-1 |
| HOG8533 | chr02 | gene_1980_tr1 | 6744707 | 6759773 | chr16 | XP_011157873.1 | 19917347 | 19933877 | SEC23-interacting protein isoform X1 |
| HOG8591 | chr02 | gene_1935_tr1 | 3671398 | 3672611 | chr16 | XP_011156801.1 | 25865185 | 25871673 | transmembrane protein 230 isoform X1 |
| HOG16016 | chr02 | gene_1855_tr1 | 1799371 | 1801890 | chr16 | XP_011164652.2 | 15717500 | 15720868 | arrestin homolog |
| HOG16856 | chr02 | gene_1455_tr1 | 855531 | 877697 | chr16 | XP_039314720.1 | 23381477 | 23411042 | dynein heavy chain 8, axonemal |
| HOG15228 | chr02 | gene_1546_tr4 | 3458611 | 3461401 | chr16 | XP_039314410.1 | 14958604 | 14972956 | hippocampus abundant transcript 1 protein isoform X1 |
| HOG7851 | chr02 | gene_1469_tr1 | 1630955 | 1636315 | chr16 | XP_039314800.1 | 15983474 | 15991932 | catenin alpha isoform X1 |

| HOG7957 | chr02 | gene_1900_tr3 | 2796458 | 2800017 | chr16 | XP_039314811.1 | 18571136 | 18574460 | LOW QUALITY PROTEIN: leucine-rich repeat-containing protein 15-like |
|---|---|---|---|---|---|---|---|---|---|
| HOG11444 | chr02 | gene_1548_tr2 | 3466345 | 3467899 | chr16 | XP_011167869.1 | 14972134 | 14977920 | methyltransferase-like protein 25 |
| HOG11151 | chr02 | gene_1882_tr1 | 2445602 | 2447802 | chr16 | XP_025989556.1 | 16306420 | 16308828 | titin |
| HOG16017 | chr02 | gene_1853_tr1 | 1733138 | 1733878 | chr16 | XP_011164666.1 | 15796782 | 15799883 | copper homeostasis protein cutC homolog |
| HOG7968 | chr02 | gene_1889_tr1 | 2536551 | 2541142 | chr16 | XP_025990923.1 | 24608618 | 24616794 | SWI/SNF complex subunit SMARCC2 |
| HOG9836 | chr02 | gene_1605_tr10 | 6783211 | 6810424 | chr16 | XP_039314308.1 | 19862587 | 19891241 | protein unc-13 homolog 4B isoform X1 |
| HOG11098 | chr02 | gene_1432_tr1 | 366596 | 369341 | chr16 | XP_011174335.1 | 22583577 | 22588088 | vesicular integral-membrane protein VIP36 |
| HOG16020 | chr02 | gene_1983_tr3 | 6785469 | 6786881 | chr16 | XP_011157834.1 | 19886682 | 19888422 | trypsin 5G1 isoform X1 |
| HOG8702 | chr02 | gene_1606_tr1 | 7121846 | 7149177 | chr16 | XP_011157760.1 | 19496127 | 19823163 | protein slit isoform X1 |
| HOG8535 | chr02 | gene_1986_tr1 | 7206491 | 7209160 | chr16 | XP_011157681.2 | 19412935 | 19415803 | flap endonuclease GEN |
| HOG7857 | chr02 | gene_1847_tr3 | 1601008 | 1605352 | chr16 | XP_011164616.2 | 15167201 | 15175918 | pickpocket protein 28 |
| HOG15089 | chr02 | gene_1481_tr1 | 2066442 | 2067463 | chr16 | XP_011170090.1 | 16951061 | 16953049 | ribulose-phosphate 3-epimerase |
| HOG7896 | chr02 | gene_1610_tr2 | 7209666 | 7211413 | chr16 | XP_011157667.1 | 19409895 | 19412851 | charged multivesicular body protein 7 |
| HOG7898 | chr02 | gene_1988_tr2 | 7216732 | 7219024 | chr16 | XP_011157570.1 | 19367690 | 19374112 | tRNA pseudouridine synthase A isoform X1 |
| HOG7900 | chr02 | gene_1611_tr1 | 7219634 | 7221695 | chr16 | XP_011157615.1 | 19369186 | 19371285 | probable 39S ribosomal protein L45, mitochondrial |
| HOG7964 | chr02 | gene_1898_tr2 | 2790895 | 2794896 | chr16 | XP_039315010.1 | 17568536 | 17574944 | cysteine sulfinic acid decarboxylase |
| HOG8217 | chr02 | gene_1488_tr1 | 2235039 | 2237092 | chr16 | XP_011164632.1 | 15448658 | 15451149 | F-box/WD repeat-containing protein 4 isoform X1 |
| HOG8129 | chr02 | gene_1831_tr4 | 748263 | 770696 | chr16 | XP_011165243.1 | 23132897 | 23176944 | attractin-like protein 1 isoform X1 |
| HOG8028 | chr02 | gene_1991_tr1 | 7234859 | 7238940 | chr16 | XP_025987316.1 | 19091243 | 19345435 | tyrosine-protein kinase transmembrane receptor Ror isoform X1 |
| HOG8472 | chr02 | gene_1535_tr2 | 3108146 | 3134625 | chr16 | XP_025994936.2 | 12436327 | 12627257 | protein lap4 isoform X1 |
| HOG8707 | chr02 | gene_1453_tr1 | 822789 | 828821 | chr16 | XP_011165091.1 | 23303851 | 23328875 | rho-related BTB domain-containing protein 1 isoform X1 |
| HOG8759 | chr02 | gene_1466_tr1 | 1589113 | 1593117 | chr16 | XP_039315121.1 | 15179363 | 15192867 | zinc finger protein 260 isoform X1 |
| HOG8106 | chr02 | gene_1994_tr2 | 7468244 | 7472789 | chr16 | XP_025987315.1 | 19079404 | 19084961 | transcription elongation factor SPT5 |
| HOG9418 | chr02 | gene_1872_tr6 | 2259064 | 2261708 | chr16 | XP_025991932.1 | 16004175 | 16006757 | farnesol dehydrogenase isoform X1 |

| HOG10889 | chr02 | gene_1435_tr3 | 544379 | 548693 | chr16 | XP_025987290.1 | 22671772 | 22789697 | dopamine receptor 1 isoform X2 |
|---|---|---|---|---|---|---|---|---|---|
| HOG8596 | chr02 | gene_1995_tr2 | 7479424 | 7487751 | chr16 | XP_011161364.1 | 19064971 | 19074495 | protein O-GlcNAcase isoform X1 |
| HOG4475 | chr02 | gene_1864_tr3 | 2116467 | 2124441 | chr16 | XP_039314456.1 | 16843476 | 16854742 | titin homolog isoform X5 |
| HOG10645 | chr02 | gene_1441_tr4 | 590415 | 596041 | chr16 | XP_025990760.1 | 22873090 | 22882736 | phosphatidylinositide phosphatase SAC2 isoform X1 |
| HOG8589 | chr02 | gene_1561_tr1 | 3810773 | 3815356 | chr16 | XP_025990962.1 | 25692531 | 25843341 | AT-rich interactive domain-containing protein 5B isoform X1 |
| HOG7954 | chr02 | gene_1470_tr1 | 1644698 | 1667692 | chr16 | XP_011164684.1 | 15942310 | 15976783 | BCL2/adenovirus E1B 19 kDa protein-interacting protein 3 |
| HOG13636 | chr02 | gene_1616_tr1 | 7553722 | 7607336 | chr16 | XP_039314493.1 | 18886118 | 19024272 | potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 3 isoform X1 |
| HOG10644 | chr02 | gene_1828_tr2 | 561841 | 570185 | chr16 | XP_011174263.3 | 22802774 | 22813157 | dynactin subunit 1 isoform X1 |
| HOG13640 | chr02 | gene_1617_tr1 | 7628100 | 7633710 | chr16 | XP_011155781.1 | 18865932 | 18880975 | carbohydrate sulfotransferase 11 |
| HOG9816 | chr02 | gene_1518_tr1 | 2817961 | 2822157 | chr16 | NP_001291533.1 | 17618379 | 17624016 | transferrin precursor |
| HOG8390 | chr02 | gene_1836_tr2 | 838903 | 853826 | chr16 | XP_039314181.1 | 23361028 | 23379858 | vacuolar protein sorting-associated protein 13B |
| HOG8551 | chr02 | gene_1618_tr1 | 7636416 | 7641713 | chr16 | XP_011155778.1 | 18840151 | 18849161 | poly [ADP-ribose] polymerase tankyrase isoform X1 |
| HOG8680 | chr02 | gene_1493_tr1 | 2305670 | 2309668 | chr16 | XP_039314217.1 | 16050663 | 16058164 | N-acetyl-D-glucosamine kinase isoform X1 |
| HOG16940 | chr02 | gene_1908_tr2 | 3054063 | 3063955 | chr16 | XP_025989515.1 | 17054056 | 17067361 | atrial natriuretic peptide receptor 1 isoform X1 |
| HOG10676 | chr02 | gene_1850_tr1 | 1627454 | 1628788 | chr16 | XP_011164612.1 | 15141050 | 15143147 | beta-1,4-galactosyltransferase 7 |
| HOG8666 | chr02 | gene_1907_tr2 | 3018332 | 3019390 | chr16 | XP_011170096.2 | 17010831 | 17012794 | tetratricopeptide repeat protein 1 |
| HOG8474 | chr02 | gene_1468_tr2 | 1625291 | 1627027 | chr16 | XP_011164611.1 | 15142997 | 15145344 | alpha-2-macroglobulin receptor-associated protein |
| HOG8563 | chr02 | gene_1619_tr2 | 7659741 | 7663127 | chr16 | XP_039315141.1 | 18807749 | 18837105 | protein phosphatase 1 regulatory subunit 3C-B |
| HOG8562 | chr02 | gene_1620_tr2 | 7676877 | 7680074 | chr16 | XP_025990739.2 | 24334297 | 24342397 | venom protease isoform X1 |
| HOG8001 | chr02 | gene_1500_tr3 | 2421470 | 2427974 | chr16 | XP_025991945.2 | 16188502 | 16201893 | cGMP-specific 3',5'-cyclic phosphodiesterase isoform X1 |
| HOG8571 | chr02 | gene_1998_tr1 | 7683628 | 7685506 | chr16 | XP_011155767.1 | 18738035 | 18741095 | interleukin enhancer-binding factor 2 |
| HOG7835 | chr02 | gene_1509_tr1 | 2545617 | 2546126 | chr16 | XP_011156760.1 | 24934835 | 24938580 | high mobility group protein HMGI-C |

| HOG7903 | chr02 | gene_1999_tr1 | 7693975 | 7736544 | chr16 | XP_039314803.1 | 18652601 | 18722783 | disintegrin and metalloproteinase domain-containing protein 10 |
|---|---|---|---|---|---|---|---|---|---|
| HOG8183 | chr02 | gene_2000_tr1 | 7749120 | 7766692 | chr16 | XP_025994007.1 | 18623415 | 18645646 | neuronal acetylcholine receptor subunit alpha-10 isoform X1 |
| HOG5340 | chr02 | gene_1485_tr2 | 2203039 | 2206996 | chr16 | XP_011170067.1 | 16753719 | 16784018 | BRISC complex subunit FAM175B |
| HOG10009 | chr02 | gene_1830_tr3 | 633905 | 642281 | chr16 | XP_025990751.1 | 22946890 | 23121595 | octopamine receptor Oamb isoform X1 |
| HOG8785 | chr02 | gene_1521_tr1 | 2844452 | 2846734 | chr16 | XP_011175179.1 | 17714742 | 17717154 | tubulin epsilon chain |
| HOG15914 | chr02 | gene_2001_tr1 | 7768659 | 7770011 | chr16 | XP_039314809.1 | 18619452 | 18623202 | F-box/LRR-repeat protein 7 |
| HOG9902 | chr02 | gene_1624_tr1 | 7773176 | 7775233 | chr16 | XP_025994002.2 | 18608344 | 18615214 | E3 ubiquitin-protein ligase COP1 isoform X1 |
| HOG8386 | chr02 | gene_1628_tr1 | 7787479 | 7788379 | chr16 | XP_011155758.1 | 18595267 | 18597609 | AN1-type zinc finger protein 1 |
| HOG8031 | chr02 | gene_2003_tr1 | 7790245 | 7791748 | chr16 | XP_011155757.1 | 18591135 | 18595161 | tetraspanin-33 |
| HOG10675 | chr02 | gene_1438_tr1 | 587446 | 588814 | chr16 | XP_011165354.1 | 22866931 | 22869701 | ubiquitin-like domain-containing CTD phosphatase 1 |
| HOG7970 | chr02 | gene_1454_tr1 | 836500 | 838102 | chr16 | XP_011165023.1 | 23358829 | 23361307 | RING finger protein 37 isoform X1 |
| HOG13937 | chr02 | gene_1528_tr1 | 3048276 | 3053169 | chr16 | XP_025989522.1 | 17046676 | 17054767 | cysteine and histidine-rich protein 1 homolog |
| HOG8965 | chr02 | gene_1895_tr1 | 2563359 | 2569774 | chr16 | XP_011156748.1 | 24904560 | 24912249 | KAT8 regulatory NSL complex subunit 3 |
| HOG7955 | chr02 | gene_1629_tr1 | 7792556 | 7796617 | chr16 | XP_011155754.1 | 18586130 | 18590977 | bromodomain-containing protein 8 isoform X1 |
| HOG7956 | chr02 | gene_1630_tr1 | 7797372 | 7801982 | chr16 | XP_025994004.1 | 18580027 | 18586032 | heparanase isoform X2 |
| HOG8212 | chr02 | gene_1871_tr1 | 2237831 | 2247764 | chr16 | XP_039315119.1 | 15451139 | 15461125 | cytoplasmic aconitate hydratase isoform X1 |
| HOG7975 | chr02 | gene_1507_tr2 | 2529255 | 2532857 | chr16 | XP_011156721.1 | 24619302 | 24626210 | protein PTCD3 homolog, mitochondrial |
| HOG16290 | chr02 | gene_1631_tr1 | 7804773 | 7807974 | chr16 | XP_039314810.1 | 18574045 | 18577426 | tektin-3-like |
| HOG8371 | chr02 | gene_2005_tr2 | 7808831 | 7815525 | chr16 | XP_011159710.1 | 18529940 | 18550235 | glycerol-3-phosphate acyltransferase 1, mitochondrial isoform X1 |
| HOG9069 | chr02 | gene_1832_tr1 | 772763 | 775761 | chr16 | XP_011165231.1 | 23178555 | 23182124 | exonuclease 3'-5' domain-containing protein 2 |
| HOG7945 | chr02 | gene_1632_tr1 | 7829322 | 7833913 | chr16 | XP_011159700.2 | 18517852 | 18528961 | biorientation of chromosomes in cell division protein 1-like 1 |
| HOG8801 | chr02 | gene_1633_tr3 | 7839020 | 7842392 | chr16 | XP_011159621.1 | 18504596 | 18511068 | mitoferrin-1 isoform X1 |

| HOG | chr | gene | start | end | chr | protein | start | end | description |
|---|---|---|---|---|---|---|---|---|---|
| HOG16229 | chr02 | gene_2006_tr1 | 7846282 | 7848004 | chr16 | XP_039314701.1 | 18494518 | 18497581 | meiosis-specific nuclear structural protein 1 |
| HOG7890 | chr02 | gene_1634_tr1 | 7848863 | 7852708 | chr16 | XP_011159689.1 | 18489958 | 18494590 | ATP-dependent RNA helicase SUV3 homolog, mitochondrial |
| HOG8949 | chr02 | gene_2008_tr3 | 7860224 | 7865015 | chr16 | XP_025989569.1 | 18470260 | 18477332 | phenoloxidase-activating factor 2 isoform X1 |
| HOG9131 | chr02 | gene_2010_tr1 | 7871199 | 7874674 | chr16 | XP_011159599.3 | 18461337 | 18466462 | phenoloxidase-activating factor 2 |
| HOG9369 | chr02 | gene_1635_tr1 | 7988690 | 7992369 | chr16 | XP_011159533.1 | 18297095 | 18301594 | conserved oligomeric Golgi complex subunit 7 |
| HOG7416 | chr02 | gene_2013_tr3 | 7995757 | 8004436 | chr16 | XP_011159511.1 | 18259777 | 18296909 | putative leucine-rich repeat-containing protein DDB_G0290503 isoform X1 |
| HOG9436 | chr02 | gene_1636_tr2 | 8259469 | 8285211 | chr16 | XP_039315011.1 | 17954582 | 18255677 | probable JmjC domain-containing histone demethylation protein 2C isoform X1 |
| HOG14901 | chr02 | gene_1637_tr3 | 8292437 | 8299997 | chr16 | XP_011175125.1 | 17933856 | 17953352 | tyrosine-protein phosphatase non-receptor type 9 |
| HOG10456 | chr02 | gene_1893_tr1 | 2554419 | 2555497 | chr16 | XP_011156752.1 | 24921737 | 24923928 | phosphoglycerate mutase 2 |
| HOG9018 | chr02 | gene_1638_tr3 | 8304929 | 8307892 | chr16 | XP_011175137.1 | 17903198 | 17907453 | cell division cycle protein 16 homolog |
| HOG8353 | chr02 | gene_2019_tr1 | 8388278 | 8391469 | chr16 | XP_039314831.1 | 24695207 | 24810271 | RNA-binding protein Musashi homolog 2 isoform X1 |
| HOG10888 | chr02 | gene_1827_tr1 | 550915 | 552731 | chr16 | XP_011174313.1 | 22790349 | 22794822 | methionine--tRNA ligase, mitochondrial |
| HOG8653 | chr02 | gene_2020_tr1 | 8478578 | 8479850 | chr16 | XP_025991944.1 | 15119956 | 15121483 | proteasome subunit alpha type-2 |
| HOG15174 | chr02 | gene_1914_tr1 | 3146965 | 3149376 | chr16 | XP_011166886.1 | 12933551 | 12971889 | aspartate aminotransferase, cytoplasmic isoform X1 |
| HOG7786 | chr02 | gene_1539_tr1 | 3231656 | 3232729 | chr16 | XP_011175281.1 | 9247439 | 9248968 | transmembrane emp24 domain-containing protein eca |
| HOG7201 | chr02 | gene_1647_tr3 | 8480522 | 8481710 | chr16 | XP_011164604.1 | 15122313 | 15124685 | sperm flagellar protein 1 |
| HOG7912 | chr02 | gene_1868_tr3 | 2184745 | 2187525 | chr16 | XP_011170060.1 | 16762066 | 16777830 | hexokinase type 2 isoform X1 |
| HOG9344 | chr02 | gene_2027_tr2 | 8520769 | 8523402 | chr16 | XP_025990966.2 | 24851559 | 24857353 | glucose dehydrogenase [FAD, quinone] |
| HOG9067 | chr02 | gene_1652_tr1 | 8525966 | 8529287 | chr16 | XP_011157594.1 | 24859458 | 24863682 | ralA-binding protein 1 isoform X1 |
| HOG9202 | chr02 | gene_2028_tr2 | 8532952 | 8538883 | chr16 | XP_039315048.1 | 24864372 | 24874524 | exportin-7 isoform X1 |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| HOG7853 | chr02 | gene_1851_tr1 | 1641742 | 1642471 | chr16 | XP_011164686.1 | 15978070 | 15983195 | polyadenylate-binding protein-interacting protein 2 |
| HOG10764 | chr02 | gene_1823_tr1 | 372066 | 375575 | chr16 | XP_011174346.1 | 22576770 | 22583464 | wolframin isoform X1 |
| HOG6280 | chr02 | gene_1653_tr2 | 8541217 | 8541938 | chr16 | XP_011170133.1 | 17335959 | 17337451 | nucleolar protein 16 |
| HOG10999 | chr02 | gene_2029_tr1 | 8542888 | 8547976 | chr16 | XP_011170131.1 | 17328625 | 17335654 | fumarylacetoacetate hydrolase domain-containing protein 2A isoform X1 |
| HOG6326 | chr02 | gene_2030_tr2 | 8555871 | 8557342 | chr16 | XP_011170130.1 | 17325990 | 17327596 | 39S ribosomal protein L35, mitochondrial |
| HOG11158 | chr02 | gene_1655_tr3 | 8557444 | 8592647 | chr16 | XP_025989546.2 | 17289159 | 17326197 | MICOS complex subunit Mic60 isoform X1 |
| HOG8966 | chr02 | gene_1512_tr1 | 2560443 | 2562269 | chr16 | XP_025990944.2 | 24912668 | 24916031 | crossover junction endonuclease EME1 |
| HOG13454 | chr02 | gene_1657_tr3 | 8679162 | 8687330 | chr16 | XP_011170127.1 | 17185335 | 17196261 | sorting nexin-27 |
| HOG5859 | chr02 | gene_2032_tr2 | 8695365 | 8696724 | chr16 | XP_025989525.2 | 17178783 | 17181074 | nucleoside diphosphate kinase homolog 5 isoform X1 |
| HOG13861 | chr02 | gene_2033_tr1 | 8709815 | 8752864 | chr16 | XP_039314262.1 | 17113859 | 17170291 | microtubule-associated protein futsch isoform X1 |
| HOG8381 | chr02 | gene_1824_tr2 | 377038 | 391543 | chr16 | XP_039314351.1 | 22545731 | 22573243 | alpha-sarcoglycan isoform X1 |
| HOG14644 | chr02 | gene_2041_tr2 | 8858673 | 8861002 | chr16 | XP_039314676.1 | 26004603 | 26021597 | flavin-containing monooxygenase FMO GS-OX-like 2 isoform X1 |
| HOG6159 | chr02 | gene_2043_tr2 | 8883074 | 8892091 | chr16 | XP_039314679.1 | 26047002 | 26062255 | G protein-activated inward rectifier potassium channel 3 isoform X1 |
| HOG3995 | chr02 | gene_2044_tr1 | 8894630 | 8894947 | chr16 | XP_039314683.1 | 26062324 | 26064224 | oxidoreductase-like domain-containing protein 1 |
| HOG8532 | chr02 | gene_2048_tr1 | 8934105 | 8942741 | chr16 | XP_039314299.1 | 9393467 | 9416297 | protein phosphatase PP2A 55 kDa regulatory subunit isoform X1 |
| HOG8475 | chr02 | gene_1913_tr2 | 3137648 | 3139191 | chr16 | XP_039314908.1 | 12609397 | 12751315 | axonemal dynein light intermediate polypeptide 1 isoform X1 |
| HOG8372 | chr02 | gene_1862_tr1 | 2076255 | 2081427 | chr16 | XP_039314429.1 | 16927260 | 16938473 | ABC transporter G family member 20 isoform X1 |
| HOG13979 | chr02 | gene_1532_tr1 | 3074660 | 3076528 | chr16 | XP_011170104.1 | 17082495 | 17085279 | m7GpppN-mRNA hydrolase |
| HOG11556 | chr02 | gene_1456_tr1 | 1127222 | 1164340 | chr16 | XP_039314721.1 | 23829468 | 23896143 | dynein heavy chain 2, axonemal |
| HOG11760 | chr02 | gene_1934_tr1 | 3478522 | 3481130 | chr16 | XP_011167863.1 | 15008257 | 15013547 | zinc finger protein 2 homolog isoform X1 |

| HOG13784 | chr02 | gene_2051_tr4 | 8974048 | 8975733 | chr16 | XP_011161972.1 | 14420379 | 14426100 | CDGSH iron-sulfur domain-containing protein 2 homolog |
|---|---|---|---|---|---|---|---|---|---|
| HOG14394 | chr02 | gene_1829_tr6 | 574059 | 576207 | chr16 | XP_011174246.2 | 22820511 | 22823223 | tubulin alpha chain |
| HOG14992 | chr02 | gene_1663_tr2 | 8977052 | 8994687 | chr16 | XP_011161963.1 | 14429099 | 14452723 | PH and SEC7 domain-containing protein isoform X1 |
| HOG8107 | chr02 | gene_1883_tr3 | 2448818 | 2451885 | chr16 | XP_011170016.1 | 16309754 | 16312862 | ATP-dependent RNA helicase abstrakt |
| HOG8366 | chr02 | gene_1503_tr2 | 2470276 | 2473546 | chr16 | XP_025989473.1 | 16331052 | 16334872 | DNA repair and recombination protein RAD54-like |
| HOG7834 | chr02 | gene_1479_tr1 | 2046560 | 2047282 | chr16 | XP_011170092.1 | 16975041 | 16977580 | U3 small nucleolar ribonucleoprotein protein IMP3 |
| HOG14931 | chr02 | gene_2053_tr1 | 9002585 | 9043915 | chr16 | XP_025993470.2 | 14458008 | 14533290 | histone-lysine N-methyltransferase 2C isoform X1 |
| HOG9408 | chr02 | gene_2054_tr1 | 9057622 | 9062025 | chr16 | XP_011161961.1 | 14555366 | 14562042 | vacuolar protein sorting-associated protein 51 homolog |
| HOG8450 | chr02 | gene_2056_tr1 | 9170851 | 9171782 | chr16 | XP_011174382.1 | 22513988 | 22515516 | probable small nuclear ribonucleoprotein G |
| HOG14655 | chr02 | gene_2068_tr2 | 9260339 | 9263319 | chr16 | XP_039314678.1 | 26000887 | 26005049 | trypsin isoform X1 |
| HOG10457 | chr02 | gene_1892_tr1 | 2552211 | 2553631 | chr16 | XP_011156754.1 | 24924414 | 24926934 | ras-related protein Rab-24 isoform X1 |
| HOG13453 | chr02 | gene_1841_tr4 | 1366591 | 1370489 | chr16 | XP_039314259.1 | 24453266 | 24462382 | mitogen-activated protein kinase p38b isoform X2 |
| HOG7910 | chr02 | gene_1499_tr1 | 2409028 | 2411611 | chr16 | XP_011164715.1 | 16178869 | 16182494 | synaptic vesicular amine transporter |
| HOG15043 | chr02 | gene_1668_tr3 | 9268769 | 9269450 | chr16 | XP_025990985.1 | 25927528 | 25929537 | guanine nucleotide exchange factor MSS4 homolog |
| HOG15088 | chr02 | gene_1860_tr1 | 2067635 | 2068741 | chr16 | XP_011170091.1 | 16949785 | 16951257 | putative methyltransferase C9orf114 homolog |
| HOG12499 | chr02 | gene_1669_tr1 | 9279870 | 9280947 | chr16 | XP_011165269.1 | 26162881 | 26164234 | coiled-coil domain-containing protein 137 |
| HOG12500 | chr02 | gene_2072_tr1 | 9281188 | 9282321 | chr16 | XP_011165270.1 | 26160306 | 26162897 | BLOC-1-related complex subunit 5 |
| HOG13927 | chr02 | gene_1529_tr1 | 3072541 | 3073664 | chr16 | XP_011170107.1 | 17080878 | 17082139 | phosphomevalonate kinase |
| HOG15180 | chr02 | gene_1536_tr1 | 3144103 | 3144456 | chr16 | XP_011173101.1 | 12828417 | 12829476 | small integral membrane protein 8 |
| HOG8491 | chr02 | gene_1877_tr1 | 2366397 | 2379139 | chr16 | XP_011164713.1 | 16112917 | 16203623 | G protein-coupled receptor kinase 5 |
| HOG16760 | chr02 | gene_1531_tr1 | 3071636 | 3072259 | chr16 | XP_011170108.1 | 17079637 | 17080688 | 60S ribosomal protein L29-1 |
| HOG9345 | chr02 | gene_1670_tr1 | 9380689 | 9384563 | chr16 | XP_011174414.1 | 22386390 | 22391289 | cilia- and flagella-associated protein 206 |

| HOG9222 | chr02 | gene_1931_tr2 | 3437143 | 3452043 | chr16 | XP_011167875.2 | 14934035 | 14957100 | serine/threonine-protein kinase WNK1 isoform X1 |
|---------|-------|---------------|---------|---------|-------|----------------|----------|----------|----|
| HOG9346 | chr02 | gene_2073_tr2 | 9386284 | 9390095 | chr16 | XP_011174406.1 | 22380812 | 22386885 | upstream activation factor subunit spp27 |
| HOG7841 | chr02 | gene_1672_tr5 | 9719909 | 9741305 | chr16 | XP_039314315.1 | 22039333 | 22090099 | mushroom body large-type Kenyon cell-specific protein 1 isoform X1 |
| HOG8606 | chr02 | gene_2075_tr1 | 9772677 | 9774527 | chr16 | XP_011174441.1 | 22017324 | 22019290 | RCC1 domain-containing protein 1 |
| HOG12176 | chr02 | gene_1533_tr1 | 3088468 | 3097185 | chr16 | XP_039314149.1 | 12368253 | 12425549 | tuberin isoform X1 |
| HOG8607 | chr02 | gene_1674_tr2 | 9776722 | 9779510 | chr16 | XP_011174433.1 | 22014511 | 22017393 | ADP-ribosylation factor-like protein 3 |
| HOG5528 | chr02 | gene_1676_tr1 | 9808188 | 9810721 | chr16 | XP_011174500.3 | 21993059 | 21997112 | odorant receptor 22c isoform X1 |
| HOG13766 | chr02 | gene_1923_tr3 | 3233620 | 3236643 | chr16 | XP_011161971.1 | 14413614 | 14419966 | STAM-binding protein-like A |
| HOG15688 | chr02 | gene_1695_tr1 | 10162993 | 10164554 | chr16 | XP_011170189.2 | 3995281 | 3997294 | cytoplasmic tRNA 2-thiolation protein 2 |
| HOG9913 | chr02 | gene_1699_tr1 | 10195772 | 10199054 | chr16 | XP_011155530.1 | 7943071 | 7946078 | scaffold protein salvador isoform X1 |
| HOG10327 | chr02 | gene_2094_tr1 | 10201899 | 10220257 | chr16 | XP_011155529.1 | 7919926 | 7939780 | serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit epsilon isoform |
| HOG10328 | chr02 | gene_1700_tr1 | 10223464 | 10224732 | chr16 | XP_039314545.1 | 7910802 | 7913008 | JNK1/MAPK8-associated membrane protein |
| HOG8448 | chr02 | gene_1492_tr1 | 2302382 | 2303597 | chr16 | XP_011164701.1 | 16048363 | 16050400 | 39S ribosomal protein L21, mitochondrial |
| HOG10326 | chr02 | gene_2095_tr1 | 10225309 | 10229586 | chr16 | XP_011155524.2 | 7905195 | 7910729 | maltase 1 isoform X1 |
| HOG13663 | chr02 | gene_1464_tr1 | 1371290 | 1373226 | chr16 | XP_011166564.1 | 24464033 | 24468153 | ATPase ASNA1 homolog |
| HOG8679 | chr02 | gene_1876_tr1 | 2317939 | 2327431 | chr16 | XP_039314208.1 | 16065271 | 16075868 | serine/arginine repetitive matrix protein 1 isoform X1 |
| HOG8382 | chr02 | gene_1874_tr8 | 2290850 | 2301671 | chr16 | XP_011164698.1 | 16035848 | 16048178 | choline/ethanolaminephosphotransferase 1 isoform X7 |
| HOG7282 | chr02 | gene_2096_tr1 | 10288709 | 10301448 | chr16 | XP_039314844.1 | 7085225 | 7094186 | LOW QUALITY PROTEIN: sodium channel protein Nach |
| HOG11768 | chr02 | gene_1549_tr1 | 3475886 | 3478196 | chr16 | XP_011167862.1 | 15005569 | 15008087 | ornithine aminotransferase, mitochondrial |
| HOG13925 | chr02 | gene_1530_tr2 | 3066697 | 3070447 | chr16 | XP_025989517.1 | 17071223 | 17076558 | calcium-independent phospholipase A2-gamma isoform X1 |
| HOG9382 | chr02 | gene_2097_tr1 | 10362251 | 10364573 | chr16 | XP_011172453.1 | 6907523 | 7038323 | leucine-rich repeat-containing protein 24 |

| HOG9203 | chr02 | gene_1514_tr1 | 2574022 | 2585803 | chr16 | XP_011156744.1 | 24880664 | 24900576 | beta-TrCP |
|---|---|---|---|---|---|---|---|---|---|
| HOG9109 | chr02 | gene_1513_tr1 | 2570850 | 2572513 | chr16 | XP_011156746.1 | 24901145 | 24903477 | phospholipid phosphatase 5 |
| HOG8758 | chr02 | gene_1845_tr2 | 1561287 | 1564013 | chr16 | XP_025991919.1 | 15221127 | 15224921 | WD repeat-containing protein 91 isoform X1 |
| HOG15284 | chr02 | gene_2098_tr1 | 10399691 | 10400295 | chr16 | XP_011171593.1 | 7777629 | 7779233 | 60S ribosomal protein L28 |
| HOG15280 | chr02 | gene_2099_tr1 | 10400938 | 10402105 | chr16 | XP_011171597.1 | 7779426 | 7780932 | S-adenosylmethionine sensor upstream of mTORC1 |
| HOG15309 | chr02 | gene_1702_tr3 | 10403239 | 10409722 | chr16 | XP_025993055.2 | 6869805 | 6878400 | STE20-like serine/threonine-protein kinase isoform X1 |
| HOG15310 | chr02 | gene_2100_tr2 | 10411652 | 10413446 | chr16 | XP_011172448.1 | 6867425 | 6871295 | WD repeat-containing protein 13 |
| HOG10953 | chr02 | gene_1884_tr1 | 2454899 | 2457123 | chr16 | XP_011170020.1 | 16315660 | 16317833 | 4-hydroxybutyrate coenzyme A transferase |
| HOG14306 | chr02 | gene_1437_tr1 | 571010 | 572742 | chr16 | XP_011174253.1 | 22813296 | 22817685 | nuclear inhibitor of protein phosphatase 1 |
| HOG15226 | chr02 | gene_2101_tr2 | 10418561 | 10427039 | chr16 | XP_011172444.1 | 6856005 | 6866241 | protein lingerer isoform X1 |
| HOG8216 | chr02 | gene_1489_tr1 | 2231292 | 2234142 | chr16 | XP_011164629.1 | 15439689 | 15443682 | major facilitator superfamily domain-containing protein 6 |
| HOG9083 | chr02 | gene_1475_tr3 | 1802955 | 1807411 | chr16 | XP_011164651.2 | 15712031 | 15717641 | erythroid differentiation-related factor 1 |
| HOG14511 | chr02 | gene_1703_tr2 | 10430301 | 10436720 | chr16 | XP_011171054.1 | 5595847 | 5602703 | mediator of RNA polymerase II transcription subunit 23 |
| HOG7002 | chr02 | gene_1704_tr2 | 10446133 | 10448575 | chr16 | XP_011171042.1 | 5592831 | 5595486 | pyrimidodiazepine synthase |
| HOG17002 | chr02 | gene_2103_tr1 | 10452747 | 10461117 | chr16 | XP_011171031.2 | 5579776 | 5591916 | neprilysin-1 isoform X1 |
| HOG10333 | chr02 | gene_2104_tr1 | 10465357 | 10474892 | chr16 | XP_025988745.1 | 5561327 | 5577535 | neuroendocrine convertase 2 |
| HOG8681 | chr02 | gene_1495_tr1 | 2328488 | 2329466 | chr16 | XP_011164706.1 | 16076484 | 16078090 | glutaredoxin-3 |
| HOG15083 | chr02 | gene_1705_tr2 | 10480844 | 10486145 | chr16 | XP_011170978.2 | 5549025 | 5555577 | poly(A)-specific ribonuclease PARN |
| HOG13963 | chr02 | gene_1842_tr1 | 1373592 | 1375505 | chr16 | XP_025990729.1 | 24467483 | 24472448 | 2-amino-3-ketobutyrate coenzyme A ligase, mitochondrial |
| HOG11167 | chr02 | gene_1707_tr1 | 10532848 | 10535480 | chr16 | XP_025988731.1 | 5459740 | 5470399 | protein dimmed |
| HOG11706 | chr02 | gene_2107_tr1 | 10543095 | 10545256 | chr16 | XP_039314193.1 | 5451045 | 5456789 | cytoglobin-2 isoform X1 |
| HOG9702 | chr02 | gene_1708_tr3 | 10550379 | 10555232 | chr16 | XP_011170905.2 | 5441471 | 5447207 | probable ATP-dependent RNA helicase spindle-E |
| HOG9703 | chr02 | gene_2108_tr1 | 10556289 | 10558712 | chr16 | XP_011170895.1 | 5437682 | 5441278 | G protein pathway suppressor 2 |
| HOG9612 | chr02 | gene_1709_tr2 | 10559350 | 10561168 | chr16 | XP_011170885.1 | 5434001 | 5437202 | CDAN1-interacting nuclease 1 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| HOG8716 | chr02 | gene_1849_tr1 | 1621609 | 1624476 | chr16 | XP_011164610.1 | 15145489 | 15150217 | X-ray repair cross-complementing protein 5 |
| HOG5399 | chr02 | gene_1446_tr1 | 627083 | 630858 | chr16 | XP_025990761.1 | 22937162 | 22942889 | facilitated trehalose transporter Tret1 |
| HOG9610 | chr02 | gene_2109_tr2 | 10561716 | 10563143 | chr16 | XP_039315102.1 | 5431828 | 5434733 | protein FAM50 homolog isoform X1 |
| HOG9828 | chr02 | gene_1710_tr2 | 10565021 | 10568187 | chr16 | XP_011170858.2 | 5427830 | 5431731 | dipeptidyl peptidase 3 isoform X1 |
| HOG10089 | chr02 | gene_1711_tr1 | 10569137 | 10577391 | chr16 | XP_011170848.1 | 5414478 | 5426727 | DNA polymerase alpha subunit B |
| HOG15311 | chr02 | gene_1920_tr3 | 3192824 | 3209943 | chr16 | XP_039314111.1 | 9731785 | 9888219 | protein TANC2 isoform X1 |
| HOG8755 | chr02 | gene_1865_tr1 | 2138813 | 2167159 | chr16 | XP_025989484.1 | 16791784 | 16841925 | max dimerization protein 1 |
| HOG10090 | chr02 | gene_1712_tr4 | 10578026 | 10594794 | chr16 | XP_039315095.1 | 5384580 | 5413651 | steroid hormone receptor ERR1 isoform X1 |
| HOG8128 | chr02 | gene_1869_tr4 | 2208105 | 2212173 | chr16 | XP_011170070.1 | 16749661 | 16753897 | Hermansky-Pudlak syndrome 1 protein homolog isoform X1 |
| HOG10184 | chr02 | gene_2110_tr1 | 10626264 | 10709958 | chr16 | XP_025988733.1 | 4988454 | 5358114 | nephrin isoform X1 |
| HOG10187 | chr02 | gene_1715_tr5 | 10840933 | 10856578 | chr16 | XP_011170774.2 | 5086065 | 5108033 | glucose dehydrogenase [FAD, quinone] |
| HOG8775 | chr02 | gene_1928_tr2 | 3296805 | 3305510 | chr16 | XP_011167883.1 | 14694405 | 14905258 | complexin isoform X1 |
| HOG9419 | chr02 | gene_1491_tr1 | 2250023 | 2251767 | chr16 | XP_011164693.1 | 15993178 | 16000203 | serine/arginine-rich splicing factor 1A |
| HOG15512 | chr02 | gene_1563_tr1 | 3832533 | 3833843 | chr16 | XP_025990935.2 | 25674362 | 25682023 | proline-rich protein 2-like |
| HOG13903 | chr02 | gene_1822_tr4 | 312935 | 341739 | chr16 | XP_039314666.1 | 25943937 | 25999258 | glutamate-gated chloride channel isoform X1 |
| HOG7100 | chr02 | gene_1896_tr3 | 2681020 | 2684878 | chr16 | XP_011170134.2 | 17437430 | 17442751 | proclotting enzyme |
| HOG9823 | chr02 | gene_2111_tr1 | 10961119 | 10964816 | chr16 | XP_011170763.1 | 4962700 | 4967344 | ecdysteroid-regulated 16 kDa protein |
| HOG10512 | chr02 | gene_2112_tr4 | 10969108 | 10978324 | chr16 | XP_011170750.1 | 4946711 | 4958954 | glutathione hydrolase 1 proenzyme |
| HOG10513 | chr02 | gene_1716_tr2 | 10980484 | 10982061 | chr16 | XP_011170739.1 | 4941982 | 4944570 | proteasome subunit alpha type-7-1 |
| HOG10515 | chr02 | gene_2113_tr1 | 10982571 | 10983285 | chr16 | XP_011170726.1 | 4940621 | 4941608 | membrane magnesium transporter 1 |
| HOG8295 | chr02 | gene_1452_tr1 | 806294 | 808604 | chr16 | XP_011165153.1 | 23290091 | 23300429 | alpha-tocopherol transfer protein-like |
| HOG9112 | chr02 | gene_1717_tr5 | 10983859 | 10987057 | chr16 | XP_011170713.2 | 4936386 | 4940192 | nucleolar complex protein 3 homolog |
| HOG10399 | chr02 | gene_1718_tr1 | 10987471 | 10990276 | chr16 | XP_011170689.2 | 4932448 | 4936043 | protein VAC14 homolog |
| HOG10386 | chr02 | gene_2114_tr1 | 10990636 | 10993321 | chr16 | XP_011170700.1 | 4928882 | 4932535 | serine/threonine-protein kinase TBK1 |
| HOG11500 | chr02 | gene_1719_tr1 | 10995214 | 11005914 | chr16 | XP_011171195.1 | 5675531 | 5688936 | alpha-2-macroglobulin-like protein 1 isoform X1 |
| HOG15030 | chr02 | gene_1838_tr1 | 1282051 | 1283833 | chr16 | XP_011164926.1 | 24065197 | 24074475 | COMM domain-containing protein 4 |

| HOG8708 | chr02 | gene_1835_tr1 | 835166 | 835865 | chr16 | XP_011165066.1 | 23347111 | 23348626 | H/ACA ribonucleoprotein complex subunit 2-like protein |
|---|---|---|---|---|---|---|---|---|---|
| HOG3928 | chr02 | gene_1721_tr1 | 11015360 | 11016962 | chr16 | XP_011171132.2 | 5640676 | 5641436 | glutamyl-tRNA(Gln) amidotransferase subunit C, mitochondrial isoform X1 |
| HOG8493 | chr02 | gene_1496_tr5 | 2342986 | 2354028 | chr16 | XP_039314212.1 | 16078243 | 16112522 | large neutral amino acids transporter small subunit 1 isoform X1 |
| HOG14737 | chr02 | gene_1720_tr1 | 11015818 | 11016864 | chr16 | XP_011171146.2 | 5639561 | 5640843 | DNA polymerase subunit gamma-2, mitochondrial |
| HOG14776 | chr02 | gene_2116_tr1 | 11017727 | 11018575 | chr16 | XP_011171124.1 | 5638121 | 5639619 | ER lumen protein-retaining receptor |
| HOG7949 | chr02 | gene_1516_tr2 | 2767965 | 2789287 | chr16 | XP_039315007.1 | 17556002 | 17569874 | semaphorin-5B isoform X1 |
| HOG10436 | chr02 | gene_1722_tr1 | 11020586 | 11023822 | chr16 | XP_011171109.1 | 5623693 | 5628008 | amino acid transporter AVT1C |
| HOG7867 | chr02 | gene_1881_tr4 | 2434383 | 2435726 | chr16 | XP_025991950.1 | 16204890 | 16208661 | golgin subfamily A member 7 isoform X1 |
| HOG15278 | chr02 | gene_2118_tr3 | 11028891 | 11031772 | chr16 | XP_025988755.2 | 5617066 | 5621371 | modifier of mdg4 isoform X1 |
| HOG10278 | chr02 | gene_2120_tr1 | 11032945 | 11034474 | chr16 | XP_011171555.1 | 5614286 | 5616241 | protein EFR3 homolog B-like |
| HOG8684 | chr02 | gene_1886_tr1 | 2486426 | 2487312 | chr16 | XP_039315036.1 | 16349400 | 16351156 | U6 snRNA-associated Sm-like protein LSm8 |
| HOG10251 | chr02 | gene_2119_tr1 | 11035203 | 11043350 | chr16 | XP_011171085.2 | 5606080 | 5614138 | transcription initiation factor TFIID subunit 1 isoform X1 |
| HOG14252 | chr02 | gene_2121_tr1 | 11045460 | 11046213 | chr16 | XP_011171067.1 | 5602845 | 5604980 | small ubiquitin-related modifier 3-like |
| HOG10444 | chr02 | gene_1517_tr1 | 2807371 | 2815016 | chr16 | XP_025989549.2 | 17607390 | 17616003 | LOW QUALITY PROTEIN: cilia- and flagella-associated protein 43-like |
| HOG10356 | chr02 | gene_1443_tr1 | 602260 | 604541 | chr16 | XP_011165293.1 | 22892412 | 22897385 | serine protease HTRA2, mitochondrial isoform X1 |
| HOG11603 | chr02 | gene_1724_tr2 | 11049948 | 11053351 | chr16 | XP_011172210.1 | 8043481 | 8046799 | cyclin-dependent kinase 8 isoform X1 |
| HOG6111 | chr02 | gene_1448_tr1 | 776848 | 779010 | chr16 | XP_011165211.1 | 23182884 | 23189089 | zinc finger protein 699 |
| HOG8763 | chr02 | gene_1465_tr3 | 1557926 | 1560610 | chr16 | XP_025991920.1 | 15222779 | 15227463 | transcription initiation protein SPT3 homolog isoform X1 |
| HOG11601 | chr02 | gene_2123_tr1 | 11078830 | 11084005 | chr16 | XP_011172207.1 | 8010781 | 8018291 | ecdysone 20-monooxygenase isoform X1 |
| HOG16199 | chr02 | gene_1726_tr1 | 11103089 | 11103517 | chr16 | XP_011172205.1 | 7983972 | 7985364 | transmembrane protein 216 |
| HOG11604 | chr02 | gene_2125_tr1 | 11104769 | 11107047 | chr16 | XP_011172202.1 | 7980975 | 7984021 | adenosine 3'-phospho 5'-phosphosulfate transporter 1 |
| HOG16309 | chr02 | gene_1888_tr2 | 2509939 | 2511890 | chr16 | XP_011156722.1 | 24656092 | 24662689 | major facilitator superfamily domain-containing protein 10 isoform X1 |

| HOG9911 | chr02 | gene_1727_tr3 | 11112259 | 11121909 | chr16 | XP_011172199.1 | 7964222 | 7972625 | RNA-binding protein Nova-1 isoform X1 |
|---------|-------|---------------|----------|----------|-------|----------------|---------|---------|--------|
| HOG9918 | chr02 | gene_2126_tr2 | 11134124 | 11136733 | chr16 | XP_039314554.1 | 7947767 | 7953232 | probable maleylacetoacetate isomerase 2 isoform X1 |
| HOG13352 | chr02 | gene_1732_tr1 | 11169595 | 11172426 | chr16 | XP_011172324.1 | 6139353 | 6141960 | E3 ubiquitin-protein ligase RNF34 |
| HOG15689 | chr02 | gene_1731_tr1 | 11175106 | 11176182 | chr16 | XP_011172321.1 | 6131662 | 6133555 | vacuolar protein-sorting-associated protein 25 |
| HOG14348 | chr02 | gene_1515_tr1 | 2610166 | 2678821 | chr16 | XP_025989465.2 | 17347717 | 17438257 | atrial natriuretic peptide receptor 1 isoform X1 |
| HOG15119 | chr02 | gene_1480_tr1 | 2058079 | 2064661 | chr16 | XP_039315041.1 | 16954588 | 16968602 | sodium-coupled neutral amino acid transporter 9 isoform X1 |
| HOG16123 | chr02 | gene_2127_tr3 | 11177318 | 11178732 | chr16 | XP_011172322.1 | 6128726 | 6131665 | cyclin-dependent kinases regulatory subunit |

**Supplementary Table 3:** List of many-to-many orthologous genes with functional annotation found on *Solenopsis* chromosome 16 and Cataglyphis chromosome 2. The highlighted cells indicate the genes found in the supergene region of the respective species.

| HOG | Cnig chrm | Cnig_id | Cnig_start | Cnig_end | Sinv chrm | Sinv_id | Sinv_start | Sinv_end | Sinv_function |
|---|---|---|---|---|---|---|---|---|---|
| HOG8698 | chr02 | gene_1641_tr1 | 8325068 | 8328804 | chr16 | XP_011164607.1 | 15136017 | 15140756 | melanotransferrin isoform X1 |
| HOG8698 | chr02 | gene_2023_tr1 | 8491765 | 8495501 | chr16 | XP_011164607.1 | 15136017 | 15140756 | melanotransferrin isoform X1 |
| HOG8827 | chr02 | gene_1589_tr1 | 5045800 | 5046548 | chr16 | XP_011161724.2 | 13267097 | 13271493 | probable E3 ubiquitin-protein ligase RHY1A |
| HOG15688 | chr02 | gene_1695_tr1 | 10162993 | 1E+07 | chr16 | XP_011170189.2 | 3995281 | 3997294 | cytoplasmic tRNA 2-thiolation protein 2 |
| HOG14644 | chr02 | gene_2041_tr2 | 8858673 | 8861002 | chr16 | XP_039314676.1 | 26004603 | 26021597 | flavin-containing monooxygenase FMO GS-OX-like 2 isoform X1 |
| HOG14988 | chr02 | gene_1922_tr1 | 3230128 | 3231288 | chr16 | XP_011164983.1 | 9360203 | 9363557 | DDRGK domain-containing protein 1 isoform X1 |
| HOG3816 | chr02 | gene_1894_tr1 | 2558786 | 2559621 | chr16 | XP_011156749.1 | 24916104 | 24918107 | histone RNA hairpin-binding protein |
| HOG8963 | chr02 | gene_1537_tr1 | 3150294 | 3154642 | chr16 | XP_039314572.1 | 12972095 | 13055442 | coiled-coil domain-containing protein R3HCC1L isoform X1 |
| HOG10455 | chr02 | gene_1510_tr1 | 2549965 | 2551574 | chr16 | XP_039315055.1 | 24927324 | 24931095 | S-adenosylmethionine mitochondrial carrier protein |
| HOG13778 | chr02 | gene_1540_tr2 | 3237179 | 3238657 | chr16 | XP_039314532.1 | 12063131 | 12112267 | DNA excision repair protein ERCC-1 isoform X1 |
| HOG5528 | chr02 | gene_1676_tr1 | 9808188 | 9810721 | chr16 | XP_011174500.3 | 21993059 | 21997112 | odorant receptor 22c isoform X1 |
| HOG9890 | chr02 | gene_1852_tr1 | 1685469 | 1700678 | chr16 | XP_039314798.1 | 15844676 | 15869563 | ABC transporter G family member 23 isoform X1 |
| HOG15358 | chr02 | gene_1930_tr1 | 3427464 | 3429176 | chr16 | XP_011167880.1 | 14926676 | 14929867 | pre-mRNA-splicing factor Slu7 |
| HOG8187 | chr02 | gene_1597_tr1 | 5443206 | 5533951 | chr16 | XP_011155884.1 | 21285578 | 21397783 | segmentation protein cap'n'collar isoform X1 |
| HOG9148 | chr02 | gene_1858_tr1 | 2051611 | 2054176 | chr16 | XP_011170086.1 | 16968597 | 16976194 | phosphoglucomutase isoform X1 |
| HOG8353 | chr02 | gene_2019_tr1 | 8388278 | 8391469 | chr16 | XP_039314831.1 | 24695207 | 24810271 | RNA-binding protein Musashi homolog 2 isoform X1 |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| HOG8606 | chr02 | gene_2075_tr1 | 9772677 | 9774527 | chr16 | XP_011174441.1 | 22017324 | 22019290 | RCC1 domain-containing protein 1 |
| HOG15068 | chr02 | gene_1569_tr1 | 3988663 | 3991066 | chr16 | XP_025990993.1 | 25495806 | 25500737 | ras-related protein Rab-27A |
| HOG13962 | chr02 | gene_1843_tr1 | 1377153 | 1379481 | chr16 | XP_011166565.2 | 24474013 | 24478058 | nicalin |
| HOG17165 | chr02 | gene_2017_tr1 | 8345501 | 8348747 | chr16 | XP_011156735.1 | 24833864 | 24841797 | fibrillin-1 isoform X1 |
| HOG17165 | chr02 | gene_1650_tr1 | 8508193 | 8511442 | chr16 | XP_011156735.1 | 24833864 | 24841797 | fibrillin-1 isoform X1 |
| HOG8446 | chr02 | gene_1844_tr1 | 1518193 | 1532996 | chr16 | XP_011164628.1 | 15263526 | 15305312 | homeotic protein empty spiracles |
| HOG9085 | chr02 | gene_1447_tr2 | 771890 | 772300 | chr16 | XP_011165219.2 | 23177534 | 23178507 | mitochondrial import inner membrane translocase subunit Tim16 |
| HOG8563 | chr02 | gene_1619_tr2 | 7659741 | 7663127 | chr16 | XP_039315141.1 | 18807749 | 18837105 | protein phosphatase 1 regulatory subunit 3C-B |
| HOG7911 | chr02 | gene_1867_tr1 | 2175099 | 2181479 | chr16 | XP_011170059.1 | 16779865 | 16787958 | protein FAM160B1 isoform X1 |
| HOG15320 | chr02 | gene_1538_tr1 | 3187547 | 3188773 | chr16 | XP_025996242.1 | 10068516 | 10130643 | tubulin polymerization-promoting protein homolog isoform X1 |
| HOG10354 | chr02 | gene_1442_tr2 | 598774 | 600601 | chr16 | XP_011165314.2 | 22885865 | 22888640 | rRNA N6-adenosine-methyltransferase ZCCHC4 |
| HOG9918 | chr02 | gene_2126_tr2 | 11134124 | 1.1E+07 | chr16 | XP_039314554.1 | 7947767 | 7953232 | probable maleylacetoacetate isomerase 2 isoform X1 |
| HOG15278 | chr02 | gene_2118_tr3 | 11028891 | 1.1E+07 | chr16 | XP_025988755.2 | 5617066 | 5621371 | modifier of mdg4 isoform X1 |
| HOG13352 | chr02 | gene_1732_tr1 | 11169595 | 1.1E+07 | chr16 | XP_011172324.1 | 6139353 | 6141960 | E3 ubiquitin-protein ligase RNF34 |
| HOG8813 | chr02 | gene_1905_tr2 | 2847105 | 2848201 | chr16 | XP_011175190.1 | 17717140 | 17718878 | charged multivesicular body protein 5 |
| HOG10946 | chr02 | gene_1501_tr2 | 2452175 | 2454290 | chr16 | XP_011170021.1 | 16312880 | 16315978 | TGF-beta-activated kinase 1 and MAP3K7-binding protein 1 |
| HOG10399 | chr02 | gene_1718_tr1 | 10987471 | 1.1E+07 | chr16 | XP_011170689.2 | 4932448 | 4936043 | protein VAC14 homolog |
| HOG8106 | chr02 | gene_1994_tr2 | 7468244 | 7472789 | chr16 | XP_025987315.1 | 19079404 | 19084961 | transcription elongation factor SPT5 |
| HOG10090 | chr02 | gene_1712_tr4 | 10578026 | 1.1E+07 | chr16 | XP_039315095.1 | 5384580 | 5413651 | steroid hormone receptor ERR1 isoform X1 |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| HOG8215 | chr02 | gene_1504_tr1 | 2476545 | 2484677 | chr16 | XP_039315026.1 | 16336210 | 16350025 | LOW QUALITY PROTEIN: serine/threonine-protein phosphatase 1 regulatory subunit 10 |
| HOG12500 | chr02 | gene_2072_tr1 | 9281188 | 9282321 | chr16 | XP_011165270.1 | 26160306 | 26162897 | BLOC-1-related complex subunit 5 |
| HOG8214 | chr02 | gene_1870_tr2 | 2218820 | 2226623 | chr16 | XP_039314336.1 | 24529290 | 24544420 | transferrin |
| HOG8450 | chr02 | gene_2056_tr1 | 9170851 | 9171782 | chr16 | XP_011174382.1 | 22513988 | 22515516 | probable small nuclear ribonucleoprotein G |
| HOG7892 | chr02 | gene_1987_tr2 | 7211682 | 7216161 | chr16 | XP_011157636.1 | 19401407 | 19409904 | modular serine protease-like |
| HOG7892 | chr02 | gene_1987_tr2 | 7211682 | 7216161 | chr16 | XP_025987236.2 | 19394578 | 19400439 | modular serine protease |
| HOG7892 | chr02 | gene_1987_tr2 | 7211682 | 7216161 | chr16 | XP_025987238.1 | 19387374 | 19394186 | clotting factor C isoform X1 |
| HOG8673 | chr02 | gene_1594_tr1 | 5111114 | 5144049 | chr16 | XP_025987255.1 | 21751143 | 21823735 | trace amine-associated receptor 9 |
| HOG14737 | chr02 | gene_1720_tr1 | 11015818 | 1.1E+07 | chr16 | XP_011171146.2 | 5639561 | 5640843 | DNA polymerase subunit gamma-2 mitochondrial |
| HOG15067 | chr02 | gene_1575_tr7 | 4520673 | 4523032 | chr16 | XP_025989498.1 | 16702011 | 16707132 | aldo-keto reductase family 1 member B1 |
| HOG10327 | chr02 | gene_2094_tr1 | 10201899 | 1E+07 | chr16 | XP_011155529.1 | 7919926 | 7939780 | serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit epsilon isoform |
| HOG8607 | chr02 | gene_1674_tr2 | 9776722 | 9779510 | chr16 | XP_011174433.1 | 22014511 | 22017393 | ADP-ribosylation factor-like protein 3 |
| HOG14573 | chr02 | gene_1859_tr1 | 2065162 | 2066044 | chr16 | XP_011170087.1 | 16952983 | 16954745 | ATP synthase subunit C lysine N-methyltransferase |
| HOG9902 | chr02 | gene_1624_tr1 | 7773176 | 7775233 | chr16 | XP_025994002.2 | 18608344 | 18615214 | E3 ubiquitin-protein ligase COP1 isoform X1 |
| HOG8077 | chr02 | gene_1502_tr1 | 2465955 | 2468779 | chr16 | XP_011170022.1 | 16326307 | 16330617 | V-type proton ATPase subunit B |
| HOG9408 | chr02 | gene_2054_tr1 | 9057622 | 9062025 | chr16 | XP_011161961.1 | 14555366 | 14562042 | vacuolar protein sorting-associated protein 51 homolog |
| HOG15232 | chr02 | gene_1932_tr1 | 3463774 | 3465313 | chr16 | XP_025993452.2 | 14979589 | 14984897 | TBC1 domain family member 7 isoform X1 |
| HOG15520 | chr02 | gene_1462_tr3 | 1350205 | 1352805 | chr16 | XP_011169804.1 | 24221257 | 24293784 | tachykinins |

| HOG15226 | chr02 | gene_2101_tr2 | 10418561 | 1E+07 | chr16 | XP_011172444.1 | 6856005 | 6866241 | protein lingerer isoform X1 |
|---|---|---|---|---|---|---|---|---|---|
| HOG8357 | chr02 | gene_1645_tr1 | 8354286 | 8354759 | chr16 | XP_011156726.2 | 24827528 | 24828835 | 39S ribosomal protein L51 mitochondrial |
| HOG8357 | chr02 | gene_2025_tr1 | 8502200 | 8502673 | chr16 | XP_011156726.2 | 24827528 | 24828835 | 39S ribosomal protein L51 mitochondrial |
| HOG15356 | chr02 | gene_1887_tr7 | 2500497 | 2507595 | chr16 | XP_011156723.1 | 24664117 | 24678278 | signal transducer and activator of transcription 5B |
| HOG13784 | chr02 | gene_2051_tr4 | 8974048 | 8975733 | chr16 | XP_011161972.1 | 14420379 | 14426100 | CDGSH iron-sulfur domain-containing protein 2 homolog |
| HOG17057 | chr02 | gene_1440_tr1 | 577319 | 581594 | chr16 | XP_011165379.1 | 22852621 | 22858466 | coiled-coil domain-containing protein 142 isoform X1 |
| HOG8794 | chr02 | gene_1904_tr2 | 2840928 | 2843949 | chr16 | XP_011175214.1 | 17709621 | 17713021 | nuclear envelope integral membrane protein 1 isoform X1 |
| HOG8492 | chr02 | gene_1497_tr1 | 2394239 | 2399165 | chr16 | XP_011164711.1 | 16158349 | 16165729 | lysyl oxidase homolog 3 |
| HOG8964 | chr02 | gene_1916_tr1 | 3156807 | 3159565 | chr16 | XP_011166899.1 | 13059468 | 13063879 | signal recognition particle subunit SRP72 |
| HOG15083 | chr02 | gene_1705_tr2 | 10480844 | 1E+07 | chr16 | XP_011170978.2 | 5549025 | 5555577 | poly(A)-specific ribonuclease PARN |
| HOG15025 | chr02 | gene_1458_tr2 | 1247917 | 1274925 | chr16 | XP_039315079.1 | 23900432 | 24065116 | protein tincar isoform X1 |
| HOG7898 | chr02 | gene_1988_tr2 | 7216732 | 7219024 | chr16 | XP_011157570.1 | 19367690 | 19374112 | tRNA pseudouridine synthase A isoform X1 |
| HOG10640 | chr02 | gene_1524_tr2 | 2991447 | 3001477 | chr16 | XP_011170093.1 | 16977851 | 16988129 | huntingtin isoform X1 |
| HOG10453 | chr02 | gene_1885_tr1 | 2458091 | 2464714 | chr16 | XP_011170017.1 | 16318164 | 16325215 | 4-hydroxybutyrate coenzyme A transferase isoform X1 |
| HOG7890 | chr02 | gene_1634_tr1 | 7848863 | 7852708 | chr16 | XP_011159689.1 | 18489958 | 18494590 | ATP-dependent RNA helicase SUV3 homolog mitochondrial |
| HOG10670 | chr02 | gene_1436_tr4 | 555219 | 560423 | chr16 | XP_011174290.1 | 22795556 | 22799567 | casein kinase I isoform X1 |
| HOG8352 | chr02 | gene_2018_tr1 | 8355667 | 8358937 | chr16 | XP_011156728.1 | 24820475 | 24827458 | dipeptidyl peptidase 9 |
| HOG8352 | chr02 | gene_1649_tr1 | 8496401 | 8501292 | chr16 | XP_011156728.1 | 24820475 | 24827458 | dipeptidyl peptidase 9 |
| HOG10436 | chr02 | gene_1722_tr1 | 11020586 | 1.1E+07 | chr16 | XP_011171109.1 | 5623693 | 5628008 | amino acid transporter AVT1C |
| HOG9022 | chr02 | gene_2014_tr1 | 8320370 | 8321475 | chr16 | XP_011175161.1 | 17889490 | 17891678 | peroxisomal membrane protein 11C |
| HOG13826 | chr02 | gene_1526_tr2 | 3021334 | 3025332 | chr16 | XP_011170099.1 | 17016110 | 17024394 | endoplasmic reticulum metallopeptidase 1 isoform X1 |
| HOG15085 | chr02 | gene_1939_tr1 | 4056694 | 4063002 | chr16 | XP_025990986.1 | 25336872 | 25415644 | protein artichoke isoform X1 |

| HOG8210 | chr02 | gene_1486_tr2 | 2212659 | 2217507 | chr16 | XP_011156707.1 | 24545232 | 24553641 | cytoplasmic dynein 2 intermediate chain 1 isoform X1 |
|---|---|---|---|---|---|---|---|---|---|
| HOG13636 | chr02 | gene_1616_tr1 | 7553722 | 7607336 | chr16 | XP_039314493.1 | 18886118 | 19024272 | potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 3 isoform X1 |
| HOG8665 | chr02 | gene_1906_tr2 | 3009188 | 3013281 | chr16 | XP_011170095.1 | 16992073 | 17009406 | probable sodium/potassium/calcium exchanger CG1090 |
| HOG8671 | chr02 | gene_1570_tr1 | 4414120 | 4420669 | chr16 | XP_011156769.2 | 24995645 | 25088183 | protein artichoke |
| HOG16969 | chr02 | gene_1603_tr3 | 6164963 | 6195172 | chr16 | XP_039314822.1 | 20496762 | 20533250 | Kv channel-interacting protein 4 isoform X1 |
| HOG8801 | chr02 | gene_1633_tr3 | 7839020 | 7842392 | chr16 | XP_011159621.1 | 18504596 | 18511068 | mitoferrin-1 isoform X1 |
| HOG8266 | chr02 | gene_1929_tr2 | 3421778 | 3426438 | chr16 | XP_039314407.1 | 14909767 | 14922009 | activin receptor type-2A isoform X1 |
| HOG13656 | chr02 | gene_1463_tr1 | 1358112 | 1362142 | chr16 | XP_011166559.1 | 24437663 | 24446348 | trimeric intracellular cation channel type 1B.1 |
| HOG8532 | chr02 | gene_2048_tr1 | 8934105 | 8942741 | chr16 | XP_039314299.1 | 9393467 | 9416297 | protein phosphatase PP2A 55 kDa regulatory subunit isoform X1 |
| HOG9131 | chr02 | gene_2010_tr1 | 7871199 | 7874674 | chr16 | XP_011159599.3 | 18461337 | 18466462 | phenoloxidase-activating factor 2 |
| HOG4469 | chr02 | gene_1439_tr4 | 582905 | 586363 | chr16 | XP_025990753.1 | 22858712 | 22865492 | peroxidase |
| HOG8118 | chr02 | gene_1596_tr1 | 5425828 | 5435317 | chr16 | XP_011155888.2 | 21405290 | 21422491 | endoplasmic reticulum aminopeptidase 1 isoform X1 |
| HOG8721 | chr02 | gene_1972_tr1 | 6201764 | 6203381 | chr16 | XP_011158029.1 | 20490961 | 20493299 | protein farnesyltransferase/geranylgeranyltransferase type-1 subunit alpha |
| HOG9823 | chr02 | gene_2111_tr1 | 10961119 | 1.1E+07 | chr16 | XP_011170763.1 | 4962700 | 4967344 | ecdysteroid-regulated 16 kDa protein |
| HOG8533 | chr02 | gene_1980_tr1 | 6744707 | 6759773 | chr16 | XP_011157873.1 | 19917347 | 19933877 | SEC23-interacting protein isoform X1 |
| HOG10187 | chr02 | gene_1715_tr5 | 10840933 | 1.1E+07 | chr16 | XP_011170774.2 | 5086065 | 5108033 | glucose dehydrogenase [FAD quinone] |

| HOG ID | Chr | Gene | Start | End | Chr | Protein | Start | End | Description |
|---|---|---|---|---|---|---|---|---|---|
| HOG8499 | chr02 | gene_1925_tr1 | 3255213 | 3256394 | chr16 | XP_025993462.1 | 14621720 | 14625536 | charged multivesicular body protein 3 |
| HOG13861 | chr02 | gene_2033_tr1 | 8709815 | 8752864 | chr16 | XP_039314262.1 | 17113859 | 17170291 | microtubule-associated protein futsch isoform X1 |
| HOG8376 | chr02 | gene_1873_tr1 | 2264436 | 2284940 | chr16 | XP_039314194.1 | 16006971 | 16033707 | MAP kinase-activating death domain protein isoform X1 |
| HOG3995 | chr02 | gene_2044_tr1 | 8894630 | 8894947 | chr16 | XP_039314683.1 | 26062324 | 26064224 | oxidoreductase-like domain-containing protein 1 |
| HOG10513 | chr02 | gene_1716_tr2 | 10980484 | 1.1E+07 | chr16 | XP_011170739.1 | 4941982 | 4944570 | proteasome subunit alpha type-7-1 |
| HOG8591 | chr02 | gene_1935_tr1 | 3671398 | 3672611 | chr16 | XP_011156801.1 | 25865185 | 25871673 | transmembrane protein 230 isoform X1 |
| HOG16016 | chr02 | gene_1855_tr1 | 1799371 | 1801890 | chr16 | XP_011164652.2 | 15717500 | 15720868 | arrestin homolog |
| HOG16856 | chr02 | gene_1455_tr1 | 855531 | 877697 | chr16 | XP_039314720.1 | 23381477 | 23411042 | dynein heavy chain 8 axonemal |
| HOG15228 | chr02 | gene_1546_tr4 | 3458611 | 3461401 | chr16 | XP_039314410.1 | 14958604 | 14972956 | hippocampus abundant transcript 1 protein isoform X1 |
| HOG7851 | chr02 | gene_1469_tr1 | 1630955 | 1636315 | chr16 | XP_039314800.1 | 15983474 | 15991932 | catenin alpha isoform X1 |
| HOG7957 | chr02 | gene_1900_tr3 | 2796458 | 2800017 | chr16 | XP_039314811.1 | 18571136 | 18574460 | LOW QUALITY PROTEIN: leucine-rich repeat-containing protein 15-like |
| HOG11444 | chr02 | gene_1548_tr2 | 3466345 | 3467899 | chr16 | XP_011167869.1 | 14972134 | 14977920 | methyltransferase-like protein 25 |
| HOG11151 | chr02 | gene_1882_tr1 | 2445602 | 2447802 | chr16 | XP_025989556.1 | 16306420 | 16308828 | titin |
| HOG16017 | chr02 | gene_1853_tr1 | 1733138 | 1733878 | chr16 | XP_011164666.1 | 15796782 | 15799883 | copper homeostasis protein cutC homolog |
| HOG7968 | chr02 | gene_1889_tr1 | 2536551 | 2541142 | chr16 | XP_025990923.1 | 24608618 | 24616794 | SWI/SNF complex subunit SMARCC2 |
| HOG16199 | chr02 | gene_1726_tr1 | 11103089 | 1.1E+07 | chr16 | XP_011172205.1 | 7983972 | 7985364 | transmembrane protein 216 |
| HOG11098 | chr02 | gene_1432_tr1 | 366596 | 369341 | chr16 | XP_011174335.1 | 22583577 | 22588088 | vesicular integral-membrane protein VIP36 |
| HOG14511 | chr02 | gene_1703_tr2 | 10430301 | 1E+07 | chr16 | XP_011171054.1 | 5595847 | 5602703 | mediator of RNA polymerase II transcription subunit 23 |
| HOG17002 | chr02 | gene_2103_tr1 | 10452747 | 1E+07 | chr16 | XP_011171031.2 | 5579776 | 5591916 | neprilysin-1 isoform X1 |
| HOG3972 | chr02 | gene_1573_tr1 | 4492366 | 4493178 | chr16 | XP_039314761.1 | 24558695 | 24560167 | nucleolin-like |
| HOG7857 | chr02 | gene_1847_tr3 | 1601008 | 1605352 | chr16 | XP_011164616.2 | 15167201 | 15175918 | pickpocket protein 28 |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| HOG15089 | chr02 | gene_1481_tr1 | 2066442 | 2067463 | chr16 | XP_011170090.1 | 16951061 | 16953049 | ribulose-phosphate 3-epimerase |
| HOG9436 | chr02 | gene_1636_tr2 | 8259469 | 8285211 | chr16 | XP_039315011.1 | 17954582 | 18255677 | probable JmjC domain-containing histone demethylation protein 2C isoform X1 |
| HOG7302 | chr02 | gene_1943_tr1 | 4481091 | 4484240 | chr16 | XP_039314108.1 | 24577632 | 24585030 | cell death specification protein 2 |
| HOG10999 | chr02 | gene_2029_tr1 | 8542888 | 8547976 | chr16 | XP_011170131.1 | 17328625 | 17335654 | fumarylacetoacetate hydrolase domain-containing protein 2A isoform X1 |
| HOG7964 | chr02 | gene_1898_tr2 | 2790895 | 2794896 | chr16 | XP_039315010.1 | 17568536 | 17574944 | cysteine sulfinic acid decarboxylase |
| HOG8217 | chr02 | gene_1488_tr1 | 2235039 | 2237092 | chr16 | XP_011164632.1 | 15448658 | 15451149 | F-box/WD repeat-containing protein 4 isoform X1 |
| HOG8129 | chr02 | gene_1831_tr4 | 748263 | 770696 | chr16 | XP_011165243.1 | 23132897 | 23176944 | attractin-like protein 1 isoform X1 |
| HOG9612 | chr02 | gene_1709_tr2 | 10559350 | 1.1E+07 | chr16 | XP_011170885.1 | 5434001 | 5437202 | CDAN1-interacting nuclease 1 |
| HOG8472 | chr02 | gene_1535_tr2 | 3108146 | 3134625 | chr16 | XP_025994936.2 | 12436327 | 12627257 | protein lap4 isoform X1 |
| HOG8707 | chr02 | gene_1453_tr1 | 822789 | 828821 | chr16 | XP_011165091.1 | 23303851 | 23328875 | rho-related BTB domain-containing protein 1 isoform X1 |
| HOG8759 | chr02 | gene_1466_tr1 | 1589113 | 1593117 | chr16 | XP_039315121.1 | 15179363 | 15192867 | zinc finger protein 260 isoform X1 |
| HOG9836 | chr02 | gene_1605_tr1 | 6783211 | 6810424 | chr16 | XP_039314308.1 | 19862587 | 19891241 | protein unc-13 homolog 4B isoform X1 |
| HOG9418 | chr02 | gene_1872_tr6 | 2259064 | 2261708 | chr16 | XP_025991932.1 | 16004175 | 16006757 | farnesol dehydrogenase isoform X1 |
| HOG8988 | chr02 | gene_2011_tr2 | 7880423 | 7884430 | chr16 | XP_011159578.1 | 18451895 | 18460427 | nanos homolog 3 |
| HOG8988 | chr02 | gene_2011_tr2 | 7880423 | 7884430 | chr16 | XP_011159747.3 | 18447780 | 18449962 | nanos homolog 3-like |
| HOG10889 | chr02 | gene_1435_tr3 | 544379 | 548693 | chr16 | XP_025987290.1 | 22671772 | 22789697 | dopamine receptor 1 isoform X2 |
| HOG10326 | chr02 | gene_2095_tr1 | 10225309 | 1E+07 | chr16 | XP_011155524.2 | 7905195 | 7910729 | maltase 1 isoform X1 |
| HOG4475 | chr02 | gene_1864_tr3 | 2116467 | 2124441 | chr16 | XP_039314456.1 | 16843476 | 16854742 | titin homolog isoform X5 |
| HOG10645 | chr02 | gene_1441_tr4 | 590415 | 596041 | chr16 | XP_025990760.1 | 22873090 | 22882736 | phosphatidylinositide phosphatase SAC2 isoform X1 |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| HOG8589 | chr02 | gene_1561_tr1 | 3810773 | 3815356 | chr16 | XP_025990962.1 | 25692531 | 25843341 | AT-rich interactive domain-containing protein 5B isoform X1 |
| HOG7954 | chr02 | gene_1470_tr1 | 1644698 | 1667692 | chr16 | XP_011164684.1 | 15942310 | 15976783 | BCL2/adenovirus E1B 19 kDa protein-interacting protein 3 |
| HOG9968 | chr02 | gene_2002_tr3 | 7783173 | 7786678 | chr16 | XP_025994005.1 | 18597564 | 18603832 | mediator of RNA polymerase II transcription subunit 28 |
| HOG9821 | chr02 | gene_1903_tr1 | 2831582 | 2836162 | chr16 | XP_011175376.2 | 17700592 | 17707876 | probable glutamate receptor isoform X1 |
| HOG9821 | chr02 | gene_1903_tr1 | 2831582 | 2836162 | chr16 | XP_025989474.2 | 17628474 | 17634977 | glutamate receptor ionotropic delta-1 isoform X1 |
| HOG9821 | chr02 | gene_1903_tr1 | 2831582 | 2836162 | chr16 | XP_039314129.1 | 17693584 | 17699958 | glutamate receptor-like |
| HOG11158 | chr02 | gene_1655_tr3 | 8557444 | 8592647 | chr16 | XP_025989546.2 | 17289159 | 17326197 | MICOS complex subunit Mic60 isoform X1 |
| HOG10644 | chr02 | gene_1828_tr2 | 561841 | 570185 | chr16 | XP_011174263.3 | 22802774 | 22813157 | dynactin subunit 1 isoform X1 |
| HOG14655 | chr02 | gene_2068_tr2 | 9260339 | 9263319 | chr16 | XP_039314678.1 | 26000887 | 26005049 | trypsin isoform X1 |
| HOG9816 | chr02 | gene_1518_tr1 | 2817961 | 2822157 | chr16 | NP_001291533.1 | 17618379 | 17624016 | transferrin precursor |
| HOG8390 | chr02 | gene_1836_tr2 | 838903 | 853826 | chr16 | XP_039314181.1 | 23361028 | 23379858 | vacuolar protein sorting-associated protein 13B |
| HOG8364 | chr02 | gene_1974_tr1 | 6210501 | 6218674 | chr16 | XP_025987284.1 | 20473483 | 20484396 | peroxidasin homolog pxn-2 |
| HOG8680 | chr02 | gene_1493_tr1 | 2305670 | 2309668 | chr16 | XP_039314217.1 | 16050663 | 16058164 | N-acetyl-D-glucosamine kinase isoform X1 |
| HOG16940 | chr02 | gene_1908_tr2 | 3054063 | 3063955 | chr16 | XP_025989515.1 | 17054056 | 17067361 | atrial natriuretic peptide receptor 1 isoform X1 |
| HOG10676 | chr02 | gene_1850_tr1 | 1627454 | 1628788 | chr16 | XP_011164612.1 | 15141050 | 15143147 | beta-14-galactosyltransferase 7 |
| HOG8666 | chr02 | gene_1907_tr2 | 3018332 | 3019390 | chr16 | XP_011170096.2 | 17010831 | 17012794 | tetratricopeptide repeat protein 1 |
| HOG8474 | chr02 | gene_1468_tr2 | 1625291 | 1627027 | chr16 | XP_011164611.1 | 15142997 | 15145344 | alpha-2-macroglobulin receptor-associated protein |
| HOG8551 | chr02 | gene_1618_tr1 | 7636416 | 7641713 | chr16 | XP_011155778.1 | 18840151 | 18849161 | poly [ADP-ribose] polymerase tankyrase isoform X1 |
| HOG8674 | chr02 | gene_1961_tr4 | 5394807 | 5397857 | chr16 | XP_011155894.2 | 21468490 | 21472522 | ferrochelatase mitochondrial |
| HOG6394 | chr02 | gene_1461_tr1 | 1292632 | 1295337 | chr16 | XP_011169823.1 | 24347395 | 24355387 | UDP-glycosyltransferase UGT5 isoform X1 |

| HOG6394 | chr02 | gene_1461_tr1 | 1292632 | 1295337 | chr16 | XP_039314135.1 | 24355716 | 24361476 | UDP-glycosyltransferase UGT5-like |
|---------|-------|---------------|---------|---------|-------|----------------|----------|----------|----------------------------------|
| HOG6394 | chr02 | gene_1460_tr2 | 1295741 | 1300031 | chr16 | XP_011169823.1 | 24347395 | 24355387 | UDP-glycosyltransferase UGT5 isoform X1 |
| HOG6394 | chr02 | gene_1460_tr2 | 1295741 | 1300031 | chr16 | XP_039314135.1 | 24355716 | 24361476 | UDP-glycosyltransferase UGT5-like |
| HOG8001 | chr02 | gene_1500_tr3 | 2421470 | 2427974 | chr16 | XP_025991945.2 | 16188502 | 16201893 | cGMP-specific 3'5'-cyclic phosphodiesterase isoform X1 |
| HOG5269 | chr02 | gene_1997_tr3 | 7671709 | 7673861 | chr16 | XP_011155770.1 | 18804054 | 18807740 | serine protease snake |
| HOG7789 | chr02 | gene_1942_tr1 | 4466584 | 4467522 | chr16 | XP_011156758.1 | 24940738 | 24942742 | ELMO domain-containing protein 2 |
| HOG7835 | chr02 | gene_1509_tr1 | 2545617 | 2546126 | chr16 | XP_011156760.1 | 24934835 | 24938580 | high mobility group protein HMGI-C |
| HOG15092 | chr02 | gene_1948_tr4 | 4566904 | 4596170 | chr16 | XP_025989494.1 | 16575605 | 16659139 | tachykinin-like peptides receptor 99D isoform X1 |
| HOG15304 | chr02 | gene_1568_tr1 | 3983445 | 3986702 | chr16 | XP_025990992.2 | 25505722 | 25510892 | molybdenum cofactor sulfurase 3 |
| HOG5340 | chr02 | gene_1485_tr2 | 2203039 | 2206996 | chr16 | XP_011170067.1 | 16753719 | 16784018 | BRISC complex subunit FAM175B |
| HOG10009 | chr02 | gene_1830_tr3 | 633905 | 642281 | chr16 | XP_025990751.1 | 22946890 | 23121595 | octopamine receptor Oamb isoform X1 |
| HOG8785 | chr02 | gene_1521_tr1 | 2844452 | 2846734 | chr16 | XP_011175179.1 | 17714742 | 17717154 | tubulin epsilon chain |
| HOG8817 | chr02 | gene_1601_tr1 | 5870774 | 5872329 | chr16 | XP_039314709.1 | 20850242 | 20864404 | glutamine amidotransferase-like class 1 domain-containing protein 3A mitochondrial |
| HOG8826 | chr02 | gene_1958_tr1 | 5052175 | 5054200 | chr16 | XP_039314418.1 | 13289459 | 13298659 | UPF0687 protein C20orf27 homolog |
| HOG6288 | chr02 | gene_1977_tr1 | 6679051 | 6679895 | chr16 | XP_011157914.1 | 20000044 | 20001599 | peritrophin-1 |
| HOG10333 | chr02 | gene_2104_tr1 | 10465357 | 1E+07 | chr16 | XP_025988745.1 | 5561327 | 5577535 | neuroendocrine convertase 2 |
| HOG10675 | chr02 | gene_1438_tr1 | 587446 | 588814 | chr16 | XP_011165354.1 | 22866931 | 22869701 | ubiquitin-like domain-containing CTD phosphatase 1 |
| HOG7970 | chr02 | gene_1454_tr1 | 836500 | 838102 | chr16 | XP_011165023.1 | 23358829 | 23361307 | RING finger protein 37 isoform X1 |
| HOG13937 | chr02 | gene_1528_tr1 | 3048276 | 3053169 | chr16 | XP_025989522.1 | 17046676 | 17054767 | cysteine and histidine-rich protein 1 homolog |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| HOG8965 | chr02 | gene_1895_tr1 | 2563359 | 2569774 | chr16 | XP_011156748.1 | 24904560 | 24912249 | KAT8 regulatory NSL complex subunit 3 |
| HOG13454 | chr02 | gene_1657_tr3 | 8679162 | 8687330 | chr16 | XP_011170127.1 | 17185335 | 17196261 | sorting nexin-27 |
| HOG10661 | chr02 | gene_1985_tr2 | 7165636 | 7171653 | chr16 | XP_039314998.1 | 19419827 | 19467597 | tyramine receptor 1 isoform X1 |
| HOG8571 | chr02 | gene_1998_tr1 | 7683628 | 7685506 | chr16 | XP_011155767.1 | 18738035 | 18741095 | interleukin enhancer-binding factor 2 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_011165193.2 | 23247076 | 23251744 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_011169811.1 | 24137908 | 24153472 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_025987297.2 | 19488326 | 19494179 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_025989496.1 | 16682503 | 16691437 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_025990768.2 | 24206933 | 24215136 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_025990771.2 | 24193882 | 24203616 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_025990773.2 | 24159797 | 24171994 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_025990783.2 | 23266922 | 23272760 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_039314235.1 | 22843883 | 22852406 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_039314243.1 | 22832891 | 22838176 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_039314247.1 | 22825641 | 22832587 | cytochrome P450 4C1-like |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_039314988.1 | 23253576 | 23261512 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1578_tr1 | 4529864 | 4534034 | chr16 | XP_039315059.1 | 16669969 | 16679811 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_011165193.2 | 23247076 | 23251744 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_011169811.1 | 24137908 | 24153472 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_025987297.2 | 19488326 | 19494179 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_025989496.1 | 16682503 | 16691437 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_025990768.2 | 24206933 | 24215136 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_025990771.2 | 24193882 | 24203616 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_025990773.2 | 24159797 | 24171994 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_025990783.2 | 23266922 | 23272760 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_039314235.1 | 22843883 | 22852406 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_039314243.1 | 22832891 | 22838176 | cytochrome P450 4C1 |

| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_039314247.1 | 22825641 | 22832587 | cytochrome P450 4C1-like |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_039314988.1 | 23253576 | 23261512 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1577_tr1 | 4538698 | 4544070 | chr16 | XP_039315059.1 | 16669969 | 16679811 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_011165193.2 | 23247076 | 23251744 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_011169811.1 | 24137908 | 24153472 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_025987297.2 | 19488326 | 19494179 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_025989496.1 | 16682503 | 16691437 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_025990768.2 | 24206933 | 24215136 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_025990771.2 | 24193882 | 24203616 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_025990773.2 | 24159797 | 24171994 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_025990783.2 | 23266922 | 23272760 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_039314235.1 | 22843883 | 22852406 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_039314243.1 | 22832891 | 22838176 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_039314247.1 | 22825641 | 22832587 | cytochrome P450 4C1-like |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_039314988.1 | 23253576 | 23261512 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1581_tr1 | 4554866 | 4559774 | chr16 | XP_039315059.1 | 16669969 | 16679811 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_011165193.2 | 23247076 | 23251744 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_011169811.1 | 24137908 | 24153472 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_025987297.2 | 19488326 | 19494179 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_025989496.1 | 16682503 | 16691437 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_025990768.2 | 24206933 | 24215136 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_025990771.2 | 24193882 | 24203616 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_025990773.2 | 24159797 | 24171994 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_025990783.2 | 23266922 | 23272760 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_039314235.1 | 22843883 | 22852406 | cytochrome P450 4C1 isoform X1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_039314243.1 | 22832891 | 22838176 | cytochrome P450 4C1 |
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_039314247.1 | 22825641 | 22832587 | cytochrome P450 4C1-like |

| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_039314988.1 | 23253576 | 23261512 | cytochrome P450 4C1 isoform X1 |
|---|---|---|---|---|---|---|---|---|---|
| HOG13743 | chr02 | gene_1693_tr1 | 10114402 | 1E+07 | chr16 | XP_039315059.1 | 16669969 | 16679811 | cytochrome P450 4C1 isoform X1 |
| HOG15821 | chr02 | gene_1644_tr2 | 8351500 | 8353144 | chr16 | XP_011156738.1 | 24831283 | 24834043 | protein eyes shut homolog |
| HOG15821 | chr02 | gene_2026_tr2 | 8503816 | 8505439 | chr16 | XP_011156738.1 | 24831283 | 24834043 | protein eyes shut homolog |
| HOG8212 | chr02 | gene_1871_tr1 | 2237831 | 2247764 | chr16 | XP_039315119.1 | 15451139 | 15461125 | cytoplasmic aconitate hydratase isoform X1 |
| HOG7975 | chr02 | gene_1507_tr2 | 2529255 | 2532857 | chr16 | XP_011156721.1 | 24619302 | 24626210 | protein PTCD3 homolog mitochondrial |
| HOG6326 | chr02 | gene_2030_tr2 | 8555871 | 8557342 | chr16 | XP_011170130.1 | 17325990 | 17327596 | 39S ribosomal protein L35 mitochondrial |
| HOG9112 | chr02 | gene_1717_tr5 | 10983859 | 1.1E+07 | chr16 | XP_011170713.2 | 4936386 | 4940192 | nucleolar complex protein 3 homolog |
| HOG9069 | chr02 | gene_1832_tr1 | 772763 | 775761 | chr16 | XP_011165231.1 | 23178555 | 23182124 | exonuclease 3'-5' domain-containing protein 2 |
| HOG5566 | chr02 | gene_1427_tr1 | 178187 | 200708 | chr16 | XP_039314324.1 | 21950247 | 21953655 | odorant receptor 63a-like |
| HOG5566 | chr02 | gene_1427_tr1 | 178187 | 200708 | chr16 | XP_039314328.1 | 21940307 | 21943310 | odorant receptor 22c-like |
| HOG8312 | chr02 | gene_1599_tr1 | 5570470 | 5579495 | chr16 | XP_011158158.1 | 21155372 | 21199818 | RNA/RNP complex-1-interacting phosphatase |
| HOG9067 | chr02 | gene_1652_tr1 | 8525966 | 8529287 | chr16 | XP_011157594.1 | 24859458 | 24863682 | ralA-binding protein 1 isoform X1 |
| HOG16061 | chr02 | gene_1959_tr1 | 5063881 | 5081427 | chr16 | XP_011155899.1 | 21745846 | 21850839 | prominin-2 isoform X1 |
| HOG9017 | chr02 | gene_1639_tr1 | 8310629 | 8316151 | chr16 | XP_039314335.1 | 17893498 | 17900904 | cytoplasmic polyadenylation element-binding protein 1 isoform X1 |
| HOG8596 | chr02 | gene_1995_tr2 | 7479424 | 7487751 | chr16 | XP_011161364.1 | 19064971 | 19074495 | protein O-GlcNAcase isoform X1 |
| HOG9703 | chr02 | gene_2108_tr1 | 10556289 | 1.1E+07 | chr16 | XP_011170895.1 | 5437682 | 5441278 | G protein pathway suppressor 2 |
| HOG7282 | chr02 | gene_2096_tr1 | 10288709 | 1E+07 | chr16 | XP_039314844.1 | 7085225 | 7094186 | LOW QUALITY PROTEIN: sodium channel protein Nach |
| HOG7956 | chr02 | gene_1630_tr1 | 7797372 | 7801982 | chr16 | XP_025994004.1 | 18580027 | 18586032 | heparanase isoform X2 |
| HOG13640 | chr02 | gene_1617_tr1 | 7628100 | 7633710 | chr16 | XP_011155781.1 | 18865932 | 18880975 | carbohydrate sulfotransferase 11 |
| HOG8562 | chr02 | gene_1620_tr2 | 7676877 | 7680074 | chr16 | XP_025990739.2 | 24334297 | 24342397 | venom protease isoform X1 |

| HOG10456 | chr02 | gene_1893_tr1 | 2554419 | 2555497 | chr16 | XP_011156752. | 24921737 | 24923928 | phosphoglycerate mutase 2 |
|---|---|---|---|---|---|---|---|---|---|
| HOG8653 | chr02 | gene_2020_tr1 | 8478578 | 8479850 | chr16 | XP_025991944. | 15119956 | 15121483 | proteasome subunit alpha type-2 |
| HOG7955 | chr02 | gene_1629_tr1 | 7792556 | 7796617 | chr16 | XP_011155754. | 18586130 | 18590977 | bromodomain-containing protein 8 isoform X1 |
| HOG10888 | chr02 | gene_1827_tr1 | 550915 | 552731 | chr16 | XP_011174313. | 22790349 | 22794822 | methionine--tRNA ligase mitochondrial |
| HOG11604 | chr02 | gene_2125_tr1 | 11104769 | 1.1E+07 | chr16 | XP_011172202. | 7980975 | 7984021 | adenosine 3'-phospho 5'-phosphosulfate transporter 1 |
| HOG8194 | chr02 | gene_1598_tr2 | 5537261 | 5538549 | chr16 | XP_011158168.2 | 21201197 | 21202834 | protein DPCD |
| HOG15174 | chr02 | gene_1914_tr1 | 3146965 | 3149376 | chr16 | XP_011166886. | 12933551 | 12971889 | aspartate aminotransferase cytoplasmic isoform X1 |
| HOG7786 | chr02 | gene_1539_tr1 | 3231656 | 3232729 | chr16 | XP_011175281. | 9247439 | 9248968 | transmembrane emp24 domain-containing protein eca |
| HOG7416 | chr02 | gene_2013_tr3 | 7995757 | 8004436 | chr16 | XP_011159511. | 18259777 | 18296909 | putative leucine-rich repeat-containing protein DDB_G0290503 isoform X1 |
| HOG7912 | chr02 | gene_1868_tr3 | 2184745 | 2187525 | chr16 | XP_011170060. | 16762066 | 16777830 | hexokinase type 2 isoform X1 |
| HOG8702 | chr02 | gene_1606_tr1 | 7121846 | 7149177 | chr16 | XP_011157760. | 19496127 | 19823163 | protein slit isoform X1 |
| HOG14992 | chr02 | gene_1663_tr2 | 8977052 | 8994687 | chr16 | XP_011161963. | 14429099 | 14452723 | PH and SEC7 domain-containing protein isoform X1 |
| HOG8183 | chr02 | gene_2000_tr1 | 7749120 | 7766692 | chr16 | XP_025994007. | 18623415 | 18645646 | neuronal acetylcholine receptor subunit alpha-10 isoform X1 |
| HOG11435 | chr02 | gene_1550_tr1 | 3471738 | 3473780 | chr16 | XP_011167858. | 15029297 | 15032037 | LOW QUALITY PROTEIN: ornithine aminotransferase mitochondrial |
| HOG11435 | chr02 | gene_1550_tr1 | 3471738 | 3473780 | chr16 | XP_039314411. | 14984631 | 14988285 | ornithine aminotransferase mitochondrial |
| HOG7853 | chr02 | gene_1851_tr1 | 1641742 | 1642471 | chr16 | XP_011164686. | 15978070 | 15983195 | polyadenylate-binding protein-interacting protein 2 |
| HOG10764 | chr02 | gene_1823_tr1 | 372066 | 375575 | chr16 | XP_011174346. | 22576770 | 22583464 | wolframin isoform X1 |
| HOG8371 | chr02 | gene_2005_tr2 | 7808831 | 7815525 | chr16 | XP_011159710. | 18529940 | 18550235 | glycerol-3-phosphate acyltransferase 1 mitochondrial isoform X1 |

| HOG | chr | gene | start1 | end1 | chr16 | XP | start2 | end2 | description |
|---|---|---|---|---|---|---|---|---|---|
| HOG7900 | chr02 | gene_1611_tr1 | 7219634 | 7221695 | chr16 | XP_011157615.1 | 19369186 | 19371285 | probable 39S ribosomal protein L45 mitochondrial |
| HOG8547 | chr02 | gene_1975_tr1 | 6219804 | 6222112 | chr16 | XP_025987285.1 | 20468781 | 20473347 | U4/U6 small nuclear ribonucleoprotein Prp3 isoform X1 |
| HOG9702 | chr02 | gene_1708_tr3 | 10550379 | 1.1E+07 | chr16 | XP_011170905.2 | 5441471 | 5447207 | probable ATP-dependent RNA helicase spindle-E |
| HOG8966 | chr02 | gene_1512_tr1 | 2560443 | 2562269 | chr16 | XP_025990944.2 | 24912668 | 24916031 | crossover junction endonuclease EME1 |
| HOG6280 | chr02 | gene_1653_tr2 | 8541217 | 8541938 | chr16 | XP_011170133.1 | 17335959 | 17337451 | nucleolar protein 16 |
| HOG11601 | chr02 | gene_2123_tr1 | 11078830 | 1.1E+07 | chr16 | XP_011172207.1 | 8010781 | 8018291 | ecdysone 20-monooxygenase isoform X1 |
| HOG8535 | chr02 | gene_1986_tr1 | 7206491 | 7209160 | chr16 | XP_011157681.2 | 19412935 | 19415803 | flap endonuclease GEN |
| HOG8381 | chr02 | gene_1824_tr2 | 377038 | 391543 | chr16 | XP_039314351.1 | 22545731 | 22573243 | alpha-sarcoglycan isoform X1 |
| HOG8386 | chr02 | gene_1628_tr1 | 7787479 | 7788379 | chr16 | XP_011155758.1 | 18595267 | 18597609 | AN1-type zinc finger protein 1 |
| HOG15150 | chr02 | gene_1950_tr1 | 4657835 | 4677811 | chr16 | XP_011170038.2 | 16538591 | 16564671 | tubby-related protein 4 |
| HOG17235 | chr02 | gene_1586_tr1 | 5014932 | 5016002 | chr16 | XP_011173007.1 | 13472076 | 13475943 | general odorant-binding protein 83a |
| HOG6159 | chr02 | gene_2043_tr2 | 8883074 | 8892091 | chr16 | XP_039314679.1 | 26047002 | 26062255 | G protein-activated inward rectifier potassium channel 3 isoform X1 |
| HOG8475 | chr02 | gene_1913_tr2 | 3137648 | 3139191 | chr16 | XP_039314908.1 | 12609397 | 12751315 | axonemal dynein light intermediate polypeptide 1 isoform X1 |
| HOG8372 | chr02 | gene_1862_tr1 | 2076255 | 2081427 | chr16 | XP_039314429.1 | 16927260 | 16938473 | ABC transporter G family member 20 isoform X1 |
| HOG13979 | chr02 | gene_1532_tr1 | 3074660 | 3076528 | chr16 | XP_011170104.1 | 17082495 | 17085279 | m7GpppN-mRNA hydrolase |
| HOG11556 | chr02 | gene_1456_tr1 | 1127222 | 1164340 | chr16 | XP_039314721.1 | 23829468 | 23896143 | dynein heavy chain 2 axonemal |
| HOG11760 | chr02 | gene_1934_tr1 | 3478522 | 3481130 | chr16 | XP_011167863.1 | 15008257 | 15013547 | zinc finger protein 2 homolog isoform X1 |
| HOG10251 | chr02 | gene_2119_tr1 | 11035203 | 1.1E+07 | chr16 | XP_011171085.2 | 5606080 | 5614138 | transcription initiation factor TFIID subunit 1 isoform X1 |
| HOG14394 | chr02 | gene_1829_tr6 | 574059 | 576207 | chr16 | XP_011174246.2 | 22820511 | 22823223 | tubulin alpha chain |

| HOG | chr | gene | start | end | chr | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|
| HOG14931 | chr02 | gene_2053_tr1 | 9002585 | 9043915 | chr16 | XP_025993470.2 | 14458008 | 14533290 | histone-lysine N-methyltransferase 2C isoform X1 |
| HOG8107 | chr02 | gene_1883_tr3 | 2448818 | 2451885 | chr16 | XP_011170016.1 | 16309754 | 16312862 | ATP-dependent RNA helicase abstrakt |
| HOG8366 | chr02 | gene_1503_tr2 | 2470276 | 2473546 | chr16 | XP_025989473.1 | 16331052 | 16334872 | DNA repair and recombination protein RAD54-like |
| HOG7834 | chr02 | gene_1479_tr1 | 2046560 | 2047282 | chr16 | XP_011170092.1 | 16975041 | 16977580 | U3 small nucleolar ribonucleoprotein protein IMP3 |
| HOG9382 | chr02 | gene_2097_tr1 | 10362251 | 1E+07 | chr16 | XP_011172453.1 | 6907523 | 7038323 | leucine-rich repeat-containing protein 24 |
| HOG13033 | chr02 | gene_1949_tr1 | 4650675 | 4652957 | chr16 | XP_011170039.1 | 16566292 | 16569457 | sphingosine-1-phosphate lyase |
| HOG14901 | chr02 | gene_1637_tr3 | 8292437 | 8299997 | chr16 | XP_011175125.1 | 17933856 | 17953352 | tyrosine-protein phosphatase non-receptor type 9 |
| HOG3928 | chr02 | gene_1721_tr1 | 11015360 | 1.1E+07 | chr16 | XP_011171132.2 | 5640676 | 5641436 | glutamyl-tRNA(Gln) amidotransferase subunit C mitochondrial isoform X1 |
| HOG10457 | chr02 | gene_1892_tr1 | 2552211 | 2553631 | chr16 | XP_011156754.1 | 24924414 | 24926934 | ras-related protein Rab-24 isoform X1 |
| HOG13453 | chr02 | gene_1841_tr4 | 1366591 | 1370489 | chr16 | XP_039314259.1 | 24453266 | 24462382 | mitogen-activated protein kinase p38b isoform X2 |
| HOG7910 | chr02 | gene_1499_tr1 | 2409028 | 2411611 | chr16 | XP_011164715.1 | 16178869 | 16182494 | synaptic vesicular amine transporter |
| HOG11500 | chr02 | gene_1719_tr1 | 10995214 | 1.1E+07 | chr16 | XP_011171195.1 | 5675531 | 5688936 | alpha-2-macroglobulin-like protein 1 isoform X1 |
| HOG15088 | chr02 | gene_1860_tr1 | 2067635 | 2068741 | chr16 | XP_011170091.1 | 16949785 | 16951257 | putative methyltransferase C9orf114 homolog |
| HOG9021 | chr02 | gene_1477_tr1 | 2023486 | 2024725 | chr16 | XP_011164648.1 | 15476624 | 15481907 | galactoside alpha-(12)-fucosyltransferase 2 |
| HOG9021 | chr02 | gene_1478_tr1 | 2033266 | 2034505 | chr16 | XP_011164648.1 | 15476624 | 15481907 | galactoside alpha-(12)-fucosyltransferase 2 |
| HOG10386 | chr02 | gene_2114_tr1 | 10990636 | 1.1E+07 | chr16 | XP_011170700.1 | 4928882 | 4932535 | serine/threonine-protein kinase TBK1 |
| HOG10884 | chr02 | gene_1938_tr1 | 3979161 | 3980572 | chr16 | XP_011156782.1 | 25514403 | 25516090 | cuticle protein 21.3 |

| HOG | chr | gene | start | end | chr | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|
| HOG7903 | chr02 | gene_1999_tr1 | 7693975 | 7736544 | chr16 | XP_039314803.1 | 18652601 | 18722783 | disintegrin and metalloproteinase domain-containing protein 10 |
| HOG13927 | chr02 | gene_1529_tr1 | 3072541 | 3073664 | chr16 | XP_011170107.1 | 17080878 | 17082139 | phosphomevalonate kinase |
| HOG15180 | chr02 | gene_1536_tr1 | 3144103 | 3144456 | chr16 | XP_011173101.1 | 12828417 | 12829476 | small integral membrane protein 8 |
| HOG8491 | chr02 | gene_1877_tr1 | 2366397 | 2379139 | chr16 | XP_011164713.1 | 16112917 | 16203623 | G protein-coupled receptor kinase 5 |
| HOG16760 | chr02 | gene_1531_tr1 | 3071636 | 3072259 | chr16 | XP_011170108.1 | 17079637 | 17080688 | 60S ribosomal protein L29-1 |
| HOG10184 | chr02 | gene_2110_tr1 | 10626264 | 1.1E+07 | chr16 | XP_025988733.1 | 4988454 | 5358114 | nephrin isoform X1 |
| HOG9222 | chr02 | gene_1931_tr2 | 3437143 | 3452043 | chr16 | XP_011167875.2 | 14934035 | 14957100 | serine/threonine-protein kinase WNK1 isoform X1 |
| HOG8120 | chr02 | gene_1964_tr2 | 5421067 | 5421712 | chr16 | XP_025987260.1 | 21422668 | 21424863 | high affinity copper uptake protein 1 isoform X1 |
| HOG8218 | chr02 | gene_2015_tr2 | 8334196 | 8336195 | chr16 | XP_025991942.1 | 15124718 | 15131029 | retinol dehydrogenase 14 |
| HOG8218 | chr02 | gene_1648_tr3 | 8484266 | 8486374 | chr16 | XP_025991942.1 | 15124718 | 15131029 | retinol dehydrogenase 14 |
| HOG14108 | chr02 | gene_1951_tr4 | 4684226 | 4690590 | chr16 | XP_011156808.3 | 25887243 | 25903896 | NADP-dependent malic enzyme isoform X1 |
| HOG10515 | chr02 | gene_2113_tr1 | 10982571 | 1.1E+07 | chr16 | XP_011170726.1 | 4940621 | 4941608 | membrane magnesium transporter 1 |
| HOG12176 | chr02 | gene_1533_tr1 | 3088468 | 3097185 | chr16 | XP_039314149.1 | 12368253 | 12425549 | tuberin isoform X1 |
| HOG8670 | chr02 | gene_1976_tr1 | 6241885 | 6280032 | chr16 | XP_011157936.1 | 20053347 | 20462470 | nucleolysin TIAR isoform X1 |
| HOG8186 | chr02 | gene_1600_tr1 | 5695305 | 5708193 | chr16 | XP_011158115.1 | 21018836 | 21079706 | fibroblast growth factor 17 isoform X1 |
| HOG13766 | chr02 | gene_1923_tr3 | 3233620 | 3236643 | chr16 | XP_011161971.1 | 14413614 | 14419966 | STAM-binding protein-like A |
| HOG11167 | chr02 | gene_1707_tr1 | 10532848 | 1.1E+07 | chr16 | XP_025988731.1 | 5459740 | 5470399 | protein dimmed |
| HOG8329 | chr02 | gene_1967_tr1 | 5744411 | 5815901 | chr16 | XP_011158097.2 | 20910154 | 20987545 | transcription factor collier isoform X1 |
| HOG9018 | chr02 | gene_1638_tr3 | 8304929 | 8307892 | chr16 | XP_011175137.1 | 17903198 | 17907453 | cell division cycle protein 16 homolog |
| HOG9345 | chr02 | gene_1670_tr1 | 9380689 | 9384563 | chr16 | XP_011174414.1 | 22386390 | 22391289 | cilia- and flagella-associated protein 206 |
| HOG8448 | chr02 | gene_1492_tr1 | 2302382 | 2303597 | chr16 | XP_011164701.1 | 16048363 | 16050400 | 39S ribosomal protein L21 mitochondrial |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| HOG14252 | chr02 | gene_2121_tr1 | 11045460 | 1.1E+07 | chr16 | XP_011171067.1 | 5602845 | 5604980 | small ubiquitin-related modifier 3-like |
| HOG13663 | chr02 | gene_1464_tr1 | 1371290 | 1373226 | chr16 | XP_011166564.1 | 24464033 | 24468153 | ATPase ASNA1 homolog |
| HOG8679 | chr02 | gene_1876_tr1 | 2317939 | 2327431 | chr16 | XP_039314208.1 | 16065271 | 16075868 | serine/arginine repetitive matrix protein 1 isoform X1 |
| HOG8382 | chr02 | gene_1874_tr8 | 2290850 | 2301671 | chr16 | XP_011164698.1 | 16035848 | 16048178 | choline/ethanolaminephosphotransferase 1 isoform X7 |
| HOG14370 | chr02 | gene_1937_tr1 | 3975447 | 3977432 | chr16 | XP_011156786.2 | 25517499 | 25519872 | DNA-directed primase/polymerase protein |
| HOG8373 | chr02 | gene_1484_tr4 | 2102539 | 2111375 | chr16 | XP_011170075.1 | 16880063 | 16892038 | ABC transporter G family member 20 isoform X1 |
| HOG8373 | chr02 | gene_1484_tr4 | 2102539 | 2111375 | chr16 | XP_011170078.1 | 16895366 | 16912466 | ABC transporter G family member 20 isoform X1 |
| HOG8373 | chr02 | gene_1484_tr4 | 2102539 | 2111375 | chr16 | XP_039314446.1 | 16865685 | 16878893 | ABC transporter G family member 20 |
| HOG8373 | chr02 | gene_1484_tr4 | 2102539 | 2111375 | chr16 | XP_039314447.1 | 16857225 | 16865520 | ABC transporter G family member 20 |
| HOG8373 | chr02 | gene_1484_tr4 | 2102539 | 2111375 | chr16 | XP_039314464.1 | 7607336 | 7622508 | ABC transporter G family member 20-like |
| HOG11768 | chr02 | gene_1549_tr1 | 3475886 | 3478196 | chr16 | XP_011167862.1 | 15005569 | 15008087 | ornithine aminotransferase mitochondrial |
| HOG13925 | chr02 | gene_1530_tr2 | 3066697 | 3070447 | chr16 | XP_025989517.1 | 17071223 | 17076558 | calcium-independent phospholipase A2-gamma isoform X1 |
| HOG9346 | chr02 | gene_2073_tr2 | 9386284 | 9390095 | chr16 | XP_011174406.1 | 22380812 | 22386885 | upstream activation factor subunit spp27 |
| HOG9203 | chr02 | gene_1514_tr1 | 2574022 | 2585803 | chr16 | XP_011156744.1 | 24880664 | 24900576 | beta-TrCP |
| HOG9109 | chr02 | gene_1513_tr1 | 2570850 | 2572513 | chr16 | XP_011156746.1 | 24901145 | 24903477 | phospholipid phosphatase 5 |
| HOG8758 | chr02 | gene_1845_tr2 | 1561287 | 1564013 | chr16 | XP_025991919.1 | 15221127 | 15224921 | WD repeat-containing protein 91 isoform X1 |
| HOG15310 | chr02 | gene_2100_tr2 | 10411652 | 1E+07 | chr16 | XP_011172448.1 | 6867425 | 6871295 | WD repeat-containing protein 13 |
| HOG7201 | chr02 | gene_1647_tr3 | 8480522 | 8481710 | chr16 | XP_011164604.1 | 15122313 | 15124685 | sperm flagellar protein 1 |
| HOG8084 | chr02 | gene_1973_tr5 | 6204904 | 6208979 | chr16 | XP_011158015.1 | 20485060 | 20490596 | heat shock 70 kDa protein cognate 5 |

| HOG7002 | chr02 | gene_1704_tr2 | 10446133 | 1E+07 | chr16 | XP_011171042.1 | 5592831 | 5595486 | pyrimidodiazepine synthase |
|---|---|---|---|---|---|---|---|---|---|
| HOG10953 | chr02 | gene_1884_tr1 | 2454899 | 2457123 | chr16 | XP_011170020.1 | 16315660 | 16317833 | 4-hydroxybutyrate coenzyme A transferase |
| HOG14306 | chr02 | gene_1437_tr1 | 571010 | 572742 | chr16 | XP_011174253.1 | 22813296 | 22817685 | nuclear inhibitor of protein phosphatase 1 |
| HOG9828 | chr02 | gene_1710_tr2 | 10565021 | 1.1E+07 | chr16 | XP_011170858.2 | 5427830 | 5431731 | dipeptidyl peptidase 3 isoform X1 |
| HOG8216 | chr02 | gene_1489_tr1 | 2231292 | 2234142 | chr16 | XP_011164629.1 | 15439689 | 15443682 | major facilitator superfamily domain-containing protein 6 |
| HOG9083 | chr02 | gene_1475_tr3 | 1802955 | 1807411 | chr16 | XP_011164651.2 | 15712031 | 15717641 | erythroid differentiation-related factor 1 |
| HOG5859 | chr02 | gene_2032_tr2 | 8695365 | 8696724 | chr16 | XP_025989525.2 | 17178783 | 17181074 | nucleoside diphosphate kinase homolog 5 isoform X1 |
| HOG10996 | chr02 | gene_1640_tr1 | 8322074 | 8324439 | chr16 | XP_011164633.2 | 15463783 | 15466820 | protein RFT1 homolog isoform X1 |
| HOG10996 | chr02 | gene_2021_tr1 | 8496126 | 8498492 | chr16 | XP_011164633.2 | 15463783 | 15466820 | protein RFT1 homolog isoform X1 |
| HOG7896 | chr02 | gene_1610_tr2 | 7209666 | 7211413 | chr16 | XP_011157667.1 | 19409895 | 19412851 | charged multivesicular body protein 7 |
| HOG9911 | chr02 | gene_1727_tr3 | 11112259 | 1.1E+07 | chr16 | XP_011172199.1 | 7964222 | 7972625 | RNA-binding protein Nova-1 isoform X1 |
| HOG10278 | chr02 | gene_2120_tr1 | 11032945 | 1.1E+07 | chr16 | XP_011171555.1 | 5614286 | 5616241 | protein EFR3 homolog B-like |
| HOG8681 | chr02 | gene_1495_tr1 | 2328488 | 2329466 | chr16 | XP_011164706.1 | 16076484 | 16078090 | glutaredoxin-3 |
| HOG9202 | chr02 | gene_2028_tr2 | 8532952 | 8538883 | chr16 | XP_039315048.1 | 24864372 | 24874524 | exportin-7 isoform X1 |
| HOG13963 | chr02 | gene_1842_tr1 | 1373592 | 1375505 | chr16 | XP_025990729.1 | 24467483 | 24472448 | 2-amino-3-ketobutyrate coenzyme A ligase mitochondrial |
| HOG9344 | chr02 | gene_2027_tr2 | 8520769 | 8523402 | chr16 | XP_025990966.2 | 24851559 | 24857353 | glucose dehydrogenase [FAD quinone] |
| HOG14365 | chr02 | gene_1567_tr2 | 3971998 | 3974172 | chr16 | XP_011156788.1 | 25520361 | 25523691 | stAR-related lipid transfer protein 7 mitochondrial isoform X1 |
| HOG7945 | chr02 | gene_1632_tr1 | 7829322 | 7833913 | chr16 | XP_011159700.2 | 18517852 | 18528961 | biorientation of chromosomes in cell division protein 1-like 1 |
| HOG15914 | chr02 | gene_2001_tr1 | 7768659 | 7770011 | chr16 | XP_039314809.1 | 18619452 | 18623202 | F-box/LRR-repeat protein 7 |

| HOG9019 | chr02 | gene_1857_tr1 | 2040226 | 2042758 | chr16 | XP_011164641.1 | 15468231 | 15473814 | pantothenate kinase 3 isoform X1 |
|---|---|---|---|---|---|---|---|---|---|
| HOG3819 | chr02 | gene_1607_tr1 | 7156690 | 7158378 | chr16 | XP_011157738.1 | 19479169 | 19481011 | pheromone-binding protein Gp-9 |
| HOG3819 | chr02 | gene_1607_tr1 | 7156690 | 7158378 | chr16 | XP_011169815.1 | 24131790 | 24136957 | pheromone-binding protein Gp-9 |
| HOG3819 | chr02 | gene_1607_tr1 | 7156690 | 7158378 | chr16 | XP_011169816.2 | 24118113 | 24119720 | pheromone-binding protein Gp-9 |
| HOG16025 | chr02 | gene_1856_tr1 | 2036593 | 2038769 | chr16 | XP_011164646.2 | 15473940 | 15477068 | dynein regulatory complex subunit 4 |
| HOG8672 | chr02 | gene_1960_tr2 | 5155982 | 5192533 | chr16 | XP_011155897.1 | 21475608 | 21744759 | leishmanolysin-like peptidase isoform X1 |
| HOG8716 | chr02 | gene_1849_tr1 | 1621609 | 1624476 | chr16 | XP_011164610.1 | 15145489 | 15150217 | X-ray repair cross-complementing protein 5 |
| HOG5399 | chr02 | gene_1446_tr1 | 627083 | 630858 | chr16 | XP_025990761.1 | 22937162 | 22942889 | facilitated trehalose transporter Tret1 |
| HOG16020 | chr02 | gene_1983_tr3 | 6785469 | 6786881 | chr16 | XP_011157834.1 | 19886682 | 19888422 | trypsin 5G1 isoform X1 |
| HOG8441 | chr02 | gene_1945_tr2 | 4495244 | 4499484 | chr16 | XP_011170055.1 | 16730385 | 16734694 | 60S ribosomal protein L4 |
| HOG15043 | chr02 | gene_1668_tr3 | 9268769 | 9269450 | chr16 | XP_025990985.1 | 25927528 | 25929537 | guanine nucleotide exchange factor MSS4 homolog |
| HOG8768 | chr02 | gene_1642_tr1 | 8331174 | 8333845 | chr16 | XP_011164602.1 | 15130921 | 15134195 | esterase FE4 |
| HOG8768 | chr02 | gene_2022_tr1 | 8486725 | 8489396 | chr16 | XP_011164602.1 | 15130921 | 15134195 | esterase FE4 |
| HOG15311 | chr02 | gene_1920_tr3 | 3192824 | 3209943 | chr16 | XP_039314111.1 | 9731785 | 9888219 | protein TANC2 isoform X1 |
| HOG8755 | chr02 | gene_1865_tr1 | 2138813 | 2167159 | chr16 | XP_025989484.1 | 16791784 | 16841925 | max dimerization protein 1 |
| HOG9913 | chr02 | gene_1699_tr1 | 10195772 | 1E+07 | chr16 | XP_011155530.1 | 7943071 | 7946078 | scaffold protein salvador isoform X1 |
| HOG8128 | chr02 | gene_1869_tr4 | 2208105 | 2212173 | chr16 | XP_011170070.1 | 16749661 | 16753897 | Hermansky-Pudlak syndrome 1 protein homolog isoform X1 |
| HOG8675 | chr02 | gene_1595_tr1 | 5397895 | 5399539 | chr16 | XP_011155895.1 | 21465338 | 21469039 | EEF1A lysine methyltransferase 2 |
| HOG10328 | chr02 | gene_1700_tr1 | 10223464 | 1E+07 | chr16 | XP_039314545.1 | 7910802 | 7913008 | JNK1/MAPK8-associated membrane protein |
| HOG8775 | chr02 | gene_1928_tr2 | 3296805 | 3305510 | chr16 | XP_011167883.1 | 14694405 | 14905258 | complexin isoform X1 |
| HOG9419 | chr02 | gene_1491_tr1 | 2250023 | 2251767 | chr16 | XP_011164693.1 | 15993178 | 16000203 | serine/arginine-rich splicing factor 1A |

| HOG | chr | gene | start | end | chr | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|
| HOG15512 | chr02 | gene_1563_tr1 | 3832533 | 3833843 | chr16 | XP_025990935.2 | 25674362 | 25682023 | proline-rich protein 2-like |
| HOG13903 | chr02 | gene_1822_tr4 | 312935 | 341739 | chr16 | XP_039314666.1 | 25943937 | 25999258 | glutamate-gated chloride channel isoform X1 |
| HOG7100 | chr02 | gene_1896_tr3 | 2681020 | 2684878 | chr16 | XP_011170134.2 | 17437430 | 17442751 | proclotting enzyme |
| HOG15309 | chr02 | gene_1702_tr3 | 10403239 | 1E+07 | chr16 | XP_025993055.2 | 6869805 | 6878400 | STE20-like serine/threonine-protein kinase isoform X1 |
| HOG16229 | chr02 | gene_2006_tr1 | 7846282 | 7848004 | chr16 | XP_039314701.1 | 18494518 | 18497581 | meiosis-specific nuclear structural protein 1 |
| HOG9610 | chr02 | gene_2109_tr2 | 10561716 | 1.1E+07 | chr16 | XP_039315102.1 | 5431828 | 5434733 | protein FAM50 homolog isoform X1 |
| HOG15284 | chr02 | gene_2098_tr1 | 10399691 | 1E+07 | chr16 | XP_011171593.1 | 7777629 | 7779233 | 60S ribosomal protein L28 |
| HOG8086 | chr02 | gene_1551_tr1 | 3617146 | 3630019 | chr16 | XP_039314948.1 | 16373005 | 16515437 | DE-cadherin isoform X1 |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164671.2 | 15877018 | 15880429 | retinol-binding protein pinta isoform X1 |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164674.1 | 15918699 | 15921707 | retinol-binding protein pinta |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164676.1 | 15898483 | 15904392 | retinol-binding protein pinta |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164678.1 | 15888580 | 15892014 | retinol-binding protein pinta isoform X1 |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164680.2 | 15884737 | 15887708 | retinol-binding protein pinta |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164681.2 | 15880761 | 15883858 | retinol-binding protein pinta |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_025991973.2 | 15922673 | 15925966 | retinol-binding protein pinta |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_039314934.1 | 24095299 | 24102479 | alpha-tocopherol transfer protein-like isoform X1 |
| HOG8295 | chr02 | gene_1452_tr1 | 806294 | 808604 | chr16 | XP_011165153.1 | 23290091 | 23300429 | alpha-tocopherol transfer protein-like |
| HOG8028 | chr02 | gene_1991_tr1 | 7234859 | 7238940 | chr16 | XP_025987316.1 | 19091243 | 19345435 | tyrosine-protein kinase transmembrane receptor Ror isoform X1 |
| HOG15280 | chr02 | gene_2099_tr1 | 10400938 | 1E+07 | chr16 | XP_011171597.1 | 7779426 | 7780932 | S-adenosylmethionine sensor upstream of mTORC1 |
| HOG10512 | chr02 | gene_2112_tr4 | 10969108 | 1.1E+07 | chr16 | XP_011170750.1 | 4946711 | 4958954 | glutathione hydrolase 1 proenzyme |
| HOG8690 | chr02 | gene_1940_tr1 | 4381399 | 4384558 | chr16 | XP_011156768.2 | 25042442 | 25044976 | alpha-glucosidase |
| HOG15030 | chr02 | gene_1838_tr1 | 1282051 | 1283833 | chr16 | XP_011164926.1 | 24065197 | 24074475 | COMM domain-containing protein 4 |

| HOG8708 | chr02 | gene_1835_tr1 | 835166 | 835865 | chr16 | XP_011165066.1 | 23347111 | 23348626 | H/ACA ribonucleoprotein complex subunit 2-like protein |
|---|---|---|---|---|---|---|---|---|---|
| HOG11706 | chr02 | gene_2107_tr1 | 10543095 | 1.1E+07 | chr16 | XP_039314193.1 | 5451045 | 5456789 | cytoglobin-2 isoform X1 |
| HOG8493 | chr02 | gene_1496_tr5 | 2342986 | 2354028 | chr16 | XP_039314212.1 | 16078243 | 16112522 | large neutral amino acids transporter small subunit 1 isoform X1 |
| HOG8949 | chr02 | gene_2008_tr3 | 7860224 | 7865015 | chr16 | XP_025989569.1 | 18470260 | 18477332 | phenoloxidase-activating factor 2 isoform X1 |
| HOG12499 | chr02 | gene_1669_tr1 | 9279870 | 9280947 | chr16 | XP_011165269.1 | 26162881 | 26164234 | coiled-coil domain-containing protein 137 |
| HOG7949 | chr02 | gene_1516_tr2 | 2767965 | 2789287 | chr16 | XP_039315007.1 | 17556002 | 17569874 | semaphorin-5B isoform X1 |
| HOG7841 | chr02 | gene_1672_tr5 | 9719909 | 9741305 | chr16 | XP_039314315.1 | 22039333 | 22090099 | mushroom body large-type Kenyon cell-specific protein 1 isoform X1 |
| HOG7867 | chr02 | gene_1881_tr4 | 2434383 | 2435726 | chr16 | XP_025991950.1 | 16204890 | 16208661 | golgin subfamily A member 7 isoform X1 |
| HOG16123 | chr02 | gene_2127_tr3 | 11177318 | 1.1E+07 | chr16 | XP_011172322.1 | 6128726 | 6131665 | cyclin-dependent kinases regulatory subunit |
| HOG2874 | chr02 | gene_1564_tr1 | 3963068 | 3963349 | chr16 | XP_025990922.1 | 25532553 | 25534584 | glutenin high molecular weight subunit DX5 |
| HOG8684 | chr02 | gene_1886_tr1 | 2486426 | 2487312 | chr16 | XP_039315036.1 | 16349400 | 16351156 | U6 snRNA-associated Sm-like protein LSm8 |
| HOG14776 | chr02 | gene_2116_tr1 | 11017727 | 1.1E+07 | chr16 | XP_011171124.1 | 5638121 | 5639619 | ER lumen protein-retaining receptor |
| HOG8190 | chr02 | gene_1965_tr2 | 5511870 | 5514331 | chr16 | XP_011155883.1 | 21312077 | 21318034 | ATP-dependent RNA helicase vasa |
| HOG8190 | chr02 | gene_1965_tr2 | 5511870 | 5514331 | chr16 | XP_025989543.2 | 17666319 | 17674498 | ATP-dependent RNA helicase vasa |
| HOG8190 | chr02 | gene_1965_tr2 | 5511870 | 5514331 | chr16 | XP_039314125.1 | 17637923 | 17644623 | ATP-dependent RNA helicase vasa |
| HOG8584 | chr02 | gene_1592_tr1 | 5061102 | 5063487 | chr16 | XP_011155907.1 | 21850661 | 21853990 | insulin-like growth factor-binding protein-related protein 1 |
| HOG10444 | chr02 | gene_1517_tr1 | 2807371 | 2815016 | chr16 | XP_025989549.2 | 17607390 | 17616003 | LOW QUALITY PROTEIN: cilia- and flagella-associated protein 43-like |

| HOG10356 | chr02 | gene_1443_tr1 | 602260 | 604541 | chr16 | XP_011165293.1 | 22892412 | 22897385 | serine protease HTRA2 mitochondrial isoform X1 |
|---|---|---|---|---|---|---|---|---|---|
| HOG15689 | chr02 | gene_1731_tr1 | 11175106 | 1.1E+07 | chr16 | XP_011172321.1 | 6131662 | 6133555 | vacuolar protein-sorting-associated protein 25 |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_011155906.1 | 21902262 | 21906351 | odorant receptor 49b isoform X1 |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_025987223.2 | 21882802 | 21885470 | odorant receptor 2a-like |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_025987227.1 | 21913118 | 21916982 | odorant receptor 82a-like |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_025987228.1 | 21918335 | 21922386 | odorant receptor 22c-like |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_039314325.1 | 21924131 | 21927504 | odorant receptor 9a-like |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_039314405.1 | 21886501 | 21891342 | odorant receptor 4-like |
| HOG5520 | chr02 | gene_1813_tr2 | 82571 | 92449 | chr16 | XP_011155906.1 | 21902262 | 21906351 | odorant receptor 49b isoform X1 |
| HOG5520 | chr02 | gene_1813_tr2 | 82571 | 92449 | chr16 | XP_025987223.2 | 21882802 | 21885470 | odorant receptor 2a-like |
| HOG5520 | chr02 | gene_1813_tr2 | 82571 | 92449 | chr16 | XP_025987227.1 | 21913118 | 21916982 | odorant receptor 82a-like |
| HOG5520 | chr02 | gene_1813_tr2 | 82571 | 92449 | chr16 | XP_025987228.1 | 21918335 | 21922386 | odorant receptor 22c-like |
| HOG5520 | chr02 | gene_1813_tr2 | 82571 | 92449 | chr16 | XP_039314325.1 | 21924131 | 21927504 | odorant receptor 9a-like |
| HOG5520 | chr02 | gene_1813_tr2 | 82571 | 92449 | chr16 | XP_039314405.1 | 21886501 | 21891342 | odorant receptor 4-like |
| HOG6111 | chr02 | gene_1448_tr1 | 776848 | 779010 | chr16 | XP_011165211.1 | 23182884 | 23189089 | zinc finger protein 699 |
| HOG8763 | chr02 | gene_1465_tr3 | 1557926 | 1560610 | chr16 | XP_025991920.1 | 15222779 | 15227463 | transcription initiation protein SPT3 homolog isoform X1 |
| HOG15839 | chr02 | gene_1582_tr1 | 4561481 | 4562970 | chr16 | XP_011170040.1 | 16662165 | 16664578 | peptide deformylase mitochondrial |
| HOG10089 | chr02 | gene_1711_tr1 | 10569137 | 1.1E+07 | chr16 | XP_011170848.1 | 5414478 | 5426727 | DNA polymerase alpha subunit B |
| HOG14128 | chr02 | gene_1584_tr1 | 4680005 | 4681037 | chr16 | XP_011156805.1 | 25882931 | 25886321 | RWD domain-containing protein 1 |
| HOG16309 | chr02 | gene_1888_tr2 | 2509939 | 2511890 | chr16 | XP_011156722.1 | 24656092 | 24662689 | major facilitator superfamily domain-containing protein 10 isoform X1 |
| HOG11603 | chr02 | gene_1724_tr2 | 11049948 | 1.1E+07 | chr16 | XP_011172210.1 | 8043481 | 8046799 | cyclin-dependent kinase 8 isoform X1 |
| HOG9369 | chr02 | gene_1635_tr1 | 7988690 | 7992369 | chr16 | XP_011159533.1 | 18297095 | 18301594 | conserved oligomeric Golgi complex subunit 7 |
| HOG8114 | chr02 | gene_1588_tr1 | 5024531 | 5025745 | chr16 | XP_039314227.1 | 13391052 | 13471488 | dopamine receptor 2 |

| HOG16290 | chr02 | gene_1631_tr1 | 7804773 | 7807974 | chr16 | XP_039314810.1 | 18574045 | 18577426 | tektin-3-like |
|---|---|---|---|---|---|---|---|---|---|
| HOG14348 | chr02 | gene_1515_tr1 | 2610166 | 2678821 | chr16 | XP_025989465.2 | 17347717 | 17438257 | atrial natriuretic peptide receptor 1 isoform X1 |
| HOG15119 | chr02 | gene_1480_tr1 | 2058079 | 2064661 | chr16 | XP_039315041.1 | 16954588 | 16968602 | sodium-coupled neutral amino acid transporter 9 isoform X1 |
| HOG8031 | chr02 | gene_2003_tr1 | 7790245 | 7791748 | chr16 | XP_011155757.1 | 18591135 | 18595161 | tetraspanin-33 |

**Supplementary Table 4**: List of one-to-one orthologous genes with functional annotation found on *Solenopsis* chromosome 16, *Cataglyphis* chromosome 2, and *Apis* chromosome 15. The highlighted rows indicate the genes found in the supergene region of both *Solenopsis* and *Cataglyphis*

| HOG | Cnig chr | Cnig_id | Cnig_start | Cnig_end | Sinv chr | Sinv_id | Sinv_start | Sinv_end | Amel_chr | Amel_id | Amel_start | Amel_end | Sinv_function |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG8827 | chr02 | 89_tr1 | 5045800 | 5046548 | chr16 | 161724. | 13267097 | 13271493 | NC_037652.1 | 8354.1 | 1901126 | 1902717 | RHY1A |
| HOG14644 | chr02 | gene_2041_tr2 | 8858673 | 8861002 | chr16 | XP_039314676.1 | 26004603 | 26021597 | NC_037652.1 | XP_016772196.2 | 7468146 | 7473237 | flavin-containing monooxygenase FMO GS-OX-like 2 isoform X1 |
| HOG14988 | chr02 | gene_1922_tr1 | 3230128 | 3231288 | chr16 | XP_011164983.1 | 9360203 | 9363557 | NC_037652.1 | XP_624506.1 | 5099646 | 5100960 | DDRGK domain-containing protein 1 isoform X1 |
| HOG8963 | chr02 | gene_1537_tr1 | 3150294 | 3154642 | chr16 | XP_039314572.1 | 12972095 | 13055442 | NC_037652.1 | XP_006558355.1 | 1967364 | 1972405 | coiled-coil domain-containing protein R3HCC1L isoform X1 |
| HOG10455 | chr02 | gene_1510_tr1 | 2549965 | 2551574 | chr16 | XP_039315055.1 | 24927324 | 24931095 | NC_037652.1 | XP_016772339.1 | 1848897 | 1851670 | S-adenosylmethionine mitochondrial carrier protein |
| HOG13778 | chr02 | gene_1540_tr2 | 3237179 | 3238657 | chr16 | XP_039314532.1 | 12063131 | 12112267 | NC_037652.1 | XP_623717.4 | 817147 | 818362 | DNA excision repair protein ERCC-1 isoform X1 |
| HOG9890 | chr02 | gene_1852_tr1 | 1685469 | 1700678 | chr16 | XP_039314798.1 | 15844676 | 15869563 | NC_037652.1 | XP_006563474.1 | 5877672 | 5913218 | ABC transporter G family member 23 isoform X1 |
| HOG15358 | chr02 | gene_1930_tr1 | 3427464 | 3429176 | chr16 | XP_011167880.1 | 14926676 | 14929867 | NC_037652.1 | XP_016772142.1 | 3754660 | 3756846 | pre-mRNA-splicing factor Slu7 |
| HOG9148 | chr02 | gene_1858_tr1 | 2051611 | 2054176 | chr16 | XP_011170086.1 | 16968597 | 16976194 | NC_037652.1 | XP_395366.2 | 2752212 | 2754786 | phosphoglucomutase isoform X1 |
| HOG8606 | chr02 | gene_2075_tr1 | 9772677 | 9774527 | chr16 | XP_011174441.1 | 22017324 | 22019290 | NC_037652.1 | XP_394080.3 | 6982941 | 6985026 | RCC1 domain-containing protein 1 |
| HOG8446 | chr02 | gene_1844_tr1 | 1518193 | 1532996 | chr16 | XP_011164628.1 | 15263526 | 15305312 | NC_037652.1 | XP_624481.3 | 3181720 | 3229507 | homeotic protein empty spiracles |

| HOG | chr | gene | start | end | chr | protein | start | end | ref | protein2 | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG9085 | chr02 | gene_1447_tr2 | 771890 | 772300 | chr16 | XP_011165219.2 | 23177534 | 23178507 | NC_037652.1 | XP_006563595.1 | 7570143 | 7570878 | mitochondrial import inner membrane translocase subunit Tim16 |
| HOG8563 | chr02 | gene_1619_tr2 | 7659741 | 7663127 | chr16 | XP_039315141.1 | 18807749 | 18837105 | NC_037652.1 | XP_026301426.1 | 3816871 | 3834574 | protein phosphatase 1 regulatory subunit 3C-B |
| HOG7911 | chr02 | gene_1867_tr1 | 2175099 | 2181479 | chr16 | XP_011170059.1 | 16779865 | 16787958 | NC_037652.1 | XP_006557650.1 | 1364081 | 1367870 | protein FAM160B1 isoform X1 |
| HOG15320 | chr02 | gene_1538_tr1 | 3187547 | 3188773 | chr16 | XP_025996242.1 | 10068516 | 10130643 | NC_037652.1 | XP_006561087.1 | 5151892 | 5159314 | tubulin polymerization-promoting protein homolog isoform X1 |
| HOG10354 | chr02 | gene_1442_tr2 | 598774 | 600601 | chr16 | XP_011165314.2 | 22885865 | 22888640 | NC_037652.1 | XP_006560154.1 | 3276780 | 3278706 | rRNA N6-adenosine-methyltransferase ZCCHC4 |
| HOG10946 | chr02 | gene_1501_tr2 | 2452175 | 2454290 | chr16 | XP_011170021.1 | 16312880 | 16315978 | NC_037652.1 | XP_006560748.1 | 3775592 | 3780805 | TGF-beta-activated kinase 1 and MAP3K7-binding protein 1 |
| HOG8215 | chr02 | gene_1504_tr1 | 2476545 | 2484677 | chr16 | XP_039315026.1 | 16336210 | 16350025 | NC_037652.1 | NP_001229447.1 | 2778947 | 2787227 | LOW QUALITY PROTEIN: serine/threonine-protein phosphatase 1 regulatory subunit 10 |
| HOG8214 | chr02 | gene_1870_tr2 | 2218820 | 2226623 | chr16 | XP_039314336.1 | 24529290 | 24544420 | NC_037652.1 | XP_001122328.3 | 1875016 | 1882188 | transferrin |
| HOG8450 | chr02 | gene_2056_tr1 | 9170851 | 9171782 | chr16 | XP_011174382.1 | 22513988 | 22515516 | NC_037652.1 | XP_625205.1 | 7247225 | 7247957 | probable small nuclear ribonucleoprotein G |
| HOG15067 | chr02 | gene_1575_tr7 | 4520673 | 4523032 | chr16 | XP_025989498.1 | 16702011 | 16707132 | NC_037652.1 | XP_624353.1 | 5174985 | 5177250 | aldo-keto reductase family 1 member B1 |
| HOG8607 | chr02 | gene_1674_tr2 | 9776722 | 9779510 | chr16 | XP_011174433.1 | 22014511 | 22017393 | NC_037652.1 | XP_392067.4 | 6981500 | 6983349 | ADP-ribosylation factor-like protein 3 |
| HOG14573 | chr02 | gene_1859_tr1 | 2065162 | 2066044 | chr16 | XP_011170087.1 | 16952983 | 16954745 | NC_037652.1 | XP_624435.2 | 2761034 | 2762120 | ATP synthase subunit C lysine N-methyltransferase |

| HOG | chr | gene | start | end | chr | XP | start | end | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG9902 | chr02 | gene_1624_tr1 | 7773176 | 7775233 | chr16 | XP_025994002.2 | 18608344 | 18615214 | NC_037652.1 | XP_026301079.1 | 3760643 | 3764379 | E3 ubiquitin-protein ligase COP1 isoform X1 |
| HOG8077 | chr02 | gene_1502_tr1 | 2465955 | 2468779 | chr16 | XP_011170022.1 | 16326307 | 16330617 | NC_037652.1 | XP_624112.1 | 3786740 | 3789477 | V-type proton ATPase subunit B |
| HOG9408 | chr02 | gene_2054_tr1 | 9057622 | 9062025 | chr16 | XP_011161961.1 | 14555366 | 14562042 | NC_037652.1 | XP_625224.4 | 6569951 | 6573839 | vacuolar protein sorting-associated protein 51 homolog |
| HOG15232 | chr02 | gene_1932_tr1 | 3463774 | 3465313 | chr16 | XP_025993452.2 | 14979589 | 14984897 | NC_037652.1 | XP_006560031.3 | 2854332 | 2857986 | TBC1 domain family member 7 isoform X1 |
| HOG15356 | chr02 | gene_1887_tr7 | 2500497 | 2507595 | chr16 | XP_011156723.1 | 24664117 | 24678278 | NC_037652.1 | XP_397181.1 | 7580555 | 7588537 | signal transducer and activator of transcription 5B |
| HOG13784 | chr02 | gene_2051_tr4 | 8974048 | 8975733 | chr16 | XP_011161972.1 | 14420379 | 14426100 | NC_037652.1 | XP_006557994.1 | 802108 | 803419 | CDGSH iron-sulfur domain-containing protein 2 homolog |
| HOG17057 | chr02 | gene_1440_tr1 | 577319 | 581594 | chr16 | XP_011165379.1 | 22852621 | 22858466 | NC_037652.1 | XP_393106.4 | 4326973 | 4331594 | coiled-coil domain-containing protein 142 isoform X1 |
| HOG8492 | chr02 | gene_1497_tr1 | 2394239 | 2399165 | chr16 | XP_011164711.1 | 16158349 | 16165729 | NC_037652.1 | XP_006560704.2 | 3569349 | 3579699 | lysyl oxidase homolog 3 |
| HOG8964 | chr02 | gene_1916_tr1 | 3156807 | 3159565 | chr16 | XP_011166899.1 | 13059468 | 13063879 | NC_037652.1 | XP_026301096.1 | 1964252 | 1968227 | signal recognition particle subunit SRP72 |
| HOG10453 | chr02 | gene_1885_tr1 | 2458091 | 2464714 | chr16 | XP_011170017.1 | 16318164 | 16325215 | NC_037652.1 | XP_016772254.2 | 3781273 | 3785782 | 4-hydroxybutyrate coenzyme A transferase isoform X1 |
| HOG10670 | chr02 | gene_1436_tr4 | 555219 | 560423 | chr16 | XP_011174290.1 | 22795556 | 22799567 | NC_037652.1 | XP_393612.2 | 7385909 | 7389473 | casein kinase I isoform X1 |
| HOG8210 | chr02 | gene_1486_tr2 | 2212659 | 2217507 | chr16 | XP_011156707.1 | 24545232 | 24553641 | NC_037652.1 | XP_016772372.2 | 1882589 | 1887117 | cytoplasmic dynein 2 intermediate chain 1 isoform X1 |

| | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG8532 | chr02 | gene_2048_tr1 | 8934105 | 8942741 | chr16 | XP_039314299.1 | 9393467 | 9416297 | NC_037652.1 | XP_026301473.1 | 6907194 | 6936706 | protein phosphatase PP2A 55 kDa regulatory subunit isoform X1 |
| HOG8499 | chr02 | gene_1925_tr1 | 3255213 | 3256394 | chr16 | XP_025993462.1 | 14621720 | 14625536 | NC_037652.1 | XP_006563526.1 | 6577026 | 6578441 | charged multivesicular body protein 3 |
| HOG8376 | chr02 | gene_1873_tr1 | 2264436 | 2284940 | chr16 | XP_039314194.1 | 16006971 | 16033707 | NC_037652.1 | XP_026301215.1 | 2636974 | 2665085 | MAP kinase-activating death domain protein isoform X1 |
| HOG16016 | chr02 | gene_1855_tr1 | 1799371 | 1801890 | chr16 | XP_011164652.2 | 15717500 | 15720868 | NC_037652.1 | XP_016772051.1 | 5985153 | 5989339 | arrestin homolog |
| HOG16856 | chr02 | gene_1455_tr1 | 855531 | 877697 | chr16 | XP_039314720.1 | 23381477 | 23411042 | NC_037652.1 | XP_026301406.1 | 3913712 | 3933292 | dynein heavy chain 8, axonemal |
| HOG15228 | chr02 | gene_1546_tr4 | 3458611 | 3461401 | chr16 | XP_039314410.1 | 14958604 | 14972956 | NC_037652.1 | XP_624637.2 | 2847740 | 2854371 | hippocampus abundant transcript 1 protein isoform X1 |
| HOG7851 | chr02 | gene_1469_tr1 | 1630955 | 1636315 | chr16 | XP_039314800.1 | 15983474 | 15991932 | NC_037652.1 | XP_006563447.1 | 5839515 | 5845614 | catenin alpha isoform X1 |
| HOG7957 | chr02 | gene_1900_tr3 | 2796458 | 2800017 | chr16 | XP_039314811.1 | 18571136 | 18574460 | NC_037652.1 | XP_016772077.1 | 7719771 | 7722791 | LOW QUALITY PROTEIN: leucine-rich repeat-containing protein 15-like |
| HOG11444 | chr02 | gene_1548_tr2 | 3466345 | 3467899 | chr16 | XP_011167869.1 | 14972134 | 14977920 | NC_037652.1 | XP_016772014.1 | 7609362 | 7610910 | methyltransferase-like protein 25 |
| HOG11151 | chr02 | gene_1882_tr1 | 2445602 | 2447802 | chr16 | XP_025989556.1 | 16306420 | 16308828 | NC_037652.1 | XP_026301082.1 | 3772985 | 3774796 | titin |
| HOG16017 | chr02 | gene_1853_tr1 | 1733138 | 1733878 | chr16 | XP_011164666.1 | 15796782 | 15799883 | NC_037652.1 | XP_001121740.1 | 5915203 | 5916217 | copper homeostasis protein cutC homolog |
| HOG7968 | chr02 | gene_1889_tr1 | 2536551 | 2541142 | chr16 | XP_025990923.1 | 24608618 | 24616794 | NC_037652.1 | XP_016772010.2 | 7603611 | 7608972 | SWI/SNF complex subunit SMARCC2 |

| HOG | chr | gene | start | end | chr | protein | start | end | accession | protein | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG11098 | chr02 | gene_1432_tr1 | 366596 | 369341 | chr16 | XP_011174335.1 | 22583577 | 22588088 | NC_037652.1 | XP_016772029.1 | 7274075 | 7277538 | vesicular integral-membrane protein VIP36 |
| HOG7857 | chr02 | gene_1847_tr3 | 1601008 | 1605352 | chr16 | XP_011164616.2 | 15167201 | 15175918 | NC_037652.1 | XP_026301152.1 | 1810844 | 1816811 | pickpocket protein 28 |
| HOG7302 | chr02 | gene_1943_tr1 | 4481091 | 4484240 | chr16 | XP_039314108.1 | 24577632 | 24585030 | NC_037652.1 | XP_026301354.1 | 1895082 | 1900039 | cell death specification protein 2 |
| HOG10999 | chr02 | gene_2029_tr1 | 8542888 | 8547976 | chr16 | XP_011170131.1 | 17328625 | 17335654 | NC_037652.1 | XP_003249587.1 | 3039818 | 3048279 | fumarylacetoacetate hydrolase domain-containing protein 2A isoform X1 |
| HOG7964 | chr02 | gene_1898_tr2 | 2790895 | 2794896 | chr16 | XP_039315010.1 | 17568536 | 17574944 | NC_037652.1 | XP_006563630.1 | 7716749 | 7719637 | cysteine sulfinic acid decarboxylase |
| HOG8217 | chr02 | gene_1488_tr1 | 2235039 | 2237092 | chr16 | XP_011164632.1 | 15448658 | 15451149 | NC_037652.1 | XP_001122354.2 | 1866081 | 1868773 | F-box/WD repeat-containing protein 4 isoform X1 |
| HOG8129 | chr02 | gene_1831_tr4 | 748263 | 770696 | chr16 | XP_011165243.1 | 23132897 | 23176944 | NC_037652.1 | XP_016772179.2 | 3448583 | 3471092 | attractin-like protein 1 isoform X1 |
| HOG8707 | chr02 | gene_1453_tr1 | 822789 | 828821 | chr16 | XP_011165091.1 | 23303851 | 23328875 | NC_037652.1 | XP_006560763.1 | 3877178 | 3891707 | rho-related BTB domain-containing protein 1 isoform X1 |
| HOG8759 | chr02 | gene_1466_tr1 | 1589113 | 1593117 | chr16 | XP_039315121.1 | 15179363 | 15192867 | NC_037652.1 | XP_006558328.1 | 1798229 | 1804590 | zinc finger protein 260 isoform X1 |
| HOG10889 | chr02 | gene_1435_tr3 | 544379 | 548693 | chr16 | XP_025987290.1 | 22671772 | 22789697 | NC_037652.1 | NP_001011595.1 | 7337792 | 7381156 | dopamine receptor 1 isoform X2 |
| HOG4475 | chr02 | gene_1864_tr3 | 2116467 | 2124441 | chr16 | XP_039314456.1 | 16843476 | 16854742 | NC_037652.1 | XP_026301456.1 | 931522 | 938104 | titin homolog isoform X5 |
| HOG10645 | chr02 | gene_1441_tr4 | 590415 | 596041 | chr16 | XP_025990760.1 | 22873090 | 22882736 | NC_037652.1 | XP_006560152.2 | 3268423 | 3274574 | phosphatidylinositide phosphatase SAC2 isoform X1 |

| HOG | chr | gene | start | end | chr | XP | start | end | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG7954 | chr02 | gene_1470_tr1 | 1644698 | 1667692 | chr16 | XP_011164684.1 | 15942310 | 15976783 | NC_037652.1 | XP_393165.3 | 5851689 | 5869613 | BCL2/adenovirus E1B 19 kDa protein-interacting protein 3 |
| HOG11158 | chr02 | gene_1655_tr3 | 8557444 | 8592647 | chr16 | XP_025989546.2 | 17289159 | 17326197 | NC_037652.1 | XP_006560038.1 | 2984878 | 3038381 | MICOS complex subunit Mic60 isoform X1 |
| HOG10644 | chr02 | gene_1828_tr2 | 561841 | 570185 | chr16 | XP_011174263.3 | 22802774 | 22813157 | NC_037652.1 | XP_026301419.1 | 7392002 | 7399902 | dynactin subunit 1 isoform X1 |
| HOG14655 | chr02 | gene_2068_tr2 | 9260339 | 9263319 | chr16 | XP_039314678.1 | 26000887 | 26005049 | NC_037652.1 | XP_016772200.1 | 7473001 | 7474470 | trypsin isoform X1 |
| HOG9816 | chr02 | gene_1518_tr1 | 2817961 | 2822157 | chr16 | NP_001291533.1 | 17618379 | 17624016 | NC_037652.1 | NP_001011572.1 | 7734777 | 7737849 | transferrin precursor |
| HOG8390 | chr02 | gene_1836_tr2 | 838903 | 853826 | chr16 | XP_039314181.1 | 23361028 | 23379858 | NC_037652.1 | XP_006560766.2 | 3898768 | 3913376 | vacuolar protein sorting-associated protein 13B |
| HOG8680 | chr02 | gene_1493_tr1 | 2305670 | 2309668 | chr16 | XP_039314217.1 | 16050663 | 16058164 | NC_037652.1 | XP_016772183.1 | 3472194 | 3475098 | N-acetyl-D-glucosamine kinase isoform X1 |
| HOG8551 | chr02 | gene_1618_tr1 | 7636416 | 7641713 | chr16 | XP_011155778.1 | 18840151 | 18849161 | NC_037652.1 | XP_026301140.1 | 3837947 | 3846341 | poly [ADP-ribose] polymerase tankyrase isoform X1 |
| HOG8001 | chr02 | gene_1500_tr3 | 2421470 | 2427974 | chr16 | XP_025991945.2 | 16188502 | 16201893 | NC_037652.1 | XP_006560707.1 | 3607609 | 3619427 | cGMP-specific 3',5'-cyclic phosphodiesterase isoform X1 |
| HOG7835 | chr02 | gene_1509_tr1 | 2545617 | 2546126 | chr16 | XP_011156760.1 | 24934835 | 24938580 | NC_037652.1 | XP_026301309.1 | 2806705 | 2809484 | high mobility group protein HMGI-C |
| HOG15092 | chr02 | gene_1948_tr4 | 4566904 | 4596170 | chr16 | XP_025989494.1 | 16575605 | 16659139 | NC_037652.1 | XP_026301054.1 | 8213522 | 8295995 | tachykinin-like peptides receptor 99D isoform X1 |
| HOG5340 | chr02 | gene_1485_tr2 | 2203039 | 2206996 | chr16 | XP_011170067.1 | 16753719 | 16784018 | NC_037652.1 | XP_006557644.1 | 1392562 | 1426650 | BRISC complex subunit FAM175B |

| | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG10009 | chr02 | gene_1830_tr3 | 633905 | 642281 | chr16 | XP_025990751.1 | 22946890 | 23121595 | NC_037652.1 | NP_001011565.1 | 3310369 | 3326664 | octopamine receptor Oamb isoform X1 |
| HOG8826 | chr02 | gene_1958_tr1 | 5052175 | 5054200 | chr16 | XP_039314418.1 | 13289459 | 13298659 | NC_037652.1 | XP_003249249.1 | 1910665 | 1913312 | UPF0687 protein C20orf27 homolog |
| HOG10675 | chr02 | gene_1438_tr1 | 587446 | 588814 | chr16 | XP_011165354.1 | 22866931 | 22869701 | NC_037652.1 | XP_001121593.3 | 3266453 | 3268150 | ubiquitin-like domain-containing CTD phosphatase 1 |
| HOG7970 | chr02 | gene_1454_tr1 | 836500 | 838102 | chr16 | XP_011165023.1 | 23358829 | 23361307 | NC_037652.1 | XP_001120581.3 | 3896389 | 3898161 | RING finger protein 37 isoform X1 |
| HOG8965 | chr02 | gene_1895_tr1 | 2563359 | 2569774 | chr16 | XP_011156748.1 | 24904560 | 24912249 | NC_037652.1 | XP_006563441.2 | 5805728 | 5812112 | KAT8 regulatory NSL complex subunit 3 |
| HOG8571 | chr02 | gene_1998_tr1 | 7683628 | 7685506 | chr16 | XP_011155767.1 | 18738035 | 18741095 | NC_037652.1 | XP_006560753.1 | 3801532 | 3803522 | interleukin enhancer-binding factor 2 |
| HOG8212 | chr02 | gene_1871_tr1 | 2237831 | 2247764 | chr16 | XP_039315119.1 | 15451139 | 15461125 | NC_037652.1 | XP_392993.2 | 1856328 | 1866310 | cytoplasmic aconitate hydratase isoform X1 |
| HOG7975 | chr02 | gene_1507_tr2 | 2529255 | 2532857 | chr16 | XP_011156721.1 | 24619302 | 24626210 | NC_037652.1 | XP_026301084.1 | 7598458 | 7601895 | protein PTCD3 homolog, mitochondrial |
| HOG6326 | chr02 | gene_2030_tr2 | 8555871 | 8557342 | chr16 | XP_011170130.1 | 17325990 | 17327596 | NC_037652.1 | XP_001121481.1 | 3038290 | 3039526 | 39S ribosomal protein L35, mitochondrial |
| HOG9112 | chr02 | gene_1717_tr5 | 10983859 | 10987057 | chr16 | XP_011170713.2 | 4936386 | 4940192 | NC_037652.1 | XP_006563442.2 | 5802212 | 5805720 | nucleolar complex protein 3 homolog |
| HOG9069 | chr02 | gene_1832_tr1 | 772763 | 775761 | chr16 | XP_011165231.1 | 23178555 | 23182124 | NC_037652.1 | XP_001121410.2 | 7569208 | 7574371 | exonuclease 3'-5' domain-containing protein 2 |
| HOG9067 | chr02 | gene_1652_tr1 | 8525966 | 8529287 | chr16 | XP_011157594.1 | 24859458 | 24863682 | NC_037652.1 | XP_006560751.3 | 3790265 | 3793768 | ralA-binding protein 1 isoform X1 |

| HOG7956 | chr02 | gene_1630_tr1 | 7797372 | 7801982 | chr16 | XP_025994004.1 | 18580027 | 18586032 | NC_037652.1 | XP_006563433.2 | 5782507 | 5786019 | heparanase isoform X2 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG10456 | chr02 | gene_1893_tr1 | 2554419 | 2555497 | chr16 | XP_011156752.1 | 24921737 | 24923928 | NC_037652.1 | XP_625114.2 | 1843783 | 1845473 | phosphoglycerate mutase 2 |
| HOG7955 | chr02 | gene_1629_tr1 | 7792556 | 7796617 | chr16 | XP_011155754.1 | 18586130 | 18590977 | NC_037652.1 | XP_006563436.1 | 5777886 | 5781715 | bromodomain-containing protein 8 isoform X1 |
| HOG10888 | chr02 | gene_1827_tr1 | 550915 | 552731 | chr16 | XP_011174313.1 | 22790349 | 22794822 | NC_037652.1 | XP_016772024.1 | 7381274 | 7385104 | methionine--tRNA ligase, mitochondrial |
| HOG15174 | chr02 | gene_1914_tr1 | 3146965 | 3149376 | chr16 | XP_011166886.1 | 12933551 | 12971889 | NC_037652.1 | XP_026301294.1 | 2741183 | 2743234 | aspartate aminotransferase, cytoplasmic isoform X1 |
| HOG7786 | chr02 | gene_1539_tr1 | 3231656 | 3232729 | chr16 | XP_011175281.1 | 9247439 | 9248968 | NC_037652.1 | XP_396009.2 | 5098612 | 5099750 | transmembrane emp24 domain-containing protein eca |
| HOG7912 | chr02 | gene_1868_tr3 | 2184745 | 2187525 | chr16 | XP_011170060.1 | 16762066 | 16777830 | NC_037652.1 | XP_006557645.1 | 1370111 | 1414274 | hexokinase type 2 isoform X1 |
| HOG14992 | chr02 | gene_1663_tr2 | 8977052 | 8994687 | chr16 | XP_011161963.1 | 14429099 | 14452723 | NC_037652.1 | XP_026301277.1 | 665447 | 801495 | PH and SEC7 domain-containing protein isoform X1 |
| HOG8183 | chr02 | gene_2000_tr1 | 7749120 | 7766692 | chr16 | XP_025994007.1 | 18623415 | 18645646 | NC_037652.1 | XP_392070.3 | 6475086 | 6484878 | neuronal acetylcholine receptor subunit alpha-10 isoform X1 |
| HOG7853 | chr02 | gene_1851_tr1 | 1641742 | 1642471 | chr16 | XP_011164686.1 | 15978070 | 15983195 | NC_037652.1 | XP_625228.3 | 5846801 | 5850960 | polyadenylate-binding protein-interacting protein 2 |
| HOG10764 | chr02 | gene_1823_tr1 | 372066 | 375575 | chr16 | XP_011174346.1 | 22576770 | 22583464 | NC_037652.1 | XP_625196.1 | 7269445 | 7273061 | wolframin isoform X1 |
| HOG8371 | chr02 | gene_2005_tr2 | 7808831 | 7815525 | chr16 | XP_011159710.1 | 18529940 | 18550235 | NC_037652.1 | XP_006563437.1 | 5790955 | 5801638 | glycerol-3-phosphate acyltransferase 1, mitochondrial isoform X1 |

| | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG8966 | chr02 | gene_1512_tr1 | 2560443 | 2562269 | chr16 | XP_025990944.2 | 24912668 | 24916031 | NC_037652.1 | XP_625230.3 | 5836240 | 5838400 | crossover junction endonuclease EME1 |
| HOG8381 | chr02 | gene_1824_tr2 | 377038 | 391543 | chr16 | XP_039314351.1 | 22545731 | 22573243 | NC_037652.1 | XP_016772284.1 | 7252464 | 7268830 | alpha-sarcoglycan isoform X1 |
| HOG8386 | chr02 | gene_1628_tr1 | 7787479 | 7788379 | chr16 | XP_011155758.1 | 18595267 | 18597609 | NC_037652.1 | XP_006563431.1 | 5773415 | 5774845 | AN1-type zinc finger protein 1 |
| HOG15150 | chr02 | gene_1950_tr1 | 4657835 | 4677811 | chr16 | XP_011170038.2 | 16538591 | 16564671 | NC_037652.1 | XP_006563712.2 | 8306616 | 8347879 | tubby-related protein 4 |
| HOG17235 | chr02 | gene_1586_tr1 | 5014932 | 5016002 | chr16 | XP_011173007.1 | 13472076 | 13475943 | NC_037652.1 | XP_016771954.1 | 6325010 | 6335391 | general odorant-binding protein 83a |
| HOG8372 | chr02 | gene_1862_tr1 | 2076255 | 2081427 | chr16 | XP_039314429.1 | 16927260 | 16938473 | NC_037652.1 | XP_026301458.1 | 818668 | 904370 | ABC transporter G family member 20 isoform X1 |
| HOG11556 | chr02 | gene_1456_tr1 | 1127222 | 1164340 | chr16 | XP_039314721.1 | 23829468 | 23896143 | NC_037652.1 | XP_026301171.1 | 4283861 | 4316980 | dynein heavy chain 2, axonemal |
| HOG11760 | chr02 | gene_1934_tr1 | 3478522 | 3481130 | chr16 | XP_011167863.1 | 15008257 | 15013547 | NC_037652.1 | XP_003250284.1 | 7619194 | 7621734 | zinc finger protein 2 homolog isoform X1 |
| HOG14394 | chr02 | gene_1829_tr6 | 574059 | 576207 | chr16 | XP_011174246.2 | 22820511 | 22823223 | NC_037652.1 | XP_006563567.1 | 7403869 | 7406083 | tubulin alpha chain |
| HOG14931 | chr02 | gene_2053_tr1 | 9002585 | 9043915 | chr16 | XP_025993470.2 | 14458008 | 14533290 | NC_037652.1 | XP_026301259.1 | 628195 | 659504 | histone-lysine N-methyltransferase 2C isoform X1 |
| HOG8107 | chr02 | gene_1883_tr3 | 2448818 | 2451885 | chr16 | XP_011170016.1 | 16309754 | 16312862 | NC_037652.1 | XP_006563505.1 | 6520331 | 6523313 | ATP-dependent RNA helicase abstrakt |
| HOG8366 | chr02 | gene_1503_tr2 | 2470276 | 2473546 | chr16 | XP_025989473.1 | 16331052 | 16334872 | NC_037652.1 | XP_026301194.1 | 2787901 | 2793072 | DNA repair and recombination protein RAD54-like |

| | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG7834 | chr02 | gene_1479_tr1 | 2046560 | 2047282 | chr16 | XP_011170092.1 | 16975041 | 16977580 | NC_037652.1 | XP_623809.3 | 2746709 | 2747683 | U3 small nucleolar ribonucleoprotein protein IMP3 |
| HOG13033 | chr02 | gene_1949_tr1 | 4650675 | 4652957 | chr16 | XP_011170039.1 | 16566292 | 16569457 | NC_037652.1 | XP_006563709.1 | 8302408 | 8305934 | sphingosine-1-phosphate lyase |
| HOG10457 | chr02 | gene_1892_tr1 | 2552211 | 2553631 | chr16 | XP_011156754.1 | 24924414 | 24926934 | NC_037652.1 | XP_625108.2 | 1847455 | 1849068 | ras-related protein Rab-24 isoform X1 |
| HOG7910 | chr02 | gene_1499_tr1 | 2409028 | 2411611 | chr16 | XP_011164715.1 | 16178869 | 16182494 | NC_037652.1 | XP_026301355.1 | 3591924 | 3595481 | synaptic vesicular amine transporter |
| HOG7903 | chr02 | gene_1999_tr1 | 7693975 | 7736544 | chr16 | XP_039314803.1 | 18652601 | 18722783 | NC_037652.1 | XP_026301182.1 | 6486653 | 6518547 | disintegrin and metalloproteinase domain-containing protein 10 |
| HOG15180 | chr02 | gene_1536_tr1 | 3144103 | 3144456 | chr16 | XP_011173101.1 | 12828417 | 12829476 | NC_037652.1 | XP_006560007.3 | 2745726 | 2746837 | small integral membrane protein 8 |
| HOG8491 | chr02 | gene_1877_tr1 | 2366397 | 2379139 | chr16 | XP_011164713.1 | 16112917 | 16203623 | NC_037652.1 | XP_394109.3 | 3540252 | 3561158 | G protein-coupled receptor kinase 5 |
| HOG9222 | chr02 | gene_1931_tr2 | 3437143 | 3452043 | chr16 | XP_011167875.2 | 14934035 | 14957100 | NC_037652.1 | XP_016772106.2 | 2830254 | 2844944 | serine/threonine-protein kinase WNK1 isoform X1 |
| HOG14108 | chr02 | gene_1951_tr4 | 4684226 | 4690590 | chr16 | XP_011156808.3 | 25887243 | 25903896 | NC_037652.1 | XP_006563718.1 | 8354130 | 8362747 | NADP-dependent malic enzyme isoform X1 |
| HOG13766 | chr02 | gene_1923_tr3 | 3233620 | 3236643 | chr16 | XP_011161971.1 | 14413614 | 14419966 | NC_037652.1 | XP_001120689.2 | 813461 | 816339 | STAM-binding protein-like A |
| HOG9345 | chr02 | gene_1670_tr1 | 9380689 | 9384563 | chr16 | XP_011174414.1 | 22386390 | 22391289 | NC_037652.1 | XP_026301414.1 | 7193694 | 7205450 | cilia- and flagella-associated protein 206 |
| HOG8448 | chr02 | gene_1492_tr1 | 2302382 | 2303597 | chr16 | XP_011164701.1 | 16048363 | 16050400 | NC_037652.1 | XP_001121076.4 | 2687555 | 2688979 | 39S ribosomal protein L21, mitochondrial |

| HOG | chr | gene | start | end | chr | XP | start | end | NC | XP2 | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG8679 | chr02 | gene_1876_tr1 | 2317939 | 2327431 | chr16 | XP_039314208.1 | 16065271 | 16075868 | NC_037652.1 | XP_006560700.1 | 3481629 | 3490486 | serine/arginine repetitive matrix protein 1 isoform X1 |
| HOG8382 | chr02 | gene_1874_tr8 | 2290850 | 2301671 | chr16 | XP_011164698.1 | 16035848 | 16048178 | NC_037652.1 | XP_016772117.1 | 2676722 | 2687337 | choline/ethanolaminephosphotransferase 1 isoform X7 |
| HOG14370 | chr02 | gene_1937_tr1 | 3975447 | 3977432 | chr16 | XP_011156786.2 | 25517499 | 25519872 | NC_037652.1 | XP_001121815.3 | 7666195 | 7668589 | DNA-directed primase/polymerase protein |
| HOG11768 | chr02 | gene_1549_tr1 | 3475886 | 3478196 | chr16 | XP_011167862.1 | 15005569 | 15008087 | NC_037652.1 | XP_394061.2 | 7617390 | 7619211 | ornithine aminotransferase, mitochondrial |
| HOG9346 | chr02 | gene_2073_tr2 | 9386284 | 9390095 | chr16 | XP_011174406.1 | 22380812 | 22386885 | NC_037652.1 | XP_392065.2 | 7189452 | 7193590 | upstream activation factor subunit spp27 |
| HOG9203 | chr02 | gene_1514_tr1 | 2574022 | 2585803 | chr16 | XP_011156744.1 | 24880664 | 24900576 | NC_037652.1 | XP_623915.2 | 5814862 | 5822003 | beta-TrCP |
| HOG9109 | chr02 | gene_1513_tr1 | 2570850 | 2572513 | chr16 | XP_011156746.1 | 24901145 | 24903477 | NC_037652.1 | XP_623887.3 | 5813096 | 5814741 | phospholipid phosphatase 5 |
| HOG8758 | chr02 | gene_1845_tr2 | 1561287 | 1564013 | chr16 | XP_025991919.1 | 15221127 | 15224921 | NC_037652.1 | XP_006558326.1 | 1753000 | 1756222 | WD repeat-containing protein 91 isoform X1 |
| HOG10953 | chr02 | gene_1884_tr1 | 2454899 | 2457123 | chr16 | XP_011170020.1 | 16315660 | 16317833 | NC_037652.1 | XP_392704.2 | 3778662 | 3781143 | 4-hydroxybutyrate coenzyme A transferase |
| HOG14306 | chr02 | gene_1437_tr1 | 571010 | 572742 | chr16 | XP_011174253.1 | 22813296 | 22817685 | NC_037652.1 | XP_003250277.1 | 7401072 | 7403555 | nuclear inhibitor of protein phosphatase 1 |
| HOG8216 | chr02 | gene_1489_tr1 | 2231292 | 2234142 | chr16 | XP_011164629.1 | 15439689 | 15443682 | NC_037652.1 | XP_006558342.1 | 1868903 | 1877664 | major facilitator superfamily domain-containing protein 6 |
| HOG9083 | chr02 | gene_1475_tr3 | 1802955 | 1807411 | chr16 | XP_011164651.2 | 15712031 | 15717641 | NC_037652.1 | XP_393162.3 | 5988608 | 5993569 | erythroid differentiation-related factor 1 |

| HOG8681 | chr02 | gene_1495_tr1 | 2328488 | 2329466 | chr16 | XP_011164706.1 | 16076484 | 16078090 | NC_037652.1 | XP_016772335.1 | 3491262 | 3492418 | glutaredoxin-3 |
|---------|-------|---------------|---------|---------|-------|----------------|----------|----------|-------------|----------------|---------|---------|----------------|
| HOG9202 | chr02 | gene_2028_tr2 | 8532952 | 8538883 | chr16 | XP_039315048.1 | 24864372 | 24874524 | NC_037652.1 | XP_006563445.1 | 5825072 | 5832919 | exportin-7 isoform X1 |
| HOG14365 | chr02 | gene_1567_tr2 | 3971998 | 3974172 | chr16 | XP_011156788.1 | 25520361 | 25523691 | NC_037652.1 | XP_006563616.1 | 7663677 | 7666565 | stAR-related lipid transfer protein 7, mitochondrial isoform X1 |
| HOG15914 | chr02 | gene_2001_tr1 | 7768659 | 7770011 | chr16 | XP_039314809.1 | 18619452 | 18623202 | NC_037652.1 | XP_003249733.2 | 3751884 | 3754310 | F-box/LRR-repeat protein 7 |
| HOG8716 | chr02 | gene_1849_tr1 | 1621609 | 1624476 | chr16 | XP_011164610.1 | 15145489 | 15150217 | NC_037652.1 | XP_026301314.1 | 1836789 | 1840008 | X-ray repair cross-complementing protein 5 |
| HOG8441 | chr02 | gene_1945_tr2 | 4495244 | 4499484 | chr16 | XP_011170055.1 | 16730385 | 16734694 | NC_037652.1 | XP_026301184.1 | 6470416 | 6473217 | 60S ribosomal protein L4 |
| HOG15043 | chr02 | gene_1668_tr3 | 9268769 | 9269450 | chr16 | XP_025990985.1 | 25927528 | 25929537 | NC_037652.1 | XP_006563593.1 | 7554461 | 7556096 | guanine nucleotide exchange factor MSS4 homolog |
| HOG15311 | chr02 | gene_1920_tr3 | 3192824 | 3209943 | chr16 | XP_039314111.1 | 9731785 | 9888219 | NC_037652.1 | XP_026301325.1 | 5106146 | 5150785 | protein TANC2 isoform X1 |
| HOG8755 | chr02 | gene_1865_tr1 | 2138813 | 2167159 | chr16 | XP_025989484.1 | 16791784 | 16841925 | NC_037652.1 | XP_026301069.1 | 951021 | 1360467 | max dimerization protein 1 |
| HOG8128 | chr02 | gene_1869_tr4 | 2208105 | 2212173 | chr16 | XP_011170070.1 | 16749661 | 16753897 | NC_037652.1 | XP_026301300.1 | 1426960 | 1430242 | Hermansky-Pudlak syndrome 1 protein homolog isoform X1 |
| HOG8775 | chr02 | gene_1928_tr2 | 3296805 | 3305510 | chr16 | XP_011167883.1 | 14694405 | 14905258 | NC_037652.1 | XP_625223.2 | 6603005 | 6841964 | complexin isoform X1 |
| HOG9419 | chr02 | gene_1491_tr1 | 2250023 | 2251767 | chr16 | XP_011164693.1 | 15993178 | 16000203 | NC_037652.1 | XP_026301223.1 | 2612739 | 2632495 | serine/arginine-rich splicing factor 1A |

| HOG | chr | gene | start | end | chr | XP | start | end | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG13903 | chr02 | gene_1822_tr4 | 312935 | 341739 | chr16 | XP_039314666.1 | 25943937 | 25999258 | NC_037652.1 | XP_006563397.1 | 7474685 | 7550203 | glutamate-gated chloride channel isoform X1 |
| HOG7100 | chr02 | gene_1896_tr3 | 2681020 | 2684878 | chr16 | XP_011170134.2 | 17437430 | 17442751 | NC_037652.1 | XP_001121888.2 | 7690528 | 7693016 | proclotting enzyme |
| HOG8295 | chr02 | gene_1452_tr1 | 806294 | 808604 | chr16 | XP_011165153.1 | 23290091 | 23300429 | NC_037652.1 | XP_393237.4 | 3871533 | 3876482 | alpha-tocopherol transfer protein-like |
| HOG8708 | chr02 | gene_1835_tr1 | 835166 | 835865 | chr16 | XP_011165066.1 | 23347111 | 23348626 | NC_037652.1 | XP_625159.2 | 3895314 | 3896376 | H/ACA ribonucleoprotein complex subunit 2-like protein |
| HOG8493 | chr02 | gene_1496_tr5 | 2342986 | 2354028 | chr16 | XP_039314212.1 | 16078243 | 16112522 | NC_037652.1 | XP_026301448.1 | 3504722 | 3533434 | large neutral amino acids transporter small subunit 1 isoform X1 |
| HOG7949 | chr02 | gene_1516_tr2 | 2767965 | 2789287 | chr16 | XP_039315007.1 | 17556002 | 17569874 | NC_037652.1 | XP_394067.4 | 7697592 | 7717297 | semaphorin-5B isoform X1 |
| HOG7841 | chr02 | gene_1672_tr5 | 9719909 | 9741305 | chr16 | XP_039314315.1 | 22039333 | 22090099 | NC_037652.1 | XP_006563386.2 | 6989419 | 7037855 | mushroom body large-type Kenyon cell-specific protein 1 isoform X1 |
| HOG7867 | chr02 | gene_1881_tr4 | 2434383 | 2435726 | chr16 | XP_025991950.1 | 16204890 | 16208661 | NC_037652.1 | XP_001122074.3 | 3621194 | 3625443 | golgin subfamily A member 7 isoform X1 |
| HOG8684 | chr02 | gene_1886_tr1 | 2486426 | 2487312 | chr16 | XP_039315036.1 | 16349400 | 16351156 | NC_037652.1 | XP_624537.1 | 2776324 | 2778184 | U6 snRNA-associated Sm-like protein LSm8 |
| HOG10444 | chr02 | gene_1517_tr1 | 2807371 | 2815016 | chr16 | XP_025989549.2 | 17607390 | 17616003 | NC_037652.1 | XP_026301341.1 | 7727328 | 7733724 | LOW QUALITY PROTEIN: cilia- and flagella-associated protein 43-like |
| HOG10356 | chr02 | gene_1443_tr1 | 602260 | 604541 | chr16 | XP_011165293.1 | 22892412 | 22897385 | NC_037652.1 | XP_624354.1 | 3279596 | 3282286 | serine protease HTRA2, mitochondrial isoform X1 |
| HOG6111 | chr02 | gene_1448_tr1 | 776848 | 779010 | chr16 | XP_011165211.1 | 23182884 | 23189089 | NC_037652.1 | XP_026301142.1 | 3846445 | 3848620 | zinc finger protein 699 |

**Supplementary Table 5**: List of many-to-many orthologous genes with functional annotation found on *Solenopsis* chromosome 16, *Cataglyphis* chromosome 2, and Apis chromosome 15. The highlighted rows indicate the genes found in the supergene region of both *Solenopsis* and *Cataglyphis*

| HOG | Cnig_chr | Cnig_id | Cnig_start | Cnig_end | Sinv_chr | Sinv_id | Sinv_start | Sinv_end | Amel_chr | Amel_id | Amel_start | Amel_end | Sinv_function |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG8698 | chr02 | gene_1641_tr1 | 8325068 | 8328804 | chr16 | XP_011164607.1 | 1.5E+07 | 1.5E+07 | NC_037644.1 | XP_396618.2 | 3003200 | 3007735 | melanotransferrin isoform X1 |
| HOG8698 | chr02 | gene_2023_tr1 | 8491765 | 8495501 | chr16 | XP_011164607.1 | 1.5E+07 | 1.5E+07 | NC_037644.1 | XP_396618.2 | 3003200 | 3007735 | melanotransferrin isoform X1 |
| HOG8827 | chr02 | gene_1589_tr1 | 5045800 | 5046548 | chr16 | XP_011161724.2 | 1.3E+07 | 1.3E+07 | NC_037652.1 | XP_006558354.1 | 1901126 | 1902717 | probable E3 ubiquitin-protein ligase RHY1A |
| HOG15688 | chr02 | gene_1695_tr1 | 1E+07 | 1E+07 | chr16 | XP_011170189.2 | 3995281 | 3997294 | NC_037642.1 | XP_016767737.2 | 1.4E+07 | 1.4E+07 | cytoplasmic tRNA 2-thiolation protein 2 |
| HOG14644 | chr02 | gene_2041_tr2 | 8858673 | 8861002 | chr16 | XP_039314676.1 | 2.6E+07 | 2.6E+07 | NC_037652.1 | XP_016772196.2 | 7468146 | 7473237 | flavin-containing monooxygenase FMO GS-OX-like 2 isoform X1 |
| HOG14988 | chr02 | gene_1922_tr1 | 3230128 | 3231288 | chr16 | XP_011164983.1 | 9360203 | 9363557 | NC_037652.1 | XP_624506.1 | 5099646 | 5100960 | DDRGK domain-containing protein 1 isoform X1 |
| HOG8963 | chr02 | gene_1537_tr1 | 3150294 | 3154642 | chr16 | XP_039314572.1 | 1.3E+07 | 1.3E+07 | NC_037652.1 | XP_006558355.1 | 1967364 | 1972405 | coiled-coil domain-containing protein R3HCC1L isoform X1 |
| HOG10455 | chr02 | gene_1510_tr1 | 2549965 | 2551574 | chr16 | XP_039315055.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_016772339.1 | 1848897 | 1851670 | S-adenosylmethionine mitochondrial carrier protein |
| HOG13778 | chr02 | gene_1540_tr2 | 3237179 | 3238657 | chr16 | XP_039314532.1 | 1.2E+07 | 1.2E+07 | NC_037652.1 | XP_623717.4 | 817147 | 818362 | DNA excision repair protein ERCC-1 isoform X1 |
| HOG9890 | chr02 | gene_1852_tr1 | 1685469 | 1700678 | chr16 | XP_039314798.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_006563474.1 | 5877672 | 5913218 | ABC transporter G family member 23 isoform X1 |
| HOG15358 | chr02 | gene_1930_tr1 | 3427464 | 3429176 | chr16 | XP_011167880.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_016772142.1 | 3754660 | 3756846 | pre-mRNA-splicing factor Slu7 |
| HOG8187 | chr02 | gene_1597_tr1 | 5443206 | 5533951 | chr16 | XP_011155884.1 | 2.1E+07 | 2.1E+07 | NC_037638.1 | XP_003251788.1 | 2.7E+07 | 2.8E+07 | segmentation protein cap'n'collar isoform X1 |
| HOG9148 | chr02 | gene_1858_tr1 | 2051611 | 2054176 | chr16 | XP_011170086.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_395366.2 | 2752212 | 2754786 | phosphoglucomutase isoform X1 |
| HOG8353 | chr02 | gene_2019_tr1 | 8388278 | 8391469 | chr16 | XP_039314831.1 | 2.5E+07 | 2.5E+07 | NC_037644.1 | XP_026297552.1 | 3089715 | 3215807 | RNA-binding protein Musashi homolog 2 isoform X1 |
| HOG8606 | chr02 | gene_2075_tr1 | 9772677 | 9774527 | chr16 | XP_011174441.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_394080.3 | 6982941 | 6985026 | RCC1 domain-containing protein 1 |
| HOG15068 | chr02 | gene_1569_tr1 | 3988663 | 3991066 | chr16 | XP_025990993.1 | 2.5E+07 | 2.6E+07 | NC_037645.1 | XP_394733.2 | 5869577 | 5873195 | ras-related protein Rab-27A |

| HOG ID | chr | gene | start | end | chr | XP | val | val | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG13962 | chr02 | gene_1843_tr1 | 1377153 | 1379481 | chr16 | XP_011166565.2 | 2.4E+07 | 2.4E+07 | NC_037645.1 | XP_394739.4 | 6326715 | 6329530 | nicalin |
| HOG17165 | chr02 | gene_2017_tr1 | 8345501 | 8348747 | chr16 | XP_011156735.1 | 2.5E+07 | 2.5E+07 | NC_037644.1 | XP_001120174.2 | 3250519 | 3256028 | fibrillin-1 isoform X1 |
| HOG17165 | chr02 | gene_1650_tr1 | 8508193 | 8511442 | chr16 | XP_011156735.1 | 2.5E+07 | 2.5E+07 | NC_037644.1 | XP_001120174.2 | 3250519 | 3256028 | fibrillin-1 isoform X1 |
| HOG8446 | chr02 | gene_1844_tr1 | 1518193 | 1532996 | chr16 | XP_011164628.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_624481.3 | 3181720 | 3229507 | homeotic protein empty spiracles |
| HOG9085 | chr02 | gene_1447_tr2 | 771890 | 772300 | chr16 | XP_011165219.2 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_006563595.1 | 7570143 | 7570878 | mitochondrial import inner membrane translocase subunit Tim16 |
| HOG8563 | chr02 | gene_1619_tr2 | 7659741 | 7663127 | chr16 | XP_039315141.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_026301426.1 | 3816871 | 3834574 | protein phosphatase 1 regulatory subunit 3C-B |
| HOG7911 | chr02 | gene_1867_tr1 | 2175099 | 2181479 | chr16 | XP_011170059.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_006557650.1 | 1364081 | 1367870 | protein FAM160B1 isoform X1 |
| HOG15320 | chr02 | gene_1538_tr1 | 3187547 | 3188773 | chr16 | XP_025996242.1 | 1E+07 | 1E+07 | NC_037652.1 | XP_006561087.1 | 5151892 | 5159314 | tubulin polymerization-promoting protein homolog isoform X1 |
| HOG10354 | chr02 | gene_1442_tr2 | 598774 | 600601 | chr16 | XP_011165314.2 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_006560154.1 | 3276780 | 3278706 | rRNA N6-adenosine-methyltransferase ZCCHC4 |
| HOG9918 | chr02 | gene_2126_tr2 | 1.1E+07 | 1.1E+07 | chr16 | XP_039314554.1 | 7947767 | 7953232 | NC_037642.1 | XP_016767705.1 | 8430697 | 8433098 | probable maleylacetoacetate isomerase 2 isoform X1 |
| HOG15278 | chr02 | gene_2118_tr3 | 1.1E+07 | 1.1E+07 | chr16 | XP_025988755.2 | 5617066 | 5621371 | NC_037644.1 | XP_016768765.1 | 1.2E+07 | 1.2E+07 | modifier of mdg4 isoform X1 |
| HOG13352 | chr02 | gene_1732_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011172324.1 | 6139353 | 6141960 | NC_037642.1 | XP_006570309.1 | 1.4E+07 | 1.4E+07 | E3 ubiquitin-protein ligase RNF34 |
| HOG8813 | chr02 | gene_1905_tr2 | 2847105 | 2848201 | chr16 | XP_011175190.1 | 1.8E+07 | 1.8E+07 | NC_037644.1 | XP_392468.3 | 3614281 | 3615740 | charged multivesicular body protein 5 |
| HOG10946 | chr02 | gene_1501_tr2 | 2452175 | 2454290 | chr16 | XP_011170021.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_006560748.1 | 3775592 | 3780805 | TGF-beta-activated kinase 1 and MAP3K7-binding protein 1 |
| HOG10399 | chr02 | gene_1718_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170689.2 | 4932448 | 4936043 | NC_037644.1 | XP_026297455.1 | 9608862 | 9611617 | protein VAC14 homolog |
| HOG8106 | chr02 | gene_1994_tr2 | 7468244 | 7472789 | chr16 | XP_025987315.1 | 1.9E+07 | 1.9E+07 | NC_037645.1 | XP_003249083.2 | 295491 | 300707 | transcription elongation factor SPT5 |
| HOG10090 | chr02 | gene_1712_tr4 | 1.1E+07 | 1.1E+07 | chr16 | XP_039315095.1 | 5384580 | 5413651 | NC_037642.1 | XP_006567657.1 | 1.1E+07 | 1.1E+07 | steroid hormone receptor ERR1 isoform X1 |
| HOG8215 | chr02 | gene_1504_tr1 | 2476545 | 2484677 | chr16 | XP_039315026.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | NP_001229447.1 | 2778947 | 2787227 | LOW QUALITY PROTEIN: serine/threonine-protein phosphatase 1 regulatory subunit 10 |

| HOG | chr | gene | start | end | chr | XP | val | val | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG12500 | chr02 | gene_2072_tr1 | 9281188 | 9282321 | chr16 | XP_011165270.1 | 2.6E+07 | 2.6E+07 | NC_037651.1 | XP_393997.1 | 8013242 | 8015091 | BLOC-1-related complex subunit 5 |
| HOG8214 | chr02 | gene_1870_tr2 | 2218820 | 2226623 | chr16 | XP_039314336.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_001122328.3 | 1875016 | 1882188 | transferrin |
| HOG8450 | chr02 | gene_2056_tr1 | 9170851 | 9171782 | chr16 | XP_011174382.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_625205.1 | 7247225 | 7247957 | probable small nuclear ribonucleoprotein G |
| HOG7892 | chr02 | gene_1987_tr2 | 7211682 | 7216161 | chr16 | XP_011157636.1 | 1.9E+07 | 1.9E+07 | NC_037643.1 | XP_001120594.2 | 5361607 | 5365983 | modular serine protease-like |
| HOG7892 | chr02 | gene_1987_tr2 | 7211682 | 7216161 | chr16 | XP_025987236.2 | 1.9E+07 | 1.9E+07 | NC_037643.1 | XP_001120594.2 | 5361607 | 5365983 | modular serine protease |
| HOG7892 | chr02 | gene_1987_tr2 | 7211682 | 7216161 | chr16 | XP_025987238.1 | 1.9E+07 | 1.9E+07 | NC_037643.1 | XP_001120594.2 | 5361607 | 5365983 | clotting factor C isoform X1 |
| HOG8673 | chr02 | gene_1594_tr1 | 5111114 | 5144049 | chr16 | XP_025987255.1 | 2.2E+07 | 2.2E+07 | NC_037638.1 | XP_026297876.1 | 2.8E+07 | 2.8E+07 | trace amine-associated receptor 9 |
| HOG14737 | chr02 | gene_1720_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011171146.2 | 5639561 | 5640843 | NC_037644.1 | XP_003250450.1 | 8712969 | 8714124 | DNA polymerase subunit gamma-2 mitochondrial |
| HOG15067 | chr02 | gene_1575_tr7 | 4520673 | 4523032 | chr16 | XP_025989498.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_624353.1 | 5174985 | 5177250 | aldo-keto reductase family 1 member B1 |
| HOG10327 | chr02 | gene_2094_tr1 | 1E+07 | 1E+07 | chr16 | XP_011155529.1 | 7919926 | 7939780 | NC_037642.1 | XP_623652.2 | 8400485 | 8424389 | serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit epsilon isoform |
| HOG8607 | chr02 | gene_1674_tr2 | 9776722 | 9779510 | chr16 | XP_011174433.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_392067.4 | 6981500 | 6983349 | ADP-ribosylation factor-like protein 3 |
| HOG14573 | chr02 | gene_1859_tr1 | 2065162 | 2066044 | chr16 | XP_011170087.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_624435.2 | 2761034 | 2762120 | ATP synthase subunit C lysine N-methyltransferase |
| HOG9902 | chr02 | gene_1624_tr1 | 7773176 | 7775233 | chr16 | XP_025994002.2 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_026301079.1 | 3760643 | 3764379 | E3 ubiquitin-protein ligase COP1 isoform X1 |
| HOG8077 | chr02 | gene_1502_tr1 | 2465955 | 2468779 | chr16 | XP_011170022.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_624112.1 | 3786740 | 3789477 | V-type proton ATPase subunit B |
| HOG9408 | chr02 | gene_2054_tr1 | 9057622 | 9062025 | chr16 | XP_011161961.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_625224.4 | 6569951 | 6573839 | vacuolar protein sorting-associated protein 51 homolog |
| HOG15232 | chr02 | gene_1932_tr1 | 3463774 | 3465313 | chr16 | XP_025993452.2 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_006560031.3 | 2854332 | 2857986 | TBC1 domain family member 7 isoform X1 |
| HOG15520 | chr02 | gene_1462_tr3 | 1350205 | 1352805 | chr16 | XP_011169804.1 | 2.4E+07 | 2.4E+07 | NC_037644.1 | NP_001011576.1 | 3706639 | 3736938 | tachykinins |
| HOG15226 | chr02 | gene_2101_tr2 | 1E+07 | 1E+07 | chr16 | XP_011172444.1 | 6856005 | 6866241 | NC_037644.1 | XP_026297411.1 | 1.2E+07 | 1.2E+07 | protein lingerer isoform X1 |

| HOG | chr | gene | start | end | chr | XP | val | val | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG8357 | chr02 | gene_1645_tr1 | 8354286 | 8354759 | chr16 | XP_011156726.2 | 2.5E+07 | 2.5E+07 | NC_037644.1 | XP_006560218.1 | 3246693 | 3247703 | 39S ribosomal protein L51 mitochondrial |
| HOG8357 | chr02 | gene_2025_tr1 | 8502200 | 8502673 | chr16 | XP_011156726.2 | 2.5E+07 | 2.5E+07 | NC_037644.1 | XP_006560218.1 | 3246693 | 3247703 | 39S ribosomal protein L51 mitochondrial |
| HOG15356 | chr02 | gene_1887_tr7 | 2500497 | 2507595 | chr16 | XP_011156723.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_397181.1 | 7580555 | 7588537 | signal transducer and activator of transcription 5B |
| HOG13784 | chr02 | gene_2051_tr4 | 8974048 | 8975733 | chr16 | XP_011161972.1 | 1.4E+07 | 1.4E+07 | NC_037652.1 | XP_006557994.1 | 802108 | 803419 | CDGSH iron-sulfur domain-containing protein 2 homolog |
| HOG17057 | chr02 | gene_1440_tr1 | 577319 | 581594 | chr16 | XP_011165379.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_393106.4 | 4326973 | 4331594 | coiled-coil domain-containing protein 142 isoform X1 |
| HOG8794 | chr02 | gene_1904_tr2 | 2840928 | 2843949 | chr16 | XP_011175214.1 | 1.8E+07 | 1.8E+07 | NC_037644.1 | XP_001120765.2 | 3618293 | 3621337 | nuclear envelope integral membrane protein 1 isoform X1 |
| HOG8492 | chr02 | gene_1497_tr1 | 2394239 | 2399165 | chr16 | XP_011164711.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_006560704.2 | 3569349 | 3579699 | lysyl oxidase homolog 3 |
| HOG8964 | chr02 | gene_1916_tr1 | 3156807 | 3159565 | chr16 | XP_011166899.1 | 1.3E+07 | 1.3E+07 | NC_037652.1 | XP_026301096.1 | 1964252 | 1968227 | signal recognition particle subunit SRP72 |
| HOG15083 | chr02 | gene_1705_tr2 | 1E+07 | 1E+07 | chr16 | XP_011170978.2 | 5549025 | 5555577 | NC_037638.1 | XP_006557413.1 | 1154365 | 1171005 | poly(A)-specific ribonuclease PARN |
| HOG15025 | chr02 | gene_1458_tr2 | 1247917 | 1274925 | chr16 | XP_039315079.1 | 2.4E+07 | 2.4E+07 | NC_037644.1 | XP_006560904.1 | 3629560 | 3685688 | protein tincar isoform X1 |
| HOG7898 | chr02 | gene_1988_tr2 | 7216732 | 7219024 | chr16 | XP_011157570.1 | 1.9E+07 | 1.9E+07 | NC_037643.1 | XP_016768353.1 | 5366144 | 5368790 | tRNA pseudouridine synthase A isoform X1 |
| HOG10640 | chr02 | gene_1524_tr2 | 2991447 | 3001477 | chr16 | XP_011170093.1 | 1.7E+07 | 1.7E+07 | NC_037644.1 | XP_001122101.2 | 2984163 | 2994696 | huntingtin isoform X1 |
| HOG10453 | chr02 | gene_1885_tr1 | 2458091 | 2464714 | chr16 | XP_011170017.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_016772254.2 | 3781273 | 3785782 | 4-hydroxybutyrate coenzyme A transferase isoform X1 |
| HOG7890 | chr02 | gene_1634_tr1 | 7848863 | 7852708 | chr16 | XP_011159689.1 | 1.8E+07 | 1.8E+07 | NC_037644.1 | XP_624007.4 | 774023 | 777486 | ATP-dependent RNA helicase SUV3 homolog mitochondrial |
| HOG10670 | chr02 | gene_1436_tr4 | 555219 | 560423 | chr16 | XP_011174290.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_393612.2 | 7385909 | 7389473 | casein kinase I isoform X1 |
| HOG8352 | chr02 | gene_2018_tr1 | 8355667 | 8358937 | chr16 | XP_011156728.1 | 2.5E+07 | 2.5E+07 | NC_037644.1 | XP_623859.3 | 3242220 | 3246635 | dipeptidyl peptidase 9 |
| HOG8352 | chr02 | gene_1649_tr1 | 8496401 | 8501292 | chr16 | XP_011156728.1 | 2.5E+07 | 2.5E+07 | NC_037644.1 | XP_623859.3 | 3242220 | 3246635 | dipeptidyl peptidase 9 |
| HOG10436 | chr02 | gene_1722_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011171109.1 | 5623693 | 5628008 | NC_037649.1 | XP_395531.2 | 3085455 | 3089187 | amino acid transporter AVT1C |

| HOG9022 | chr02 | gene_2014_tr1 | 8320370 | 8321475 | chr16 | XP_011175161.1 | 1.8E+07 | 1.8E+07 | NC_037641.1 | XP_006558776.1 | 3403052 | 3404489 | peroxisomal membrane protein 11C |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG13826 | chr02 | gene_1526_tr2 | 3021334 | 3025332 | chr16 | XP_011170099.1 | 1.7E+07 | 1.7E+07 | NC_037645.1 | XP_006563910.1 | 6298874 | 6305232 | endoplasmic reticulum metallopeptidase 1 isoform X1 |
| HOG15085 | chr02 | gene_1939_tr1 | 4056694 | 4063002 | chr16 | XP_025990986.1 | 2.5E+07 | 2.5E+07 | NC_037645.1 | XP_006561540.1 | 5724589 | 5794463 | protein artichoke isoform X1 |
| HOG8210 | chr02 | gene_1486_tr2 | 2212659 | 2217507 | chr16 | XP_011156707.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_016772372.2 | 1882589 | 1887117 | cytoplasmic dynein 2 intermediate chain 1 isoform X1 |
| HOG13636 | chr02 | gene_1616_tr1 | 7553722 | 7607336 | chr16 | XP_039314493.1 | 1.9E+07 | 1.9E+07 | NC_037645.1 | XP_026297802.1 | 323612 | 504123 | potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 3 isoform X1 |
| HOG8665 | chr02 | gene_1906_tr2 | 3009188 | 3013281 | chr16 | XP_011170095.1 | 1.7E+07 | 1.7E+07 | NC_037644.1 | XP_026297409.1 | 3691625 | 3700285 | probable sodium/potassium/calcium exchanger CG1090 |
| HOG8671 | chr02 | gene_1570_tr1 | 4414120 | 4420669 | chr16 | XP_011156769.2 | 2.5E+07 | 2.5E+07 | NC_037645.1 | XP_006561538.1 | 5311778 | 5511371 | protein artichoke |
| HOG16969 | chr02 | gene_1603_tr3 | 6164963 | 6195172 | chr16 | XP_039314822.1 | 2E+07 | 2.1E+07 | NC_037638.1 | XP_026302114.1 | 2.7E+07 | 2.7E+07 | Kv channel-interacting protein 4 isoform X1 |
| HOG8801 | chr02 | gene_1633_tr3 | 7839020 | 7842392 | chr16 | XP_011159621.1 | 1.9E+07 | 1.9E+07 | NC_037644.1 | XP_625179.1 | 762056 | 766637 | mitoferrin-1 isoform X1 |
| HOG8266 | chr02 | gene_1929_tr2 | 3421778 | 3426438 | chr16 | XP_039314407.1 | 1.5E+07 | 1.5E+07 | NC_037641.1 | XP_006558773.1 | 3432609 | 3436942 | activin receptor type-2A isoform X1 |
| HOG13656 | chr02 | gene_1463_tr1 | 1358112 | 1362142 | chr16 | XP_011166559.1 | 2.4E+07 | 2.4E+07 | NC_037645.1 | XP_395200.2 | 6305789 | 6309954 | trimeric intracellular cation channel type 1B.1 |
| HOG8532 | chr02 | gene_2048_tr1 | 8934105 | 8942741 | chr16 | XP_039314299.1 | 9393467 | 9416297 | NC_037652.1 | XP_026301473.1 | 6907194 | 6936706 | protein phosphatase PP2A 55 kDa regulatory subunit isoform X1 |
| HOG9131 | chr02 | gene_2010_tr1 | 7871199 | 7874674 | chr16 | XP_011159599.3 | 1.8E+07 | 1.8E+07 | NC_037645.1 | XP_016769181.1 | 1195970 | 1199949 | phenoloxidase-activating factor 2 |
| HOG4469 | chr02 | gene_1439_tr4 | 582905 | 586363 | chr16 | XP_025990753.1 | 2.3E+07 | 2.3E+07 | NC_037644.1 | XP_026297653.1 | 885875 | 906429 | peroxidase |
| HOG8118 | chr02 | gene_1596_tr1 | 5425828 | 5435317 | chr16 | XP_011155888.2 | 2.1E+07 | 2.1E+07 | NC_037638.1 | XP_006571773.2 | 2.8E+07 | 2.8E+07 | endoplasmic reticulum aminopeptidase 1 isoform X1 |
| HOG8721 | chr02 | gene_1972_tr1 | 6201764 | 6203381 | chr16 | XP_011158029.1 | 2E+07 | 2E+07 | NC_037638.1 | XP_624123.1 | 2.7E+07 | 2.7E+07 | protein farnesyltransferase/geranylgeranyltransferase type-1 subunit alpha |
| HOG9823 | chr02 | gene_2111_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170763.1 | 4962700 | 4967344 | NC_037642.1 | XP_001120140.3 | 1.2E+07 | 1.2E+07 | ecdysteroid-regulated 16 kDa protein |
| HOG8533 | chr02 | gene_1980_tr1 | 6744707 | 6759773 | chr16 | XP_011157873.1 | 2E+07 | 2E+07 | NC_037638.1 | XP_003251782.2 | 2.6E+07 | 2.6E+07 | SEC23-interacting protein isoform X1 |

| HOG | chr | gene | start | end | chr | XP | start | end | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG10187 | chr02 | gene_1715_tr5 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170774.2 | 5086065 | 5108033 | NC_037642.1 | XP_006567694.1 | 1.2E+07 | 1.2E+07 | glucose dehydrogenase [FAD quinone] |
| HOG8499 | chr02 | gene_1925_tr1 | 3255213 | 3256394 | chr16 | XP_025993462.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_006563526.1 | 6577026 | 6578441 | charged multivesicular body protein 3 |
| HOG13861 | chr02 | gene_2033_tr1 | 8709815 | 8752864 | chr16 | XP_039314262.1 | 1.7E+07 | 1.7E+07 | NC_037644.1 | XP_026297481.1 | 3758652 | 3790433 | microtubule-associated protein futsch isoform X1 |
| HOG8376 | chr02 | gene_1873_tr1 | 2264436 | 2284940 | chr16 | XP_039314194.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301215.1 | 2636974 | 2665085 | MAP kinase-activating death domain protein isoform X1 |
| HOG10513 | chr02 | gene_1716_tr2 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170739.1 | 4941982 | 4944570 | NC_037642.1 | XP_006570292.1 | 1.4E+07 | 1.4E+07 | proteasome subunit alpha type-7-1 |
| HOG8591 | chr02 | gene_1935_tr1 | 3671398 | 3672611 | chr16 | XP_011156801.1 | 2.6E+07 | 2.6E+07 | NC_037645.1 | XP_006558272.1 | 1180949 | 1183291 | transmembrane protein 230 isoform X1 |
| HOG16016 | chr02 | gene_1855_tr1 | 1799371 | 1801890 | chr16 | XP_011164652.2 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_016772051.1 | 5985153 | 5989339 | arrestin homolog |
| HOG16856 | chr02 | gene_1455_tr1 | 855531 | 877697 | chr16 | XP_039314720.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_026301406.1 | 3913712 | 3933292 | dynein heavy chain 8 axonemal |
| HOG15228 | chr02 | gene_1546_tr4 | 3458611 | 3461401 | chr16 | XP_039314410.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_624637.2 | 2847740 | 2854371 | hippocampus abundant transcript 1 protein isoform X1 |
| HOG7851 | chr02 | gene_1469_tr1 | 1630955 | 1636315 | chr16 | XP_039314800.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_006563447.1 | 5839515 | 5845614 | catenin alpha isoform X1 |
| HOG7957 | chr02 | gene_1900_tr3 | 2796458 | 2800017 | chr16 | XP_039314811.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_016772077.1 | 7719771 | 7722791 | LOW QUALITY PROTEIN: leucine-rich repeat-containing protein 15-like |
| HOG11444 | chr02 | gene_1548_tr2 | 3466345 | 3467899 | chr16 | XP_011167869.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_016772014.1 | 7609362 | 7610910 | methyltransferase-like protein 25 |
| HOG11151 | chr02 | gene_1882_tr1 | 2445602 | 2447802 | chr16 | XP_025989556.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301082.1 | 3772985 | 3774796 | titin |
| HOG16017 | chr02 | gene_1853_tr1 | 1733138 | 1733878 | chr16 | XP_011164666.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_001121740.1 | 5915203 | 5916217 | copper homeostasis protein cutC homolog |
| HOG7968 | chr02 | gene_1889_tr1 | 2536551 | 2541142 | chr16 | XP_025990923.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_016772010.2 | 7603611 | 7608972 | SWI/SNF complex subunit SMARCC2 |
| HOG16199 | chr02 | gene_1726_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011172205.1 | 7983972 | 7985364 | NC_037642.1 | XP_006570296.1 | 1.4E+07 | 1.4E+07 | transmembrane protein 216 |
| HOG11098 | chr02 | gene_1432_tr1 | 366596 | 369341 | chr16 | XP_011174335.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_016772029.1 | 7274075 | 7277538 | vesicular integral-membrane protein VIP36 |
| HOG14511 | chr02 | gene_1703_tr2 | 1E+07 | 1E+07 | chr16 | XP_011171054.1 | 5595847 | 5602703 | NC_037644.1 | XP_026297386.1 | 4722736 | 4729199 | mediator of RNA polymerase II transcription subunit 23 |

| HOG | chr | gene | start | end | chr | XP | start | end | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG17002 | chr02 | gene_2103_tr1 | 1E+07 | 1E+07 | chr16 | XP_011171031.2 | 5579776 | 5591916 | NC_037642.1 | XP_001120289.2 | 8577355 | 8643888 | neprilysin-1 isoform X1 |
| HOG7857 | chr02 | gene_1847_tr3 | 1601008 | 1605352 | chr16 | XP_011164616.2 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_026301152.1 | 1810844 | 1816811 | pickpocket protein 28 |
| HOG15089 | chr02 | gene_1481_tr1 | 2066442 | 2067463 | chr16 | XP_011170090.1 | 1.7E+07 | 1.7E+07 | NC_037648.1 | XP_026299779.1 | 8303837 | 8304960 | ribulose-phosphate 3-epimerase |
| HOG9436 | chr02 | gene_1636_tr2 | 8259469 | 8285211 | chr16 | XP_039315011.1 | 1.8E+07 | 1.8E+07 | NC_037644.1 | XP_006560193.2 | 2679151 | 2933459 | probable JmjC domain-containing histone demethylation protein 2C isoform X1 |
| HOG7302 | chr02 | gene_1943_tr1 | 4481091 | 4484240 | chr16 | XP_039314108.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_026301354.1 | 1895082 | 1900039 | cell death specification protein 2 |
| HOG10999 | chr02 | gene_2029_tr1 | 8542888 | 8547976 | chr16 | XP_011170131.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_003249587.1 | 3039818 | 3048279 | fumarylacetoacetate hydrolase domain-containing protein 2A isoform X1 |
| HOG7964 | chr02 | gene_1898_tr2 | 2790895 | 2794896 | chr16 | XP_039315010.1 | 1.8E+07 | 1.8E+07 | NC_037652.1 | XP_006563630.1 | 7716749 | 7719637 | cysteine sulfinic acid decarboxylase |
| HOG8217 | chr02 | gene_1488_tr1 | 2235039 | 2237092 | chr16 | XP_011164632.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_001122354.2 | 1866081 | 1868773 | F-box/WD repeat-containing protein 4 isoform X1 |
| HOG8129 | chr02 | gene_1831_tr4 | 748263 | 770696 | chr16 | XP_011165243.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_016772179.2 | 3448583 | 3471092 | attractin-like protein 1 isoform X1 |
| HOG9612 | chr02 | gene_1709_tr2 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170885.1 | 5434001 | 5437202 | NC_037642.1 | XP_006564861.1 | 8536672 | 8538007 | CDAN1-interacting nuclease 1 |
| HOG8472 | chr02 | gene_1535_tr2 | 3108146 | 3134625 | chr16 | XP_025994936.2 | 1.2E+07 | 1.3E+07 | NC_037641.1 | XP_026295782.1 | 3355479 | 3402249 | protein lap4 isoform X1 |
| HOG8707 | chr02 | gene_1453_tr1 | 822789 | 828821 | chr16 | XP_011165091.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_006560763.1 | 3877178 | 3891707 | rho-related BTB domain-containing protein 1 isoform X1 |
| HOG8759 | chr02 | gene_1466_tr1 | 1589113 | 1593117 | chr16 | XP_039315121.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_006558328.1 | 1798229 | 1804590 | zinc finger protein 260 isoform X1 |
| HOG9836 | chr02 | gene_1605_tr10 | 6783211 | 6810424 | chr16 | XP_039314308.1 | 2E+07 | 2E+07 | NC_037645.1 | XP_006561491.3 | 4509082 | 4537706 | protein unc-13 homolog 4B isoform X1 |
| HOG8988 | chr02 | gene_2011_tr2 | 7880423 | 7884430 | chr16 | XP_011159578.1 | 1.8E+07 | 1.8E+07 | NC_037645.1 | XP_026297805.1 | 1204681 | 1212052 | nanos homolog 3 |
| HOG8988 | chr02 | gene_2011_tr2 | 7880423 | 7884430 | chr16 | XP_011159747.3 | 1.8E+07 | 1.8E+07 | NC_037645.1 | XP_026297805.1 | 1204681 | 1212052 | nanos homolog 3-like |
| HOG10889 | chr02 | gene_1435_tr3 | 544379 | 548693 | chr16 | XP_025987290.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | NP_001011595.1 | 7337792 | 7381156 | dopamine receptor 1 isoform X2 |
| HOG10326 | chr02 | gene_2095_tr1 | 1E+07 | 1E+07 | chr16 | XP_011155524.2 | 7905195 | 7910729 | NC_037642.1 | XP_624758.3 | 8394442 | 8398255 | maltase 1 isoform X1 |

| HOG | chr | gene | start | end | chr | XP | E1 | E2 | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG4475 | chr02 | gene_1864_tr3 | 2116467 | 2124441 | chr16 | XP_039314456.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_026301456.1 | 931522 | 938104 | titin homolog isoform X5 |
| HOG10645 | chr02 | gene_1441_tr4 | 590415 | 596041 | chr16 | XP_025990760.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_006560152.2 | 3268423 | 3274574 | phosphatidylinositide phosphatase SAC2 isoform X1 |
| HOG8589 | chr02 | gene_1561_tr1 | 3810773 | 3815356 | chr16 | XP_025990962.1 | 2.6E+07 | 2.6E+07 | NC_037645.1 | XP_003249244.1 | 979012 | 1164689 | AT-rich interactive domain-containing protein 5B isoform X1 |
| HOG7954 | chr02 | gene_1470_tr1 | 1644698 | 1667692 | chr16 | XP_011164684.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_393165.3 | 5851689 | 5869613 | BCL2/adenovirus E1B 19 kDa protein-interacting protein 3 |
| HOG9968 | chr02 | gene_2002_tr3 | 7783173 | 7786678 | chr16 | XP_025994005.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_001122387.2 | 3768277 | 3772417 | mediator of RNA polymerase II transcription subunit 28 |
| HOG9821 | chr02 | gene_1903_tr1 | 2831582 | 2836162 | chr16 | XP_011175376.2 | 1.8E+07 | 1.8E+07 | NC_037652.1 | XP_026301346.1 | 7740600 | 7745432 | probable glutamate receptor isoform X1 |
| HOG9821 | chr02 | gene_1903_tr1 | 2831582 | 2836162 | chr16 | XP_025989474.2 | 1.8E+07 | 1.8E+07 | NC_037652.1 | XP_026301346.1 | 7740600 | 7745432 | glutamate receptor ionotropic delta-1 isoform X1 |
| HOG9821 | chr02 | gene_1903_tr1 | 2831582 | 2836162 | chr16 | XP_039314129.1 | 1.8E+07 | 1.8E+07 | NC_037652.1 | XP_026301346.1 | 7740600 | 7745432 | glutamate receptor-like |
| HOG11158 | chr02 | gene_1655_tr3 | 8557444 | 8592647 | chr16 | XP_025989546.2 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_006560038.1 | 2984878 | 3038381 | MICOS complex subunit Mic60 isoform X1 |
| HOG10644 | chr02 | gene_1828_tr2 | 561841 | 570185 | chr16 | XP_011174263.3 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_026301419.1 | 7392002 | 7399902 | dynactin subunit 1 isoform X1 |
| HOG14655 | chr02 | gene_2068_tr2 | 9260339 | 9263319 | chr16 | XP_039314678.1 | 2.6E+07 | 2.6E+07 | NC_037652.1 | XP_016772200.1 | 7473001 | 7474470 | trypsin isoform X1 |
| HOG9816 | chr02 | gene_1518_tr1 | 2817961 | 2822157 | chr16 | NP_001291533.1 | 1.8E+07 | 1.8E+07 | NC_037652.1 | NP_001011572.1 | 7734777 | 7737849 | transferrin precursor |
| HOG8390 | chr02 | gene_1836_tr2 | 838903 | 853826 | chr16 | XP_039314181.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_006560766.2 | 3898768 | 3913376 | vacuolar protein sorting-associated protein 13B |
| HOG8364 | chr02 | gene_1974_tr1 | 6210501 | 6218674 | chr16 | XP_025987284.1 | 2E+07 | 2E+07 | NC_037638.1 | NP_001207416.1 | 2.7E+07 | 2.7E+07 | peroxidasin homolog pxn-2 |
| HOG8680 | chr02 | gene_1493_tr1 | 2305670 | 2309668 | chr16 | XP_039314217.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_016772183.1 | 3472194 | 3475098 | N-acetyl-D-glucosamine kinase isoform X1 |
| HOG16940 | chr02 | gene_1908_tr2 | 3054063 | 3063955 | chr16 | XP_025989515.1 | 1.7E+07 | 1.7E+07 | NC_037645.1 | XP_001120461.3 | 6259768 | 6268545 | atrial natriuretic peptide receptor 1 isoform X1 |
| HOG10676 | chr02 | gene_1850_tr1 | 1627454 | 1628788 | chr16 | XP_011164612.1 | 1.5E+07 | 1.5E+07 | NC_037644.1 | XP_624054.3 | 2999887 | 3001894 | beta-14-galactosyltransferase 7 |
| HOG8666 | chr02 | gene_1907_tr2 | 3018332 | 3019390 | chr16 | XP_011170096.2 | 1.7E+07 | 1.7E+07 | NC_037644.1 | XP_006560910.1 | 3700856 | 3702254 | tetratricopeptide repeat protein 1 |

| HOG | chr | gene | start | end | chr | XP | val | val | NC | XP | val | val | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG8474 | chr02 | gene_14 68_tr2 | 1625291 | 1627027 | chr16 | XP_0111 64611.1 | 1.5E+07 | 1.5E+07 | NC_0376 41.1 | XP_3950 04.3 | 3349436 | 3351282 | alpha-2-macroglobulin receptor-associated protein |
| HOG8551 | chr02 | gene_16 18_tr1 | 7636416 | 7641713 | chr16 | XP_0111 55778.1 | 1.9E+07 | 1.9E+07 | NC_0376 52.1 | XP_0263 01140.1 | 3837947 | 3846341 | poly [ADP-ribose] polymerase tankyrase isoform X1 |
| HOG8674 | chr02 | gene_19 61_tr4 | 5394807 | 5397857 | chr16 | XP_0111 55894.2 | 2.1E+07 | 2.1E+07 | NC_0376 38.1 | XP_0065 71767.1 | 2.8E+07 | 2.8E+07 | ferrochelatase mitochondrial |
| HOG8001 | chr02 | gene_15 00_tr3 | 2421470 | 2427974 | chr16 | XP_0259 91945.2 | 1.6E+07 | 1.6E+07 | NC_0376 52.1 | XP_0065 60707.1 | 3607609 | 3619427 | cGMP-specific 3'5'-cyclic phosphodiesterase isoform X1 |
| HOG7789 | chr02 | gene_19 42_tr1 | 4466584 | 4467522 | chr16 | XP_0111 56758.1 | 2.5E+07 | 2.5E+07 | NC_0376 45.1 | XP_0065 61531.1 | 5273232 | 5274838 | ELMO domain-containing protein 2 |
| HOG7835 | chr02 | gene_15 09_tr1 | 2545617 | 2546126 | chr16 | XP_0111 56760.1 | 2.5E+07 | 2.5E+07 | NC_0376 52.1 | XP_0263 01309.1 | 2806705 | 2809484 | high mobility group protein HMGI-C |
| HOG1509 2 | chr02 | gene_19 48_tr4 | 4566904 | 4596170 | chr16 | XP_0259 89494.1 | 1.7E+07 | 1.7E+07 | NC_0376 52.1 | XP_0263 01054.1 | 8213522 | 8295995 | tachykinin-like peptides receptor 99D isoform X1 |
| HOG1530 4 | chr02 | gene_15 68_tr1 | 3983445 | 3986702 | chr16 | XP_0259 90992.2 | 2.6E+07 | 2.6E+07 | NC_0376 45.1 | XP_3947 34.5 | 5873379 | 5876958 | molybdenum cofactor sulfurase 3 |
| HOG5340 | chr02 | gene_14 85_tr2 | 2203039 | 2206996 | chr16 | XP_0111 70067.1 | 1.7E+07 | 1.7E+07 | NC_0376 52.1 | XP_0065 57644.1 | 1392562 | 1426650 | BRISC complex subunit FAM175B |
| HOG1000 9 | chr02 | gene_18 30_tr3 | 633905 | 642281 | chr16 | XP_0259 90751.1 | 2.3E+07 | 2.3E+07 | NC_0376 52.1 | NP_0010 11565.1 | 3310369 | 3326664 | octopamine receptor Oamb isoform X1 |
| HOG8785 | chr02 | gene_15 21_tr1 | 2844452 | 2846734 | chr16 | XP_0111 75179.1 | 1.8E+07 | 1.8E+07 | NC_0376 44.1 | XP_3947 00.4 | 3615704 | 3618215 | tubulin epsilon chain |
| HOG8817 | chr02 | gene_16 01_tr1 | 5870774 | 5872329 | chr16 | XP_0393 14709.1 | 2.1E+07 | 2.1E+07 | NC_0376 38.1 | XP_0263 02115.1 | 2.7E+07 | 2.7E+07 | glutamine amidotransferase-like class 1 domain-containing protein 3A mitochondrial |
| HOG8826 | chr02 | gene_19 58_tr1 | 5052175 | 5054200 | chr16 | XP_0393 14418.1 | 1.3E+07 | 1.3E+07 | NC_0376 52.1 | XP_0032 49249.1 | 1910665 | 1913312 | UPF0687 protein C20orf27 homolog |
| HOG6288 | chr02 | gene_19 77_tr1 | 6679051 | 6679895 | chr16 | XP_0111 57914.1 | 2E+07 | 2E+07 | NC_0376 38.1 | XP_0032 51779.1 | 2.6E+07 | 2.6E+07 | peritrophin-1 |
| HOG6288 | chr02 | gene_19 77_tr1 | 6679051 | 6679895 | chr16 | XP_0111 57914.1 | 2E+07 | 2E+07 | NC_0376 38.1 | XP_0032 51791.2 | 2.6E+07 | 2.6E+07 | peritrophin-1 |
| HOG1033 3 | chr02 | gene_21 04_tr1 | 1E+07 | 1E+07 | chr16 | XP_0259 88745.1 | 5561327 | 5577535 | NC_0376 42.1 | XP_3923 66.2 | 8541707 | 8573378 | neuroendocrine convertase 2 |
| HOG1067 5 | chr02 | gene_14 38_tr1 | 587446 | 588814 | chr16 | XP_0111 65354.1 | 2.3E+07 | 2.3E+07 | NC_0376 52.1 | XP_0011 21593.3 | 3266453 | 3268150 | ubiquitin-like domain-containing CTD phosphatase 1 |
| HOG7970 | chr02 | gene_14 54_tr1 | 836500 | 838102 | chr16 | XP_0111 65023.1 | 2.3E+07 | 2.3E+07 | NC_0376 52.1 | XP_0011 20581.3 | 3896389 | 3898161 | RING finger protein 37 isoform X1 |

| HOG | chr | gene | start | end | chr | XP | val1 | val2 | NC | XP2 | start2 | end2 | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG13937 | chr02 | gene_1528_tr1 | 3048276 | 3053169 | chr16 | XP_025989522.1 | 1.7E+07 | 1.7E+07 | NC_037645.1 | XP_396554.4 | 6268589 | 6273975 | cysteine and histidine-rich protein 1 homolog |
| HOG8965 | chr02 | gene_1895_tr1 | 2563359 | 2569774 | chr16 | XP_011156748.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_006563441.2 | 5805728 | 5812112 | KAT8 regulatory NSL complex subunit 3 |
| HOG13454 | chr02 | gene_1657_tr3 | 8679162 | 8687330 | chr16 | XP_011170127.1 | 1.7E+07 | 1.7E+07 | NC_037644.1 | XP_006560197.1 | 2950053 | 2965162 | sorting nexin-27 |
| HOG10661 | chr02 | gene_1985_tr2 | 7165636 | 7171653 | chr16 | XP_039314998.1 | 1.9E+07 | 1.9E+07 | NC_037638.1 | NP_001032395.1 | 2.6E+07 | 2.6E+07 | tyramine receptor 1 isoform X1 |
| HOG8571 | chr02 | gene_1998_tr1 | 7683628 | 7685506 | chr16 | XP_011155767.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_006560753.1 | 3801532 | 3803522 | interleukin enhancer-binding factor 2 |
| HOG15821 | chr02 | gene_1644_tr2 | 8351500 | 8353144 | chr16 | XP_011156738.1 | 2.5E+07 | 2.5E+07 | NC_037644.1 | XP_006560219.1 | 3248508 | 3250496 | protein eyes shut homolog |
| HOG15821 | chr02 | gene_2026_tr2 | 8503816 | 8505439 | chr16 | XP_011156738.1 | 2.5E+07 | 2.5E+07 | NC_037644.1 | XP_006560219.1 | 3248508 | 3250496 | protein eyes shut homolog |
| HOG8212 | chr02 | gene_1871_tr1 | 2237831 | 2247764 | chr16 | XP_039315119.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_392993.2 | 1856328 | 1866310 | cytoplasmic aconitate hydratase isoform X1 |
| HOG7975 | chr02 | gene_1507_tr2 | 2529255 | 2532857 | chr16 | XP_011156721.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_026301084.1 | 7598458 | 7601895 | protein PTCD3 homolog mitochondrial |
| HOG6326 | chr02 | gene_2030_tr2 | 8555871 | 8557342 | chr16 | XP_011170130.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_001121481.1 | 3038290 | 3039526 | 39S ribosomal protein L35 mitochondrial |
| HOG9112 | chr02 | gene_1717_tr5 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170713.2 | 4936386 | 4940192 | NC_037652.1 | XP_006563442.2 | 5802212 | 5805720 | nucleolar complex protein 3 homolog |
| HOG9069 | chr02 | gene_1832_tr1 | 772763 | 775761 | chr16 | XP_011165231.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_001121410.2 | 7569208 | 7574371 | exonuclease 3'-5' domain-containing protein 2 |
| HOG8312 | chr02 | gene_1599_tr1 | 5570470 | 5579495 | chr16 | XP_011158158.1 | 2.1E+07 | 2.1E+07 | NC_037638.1 | XP_026302096.1 | 2.7E+07 | 2.7E+07 | RNA/RNP complex-1-interacting phosphatase |
| HOG9067 | chr02 | gene_1652_tr1 | 8525966 | 8529287 | chr16 | XP_011157594.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_006560751.3 | 3790265 | 3793768 | ralA-binding protein 1 isoform X1 |
| HOG16061 | chr02 | gene_1959_tr1 | 5063881 | 5081427 | chr16 | XP_011155899.1 | 2.2E+07 | 2.2E+07 | NC_037650.1 | XP_026300572.1 | 1236508 | 1297026 | prominin-2 isoform X1 |
| HOG9017 | chr02 | gene_1639_tr1 | 8310629 | 8316151 | chr16 | XP_039314335.1 | 1.8E+07 | 1.8E+07 | NC_037641.1 | XP_395376.3 | 3406438 | 3414622 | cytoplasmic polyadenylation element-binding protein 1 isoform X1 |
| HOG8596 | chr02 | gene_1995_tr2 | 7479424 | 7487751 | chr16 | XP_011161364.1 | 1.9E+07 | 1.9E+07 | NC_037645.1 | XP_395374.2 | 307742 | 316737 | protein O-GlcNAcase isoform X1 |
| HOG9703 | chr02 | gene_2108_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170895.1 | 5437682 | 5441278 | NC_037642.1 | XP_016767840.1 | 8490274 | 8493005 | G protein pathway suppressor 2 |

| HOG | chr | gene | start | end | chr | XP query | val | val | NC | XP subject | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG7956 | chr02 | gene_1630_tr1 | 7797372 | 7801982 | chr16 | XP_025994004.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_006563433.2 | 5782507 | 5786019 | heparanase isoform X2 |
| HOG13640 | chr02 | gene_1617_tr1 | 7628100 | 7633710 | chr16 | XP_011155781.1 | 1.9E+07 | 1.9E+07 | NC_037645.1 | XP_026298277.1 | 1785365 | 1797553 | carbohydrate sulfotransferase 11 |
| HOG8562 | chr02 | gene_1620_tr2 | 7676877 | 7680074 | chr16 | XP_025990739.2 | 2.4E+07 | 2.4E+07 | NC_037652.1 | XP_001120043.2 | 3812648 | 3815066 | venom protease isoform X1 |
| HOG8562 | chr02 | gene_1620_tr2 | 7676877 | 7680074 | chr16 | XP_025990739.2 | 2.4E+07 | 2.4E+07 | NC_037652.1 | XP_016772143.2 | 3808079 | 3810354 | venom protease isoform X1 |
| HOG10456 | chr02 | gene_1893_tr1 | 2554419 | 2555497 | chr16 | XP_011156752.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_625114.2 | 1843783 | 1845473 | phosphoglycerate mutase 2 |
| HOG8653 | chr02 | gene_2020_tr1 | 8478578 | 8479850 | chr16 | XP_025991944.1 | 1.5E+07 | 1.5E+07 | NC_037644.1 | XP_393294.2 | 3030597 | 3031897 | proteasome subunit alpha type-2 |
| HOG7955 | chr02 | gene_1629_tr1 | 7792556 | 7796617 | chr16 | XP_011155754.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_006563436.1 | 5777886 | 5781715 | bromodomain-containing protein 8 isoform X1 |
| HOG10888 | chr02 | gene_1827_tr1 | 550915 | 552731 | chr16 | XP_011174313.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_016772024.1 | 7381274 | 7385104 | methionine--tRNA ligase mitochondrial |
| HOG11604 | chr02 | gene_2125_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011172202.1 | 7980975 | 7984021 | NC_037642.1 | XP_395892.2 | 1.4E+07 | 1.4E+07 | adenosine 3'-phospho 5'-phosphosulfate transporter 1 |
| HOG8194 | chr02 | gene_1598_tr2 | 5537261 | 5538549 | chr16 | XP_011158168.2 | 2.1E+07 | 2.1E+07 | NC_037638.1 | XP_001120941.3 | 2.7E+07 | 2.7E+07 | protein DPCD |
| HOG15174 | chr02 | gene_1914_tr1 | 3146965 | 3149376 | chr16 | XP_011166886.1 | 1.3E+07 | 1.3E+07 | NC_037652.1 | XP_026301294.1 | 2741183 | 2743234 | aspartate aminotransferase cytoplasmic isoform X1 |
| HOG7786 | chr02 | gene_1539_tr1 | 3231656 | 3232729 | chr16 | XP_011175281.1 | 9247439 | 9248968 | NC_037652.1 | XP_396009.2 | 5098612 | 5099750 | transmembrane emp24 domain-containing protein eca |
| HOG7416 | chr02 | gene_2013_tr3 | 7995757 | 8004436 | chr16 | XP_011159511.1 | 1.8E+07 | 1.8E+07 | NC_037644.1 | XP_026297374.1 | 2641369 | 2677023 | putative leucine-rich repeat-containing protein DDB_G0290503 isoform X1 |
| HOG7912 | chr02 | gene_1868_tr3 | 2184745 | 2187525 | chr16 | XP_011170060.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_006557645.1 | 1370111 | 1414274 | hexokinase type 2 isoform X1 |
| HOG8702 | chr02 | gene_1606_tr1 | 7121846 | 7149177 | chr16 | XP_011157760.1 | 1.9E+07 | 2E+07 | NC_037645.1 | XP_006561500.1 | 4556616 | 5047959 | protein slit isoform X1 |
| HOG14992 | chr02 | gene_1663_tr2 | 8977052 | 8994687 | chr16 | XP_011161963.1 | 1.4E+07 | 1.4E+07 | NC_037652.1 | XP_026301277.1 | 665447 | 801495 | PH and SEC7 domain-containing protein isoform X1 |
| HOG8183 | chr02 | gene_2000_tr1 | 7749120 | 7766692 | chr16 | XP_025994007.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_392070.3 | 6475086 | 6484878 | neuronal acetylcholine receptor subunit alpha-10 isoform X1 |
| HOG11435 | chr02 | gene_1550_tr1 | 3471738 | 3473780 | chr16 | XP_011167858.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_016772012.1 | 7613797 | 7616784 | LOW QUALITY PROTEIN: ornithine aminotransferase mitochondrial |

| HOG | chr | gene | start | end | chr | XP | val | val | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG11435 | chr02 | gene_1550_tr1 | 3471738 | 3473780 | chr16 | XP_039314411.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_016772012.1 | 7613797 | 7616784 | ornithine aminotransferase mitochondrial |
| HOG7853 | chr02 | gene_1851_tr1 | 1641742 | 1642471 | chr16 | XP_011164686.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_625228.3 | 5846801 | 5850960 | polyadenylate-binding protein-interacting protein 2 |
| HOG10764 | chr02 | gene_1823_tr1 | 372066 | 375575 | chr16 | XP_011174346.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_625196.1 | 7269445 | 7273061 | wolframin isoform X1 |
| HOG8371 | chr02 | gene_2005_tr2 | 7808831 | 7815525 | chr16 | XP_011159710.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_006563437.1 | 5790955 | 5801638 | glycerol-3-phosphate acyltransferase 1 mitochondrial isoform X1 |
| HOG7900 | chr02 | gene_1611_tr1 | 7219634 | 7221695 | chr16 | XP_011157615.1 | 1.9E+07 | 1.9E+07 | NC_037644.1 | XP_625034.1 | 10870 | 12898 | probable 39S ribosomal protein L45 mitochondrial |
| HOG8547 | chr02 | gene_1975_tr1 | 6219804 | 6222112 | chr16 | XP_025987285.1 | 2E+07 | 2E+07 | NC_037638.1 | NP_001207415.1 | 2.7E+07 | 2.7E+07 | U4/U6 small nuclear ribonucleoprotein Prp3 isoform X1 |
| HOG9702 | chr02 | gene_1708_tr3 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170905.2 | 5441471 | 5447207 | NC_037642.1 | XP_001122827.2 | 8492994 | 8499083 | probable ATP-dependent RNA helicase spindle-E |
| HOG8966 | chr02 | gene_1512_tr1 | 2560443 | 2562269 | chr16 | XP_025990944.2 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_625230.3 | 5836240 | 5838400 | crossover junction endonuclease EME1 |
| HOG6280 | chr02 | gene_1653_tr2 | 8541217 | 8541938 | chr16 | XP_011170133.1 | 1.7E+07 | 1.7E+07 | NC_037645.1 | XP_624594.1 | 5882799 | 5883927 | nucleolar protein 16 |
| HOG11601 | chr02 | gene_2123_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011172207.1 | 8010781 | 8018291 | NC_037642.1 | XP_026296306.1 | 1.4E+07 | 1.4E+07 | ecdysone 20-monooxygenase isoform X1 |
| HOG8535 | chr02 | gene_1986_tr1 | 7206491 | 7209160 | chr16 | XP_011157681.2 | 1.9E+07 | 1.9E+07 | NC_037638.1 | XP_026299704.1 | 2.6E+07 | 2.6E+07 | flap endonuclease GEN |
| HOG8381 | chr02 | gene_1824_tr2 | 377038 | 391543 | chr16 | XP_039314351.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_016772284.1 | 7252464 | 7268830 | alpha-sarcoglycan isoform X1 |
| HOG8386 | chr02 | gene_1628_tr1 | 7787479 | 7788379 | chr16 | XP_011155758.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_006563431.1 | 5773415 | 5774845 | AN1-type zinc finger protein 1 |
| HOG15150 | chr02 | gene_1950_tr1 | 4657835 | 4677811 | chr16 | XP_011170038.2 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_006563712.2 | 8306616 | 8347879 | tubby-related protein 4 |
| HOG17235 | chr02 | gene_1586_tr1 | 5014932 | 5016002 | chr16 | XP_011173007.1 | 1.3E+07 | 1.3E+07 | NC_037652.1 | XP_016771954.1 | 6325010 | 6335391 | general odorant-binding protein 83a |
| HOG6159 | chr02 | gene_2043_tr2 | 8883074 | 8892091 | chr16 | XP_039314679.1 | 2.6E+07 | 2.6E+07 | NC_037651.1 | XP_026300985.1 | 8019082 | 8032890 | G protein-activated inward rectifier potassium channel 3 isoform X1 |
| HOG8475 | chr02 | gene_1913_tr2 | 3137648 | 3139191 | chr16 | XP_039314908.1 | 1.3E+07 | 1.3E+07 | NC_037641.1 | XP_623486.1 | 3351594 | 3352933 | axonemal dynein light intermediate polypeptide 1 isoform X1 |
| HOG8372 | chr02 | gene_1862_tr1 | 2076255 | 2081427 | chr16 | XP_039314429.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_026301458.1 | 818668 | 904370 | ABC transporter G family member 20 isoform X1 |

| HOG | chr | gene | start | end | chr | XP | val1 | val2 | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG13979 | chr02 | gene_1532_tr1 | 3074660 | 3076528 | chr16 | XP_011170104.1 | 1.7E+07 | 1.7E+07 | NC_037644.1 | XP_016768748.1 | 3798288 | 3800919 | m7GpppN-mRNA hydrolase |
| HOG11556 | chr02 | gene_1456_tr1 | 1127222 | 1164340 | chr16 | XP_039314721.1 | 2.4E+07 | 2.4E+07 | NC_037652.1 | XP_026301171.1 | 4283861 | 4316980 | dynein heavy chain 2 axonemal |
| HOG11760 | chr02 | gene_1934_tr1 | 3478522 | 3481130 | chr16 | XP_011167863.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_003250284.1 | 7619194 | 7621734 | zinc finger protein 2 homolog isoform X1 |
| HOG10251 | chr02 | gene_2119_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011171085.2 | 5606080 | 5614138 | NC_037644.1 | XP_026297442.1 | 1.2E+07 | 1.3E+07 | transcription initiation factor TFIID subunit 1 isoform X1 |
| HOG14394 | chr02 | gene_1829_tr6 | 574059 | 576207 | chr16 | XP_011174246.2 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_006563567.1 | 7403869 | 7406083 | tubulin alpha chain |
| HOG14931 | chr02 | gene_2053_tr1 | 9002585 | 9043915 | chr16 | XP_025993470.2 | 1.4E+07 | 1.5E+07 | NC_037652.1 | XP_026301259.1 | 628195 | 659504 | histone-lysine N-methyltransferase 2C isoform X1 |
| HOG8107 | chr02 | gene_1883_tr3 | 2448818 | 2451885 | chr16 | XP_011170016.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_006563505.1 | 6520331 | 6523313 | ATP-dependent RNA helicase abstrakt |
| HOG8366 | chr02 | gene_1503_tr2 | 2470276 | 2473546 | chr16 | XP_025989473.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301194.1 | 2787901 | 2793072 | DNA repair and recombination protein RAD54-like |
| HOG7834 | chr02 | gene_1479_tr1 | 2046560 | 2047282 | chr16 | XP_011170092.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_623809.3 | 2746709 | 2747683 | U3 small nucleolar ribonucleoprotein protein IMP3 |
| HOG9382 | chr02 | gene_2097_tr1 | 1E+07 | 1E+07 | chr16 | XP_011172453.1 | 6907523 | 7038323 | NC_037648.1 | XP_026299571.1 | 3583920 | 4087918 | leucine-rich repeat-containing protein 24 |
| HOG13033 | chr02 | gene_1949_tr1 | 4650675 | 4652957 | chr16 | XP_011170039.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_006563709.1 | 8302408 | 8305934 | sphingosine-1-phosphate lyase |
| HOG14901 | chr02 | gene_1637_tr3 | 8292437 | 8299997 | chr16 | XP_011175125.1 | 1.8E+07 | 1.8E+07 | NC_037644.1 | XP_394701.3 | 2937581 | 2948344 | tyrosine-protein phosphatase non-receptor type 9 |
| HOG10457 | chr02 | gene_1892_tr1 | 2552211 | 2553631 | chr16 | XP_011156754.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_625108.2 | 1847455 | 1849068 | ras-related protein Rab-24 isoform X1 |
| HOG13453 | chr02 | gene_1841_tr4 | 1366591 | 1370489 | chr16 | XP_039314259.1 | 2.4E+07 | 2.4E+07 | NC_037645.1 | XP_003250351.1 | 6317172 | 6321552 | mitogen-activated protein kinase p38b isoform X2 |
| HOG7910 | chr02 | gene_1499_tr1 | 2409028 | 2411611 | chr16 | XP_011164715.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301355.1 | 3591924 | 3595481 | synaptic vesicular amine transporter |
| HOG11500 | chr02 | gene_1719_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011171195.1 | 5675531 | 5688936 | NC_037644.1 | XP_006565503.2 | 9617758 | 9627630 | alpha-2-macroglobulin-like protein 1 isoform X1 |
| HOG15088 | chr02 | gene_1860_tr1 | 2067635 | 2068741 | chr16 | XP_011170091.1 | 1.7E+07 | 1.7E+07 | NC_037648.1 | XP_003250387.2 | 8302490 | 8303724 | putative methyltransferase C9orf114 homolog |
| HOG9021 | chr02 | gene_1477_tr1 | 2023486 | 2024725 | chr16 | XP_011164648.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_001121862.1 | 6204770 | 6207174 | galactoside alpha-(12)-fucosyltransferase 2 |

| HOG | chr | gene | start | end | chr | XP | val1 | val2 | NC | XP | val3 | val4 | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG9021 | chr02 | gene_1478_tr1 | 2033266 | 2034505 | chr16 | XP_011164648.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_001121862.1 | 6204770 | 6207174 | galactoside alpha-(12)-fucosyltransferase 2 |
| HOG10386 | chr02 | gene_2114_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170700.1 | 4928882 | 4932535 | NC_037644.1 | XP_396937.4 | 9612150 | 9614995 | serine/threonine-protein kinase TBK1 |
| HOG10884 | chr02 | gene_1938_tr1 | 3979161 | 3980572 | chr16 | XP_011156782.1 | 2.6E+07 | 2.6E+07 | NC_037645.1 | XP_003249871.1 | 5879791 | 5881685 | cuticle protein 21.3 |
| HOG7903 | chr02 | gene_1999_tr1 | 7693975 | 7736544 | chr16 | XP_039314803.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_026301182.1 | 6486653 | 6518547 | disintegrin and metalloproteinase domain-containing protein 10 |
| HOG13927 | chr02 | gene_1529_tr1 | 3072541 | 3073664 | chr16 | XP_011170107.1 | 1.7E+07 | 1.7E+07 | NC_037644.1 | XP_001120920.1 | 3802778 | 3803824 | phosphomevalonate kinase |
| HOG15180 | chr02 | gene_1536_tr1 | 3144103 | 3144456 | chr16 | XP_011173101.1 | 1.3E+07 | 1.3E+07 | NC_037652.1 | XP_006560007.3 | 2745726 | 2746837 | small integral membrane protein 8 |
| HOG8491 | chr02 | gene_1877_tr1 | 2366397 | 2379139 | chr16 | XP_011164713.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_394109.3 | 3540252 | 3561158 | G protein-coupled receptor kinase 5 |
| HOG16760 | chr02 | gene_1531_tr1 | 3071636 | 3072259 | chr16 | XP_011170108.1 | 1.7E+07 | 1.7E+07 | NC_037644.1 | XP_006560912.1 | 3804858 | 3806666 | 60S ribosomal protein L29-1 |
| HOG10184 | chr02 | gene_2110_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_025988733.1 | 4988454 | 5358114 | NC_037642.1 | XP_026296824.1 | 1.1E+07 | 1.2E+07 | nephrin isoform X1 |
| HOG9222 | chr02 | gene_1931_tr2 | 3437143 | 3452043 | chr16 | XP_011167875.2 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_016772106.2 | 2830254 | 2844944 | serine/threonine-protein kinase WNK1 isoform X1 |
| HOG8120 | chr02 | gene_1964_tr2 | 5421067 | 5421712 | chr16 | XP_025987260.1 | 2.1E+07 | 2.1E+07 | NC_037638.1 | XP_006571776.1 | 2.8E+07 | 2.8E+07 | high affinity copper uptake protein 1 isoform X1 |
| HOG8218 | chr02 | gene_2015_tr2 | 8334196 | 8336195 | chr16 | XP_025991942.1 | 1.5E+07 | 1.5E+07 | NC_037644.1 | XP_006560213.1 | 3023127 | 3027981 | retinol dehydrogenase 14 |
| HOG8218 | chr02 | gene_1648_tr3 | 8484266 | 8486374 | chr16 | XP_025991942.1 | 1.5E+07 | 1.5E+07 | NC_037644.1 | XP_006560213.1 | 3023127 | 3027981 | retinol dehydrogenase 14 |
| HOG14108 | chr02 | gene_1951_tr4 | 4684226 | 4690590 | chr16 | XP_011156808.3 | 2.6E+07 | 2.6E+07 | NC_037652.1 | XP_006563718.1 | 8354130 | 8362747 | NADP-dependent malic enzyme isoform X1 |
| HOG10515 | chr02 | gene_2113_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170726.1 | 4940621 | 4941608 | NC_037642.1 | XP_001121144.1 | 1.4E+07 | 1.4E+07 | membrane magnesium transporter 1 |
| HOG12176 | chr02 | gene_1533_tr1 | 3088468 | 3097185 | chr16 | XP_039314149.1 | 1.2E+07 | 1.2E+07 | NC_037648.1 | XP_006572241.1 | 6999104 | 7009164 | tuberin isoform X1 |
| HOG8670 | chr02 | gene_1976_tr1 | 6241885 | 6280032 | chr16 | XP_011157936.1 | 2E+07 | 2E+07 | NC_037638.1 | XP_026301558.1 | 2.6E+07 | 2.7E+07 | nucleolysin TIAR isoform X1 |
| HOG8186 | chr02 | gene_1600_tr1 | 5695305 | 5708193 | chr16 | XP_011158115.1 | 2.1E+07 | 2.1E+07 | NC_037638.1 | XP_026302091.1 | 2.7E+07 | 2.7E+07 | fibroblast growth factor 17 isoform X1 |

| HOG | chr | gene | start | end | chr | XP | val | val | NC | XP | val | val | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG13766 | chr02 | gene_1923_tr3 | 3233620 | 3236643 | chr16 | XP_011161971.1 | 1.4E+07 | 1.4E+07 | NC_037652.1 | XP_001120689.2 | 813461 | 816339 | STAM-binding protein-like A |
| HOG11167 | chr02 | gene_1707_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_025988731.1 | 5459740 | 5470399 | NC_037638.1 | XP_006558617.1 | 1491999 | 1643433 | protein dimmed |
| HOG8329 | chr02 | gene_1967_tr1 | 5744411 | 5815901 | chr16 | XP_011158097.2 | 2.1E+07 | 2.1E+07 | NC_037638.1 | XP_026299443.1 | 2.7E+07 | 2.7E+07 | transcription factor collier isoform X1 |
| HOG9018 | chr02 | gene_1638_tr3 | 8304929 | 8307892 | chr16 | XP_011175137.1 | 1.8E+07 | 1.8E+07 | NC_037641.1 | XP_623316.3 | 3415321 | 3418434 | cell division cycle protein 16 homolog |
| HOG9345 | chr02 | gene_1670_tr1 | 9380689 | 9384563 | chr16 | XP_011174414.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_026301414.1 | 7193694 | 7205450 | cilia- and flagella-associated protein 206 |
| HOG8448 | chr02 | gene_1492_tr1 | 2302382 | 2303597 | chr16 | XP_011164701.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_001121076.4 | 2687555 | 2688979 | 39S ribosomal protein L21 mitochondrial |
| HOG14252 | chr02 | gene_2121_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011171067.1 | 5602845 | 5604980 | NC_037644.1 | XP_623227.1 | 1.3E+07 | 1.3E+07 | small ubiquitin-related modifier 3-like |
| HOG13663 | chr02 | gene_1464_tr1 | 1371290 | 1373226 | chr16 | XP_011166564.1 | 2.4E+07 | 2.4E+07 | NC_037645.1 | XP_006563902.1 | 6322133 | 6323893 | ATPase ASNA1 homolog |
| HOG8679 | chr02 | gene_1876_tr1 | 2317939 | 2327431 | chr16 | XP_039314208.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_006560700.1 | 3481629 | 3490486 | serine/arginine repetitive matrix protein 1 isoform X1 |
| HOG8382 | chr02 | gene_1874_tr8 | 2290850 | 2301671 | chr16 | XP_011164698.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_016772117.1 | 2676722 | 2687337 | choline/ethanolaminephosphotransferase 1 isoform X7 |
| HOG14370 | chr02 | gene_1937_tr1 | 3975447 | 3977432 | chr16 | XP_011156786.2 | 2.6E+07 | 2.6E+07 | NC_037652.1 | XP_001121815.3 | 7666195 | 7668589 | DNA-directed primase/polymerase protein |
| HOG8373 | chr02 | gene_1484_tr4 | 2102539 | 2111375 | chr16 | XP_011170075.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_026301459.1 | 877167 | 923047 | ABC transporter G family member 20 isoform X1 |
| HOG8373 | chr02 | gene_1484_tr4 | 2102539 | 2111375 | chr16 | XP_011170078.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_026301459.1 | 877167 | 923047 | ABC transporter G family member 20 isoform X1 |
| HOG8373 | chr02 | gene_1484_tr4 | 2102539 | 2111375 | chr16 | XP_039314446.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_026301459.1 | 877167 | 923047 | ABC transporter G family member 20 |
| HOG8373 | chr02 | gene_1484_tr4 | 2102539 | 2111375 | chr16 | XP_039314447.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_026301459.1 | 877167 | 923047 | ABC transporter G family member 20 |
| HOG8373 | chr02 | gene_1484_tr4 | 2102539 | 2111375 | chr16 | XP_039314464.1 | 7607336 | 7622508 | NC_037652.1 | XP_026301459.1 | 877167 | 923047 | ABC transporter G family member 20-like |
| HOG11768 | chr02 | gene_1549_tr1 | 3475886 | 3478196 | chr16 | XP_011167862.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_394061.2 | 7617390 | 7619211 | ornithine aminotransferase mitochondrial |
| HOG13925 | chr02 | gene_1530_tr2 | 3066697 | 3070447 | chr16 | XP_025989517.1 | 1.7E+07 | 1.7E+07 | NC_037644.1 | XP_001122377.3 | 401534 | 405422 | calcium-independent phospholipase A2-gamma isoform X1 |

| HOG | chr | gene | start | end | chr | XP | start | end | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG9346 | chr02 | gene_2073_tr2 | 9386284 | 9390095 | chr16 | XP_011174406.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_392065.2 | 7189452 | 7193590 | upstream activation factor subunit spp27 |
| HOG9203 | chr02 | gene_1514_tr1 | 2574022 | 2585803 | chr16 | XP_011156744.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_623915.2 | 5814862 | 5822003 | beta-TrCP |
| HOG9109 | chr02 | gene_1513_tr1 | 2570850 | 2572513 | chr16 | XP_011156746.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_623887.3 | 5813096 | 5814741 | phospholipid phosphatase 5 |
| HOG8758 | chr02 | gene_1845_tr2 | 1561287 | 1564013 | chr16 | XP_025991919.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_006558326.1 | 1753000 | 1756222 | WD repeat-containing protein 91 isoform X1 |
| HOG15310 | chr02 | gene_2100_tr2 | 1E+07 | 1E+07 | chr16 | XP_011172448.1 | 6867425 | 6871295 | NC_037644.1 | XP_396208.4 | 1.2E+07 | 1.2E+07 | WD repeat-containing protein 13 |
| HOG7201 | chr02 | gene_1647_tr3 | 8480522 | 8481710 | chr16 | XP_011164604.1 | 1.5E+07 | 1.5E+07 | NC_037644.1 | XP_026297479.1 | 3028666 | 3030023 | sperm flagellar protein 1 |
| HOG8084 | chr02 | gene_1973_tr5 | 6204904 | 6208979 | chr16 | XP_011158015.1 | 2E+07 | 2E+07 | NC_037638.1 | NP_001153520.1 | 2.7E+07 | 2.7E+07 | heat shock 70 kDa protein cognate 5 |
| HOG7002 | chr02 | gene_1704_tr2 | 1E+07 | 1E+07 | chr16 | XP_011171042.1 | 5592831 | 5595486 | NC_037644.1 | XP_026297391.1 | 4729272 | 4731424 | pyrimidodiazepine synthase |
| HOG10953 | chr02 | gene_1884_tr1 | 2454899 | 2457123 | chr16 | XP_011170020.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_392704.2 | 3778662 | 3781143 | 4-hydroxybutyrate coenzyme A transferase |
| HOG14306 | chr02 | gene_1437_tr1 | 571010 | 572742 | chr16 | XP_011174253.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_003250277.1 | 7401072 | 7403555 | nuclear inhibitor of protein phosphatase 1 |
| HOG9828 | chr02 | gene_1710_tr2 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170858.2 | 5427830 | 5431731 | NC_037642.1 | XP_395694.2 | 1.1E+07 | 1.1E+07 | dipeptidyl peptidase 3 isoform X1 |
| HOG8216 | chr02 | gene_1489_tr1 | 2231292 | 2234142 | chr16 | XP_011164629.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_006558342.1 | 1868903 | 1877664 | major facilitator superfamily domain-containing protein 6 |
| HOG9083 | chr02 | gene_1475_tr3 | 1802955 | 1807411 | chr16 | XP_011164651.2 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_393162.3 | 5988608 | 5993569 | erythroid differentiation-related factor 1 |
| HOG5859 | chr02 | gene_2032_tr2 | 8695365 | 8696724 | chr16 | XP_025989525.2 | 1.7E+07 | 1.7E+07 | NC_037644.1 | XP_026297736.1 | 2970360 | 2973638 | nucleoside diphosphate kinase homolog 5 isoform X1 |
| HOG10996 | chr02 | gene_1640_tr1 | 8322074 | 8324439 | chr16 | XP_011164633.2 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_026301199.1 | 1851747 | 1854630 | protein RFT1 homolog isoform X1 |
| HOG10996 | chr02 | gene_2021_tr1 | 8496126 | 8498492 | chr16 | XP_011164633.2 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_026301199.1 | 1851747 | 1854630 | protein RFT1 homolog isoform X1 |
| HOG7896 | chr02 | gene_1610_tr2 | 7209666 | 7211413 | chr16 | XP_011157667.1 | 1.9E+07 | 1.9E+07 | NC_037643.1 | XP_006560095.2 | 3695746 | 3698098 | charged multivesicular body protein 7 |
| HOG9911 | chr02 | gene_1727_tr3 | 1.1E+07 | 1.1E+07 | chr16 | XP_011172199.1 | 7964222 | 7972625 | NC_037642.1 | XP_003250531.1 | 8442736 | 8450993 | RNA-binding protein Nova-1 isoform X1 |

| HOG | chr | gene | start | end | chr | XP | start | end | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG10278 | chr02 | gene_2120_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011171555.1 | 5614286 | 5616241 | NC_037644.1 | XP_006568357.1 | 1.2E+07 | 1.2E+07 | protein EFR3 homolog B-like |
| HOG8681 | chr02 | gene_1495_tr1 | 2328488 | 2329466 | chr16 | XP_011164706.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_016772335.1 | 3491262 | 3492418 | glutaredoxin-3 |
| HOG9202 | chr02 | gene_2028_tr2 | 8532952 | 8538883 | chr16 | XP_039315048.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_006563445.1 | 5825072 | 5832919 | exportin-7 isoform X1 |
| HOG13963 | chr02 | gene_1842_tr1 | 1373592 | 1375505 | chr16 | XP_025990729.1 | 2.4E+07 | 2.4E+07 | NC_037645.1 | XP_395385.3 | 6323956 | 6326061 | 2-amino-3-ketobutyrate coenzyme A ligase mitochondrial |
| HOG9344 | chr02 | gene_2027_tr2 | 8520769 | 8523402 | chr16 | XP_025990966.2 | 2.5E+07 | 2.5E+07 | NC_037644.1 | XP_396549.3 | 3266949 | 3271873 | glucose dehydrogenase [FAD quinone] |
| HOG14365 | chr02 | gene_1567_tr2 | 3971998 | 3974172 | chr16 | XP_011156788.1 | 2.6E+07 | 2.6E+07 | NC_037652.1 | XP_006563616.1 | 7663677 | 7666565 | stAR-related lipid transfer protein 7 mitochondrial isoform X1 |
| HOG7945 | chr02 | gene_1632_tr1 | 7829322 | 7833913 | chr16 | XP_011159700.2 | 1.9E+07 | 1.9E+07 | NC_037644.1 | XP_016768580.2 | 754098 | 758989 | biorientation of chromosomes in cell division protein 1-like 1 |
| HOG15914 | chr02 | gene_2001_tr1 | 7768659 | 7770011 | chr16 | XP_039314809.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_003249733.2 | 3751884 | 3754310 | F-box/LRR-repeat protein 7 |
| HOG9019 | chr02 | gene_1857_tr1 | 2040226 | 2042758 | chr16 | XP_011164641.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_006563482.1 | 6212313 | 6215474 | pantothenate kinase 3 isoform X1 |
| HOG16025 | chr02 | gene_1856_tr1 | 2036593 | 2038769 | chr16 | XP_011164646.2 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_006563481.1 | 6208825 | 6211211 | dynein regulatory complex subunit 4 |
| HOG8672 | chr02 | gene_1960_tr2 | 5155982 | 5192533 | chr16 | XP_011155897.1 | 2.1E+07 | 2.2E+07 | NC_037638.1 | XP_026297862.1 | 2.8E+07 | 2.8E+07 | leishmanolysin-like peptidase isoform X1 |
| HOG8716 | chr02 | gene_1849_tr1 | 1621609 | 1624476 | chr16 | XP_011164610.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_026301314.1 | 1836789 | 1840008 | X-ray repair cross-complementing protein 5 |
| HOG16020 | chr02 | gene_1983_tr3 | 6785469 | 6786881 | chr16 | XP_011157834.1 | 2E+07 | 2E+07 | NC_037645.1 | XP_026298216.1 | 4511467 | 4555900 | trypsin 5G1 isoform X1 |
| HOG8441 | chr02 | gene_1945_tr2 | 4495244 | 4499484 | chr16 | XP_011170055.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_026301184.1 | 6470416 | 6473217 | 60S ribosomal protein L4 |
| HOG15043 | chr02 | gene_1668_tr3 | 9268769 | 9269450 | chr16 | XP_025990985.1 | 2.6E+07 | 2.6E+07 | NC_037652.1 | XP_006563593.1 | 7554461 | 7556096 | guanine nucleotide exchange factor MSS4 homolog |
| HOG8768 | chr02 | gene_1642_tr1 | 8331174 | 8333845 | chr16 | XP_011164602.1 | 1.5E+07 | 1.5E+07 | NC_037644.1 | XP_393293.2 | 3017741 | 3023131 | esterase FE4 |
| HOG8768 | chr02 | gene_2022_tr1 | 8486725 | 8489396 | chr16 | XP_011164602.1 | 1.5E+07 | 1.5E+07 | NC_037644.1 | XP_393293.2 | 3017741 | 3023131 | esterase FE4 |
| HOG15311 | chr02 | gene_1920_tr3 | 3192824 | 3209943 | chr16 | XP_039314111.1 | 9731785 | 9888219 | NC_037652.1 | XP_026301325.1 | 5106146 | 5150785 | protein TANC2 isoform X1 |

| HOG | chr | gene | start | end | chr | XP | val1 | val2 | NC | XP | val3 | val4 | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG8755 | chr02 | gene_1865_tr1 | 2138813 | 2167159 | chr16 | XP_025989484.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_026301069.1 | 951021 | 1360467 | max dimerization protein 1 |
| HOG9913 | chr02 | gene_1699_tr1 | 1E+07 | 1E+07 | chr16 | XP_011155530.1 | 7943071 | 7946078 | NC_037642.1 | XP_016767933.1 | 8426297 | 8428192 | scaffold protein salvador isoform X1 |
| HOG8128 | chr02 | gene_1869_tr4 | 2208105 | 2212173 | chr16 | XP_011170070.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_026301300.1 | 1426960 | 1430242 | Hermansky-Pudlak syndrome 1 protein homolog isoform X1 |
| HOG8675 | chr02 | gene_1595_tr1 | 5397895 | 5399539 | chr16 | XP_011155895.1 | 2.1E+07 | 2.1E+07 | NC_037638.1 | XP_624562.2 | 2.8E+07 | 2.8E+07 | EEF1A lysine methyltransferase 2 |
| HOG10328 | chr02 | gene_1700_tr1 | 1E+07 | 1E+07 | chr16 | XP_039314545.1 | 7910802 | 7913008 | NC_037642.1 | XP_393310.1 | 8398332 | 8399874 | JNK1/MAPK8-associated membrane protein |
| HOG8775 | chr02 | gene_1928_tr2 | 3296805 | 3305510 | chr16 | XP_011167883.1 | 1.5E+07 | 1.5E+07 | NC_037652.1 | XP_625223.2 | 6603005 | 6841964 | complexin isoform X1 |
| HOG9419 | chr02 | gene_1491_tr1 | 2250023 | 2251767 | chr16 | XP_011164693.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301223.1 | 2612739 | 2632495 | serine/arginine-rich splicing factor 1A |
| HOG13903 | chr02 | gene_1822_tr4 | 312935 | 341739 | chr16 | XP_039314666.1 | 2.6E+07 | 2.6E+07 | NC_037652.1 | XP_006563397.1 | 7474685 | 7550203 | glutamate-gated chloride channel isoform X1 |
| HOG7100 | chr02 | gene_1896_tr3 | 2681020 | 2684878 | chr16 | XP_011170134.2 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_001121888.2 | 7690528 | 7693016 | proclotting enzyme |
| HOG15309 | chr02 | gene_1702_tr3 | 1E+07 | 1E+07 | chr16 | XP_025993055.2 | 6869805 | 6878400 | NC_037644.1 | XP_026297764.1 | 1.2E+07 | 1.2E+07 | STE20-like serine/threonine-protein kinase isoform X1 |
| HOG16229 | chr02 | gene_2006_tr1 | 7846282 | 7848004 | chr16 | XP_039314701.1 | 1.8E+07 | 1.8E+07 | NC_037644.1 | XP_016768435.2 | 769780 | 772899 | meiosis-specific nuclear structural protein 1 |
| HOG9610 | chr02 | gene_2109_tr2 | 1.1E+07 | 1.1E+07 | chr16 | XP_039315102.1 | 5431828 | 5434733 | NC_037642.1 | XP_006564855.1 | 8531552 | 8532956 | protein FAM50 homolog isoform X1 |
| HOG15284 | chr02 | gene_2098_tr1 | 1E+07 | 1E+07 | chr16 | XP_011171593.1 | 7777629 | 7779233 | NC_037648.1 | XP_016773031.1 | 4260226 | 4262644 | 60S ribosomal protein L28 |
| HOG8086 | chr02 | gene_1551_tr1 | 3617146 | 3630019 | chr16 | XP_039314948.1 | 1.6E+07 | 1.7E+07 | NC_037652.1 | XP_006563612.1 | 7625070 | 7661948 | DE-cadherin isoform X1 |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164671.2 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301363.1 | 5875593 | 5877977 | retinol-binding protein pinta isoform X1 |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164674.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301363.1 | 5875593 | 5877977 | retinol-binding protein pinta |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164676.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301363.1 | 5875593 | 5877977 | retinol-binding protein pinta |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164678.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301363.1 | 5875593 | 5877977 | retinol-binding protein pinta isoform X1 |

| HOG | chr | gene | start | end | chr16 | XP | val1 | val2 | NC | XP2 | start2 | end2 | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164680.2 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301363.1 | 5875593 | 5877977 | retinol-binding protein pinta |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_011164681.2 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301363.1 | 5875593 | 5877977 | retinol-binding protein pinta |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_025991973.2 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301363.1 | 5875593 | 5877977 | retinol-binding protein pinta |
| HOG9892 | chr02 | gene_1471_tr3 | 1680476 | 1683069 | chr16 | XP_039314934.1 | 2.4E+07 | 2.4E+07 | NC_037652.1 | XP_026301363.1 | 5875593 | 5877977 | alpha-tocopherol transfer protein-like isoform X1 |
| HOG8295 | chr02 | gene_1452_tr1 | 806294 | 808604 | chr16 | XP_011165153.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_393237.4 | 3871533 | 3876482 | alpha-tocopherol transfer protein-like |
| HOG8028 | chr02 | gene_1991_tr1 | 7234859 | 7238940 | chr16 | XP_025987316.1 | 1.9E+07 | 1.9E+07 | NC_037645.1 | XP_026297809.1 | 13090 | 289186 | tyrosine-protein kinase transmembrane receptor Ror isoform X1 |
| HOG15280 | chr02 | gene_2099_tr1 | 1E+07 | 1E+07 | chr16 | XP_011171597.1 | 7779426 | 7780932 | NC_037648.1 | XP_001123012.2 | 4261400 | 4262894 | S-adenosylmethionine sensor upstream of mTORC1 |
| HOG10512 | chr02 | gene_2112_tr4 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170750.1 | 4946711 | 4958954 | NC_037642.1 | XP_026296543.1 | 1.4E+07 | 1.4E+07 | glutathione hydrolase 1 proenzyme |
| HOG8690 | chr02 | gene_1940_tr1 | 4381399 | 4384558 | chr16 | XP_011156768.2 | 2.5E+07 | 2.5E+07 | NC_037645.1 | NP_001035349.1 | 5359323 | 5361065 | alpha-glucosidase |
| HOG15030 | chr02 | gene_1838_tr1 | 1282051 | 1283833 | chr16 | XP_011164926.1 | 2.4E+07 | 2.4E+07 | NC_037644.1 | XP_393404.2 | 3739216 | 3754819 | COMM domain-containing protein 4 |
| HOG8708 | chr02 | gene_1835_tr1 | 835166 | 835865 | chr16 | XP_011165066.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_625159.2 | 3895314 | 3896376 | H/ACA ribonucleoprotein complex subunit 2-like protein |
| HOG11706 | chr02 | gene_2107_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_039314193.1 | 5451045 | 5456789 | NC_037638.1 | XP_026298774.1 | 1672100 | 1674611 | cytoglobin-2 isoform X1 |
| HOG8493 | chr02 | gene_1496_tr5 | 2342986 | 2354028 | chr16 | XP_039314212.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_026301448.1 | 3504722 | 3533434 | large neutral amino acids transporter small subunit 1 isoform X1 |
| HOG8949 | chr02 | gene_2008_tr3 | 7860224 | 7865015 | chr16 | XP_025989569.1 | 1.8E+07 | 1.8E+07 | NC_037645.1 | XP_016769182.2 | 1189085 | 1193536 | phenoloxidase-activating factor 2 isoform X1 |
| HOG12499 | chr02 | gene_1669_tr1 | 9279870 | 9280947 | chr16 | XP_011165269.1 | 2.6E+07 | 2.6E+07 | NC_037651.1 | XP_026300986.1 | 8014984 | 8016212 | coiled-coil domain-containing protein 137 |
| HOG7949 | chr02 | gene_1516_tr2 | 2767965 | 2789287 | chr16 | XP_039315007.1 | 1.8E+07 | 1.8E+07 | NC_037652.1 | XP_394067.4 | 7697592 | 7717297 | semaphorin-5B isoform X1 |
| HOG7841 | chr02 | gene_1672_tr5 | 9719909 | 9741305 | chr16 | XP_039314315.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_006563386.2 | 6989419 | 7037855 | mushroom body large-type Kenyon cell-specific protein 1 isoform X1 |
| HOG7867 | chr02 | gene_1881_tr4 | 2434383 | 2435726 | chr16 | XP_025991950.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_001122074.3 | 3621194 | 3625443 | golgin subfamily A member 7 isoform X1 |

| HOG16123 | chr02 | gene_2127_tr3 | 1.1E+07 | 1.1E+07 | chr16 | XP_011172322.1 | 6128726 | 6131665 | NC_037642.1 | XP_006562189.1 | 5906010 | 5908473 | cyclin-dependent kinases regulatory subunit |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG8684 | chr02 | gene_1886_tr1 | 2486426 | 2487312 | chr16 | XP_039315036.1 | 1.6E+07 | 1.6E+07 | NC_037652.1 | XP_624537.1 | 2776324 | 2778184 | U6 snRNA-associated Sm-like protein LSm8 |
| HOG14776 | chr02 | gene_2116_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011171124.1 | 5638121 | 5639619 | NC_037644.1 | XP_026297475.1 | 8713619 | 8715589 | ER lumen protein-retaining receptor |
| HOG8190 | chr02 | gene_1965_tr2 | 5511870 | 5514331 | chr16 | XP_011155883.1 | 2.1E+07 | 2.1E+07 | NC_037638.1 | XP_006571765.2 | 2.7E+07 | 2.7E+07 | ATP-dependent RNA helicase vasa |
| HOG8190 | chr02 | gene_1965_tr2 | 5511870 | 5514331 | chr16 | XP_025989543.2 | 1.8E+07 | 1.8E+07 | NC_037638.1 | XP_006571765.2 | 2.7E+07 | 2.7E+07 | ATP-dependent RNA helicase vasa |
| HOG8190 | chr02 | gene_1965_tr2 | 5511870 | 5514331 | chr16 | XP_039314125.1 | 1.8E+07 | 1.8E+07 | NC_037638.1 | XP_006571765.2 | 2.7E+07 | 2.7E+07 | ATP-dependent RNA helicase vasa |
| HOG10444 | chr02 | gene_1517_tr1 | 2807371 | 2815016 | chr16 | XP_025989549.2 | 1.8E+07 | 1.8E+07 | NC_037652.1 | XP_026301341.1 | 7727328 | 7733724 | LOW QUALITY PROTEIN: cilia- and flagella-associated protein 43-like |
| HOG10356 | chr02 | gene_1443_tr1 | 602260 | 604541 | chr16 | XP_011165293.1 | 2.3E+07 | 2.3E+07 | NC_037652.1 | XP_624354.1 | 3279596 | 3282286 | serine protease HTRA2 mitochondrial isoform X1 |
| HOG15689 | chr02 | gene_1731_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011172321.1 | 6131662 | 6133555 | NC_037642.1 | XP_395839.1 | 5898782 | 5899801 | vacuolar protein-sorting-associated protein 25 |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_011155906.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_016772002.2 | 6967145 | 6969218 | odorant receptor 49b isoform X1 |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_011155906.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_016772004.2 | 6964546 | 6966223 | odorant receptor 49b isoform X1 |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_025987223.2 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_016772002.2 | 6967145 | 6969218 | odorant receptor 2a-like |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_025987223.2 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_016772004.2 | 6964546 | 6966223 | odorant receptor 2a-like |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_025987227.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_016772002.2 | 6967145 | 6969218 | odorant receptor 82a-like |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_025987227.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_016772004.2 | 6964546 | 6966223 | odorant receptor 82a-like |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_025987228.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_016772002.2 | 6967145 | 6969218 | odorant receptor 22c-like |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_025987228.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_016772004.2 | 6964546 | 6966223 | odorant receptor 22c-like |
| HOG5520 | chr02 | gene_1808_tr1 | 22721 | 45107 | chr16 | XP_039314325.1 | 2.2E+07 | 2.2E+07 | NC_037652.1 | XP_016772002.2 | 6967145 | 6969218 | odorant receptor 9a-like |

| | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG5520 | chr02 | gene_18 08_tr1 | 22721 | 45107 | chr16 | XP_0393 14325.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72004.2 | 6964546 | 6966223 | odorant receptor 9a-like |
| HOG5520 | chr02 | gene_18 08_tr1 | 22721 | 45107 | chr16 | XP_0393 14405.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72002.2 | 6967145 | 6969218 | odorant receptor 4-like |
| HOG5520 | chr02 | gene_18 08_tr1 | 22721 | 45107 | chr16 | XP_0393 14405.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72004.2 | 6964546 | 6966223 | odorant receptor 4-like |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0111 55906.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72002.2 | 6967145 | 6969218 | odorant receptor 49b isoform X1 |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0111 55906.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72004.2 | 6964546 | 6966223 | odorant receptor 49b isoform X1 |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0259 87223.2 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72002.2 | 6967145 | 6969218 | odorant receptor 2a-like |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0259 87223.2 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72004.2 | 6964546 | 6966223 | odorant receptor 2a-like |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0259 87227.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72002.2 | 6967145 | 6969218 | odorant receptor 82a-like |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0259 87227.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72004.2 | 6964546 | 6966223 | odorant receptor 82a-like |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0259 87228.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72002.2 | 6967145 | 6969218 | odorant receptor 22c-like |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0259 87228.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72004.2 | 6964546 | 6966223 | odorant receptor 22c-like |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0393 14325.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72002.2 | 6967145 | 6969218 | odorant receptor 9a-like |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0393 14325.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72004.2 | 6964546 | 6966223 | odorant receptor 9a-like |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0393 14405.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72002.2 | 6967145 | 6969218 | odorant receptor 4-like |
| HOG5520 | chr02 | gene_18 13_tr2 | 82571 | 92449 | chr16 | XP_0393 14405.1 | 2.2E+07 | 2.2E+07 | NC_0376 52.1 | XP_0167 72004.2 | 6964546 | 6966223 | odorant receptor 4-like |
| HOG6111 | chr02 | gene_14 48_tr1 | 776848 | 779010 | chr16 | XP_0111 65211.1 | 2.3E+07 | 2.3E+07 | NC_0376 52.1 | XP_0263 01142.1 | 3846445 | 3848620 | zinc finger protein 699 |
| HOG8763 | chr02 | gene_14 65_tr3 | 1557926 | 1560610 | chr16 | XP_0259 91920.1 | 1.5E+07 | 1.5E+07 | NC_0376 52.1 | XP_0263 01246.1 | 1751161 | 1752770 | transcription initiation protein SPT3 homolog isoform X1 |
| HOG1583 9 | chr02 | gene_15 82_tr1 | 4561481 | 4562970 | chr16 | XP_0111 70040.1 | 1.7E+07 | 1.7E+07 | NC_0376 52.1 | XP_0065 63707.1 | 8209961 | 8211902 | peptide deformylase mitochondrial |

| HOG | chr | gene | start | end | chr | XP | start | end | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG10089 | chr02 | gene_1711_tr1 | 1.1E+07 | 1.1E+07 | chr16 | XP_011170848.1 | 5414478 | 5426727 | NC_037642.1 | XP_006567697.2 | 1.1E+07 | 1.1E+07 | DNA polymerase alpha subunit B |
| HOG14128 | chr02 | gene_1584_tr1 | 4680005 | 4681037 | chr16 | XP_011156805.1 | 2.6E+07 | 2.6E+07 | NC_037652.1 | XP_393179.1 | 8349465 | 8350810 | RWD domain-containing protein 1 |
| HOG16309 | chr02 | gene_1888_tr2 | 2509939 | 2511890 | chr16 | XP_011156722.1 | 2.5E+07 | 2.5E+07 | NC_037652.1 | XP_006563601.1 | 7588891 | 7591104 | major facilitator superfamily domain-containing protein 10 isoform X1 |
| HOG11603 | chr02 | gene_1724_tr2 | 1.1E+07 | 1.1E+07 | chr16 | XP_011172210.1 | 8043481 | 8046799 | NC_037642.1 | XP_026296394.1 | 1.4E+07 | 1.4E+07 | cyclin-dependent kinase 8 isoform X1 |
| HOG9369 | chr02 | gene_1635_tr1 | 7988690 | 7992369 | chr16 | XP_011159533.1 | 1.8E+07 | 1.8E+07 | NC_037645.1 | XP_394713.3 | 1297182 | 1300775 | conserved oligomeric Golgi complex subunit 7 |
| HOG8114 | chr02 | gene_1588_tr1 | 5024531 | 5025745 | chr16 | XP_039314227.1 | 1.3E+07 | 1.3E+07 | NC_037652.1 | XP_026301043.1 | 6339314 | 6402471 | dopamine receptor 2 |
| HOG16290 | chr02 | gene_1631_tr1 | 7804773 | 7807974 | chr16 | XP_039314810.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_395193.1 | 5788514 | 5790792 | tektin-3-like |
| HOG14348 | chr02 | gene_1515_tr1 | 2610166 | 2678821 | chr16 | XP_025989465.2 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_026301090.1 | 7669569 | 7689024 | atrial natriuretic peptide receptor 1 isoform X1 |
| HOG15119 | chr02 | gene_1480_tr1 | 2058079 | 2064661 | chr16 | XP_039315041.1 | 1.7E+07 | 1.7E+07 | NC_037652.1 | XP_624412.2 | 2755253 | 2760672 | sodium-coupled neutral amino acid transporter 9 isoform X1 |
| HOG8031 | chr02 | gene_2003_tr1 | 7790245 | 7791748 | chr16 | XP_011155757.1 | 1.9E+07 | 1.9E+07 | NC_037652.1 | XP_026301239.1 | 5776053 | 5777714 | tetraspanin-33 |

| HOG | chr | gene | start | end | chr | XP | start | end | NC | XP | start | end | description |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| HOG8763 | chr02 | gene_1465_tr3 | 1557926 | 1560610 | chr16 | XP_025991920.1 | 15222779 | 15227463 | NC_037652.1 | XP_026301246.1 | 1751161 | 1752770 | transcription initiation protein SPT3 homolog isoform X1 |
| HOG15839 | chr02 | gene_1582_tr1 | 4561481 | 4562970 | chr16 | XP_011170040.1 | 16662165 | 16664578 | NC_037652.1 | XP_006563707.1 | 8209961 | 8211902 | peptide deformylase, mitochondrial |
| HOG14128 | chr02 | gene_1584_tr1 | 4680005 | 4681037 | chr16 | XP_011156805.1 | 25882931 | 25886321 | NC_037652.1 | XP_393179.1 | 8349465 | 8350810 | RWD domain-containing protein 1 |
| HOG16309 | chr02 | gene_1888_tr2 | 2509939 | 2511890 | chr16 | XP_011156722.1 | 24656092 | 24662689 | NC_037652.1 | XP_006563601.1 | 7588891 | 7591104 | major facilitator superfamily domain-containing protein 10 isoform X1 |
| HOG8114 | chr02 | gene_1588_tr1 | 5024531 | 5025745 | chr16 | XP_039314227.1 | 13391052 | 13471488 | NC_037652.1 | XP_026301043.1 | 6339314 | 6402471 | dopamine receptor 2 |
| HOG16290 | chr02 | gene_1631_tr1 | 7804773 | 7807974 | chr16 | XP_039314810.1 | 18574045 | 18577426 | NC_037652.1 | XP_395193.1 | 5788514 | 5790792 | tektin-3-like |
| HOG14348 | chr02 | gene_1515_tr1 | 2610166 | 2678821 | chr16 | XP_025989465.2 | 17347717 | 17438257 | NC_037652.1 | XP_026301090.1 | 7669569 | 7689024 | atrial natriuretic peptide receptor 1 isoform X1 |
| HOG15119 | chr02 | gene_1480_tr1 | 2058079 | 2064661 | chr16 | XP_039315041.1 | 16954588 | 16968602 | NC_037652.1 | XP_624412.2 | 2755253 | 2760672 | sodium-coupled neutral amino acid transporter 9 isoform X1 |
| HOG8031 | chr02 | gene_2003_tr1 | 7790245 | 7791748 | chr16 | XP_011155757.1 | 18591135 | 18595161 | NC_037652.1 | XP_026301239.1 | 5776053 | 5777714 | tetraspanin-33 |