1

Repeated evolution of supergenes on an ancient social chromosome

Aparna Lajmi[1,2,*], Pnina Cohen[1,3], Chih-Chi Lee[1,4,5], Zeev Frenkel[1], Yoann Pellen[1], Eyal Privman[1,*]

[1] Department of Evolutionary and Environmental Biology, Institute of Evolution,
University of Haifa, Haifa, Israel

[2] Department of Life Sciences, Gandhi Institute of Technology and Management, Visakhapatnam, India

[3] Department of Archaeology, Zinman Institute, University of Haifa, Haifa, Israel.

[4] Agricultural Biotechnology Research Center, Academia Sinica, Taipei, Taiwan

[5] Biotechnology Center in Southern Taiwan, Academia Sinica, Tainan, Taiwan

* Co-corresponding authors

Abstract

Supergenes are non-recombining chromosomal regions that code for complex polymorphic traits. Advances in population genomics have uncovered supergenes associated with diverse traits, ranging from butterfly wing patterns to floral morphs. In ants, two supergenes on non-homologous "social chromosomes" in *Solenopsis* and *Formica* are associated with social polymorphism, with either single queen (monogyne) or multiple queens (polygyne) colonies. We discovered a new supergene associated with similar polymorphism in the desert ant *Cataglyphis niger*. Despite *Cataglyphis* being more closely related to *Formica* than *Solenopsis*, its social chromosome is homologous to that of *Solenopsis*, with conservation of synteny in gene content and order. This suggests that the social chromosome is ancient, dating back to the common ancestor of *Solenopsis* and *Cataglyphis*, at least 90 million years ago. Previous work dated the *Solenopsis* supergene to be only one million years old, therefore, we propose that the two supergenes evolved independently in the two diverged lineages on this ancient social chromosome. Comparative analysis of hymenopteran genomes further revealed a bee chromosome homologous to the ants' social chromosome. The conservation of the gene set on these homologous chromosomes in ants and bees suggests that this chromosome plays an important role in social behavior across social Hymenoptera. We propose that the conserved gene set in the social chromosome was repeatedly used as a pre-adapted toolkit for the evolution of social traits in general, and specifically in the evolution of polygyne social structure in ants.

Keywords: Supergene, social chromosome, polygyne, complex trait, Hymenoptera, social insects, phylogenomics, population genomics

1. Introduction

Convergent evolution is a common phenomenon across the tree of life[1], yet the genomic architectures that facilitate the repeated evolution of complex traits remain poorly understood. Social behavior and the underlying sociobiological traits, particularly in Hymenoptera (ants, bees, and wasps), exemplify complex phenotypes that have evolved repeatedly in multiple lineages. Previous literature suggests that the evolution of sociality in diverse hymenopteran lineages involved the reuse of a core set of conserved genes that were originally involved in other biological processes, a hypothesis termed the genetic "toolkit" hypothesis[2]. This hypothesis suggests that genes originally involved in solitary behaviors or other biological processes were co-opted to facilitate sociality, in line with the notion that evolutionary innovations often emerge from preexisting genetic frameworks. For example, the *foraging* gene that is conserved across insects and regulates foraging behavior in *Drosophila melanogaster*[3] has been co-opted in bees and ants to regulate caste-specific behaviors[4,5]. Foraging and brood care behaviors in the worker caste are key to the success of social insect colonies. Maternal care, which is not common in insects but found in many solitary wasps, has been proposed to be a pre-adaptation for the evolution of sociality[6,7]. Such pre-adaptations in solitary Hymenoptera may have facilitated the repeated evolution of sociobiological traits in this clade.

Ants have repeatedly evolved the trait of multiple-queen colonies (polygyny) from an ancestral phenotype of a single-queen colony (monogyny)[8,9]. Polygyny is a complex trait involving a suite of life-history, behavioral, physiological, and morphological modifications[10,11,12]. Moreover, it results in nestmates that have very low relatedness, which undermines the evolution of cooperation by means of kin selection[9,12]. Hughes *et al.*[8] reported that polygyne colonies were observed for 54 out of 130 (42%) ant species included in that study. Some species of ants are socially polymorphic, where both monogyne and polygyne colonies coexist in the same population. A genetic basis of this social polymorphism has been documented and examined in detail in two distantly related ant lineages, *Solenopsis* fire ants and *Formica* wood ants. In both lineages, a supergene was found to be associated with this polymorphism[13]. These supergenes are

chromosomal regions with suppressed recombination, containing multiple tightly linked loci that code for a complex trait. The most well-known type of supergenes are sex chromosomes, but virtually any other complex polymorphic trait may be coded by a supergene[14]. The supergenes responsible for variation in queen number in ants offer an opportunity to investigate the genomic basis of the recurrent evolution of complex traits.

Independently evolved supergenes on non-homologous chromosomes underlie similar social polymorphism in both *Solenopsis* and *Formica*[15,16]. However, there are several important similarities and differences in how the supergene is maintained. The first social supergene was discovered by Wang and colleagues in the red fire ant *Solenopsis invicta*, which is native to tropical South America[15]. In monogyne colonies of this species, the queen and the workers are MM homozygotes at the supergene (the *Solenopsis* supergene haplotypes were originally named *SB* and *Sb*, but we are using M for monogyne-associated and P for polygyne-associated haplotypes across all species for consistency). On the other hand, the polygyne queens are always MP heterozygous at the supergene, and produce workers of all three genotypes: MM, MP, and rarely PP, by mating with both M and P haploid males[17]. The absence of the PP genotype in queens led Wang and his colleagues to refer to this chromosome as a "Y-like social chromosome". This supergene was found across a few closely related, socially polymorphic *Solenopsis* species that speciated approximately one million years ago (MYA)[18]. The formation of the supergene was dated approximately to the same time based on the divergence of the M and P haplotype sequences[19,20]. In the Alpine wood ant *Formica selysi*, where the second social supergene was discovered, monogyne queens and workers are MM homozygotes at the supergene while the polygyne queens and workers are either PP homozygotes or MP heterozygotes. Unlike in *S. invicta*, MM genotypes in polygyne colonies are eliminated in this species, which was suggested to be due to a maternal effect killing[21]. Supergenes homologous to the *F. selysi* supergene were found in numerous *Formica* species and therefore evolved in their common ancestor at least 30 MYA[22]. *Formica* and *Solenopsis* belong to two different ant subfamilies (Formicinae and Myrmicinae) that diverged more than 90 MYA[23]. The young age of the two supergenes relative to the lineage divergence, and

the fact that they reside on non-homologous chromosomes, imply that the two

supergenes have independent origins.

The desert ant *Cataglyphis niger* is a socially polymorphic species, with populations

consisting of both monogyne and polygyne colonies. These populations were initially

thought to consist of multiple species, based on divergent mitochondrial sequences

(mitotypes) that are associated with the social form[24]. However, nuclear genomic

sequencing revealed no genetic divergence between monogyne and polygyne colonies,

demonstrating that *C. niger* is a socially polymorphic species[25]. The genera *Cataglyphis*

and *Formica* belong to the same tribe Formicini within the Formicinae subfamily, which

diverged approximately 40 MYA[23]. Therefore, *C. niger* is an interesting target species

for the study of the evolution of social polymorphism across the ant phylogeny.

Here, we examined the genomic basis of social polymorphism in *C. niger* by generating

reduced representation genome sequencing data for 672 individuals from 30 nests

representing both social forms from a single population. The social form of the nests

was ascertained by estimating relatedness among nestmates, as well as other lines of

evidence, including aggressive behavior between colonies. We discovered a novel "Y-

like" social chromosome in *C. niger* harboring a supergene that is strictly maternally

inherited. Surprisingly, we find that the *Cataglyphis* social chromosome is homologous

to the *Solenopsis* social chromosome, even though the two lineages diverged more

than 90 MYA. A wider survey across multiple hymenopteran genomes revealed

conserved synteny of the social chromosome across bees and ants. Our study revealed

a conserved gene set on an ancient chromosome that was repeatedly used in the

evolution of supergenes coding for alternative social structures in diverged ant lineages.

We hypothesize that this gene set represents a genetic "toolkit" that may have been

involved in the evolution of various sociobiological traits in Hymenoptera. This gene set

evolved in a common ancestor of ants and bees, where it may have originally coded for

a different trait. This ancestral trait may have been a pre-adaptation for sociality, such

as regulation of maternal care.

2. Results and discussion

2.1 Social polymorphism in *Cataglyphis niger* is associated with chromosome 2

We fully excavated 30 nests from a single population in Tel Baruch, Israel, where both monogyne and polygyne nests were known to co-exist. Genomic data was generated for 672 samples (28 queens and 644 workers) from the 30 nests (20-24 individuals each) using double-digest restriction-site associated DNA sequencing (RAD-seq), yielding genotype data for 30,955 high-confidence single nucleotide polymorphisms (SNPs). As queens were found only in 14 nests (7 had one queen and 7 had more than one queen), the social form of the sampled nests was assigned based on the relatedness within the nest.

A pairwise kinship matrix constructed based on the total length of shared identity-by-descent (IBD) haplotypes across 672 sampled individuals showed some nests with relatedness between all nestmates and some where most pairs were unrelated (Figure 1a), as expected for monogyne and polygyne nests, respectively. In monogyne nests, we expected three clusters of relatedness scores— full siblings or super sisters sharing 75% of their genome, mother-daughter pairs with 50% relatedness, and half-sisters (from different fathers) with 25% relatedness. The nests with higher relatedness had broadly three clusters of relatedness scores consistent with our expectations in a monogyne nest (Figure 1b). However, we did find several worker pairs with 50% relatedness, which could result from the queen mating with brothers. As expected for polygyne nests, most pairs of workers had relatedness scores lower than half-sister relatedness (25%). Based on the distribution of kinship scores, 13 nests were classified as monogyne and 17 as polygyne for downstream analyses. Aggression assays also corroborated this classification of the nests: aggression scores were significantly higher in interactions between two monogyne nests and between a monogyne and a polygyne nest, compared to between two polygyne nests (Figure 1c; *t*-test *p*-value << 0.05 for all comparisons). The number of queens found during sampling was also in line with these results (full details provided in Supplementary Figures S1 & S2).

Figure 1: Identification of social form using kinship inference based on the total length of identity-by-descent (IBD) haplotypes shared between nestmates, **(a)** plotted as a kinship matrix demarketing monogyne nests in blue and polygyne nests in green. **(b)** Distributions of relatedness within nests. **(c)** Aggression scores from the dyadic assay showed significantly higher aggression between monogyne and monogyne-polygyne nests compared to between polygyne nests.

We examined F_{ST} at each SNP between monogyne and polygyne samples (Figure 2a). The average weighted F_{ST} across the genome was low (0.019) as expected due to gene flow and lack of genetic differentiation between the two social forms[25]. However, the average weighted F_{ST} across chromosome 2 was much higher (0.12) than the rest of

the genome. The high F_{ST} (>0.2) region was a ~5.8Mb stretch extending from position
3,930,062 bp to 9,747,310 bp in the Cnig_gn3.1 genome assembly.

Figure 2: Genetic divergence and high linkage disequilibrium on chromosome 2 suggest the presence of a supergene. **(a)** Plot of F_{ST} between social forms (defined based on kinship) showing high F_{ST} values on chromosome 2 . **(b)** Heatmaps of linkage disequilibrium in monogyne (above the diagonal) and polygyne (below the diagonal) population samples (displaying only r^2>0.2) across all chromosomes. **(c)** Heterozygosity along chromosome 2 in 69 full sisters from a polygyne colony. **(d)** Comparison of the boundaries of the supergene region according to the different criteria.

2.2 Suppression of recombination in chromosome 2

Supergenes maintain complex polymorphic traits despite high gene flow between morphs by suppression of recombination that prevents the shuffling of the alleles associated with the trait at different loci. We tested for suppression of recombination using two approaches— (1) testing for linkage disequilibrium (LD) in the population sample and (2) testing for linkage in progeny of a single polygyne queen. We conducted LD analyses separately on monogyne and polygyne samples to examine if the supergene region on chromosome 2 showed suppression of recombination. The analysis revealed high LD ($r^2 \geq 0.2$) in polygynes but not in monogynes across most SNP pairs in the supergene region (chromosome 2 positions 3,813,975 – 10,147,746

bp), aligning with the region of high F_{ST} SNPs (Figure 2b). We also noted LD decaying rapidly with SNP distance across all chromosomes of monogyne and polygyne samples except in chromosome 2 in polygynes, where LD did not decay over long distances (Supplementary Figure S3). Adjacent to the large 5.7Mb LD block, a smaller 270 kb LD block can be observed in both monogyne and polygyne samples (Supplementary Figure S4). Similarly, four additional small LD blocks spanning over 249 kb were observed in other chromosomes of both the social forms (chr07: 298 kb; chr11: 249 kb; chr13: 434 kb; chr17: 297 kb) (Figure 2b).

Additionally, a linkage analysis was conducted based on RAD sequencing of 225 brood samples collected from a single polygyne queen. The mother was also sequenced. Six patrilines were identified in this data set and the largest patriline (69 full sisters) was used to build a linkage map. However, the genotype data did not allow for building a map of chromosome 2 because of an unusual pattern. All sisters had the same genotypes across the supergene region: 131 SNPs (70%) were always heterozygous, and 44 SNPs (24%) always had the same homozygous genotype, out of 186 SNPs that were heterozygous in the mother (Supplementary Figure S5). Plotting heterozygosity along chromosome 2 (Figure 2c) showed that SNPs in the supergene region were either 100% or 0% heterozygous. The extent of markers with heterozygosity of 100% spanned chromosome position 3,986,390 - 10,237,665bp. This pattern can be explained by a combination of two effects: (1) there were no recombination events within the supergene region, and (2) all the daughters received the same haplotype of the supergene from their mother. That is, recombination in the supergene was fully suppressed and one of the two maternal haplotypes is lethal. We note that this pattern is reminiscent of maternal effect killing of homozygous MM daughters in polygyne *F. selysi* colonies.

2.3 Maternally inherited 'Y-like' supergene

We examined the genotypes of 164 high F_{ST} (>0.2) SNPs on chromosome 2, by generating a heatmap (Figure 3a). The heatmap showed that polygyne samples were predominantly heterozygous (127/164 loci on average), whereas monogyne samples

were largely homozygous for one allele (132/164 loci on average). This suggests that

the polygyne social form is associated with a 'Y-like' supergene haplotype, which we

denote as the P haplotype. That is, polygyne samples carry a heterozygous MP

supergene genotype, while monogyne samples are MM homozygotes. This pattern

holds for all queens and most of the workers. One conspicuous exception to this rule

was nest M31, which was designated as a monogyne nest based on high relatedness,

and yet, all samples in this nest had the MP genotype. The phenotype of high

relatedness in this nest could have been observed if polygyne nests in *Cataglyphis* were

also founded independently by queens and not, as commonly assumed, through nest

budding. We speculate that some polygyne nests may be founded by a single MP

queen, which is subsequently joined by additional MP queens. Indeed, this nest had a

relatively small number of workers, as expected for a newly established nest. This is

also in line with the geographic location of this nest among other polygyne nests (Figure

3c). The mitotype (discussed below) and aggression scores of this nest are also as

expected for the polygyne form. The aggression scores of this nest against polygyne

nests (Supplementary Figure S1) were relatively low (0.2 – 0.9) compared to its score

against a monogyne nest (1.3). There were seven other exceptions to the broad pattern

observed in the heat map, all of which were workers: five samples from polygyne nests

had an MM genotype (four from P34 and one from P35) and two samples from

monogyne nests had an MP genotype (M19-44 and M27-19). Nest details along with

supergene genotypes and mitotypes are given in Supplementary Table S1.

Figure 3: (a) Heatmap showing the genotypes at SNPs with high F_{ST} (>0.2) on chromosome 2. Generally, one allele is only found in heterozygous genotypes in polygyne colonies, whereas the other allele is homozygous in monogyne colonies. Exceptions to the rule are highlighted in nests M19, M27, M31, P34. **(b)** Mitochondrial phylogeny with a deep divergence between monogyne and polygyne samples indicating maternal inheritance of the social chromosome. **(c)** Geographic location of the nests where the social form was identified based on kinship and supergene genotype.

Previous work on this species suggested deep divergence between mitotypes associated with monogyne and polygyne social forms based on samples collected across Israel[24]. To ascertain if this pattern was consistent across the nests sampled in this study, we sequenced part of the mitochondrial gene Cytochrome *b* from two to three individuals from each nest. The mitochondrial phylogeny (Figure 3b) resulted in two deeply divergent well-supported monophyletic clades largely corresponding to monogyne and polygyne samples. Previously published mitotypes associated with monogyne nests (previously named the "savignyi" and "drusus" mitotypes) were nested within the monogyne clade and mitotypes associated with polygyne nests (the "niger" mitotype) were nested within the polygyne clade. This divergence in mitotypes

associated with social form indicates a strict, non-Mendelian, maternal inheritance of the

P chromosome. Five samples were an exception to this rule. Both samples sequenced

from nest M31 (a queen and a worker), which had the MP supergene genotype, also

carried a polygyne-associated mitotype. Two worker samples from polygyne nests with

MP supergene genotypes carried a monogyne-associated mitotype.

One conspicuous result from the heat map is the absence of both homozygous PP and

MM genotypes from polygyne nests, where virtually all individuals have a heterozygous

MP genotype. The absence of MM genotypes suggests negative selection, possibly a

result of mechanisms such as meiotic drive, maternal effect killer, or green beard effect,

which were suggested for the social supergenes in other species. Maternal effect killing

was proposed to eliminate MM genotypes in polygyne colonies of *F. selysi*[21], whereby

the presence of the P chromosome in the mother results in the death of progeny that did

not inherit at least one P chromosome. Another possibility is a green beard effect,

whereby workers carrying the P chromosome would kill brood that do not, similar to the

killing of MM queens in polygyne colonies of *S. invicta*[26]. The absence of PP genotypes

also indicates negative selection, potentially due to deleterious mutations, which may be

similar to the low fitness of PP queens in *S. invicta*. Such deleterious mutations may

also result in P being lethal in males or in P males being sterile (although this is not the

case in *S. invicta*). If so, MP daughters would always inherit their P chromosome from

their mother (and M from their father), in line with the mitochondrial phylogeny. Another

possible explanation for the maternal inheritance of the P chromosome is that P males

do exist, but monogyne queens do not mate with them.

2.4 The supergene is distributed across multiple populations in Israel

We used a previously published RAD-seq dataset of worker samples collected across

Israel to verify that the supergene is distributed beyond our study population of Tel

Baruch[25]. This study used different enzymes from our study to generate genomic

libraries resulting in a different set of genomic loci. We used the mitotype of the samples

as a proxy to identify their social form and calculated F_{ST} between inferred monogyne

and polygyne samples at each SNP across the genome. High F_{ST} SNPs were detected

on chromosome 2 between positions 3,985,614-9,804,010, which overlapped the

supergene region identified in the Tel Baruch population (Supplementary Figures S6a

and S6b). Similar to the pattern reported above (Figure 3a), the 164 high F_{ST} SNPs

(>0.4) in this dataset were mostly homozygous MM (52/56) in the monogyne mitotype

samples and heterozygous MP (13/14) in the polygyne mitotype samples

(Supplementary Figure S6c). This sampling across Israel shows that most of the range

consists of pure MM populations, whereas the P haplotypes are always found in

polymorphic populations (MM and MP) limited to the central coastline region between

Ashkelon and Netanya (Supplementary Figure S6d).

2.5 Repeated evolution of social supergenes on an ancient chromosome

Non-homologous social chromosomes are known to be associated with social

polymorphism in *Solenopsis* and *Formica*. We wanted to examine whether the

Cataglyphis social chromosome is homologous to either of these chromosomes.

Cataglyphis and *Formica* both belong to the subfamily Formicinae, whereas *Solenopsis*

belongs to the subfamily Myrmicinae. The two subfamilies diverged around 93 MYA[23]

(95% credible interval 91–95 MYA). Synteny dot plots based on one-to-one orthologs

showed that the social chromosomes of *Cataglyphis* and *Formica* were not homologous

(Figure 4a)*. Surprisingly, despite belonging to different subfamilies, the social

chromosomes of *Cataglyphis* and *Solenopsis* are homologous (Figure 4b). Synteny is

not only conserved in terms of gene content (macrosynteny) but also gene order

(microsynteny), especially in the supergene region (red points in Figure 4c).

Interestingly, the largest region with conserved gene order is inside both supergenes,

while the non-overlapping regions of the supergenes show conservation of gene order

to a lesser extent.

Figure 4: Conserved synteny of the social chromosome across ants and bees. Dot plots of one-to-one

orthologs **(a)** between *Cataglyphis* and *Solenopsis* and **(b)** between *Cataglyphis* and *Formica*. Orange

arrows indicate the social chromosome in *Solenopsis* and *Formica*, and the orange box indicates the

social chromosome in *Cataglyphis*. **(c)** Social chromosomes with the supergene regions of *Cataglyphis* in

blue and *Solenopsis* in green. Supergene coordinates in *Cataglyphis* are defined based on the high F_{ST}

region (3,930,062 to 9,747,310bp), and in *Solenopsis* based on the inversion breakpoints (Yan *et al.*[20]).

The overlapping supergene region highlighted in red shows remarkable conservation in gene order. The

yellow triangles indicate odorant receptors (ORs) and red stars indicate odorant-binding proteins (OBPs).

(d) *Cataglyphis* and *Apis* dot plot. *Cataglyphis* chromosome 2 shows notable conservation of gene

content and homology to *Apis* chromosome 15.

(a)

CHRM	Length (bp)	Cataglyphis niger	Cataglyphis hispanica	Formica aquilonia Formica polyctena	Camponotus floridanus	Lasius platythorax	Pogonomyrmex californicus	Monomorium pharaonis	Solenopsis invicta	Trachymyrmex septentrionalis	Ooceraea biroi	Nysson spinosus	Pemphredon lugubris	Ectemnius continuus	Colletes gigas	Osmia bicornis	Apis mellifera	Anoplius nigerrimus	Vespula vulgaris	Ancistrocerus nigricornis	Nasonia vitripennis	Microplitis mediator	Venturia canescens
chr01	15,508,086	1.00	1.00	0.74	0.33	0.93	0.73	0.85	0.93	0.43	0.24	0.40	0.40	0.28	0.29	0.28	0.27	0.14	0.20	0.21	0.46	0.27	0.28
chr02	12,681,539	1.00	0.65	0.65	0.62	0.65	0.62	0.76	0.75	0.44	0.31	0.24	0.65	0.46	0.41	0.43	0.41	0.35	0.13	0.28	0.37	0.22	0.30
chr03	12,470,023	1.00	1.00	0.70	0.40	0.47	0.34	0.31	0.31	0.35	0.19	0.28	0.35	0.23	0.20	0.20	0.23	0.17	0.17	0.22	0.25	0.22	0.25
chr04	10,525,047	1.00	1.00	0.66	0.40	0.41	0.40	0.47	0.42	0.25	0.20	0.30	0.50	0.27	0.21	0.25	0.39	0.15	0.14	0.25	0.31	0.27	0.23
chr05	12,244,759	1.00	1.00	0.70	0.32	0.52	0.53	0.65	0.51	0.26	0.22	0.13	0.32	0.14	0.12	0.12	0.15	0.10	0.14	0.28	0.29	0.16	0.15
chr06	10,139,329	1.00	1.00	0.72	0.28	0.53	0.42	0.54	0.53	0.26	0.25	0.23	0.40	0.29	0.21	0.21	0.21	0.14	0.13	0.26	0.33	0.30	0.21
chr07	8,573,820	1.00	1.00	0.96	0.99	0.46	0.39	0.71	0.72	0.23	0.40	0.25	0.31	0.27	0.30	0.27	0.26	0.28	0.19	0.23	0.24	0.20	0.20
chr08	7,919,391	1.00	1.00	1.00	0.39	0.35	0.29	0.35	0.29	0.38	0.48	0.15	0.36	0.18	0.12	0.21	0.19	0.18	0.11	0.21	0.27	0.19	0.17
chr09	13,178,881	1.00	0.53	0.51	0.28	0.29	0.29	0.41	0.29	0.20	0.34	0.14	0.47	0.23	0.11	0.21	0.20	0.24	0.14	0.25	0.23	0.16	0.26
chr10	7,568,083	1.00	1.00	0.99	0.57	0.99	0.67	0.67	0.66	0.53	0.39	0.18	0.42	0.31	0.17	0.19	0.19	0.16	0.14	0.34	0.24	0.19	0.26
chr11	6,647,852	1.00	0.99	0.90	0.99	0.98	0.97	0.99	0.94	0.79	0.48	0.26	0.66	0.47	0.35	0.48	0.30	0.42	0.28	0.33	0.71	0.57	0.70
chr12	5,320,879	1.00	1.00	0.52	0.63	0.36	0.36	0.51	0.37	0.37	0.36	0.29	0.36	0.26	0.16	0.25	0.30	0.26	0.18	0.22	0.45	0.24	0.26
chr13	7,063,411	1.00	0.99	0.83	0.43	0.75	0.97	0.54	0.77	0.74	0.60	0.43	0.40	0.29	0.09	0.32	0.31	0.21	0.12	0.28	0.35	0.31	0.42
chr14	6,358,800	1.00	1.00	0.99	0.67	0.63	0.58	0.77	0.59	0.33	0.21	0.36	0.38	0.26	0.23	0.28	0.29	0.19	0.19	0.33	0.34	0.17	0.22
chr15	6,488,307	1.00	0.99	0.98	0.98	0.99	0.96	0.98	0.98	0.54	0.98	0.30	0.48	0.48	0.27	0.39	0.40	0.43	0.13	0.24	0.37	0.23	0.23
chr16	4,535,747	1.00	1.00	0.99	0.99	0.81	0.78	0.65	0.78	0.44	0.61	0.17	0.63	0.57	0.23	0.31	0.40	0.30	0.22	0.38	0.34	0.45	0.40
chr17	4,997,640	1.00	0.98	0.87	0.40	0.76	0.74	0.74	0.75	0.33	0.22	0.43	0.60	0.55	0.39	0.55	0.56	0.24	0.16	0.20	0.35	0.22	0.26
chr18	6,198,604	1.00	1.00	0.99	0.86	0.93	0.67	0.67	0.69	0.73	0.22	0.25	0.41	0.26	0.17	0.20	0.22	0.19	0.13	0.25	0.30	0.20	0.18
chr19	4,963,000	1.00	1.00	1.00	0.45	0.49	0.30	0.37	0.37	0.32	0.40	0.15	0.42	0.27	0.13	0.21	0.17	0.24	0.20	0.28	0.21	0.17	0.28
chr21	5,028,316	1.00	1.00	0.97	1.00	1.00	0.34	0.98	0.99	0.96	0.90	0.23	0.39	0.35	0.18	0.39	0.37	0.26	0.15	0.28	0.34	0.29	0.33
chr22	4,533,825	1.00	1.00	0.98	0.55	0.99	0.54	0.46	0.52	0.98	0.34	0.27	0.37	0.40	0.36	0.29	0.30	0.29	0.17	0.23	0.29	0.22	0.25
chr23	2,743,847	1.00	1.00	0.94	1.00	0.66	0.68	0.96	0.66	0.59	0.32	0.38	0.44	0.28	0.25	0.26	0.25	0.40	0.36	0.35	0.25	0.34	0.36
chr24	3,958,350	1.00	1.00	0.92	0.69	0.64	0.76	0.67	0.65	0.35	0.41	0.30	0.44	0.43	0.28	0.25	0.34	0.27	0.18	0.19	0.37	0.26	0.26
chr25	2,989,821	1.00	1.00	0.84	0.58	0.54	0.56	0.47	0.50	0.72	0.27	0.20	0.38	0.29	0.22	0.23	0.28	0.39	0.13	0.26	0.35	0.27	0.22
chr26	2,288,207	1.00	1.00	0.93	0.90	0.92	0.83	0.89	0.88	0.63	0.51	0.78	0.51	0.91	0.31	0.45	0.56	0.46	0.32	0.33	0.41	0.39	0.58

(b)

Figure 5: Conservation of synteny of all *Cataglyphis* chromosomes. **(a)** Percent of one-to-one

orthologous genes in each *Cataglyphis* chromosome that reside in one chromosome in other

Hymenoptera species. **(b)** Synteny plot showing three representative species of ants, two Apoidea

species, and one wasp outgroup. The figure highlights three broad regions of *Cataglyphis* chromosome 2,

the supergene region (red), the region upstream (green), and the region downstream (blue) of the

supergene. The blue box marks a cluster of genes that appears to have been convergently fused to the

social chromosome in both *Cataglyphis* and *Solenopsis* lineages (see Supplementary Figure S7).

To investigate the extent of the conservation of synteny of the social chromosome, we

conducted a comparative genomic analysis of all published annotated, chromosome-

level ant assemblies. We also added representatives of bees and wasps to this analysis

amounting to 22 chromosome-level assemblies that allowed us to evaluate the

conservation of synteny across Hymenoptera. As an estimate of macrosynteny, we

calculated the percentage of *Cataglyphis* genes found on the homologous chromosome

(the chromosome carrying the largest number of homologous genes) in each of these

species (one-to-one orthologs only). Figure 5a shows that *Cataglyphis* chromosome 2 is

one of the more highly conserved chromosomes across Hymenoptera, especially

relative to other large chromosomes (chromosomes are numbered from largest to

smallest). This chromosome is notably more conserved than most other chromosomes

in distant lineages such as *Solenopsis* (75% of one-to-one orthologs on the same

chromosome) and even in the honey bee *Apis mellifera* (41%).

Figure 5b shows the conservation of microsynteny of genes on the social chromosome

in three ants, two Apoidea species (the honey bee and a digger wasp), and the jewel

wasp *Nasonia* as an outgroup (Supplementary Figure S7 shows all 22 Hymenoptera).

Microsynteny in the supergene and the region upstream to the supergene in

Cataglyphis appear to be conserved to a certain extent across the available genomes

from Myrmicinae and Formicinae. In the clonal raider ant *Ooceraea biroi,* however, this

region is broken up and represented on five chromosomes. The conservation of the

chromosome in bees suggests that it was broken up in the lineage leading to *O. biroi*

after it diverged from the lineage leading to Myrmicinae and Formicinae. It remains to be

determined whether this is common to other members of the subfamily Dorylinae or

specific to the clonal raider ant. The downstream region of the *Cataglyphis* chromosome

is broken in many formicine species (e.g. *Formica*) and myrmicine species (e.g.

Pogonomyrmex), and also in bees. Either the downstream region was part of the

ancestral chromosome and broke off in multiple myrmicine and formicine lineages, or it

was not part of the ancestral chromosome, and the same synteny group of genes was

fused to the social chromosome in both *Cataglyphis* and *Solenopsis* lineages (marked

in Figure 5b; see also Supplementary Figure S7).

Most genes in the upstream region of the *Cataglyphis* social chromosome and the supergene region (red and green lines in Figure 5b) are found on chromosome 15 of *Apis*, followed by chromosomes 1, 7, 8, and 14. The downstream *Cataglyphis* genes (blue lines) are not on *Apis* chromosome 15. This conservation of macrosynteny is evident across all the six representatives of the superfamily Apoidea, which includes the bees and the crabronid wasps (Supplementary Figure S7). These results indicate deep conservation of the social chromosome, dating back to the divergence of Apoidea and Formicidae (ants) around 175 million years ago[23]. This exceptional conservation of the social chromosome suggests that its genomic architecture has adaptive significance. We propose that the conserved gene content in the supergene region is an ancestral genetic "toolkit" that has been repeatedly used in the evolution of sociobiological traits in hymenopteran lineages.

The association of social polymorphism with a supergene on the social chromosome in both *Cataglyphis* and *Solenopsis* can be interpreted in two ways. One possibility is that the supergene was present in the common ancestor and conserved in both lineages. However, the sequence divergence between the two *Solenopsis* supergene haplotypes —corresponding to only one million years ago— suggests a more recent origin[15,19]. Although the age of the *Cataglyphis niger* supergene remains unknown, no polygyne colonies have been reported from closely related *Cataglyphis* species (the bicolor group). This suggests that the two lineages independently evolved analogous supergenes on the same chromosome. While this may seem surprising, studies on sex chromosome turnover show that certain chromosomes repeatedly transition into sex chromosomes, likely due to the genes they carry[27,28]. Similarly, if the social chromosome harbors an ancient genetic toolkit for sociobiological traits, this would explain the repeated evolution of supergenes on this chromosome. Examining the genomic basis of multiple socially polymorphic species in other ant lineages could help shed light on this question.

2.5 Conserved gene content of the social chromosome across ants and bees

We examined the gene content of the *Cataglyphis* and *Solenopsis* social chromosomes to find 786 and 1088 putative protein-coding genes, respectively. The two supergene regions contain 248 and 441 putative protein-coding genes in *Cataglyphis* and *Solenopsis*, respectively. Next, we identified orthologous genes between the two chromosomes. In the *Solenopsis* supergene region, there are 207 functionally-annotated protein-coding genes with one-to-one orthologs in *Cataglyphis*, and 112 such genes in the *Cataglyphis* supergene, with 84 of these being found in both the supergene regions. All but one of the one-to-one orthologs of genes in the *Cataglyphis* supergene are in the *Solenopsis* social chromosome, and one gene is in an unassembled scaffold (Supplementary Table S2), demonstrating the exceptional degree of gene content conservation of this chromosomal region. Other genes have many-to-many orthology relationships between the two species (Supplementary Table S3; genes with functional annotation). The *Cataglyphis* social chromosome contains 492 such genes in total, of which 149 are in the supergene region. The *Solenopsis* social chromosome contains 665 such genes, 279 of which are in the supergene region.

The homologous supergene regions contain a large and diverse set of molecular functions. It is difficult to identify specific candidate genes without additional functional information. Nevertheless, we reviewed the list and looked for gene functions that may be implicated in the sociobiology of ants (Supplementary Tables S2 and S3 – one-to-one and many-to-many; Supplementary Figure S8 – dot plot with gene annotations). One notable example is a cluster of three cytochrome P450 4C1 genes in the *Cataglyphis* supergene, which are orthologous to nine genes in the *Solenopsis* supergene. Cytochrome P450 genes are a large and diverse family of enzymes, which was noted already when the first ant genomes were sequenced. The genome of the harvester ant *Pogonomyrmex barbatus* was found to contain 72 cytochrome P450 genes, which is a significant expansion relative to other insect species[29]. Enzymes of this gene family participate in a wide range of metabolic processes[30], including synthesis and degradation of hormones such as juvenile hormone, a major regulator of insect development that plays a key role in caste determination in social insects[31–34].

We also took note of the downstream region of the *Cataglyphis* chromosome, which is

homologous to the upstream region (short arm) of the *Solenopsis* chromosome,

because this region may have been convergently fused to the chromosome in both

lineages (Figure 5b; Supplementary Figure S7). This region contains 121 functionally

annotated one-to-one orthologous genes, including a steroid hormone receptor

(*Cataglypis* gene_1712 / *Solenopsis* XP_039315095.1), an ecdysone 20-

monooxygenase (*Cataglyphis* gene_2123 / *Solenopsis* XP_011172207.1), and an

ecdysteroid-regulated 16 kDa protein (*Cataglyphis* gene_2111 / *Solenopsis*

XP_011170763.1). Genes implicated in such functions may have significance for the

sociobiology of ants, as different levels of ecdysteroids and ecdysone receptors were

found in the brains of foragers versus nurses in *Camponotus* and *Harpegnathos*[35,36].

The total number of functionally annotated one-to-one orthologs shared between the

Cataglyphis and *Solenopsis* social chromosomes and *Apis mellifera* (anywhere in the

bee genome) is 305. Of these, 164 (54%) are located on *Apis* chromosome 15

(Supplementary Table S4), and 34 out of the 164 are located in the supergenes of both

Solenopsis and *Cataglyphis*. These 34 genes include dopamine receptor 2 (*Cataglypis*

gene_1588 / *Solenopsis* XP_039314227.1 / *Apis* XP_026301043.1), odorant binding

protein-9 (*Cataglyphis* gene_1588 / *Solenopsis* XP_039314227.1 / *Apis*

XP_026301043.1), and neuronal acetylcholine receptor subunit alpha-10 (*Cataglyphis*

gene_2000 / *Solenopsis* XP_025994007.1 / *Apis* XP_392070.3). Dopamine is

implicated in modulating social and reproductive behavior in Hymenoptera[37–41].

2.6 Enrichment in olfactory and neural signaling functions in the social

chromosome across ants and bees

We tested for enrichment of 13,317 Gene Ontology (GO) functional categories in the

conserved gene set on the social chromosome. We analyzed representative genomes

from eight distantly related Apoidea lineages (including ants and bees) that showed

conserved synteny of the social chromosome. We defined a conserved set of 187

orthologous gene groups (OGs) where the ortholog is found in the chromosome

homologous to the social chromosome in at least six out of the eight genomes. Table 1

lists the GO categories that were significantly enriched (FDR corrected *p*-value < 0.05).

The full table with additional information can be found in Supplementary Table S6. Most

notably, the top GO categories include olfactory learning and response to odorants (GO:

0090328, GO: 1990834). Other enriched GO categories are related to neural functions,

especially dopamine signaling (GO:0007191, GO:0004952, GO:0001588). These

results indicate that the conserved gene set in the social chromosome is involved in

chemical communication and neural signaling. Other GO categories that were

significantly enriched were involved in metabolic regulation and immune function.

Table 1: Functional categories enriched in the conserved gene set of the social chromosome

GO category	OGs with GO*	FDR corrected *p*-value	*Cataglyphis* gene in each conserved OG
BP: Regulation of olfactory learning (GO:0090328)	3/5	0.019	Dopamine receptor 1 (gene_1435), Octopamine receptor in mushroom bodies *Oamb* (gene_1830), Uncharacterized protein (gene_1662)
BP: Response to odorant (GO:1990834)	3/7	0.019	Dopamine receptor 1 (gene_1435), Dopamine receptor 2 (gene_1588), Uncharacterized protein (gene_1662)
BP: Adenylate cyclase-activating dopamine receptor signaling pathway (GO:0007191)	2/2	0.019	Dopamine receptor 1 (gene_1435), Dopamine receptor 2 (gene_1588)
MF: Dopamine neurotransmitter receptor activity coupled via Gs (GO:0001588)	2/2	0.019	Dopamine receptor 1 (gene_1435), Dopamine receptor 2 (gene_1588)
MF: Dopamine neurotransmitter receptor activity (GO:0004952)	2/2	0.019	Dopamine receptor 1 (gene_1435), Dopamine receptor 2 (gene_1588)
BP: Positive regulation of sensory perception of pain (GO:1904058)	2/2	0.019	Tachykinin-like peptides receptor 99D (gene_1948), ATP synthase subunit C lysine N-methyltransferase (gene_1859)
BP: Propionate metabolic process methylcitrate cycle (GO:0019679)	2/2	0.019	4-hydroxybutyrate coenzyme A transferase (gene_1884, gene_1885)
BP: Acetate metabolic process (GO:0006083)	2/2	0.019	4-hydroxybutyrate coenzyme A transferase (gene_1884, gene_1885)
MF: Acetate CoA-transferase activity (GO:0008775)	2/2	0.019	4-hydroxybutyrate coenzyme A transferase (gene_1884, gene_1885)
BP: Positive regulation of activated T cell proliferation (GO:0042104)	2/2	0.019	Signal transducer and activator of transcription 5B (gene_1887), Glycerol-3-phosphate acyltransferase 1 mitochondrial (gene_2005)

	*		
BP: Positive regulation of programmed cell death (GO:0043068)	2/2	0.019	Cell death specification protein 2 (gene_1943), Uncharacterized protein (gene_1662)
CC: Integral component of membrane (GO:0016021)	33/1241	0.019	33 genes including Dopamine receptor 1 (gene_1435), Dopamine receptor 2 (gene_1588), Octopamine receptor in mushroom bodies *Oamb* (gene_1830), Tachykinin-like peptides receptor 99D (gene_1948), ATP synthase subunit C lysine N-methyltransferase (gene_1859), Serine protease HTRA2 mitochondrial (gene_1443), Attractin-like protein 1 (gene_1831), Pickpocket protein 28 (gene_1847), Beta-14-galactosyltransferase 7 (gene_1850), ABC transporter G family member 20 (gene_1862), MAP kinase-activating death domain protein (gene_1873), probable sodium/potassium/calcium exchanger CG1090 (gene_1906), Calcium-independent phospholipase A2-gamma (gene_1530), Sphingosine-1-phosphate lyase (gene_1949), Tetraspanin-33 (gene_2003), Sodium-coupled neutral amino acid transporter 9 (gene_1480)
MF: Manganese ion binding (GO:0030145)	4/24	0.027	Beta-14-galactosyltransferase 7 (gene_1850), m7GpppN-mRNA hydrolase (gene_1532), NADP-dependent malic enzyme (gene_1951), DNA-directed primase/polymerase protein (gene_1937)'
MF: Hydrolase activity (GO:0016787)	5/49	0.041	V-type proton ATPase subunit B (gene_1502), DNA repair and recombination protein RAD54-like (gene_1503), Fumarylacetoacetate hydrolase domain-containing protein 2A (gene_2029), Phosphoglycerate mutase 2 (gene_1893), Uncharacterized protein (gene_2004)
MF: Acetyl-CoA hydrolase activity (GO:0003986)	2/3	0.041	4-hydroxybutyrate coenzyme A transferase (gene_1884, gene_1885)
BP: Regulation of presynaptic cytosolic calcium ion concentration (GO:0099509)	2/3	0.041	Dopamine receptor 1 (gene_1435), Dopamine receptor 2 (gene_1588)

Results of the hypergeometric test for enrichment of Gene Ontology (GO) categories in the conserved set

of Ortholog Groups (OGs) that are on the social chromosome in at least six out of the eight species

included in the analysis.

BP = biological process; MF = molecular function; CC = cellular component.

* The number of OGs in the conserved gene set out of the total number of OGs in the dataset that were

annotated with a specific GO term.

We were particularly interested in genes implicated in olfactory functions, which were

extensively studied in *Solenopsis* because of their potential role in chemical signaling

and the regulation of social structure. Previous studies reported differences between

monogyne and polygyne queens in cuticular hydrocarbons and other compounds that

were suggested to function as queen pheromones involved in the regulation of social

form and the green beard effect[26,42,43]. Other studies focused on odorant binding

proteins (OBPs) and odorant receptors (ORs) in the *Solenopsis* supergene, which may

be involved in olfactory response to such chemical signals[44–47]. The *Solenopsis*

supergene contains a large array of 23 ORs and 3 more separate ORs[47]. We annotated

ORs in the *Cataglyphis niger* genome and constructed a gene tree including ORs from

C. niger, Solenopsis invicta, Camponotus floridanus, and *Linepithema humile* (Supplementary Figure S8). The 23 *Solenopsis* ORs and their orthologs in *Cataglyphis* belong to a single subfamily of the OR gene superfamily. The gene tree shows that the 23 *Solenopsis* ORs have one-to-one or many-to-many orthology relationships with ORs in two regions of the *Cataglyphis* social chromosome, but outside of the supergene (marked by red stars on the dot plot in Figure 4c). Three more ORs in the *Solenopsis* supergene (belonging to a different OR subfamily) are orthologs of two *Cataglyphis* ORs, which also reside outside of the supergene. Given that the ORs are a fast-evolving family, there is no clear orthology relationship between ant and bee ORs.

The *Solenopsis* supergene also contains 9 out of a total of 23 OBPs in the genome[46]. One of the OBPs is *Gp-9*, the first gene discovered in the *Solenopsis* supergene in the 1990s and the one protein that shows the largest number of amino acid substitutions between the M and the P chromosomes, indicating positive selection[44–46]. Interestingly, the homologous chromosome in the honey bee, chromosome 15, also contains 9 out of 21 OBPs, but only one of these (AmelOBP9) is orthologous to one of the supergene OBPs (SiOBP9). The 9 OBPs in the *Solenopsis* supergene are orthologous to 3 OBPs in the *Cataglyphis* supergene (Fig. 4c). Cnig_gene_1607 is orthologous to 7 of the *Solenopsis* OBPs; Cnig_gene_1608 is the ortholog of the *Solenopsis Gp-9* gene (SiOBP3); and Cnig_gene_1586 is orthologous to SiOBP9 and AmelOBP9 (Fig. 6). We note that the OBP cluster containing *Gp-9* is in the largest block of conserved microsynteny between *Solenopsis* and *Cataglyphis*. Altogether, these gene families implicated in olfaction are promising candidates for the toolkit that may have been reused for social evolution in multiple hymenopteran lineages.

2 **Figure 6:** Gene tree of the odorant binding protein (OBP) gene family in the ants *Cataglyphis niger* and

3 *Solenopsis invicta* and the honey bee *Apis mellifera*. OBPs marked by rectangles reside on the social

4 chromosomes (*Cataglyphis* chromosome 2, *Solenopsis* chromosome 16, *Apis* chromosome 15).

3. Conclusion

The complex trait of multi-queen (polygyne) colonies has evolved repeatedly from single-queen (monogyne) colonies in ants[8,9]. Multiple ant species display polymorphism in social form, providing a unique opportunity to investigate the evolutionary genomic basis of this trait. We investigated the genomic architecture associated with social form in a polymorphic population of *Cataglyphis niger*. Our data show that social polymorphism is associated with a large supergene— a region of suppressed recombination across at least 5.8Mb of chromosome 2. This supergene is virtually always heterozygous (MP) in polygyne individuals and homozygous (MM) in monogynes. Surprisingly, this supergene is homologous to the supergene of the fire ant *Solenopsis invicta*, even though these two lineages diverged more than 90 million years ago. These species are representatives of the two largest ant subfamilies Formicinae and Myrmicinae. Most of *Cataglyphis* chromosome 2 is homologous to most of *Solenopsis* chromosome 16, with a substantial overlap of the supergene regions where the gene order is highly conserved. The specific structure of this set of genes suggests that this chromosome has conserved functions related to social colony structure and the sociobiology of ants, which is likely to be shared across many ant species. Conservation of synteny in gene content of the social chromosome is also observed in representatives of Apoidea, including the honeybee *Apis mellifera*. The observation of conserved synteny dating back to around 175 MYA indicates that this set of genes acted in synergy to produce some traits with fitness consequences. Specifically in ants, this gene set facilitated the repeated evolution of polygyne social structure. The conservation of synteny before the evolution of social polymorphism in *Solenopsis* and *Cataglyphis* suggests that this gene set was responsible for some traits of consequence over 175 million years, possibly also related to sociobiology. The conserved gene set could have served as a genetic "toolkit" for evolving sociobiological traits across bee and ant lineages. This discovery opens the way for both evolutionary and mechanistic studies of the genetic basis and genomic architecture of complex phenotypes related to insect sociality.

4. Methods

4.1 Sample collection

Thirty nests of *Cataglyphis niger* were collected from a single population in Tel Baruch beach, Israel, between November and December 2019. This location is known to harbor both monogyne and polygyne colonies based on Reiner Brodetzki *et al.*[25]. We targeted at least two monogyne and two polygyne nests in every sampling trip based on their geographical distribution in that previous study. This also allowed us to conduct aggression assays (see below) between monogyne and polygyne nests after every batch of sampling. A distance of at least 100m was maintained between two sampled nests as these ants are known to be polydomous. An attempt was made to collect the queens by excavating the entire nest. Nest samples were carried back to the lab (TBA14 - 44) to perform behavioral experiments and extract genomic DNA. An additional polygyne nest (TBA67) with several queens was collected in January 2021 and was used for building a linkage map. The queens from this nest were isolated and 225 samples from the brood of a single queen were collected for RAD sequencing.

4.2 Molecular work

Samples not used for the behavioral assay were stored at -80C immediately after sampling. 22-24 workers from each nest in addition to the queens amounting to 714 samples were used for the molecular work from sampling carried out in 2019, henceforth referred to as the SC (social chromosome) dataset. An additional 225 brood samples from a single polygyne mother were used to construct a linkage map (linkage map dataset). DNA was extracted from the workers' abdomen and the queen's thorax using a liquid handling robot (Tecan Freedom EVO 150 NGS) and DNeasy (QIAGEN) protocol. The extracts were cleaned using ethanol precipitation and stored at -20C. A partial fragment of 450bp of Cytochrome b (*cyt b*) was PCR amplified and Sanger sequenced for 2-3 workers and queens from each nest. Double-digest Restriction-site Associated DNA sequencing (RAD-seq) was used to generate genomic libraries following Brelsford *et al.*[48]. Briefly, the genomic DNA was cut using restriction enzymes EcoRI and BfaI. Adaptors with individual barcodes were ligated to each sample for multiplexing. The ligated products were PCR amplified using barcoded primers and

pooled. Size selection of DNA fragments between 330–1000bp was carried out using a

Bluepippin instrument (Sage Science). A detailed protocol can be found in Lajmi *et al.*[49].

Libraries were sequenced on an Illumina HiSeq X machine with paired-end 150bp

reads.

4.3 Molecular Analysis

4.3.1 Mitochondrial phylogeny

The Sanger sequences were aligned in MEGA X[50] using the Muscle algorithm[51]. To

assign our samples to previously described mitotype clades, published *cyt b* sequences

of monogyne and polygyne colonies were also included from Reiner Brodetzki *et al.*[25].

Mitotypes of *Cataglyphis israelensis* were used as an outgroup. A maximum likelihood

tree was constructed using raxmlGUI 2.0[52] using a GTR+G model of sequence

evolution and 1000 bootstraps.

4.3.2 RAD-seq pipeline

The bioinformatics pipeline was run separately for the SC data and linkage map, as well

as the two datasets combined. The raw sequencing data were examined using

FastQC[53] for quality. Adaptors and low-quality reads were then trimmed/filtered using

Trimmomatic[54]. Two mismatches were allowed in identifying the adaptors, with eight

being the minimum length for identifying an adaptor, a sliding window size of 4bp with

an average required quality of 15 was used. FLASH[55] was used to combine overlapping

paired reads. The data were then demultiplexed using the command process_radtags.pl

from the Stacks2 package[56]. The reads were then mapped to the *Cataglyphis niger*

genome (assembly version Cnig_gn3.1 [57]; NCBI BioProject accession PRJNA1168153)

using Bowtie2[58] with the --end-to-end and --very-sensitive options. SNP calling was

conducted using Ref_map.pl from Stacks2 to obtain a VCF file. Samples with more than

50% missing data were dropped from the analyses. The SNP calling was done

separately for the SC and linkage map datasets, as well as the combined dataset. A

blacklist of loci with potential artifacts was created based on the heterozygous genotype

calls in three or more samples out of 64 haploid males. These loci were excluded from

the VCF. Such artifacts may be the result, for example, of errors in the assembly of the

reference genome, where repetitive sequences are collapsed into a single copy.

Along with removing the blacklisted loci, the following loci were removed from the VCF

file for the SC dataset using VCFtools version 0.1.14[59]: i) genotype calls based on less

than 12 reads; ii) loci with more than 30% missing data; iii) loci with a minor allele

frequency of less than 0.4% or 3 allele copies; iv) loci with a mean depth greater than

175 reads per individual. The final VCF file for the SC dataset consisted of 672

individuals comprising 28 queens and 644 workers. This resulted in a catalog of 30,955

high-confidence SNPs that were used for most of our analyses. The linkage map data

VCF file varied slightly in the filtration in that we required only 8 reads for a genotype

call, a maximum mean depth of 150 reads per individual (because of lower average

depth), and a minor allele frequency of 1% or 3 allele copies (because of the smaller

number of samples). The final VCF file for the linkage map consisted of 225 offspring

and their mother and amounted to 17,984 high-confidence SNPs.

4.4 Identification of the social form

Cataglyphis niger is a polydomous species, i.e., a colony may have multiple nests. This

made finding the queens of each colony difficult. Therefore, we relied on kinship

estimates among worker samples to identify the colony's social form. Monogyne nests

were expected to show high kinship between nestmates as they are expected to be full

or half-siblings, compared to polygyne nests where workers are not necessarily related.

We also conducted behavioral assays, as additional supporting evidence for the

colony's social structure.

4.4.1 Kinship

Kinship inference was carried out for the two genomic datasets separately. For the SC

dataset, kinship was used to identify the social form of the nests. Whereas in the linkage

map dataset, kinship was used to identify the patrilines. The RAD-seq SNP catalog was

imputed and phased using Beagle version 5.0[60] and refined-IBD[61] was used to identify

shared identity-by-descent (IBD) haplotypes between samples. The kinship between a

pair of samples was estimated based on the combined lengths of their shared IBD

haplotypes, resulting in a kinship matrix. To obtain an estimate of the relatedness

coefficient we divided the total IBD length by twice the total genome assembly length

(because the samples are diploid). A total of 7,651,126 and 6,238,277 IBD haplotypes

were identified in the SC dataset (among the 672 samples) and the linkage map dataset

(among the 225 samples), respectively. Social form was assigned to each nest based

on the distribution of kinship scores. Nests were classified as polygyne if more than

75% of pairs of nestmates had less than 20% relatedness, and as monogyne otherwise.

4.4.2 Behavioral experiment

Behavioral assays were conducted within a week of collection from the field. These

assays were based on the assumption that workers from different colonies would show

high aggression towards each other. Therefore, workers from two different monogyne

nests were expected to show higher aggression than workers from two polygyne nests,

as we expect all polygyne ants to be part of a supercolony[25]. Dyadic aggression assays

were performed, where two ants from different nests (test) or the same nest (control)

were introduced into a round 12 cm diameter arena and their behavior was video

recorded for three minutes. Behavior aggression scores were coded as antenation=0,

mandibular threat=1, short bite with jump=2, bite=3, spraying=4 and the total aggression

score was calculated as

$$\sum_{i=1}^{n} \frac{A_i * t_i}{T}$$

where A_i and t_i are the aggression score and duration of each behavior over the total

time T, respectively. Ten trials were conducted for each pair of nests that was tested

and an average score was calculated. A total of 1029 assays were carried out to get

104 averaged scores between pairs of nests. A two-sample *t*-test was conducted to

compare the aggression scores between nestmates, polygyne-polygyne nests, and

monogyne/polygyne-monogyne nest. The test was performed in R using command

t.test.

4.5 Identifying genomic regions associated with social form

Weir and Cockerham (1984)[62] F_{ST} was calculated for every SNP using VCFtools version

0.1.14[59] between 282 monogyne samples and 388 polygyne samples, where the social

form was defined based on the distribution of kinship score within the nest. Linkage

disequilibrium (LD; r^2) analysis was performed among SNPs in each chromosome for

monogyne and polygyne samples separately, using Plink v1.9[63]. Only SNPs exhibiting a

minor allele frequency surpassing 0.05 for each social form were included in this

analysis.

4.5.1 Linkage analysis in the progeny of a polygyne queen

To test for recombination in the supergene in polygyne colonies, we isolated a single

polygyne queen from nest TBA67 with workers (TBA67c). 269 samples consisting of

larvae and pupae were collected in a span of 60 days after the queen was isolated.

RAD-seq libraries were constructed for the queen and her offspring using the same

protocol as above. Patrilines were identified using kinship analysis (as above) and a

linkage map was constructed for the patriline with the largest number of samples (69

samples). A linkage map was constructed based on 3,149 bi-allelic loci where the minor

allele was present in at least two individuals from the same patriline. However, this

analysis failed in the supergene region of chromosome 2 because all 69 samples had

the same genotype in every SNP. Therefore, it was not possible to estimate the

frequency of recombination and test for linkage between neighboring SNPs. To illustrate

this unusual genetic pattern, we plotted the heterozygosity in the 69 progeny along the

chromosome for all SNPs that were heterozygous in the mother queen.

4.6 Synteny analysis

Synteny analysis was carried out to evaluate the conservation of synteny in the

chromosome carrying the supergene across 22 hymenopteran species with the oldest

divergence dating to ~230 million years ago. The analysis included the ant subfamilies

Formicinae (*Cataglyphis niger*, *C. hispanica*, a *Formica aquilonia x polyctena* hybrid*,

Camponotus floridanus,* and *Lasius platythorax*), Myrmicinae (*Monomorium pharaonis*,

Solenopsis invicta, *Pogonomyrmex californicus,* and *Trachymyrmex septentrionalis*) and

Dorylinae (*Ooceraea biroi*). Six species belonging to the Apoidea superfamily, the sister

clade to ants, were also included in this analysis. This included bee families Apidae

(*Apis mellifera*) and Colletidae (*Colletes gigas*), a solitary bee belonging to Megachilidae

(*Osmia bicornis*), and solitary wasps belonging to Crabronidae (*Pemphredon lugubris*,

Nysson spinosus, and *Ectemnius continuus*). Six additional wasp lineages were

included— families Vespidae (*Vespula vulgaris* and *ancistrocerus nigricornis*),

Chalcidoidea (*Nasonia vitripennis*), Ichneumonidae (*Venturia canescens*),

Ichneumonoidea (*Mocroplitis mediator*), and Pompilidae (*Anoplius nigerrimus)*. The

genomes for this analysis were chosen based on their availability on NCBI, a scaffold

N50 of more than 1MB, and gene annotation for the genome assembly. OMA v2.6.0[64]

was used to infer orthogroups and transfer functional annotations from *Solenopsis*

invicta orthologs to *Cataglyphis niger* genes.

4.7 GO enrichment analysis

GO categories were assigned to *C. niger* genes using Trinotate[65] and used to test for

enrichment on the social chromosome using the hypergeometric test implemented in

the python module scipy[66]. We used Hierarchical Orthologous Groups (HOGs) defined

by OMA for chromosome-level assemblies of eight representatives of diverged Apoidea

lineages: *Ectemnius continuus*, *Pemphredon lugubris*, *Colletes gigas*, *Apis mellifera*,

Pogonomyrmex californicus, *Solenopsis invicta*, *Cataglyphis niger*, *Lasius platythorax*.

From the OMA results, HOGs uniqely presented on at least on six out of the eight

genomes in the chromosome homologous to *C. niger* chromosome 2 were designated

as our sample population for the test. For each GO category present on these HOGs,

we counted how often they are present in these compared to the whole genome.

4.8 OR and OBP phylogenetic analysis

Odorant receptors (ORs) were annotated using HAPpy-ABCENTH[67], based on

manually curated OR gene models from 5 ant species. The first part of the program

locates putative genetic loci through homology, and the second part, ABCENTH, is a

gene finder specifically built for multigene families like ORs, with high sequence

divergence but a very conserved exon structure. OR protein sequences were compiled

from *Cataglyphis niger, Solenopsis invicta*, *Linepithema humile,* and *Camponotus*

floridanus. Odorant Binding Protein (OBP) sequences for *Apis mellifera*[68] and

Solenopsis invicta[46] were downloaded from NCBI and *Cataglyphis* annotations were

used from Inbar *et al.*[57] and analyzed separately. For each of these datasets, MAFFT[69]

was used to build a multiple sequence alignment. OR and OBP gene trees were

reconstructed using RAxML v8.2.12[70], with the PROTCALG model, the LG rate matrix, and 100 bootstraps repeats.

5. Acknowledgments

We thank Abraham Hefetz for introducing us to the polymorphic study population and for background knowledge on this species and its social forms. We are grateful to Jessica Purcell and Alan Brelsford for their advice on RAD sequencing and the study of supergenes. EP would like to thank Laurent Keller for inspiring and motivating the study of social supergenes. AL would like to thank Praveen Karanth for his support during the analysis and writing of this paper. AL was supported by the "Study in Israel" Fellowship provided by the Planning & Budgeting Committee (PBC) of the Council for Higher Education (CHE). This study was funded by Israel Science Foundation grants 646/15 and 1845/22.

6. References

1. Blount, Z. D., Lenski, R. E. & Losos, J. B. Contingency and determinism in evolution: Replaying life's tape. *Science* **362**, eaam5979 (2018).
2. Toth, A. L. & Robinson, G. E. Evo-devo and the evolution of social behavior. *Trends in Genetics* **23**, 334–341 (2007).
3. Osborne, K. A. *et al.* Natural Behavior Polymorphism Due to a cGMP-Dependent Protein Kinase of Drosophila. *Science* **277**, 834–836 (1997).
4. Ben-Shahar, Y., Robichon, A., Sokolowski, M. B. & Robinson, G. E. Influence of Gene Action Across Different Time Scales on Behavior. *Science* **296**, 741–744 (2002).
5. Ingram, K. K., Oefner, P. & Gordon, D. M. Task-specific expression of the *foraging* gene in harvester ants. *Molecular Ecology* **14**, 813–818 (2005).
6. Toth, A. L. *et al.* Wasp Gene Expression Supports an Evolutionary Link Between Maternal Behavior and Eusociality. *Science* **318**, 441–444 (2007).
7. Linksvayer, T. A. & Wade, M. J. The Evolutionary Origin and Elaboration of Sociality in the Aculeate Hymenoptera: Maternal Effects, Sib-Social Effects, and Heterochrony. *The Quarterly Review of Biology* **80**, 317–336 (2005).
8. Hughes, W. O. H., Ratnieks, F. L. W. & Oldroyd, B. P. Multiple paternity or multiple queens: two routes to greater intracolonial genetic diversity in the eusocial Hymenoptera. *J of Evolutionary Biology* **21**, 1090–1095 (2008).

9. Hughes, W. O. H., Oldroyd, B. P., Beekman, M. & Ratnieks, F. L. W. Ancestral Monogamy Shows Kin Selection Is Key to the Evolution of Eusociality. *Science* **320**, 1213–1216 (2008).

10. Ross, K. G. & Keller, L. Ecology and Evolution of Social Organization: Insights from Fire Ants and Other Highly Eusocial Insects. *Annual Review of Ecology and Systematics* **26**, 631–656 (1995).

11. Keller, L. Queen number, mode of colony founding, and queen reproductive success in ants (Hymenoptera Formicidae). *Ethology Ecology & Evolution* **3**, 307–316 (1991).

12. Keller, L. Social life: the paradox of multiple-queen colonies. *Trends in Ecology & Evolution* **10**, 355–360 (1995).

13. Kay, T., Helleu, Q. & Keller, L. Iterative evolution of supergene-based social polymorphism in ants. *Philosophical Transactions of the Royal Society B: Biological Sciences* **377**, 20210196 (2022).

14. Schwander, T., Libbrecht, R. & Keller, L. Supergenes and Complex Phenotypes. *Current Biology* **24**, R288–R294 (2014).

15. Wang, J. *et al.* A Y-like social chromosome causes alternative colony organization in fire ants. *Nature* **493**, 664–668 (2013).

16. Purcell, J., Brelsford, A., Wurm, Y., Perrin, N. & Chapuisat, M. Convergent Genetic Architecture Underlies Social Organization in Ants. *Current Biology* **24**, 2728–2732 (2014).

17. Ross, K. G. Multilocus Evolution in Fire Ants: Effects of Selection, Gene Flow and Recombination. *Genetics* **145**, 961–974 (1997).

18. Cohen, P. & Privman, E. Speciation and hybridization in invasive fire ants. *BMC Evol Biol* **19**, 111 (2019).

19. Cohen, P. & Privman, E. The social supergene dates back to the speciation time of two Solenopsis fire ant species. *Sci Rep* **10**, 11538 (2020).

20. Yan, Z. *et al.* Evolution of a supergene that regulates a trans-species social polymorphism. *Nat Ecol Evol* **4**, 240–249 (2020).

21. Avril, A., Purcell, J., Béniguel, S. & Chapuisat, M. Maternal effect killing by a supergene controlling ant social organization. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 17130–17134 (2020).

22. Brelsford, A. *et al.* An Ancient and Eroded Social Supergene Is Widespread across Formica Ants. *Current Biology* **30**, 304-311.e4 (2020).

23. Moreau, C. S. & Bell, C. D. Testing the museum versus cradle tropical biological diversity hypothesis: Phylogeny, diversification, and ancestral biogeographic range evolution of the ants. *Evolution* **67**, 2240–2257 (2013).

24. Eyer, P. A., Seltzer, R., Reiner-Brodetzki, T. & Hefetz, A. An integrative approach to untangling species delimitation in the Cataglyphis bicolor desert ant complex in Israel. *Molecular Phylogenetics and Evolution* **115**, 128–139 (2017).

25. Reiner Brodetzki, T. *et al.* The Interplay between Incipient Species and Social Polymorphism in the Desert Ant Cataglyphis. *Sci Rep* **9**, 9495 (2019).

26. Keller, L. & Ross, K. G. Selfish genes: a green beard in the red fire ant. *Nature* **394**, 573–575 (1998).

27. Marshall Graves, J. A. & Peichel, C. L. Are homologies in vertebrate sex determination due to shared ancestry or to limited options? *Genome Biol* **11**, 205 (2010).

28. Jeffries, D. L. *et al.* A rapid rate of sex-chromosome turnover and non-random transitions in true frogs. *Nat Commun* **9**, 4088 (2018).

29. Smith, C. R. *et al.* Draft genome of the red harvester ant *Pogonomyrmex barbatus*. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 5667–5672 (2011).

30. Feyereisen, R. INSECT P450 ENZYMES. *Annu. Rev. Entomol.* **44**, 507–533 (1999).

31. Robinson, G. E. & Vargo, E. L. Juvenile hormone in adult eusocial hymenoptera: Gonadotropin and behavioral pacemaker. *Arch. Insect Biochem. Physiol.* **35**, 559–583 (1997).

32. Hartfelder, K. Insect juvenile hormone: from 'status quo' to high society. *Braz J Med Biol Res* **33**, 157–177 (2000).

33. Lillico-Ouachour, A. & Abouheif, E. Regulation, development, and evolution of caste ratios in the hyperdiverse ant genus Pheidole. *Current Opinion in Insect Science* **19**, 43–51 (2017).

34. Nijhout, H. F. & Wheeler, D. E. Juvenile Hormone and the Physiological Basis of Insect Polymorphisms. *The Quarterly Review of Biology* **57**, 109–133 (1982).

35. Nemoto, M. & Hara, K. Ecdysone receptor expression in developing and adult mushroom bodies of the ant Camponotus japonicus. *Dev Genes Evol* **217**, 619–627 (2007).

36. Penick, C. A., Liebig, J. & Brent, C. S. Reproduction, dominance, and caste: endocrine profiles of queens and workers of the ant Harpegnathos saltator. *J Comp Physiol A* **197**, 1063–1071 (2011).

37. Fletcher, D. J. C. & Blum, M. S. Pheromonal Control of Dealation and Oogenesis in Virgin Queen Fire Ants. *Science* **212**, 73–75 (1981).

38. Harris, J. W. & Woodring, J. Elevated brain dopamine levels associated with ovary development in queenless worker honey bees (Apis mellifera L.). *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology* **111**, 271–279 (1995).

39. Sasaki, K. & Nagao, T. Distribution and levels of dopamine and its metabolites in brains of reproductive workers in honeybees. *Journal of Insect Physiology* **47**, 1205–1216 (2001).

40. Boulay, R., Hooper-Bui, L. M. & Woodring, J. Oviposition and oogenesis in virgin fire ant females *Solenopsis invicta* are associated with a high level of dopamine in the brain. *Physiological Entomology* **26**, 294–299 (2001).

41. Beggs, K. T. *et al.* Queen pheromone modulates brain dopamine function in worker honey bees. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 2460–2464 (2007).

42. Eliyahu, D., Ross, K. G., Haight, K. L., Keller, L. & Liebig, J. Venom Alkaloid and Cuticular Hydrocarbon Profiles Are Associated with Social Organization, Queen Fertility Status, and Queen Genotype in the Fire Ant Solenopsis invicta. *J Chem Ecol* **37**, 1242–1254 (2011).

43. Zeng, H., Millar, J. G., Chen, L., Keller, L. & Ross, K. G. Characterization of Queen Supergene Pheromone in the Red Imported Fire Ant Using Worker Discrimination Assays. *J Chem Ecol* **48**, 109–120 (2022).

44. Krieger, M. J. B. & Ross, K. G. Identification of a Major Gene Regulating Complex Social Behavior. *Science* **295**, 328–332 (2002).

45. Gotzek, D., Robertson, H. M., Wurm, Y. & Shoemaker, D. Odorant Binding Proteins of the Red Imported Fire Ant, Solenopsis invicta: An Example of the Problems Facing the Analysis of Widely Divergent Proteins. *PLoS ONE* **6**, e16289 (2011).

46. Pracana, R. *et al.* Fire ant social chromosomes: Differences in number, sequence and expression of odorant binding proteins. *Evolution Letters* **1**, 199–210 (2017).

47. Cohanim, A. B., Amsalem, E., Saad, R., Shoemaker, D. & Privman, E. Evolution of olfactory functions on the fire ant social chromosome. *Genome Biology and Evolution* (2018) doi:10.1093/gbe/evy204.

48. Brelsford, A., Dufresnes, C. & Perrin, N. High-density sex-specific linkage maps of a European tree frog (Hyla arborea) identify the sex chromosome without information on offspring sex. *Heredity* **116**, 177–181 (2016).

49. Lajmi, A., Glinka, F. & Privman, E. Optimizing ddRAD sequencing for population genomic studies with ddgRADer. *Molecular Ecology Resources* 1755–0998.13870 (2023) doi:10.1111/1755-0998.13870.

50. Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and Evolution* **35**, 1547–1549 (2018).

51. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research* **32**, 1792–1797 (2004).

52. Edler, D., Klein, J., Antonelli, A. & Silvestro, D. raxmlGUI 2.0: A graphical interface and toolkit for phylogenetic analyses using RAxML. *Methods Ecol Evol* **12**, 373–377 (2021).

53. Andrews, S. FastQC: a quality control tool for high throughput sequence data. (2010).

54. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

55. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* **27**, 2957–2963 (2011).

56. Rochette, N. C., Rivera-Colón, A. G. & Catchen, J. M. Stacks 2: Analytical methods for paired-end sequencing improve RADseq-based population genomics. *Mol Ecol* **28**, 4737–4754 (2019).

57. Inbar, S. *et al.* Genetic basis of cuticular hydrocarbons in the desert ant. *In prep*.

58. Langmead B & Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9**, 357–359 (2012).

59. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).

60. Browning, B. L., Zhou, Y. & Browning, S. R. A One-Penny Imputed Genome from Next-Generation Reference Panels. *The American Journal of Human Genetics* **103**, 338–348 (2018).

61. Browning, B. L. & Browning, S. R. Improving the Accuracy and Efficiency of Identity-by-Descent Detection in Population Data. *Genetics* **194**, 459–471 (2013).

62. Weir, B. S. & Cockerham, C. C. Estimating F-Statistics for the Analysis of Population Structure. *Evolution* **38**, 1358 (1984).

63. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *The American Journal of Human Genetics* **81**, 559–575 (2007).

64. Altenhoff, A. M. *et al.* OMA standalone: orthology inference among public and custom genomes and transcriptomes. *Genome Res.* **29**, 1152–1163 (2019).

65. Bryant, D. M. *et al.* A Tissue-Mapped Axolotl De Novo Transcriptome Enables Identification of Limb Regeneration Factors. *Cell Reports* **18**, 762–776 (2017).

66. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* **17**, 261–272 (2020).

67. HAPpy-ABCENTH https://github.com/biorover/HAPpy-ABCENTH.

68. Forêt, S. & Maleszka, R. Function and evolution of a gene family encoding odorant binding-like proteins in a social insect, the honey bee (*Apis mellifera*). *Genome Res.* **16**, 1404–1413 (2006).

69. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**, 772–780 (2013).

70. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

7. Supplementary material

Supplementary Figure S1: Matrix of average aggression scores among nests. The intensity of cell color is proportional to the aggression score. The first four rows provide nest IDs, kinship, genotype at the supergene, number of queens found, and mitotypes (monogyne/polygyne). The monogyne form is denoted in blue and polygyne in green.

1   Supplementary Figure S2: (a) Histogram of aggression scores. (b) Aggression scores

2   across social forms assigned based on mitotype.

Supplementary Figure S3: LD decay plot for *Cataglyphis niger* chromosomes. Blue dots

2   represent the LD at corresponding SNP distances in monogyne samples, while green

3   dots represent polygyne samples. The blue dashed and solid green lines indicate the

4   average LD at each SNP distance in monogyne and polygyne samples, respectively.

1   Supplementary Figure S4: A plot of linkage disequilibrium (LD) along chromosome 2 in

2   (a) polygyne and (b) monogyne samples.

3   (a)

5   (b)

1 Supplementary Figure S5: Phased genotypes of 65 daughters of a single polygyne

2 queen and a single father (full-sisters). Only sites that were polymorphic in these

3 samples are shown. The supergene region in chromosome 2 stands out for having

4 100% heterozygosity.

Supplementary Figure S6: Genomic data of broad scale sampling of populations of *C.*

niger across Israel from Reiner Brodetzki et al. (2019), showing the presence of the

supergene on chromosome 2. Monogyne and polygyne samples were assigned based

on the mitotype. (a) Genome-wide F_{ST} analysis using broad sampling of *C. niger* also

shows a large number of high F_{ST} SNPs on chromosome 2. (b) The high F_{ST} region on

chromosome 2 is consistent with our data from a single population. (c) Genotypes at the

high F_{ST} SNPs show the association between mitotypes and supergene genotypes.

Samples bearing the polygyne mitotype are mostly heterozygous MP and monogyne

samples are homozygous MM at the supergene. (d) Distribution of supergene

genotypes across Israel.

(a)

(b)

1   (c)

1   (d)

Supplementary Figure S7: Synteny plot with 22 hymenopteran species based on one-

to-one orthologous genes on the social chromosome of *Cataglyphis niger*.

Supplementary Figure S8: Dotplot of *Solenopsis* and *Cataglyphis* social chromosomes
with gene names for one-to-one orthologs. Red dots indicate genes in the supergene
region in both species.

Supplementary Figure S9: Subtree of the Odorant Receptor (OR) gene tree including genes in the *Cataglyphis* and *Solenopsis* social chromosomes. CnigOR = *Cataglyphis niger* OR; CfOR = *Camponotus floridanus* OR; SiOR = *Solenopsis invicta* OR; LhOR = *Linepithema humile* OR.