1
2

Title: Cortical microtubules act as a template to organize nano-scale patterning of exocytosis

Authors: Jelmer J. Lindeboom[1,2]*, Ryan Gutierrez[1,3], Viktor Kirik[4] and David W. Ehrhardt[1,3]*

Affiliations:
[1]Department of Plant Biology, Carnegie Institution for Science; Stanford, CA 94305, USA.

[2]Current Address: Novo Nordisk Foundation Center for Genomic Mechanism of Disease, Broad
Institute of MIT and Harvard, MA 02142, USA.

[3]Department of Biology, Stanford University; Stanford, CA 94305, USA.

[4]School of Biological Sciences, Illinois State University; Normal, IL 61790, USA.

*Corresponding authors
Email: ehrhardt@stanford.edu
14   jelmer@broadinstitute.org

Abstract

The microtubule cytoskeleton plays an important role in targeting and organizing exocytosis, enabling cellular morphogenesis, motility and polarized transport. However, in many organisms and cell types, the mechanisms by it does so are not well understood. Here, we identify the exocytotic protein KEULE, an essential SEC/MUNC, as a dynamic marker for exocytosis that forms clusters that are associated with and required for discrete exocytotic events in growing *Arabidopsis* cells. Live cell imaging revealed that these clusters, including those associated with verified exocytotic events, have stereotyped and distinct patterns of protein accumulate and loss, suggesting discreet phases in the assembly, function and of exocytotic protein complexes. Super-resolution image analysis revealed that exocytotic events marked by these KEULE clusters are not randomly distributed at the plasma membrane but are significantly enriched in ~180 nm wide linear domains that flank cortical microtubules associated with the plasma membrane, while being under-represented in adjacent domains ~520 nm wide. Thus, the cortical microtubule array acts as a highly organized and dynamic two-dimensional template that effectively redistributes and patterns a significant portion of exocytotic traffic. This positioning mechanism differs from previously reported mechanisms based on directed transport to microtubule ends, allowing for linear and ordered deposition of cargoes to the cell membrane and extracellular matrix to support functions such as cell morphogenesis. The role microtubule arrays near or at the plasma membrane play in organizing exocytosis may be conserved in other cell types and organisms.

Significance Statement

Cells deliver transmembrane proteins and extracellular matrix material through exocytosis, the process of vesicular fusion to the plasma membrane. The plasma membrane contains specialized subdomains crucial for cell shape and function, requiring precise spatial control over exocytosis. Microtubules organize exocytosis partly by transporting vesicles from the cell body to the periphery along oriented centrosomal arrays, but how acentrosomal microtubule arrays organize exocytosis remains poorly understood. We found that in plant cells, which completely lack centrosomes, microtubules define linear and highly structured plasma membrane domains of increased and decreased activity of individual exocytotic events. This demonstrates a means by which ordered microtubule organization may generate functional anisotropy crucial for morphogenesis in plant cells, a mechanism possibly shared by other acentrosomal arrays.

Main Text

Introduction

The ability to organize biomolecules at the cell cortex was a fundamental innovation in the evolution of cellular life, enabling essential biological functions including cellular morphogenesis, motility and polarized transport. Targeted exocytosis, non-random positioning and fusion of exocytotic vesicles at the plasma membrane, is a crucial mechanism for creating such cortical organization. Across phyla, the actin and microtubule cytoskeletons have been implicated in spatial organization of exocytosis. This functionality is exemplified in plant and fungal cells by cellular tip growth, and in animal cells by polarization of mammalian epithelia, cell migration and the positioning of focal adhesions[1,2]. The molecular polarity of microtubules plays an important role in how their function in exocytotic targeting is understood. For example, polarized arrays of microtubules created by nucleation from centralized organizing bodies - such as centrosomes or Golgi - can serve as directional tracks for motor-driven transport of exocytotic vesicles from where they are created in the cell interior to peripheral destinations at the plasma membrane[2,3]. More refined positioning of exocytosis is proposed to be accomplished by association of microtubule plus ends with specific locations at the plasma membrane, including focal adhesions[2]. Polarized arrays of microtubules can also organize exocytosis through directed transport of

factors that establish membrane domains that serve to target exocytotic traffic, as exemplified during tip
growth in fission yeast[4]. However, many cells across diverse organisms feature interphase microtubule
arrays that are not centrally organized and have architectures where microtubules are not clearly polarized
with respect to the plasma membrane[5]. Acentrosomal arrays are required in important biological
processes that require exocytosis, such as morphogenesis, but relatively little is understood about how
they may function to target and pattern exocytotic traffic.

Gaining an understanding of exocytotic targeting mechanisms requires an ability to observe
individual exocytotic events with respect to targeting machineries and a means to verify successful
delivery of exocytotic cargo. Observing individual exocytosis events has been challenging in many cell
models because microtubules, exocytotic vesicles and the machineries of exocytosis are often densely
concentrated. Axially growing cells in the epidermis of higher plants have features that provide an
excellent experimental platform to make such observations and for investigating how acentrosomal
microtubule arrays may target and pattern exocytosis. In contrast to cells with microtubule arrays that are
polarized with respect to the cell interior and the plasma membrane, these cells feature highly ordered
arrays of microtubules that lie at the cell cortex in parallel to and in contact with the plasma membrane
along their lattices. Large regions of the plasma membrane lie in a single focal plane, where the individual
microtubules and microtubule bundles that constitute these arrays are easily visualized due to their quasi-
2-dimensional nature and comparatively low density[6]. In previous studies, we determined that cellulose
synthase complexes (CSCs), large integral membrane protein complexes that synthesize and extrude
cellulose into the cell wall, can be used to observe individual exocytotic events at the plasma membrane
by allowing dynamic tracking of both exocytotic vesicles and the successful delivery of cargo to the
plasma membrane[7–9]. By tracking CSCs while simultaneously imaging dynamic cortical arrays, we
discovered that CSCs were preferentially delivered to the plasma membrane at locations defined by
microtubules. This targeting did not involve vesicular transport along microtubules, nor did it involve
defining exocytotic domains by end-on association of microtubule ends with the plasma membrane[7], thus
suggesting a previously undescribed mechanism for microtubule-dependent exocytotic targeting.
However, CSC's themselves have been shown to interact with microtubules via a specific protein,
Cellulose Synthase-Interaction 1 (CSI1), an interaction that can also tether CSC-containing vesicles and
organelles to microtubules[10–13]. Thus, it was not clear if this exocytotic patterning mechanism may be
widely employed in plant cells, or if it is unique to delivery of CSCs to the plasma membrane.

Microtubule targeting of exocytosis is of special interest in the context of plant cell
morphogenesis because ordered cortical microtubule arrays are essential to create functional cell shape.
Depolymerization of microtubules by the drug oryzalin causes isotropic growth and a complete loss of
genetically determined cell shape. The predominant hypothesis for the function of cortical microtubules in
directing cell growth is that they guide and organize the trajectories of CSCs as they synthesize and
extrude cellulose microfibrils into the cell wall, thus creating material and mechanical anisotropy that
directs expansion of the cell wall (which is under tension generated by high turgor pressure). We
confirmed that individual CSC trajectories are in fact guided by microtubules by live cell imaging
studies[14]. However, disruption of both microtubule guidance of CSCs and delivery of CSCs to the plasma
membrane in *csi1* null mutants causes only a modest loss of cell growth anisotropy[10,11,13]. In addition,
although complete loss of growth anisotropy is caused by microtubule depolymerization, this phenotype
is not due to a loss of CSC delivery to the plasma membrane[7] nor a lack of ordering in CSC
trajectories[7,15,16]. Further, cell growth rate does not appear to be impaired by oryzalin treatment and loss of
cortical microtubules[17]. These observations suggest that cortical microtubules arrays have additional
important functions in establishing cell shape, prompting us to ask if cortical microtubules may guide
cellular morphogenesis by acting as a general targeting mechanism, beyond CSC's, to pattern exocytosis
during cell growth.

To determine if individual exocytosis events in growing plant cells are broadly patterned by
cortical microtubule targeting, we require a dynamic biological marker that identifies single exocytotic

events and is involved in a wide range of exocytotic traffic to the plasma membrane. To date, such a
marker has not been identified. Imaging studies of core exocytotic proteins, including soluble *N*-
ethylmaleimide-sensitive factor attachment protein receptors (SNARE proteins)[18–20] and components of
the exocyst complex[21,22], have not revealed a clear relationship between the presence and dynamics of
these proteins with verified exocytosis events nor have they shown association of these proteins with
microtubules[23].

To address this gap in tools for studying individual exocytotic events, we investigated use of the
essential Sec/Munc (SM) protein KEULE. SM proteins are essential regulators of SNARE-mediated
vesicle fusion. In exocytosis, SNARE proteins anchored to the vesicle interact with those on the plasma
membrane and form a membrane bridging complex with other partners that enables membrane fusion[23–25].
The soluble SM proteins are known to bind to SNAREs, and evidence from genetic studies together with
in vivo and *in vitro* experiments suggest they function in vesicle docking, SNARE complex assembly and
vesicle fusion[26–31]. In plants, the *Arabidopsis* Sec1 homolog KEULE was identified in a genetic screen for
embryonic pattern mutants[32], where loss of function mutations are lethal, causing bloated and
multinucleate cells[33]. Consistent with other SM proteins, KEULE interacts genetically and biochemically
with Qa-SNAREs, specifically KNOLLE (SYP111) and SYP121, and has been shown to play essential
roles in cell plate assembly during cytokinesis and exocytotic traffic to the plasma membrane[18,33–37]. The
demonstrated function of KEULE in exocytosis, combined with its severe loss of function phenotype due
to lack of exocytosis suggested that KEULE might serve as an excellent candidate for a generalized
reporter of exocytosis.

The molecular and structural mechanisms by which SM proteins interact with and regulate
SNARE complex assembly have been investigated intensively by genetic, biochemical*, in vitro* and
structural studies[26–31], and SM proteins have been imaged in live cells in several systems[38–45]. However,
important features of their behaviors and biological function in living cells remain to be uncovered. These
questions include the fine scale cellular distributions of SM proteins and clusters, the association of SM
clusters with exocytosis events, the relationship of these exocytosis events to potential organizing
machineries and the dynamics of SM assembly and loss at exocytosis sites, especially prior to and during
vesicle fusion.

Here, we develop and apply automated and high precision analysis methods to detect and track
tens of thousands of individual and fluorescently tagged KEULE clusters to investigate their dynamics
and to map their positions at the plasma membrane in living cells. We evaluate these clusters as
exocytotic events by use of CSCs as a validated cargo and employ genetic analysis to test for the function
of KEULE in CSC delivery. Finally, we use high precision mapping of KEULE clusters and
microtubules, combined with Bayesian methods, to quantitatively determine if the cortical microtubule
cytoskeleton as a template in patterning exocytosis.

Results
149
Functional dynamics of tdTomato-KEULE cluster formation

To visualize KEULE protein *in vivo*, we created a tdTomato-KEULE fluorescent reporter
constructed from the chromosomal gene with native upstream regulatory sequence and introduced it into
Arabidopsis plants heterozygous for a *keule* deletion allele. To obtain plants for study of KEULE
localization and dynamics, we screened for self-fertilized progeny of these transgenic plants that exhibited
rescue of *keule* loss of function phenotype by the tagged KEULE protein. These plants exhibited full
rescue of plant growth and development through the 5-day seedling stage used for our further studies
(Fig. S1, Methods).

In dark-grown *Arabidopsis* hypocotyl cells expressing the tdTomato-KEULE construct, we
observed KEULE signal appear as densely distributed and temporally dynamic puncta at the plasma
membrane (Fig. 1A). The dynamics of signal increase and decay were apparent in time-lapse movies

(Movie S1), kymographs of signal intensity over time along transect lines (Fig. 1B) and in sequential
image samples of individual puncta (Fig. 1C). In all these events, a steady increase in signal intensity was
observed, followed by an abrupt decrease and disappearance of signal. This behavior indicates the
accumulation and loss of many fluorescently labeled KEULE proteins, which we will refer to hereafter as
clusters.

The dynamics of tdTomato-KEULE accumulation and disappearance are presumably related to
its function. Therefore, we reasoned that a quantitative and robust picture of KEULE cluster dynamics
would be useful in gaining insight into the underlying molecular mechanisms of exocytosis. To
accomplish this, we collected a large-scale and high-resolution dataset comprising thousands of individual
events. We acquired streamed images at 200 ms intervals and performed automated detection and frame-
to-frame linking of KEULE cluster signal using μTrack[46]. These track data were then used to extract
position and intensity information by Maximum Likelihood Estimate 2D Gaussian fitting[47] (see Methods,
Movie S2). To define events of signal appearance, accumulation and loss, we used a minimum sample of
4 sequential acquisitions, which made our shortest measured lifetimes approximately 800 ms. The
lifetime distribution of tdTomato-KEULE clusters showed a large range, from this minimal event length
of 800 ms to over half a minute. These data followed a roughly exponential distribution with a mean of
2.34 s and ~95% of lifetimes being less than 7 seconds long (Fig. 1D).

The large lifetime range of observed KEULE cluster events might represent a collection of
functional classes. If so, such classes may by differentiated by their kinetics. To assess the kinetics of
KEULE signal accumulation and loss, and to ask if these kinetics might vary as a function of cluster
lifetime, we leveraged our large data set to calculate KEULE kinetics for each 200 ms lifetime increment.
Fig. 1E shows average intensities for all observed clusters with lifetimes of 4.2, 4.4 and 4.6 s. All these
plots show a similar pattern: an initial steep rise in intensity, followed by a steady increase, then a short
plateau phase, and ending with a sharp decrease. When we extended this analysis over a larger range of
lifetimes, we observed evidence that the shortest-lived events accumulated tdTomato-KEULE intensity at
a slower rate, suggesting they might be a functionally different class (Fig. 1F). At longer lifetimes, the
plateau phase became more extended, suggesting that these events may be paused or stalled (Fig. 1F). To
quantitatively compare differences in accumulation kinetics as a function of cluster lifetime, we grouped
events into three duration classes; short (0.8 to 1.6s), medium (1.8 to 6.2s) and long (6.4+s) – and then
averaged the intensities at each 200 msec timepoint (Fig. 1G, for details see Methods). The timepoints
following the plateau phase were omitted to aid comparison of signal accumulation trajectories. This
analysis confirmed that, indeed, the short events show significantly slower signal accumulation (Fig. 1G).
Interestingly, the medium and long cluster lifetimes show nearly identical trends in signal intensity
accumulation over time, indicating that they might be part of the same functional population during
KEULE recruitment/assembly. It is noteworthy that for events up to ~5s, maximum cluster signal
intensity increased with cluster lifetime. A possible explanation for this relationship is that longer
accumulation events represent the assembly of exocytotic apparatus at larger vesicles. SEM studies of *in*
vitro vesicle fusion, for example, have shown that more SNARE proteins are present at the fusion sites of
larger vesicles as compared to smaller vesicles[48,49] In events lasting longer than about 5 seconds
maximum signal no longer increased with lifetime but instead showed a prolonged plateau phase,
possibly indicating stalled exocytosis events or alternatively, an extended checkpoint.

Clusters of tdTomato-KEULE coincide with CSC delivery to the plasma membrane

To place SM protein accumulation in context of functional exocytosis, we took advantage of our
previous work on delivery of CSCs to the plasma membrane[7]. CSCs are assembled in the Golgi system
from multiple CESA catalytic subunits[50,51], move through the trans-Golgi network (TGN) and are budded
off into vesicles that fuse with the plasma membrane[7]. Because CSCs are formed of multiple CESA
subunits, they can be labeled by fluorescent proteins in sufficient number to allow their detection and the

tracking of individual complexes during the process of exocytosis. Labeled vesicles display rapid and
erratic dynamics that abruptly become stabilized at the optical plane of the plasma membrane with a mean
duration of about 60 seconds, followed by a transition to slow, steady and typically linear motility of
between 250 and 450 nm/min. The transition erratic to static movement is interpreted as evidence for
vesicular capture and tethering at the plasma membrane, and the transition to slow and steady movement
indicates successful vesicle fusion and delivery of CSCs, which are propelled by the synthesis and
extrusion of cellulose microfibrils[7,14]. This stereotyped behavior makes CSCs an excellent reporter for
successful exocytosis, revealing a time and place for vesicle tethering, and distinguishing vesicle tethering
at the plasma membrane from successful vesicle fusion and cargo delivery.

To visualize KEULE together with CSCs, we generated plants that express tdTomato-KEULE
and CIT-CESA6, a catalytic CSC subunit labeled with Citrine fluorescent protein[14] in a *keule*/*cesa6*
double mutant background (Methods). CIT-CESA6 signal is relatively dense in the plasma membrane of
rapidly growing cells so we photobleached the CIT-CESA6 signal over a region of the plasma membrane
to aid detection of new CSC arrivals and their relation to td-Tomato-KEULE signal (Fig. 2A, Movie S3).
For each CSC arrival at the optical plane of the plasma membrane, we determined the first frame in which
the CESA6 signal became stationary (an indication of interaction and tethering at the plasma membrane).
We then identified the nearest KEULE cluster in a range of -10 to 50 seconds of the first stable CESA6
frame and determined the distance between the centroid of the CESA6 signal and the centroid of the
nearest KEULE signal by maximum likelihood 2D Gaussian fitting (See Methods, Fig. 2B and D). A
search range starting at -10 seconds was chosen because the exact frame of stationary transitions was in
some cases obscured by signal from clusters of other labeled organelles such as Golgi bodies and TGN. If
KEULE is involved in the exocytotic delivery of CSCs to the plasma membrane, we expect to observe co-
localization between CSC delivery events and KEULE clusters. However, CSC deliveries, as defined by
static tracks, are relatively sparse and long lived as compared to KEULE cluster events, so some degree of
colocalization in space and time is expected by chance alone. To assess statistically the spatial and
temporal association of CSC delivery events and KEULE clusters, we generated a simulated dataset of
random locations within the photobleaching areas and identified the nearest KEULE spot to the random
locations within the same time window we used for our experimental observations. We then compared the
distributions of distances to the nearest KEULE cluster between the observed CSC deliveries and the
simulated data (Fig. 2B) and found that the experimental data were strongly skewed to shorter distances
(p < 1E-06, Kolmogorov-Smirnov test). The pattern of over- and under-representation of observed with
respect to simulated data is more easily seen in a plot of the bin-by bin differences where shorter KEULE-
microtubule distances are seen to be over-represented in bins below 200 nm and underrepresented in bins
of longer values (Fig. 2C). Using a Bayesian method (See Methods), we found that the most likely values
of the experimental biases from the null hypothesis of a random association were significantly above that
expected by chance alone for distance bins below 120 nm (Fig 2E and F) and significantly lower than
expected in bins of longer distances from 200 to 480 nm. Based on these data we defined colocalization
as observing a KEULE cluster centroid within 200 nm from the position of CSC signal in the first frame
in which it was observed to stabilize (and in a time window of -10 to 50 seconds around that frame). By
these criteria, colocalization occurred in 81% (157 out of 194) of observed delivery events compared to
29% (13035 out of 45051) of random locations at the plasma membrane (p < 1.4E-50, Fisher's exact test).

We considered that our automated experimental protocol may allow for underestimation of
colocalization between CSC delivery and KEULE. For example, the time interval used for these
experiments is 2.11 s, of which 400 ms is used for tdTomato-KEULE excitation and image acquisition.
Since a substantial portion of KEULE cluster lifetimes is shorter than the ~1.7 s gaps between image
acquisitions, we do not expect to detect all KEULE clusters within the -10 to 50 second experimental
window. In addition, not all excited KEULE foci are successfully identified by the spot detection and
filtering routines used for large-scale data collection. To understand how these two cases, no excitation
and no detection, affect our colocalization results, we performed a hand-curated analysis using a custom
MATLAB GUI. For each CSC delivery, we examined whether there was overlap of KEULE signal with

the centroid of the CSC signal in the first frame of positionally stable KEULE signal (tethering frame).
We found that in 92% (159 out of 172) of the cases, KEULE signal was observed to overlap the CSC
position in the tethering frame within -10 to 50 s of CSC stabilization, suggesting that our automated
analysis, while free of possible observer bias, may underestimate the true degree of colocalization
frequency of CSC exocytosis events and KEULE.

If KEULE clusters function in CSC exocytosis after vesicle tethering, we expect that not only
should they colocalize with CSC signal, but that the time intervals between vesicle tethering and the
recruitment of KEULE should have a non-random distribution. We tested this prediction by comparing
the delay times for arrival of the nearest KEULE spot following the tethering frame in the observed vs the
simulated KEULE data sets. The observed delay time distribution was significantly different from the
random distribution of the simulated data (p < 7.8E-04 Mann-Whitney U test) (Fig. 3A), having a peaked
distribution with a median of 2.111 s and a mean of 7.92 s, suggesting that KEULE is recruited and
functions within several seconds after the CSC vesicle is stabilized and tethered at the plasma membrane.

The durations of KEULE clusters associated with CES deliveries had a distribution similar in
character to that observed For KEULE clusters in general (Fig. 3B, compare to Fig.1D), with both
appearing roughly exponential but long tailed. While lifetimes of clusters associated with CESA
deliveries spanned a similar range of durations, those associated with CESA delivery were observed to
have a significant bias towards longer values, with clusters associated with CESA having a median
duration of 4.3 seconds, as compared to 2.3 seconds for clusters in general. To test the significant of this
observed bias, we compared the relative frequencies of durations longer than two seconds and found a
highly significant difference - 59% vs 37% for CESA delivery-associated KEULE clusters events vs
clusters in general (Fig. 3C, p=6.07E-09 Fisher's exact test). These results indicate that both short-lived
and long-lived KEULE clusters are associated with verified exocytosis events and that those associated
with CESA complexes are modestly biased towards longer lifetimes. Taken together, our data now
provide insight into the kinetics of SM protein recruitment to sites of exocytosis and reveal the time of
vesicle fusion following exocytotic vesicle stabilization and tethering at the plant cell plasma membrane.

Our data show that KEULE clusters are associated with confirmed sites of CSC exocytosis in
both space and time, but is KEULE function required for CSC exocytosis? To answer this question, we
quantified CSC delivery in selfed progeny of parental plants homozygous for a transgene expressing CIT-
CESA6, heterozygous for a *tdTomato-KEULE* allele, and homozygous for *cesa6* and *keule* loss of
function alleles (See Methods). Seedlings tested for loss of KEULE function were identified by
morphological *keule* phenotype (severely reduced in size, large and misshaped cells) and little to no
tdTomato-KEULE signal. In these mutant seedlings, we found that the CSC delivery rate was reduced by
over 17-fold as compared to that in wildtype (p<0.0001, Wilcoxon two sample test, Fig. 2G). Taken
together our results indicate that KEULE plays an essential role in plasma membrane delivery of primary
CSCs, and that CSC delivery is spatially and temporally associated with KEULE clusters which display
stereotyped dynamics of formation and loss.

KEULE dynamics are sensitive to drugs and treatments that reduce exocytosis

We previously reported that the number of active CSCs in the plasma membrane were rapidly and
significantly reduced by treatment with mannitol (osmotic stress) and the herbicide isoxaben[7].
Measurements of CSC delivery revealed that mannitol treatment also caused a 100-fold reduction in the
delivery rate of CSC's, indicating that CSC loss at the plasma membrane was likely caused primarily by
inhibition of exocytosis. Likewise, isoxaben has been characterized as a specific cellulose synthase
complex inhibitor[52], but its mode of action is also not well understood. Here we (1) determine if isoxaben
treatment also reduces CSC exocytosis and (2) ask if KEULE dynamics may provide insight into how
osmotic stress and isoxaben act to inhibit exocytosis.

To address the above questions, we treated seedlings expressing tdTomato-KEULE and Cit-
CESA6 with either 200 mM mannitol, 100nM isoxaben or carrier only (0.01% DMSO, mock) and

assayed CSC delivery to the plasma membrane as described above. We found that, like osmotic stress caused by mannitol, isoxaben is also a potent inhibitor of CSC exocytosis, reducing CSC delivery rate by ~70 fold (Fig. S2). By contrast to CSC delivery, the overall frequencies of KEULE cluster formation after both mannitol and isoxaben treatment were not significantly lower than that observed for mock treatment, indicating that neither acts to absolutely block initial KEULE recruitment and cluster formation rates (Fig. S3A). However, when we examined KEULE accumulation dynamics we observed striking differences among treatments (Fig. S3B, C). In cells treated with mannitol KEULE signal accumulated more slowly and to lower levels in the short and intermediate lifetime classes events (Fig. S3B). Intriguingly, the trajectories of long duration clusters (6.4 s or longer) exhibited a markedly different pattern. KEULE signal trajectories for both mock and mannitol treated cells tracked each other closely for the first 5 s or so, but rather than reaching a plateau after 5 seconds, signal in mannitol treated cells continued to increase steadily until the transition to rapid signal loss (Fig. S3B). Taken together, these data suggest that there may be two different mechanisms by which mannitol treatment and osmotic stress affects KEULE dynamics, depending on their duration class: one mechanism that acts negatively and from cluster inception that reduces the rate and level of KEULE accumulation in short and intermediate events, and one which acts at a later stage in long duration events that prevents transition to the normally occurring plateau stage. By contrast, we observed that isoxaben treatment slowed accumulation of tdTomato-KEULE across events of all durations (Fig. S3C).

We observed related differences among treatments in KEULE cluster lifetimes. In comparing the distributions of cluster lifetimes in mock and mannitol treated cells, a bimodal effect was again observed, with both shorter and longer lifetimes being over-represented in comparison with mock treatment (and intermediate events being under-represented) (Fig S4A, B, D). As was the case with intensity dynamics, isoxaben treatment showed more of a unimodal effect, with the lifetime distribution being shifted to the left (Fig. S4B, C, E). Mannitol and isoxaben also caused diverging effects on persistence of KEULE clusters, as assessed by survival analysis (Fig. S4F, G). Taken together, our data show that KEULE cluster dynamics are sensitive to perturbations that affect exocytosis and reveal evidence that mannitol and isoxaben treatments affect these dynamics in distinct ways, suggesting they may perturb exocytosis by different mechanisms.

Cortical microtubules pattern KEULE cluster distribution and exocytosis

Having established dynamic KEULE clusters as markers of exocytotic events, we noted that the flux of KEULE cluster formation, 144 events per μm^2 per h (Fig. S3A), is ~85-fold higher than that of CSC delivery (1.59 evens per µm2 per h, Fig. 2G). We therefore utilized dynamic imaging of KEULE and microtubules together to ask if cortical microtubules pattern the distribution of KEULE protein and to determine if targeting of exocytosis is specific for CSCs or if it is a broad and fundamental function for the cortical microtubule cytoskeleton in growing plant cells. Further, we took advantage of the ability to detect thousands of KEULE clusters to map the spatial range of influence of microtubule targeting with super-resolved precision.

To visualize KEULE cluster events together with cortical microtubules, we performed live cell imaging at 1 second intervals in hypocotyl cells of *Arabidopsis* seedlings expressing GFP-TUA5 and tdTomato-KEULE in a *keule* knock-out background (Fig. 4A, Movie S4). We tracked individual KEULE events at the plasma membrane and used MLEwG fitting to extract precise positional information (Fig. 4B, Methods), after which we identified the frame with peak signal intensity as the reference point for each detected KEULE cluster position (Fig. 4C panels 1-3). We then measured signal intensity in the microtubule channel along a series of scan lines that were rotated at 9° intervals around the centroid of the KEULE cluster (Methods and Fig. 4C panel 4). On each of these scan lines, we detected all microtubule signal peaks and fit them with a modified quasi-1-dimensional MLEwG fit (Methods, Fig. 4D). Since the microtubule signal is best represented as a quasi-1-dimenional Gaussian orthogonal to the microtubule

axis, we determined the orientation of the nearest microtubule as identified by peak analysis and used the scan line most orthogonal to the microtubule angle for determining the location of the microtubule central axis closest to the KEULE spot in question (Fig. 4C, panel 5). For example, the intensity of KEULE signal in photoelectrons for the most orthogonal scan line in Fig. 4C panel 5 is shown in Fig. 4D panel 1, the triple Gaussian 1D MLEwG fit in panel 2, and the decomposed individual peak fits in panel 3. We used the fit positional coordinates of the nearest microtubule signal peak and the KEULE cluster to calculate the distance and the photoelectron fit values to estimate the standard error of the measurement (See Methods and Fig. 4C panel 6). We repeated this process for all detected KEULE tracks in each movie to assess the spatial relationship between KEULE clusters and the nearest microtubules to those clusters (Fig. 4E, panel 1).

To ask if the measured distribution of KEULE to microtubule distances differs from that expected if KEULE cluster positions are independent of microtubule locations (the null hypothesis), we first needed to determine what the expected distributions are under the null hypothesis when our detection and analysis routines are applied to the image datasets we used for our experimental observations. For example, microtubule array structure, which varies within and among cells, will have a strong influence on this distribution, as will the dimensions of our detection kernels. Therefore, we generated a simulated set of random locations within each region of interest across all image frames in the movies and calculated the distances from these synthetic KEULE foci to the nearest microtubule (Fig. 4E, panel 2). When the experimental and expected distributions were plotted together as relative frequencies, they showed evidence for distinct differences (Fig. 5A). The spatial pattern of under- and over-representation is more easily visualized in a plot of the differences between observed and expected frequencies (Fig. 5B). To quantify over- or under-representation of the experimental data given the expected data under the null hypotheses, we used a Bayesian method to determine the most likely positive or negative bias at each distance bin (Fig. 5C and D and Methods). We found that at KEULE-microtubule distances smaller than 180 nm, the maximum likelihood bias was significantly positive (more frequent than expected) and between 180 and 700 nm this bias was significantly negative (less frequent than expected). At distances > 700 nm we again found a positive, but smaller, bias (Fig. 5D).

When examining patterns of KEULE foci, we also observed that sites of previous exocytosis events were 3.4 times more likely to show consecutive exocytosis events within 60 seconds than random sites at the plasma membrane (Fig. S5A-E,G). Such "hot spots" were also significantly over-represented within 180 nm of microtubules, but only modestly so (1.25 times more likely) (Fig. S5F,G).

Taken together, our results reveal that microtubules spatially organize exocytosis machinery by increasing the frequency of KEULE cluster formation within about 180 nm of the center of the microtubules, a distance about 7 times the dimension of a single microtubule, while at the same time decreasing the probability of KEULE cluster formation distally over a region that is about 3 times as wide. This acts effectively as a concentrating mechanism for KEULE cluster events that is patterned by cortical microtubule organization. As these KEULE clusters overlap heavily in their dynamic characteristics with those observed associated with validated exocytosis events of CSCs, this is strong evidence that microtubule organization of exocytosis is not confined to guiding insertion of CSCs but is a more generalized feature of exocytosis in growing plant cells.

Discussion

To assess imaging of SM/KEULE clusters as a general marker for exocytosis it was important to answer two questions; if there are different types of signal foci that may represent multiple types of cellular structures or functional events, and what features of signal dynamics are associated with exocytotic events. For example, some exocytotic proteins, including V-SNARES and constituents of the exocyst, have been shown to arrive at the site of exocytosis on exocytotic vesicles themselves, while others reside in the cytosol or associate with the plasma membrane and are recruited to the exocytosis site. While we found some evidence for KEULE signal on cytosolic organelles, these foci were relatively rare

and were easily identified by their positional dynamics. Further, in experiments using labeled CSCs as a
trackable exocytotic cargo, no KEULE signal was detected arriving at validated exocytosis sites on CSC-
containing vesicles, indicating that KEULE clusters at the plasma membrane form by recruitment of
KEULE to sites of exocytosis after arrival and interaction of cargo-carrying organelles with the plasma
membrane. Assembly dynamics of KEULE clusters associated with CSC delivery had similar
characteristics as the KEULE cluster population in general, including a wide range of cluster lifetimes
ranging from a couple of seconds to half a minute and stereotyped dynamics of signal accumulation and
loss. The demonstrated functions of SM proteins in mediating SNARE complex formation and membrane
fusion, together with their stereotyped dynamics and tight association with CSC delivery, indicate that
dynamic KEULE clusters are valuable markers of individual exocytotic events.

Our measurements of thousands of KEULE clusters at the plasma allowed us to analyze patterns
of signal accumulation and loss as a function of varying cluster duration - revealing features of, and
raising new questions about, the underlying mechanisms of exocytotic machinery assembly, maturation
and resolution. Over all durations, KEULE recruitment features an initial phase of rapid accumulation,
followed by continued recruitment at a slower rate. This pattern suggests two phases of recruitment with
different kinetics. The second rising phase is near linear, indicating a near constant rate of addition over
time, characteristic of mechanisms such as polymerization, rather than recruitment of SM proteins to a
fixed number of pre-existing binding sites. Such a kinetic might be consistent with the sequential addition
of SM-bound SNARE-PINs to a growing chain of laterally-interacting SNARE-PINs at the vesicle-
plasma membrane interface[53–55]. This accumulation phase ends and transitions to a plateau, after which
signal abruptly decays to non-detectible levels. Both the final signal level and the length of the pause state
correlate positively with cluster assembly time, with longer assembly times having higher signal levels
and longer pause durations. Over the population of observed events, global peak signal is reached at
durations of about 6 seconds, after which longer events involve no further signal accumulation but feature
only lengthened plateau times. The plateau phase could represent a state where all the SNARE complexes
required have been recruited to the fusion site and the components of the fusion super complex are
transitioning to a functionally competent state prior to the enactment of membrane fusion. The plateau
could also reflect an assembly checkpoint. Previous *in vitro* studies have found that greater numbers of
SNARE proteins are recruited to interfaces of large vesicles and planar membranes[48,49]. It is possible,
therefore, that our measurements of KEULE dynamics are shaped in part by a distribution of vesicle sizes,
with longer assembly phases that build a brighter SM signal representing assembly of exocytotic
machinery at larger vesicles.

We found that KEULE dynamics were changed when we disrupted exocytosis with osmotic
stress by treatment with mannitol, or by application of a drug, isoxaben. The changes in dynamics were
qualitatively different for the two antagonists, indicating that they likely act on different processes
affecting exocytosis. These observations indicate that KEULE dynamics, which indicate functional
aspects of vesicle fusion, are a sensitive and useful readout, or marker, for studying exocytosis and agents
that disrupt or alter exocytosis.

High resolution mapping and Bayesian statistical analysis of individual exocytosis events marked
by KEULE clusters revealed that these events are not randomly distributed in growing hypocotyl cells but
are significantly over-represented in domains defined by the cortical microtubule array. This patterning of
exocytosis differs from previously described mechanisms for exocytosis targeting by microtubules. In the
latter, the positioning of microtubule plus ends define the locations for motor-transported delivery of
either the exocytotic vesicles themselves or proteins that act in defining exocytotic domains. By contrast,
our data indicate that the whole microtubule is associated with exocytosis targeting, making the entire
cortical array an ordered, dynamic and steerable template for exocytotic patterning and organization of
the cell cortex (Fig, 6A). As discussed further below, such highly organized patterning of exocytotic
traffic may be useful for creating functional and mechanical asymmetries to support crucial functions like
morphogenesis.

Our high precision measurements of both microtubule and KEULE cluster locations revealed that
microtubules pattern exocytosis both positively and negatively and do so over spatial domains much
larger than the scale of microtubules and microtubule bundles. Single microtubules and bundles of two
polymers are ~25 nm and ~75 nm wide at the plant cell cortex, respectively[56], whereas the combined
width of over-represented and under-represented domains was measured to be ~700 nm. The mean
spacing between ordered microtubules and microtubules bundles in elongating hypocotyl cells is about
twice this distance[57], indicating that microtubules influence exocytosis over most of the observed plasma
membrane these cells (Fig. 6B). In addition, the combination of narrower elevated domains coupled with
wider depleted domains of elevated exocytosis frequency suggests a concentrating effect that further
enhances the relative difference in exocytosis adjacent to microtubules as compared to distances several
hundred nanometers away.

One mechanism that could explain the spatial scales of these microtubule-associated domains is
direct interaction of exocytotic vesicles with cortical microtubules, such as we previously observed for
vesicles containing CSC's[7,58]. Microtubule-tethered vesicles might capture local KEULE associated with
the plasma membrane, perhaps as bound to SNARE partners such as SYP121[34], depleting KEULE
concentration in the nearby plasma membrane and making exocytosis there less likely, thus creating the
observed zones of reduced exocytosis (Fig. 6B). Alternatively, upstream exocytotic machinery, such as
plasma-membrane associated components of the exocyst complex, might be recruited by a microtubule-
organized membrane domain, elevating location concentration while causing depletion and thus vesicle
capture at a distance (Fig. 6B). Evidence that such microtubule-organized membrane domains might exist
has been suggested by observations of the plasma membrane associated actin polymerase Formin1, which
reveals narrow "dark" linear domains defined by cortical microtubules where fluorescently-tagged
Formin1 is excluded[59].

The observed flux of KEULE clusters per unit area and time was about 90-fold higher than that
measured for CSC delivery, therefore far more exocytosis events are associated with domains defined by
microtubules than can be accounted for by CSC delivery alone, implying either that many or most
exocytosis events in these cells are influenced by microtubules to some degree, or that a quantitatively
significant subpopulation of specialized traffic is redistributed, ~13% of KEULE associated events as
measured by our analysis. One possibility for specialized exocytotic traffic targeted by cortical
microtubules are cargoes involved in cell wall biosynthesis and cell wall expansion. Supporting this idea
is that KEULE has been shown to be an essential partner of the Qa-SNARE SYP121, which is implicated
by a growing body of evidence in mediating exocytosis required for cell growth and expansion[19,34,35,60,61].
Cellulose is synthesized at the plasma membrane, while many other structural oligosaccharides, such as
pectins[62], and wall-active proteins, such as expansins[63], are secreted. Targeting this exocytotic traffic by
cortical microtubule arrays to the same locations where newly synthesized cellulose microfibrils are also
being laid down could help coordinate the assembly of these diverse components to build a functional cell
wall. In addition to coordination of wall assembly, it is possible that the patterned delivery of cell wall
oligosaccharides and cell wall active proteins could itself act to support the anisotropic expansion of cell
walls required to create specific cell shape. Targeting the delivery of proteins and other molecules that
promote wall remodeling by ordered microtubule arrays could create linear and parallel microdomains of
increased wall extendibility that lie perpendicular to the main axis of the cell growth - shaping the
anisotropy of cell wall expansion in the same direction as proposed in prevailing models for constraint of
expansion by cellulose microfibrils organized by microtubules[64]. As discussed in the introduction of this
manuscript, an important reason to consider alternative functions for cortical microtubules arrays in
guiding cell wall expansion and cell shape is that experimental evidence suggests that guidance of
cellulose deposition and CSC delivery alone is not sufficient to explain the anisotropy of cell wall
expansion and cell growth. Microtubule targeting of cell wall-active factors by exocytosis could also help
explain redirection of cell growth in response to signals, such as blue-light stimulation of phototrophic
growth. Perception of blue light by Phototropin receptors causes a rapid 90° reorientation of cortical
microtubule arrays[65], but the following redirection of expansion cannot be easily explained by directed

cellulose deposition model since previously synthesized cellulose is deposited perpendicular to the
direction of expansion. Microtubule patterning of cell wall-active factors might allow for redirection of
cell expansion regardless of cellulose orientation.

The insights we gained into patterning of exocytosis by cortical microtubule arrays were enabled
by utilizing an experimental platform and methods that made detecting, measuring and assessing many
thousands of discreet molecular events possible. A protein implicated in mediating a broad swath of a
cell's exocytotic traffic might be expected to be subject to a range of targeting mechanisms, including no
targeting at all. Such a background could easily obscure a meaningful association between the protein and
a hypothesized targeting machinery. Very large sample sizes and appropriate methods to test for
meaningful associations, such as we employed here, can robustly reveal such relationships.

The fact that growing plant cells have a planar microtubule array that is associated with the
plasma membrane, and that exocytotic events as marked by KEULE protein are of relatively low density,
has provided an excellent platform to study how microtubules pattern exocytosis, and do so at the level of
individual exocytotic events. When we consider cellular microtubule function in cellular trafficking, we
typically think of directed transport of cargos by kinesins and dynein. In contrast, this work highlights
that microtubules can also function as scaffolds for organizing exocytosis. This scaffolding function for
microtubules might be considered in other systems where there is indirect evidence that microtubules
regulate exocytosis where they interact with the plasma membrane, such as in focal adhesions and
neuronal growth cones[1,66,67]. In addition, scaffolded guidance of exocytosis should also be considered in
the context of disease. For example, amyotrophic lateral sclerosis (ALS) risk is associated with genetic
variants which reduce levels of functional UNC13A[68,69], and with genetic variation within several
microtubule associated proteins[68–71]. It is possible that these proteins all converge on the same biological
process of microtubule dependent patterning of exocytosis.

Materials and Methods

Generation of transgenic lines
The *pKEULE-tdTomato-KEULE* construct was generated by modifying the Gateway-compatible
vector pMDC43[72]. A 2049-bp region immediately upstream of the start codon of *KEULE* (At1g12360)
was amplified with primers CGTTTAAACTACAGTTCCATTGGAGCAAC and
TAGGTACCTGCGATCTAACAATTT CAGACTTC. The tdTomato gene was amplified from a donor
vector[73] with primers CAGGTACCATGGTGAGCAAGGGCGAGGAGGT and TGGCGCGCCCTT
GTACAGCTCCTCCATGCCGT. The 35S promoter of pMDC43 was exchanged with the *KEULE*
upstream regulatory region via PmeI and Acc65I sites, and GFP in pMDC43 was exchanged with
tdTomato via KpnI and AscI sites. The full-length *KEULE* gene was amplified from genomic DNA using
primers GGGGACAAGTTTGTACAAAAAAGCAG GCTTGACCATGTCGTACTCTGACTCC and
GGGGACCACTTTGTACAAGAAAGCTGGG TTTATTTGGAGATCGTCTA and then inserted into
the modified pMDC43 vector using Gateway cloning technology (Invitrogen) as described previously[65].
The construct was introduced into plants heterozygous for the *keule* MM125 deletion allele[37] via
Agrobacterium-mediated transformation[74,75]. T2 plants were genotyped for the *KEULE* locus using PCR:
primers CGACCTTATCAGAGAGCAAGG and Keu-as0 (ACAGGGATTTGATTTGAGATCTGG)
amplified a 1661-bp fragment from the wild-type allele and a 1505-bp allele from the deletion allele.
Additionally, the primer pair GTCTGTTAACGGCTGCAGAA and Keu-as0 was used to amplify a
fragment (1103 bp) only present in the wild-type allele. A T2 plant homozygous for the deletion allele
and hemizygous for the *pKEULE-tdTomato-KEULE* transgene was isolated. A doubly homozygous line
was subsequently isolated in the T3 generation and used for experiments.
To generate double-labeled plants expressing markers for KEULE protein and cellulose synthase,
homozygous *pKEULE-tdTomato-KEULE/ keule* MM125 plants (ecotype Landsberg) were crossed with
homozygous *pCESA6-Citrine-CESA6/cesa6prc1-1* plants (ecotype Columbia[14]. An F2 line was selected

that was homozygous for both mutations and hemizygous for both markers. In the F3 generation, a
quadruply homozygous plant was isolated together with a plant homozygous for both mutations,
homozygous for *pCESA6-Citrine-CESA6* and hemizygous for *pKEULE-tdTomato-KEULE*. Seed from the
latter plant thus show segregation for the *keule* mutant phenotype.
To generate double-labeled plants expressing markers for KEULE protein and microtubules, *p35S-*
EGFP-TUA5 was introduced into *pKEULE-tdTomato-KEULE/keule*-MM125 plants via Agrobacterium-
mediated transformation. T2 seedlings expressing the microtubule label were used for experiments.

Growth
Seedlings were grown in the dark on vertical agar plates as described in Lindeboom et al.[65]. For
hypocotyl elongation assays, seedlings were grown for 5 days. For live cell imaging experiments,
seedlings were imaged 3 days after germination.

Drug treatments
Oryzalin, taxol and isoxaben were dissolved in DMSO for 100x stock solutions. Oryzalin was
used at a final concentration of 20 µM, taxol (Sigma) at 20 µM and isoxaben (Sigma) at 100 nM in plant
growth media (Hoagland's No. 2). Mannitol was dissolved in plant growth media directly at a
concentration of 200 mM. For control treatment (mock), DMSO was added to the growth media to a final
concentration of 0.01%. Incubation of the seedlings was performed in 6-well plates (WVR, nontreated)
with 2 mL of plant growth media in each well. The plates were wrapped in aluminum foil to block light
and incubated at 22 °C. The drug treatment durations were 16 hours for oryzalin, and 2 hours for all other
treatments. Likewise, the mock treatments were 16 hours for comparison to oryzalin, and 2 hours for
comparison to the remaining treatments.

Confocal microscopy
Imaging was performed on an instrument featuring a CSU-X1 spinning disk head (Yokogawa
Electric), a DMI6000 B inverted microscope (Leica), a 100X/1.4 NA oil immersion objective (Leica), an
Evolve 512 EMCCD camera (Photometrics) and a 1.2X telescope between the spinning disk unit and
camera. EGFP/CITRINE and tdTomato were excited at 488 nm and 561 nm respectively, by solid-state
lasers (Coherent Cube). Switching, shuttering and intensity of these lasers was performed by direct
electronic control. Emission filtering was accomplished with bandpass filters, (525/50 nm for EGFP and
CITRINE and 636/37 nm for tdTomato (SemRock), together with a 405/488/568 nm dichroic filter
(SemRock). Exposure times were 400 ms for tdTomato-KEULE, 200 ms for EGFP-TUA5 and 800 ms for
Citrine-CESA6. Images were acquired at 1 second intervals in experiments in which both tdTomato-
KEULE and EGFP-TUA5 were imaged and 2 second intervals in experiments featuring tdTomato-
KEULE together with Citrine-CESA6. All light energies were approximately 4 mW as measured at the
end of the optical fiber entering the confocal head. For high-speed acquisitions of tdTomato-KEULE,
image streaming was performed at an acquisition rate of 5 Hz (200 ms exposures). The focal plane in all
experiments was stabilized by hardware Adaptive Focus Control (Leica) to mitigate changes in specimen
intensity over time caused by thermal and mechanical drift in focus.
Photobleaching was performed by a Vector laser scanning unit (Intelligent Imaging Innovations)
integrated into the setup above at the right camera port. Focused laser light was pulsed and scanned (pulse
duration = 10 ms, raster block size = 20 units) on the image plane by a pair of galvanometer-driven
mirrors inside the Vector unit. Scanner positions were calibrated to image coordinates using an automated
procedure in Slidebook software (Intelligent Imaging Innovations). Citrine photobleaching was performed
using the 488 nm laser at the same power as used for imaging but focused on the image plane. Areas of
~200 µm^2 were bleached in 20-30 s.
All imaging was performed on the upper hypocotyl cells (1-2 mm below the apical hook).
Specimen mounting was performed as described previously[65].

Image registration

To correct for lateral specimen drift in the movies, we performed image registration in MATLAB.
We first performed a running average on the image sequence using a three-frame window (yielding an
image series 2 frames shorter than the original series). We then performed an intensity-based image
registration algorithm using the MATLAB function imregtform, using tformtype "translation" and
conserving the pixel size. To minimize interpolation artifacts, we first determined the transformation
matrix that yielded the best intensity match for each consecutive image pair, then calculated the
cumulative transformation matrix for the whole averaged image series and applied it to the original movie
that was not frame averaged, minus the first and last frame that were lost because of the frame averaging.
Some of the KEULE- tdTomato and Citrine-CESA colocalization movies (3 out of 8) still exhibited
sample drift after intensity-based registration. In those cases, we applied a feature-based method
employing the locations of detected KEULE spots (See Methods: Particle Tracking for details) as
registration features and using the MATLAB function estgeotform2d and a rigid transform.
613
<u>KEULE particle tracking, streaming experiments, image series with labeled microtubules</u>
Particle detection and tracking was performed on the registered movies. For streaming
experiments and experiments featuring labeled microtubules (1 second time intervals) we used µTrack[46]
to detect and estimate the positions of tdTomato-KEULE clusters in each frame and subsequently link
detected foci from frame to frame to track individual KEULE clusters over time. For each experiment
type, we used the following parameters:
620
621   <u>Streaming image series</u>.
622   For detection: initial psfSigma = 1.2 pixels, alpha for local max 0.05 and 10 max iterations for
623   psfSigma estimation.
624   For tracking: max detection gap = 2 frames, min track length = 4 frames, max search radius = 2
625   pixels, gap penalty = 4.
626
627   <u>1-second interval image series.</u>
628   For detection: initial psfSigma = 1.2 pixels, alpha for local max 0.05 and 10 max iterations for
629   psfSigma estimation.
630   For tracking: max detection gap = 1 frame, min track length = 1 frames, max search radius = 3
631   pixels, gap penalty = 4.
632
<u>KEULE and CESA particle tracking in dual labeled movies</u>
In experiments assessing KEULE localization and dynamics with respect to cellulose synthase
complex (CSC) trafficking and we used the Spots module in Imaris software (Bitplane) to detect and track
both tdTomato-KEULE clusters and Citrine-CESA6 particles. The Imaris Spots detector and filters
performed better for Citrine-CESA6 signals, in particular. CSCs as labeled by Citrine-CESA6 were
detected using a size value of 0.3 µm and quality value of 110. Tracking was performed by applying the
Autoregressive Motion algorithm (maximum distance = 0.35 µm, maximum gap size = 2 frames).
Successful deliveries of CSC complexes to the plasma membrane were determined by dynamic criteria as
described previously[7]. Briefly, a delivery event was defined when puncta that were mobile in the
streaming cytosol (1) stabilized positionally in the optical plane of the plasma membrane, and (2) showed
a transition to slow, steady and linear movement in the velocity range (~200-450 nm/min) we previously
have determined be characteristic of CSC's actively synthesizing and extruding cellulose fibers into the
cell wall[14]. To make this assessment, we first filtered CSC tracks by duration (>30 seconds), then used a
custom GUI in MATLAB to manually inspect and verify each Spots track and associated image data for
these characteristics. As a part of this process, the frame in which a tracked particle first showed evidence
of stabilized lateral position was marked as the "tethering" frame – the frame showing the time and
location of first stable interaction with the cell cortex of a presumed vesicle carrying one or more labeled
CSC's.

Clusters of tdTomato-KEULE were detected with Spots module using a size parameter value of
0.3 µm, a quality value of 100 and were tracked using the Connected Components algorithm (frame to
frame displacements of KEULE signal are small and not auto-correlated). These settings adequately
detected and tracked KEULE clusters based on visual inspection of the results.
655
<u>Maximum Likelihood Gaussian fitting for particles</u>
We employed maximum likelihood Gaussian fitting as described in Mortensen et al.[47] to use
photon counts in estimating the accuracy of particle position and shape. To determine the number of
photons collected in each image pixel, we first needed to calibrate our camera gain (photoelectrons per
grey value or ADU). Photon detection can be modeled as a Poisson process, which creates a distribution
in which the mean equals the variance. Therefore, the camera gain is equal to the coefficient between the
mean and variance of the gray values in the image. To calculate this coefficient, we first measured the
noise floor of the camera by taking the average of the gray values (intensity) of images acquired with the
camera capped. We then determined the grey value variance by subtracting one image from the other then
calculating the variance among these pixel-by-pixel difference values. To obtain the final variance, we
divided this number by 2 to account for the fact we are using two images for determining the differences.
Finally, we calculated the mean intensity of the images, subtracted the noise floor, and divided by the
variance to estimate the camera gain.
For each time point in every track, we cropped out a pixel grid with the estimated center of the
detected KEULE spot in the middle pixel (9 x 9 pixels). Along with a cropped image, the fitting routine
requires a set of initial parameters: xCoordinate, yCoordinate, point spread function standard deviation
(PSF SD), background, and an estimate of number of photons that comprise the imaged PSF. The x- and
y-coordinates we used are the estimates of position from the particle tracking described above. Based on
the approximate emission wavelength and objective numerical aperture, we used 133 nm for the initial
standard deviation of the PSF. We estimated the background by calculating the median intensity of the
outer border of the pixel grid used for fitting after subtraction of the camera noise floor. If the center of
the point source is in the middle of the pixel and the PSF SD is 133 nm, we expect 0.16 of the
photoelectrons to be in the center pixel. Therefore, we estimated the number of photoelectrons in the
imaged spot by first subtracting the noise floor and pixel background from the intensity in the center pixel
and multiplying the result by 1/0.16 (6.25). Before performing fitting using these initial data, we needed
to address noise that could have been introduced by bilinear interpolation during the image registration
that was employed to mitigate specimen drift. To accomplish this, we used the transformation matrix
from registration to map the locations of the tracked particles back to the locations in the original image
sequence, we then performed MLE fitting on the non-interpolated images by employing the method of
Mortenson et al.[47] in MATLAB.
Since we used free fit parameters and there is noise in the data from photon counting, free
fluorescently tagged proteins, high local particle densities and unresolved distributions of labeled
structures, we excluded distance measurement from fits that were outside the expected parameter ranges.
In the case of the SD of the 2D Gaussian, we found that there was a distinct population where the SD fit
parameter was < 50 nm. Since our pixels correspond to 133 nm, in these fits all the photoelectrons are
contained in one pixel, which is not consistent for a diffraction-limited spot at the magnification onto our
camera (roughly 4 pixels). Similarly, we found that fit SD values above 2x the estimated PSF SD in most
cases represented a fit containing more than one particle. Therefore, SD fit results with SD values outside
50 to 266 nm were excluded from position measurements. Fit results with estimated numbers of
photoelectrons > 1000 were also considered outliers and excluded. Finally, high local particle densities
the distance in some cases caused the MLEwG fit to center on a nearby particle and not the particle
marked by the starting position. We therefore also did not include position information for fits where the
distance between the original estimate and the MLEwG position differed more than 2 x 133 nm. See Fig.
S6 for fit parameters for the tdTomato-KEULE clusters and CIT-CESA6 labeled CSCs.
700
<u>KEULE to microtubule distance measurements</u>

To determine the position of each KEULE spot to assess the spatial relationship of KEULE
accumulation with the microtubule cytoskeleton we first determined the single frame with the highest
number of measured photoelectrons for each KEULE track. For tracks that were longer than 4 frames we
determined the max number of photoelectrons using a running three frames average, for shorter tracks we
did not use frame averaging. The rational for this selection was two-fold. First, we reasoned that
maximum signal might best be associated with the site of KEULE function, with initial signal rise being
indicative of KEULE recruitment and/or assembly into a membrane fusion structure, and signal loss
indicating KEULE disassociation or disassembly of the fusion machinery. Second, the precision of
estimated position is proportional to photon sampling. It is also relevant to note that frame-to-frame
movement of KEULE tracks were small (Fig. S7).
To locate positions of microtubule signal relative to each KEULE track location, we used a
rotating linear filter combined with analysis of signal peaks and a two-step fitting procedure. At the
coordinates of KEULE signal in each max value frame we drew a line 21 pixels long and 1 pixel wide
extending 10.5 pixels to each side of the KEULE position and repeated this process in increments of 9
degrees to cover 360 degrees (Fig. 4C). We then projected this array of lines on the microtubule channel
and extracted the intensities (in photons) along each line at 1-pixel (133 nm) intervals by bilinear
interpolation. To locate potential positions of microtubule signal along each sampling angle we used the
MATLAB function Findpeaks with a MinPeakProminence of 25 to identify signal peaks. The 10.5-pixel
distance is typically long enough to contain the peak intensity of the nearest microtubule. In fact, in many
cases this length intersects multiple microtubules, but it is also possible for the end of the lines to partially
intersect a microtubule with a peak outside the 10.5 pixels. For subsequent fitting steps, we included
pixels from the center until the furthest peak on each side plus 3 pixels to catch the shoulder of these
peaks, and at the same time, exclude signal from microtubules whose peaks are just outside the 10.5 pixel
range from influencing the fitting. We then estimated the center position of each signal peak by
modifying the 2D MLEwG[47] to a quasi-1D version, modeling the microtubule as a Gaussian wall[76] that is
one pixel wide, as shown in formula 1 below.
728
729   (Formula 1)
730

731
$$E = N\left(\frac{1}{\sqrt{2\pi\sigma^2}}\right)\left(\frac{1}{w}\right)\exp\left(-\frac{(x - \mu_x)^2}{2\sigma^2}\right) + b^2$$

732
Where:
E = expected number of photoelectrons per pixel
N = total number of photoelectrons
σ = Standard deviation
w = line width in nm
x = position on the line
μ_x = center position
b^2 = background photoelectrons per pixel
741
742   We found up to four peaks on a single line and fitted all peaks simultaneously by adding an
743   additional term describing each additional peak. Formula 2 describes the case for fitting the case for two
744   peaks:
745
746   (Formula 2)

747
$$E = N_{\text{first}}\left(\frac{1}{\sqrt{2\pi\sigma_{\text{first}}^2}}\right)\left(\frac{1}{w}\right)\exp\left(-\frac{(x - \mu_{x\text{ first}})^2}{2\sigma_{\text{first}}^2}\right) +$$

$$N_{\text{second}} \left(\frac{1}{\sqrt{2\pi\sigma_{\text{second}}^2}} \right) \left(\frac{1}{w} \right) \exp \left(-\frac{(x - \mu_{x\,\text{second}})^2}{2\sigma_{\text{second}}^2} \right) + b_{\text{global}}^2$$

751   For estimating the seeding parameters for the fit, we used the center of the maximum value pixel
752 for each peak identified by Findpeaks. We used 160 nm (median value of the peak population) as the
753 estimate for the standard deviation and we approximated the background by taking the square root of the
754 lowest 10 percentile of all the pixels for all angles after subtracting the noise floor.
755   After fitting all microtubule peaks along all angular steps, we determined the distance from the
756 KEULE location to the center of the nearest microtubule signal. We did this in three steps. First, we
757 determined the nearest microtubule signal peak to the KEULE location from among the initial angular
758 samples. Second, we determined the angle of the microtubule identified by the closest peak. Measuring
759 the distance from the KEULE location to the center of the microtubule signal is most accurate along a line
760 perpendicular from the KEULE location to the microtubule. By determining the local microtubule signal
761 angle, we could then determine the optimal perpendicular sample angle. To find the local orientation of
762 the microtubule we rotated a line of 21 pixels long around the center of the location where we initially
763 found the closest microtubule peak. We extracted the summed signal intensities along these lines and
764 determined the angle at which we found the maximum intensity, corresponding to the microtubule
765 angle[69]. Finally, once we established the microtubule angle, we used the angle perpendicular to that
766 microtubule angle to extract the nearest microtubule fit for the representative KEULE position. Distance
767 values where the fit parameters for the microtubule positions were the number of photoelectrons were <
768 50 or > 10000, as well as SD fit parameters < 50 or > 250 nm were excluded from distance
769 measurements. We required all the microtubule fits on the line to be within these parameters to be
770 included in the analysis. See Fig. S8A-C for fit parameters for KEULE (A) and microtubule signal (B) fits
771 after selection as well as the standard error of the distance measurement between them (C).

773 <u>Analysis of KEULE localization with CESA6 delivery sites</u>
774   KEULE and CESA6 spots were detected and tracked in Imaris as described above and maximum
775 likelihood 2D Gaussian fitting was applied to each set of tracks. CESA tracks tend to be relatively long
776 and can be confounded by a dynamic background produced by highly motile CESA containing organelles
777 including Golgi bodies and Trans-Golgi network compartments. This background makes fully automated
778 detection and quantitation of CSC delivery events challenging. We therefore designed a graphical user
779 interface in MATLAB that allowed us to efficiently curate potential CESA delivery events identified in
780 Imaris and verify whether individual tracks were indeed depicting a CESA delivery. We have shown
781 previously that CESA delivery to the plasma membrane typically shows three distinct phases; (1) erratic
782 movement of CESA containing vesicles as they appear in the optical plane at the cell cortex, (2)
783 positional-stabilization and (3) slow and steady movement with typically linear trajectories. We speculate
784 that the second phase represents molecular tethering at the plasma membrane and the events that lead to
785 vesicle fusion and cargo delivery. The third phase is behavior that we have shown is characteristic of
786 active CSC complexes in the plasma membrane as they synthesize cellulose[14]. Thus, observation of this
787 latter behavior indicates successful delivery of CESA cargo to the plasma membrane. For each CESA
788 track that showed these three distinct phases, we marked the first frame of CESA stabilization at the
789 plasma membrane. From this position, we determined the distance to the nearest KEULE spot in a 30-
790 frame time range (60 seconds). Some degree of coincidence between CESA deliveries and KEULE spots
791 is expected by chance alone, especially since the density of KEULE clusters at the plasma membrane is
792 relatively high. To evaluate whether the observed distribution of KEULE locations with sites of CESA
793 stabilization might be due to chance alone, we generated random positions in the same regions of interest
794 used for the experimental data and calculated the distances to the nearest KEULE spot. We used these
795 simulated data to model the expected distribution of distances and compared this distribution to the

796 distribution of distances found for confirmed CESA delivery events. First, we divided both the
797 experimental and simulated distance distributions up into 40 nm distance bins and calculated the bin-by-
798 bin count frequencies by the total number of simulated counts. We then introduced a bias parameter δ that
799 equates an observed frequency in each distance bin to the expected value from the simulated distribution
800 as shown in formula (4).

801
802   (Formula 4)

803
$$P_{inDistBin}(\delta) = \frac{\text{Expected}_{inDistBin}\,\delta}{\text{Expected}_{\text{inDistBin}}\,\delta + \text{Expected}_{\text{outsideDistBin}}(1-\delta)}$$

804
The value of δ may vary between 0 and 1. For a possible observed frequency that equals the
expected frequency, δ is equal 0.5. If δ is greater than 0.5, an observed frequency is higher than expected.
Likewise, if δ is less than 0.5 the observed frequency lower than expected under the null hypothesis. Use
of the parameter allows us to use a Bayesian approach to test multiple hypotheses to ask what bias value
best may best explain the actual observed frequencies. To do this, we calculated the Likelihood Ratio of
observing an experimental result given a range of values for δ (in increments of 0.001, excluding 0) for
each frequency bin:

812
813   (Formula 5)

814
$$\frac{L(\delta)}{L\frac{1}{2}} = \frac{P_{inDistBin}(\delta)^M\,(1 - P_{inDistBin}(\delta)^{N-M})}{Expected^N_{inDistBin}\,Expected^{N-M}_{outsideDistBin}}$$

815 Where:
816 $L(\delta)$ = likelihood of the data given the bias δ,
817 $L\frac{1}{2}$ = likelihood of the data in the neutral case (no positive or negative bias)
818 N = total number of measurements
819 M = number of measurements in this specific distance bin

820
821   We then found the maximum a posteriori probability (MAP) for δ and computed the shortest
822 range of δ values for the highest posterior density (HPD) that contains 95% of the normalized probability
823 density. Maximum likelihood estimates for δ below 0.5 indicate underrepresentation at the distance of
824 interest and above 0.5 indicate overrepresentation of experimental data compared to the unbiased case.

825
826 Analysis of KEULE localization with microtubules

827
828   We used the same Bayesian approach describe above for analyzing the localization of KEULE
829 clusters with respect to the estimated center of the nearest microtubule. We changed the distance bin size
830 from 40 to 20 nm.

831
Hot spot analysis
To investigate if some KEULE events are independent of each other each or not (e.g. if "hot
spots" for KEULE accumulation may exist), we examined colocalization between sequential KEULE
spots in time series collected from cells expressing both labeled KEULE and tubulin proteins. Within a
region of interest in each time series, we selected all KEULE tracks that were first detected between
frames 2 and 31 and calculated the distance to all other KEULE tracks that started subsequently up to and
including frame 301. For position estimates we used the frame in the track that showed the highest
number of fitted photoelectrons. To compare the degree of colocalization of the initial KEULE tracks
with future KEULE tracks to that expected from chance alone, we created a simulated initial dataset with
random locations within the same region of interest and calculated the distances to all KEULE tracks in
the same time range.

Since KEULE foci are positionally stable (Fig. S7) we chose a stringent distance threshold of 133

nm (1 pixel) to declare two spots or locations to be co-localized. Because of the relatively high density of

KEULE spots at the plasma membrane, colocalization is expected at some point in time, therefore the

time frame of colocalization analysis is also important. Indeed, we observe a large degree of

colocalization in the whole dataset for both the experimental and simulated data, but with very distinct

waiting time distributions (Fig. S5D). Based on the differences in the distribution of waiting times

between random simulated positions and KEULE positions until the next KEULE spot arrives, we

determined that a 40 second time delay window captures most biologically relevant colocalizations while

excluding most random colocalizations due to KEULE density and chance.

Bibliography

1. Noordstra, I. & Akhmanova, A. Linking cortical microtubule attachment and exocytosis. *F1000Research* **6**, 469 (2017).

2. Spiliotis, E. T. & Nelson, W. J. Spatial control of exocytosis. *Curr. Opin. Cell Biol* **15**, 430–437 (2003).

3. Fourrière, L., Jimenez, A. J., Perez, F. & Boncompain, G. The role of microtubules in protein transport. *Journal of Cell Science* **133**, 237016 (2020).

4. Chang, F. & Martin, S. G. Shaping Fission Yeast with Microtubules. *Cold Spring Harb. Perspect. Biol* **1**, 001347 (2009).

5. Bartolini, F. & Gundersen, G. G. Generation of noncentrosomal microtubule arrays. *J. Cell Sci* **119**, 4155–4163 (2006).

6. Shaw, S. L., Kamyar, R. & Ehrhardt, D. W. Sustained Microtubule Treadmilling in Arabidopsis Cortical Arrays. *Science* **300**, 1715–1718 (2003).

7. Gutierrez, R., Lindeboom, J. J., Paredez, A. R., Emons, A. M. C. & Ehrhardt, D. W. Arabidopsis cortical microtubules position cellulose synthase delivery to the plasma membrane and interact with cellulose synthase trafficking compartments. *Nat. Cell Biol* **11**, 797–806 (2009).

8. Sampathkumar, A. *et al.* Patterning and Lifetime of Plasma Membrane-Localized Cellulose Synthase Is Dependent on Actin Organization in Arabidopsis Interphase Cells. *Plant Physiol* **162**, 675–688 (2013).

9. Vellosillo, T., Dinneny, J. R., Somerville, C. R. & Ehrhardt, D. W. TRANVIA (TVA) facilitates cellulose synthase trafficking and delivery to the plasma membrane. *Proc. Natl. Acad. Sci* **118**, 2021790118 (2021).

10. Li, S., Lei, L., Somerville, C. R. & Gu, Y. Cellulose synthase interactive protein 1 (CSI1) links microtubules and cellulose synthase complexes. *Proc. Natl. Acad. Sci* **109**, 185–190 (2012).

11. Bringmann, M. *et al.* POM-POM2/CELLULOSE SYNTHASE INTERACTING1 Is Essential for the Functional Association of Cellulose Synthase and Microtubules in Arabidopsis. *Plant Cell* **24**, 163–177 (2012).

12. Zhu, X., Li, S., Pan, S., Xin, X. & Gu, Y. CSI1, PATROL1, and exocyst complex cooperate in delivery of cellulose synthase complexes to the plasma membrane. *Proc. Natl. Acad. Sci* **115**, 3578–3587 (2018).

13. Lei, L. *et al.* CELLULOSE SYNTHASE INTERACTIVE1 Is Required for Fast Recycling of Cellulose Synthase Complexes to the Plasma Membrane in Arabidopsis. *The Plant Cell* **27**, 2926–2940 (2015).

14. Paredez, A. R., Somerville, C. R. & Ehrhardt, D. W. Visualization of Cellulose Synthase Demonstrates Functional Association with Microtubules. *Science* **312**, 1491–1495 (2006).

15. DeBolt, S. *et al.* Morlin, an inhibitor of cortical microtubule dynamics and cellulose synthase movement. *Proc. Natl. Acad. Sci* **104**, 5854–5859 (2007).

16. Emons, A. M. C., Mulder, B. M., Emons, A. M. C. & Mulder, B. M. How the deposition of cellulose microfibrils builds cell wall architecture. *Trends Plant Sci* **5**, 35–40 (2000).

17. Hamant, O. *et al.* Developmental Patterning by Mechanical Signals in Arabidopsis. *Science* **322**, 1650–1655 (2008).

18. Zhang, B., Karnik, R., Alvim, J., Donald, N. & Blatt, M. R. Dual Sites for SEC11 on the SNARE SYP121 Implicate a Binding Exchange during Traffic1. *Plant Physiol* **180**, 228–239 (2019).

19. Honsbein, A. *et al.* A tripartite SNARE-K+ channel complex mediates in channel-dependent K+ nutrition in Arabidopsis. *Plant Cell* **21**, 2859–2877 (2009).

20. Assaad, F. F. *et al.* The PEN1 Syntaxin Defines a Novel Cellular Compartment upon Fungal Attack and Is Required for the Timely Assembly of Papillae. *Mol. Biol. Cell* **15**, 5118–5129 (2004).

21. Bloch, D. *et al.* Exocyst SEC3 and Phosphoinositides Define Sites of Exocytosis in Pollen Tube Initiation and Growth. *Plant Physiol* **172**, 980–1002 (2016).

22. Fendrych, M. *et al.* Visualization of the exocyst complex dynamics at the plasma membrane of Arabidopsis
thaliana. *MBoC* **24**, 510–520 (2013).

23. Barg, S., Knowles, M. K., Chen, X., Midorikawa, M. & Almers, W. Syntaxin clusters assemble reversibly at
sites of granules in live cells. in *Proceedings of the National Academy of Sciences* vol. 107 20804–20809
(2010).

24. Knowles, M. K. *et al.* Single granules of live cells recruit syntaxin-1 and synaptosomal associated protein 25
(SNAP-25) in large copy numbers. in *Proceedings of the National Academy of Sciences* vol. 107 20810–20815
(2010).

25. McNew, J. A. *et al.* Close Is Not Enough. *The Journal of Cell Biology* **150**, 105–118 (2000).

26. Südhof, T. C. & Rothman, J. E. Membrane Fusion: Grappling with SNARE and SM Proteins. *Science* **323**,
474–477 (2009).

27. Burgoyne, R. D. *et al.* The Functions of Munc18-1 in Regulated Exocytosis. *Ann. N. York Acad. Sci* **1152**, 76–
86 (2009).

28. Baker, R. W. *et al.* A direct role for the Sec1/Munc18-family protein Vps33 as a template for SNARE
assembly. *Science* **349**, 1111–1114 (2015).

29. Stepien, K. P., Prinslow, E. A. & Rizo, J. Munc18-1 is crucial to overcome the inhibition of synaptic vesicle
fusion by αSNAP. *Nat. Commun* **10**, 4326 (2019).

30. Peng, R.-W., Guetg, C., Abellan, E. & Fussenegger, M. Munc18b regulates core SNARE complex assembly
and constitutive exocytosis by interacting with the N-peptide and the closed-conformation C-terminus of
syntaxin 3. *Biochem. J* **431**, 353–361 (2010).

31. Hashizume, K., Cheng, Y.-S., Hutton, J. L., Chiu, C. & Carr, C. M. Yeast Sec1p Functions before and after
Vesicle Docking. *Mol. Biol. Cell* **20**, 4673–4685 (2009).

32. Mayer, U., Ruiz, R. A. T., Berleth, T., Miséra, S. & Jürgens, G. Mutations affecting body organization in the
Arabidopsis embryo. *Nature* **353**, 402–407 (1991).

33. Assaad, F. F., Mayer, U., Wanner, G. & Jürgens, G. The KEULE gene is involved in cytokinesis in
Arabidopsis. *Mol. Gen. Genet. MGG* **253**, 267–277 (1996).

34. Karnik, R. *et al.* Arabidopsis Sec1/Munc18 protein SEC11 is a competitive and dynamic modulator of SNARE
binding and SYP121-dependent vesicle traffic. *Plant Cell* **25**, 1368–1382 (2013).

35. Karnik, R. *et al.* Binding of SEC11 indicates its role in SNARE recycling after vesicle fusion and identifies two
pathways for vesicular traffic to the plasma membrane. *Plant Cell* **27**, 675–694 (2015).

36. Karnahl, M. *et al.* Functional diversification of Arabidopsis SEC1-related SM proteins in cytokinetic and
secretory membrane fusion. *Proceedings of the National Academy of Sciences* **115**, 6309–6314 (2018).

37. Assaad, F. F., Huet, Y., Mayer, U. & Jürgens, G. The Cytokinesis Gene KEULE Encodes a Sec1 Protein That
Binds the Syntaxin Knolle. *The Journal of Cell Biology* **152**, 531–544 (2001).

38. Carr, C. M., Grote, E., Munson, M., Hughson, F. M. & Novick, P. J. Sec1p Binds to SNARE Complexes and
Concentrates at Sites of Secretion. *J. Cell Biol* **146**, 333–344 (1999).

39. Scott, B. L. *et al.* Sec1p directly stimulates SNARE-mediated membrane fusion in vitro. *J. Cell Biol* **167**, 75–85
(2004).

40. Zilly, F. E., Sørensen, J. B., Jahn, R. & Lang, T. Munc18-Bound Syntaxin Readily Forms SNARE Complexes
with Synaptobrevin in Native Plasma Membranes. *PLoS Biol* **4**, 330 (2006).

41. Medine, C. N., Rickman, C., Chamberlain, L. H. & Duncan, R. R. Munc18-1 prevents the formation of ectopic
SNARE complexes in living cells. *J. Cell Sci* **120**, 4407–4415 (2007).

42. Rickman C. & Duncan R. R. Munc18/Syntaxin Interaction Kinetics Control Vesicle Dynamics2. *J. Biol. Chem*
285, 3965–3972 (2010).

43. Smyth, A. M., Rickman, C. & Duncan, R. R. Vesicle Fusion Probability Is Determined by the Specific
Interactions of Munc18*. *J. Biol. Chem* **285**, 38141–38148 (2010).

44. Trexler, A. J., Sochacki, K. A. & Taraska, J. W. Imaging the recruitment and loss of proteins and lipids at
single sites of calcium-triggered exocytosis. *Mol. Biol. Cell* **27**, 2423–2434 (2016).

45. Somasundaram, A. & Taraska, J. W. Local protein dynamics during microvesicle exocytosis in neuroendocrine
cells. *Mol. Biol. Cell* **29**, 1891–1903 (2018).

46. Jaqaman, K. *et al.* Robust single-particle tracking in live-cell time-lapse sequences. *Nat. Methods* **5**, 695–702
(2008).

47. Mortensen, K. I., Churchman, L. S., Spudich, J. A. & Flyvbjerg, H. Optimized localization analysis for single-
molecule tracking and super-resolution microscopy. *Nat. Methods* **7**, 377–381 (2010).

48. Cho, W. J., Jeremic, A. & Jena, B. P. Size of Supramolecular SNARE Complex: Membrane-Directed Self-
Assembly. *J. Am. Chem. Soc* **127**, 10156–10157 (2005).

49. Cho, W. J. *et al.* Membrane-directed molecular assembly of the neuronal SNARE complex. *J. Cell. Mol. Med*
15, 31–37 (2011).

50. Haigler, C. H. & Brown, R. M. Transport of rosettes from the golgi apparatus to the plasma membrane in
isolated mesophyll cells ofZinnia elegans during differentiation to tracheary elements in suspension culture.
Protoplasma **134**, 111–120 (1986).

51. Zhang, Y. *et al.* Golgi-localized S^TEL^LO proteins regulate the assembly and trafficking of cellulose synthase
complexes in Arabidopsis. *Nat. Commun* **7**, 11656 (2016).

52. Scheible, W.-R., Eshed, R., Richmond, T., Delmer, D. & Somerville, C. Modifications of cellulose synthase
confer resistance to isoxaben and thiazolidinone herbicides in Arabidopsis Ixr1 mutants. in *Proceedings of the*
National Academy of Sciences vol. 98 10079–10084 (2001).

53. Zhou, Q. *et al.* Architecture of the synaptotagmin–SNARE machinery for neuronal exocytosis. *Nature* **525**, 62–
67 (2015).

54. Brunger, A. T. Structure and function of SNARE and SNARE-interacting proteins. *Q Rev Biophys* **38**, 1–47
(2005).

55. Rothman, J. E., Krishnakumar, S. S., Grushin, K. & Pincet, F. Hypothesis – buttressed rings assemble, clamp,
and release SNAREpins for synaptic transmission. *FEBS Lett* **591**, 3459–3480 (2017).

56. Giddings, T. H. & Staehelin, L. A. Spatial relationship between microtubules and plasma-membrane rosettes
during the deposition of primary wall microfibrils in Closterium sp. *Planta* **173**, 22–30 (1988).

57. Lindeboom, J. J. *et al.* CLASP stabilization of plus ends created by severing promotes microtubule creation and
reorientation. *J. Cell Biol* **218**, 190–205 (2019).

58. Crowell, E. F. *et al.* Pausing of Golgi Bodies on Microtubules Regulates Secretion of Cellulose Synthase
Complexes in Arabidopsis. *Plant Cell* **21**, 1141–1154 (2009).

59. Martinière, A., Gayral, P., Hawes, C. & Runions, J. Building bridges: formin1 of Arabidopsis forms a
connection between the cell wall and the actin cytoskeleton. *Plant J* **66**, 354–365 (2011).

60. Geelen, D. *et al.* The abscisic acid-related SNARE homolog NtSyr1 contributes to secretion and growth:
evidence from competition with its cytosolic domain. *Plant Cell* **14**, 387–406 (2002).

61. Grefen, C. *et al.* A novel motif essential for SNARE interaction with the K(+) channel KC1 and channel gating
in Arabidopsis. *Plant Cell* **22**, 3076–3092 (2010).

62. Hoffmann, N., King, S., Samuels, A. L. & McFarlane, H. E. Subcellular coordination of plant cell wall
synthesis. *Dev. Cell* **56**, 933–948 (2021).

63. Cosgrove, D. J. Plant expansins: diversity and interactions with plant cell walls. *Curr. Opin. Plant Biol* **25**,
162–172 (2015).

64. Baskin, T. I. ANISOTROPIC EXPANSION OF THE PLANT CELL WALL. *Annu. Rev. Cell Dev. Biol* **21**, 203–222 (2005).

65. Lindeboom, J. J. *et al.* A Mechanism for Reorientation of Cortical Microtubule Arrays Driven by Microtubule Severing. *Science* **342**, 1245533 (2013).

66. Baas, P. W., Rao, A. N., Matamoros, A. J. & Leo, L. Stability properties of neuronal microtubules. *Cytoskeleton* **73**, 442–460 (2016).

67. Tojima, T., Itofusa, R. & Kamiguchi, H. Steering Neuronal Growth Cones by Shifting the Imbalance between Exocytosis and Endocytosis. *The Journal of Neuroscience* **34**, 7165–7178 (2014).

68. Brown, A.-L. *et al.* TDP-43 loss and ALS-risk SNPs drive mis-splicing and depletion of UNC13A. *Nature* **603**, 131–137 (2022).

69. Ma, X. R. *et al.* TDP-43 represses cryptic exon inclusion in the FTD–ALS gene UNC13A. *Nature* **603**, 124–130 (2022).

70. Liu, X. & Henty-Ridilla, J. L. Multiple roles for the cytoskeleton in ALS. *Exp. Neurol* **355**, 114143 (2022).

71. Zhang, S. *et al.* Genome-wide identification of the genetic basis of amyotrophic lateral sclerosis. *Neuron* **110**, 992-1008 11 (2022).

72. Curtis, M. D. & Grossniklaus, U. A Gateway Cloning Vector Set for High-Throughput Functional Analysis of Genes in Planta. *Plant Physiol* **133**, 462–469 (2003).

73. Shaner, N. C. *et al.* Improved monomeric red, orange and yellow fluorescent proteins derived from Discosoma sp. red fluorescent protein. *Nat. Biotechnol* **22**, 1567–1572 (2004).

74. Clough, S. J. & Bent, A. F. Floral dip: a simplified method for Agrobacterium -mediated transformation of Arabidopsis thaliana. *Plant J* **16**, 735–743 (1998).

75. Logemann, E., Birkenbihl, R. P., Ülker, B. & Somssich, I. E. An improved method for preparing Agrobacterium cells that simplifies the Arabidopsis transformation protocol. *Plant Methods* **2**, 16 (2006).

76. Ruhnow, F., Zwicker, D. & Diez, S. Tracking Single Particles and Elongated Filaments with Nanometer Precision. *Biophys. J* **100**, 2820–2828 (2011).

Acknowledgments: We thank Michael Davidson for the kind gift of the tdTomato construct. We thank Zdeněk Lánský, Johnathan Cooper-Knock and Masayoshi Nakamura for advice and insightful comments.

Funding: National Institutes of Health grant R15GM137247-01 (VK), National Institutes of Health grant R01GM123259-01 (DWE)

Author contributions:

Conceptualization: JJL, RG, VK, DWE

Methodology: JJL, RG, VK, DWE

Investigation: JJL, RG, VK, DWE

Visualization: JJL, RG, DWE

Funding acquisition: VK, DWE

Writing – original draft: JJL, RG, DWE

Writing – review & editing: JJL, RG, VK, DWE

1024 **Competing interests:** The authors declare no competing interests.

1025 **Data and materials availability:** Biological material, raw data, processed data and MATLAB scripts
1026 used for the analysis are available upon request.

Figures

Figure 1

Fig. 1. KEULE assembly dynamics at the plasma membrane. (**A**) Distribution of tdTomato-KEULE signal at the plasma membrane of etiolated hypocotyl cells in *Arabidopsis*. Arrowheads point out individual clusters of labeled KEULE protein. (**B**) Kymograph of the tdTomato-KEULE signal intensity over time as sampled over the transect line shown in (**A**) (blue dashes). The yellow brackets highlight several individual tdTomato-KEULE clusters. (**C**) Example of tdTomato-KEULE cluster intensity over time. (**D**) Histogram of KEULE cluster lifetimes in 200 ms bins. n = 15,130 clusters. (**E**) Average tdTomato-KEULE cluster signal intensity with a lifetime of 4.2, 4.4 and 4.6 s. Note the initial rapid increase in signal over the first acquisition step (darker green), the more gradual, near linear, ramp up (green) to a plateau (yellow), followed by abrupt loss to non-detection (red). n = 231, 213,182 clusters for 4.2, 4.4 4.8 s, respectively. (**F**) Average tdTomato-KEULE cluster signal intensity with lifetimes ranging from 0.8 to 6.8 s. n = 15,397 clusters in total, ranging from 4992 clusters for 0.8 s to 122 for 6.8 s. (**G**) Average signal intensities of tdTomato-KEULE clusters in three cumulative lifetime ranges, 0.8 to 1.6 s, 1.8 to 6.2 s and 6.4+ s. The inset shows the lifetime distribution histogram of tdTomato-KEULE cluster lifetimes with the lifetime ranges color coded. n = 15,130 clusters. All data in panels D-G acquired from 12 cells from 12 seedlings. Error bars are S.E. Scale bars are 2 μm.

Figure 2

Fig. 2. KEULE is spatially associated with and required for CSC delivery to the plasma membrane.
(**A**) CESA exocytosis event. (Top panel) Time series (top panel) and kymograph (bottom panel) of
tdTomato-KEULE and CIT-CESA6 during a CESA exocytosis event. The kymograph shows evidence for
CSC-vesicle tethering at the plasma membrane (start of static phase) and successful CSC delivery to the
plasma membrane as an active complex (steady movement phase, see text). (**B**) Distribution of distances
between the positions of CIT-CESA6 signals at start of static phase and the nearest tdTomato-KEULE
cluster, as estimated by MLE PSF fitting. Data shown both for observed CIT-CESA6 locations and
simulated random locations. (**C**) Differences of the observed vs expected CESA6-KEULE distance
frequencies in (**B**), portraying over- and under-representation (green and red respectively). (**D**) Examples
of colocalization and fitted positions for events in (**B**). (**E**) Bayesian likelihood ratios of the observed
frequency of distances between 0 and 40 nm given bias δ over $\delta(0.5)$. The position of the peak value
indicates the most likely, or maximum a posteriori probability (MAP), value for the bias δ for the
observed frequency given the expected frequency. (**F**) Bayesian analysis of the MAP(δ) for each distance
bin, MAP(δ) = 0.5 indicates no difference from expected. Error bars are 95% confidence intervals
expressed as the highest posterior density (HPD, the shortest interval containing 95% of the probability
density). (**G**) CESA6 delivery rates in wild type and *keule* mutant cells. Whiskers are quartiles. ****
p<0.0001 Wilcoxon 2-sample test. In panels B-F, n = 194 CESA exocytosis events in 8 cells from 8
seedlings for observed data. For simulated data, n = 45051 randomized CESA exocytosis locations in the
same KEULE channel data sets. In panel G, n = 8 cells from 8 seedlings for WT and 16 cells from 16
plants for *keule*. Scale bars are 1 μm.

Figure 3

1069

Fig. 3. KEULE cluster appearance is temporally associated with CSC appearance at the plasma
membrane. (**A**) Comparison of the waiting time distribution from CSC signal stabilization until the
detection of the nearest tdTomato-KEULE cluster in observed and simulated data. n = 194 CSC delivery
events in 8 cells from 8 plants. (**B**) KEULE dynamics in the context of CSC delivery to the plasma
membrane. Histograms of waiting times (left pane) and lifetimes of tdTomato-KEULE clusters co-
localized with CSC delivery events (right pane). n = 161 CSC delivery events in 8 cells from 8 plants for
observed data, n = 45,051 simulated CSC delivery events applied to the same data sets. (**C**) KEULE
cluster durations for all events and those associated with CSC delivery, compared by relative frequency of
durations over 2 seconds. n = 13766 for all KEULE clusters, n = 180 for KEULE clusters associated with
CSC delivery. Whiskers are exact binomial 95% confidence intervals, p = 6.1E-09 Fishers' exact test.

Figure 4

1081
Fig. 4. Microtubule dependent distribution of exocytosis events. (**A**) Images of GFP-TUA5 and
tdTomato-KEULE signal in epidermal cells. Arrowheads indicate examples of apparent co-localization of
signal. (**B**) Spatial patterning of tdTomato-KEULE clusters over the range of 36.8 seconds. Crosses show
positions of MLEwG fits. (**C**) Steps in the routine employed to calculation the distance from a KEULE
cluster to the center of the nearest microtubule distance. (**D**) Steps in quasi-1-dimensional MLEwG fitting
of microtubule signal along the most-orthogonal scan line of a rotating filter. (**E**) Overlay of a
microtubule channel image and pairs of markers for MLE fit locations of KEULE clusters and the center
of the nearest microtubule in observed data (left) and for simulated random KEULE locations (right).
KEULE cluster detects from -5 to + 5 frames from the microtubule image are shown. Scale bars are 2 μm.

Figure 5

1092

Fig. 5. Bayesian analysis of microtubule bias for exocytosis events. (**A**) Comparison of the
distributions of KEULE-nearest microtubule distances for observed and simulated data. n = 115,963
KEULE clusters in 14 cells from 14 plants. (**B**) Differences of the observed vs expected KEULE-
microtubule distance frequencies shown in (**A**), portraying the pattern of over- and under-representation
(green and red respectively). (**C**) Bayesian likelihood ratios of the observed frequency of KEULE-
microtubule distances between 0 and 20 nm given bias δ over δ(0.5). (**D**) Bayesian analysis of the
MAP(δ) of each distance bin. Data acquired from 14 cells in 14 seedlings. n = 115,963 KEULE clusters
in 14 cells from 14 plants. Error bars are 95% confidence intervals expressed as HPD.

Figure 6

Polarized tansport to plus end locations   Templated patterning

A

— Microtubule
Golgi body
● Secretory vesicle
Kinesin
Targeted domain
Focal adhesions
Adherens junction

B

Membrane domain model   Vesicle-MT interaction model

400 nm
200 nm
100 nm
70 nm

max +

Bias of observed vs
expected frequency

max -

— Plasma membrane
· Microtubule
Plasma membrane linker
Vesicle linker
● KEULE/SNAREs

Microtubule-directed
vesicle

200   500 nm

1400 nm

95% CI

Fig. 6. Patterning of exocytosis by microtubule arrays. (A) Schematic diagrams of microtubule-
directed patterning of exocytosis. The three cells on the left show targeting of exocytosis by motor-driven
transport of vesicles along polarized microtubule tracks to locations defined by their plus ends[3] .
Clockwise from upper left: migrating fibroblast, epithelial cell, neuronal cell. By contrast, the diagram at
far right shows an epidermal cell in the plant axis, where a 2-D array of microtubules at the cell cortex
acts as a template that directs exocytosis in a complex but ordered pattern. The organization of the
microtubule lattices defines this pattern. **(B)** Alternative models for spatial patterning of exocytosis by
cortical microtubule arrays in plant cells. All structures are abstracted but depicted at approximate scale.
Microtubules (grey) are shown spaced at the mean spacing that has been measured in the cortical arrays
of growing cells in the hypocotyl epidermis of dark grown *Arabidopsis* seedings[61]. The likelihood of
exocytosis marked by dynamic KEULE clusters is shown as a red-green heat map as scaled from the
observed over-representation and under-representation of KEULE clusters compared to that expected
from the null hypothesis of a random distribution (depicted in Fig. 5B). The two sides of the diagram
depict alternative models to explain the measured spatial dimensions of the domains enriched in
exocytosis. On the right, the measured extent of the enriched domain is determined by the interaction
exocytotic vesicle with microtubules, where those vesicles have a wide range of sizes. On the left,
microtubules establish a membrane domain that extends approximately 180 nm on either side where
proteins needed for vesicle capture or exocytosis are concentrated. These models are not mutually
exclusive. The grey-scale bar at lower right depicts the mean 95% confidence interval for the measured
distances of KEULE clusters to microtubule centers.
1123

Supplemental Materials

1125   **Figure S1**

A

1126

Fig. S1. Complementation of the *keule* mutation. (A) Seedlings of indicated genotypes were grown in
the dark for 5 d. The recessive *keule* MM125 allele is seedling lethal; mutant seedlings have bloated,
multinucleate cells. Expression of a tdTomato-tagged version of KEULE, expressed under the control of
the native upstream regulatory element, complemented the *keule* mutation in dark-grown seedlings. Scale
bar, 5 mm. **(B)** Hypocotyl elongation in 5-day-old dark-grown seedlings. Rescued seedlings were
comparable to wild-type ($n = 50$ plants per genotype, $P = 0.72$ by *t*-test). Error bars represent SEM.

Figure S2

1134

Fig. S2. CSC delivery rates. CSC delivery rates to the plasma membrane of cells expressing tdTomato-
KEULE and CIT-CESA6 after treatment with 0.01% DMSO (mock), 200 mM mannitol or 100 nM
isoxaben. See Methods for treatment durations. Values are means, whiskers are quartiles. Mock: n = 8
cells from 8 seedlings, 208 delivery events observed in 1570 μm^2 total area over 0.0828 hr. Mannitol: n =
15 cells from 15 seedlings, 11 delivery events observed in 880 μm^2 total area over 0.0793 hr. Isoxaben: n
= 12 cells from 12 seedlings, 2 delivery events observed in 996 μm^2 total area over 0.0793 hr. ****
p<0.0001, Wilcoxon two sample test.

Figure S3

Intensity rising phase grouped by lifetime class

1143
1144

Fig. S3. Intensity of dynamics of tdTomato-KEULE clusters by lifetime. (A) Average tdTomato-
KEULE cluster signal intensity with lifetimes ranging from 0.8 to 6.8 s for mock, mannitol treatment. and
isoxaben treatment. Whiskers are means and quartiles. n.s. p>0.05 Wilcoxon two sample test. (**B** and **C**)
Comparison of average signal intensities of tdTomato-KEULE clusters in three cumulative lifetime
ranges, 0.8 to 1.6 s, 1.8 to 6.2 s and 6.4 to 12.2 s, between (**B**) mock vs mannitol treatment, and (**C**) mock
vs isoxaben treatment. Mock: n = 15,130 clusters observed in 12 cells from 12 plants. Mannitol: n =
35,496 clusters observed in 15 cells from 15 plants. Isoxaben: n = 25,251 clusters observed in 12 cells
from 12 plants. Error bars are SEM.

1153

1154   **Figure S4**

1155

Fig. S4. KEULE cluster lifetimes. (**A** to **C**) Lifetime distribution of tdTomato-KEULE clusters in mock
(**A**), mannitol (**B**) and isoxaben (**C**) treated cells. (**D** and **E**) Difference in relative lifetime frequency
between mannitol and mock (**D**), and isoxaben and mock (**E**). Vertical lines indicate the lifetime classes
in Figure 1G. (**F**) Formation rates of new tdTomato-KEULE clusters. Values are means. (**G** and **H**)
Comparison of the fraction of remaining tdTomato-KEULE clusters that were present in the first frame of
the movie for mannitol to mock (**G**) and isoxaben to mock. Mock: n = 763 clusters observed in 12 cells
from 12 plants. Mannitol: n = 1149 clusters observed in 15 cells from 15 plants. Isoxaben: n = 724
clusters observed in 12 cells from 12 plants.

1164

1165   **Figure S5**

1166

1167

1168

Fig. S5 KEULE cluster hot spots. (**A**) Map of tdTomato-KEULE cluster detections over x minutes.
Crosses are positions determined by MLEwG fitting. Arrowheads indicate examples of repeat detects at
the same location, or hot spots. (**B**) Kymograph of tdTomato-KEULE signal along a 3 pixel wide transect
line. Yellow brackets indicate examples of repeated KEULE cluster formation at the same location. (**C**)
Individual image frames at a hot spot, 200 msec intervals over 20 seconds. (**D**) Distributions of waiting
times for a second cluster to appear at the same location as a previous cluster, as defined by a stringent
co-localization cutoff distance of 133 nanometers. The distributions for observed data and randomly
simulated data are shown in purple and grey, respectively. Note that the observed intervals are
significantly skewed to shorter intervals as compared to those in the simulated data (p<3.27E-8
Kolmogorov-Smirnov test). n = 12,719 clusters in 17 cells from 17 plants. (**E**) Distributions of the
number of co-localized events observed at each KEULE cluster location within 60 seconds of the
formation of the first cluster. Colocalization events were 3.36 more frequent in observed data (green) than
expected by chance alone (grey), as determined by odds ratio analysis (see panel (**G**)). n = 9458 clusters
in 17 cells from 17 plants. (F) Distributions of KEULE colocalization events either on or off microtubules
on observed data. The distance threshold for microtubule association was < 200 nm. Sequential KEULE
cluster co-localization was 1.25 times more frequent for KEULE events associated with microtubules than
for those not associated with microtubules, as determined by odds ratio analysis (see panel (G)). n = 4000
clusters on microtubules and 4866 clusters off microtubules in 17 cells from 17 plants. (G) Odds ratios
and 95% confidence intervals for the data shown in panels (G) and (F). The dotted line indicates the odds
ratio expected for the null hypothesis of no different from random.
1189

1190

1191   **Figure S6**

1192

1193   **Fig. S6. Fit parameters for CSC vesicles in the tethering frame and the nearest KEULE clusters to**
1194   **these tethering sites.** (**A**) Distributions of fit parameters for MLEwG fits of tdTomato-KEULE clusters.
1195   (**B**) Distributions of fit parameters for MLEwG fits of CIT-CESA6 signal. (**C**) Distribution of standard
1196   errors for KEULE to microtubule distance measurements. Black lines and values indicate means. n = 194
1197

1198   **Figure S7**

1199

1200   **Fig. S7.** Frame-to-frame displacements of tdTomato-KEULE clusters in 200 ms streaming acquisitions. n
1201   = 283,127 displacements for 15,130 clusters observed in 12 cells from 12 plants.

1202
1203

Figure S8

Fig. S8. Fit parameters for KEULE to microtubule distances used for localization analysis. (**A**) Fit parameters for MLEwG fits of tdTomato-KEULE clusters. (**B**) Fit parameters for MLEw1dG fitting of microtubule signal. (**C**) Standard error of KEULE to microtubule distance measurements. Black lines and values indicate means. *Truncations of standard deviation distributions are due to cutoff values used to filter and exclude bad fits due to signal patterns that complicate localization analysis. Bad fits for tdTomato-KEULE signal occur where more than one tdTomato-KEULE cluster are close in proximity, and bad fits for microtubule signal occur where microtubules are dense and where they intersect in complex patterns. n = 115,963 KEULE clusters, nearest microtubule locations and KEULE-microtubule distances observed in 14 cells from 14 plants.

Movie S1.

tdTomato-KEULE cluster dynamics. Spinning disk confocal images of tdTomato cluster formation and loss at the plasma membrane of an epidermal hypocotyl cell in a dark grown *Arabidopsis* seedling. To reduce noise, images were averaged with a running 3-frame window. Streaming acquisition at 5 frames/sec. Playback at 4x real time.

Movie S2.

MLE Gaussian fits of tdTomato-KEULE clusters. Spinning disk confocal images of tdTomato-KEULE clusters in an epidermal hypocotyl cell of a dark grown *Arabidopsis* seedling. Magenta circles indicate an automated detection of a KEULE cluster. For each of these detects the coordinates of the signal centroid number of detected photons were estimated by MLE Gaussian fitting (see methods). Streaming acquisition at 5 frames/sec. Playback at 10x real time.

Movie S3.

KEULE localization to CSC delivery events. Spinning disk confocal images of co-expressed tdTomato-KEULE and Cit-CESA6. Filled arrowheads point to locations of tracked CSC signal. Empty arrowheads point to co-localized KEULE clusters that appear shortly after appearance and positional stabilization of CESA6 signal. 2 second acquisition intervals. 3-frame running average. Playback at 10x real time.

Movie S4.

MLE Gaussian fits of tdTomato-KEULE clusters and EGFP-TUA5 labeled microtubules. Spinning disk confocal images of co-expressed tdTomato-KEULE and EGFP-TUA5 labeled microtubules. White circles indicate automated KEULE signal detects. The position of each cluster at the first frame of stabilization was estimated at a sub-pixel level if MLE Gaussian fitting. 1 second acquisition intervals. Playback at 20x real time.