Title: Cortical microtubules act as a template to organize nano-scale patterning of exocytosis

Authors: Jelmer J. Lindeboom[1,2]*, Ryan Gutierrez[1,3], Viktor Kirik[4] and David W. Ehrhardt[1,3]*

Affiliations:

[1]Department of Plant Biology, Carnegie Institution for Science; Stanford, CA 94305, USA.

[2]Current Address: Novo Nordisk Foundation Center for Genomic Mechanism of Disease, Broad Institute of MIT and Harvard, MA 02142, USA.

[3]Department of Biology, Stanford University; Stanford, CA 94305, USA.

[4]School of Biological Sciences, Illinois State University; Normal, IL 61790, USA.

*Corresponding authors

Email: ehrhardt@stanford.edu

jelmer@broadinstitute.org

Abstract

Targeting of exocytosis enables cellular morphogenesis, motility and polarized transport, yet relatively little is known about the targeting mechanisms in cellular systems. Here we show that the SEC/MUNC protein KEULE is a dynamic marker for individual secretory events and employ it as a live cell probe, that together with high-precision image analysis of thousands of events, reveal that cortical microtubule arrays act as two-dimensional templates that pattern exocytosis at the nano-scale in higher plant cells. This mechanism is distinct from previously described mechanisms involving motor-driven transport and defines ordered and adjacent linear domains where secretory events are higher and lower than expected, effectively redistributing exocytosis over most of the cell membrane. In addition, analysis of KEULE kinetics revealed distinct phases of assembly/disassembly that are differentially sensitive to experimental treatments that reduce exocytosis, revealing SEC/MUNC dynamics as a versatile and information rich read-out of exocytosis *in vivo*.

Main Text

Introduction

The ability to organize biomolecules at the cell cortex was a fundamental innovation in
the evolution of cellular life, enabling essential biological functions including cellular
morphogenesis, motility and polarized transport. Targeted exocytosis, non-random positioning
and fusion of secretory vesicles at the plasma membrane, is a crucial mechanism for creating
such cortical organization.

Across phyla, the microtubule cytoskeleton has been implicated in spatial organization of
exocytosis. This functionality is exemplified in plant and fungal cells by cellular tip growth, and
in animal cells by examples such as polarization of mammalian epithelia, cell migration and the
positioning of focal adhesions[1,2]. The molecular polarity of microtubules plays an important role
in how their function in secretory targeting is understood. For example, polarized arrays of
microtubules created by nucleation from centralized organizing bodies - such as centrosomes or
Golgi - can serve as directional tracks for motor-driven transport of secretory vesicles from
where they are created to destinations at the plasma membrane[2,3]. This process sometimes aided
by specific association of microtubule ends with the cell membrane, such as at or near targets
such as focal adhesions [1]. Centrally polarized arrays of microtubules can also organize secretion
through directed transport of factors that establish domains at the plasma membrane where
secretion is targeted to, such as for tip growth in fission yeast[4]. However, many cells across
diverse organisms feature interphase microtubule arrays that are not centrally organized and have
architectures where microtubules are not clearly polarized with respect to the plasma membrane[5].
Many of these arrays are implicated in important biological functions, such as morphogenesis,
but relatively little is understood about whether they act to support these functions by targeting
secretion, and what the mechanisms may be.

Gaining an understanding of secretory targeting mechanisms requires the ability to
observe individual secretory events with respect to targeting machineries and a means to verify
successful delivery of secretory cargo. Observing individual exocytosis events has been
challenging in many cell models because microtubules, secretory vesicles and the machineries of
secretion are often densely concentrated. Axially growing cells in the epidermis of higher plants
have features that provide an excellent experimental platform to make such observations and for
investigating how acentrosomal microtubule arrays may target and pattern secretion. In contrast
to cells with microtubule arrays that are polarized with respect to the cell interior and the plasma
membrane, these cells feature highly ordered arrays of microtubules that lie at the cell cortex in
parallel to and in contact with the plasma membrane along their lattices. Large regions of the
plasma membrane lie in a single focal plane, where the individual microtubules and bundles that
comprise these arrays are easily visualized due to their quasi-2-dimensional nature and
comparatively low density[6]. In previous studies, we found that cellulose synthase complexes
(CSCs), large integral membrane protein complexes which synthesize and extrude cellulose into
the cell wall, can be used to observe individual secretory events at the plasma membrane by
allowing dynamic tracking of both secretory vesicles and successfully delivered cargo in the
plasma membrane[7–9]. By tracking CSCs while simultaneously imaging dynamic cortical arrays,
we discovered that CSCs were preferentially delivered to the cell membrane at locations defined
by microtubules. This targeting did not involve vesicular transport along microtubules, not did it
involve defining secretory domains by end-on association of microtubule ends with the cell
membrane[10], thus suggesting a previously undescribed mechanism for microtubule-dependent

secretory targeting. CSC's themselves have been shown to interact with microtubules via a
specific linker protein (CSI1), a mechanism that can also tether CSC-containing vesicles and
organelles to microtubules[11–14], so it is not clear if this secretory targeting mechanism might be
unique only to delivery of CSC's to the plasma membrane.

The question of microtubule targeting of secretion is of special interest in the context of
plant cell growth because ordered cortical microtubule arrays are essential to create functional
cell shape. Depolymerization of microtubules by the drug oryzalin causes isotropic growth and a
complete loss of genetically determined cell shape. The predominant hypothesis for the function
of cortical microtubules in directing cell growth is that they guide the trajectories of CSCs as
they synthesize and extrude cellulose microfibrils into the cell wall, thus creating material and
mechanical anisotropy to direct expansion of the cell wall under tension by high in turgor. The
hypothesis that cortical microtubules guide the trajectories of CSCs was microtubules was
confirmed by live cell imaging studies[15]. However, disruption of both microtubule guidance of
CSCs and CSC secretion in *csi1* null mutants causes only a modest loss of cell growth
anisotropy[11,12,14]. In addition, although complete loss of growth anisotropy is caused by
microtubule depolymerization, this phenotype is not due to a loss of CSC secretion[10] nor a lack
of ordering in CSC trajectories[16–18], and oryzalin treated cells continue to grow at near normal
rates[19]. These observations suggest that cortical microtubules arrays have additional important
functions in directing cell shape, prompting us to ask if cortical microtubules may guide cellular
morphogenesis by acting as a general targeting mechanism, beyond CSC's, to pattern secretion
during cell growth.

To determine if secretion in growing plant cells is broadly patterned by cortical
microtubule targeting, we require a dynamic biological marker that identifies single exocytotic
events and is involved in a wide range of exocytotic traffic to the plasma membrane. To date,
such a marker has not been identified. Imaging studies of core secretory proteins, including
soluble *N*-ethylmaleimide-sensitive factor attachment protein receptors (SNARE proteins)[20–22]
and components of the exocyst complex[23,24], have not revealed a clear relationship between the
presence and dynamics of these proteins with verified exocytosis events nor have they shown
association of these proteins with microtubules[24].

SM proteins are essential regulators of SNARE-mediated vesicle fusion. In exocytosis,
SNARE proteins anchored to the vesicle interact with those on the plasma membrane and form a
membrane bridging complex with other partners that enables membrane fusion[25–27]. The soluble
SM proteins are known to bind to SNAREs, and evidence from genetic studies together with *in
vivo* and *in vitro* experiments suggest they function in vesicle docking, SNARE complex
assembly and vesicle fusion [28–33]. In plants, the *Arabidopsis* Sec1 homolog KEULE was
identified in a genetic screen for embryonic pattern mutants[34], where loss of function mutations
are lethal, causing bloated and multinucleate cells[35]. Consistent with other SM proteins, KEULE
interacts genetically and biochemically with Qa-SNAREs, specifically KNOLLE (SYP111) and
SYP121, and has been shown to play essential roles in cell plate assembly during cytokinesis and
secretory traffic to the cell membrane[20,36–40]. The demonstrated function of KEULE in secretion
to the plasma membrane, combined with its severe loss of function phenotype due to lack of
secretion suggested that KEULE might serve as an excellent candidate for a generalized reporter
of secretion.

The molecular and structural mechanisms by which SM proteins interact with and
regulate SNARE complex assembly have investigated intensively by genetic, biochemical, *in*

vitro and structural studies[28–33], and SM proteins have been imaged in live cells in several
systems[41–48]. However, important features of their behaviors and biological function in living
cells remain to be uncovered. These questions include the fine scale cellular distributions of SM
proteins and clusters, the association of SM clusters with exocytosis events, the relationship of
these exocytosis events to potential organizing machineries and the dynamics of SM assembly
and loss at exocytosis sites, especially prior to and during vesicle fusion.

Here, we develop and apply automated and high precision analysis methods to detect and
track tens of thousands of individual fluorescently-tagged KEULE clusters to investigate their
dynamics and to map their positions at the plasma membrane in living cells. We evaluate these
clusters as secretory events by use of CSC's as a validated secretory cargo and employ genetic
analysis to test for the function of KEULE in CSC delivery. Finally, we employ high precision
mapping of KEULE clusters and microtubules, combined with Baysian statistical methods, to
quantitatively investigate the cortical microtubule cytoskeleton as a template in patterning
general exocytosis.

Results

Functional dynamics of tdTomato-KEULE cluster formation

To visualize KEULE protein *in vivo*, we created a tdTomato-KEULE fluorescent reporter
constructed from the chromosomal gene with native upstream regulatory sequence and
introduced it into *Arabidopsis* plants heterozygous for a *keule* loss of function allele. To obtain
plants for study of KEULE localization and dynamics, we screened for self-fertilized progeny of
these transgenic plants that exhibited rescue of *keule* mutant phenotype by the tagged KEULE
protein. These plants exhibited full rescue of plant growth and development through the 5-day
seedling stage used for our further studies (Fig. S1, See Methods).

In dark-grown *Arabidopsis* hypocotyl cells expressing the tdTomato-KEULE construct,
we observed KEULE signal appear as densely distributed and temporally dynamic puncta at the
plasma membrane (See Fig. 1A). The dynamics of signal increase and decay were apparent in
time-lapse movies (Movie S1), kymographs of signal intensity over time along transect lines
(Fig. 1B) and in sequential image samples of individual puncta (See Fig. 1C). In all, a steady
increase in signal intensity was observed, followed by an abrupt decrease and disappearance of
signal. This behavior indicates the accumulation and loss of many fluorescently labeled KEULE
proteins, which we will refer to hereafter as clusters.

The dynamics of tdTomato-KEULE accumulation and disappearance are presumably
related to its function. Therefore, we reasoned that a quantitative and robust picture of KEULE
cluster dynamics would be useful in gaining insight into the underlying molecular mechanisms
of secretion. To accomplish this, we collected a large-scale and high-resolution dataset
comprising thousands of individual events. We acquired streamed images at 200 ms intervals and
performed automated detection and frame-to-frame linking of KEULE cluster signal using
µTrack[49]. This track information was then used to extract position and intensity information by
Maximum Likelihood Estimate 2D Gaussian fitting[50] (for details see Methods, Movie S2). To
define events of signal appearance, accumulation and loss, we used a minimum sample of 4
sequential acquisitions, which made our shortest measured lifetimes approximately 800 ms. The
lifetime distribution of tdTomato-KEULE clusters showed a large range, from this minimal event

length of 800 ms to over half a minute. The data followed a roughly exponential distribution with
a mean of 2.34 s and ~95% of lifetimes being less than 7 seconds long (See Fig. 1D).

The large lifetime range of observed KEULE cluster events might represent a collection
of functional classes. If so, such classes may by differentiated by their kinetics. To assess the
kinetics KEULE signal accumulation and loss, and to ask if these kinetics might vary as a
function of cluster lifetime, we leveraged our large data set to calculate KEULE kinetics for each
200 ms lifetime increment. Fig. 1E shows average intensities for all observed clusters with
lifetimes of 4.2, 4.4 and 4.6 s. All these plots show a similar pattern: an initial steep rise in
intensity, followed by a steady increase, then a short plateau phase, and ending with a sharp
decrease. When we extended this analysis over a larger range of lifetimes, we observed evidence
that the shorter-lived events accumulated tdTomato-KEULE intensity at a slower rate, suggesting
they might be a functionally different class (See Fig. 1F). At longer lifetimes, the plateau phase
became more extended, suggesting that these events may be paused or stalled (See Fig. 1F). To
examine further these differences in accumulation kinetics as a function of cluster lifetime, we
grouped events into three classes; short (0.8 to 1.6s), medium (1.8 to 6.2s) and long (6.4+s) – and
then averaged the intensities at each 200 msec timepoint (Fig. 1G, for details see Methods). The
timepoints following the plateau phase were omitted to reveal better differences or similarities in
signal accumulation. This analysis confirmed that, indeed, the short events show significantly
slower signal accumulation (See Fig. 1G). Interestingly, the medium and long tdTomato-KEULE
cluster lifetimes show nearly identical trends in signal intensity accumulation over time,
indicating that they might be part of the same functional population during KEULE
recruitment/assembly. It is noteworthy that for events up to ~5s, longer-lasting events reach a
higher maximum cluster signal intensity. A possible explanation for this phenomenon is that
larger vesicles require a larger area for membrane fusion and therefore more KEULE. Unlike the
medium class, the long class exhibited a prolonged plateau phase, possibly indicating stalled
exocytosis events or alternatively, an extended checkpoint for a class of exocytosis events
involving a large amount of KEULE, and therefore perhaps larger secretory vesicles.

Clusters of tdTomato-KEULE coincide with CSC delivery to the plasma membrane

To place SM protein accumulation in context of functional exocytosis, we took advantage
of our previous work on delivery of CSCs to the plasma membrane[7]. CSCs are assembled in the
Golgi system from multiple CESA catalytic subunits[51,52], move through the trans-Golgi network
(TGN) and are budded off into secretory vesicles that fuse with the plasma membrane[7]. Because
CSCs are formed of multiple CESA subunits, they can be labeled by fluorescent proteins in
sufficient number to allow their detection and the tracking of individual complexes during the
process of exocytosis. Labeled vesicles display rapid and erratic dynamics that abruptly become
stabilized at the optical plane of the plasma membrane with a mean duration of about 60 seconds,
followed by a transition to slow, steady and typically linear motility of between 250 and 450
nm/min. The transition erratic to static movement is interpreted as evidence for vesicular capture
and tethering at the plasma membrane, and the transition to slow and steady movement indicates
successful vesicle fusion and delivery of CSCs, which are propelled by the synthesis and
extrusion of cellulose microfibrils[15,17]. This stereotyped behavior makes CSCs an excellent
reporter for successful exocytosis, revealing a time and place for vesicle tethering, and
distinguishing vesicle tethering at the plasma membrane from successful vesicle fusion and cargo
delivery.

To visualize KEULE together with CSCs, we generated plants that express tdTomato-
KEULE and CIT-CESA6, a catalytic CSC subunit labeled with Citrine fluorescent protein[53] in a
keule/cesa6 double mutant background (See Methods). CIT-CESA6 signal is relatively dense in
the plasma membrane of rapidly growing cells so we photobleached the CIT-CESA6 signal over
a region of the plasma membrane to aid detection of new CSC arrivals and their relation to td-
Tomato-KEULE signal (See Fig. 2A, Movie S3). For each CSC arrival at the optical plane of the
plasma membrane, we determined the first frame in which the CESA6 signal became stationary
(an indication of interaction and tethering at the plasma membrane). We then identified the
nearest KEULE cluster in a range of -10 to 50 seconds of the first stable CESA6 frame and
determined the distance between the centroid of the CESA6 signal and the centroid of the nearest
KEULE signal by maximum likelihood 2D Gaussian fitting (See Methods). A search range
starting at -10 seconds was chosen because the exact frame of stationary transitions was in some
cases obscured by signal from clusters of other labeled organelles such as Golgi bodies and
TGN. If KEULE is involved in the exocytotic delivery of CSCs to the plasma membrane, we
expect to observe co-localization between CSC delivery events and KEULE clusters. However,
CSC deliveries, as defined by static tracks, are relatively sparse and long lived as compared to
KEULE cluster events, so some degree of colocalization in space and time is expected by chance
alone. To assess statistically the spatial and temporal association of CSCs delivery events and
KEULE clusters, we generated a simulated dataset of random locations within the
photobleaching areas and identified the nearest KEULE spot to those random spots within the
same time window we used for our experimental observations. We then compared the
distribution of distances to the nearest KEULE cluster between the CSC deliveries and simulated
spots (See Fig. 2B and D) and found that the experimental data were strongly skewed to shorter
distances ($p < 3.4$ e-56, Mann-Whitney U test). The pattern of over- and under-representation of
observed with respect to simulated data is more easily seen in a plot of the bin-by bin differences
where shorter KEULE to microtubule distances are over-represented and longer distances are
underrepresented (See Fig. 2C). Using a Bayesian method (See Methods), we show that the most
likely value of the experimental bias from the null hypothesis of a random association is
significantly above that expected by chance alone for CSCs with distances < 200 nm (See Fig 2E
and F) and longer distances are significantly (and strikingly) underrepresented in the
experimental data. Based on these data we define colocalization as observing a KEULE cluster
centroid within 200 nm from the position of CSC signal in the first frame in which it was
observed to stabilize (and in a time window of -10 to 50 seconds around that frame). By these
criteria, we found colocalization in 81% (157 out of 194) of observed delivery events compared
to 29% (13035 out of 45051) of random locations at the plasma membrane ($p < 1.4$e-50, Fisher's
exact test).

We considered that our automated experimental protocol may allow for underestimation
of colocalization between CSC delivery and KEULE. For example, the time interval used for
these experiments is 2.11 s, of which 400 ms is used for KEULE illumination. Since a substantial
portion of KEULE spot lifetimes at the plasma membrane is shorter than the ~1.7 s gap in each
acquisition cycle when KEULE is not illuminated, we do not expect to detect all KEULE spots
within the -10 to 50 second experimental window. In addition, not all illuminated KEULE foci
are successfully detected by the detection and filtering routines used for large-scale data
collection. To understand how these two cases, no illumination and no detection, affect our
colocalization results, we performed a hand-curated analysis using a custom MATLAB GUI. For
each CSC delivery, we examined whether there was overlap of KEULE signal with the centroid

of the CSC signal in the first stable frame (tethering frame). We found that in 92% (159 out of 172) of the cases, KEULE signal was overlapping with the position of CSC in the tethering frame within -10 to 50 s of CSC stabilization, suggesting that our automated analysis may indeed underestimate the true degree of colocalization frequency of CSC exocytosis events and KEULE.

The durations of KEULE clusters associated with CES deliveries had a distribution similar in character to that observed For KEULE clusters in general (Fig. 3A, compare to Fig.1D), with both appearing roughly exponential but long tailed. While lifetimes of clusters associated with CESA deliveries spanned a similar range of durations, those associated with CESA delivery were observed to have a significant bias towards longer values, with clusters associated with CESA having a median duration of 4.3 seconds, as compared to 2.3 seconds for clusters in general. To test the significant of this observed bias, we compared the proportions of durations longer than two seconds and found a highly significant difference - 59% vs 37% for CESA delivery-associated KEULE clusters events vs clusters in general (p=6.07E-09, Fisher's exact test). These results indicate that KEULE clusters with a wide range of lifetimes are associated with verified exocytosis events, with those associated with CESA complexes, which is a relatively large complex, biased towards longer lifetimes.

If KEULE has a function in CSC exocytosis, we also expect that KEULE clusters both are colocalized with CSC static phases and that the time between the start of the static CSC phase and the first detected KEULE spot has a non-random distribution. Indeed, the time offset between stable CSC and nearest KEULE spot observed is non-random and significantly differs from the random distribution of the simulated data (p < 7.8e-04 Mann-Whitney U test) (See Fig. 3B). These data show that the appearance of the nearest KEULE cluster correlates in time with the CSC signal becoming stabilized at the plasma membrane with a median time delay of 2.111 s (mean = 7.92 s), suggesting that KEULE is typically recruited and functions within several seconds after the time the CSC vesicle becomes stationary, and is presumably tethered, at the plasma membrane. These data now provide insight into the kinetics of SM protein recruitment and vesicle fusion relative to secretory vesicle stabilization and tethering at the plant cell membrane.

KEULE abundance is significantly increased at confirmed sites of CSC exocytosis in both space and time, but is it required for CSC exocytosis? To answer this question, we quantified CSC delivery in seedlings of parental plants expressing CIT-CESA6 and segregating a tdTomato-KEULE allele in a background homozygous for both *cesa6* and *keule* loss of function (See Methods). Seedlings examined for loss of KEULE function were identified by morphological *keule* phenotype and little to no tdTomato-KEULE signal. We found that the CSC delivery rate was reduced by over 17-fold, which is a significant (p<0.0001, Wilcoxon two sample test) and quantitatively substantial reduction (See Fig. 2G). The seedlings that showed little to no tdTomato-KEULE fluorescence were severely reduced in size, showed large, misshaped cells and did not grow beyond the size of a 2-day old wild type seedling. Taken together our results indicate that KEULE plays an essential role in exocytosis of primary CSCs, and that CSC secretion is spatially and temporally associated with KEULE clusters, which have stereotyped dynamics of assembly and loss.

KEULE cluster dynamics are sensitive to perturbations that reduce exocytosis

We previously reported that the number of active CSCs in the plasma membrane were
rapidly and significantly reduced by treatment with mannitol (osmotic stress) and the herbicide
isoxaben[7]. Measurements of CSC delivery revealed that mannitol treatment also caused a 100-
fold reduction in the delivery rate of CSC's, indicating that CSC loss at the cell membrane was
likely caused primarily by inhibition of secretion. Isoxaben has been characterized as a specific
cellulose synthase complex inhibitor[54], but its mode of action is not well understood. Here we (1)
determine if isoxaben treatment also reduces CSC exocytosis and (2) ask if analysis of KEULE
dynamics may provide insight into how these treatments act to inhibit exocytosis.

To address the questions above, we imaged cells expressing tdTomato-KEULE and Cit-
CESA6 following treatment with mannitol, isoxaben or carrier only (mock). We assayed CSC
delivery to the plasma membrane as described above and found that, like mannitol, isoxaben is
also a potent inhibitor of CSC secretion (Fig. S2).

Following mannitol and isoxaben treatment tdTomato-KEULE cluster intensity
dynamics overall displayed a similar signature as observed in mock treatment - a steady increase
in signal and a plateau phase followed by an abrupt disappearance of signal (See Fig. S3 A).
These results suggest that mannitol and isoxaben treatment do not inhibit exocytosis of CSCs in
a manner that absolutely blocks KEULE accumulation. However, when comparing accumulation
dynamics grouped for short, intermediate and long events, we see striking differences among
treatments (See Fig. S3B). Both short- (0.8-1.6s) and intermediate-lived (1.8-6.2s) events in the
mannitol treated cells accumulated tdTomato-KEULE signal more slowly and to lower levels
than in the mock treated cells. Intriguingly, the dynamics for tdTomato-KEULE clusters that last
for 6.4 s or more exhibited a markedly different pattern. Initially, from 0 to ~6 s, signal
accumulation was indistinguishable between mannitol and mock treated cells. After ~6 s,
intensity gain was stalled in mock treated cells but continued to steadily increase in those treated
with mannitol. Taken together, these data suggest that mannitol application, and the osmotic
stress it causes, has bimodal effects on KEULE dynamics and may impact exocytosis by more
than one mechanism. By contrast, we observed that isoxaben treatment slowed accumulation of
tdTomato-KEULE across events of all durations (Fig. S3 C).

We observed related differences among treatments in KEULE cluster lifetimes. In comparing the
distributions of cluster lifetimes in mock and mannitol treated cells, a bimodal effect was again
observed, with both shorter and longer lifetimes being over-represented in comparison with
mock treatment (and intermediate events being under-represented) (Fig S4A, B, D). As was the
case with intensity dynamics, isoxoben treatment showed more of a unimodal effect, with the
lifetime distribution being shifted to the left (Fig. S4B, C, E). Mannitol and isoxaben also caused
diverging effects on persistence of KEULE clusters, as assessed by survival analysis (Fig.
S4F,G). Interestingly, the rates of new KEULE cluster formation under both treatments were
similar to that observed in the mock control (Fig. S4H), indicating that initiation of cluster
formation was not affected. Taken all together, our data show that KEULE cluster dynamics are
sensitive to perturbations that affect exocytosis and reveal evidence that mannitol and isoxoben
treatments affect these dynamics in distinct ways, suggesting that may act on different aspects of
exocytosis.

Microtubules define spatial patterns of exocytosis

In this study we showed that exocytosis of cellulose synthase complexes (CSCs) is
marked by formation of KEULE clusters (Fig. 2) and we measured the mean flux of CSC
deliveries as 1.59 events per μm^2 per h (Fig. 2G). However, the mean flux of KEULE cluster
formation was 134.4 events per μm^2 per h (Fig. S4H), or 85-fold higher. Therefore, CSC
delivery only accounts for a small fraction of the KEULE foci at the plasma membrane. Here, we
utilize KEULE clusters as a more generalized marker for secretion to ask if microtubules pattern
CSC secretion specifically or if microtubule positioning of secretory machineries is a broader
and fundamental function for the interphase microtubule cytoskeleton. It is important to point out
that another possibility is that cortical microtubules might in fact interfere sterically with many
secretory events, yielding an overall antagonistic relationship. To differentiate between these
alternatives, we performed a large-scale quantitative analysis of the positions of KEULE events
relative to the locations of cortical microtubules.

To visualize KEULE cluster events together with cortical microtubules, we performed
live cell imaging at 1 second intervals in hypocotyl cells of *Arabidopsis* seedlings expressing
GFP-TUA5 and tdTomato-KEULE in a *keule* knock-out background (See Fig. 4A, Movie S4).
After tracking individual KEULE events at the plasma membrane and using MLEwG fitting to
extract positional information (Fig. 4B, See Methods), we identified the frame with peak
tdTomato-KEULE intensity as the reference point for each detected KEULE cluster position
(Fig. 4C panels 1-3). We then measured signal intensity in the microtubule channel along a series
of scan lines that were rotated at 9° intervals around the centroid of the KEULE cluster (See
Methods and Fig. 4C panel 4). On each of these scan lines, we detected all microtubule signal
peaks and fit them with a modified quasi-1-dimensional MLEwG fit (See Methods, Fig. 4D).
Since the microtubule signal is best represented as a quasi-1-dimenional Gaussian orthogonal to
the microtubule axis, we determined the orientation of the nearest microtubule as identified by
peak analysis and used the scan line most orthogonal to the microtubule angle for determining
the location of the microtubule central axis closest to the KEULE spot in question (Fig. 4C, panel
5). For the example, for the most orthogonal scan line shown in Fig. 4C panel 5, the intensity of
KEULE signal in photoelectrons is shown in Fig. 4D panel 1, the triple Gaussian MLEw1dG fit
in panel 2, and the decomposed individual peak fits in panel 3. We used the fit positional
coordinates of the nearest microtubule signal peak and the KEULE cluster to calculate the
distance and the photoelectron fit values to estimate the standard error of the measurement (See
Methods and Fig. 4C panel 6). We repeated this process for all detected KEULE tracks in each
movie to assess the spatial relationship between KEULE clusters and the nearest microtubules to
those clusters (Fig. 4E, panel 1).

To ask if the measured distribution of KEULE to microtubule distances differs from that
expected if KEULE cluster positions are independent of microtubule locations (the null
hypothesis), we first needed to determine what the expected distributions are under the null
hypothesis using our detection and analysis routines within the regions of interest in which we
made our experimental observations. For example, microtubule array structure, which varies
within and among cells, will have a strong influence on this distribution, as will the dimensions
of our detection kernels. Therefore, we generated a simulated set of random locations within
each region of interest across the length of the movies and calculated the distances from these
synthetic KEULE foci to the nearest microtubule (Fig. 4E, panel 2). This allows us to compare
the distribution of experimentally observed data to that expected if KEULE cluster positions are
independent of microtubule locations. If KEULE cluster positions are distributed at random with
respect to microtubule array structure, these two distributions should converge on identity at

large sample sizes. When the experimental and expected distributions are plotted together as
relative frequencies, they show evidence for distinct differences (Fig. 5A and B). To quantify
over- or under-representation of the experimental data given the expected data under the null
hypotheses, we used a Bayesian method to determine the most likely positive or negative bias at
each distance bin (See Fig. 5C and D and Methods). We found that at KEULE-microtubule
distances smaller than 180 nm, the maximum likelihood bias was significantly positive (more
frequent than expected) and between 180 and 700 nm this bias was significantly negative (less
frequent than expected). At distances > 700 nm we again found a positive, but smaller, bias (Fig.
5D). This spatial pattern of under- and over-representation is more easily visualized in a plot of
the differences between observed and expected frequencies (Fig. 5B).

When examining patterns of KEULE foci, we also observed that sites of previous
exocytosis events were 3.4 times more likely to show consecutive exocytosis events than random
sites at the plasma membrane (Fig. S5). Such "hot spots" were also significantly over-
represented within 180 nm of microtubules, but only modestly so (1.25 times more likely) (Fig.
S5F).

Taken together, our results reveal that microtubules spatially organize exocytosis
machinery by increasing the frequency of KEULE cluster formation within about 180 nm of the
center of the microtubules, a distance about 7 times the dimension of a single microtubule, while
at the same time decreasing the probability of KEULE cluster formation distally over a region
that is about 3 times as wide. This acts effectively as a concentrating mechanism for KEULE
cluster events that is patterned by cortical microtubule organization. As these KEULE clusters
overlap heavily in their dynamic characteristics with those observed associated with validated
exocytosis events of CSCs, this is strong evidence that microtubule organization of secretion is
not confined to guiding insertion of CSCs but is a more generalized feature of secretion in
growing plant cells.

Discussion

To assess imaging of SM/KEULE clusters as a general marker for exocytosis it was
important to answer two questions; if there are different types of signal foci that may represent
multiple types of cellular structures or functional events, and what features of signal dynamics
are associated with exocytotic events. For example, some exocytotic proteins, including V-
SNARES and constituents of the exocyst, have been shown to arrive at the site of exocytosis on
exocytotic vesicles themselves, while others reside in the cytosol or associate with the plasma
membrane and are recruited to the exocytosis site. While we found some evidence for KEULE
signal on cytosolic organelles, these foci were relatively infrequent and were easily identified by
their positional dynamics. Further, in experiments using labeled CSCs as a trackable secretory
cargo, no KEULE signal was detected arriving at validated secretory sites on CSC containing
vesicles, indicating that KEULE clusters at the cell membrane form by recruitment of KEULE to
sites of secretion after arrival and interaction of cargo-carrying organelles with the plasma
membrane. KEULE assembly dynamics of clusters associated with CSC secretion had similar
characteristics as the KEULE population in general, including a wide range of cluster lifetimes
from a couple of seconds to half a minute and stereotyped dynamics of signal accumulation and
loss. The demonstrated functions of SM proteins in mediating SNARE complex formation and
membrane fusion, together with their stereotyped dynamics and tight association with CSC
secretion, indicate that dynamic KEULE clusters are valuable markers of individual secretory
events.

Our measurements of thousands of KEULE clusters at the plasma allowed us to analyze
patterns of signal accumulation and loss as a function of varying cluster duration - revealing
features of, and raising new questions about, the underlying mechanisms of secretory machinery
assembly, maturation and resolution. Over all durations, KEULE recruitment features an initial a
phase of rapid accumulation, followed by continued recruitment at a slower rate. This pattern
suggests two phases of recruitment with different kinetics. The second rising phase is near linear,
indicating a near constant rate of addition over time, characteristic of mechanism such as
polymerization, rather than recruitment of SM proteins to a fixed number of pre-existing binding
sites. Such a kinetic might be consistent with the sequential addition of SM-bound SNARE-PINs
to a growing chain of laterally-interacting SNARE-PINs at the vesicle-plasma membrane
interface[55–57]. This accumulation phase ends and transitions to a plateau, after which signal
abruptly decays to non-detectible levels. Both the final signal level and the length of the pause
state correlate positively with cluster assembly time, with longer assembly times having higher
signal levels and longer pause durations. Over the population of observed events, global peak
signal is reached at durations of about 6 seconds, after which longer events involve no further
signal accumulation but feature only lengthened plateau times. The plateau phase could represent
a state where all the SNARE complexes required have been recruited to the fusion site and the
components of the fusion super complex are transitioning to a functionally competent state prior
to the enactment of membrane fusion. The plateau could also reflect an assembly checkpoint.
Previous *in vitro* studies have found that greater numbers of SNARE proteins are recruited to
interfaces of large vesicles and planar membranes[58,59]. It is possible, therefore, that our
measurements of KEULE dynamics are shaped in part by a distribution of vesicle sizes, with
longer assembly phases that build a brighter SM signal representing assembly of secretory
machinery at larger vesicles.

We found that KEULE dynamics were changed when we disrupted exocytosis with
osmotic stress by treatment with mannitol, or by application of a drug, isoxaben. The changes in
dynamics were qualitatively different for the two antagonists, indicating that they likely act on
different processes affecting secretion. These observations indicate that KEULE dynamics,
which indicate functional aspects of vesicle fusion, are a sensitive and useful readout, or marker,
for studying exocytosis and agents that disrupt or alter exocytosis.

High resolution mapping and Baysian statistical analysis of individual secretory events
marked by KEULE clusters revealed that these events are not randomly distributed in growing
hypocotyl cells but are significantly over-represented in domains defined by the cortical
microtubule array. This patterning of secretion differs from previously described mechanisms for
secretory targeting by microtubules. In the latter, the positioning of microtubule plus ends define
the locations for motor-transported delivery of either the secretory vesicles themselves or
proteins that act in defining secretory domains. By contrast, our data indicate that every part of
the microtubule is associated with secretory targeting, making the entire cortical array a dynamic
and steerable template for secretory patterning and organization of the cell cortex (Fig, 6A).

High precision measurement of both microtubule and KEULE cluster locations revealed
that the spatial range of influence for microtubule secretory targeting is significantly larger than
the scale of microtubules themselves. These measurements revealed that domains of higher
secretion frequency are approximately 180 nm wide on either side of the center of microtubules
or microtubule bundles, creating ~360 nm wide liner domains on average. Our analysis also
revealed adjacent domains, from approximately 180-700 nm from microtubule centers, where

KEULE cluster density was lower than expected by chance alone (Fig. 5). These results indicate
microtubule pattern secretion both positively and negatively and do so over spatial domains
larger than a micron wide, whereas the scale of microtubules are ~25 nm and bundles of two
microtubules in the range of 75 nm wide [60]. The width of the combined over-represented and
under-represented domains are about half the mean distance between ordered microtubules and
microtubules bundles in these cells [61], suggesting that secretion in most of the observed plasma
membrane is influenced by microtubules in these cells (Fig. 6B). In addition, the combination of
wider depleted domains coupled with narrower domains of elevated KEULE cluster density
suggests a concentrating effect that further enhances the relative difference in secretory
trafficking adjacent to microtubules as compared to distances several hundred nanometers away.

One mechanism that could explain the spatial scales of these microtubule-associated
secretory domains is direct interaction of secretory organelles with cortical microtubules, such as
we previously observed for vesicles containing CSC's[7,62]. Microtubule-tethered secretory
vesicles might capture local KEULE associated with the cell membrane, perhaps as bound to
SNARE partners such as SYP121[36], depleting KEULE concentration in the nearby plasma
membrane and making secretion there less likely, thus creating the observed zones of reduced
secretion (Fig. 6B). Alternatively, upstream exocytotic machinery, such as plasma-membrane
associated components of the exocyst complex, might be recruited by a microtubule-organized
membrane domain, elevating location concentration while causing depletion and thus vesicle
capture at a distance (Fig. 6B). Evidence that such microtubule-organized membrane domains
might exist has been suggested by observations of the plasma membrane associated actin
polymerase Formin1, which reveals narrow "dark" linear domains defined by cortical
microtubules where fluorescently-tagged Formin1 is excluded[63].

The observed flux of KEULE clusters per unit area and time was about 85-fold higher than that
measured for CSC delivery, therefore far more exocytosis events are associated with domains
defined by microtubules than can be accounted for by CSC secretion alone, implying either that
many or most exocytosis events in these cells are influenced by microtubules to some degree, or
that a quantitatively significant subpopulation of specialized traffic is redistributed, ~13% of
KEULE clusters as measured by our analysis. One possibility for specialized secretory traffic
targeted by cortical microtubules are cargoes involved in cell wall biosynthesis and cell wall
expansion. Supporting this idea is that KEULE has been shown to be an essential partner of the
Qa-SNARE SYP121, which is implicated by a growing body of evidence in mediating secretion
required for cell growth and expansion[21,36,37,64,65]. Cellulose is synthesized at the plasma
membrane, while many other structural oligosaccharides, such as pectins[66], and wall-active
proteins, such as expansins[67], are secreted. Targeting this secretory traffic by cortical
microtubule arrays to the same locations where newly synthesized cellulose microfibrils are also
being laid down could help coordinate the assembly of these diverse components to build a
functional cell wall. In addition to coordination of wall assembly, it is possible that the patterned
delivery of cell wall oligosaccharides and cell wall active proteins could itself act to support the
anisotropic expansion of cell walls required to create specific cell shape. Targeting the delivery
of proteins and other molecules that promote wall remodeling by ordered microtubule arrays
could create linear and parallel microdomains of increased wall extendibility that lie
perpendicular to the main axis of the cell growth - shaping the anisotropy of cell wall expansion
in the same direction as proposed in prevailing models for constraint of expansion by cellulose
microfibrils organized by microtubules[68]. As discussed in the introduction of this manuscript, an
important reason to consider alternative functions for cortical microtubules arrays in guiding cell

wall expansion and cell shape is that experimental evidence suggests that guidance of cellulose
deposition and CSC secretion alone is not sufficient to explain the anisotropy of cell wall
expansion and cell growth. Microtubule targeting of wall-active secretion could also help explain
redirection of cell growth in response to signals, such as blue-light stimulation of phototrophic
growth. Perception of blue light by Phototropin receptors causes a rapid 90° reorientation of
cortical microtubule arrays[69], but the following redirection of expansion cannot be easily
explained by directed cellulose deposition model since previously synthesized cellulose is
deposited perpendicular to the direction of expansion. Microtubule targeting and patterning of
wall-active secretion might allow for redirection of cell expansion regardless of cellulose
orientation.

The insights we gained into patterning of secretion by cortical microtubule arrays was
enabled by having an experimental platform and methods that made measuring and assessing
many thousands of discreet molecular events possible. A secretory protein, such as KEULE, that
is implicated in mediating a wide range of a cell's secretory traffic might be expected to be
subject to a range of targeting mechanisms, including no targeting at all. Such a background can
easily obscure a meaningful association between the secretory marker and a hypothesized
secretory machinery. Very large sample sizes and appropriate methods to test for meaningful
associations, such as we employed here, can robustly reveal such otherwise hidden relationships.

The fact that growing plant cells have a planar microtubule array that is associated with
the cell membrane, and that exocytotic events as marked by KEULE protein are of relatively low
density, has provided an excellent platform to study how microtubules pattern exocytosis, and do
so at the level of individual exocytotic events. When we consider cellular microtubule function
in cellular trafficking, we typically think of directed transport of cargos by kinesins and dynein.
In contrast, this work highlights that microtubules can also function as scaffolds for organizing
exocytosis. This scaffolding function for microtubules might be considered in other systems
where there is indirect evidence that microtubules regulate exocytosis where they interact with
the plasma membrane, such as in focal adhesions and neuronal growth cones[1,70,71]. In addition,
scaffolded guidance of secretion should also be considered in the context of disease. For
example, amyotrophic lateral sclerosis (ALS) risk is associated with genetic variants which
reduce levels of functional UNC13A[72,73], and with genetic variation within several microtubule
associated proteins[72–75]. It is possible that these proteins all converge on the same biological
process of microtubule dependent patterning of exocytosis.

Materials and Methods

Generation of transgenic lines

The *pKEULE-tdTomato-KEULE* construct was generated by modifying the Gateway-compatible vector pMDC43[76]. A 2049-bp region immediately upstream of the start codon of *KEULE* (At1g12360) was amplified with primers CGTTTAAACTACAGTTCCATTGGAGCAAC and TAGGTACCTGCGATCTAACAATTT CAGACTTC. The tdTomato gene was amplified from a donor vector[77] with primers CAGGTACCATGGTGAGCAAGGGCGAGGAGGT and TGGCGCGCCCTT GTACAGCTCCTCCATGCCGT. The 35S promoter of pMDC43 was exchanged with the *KEULE* upstream regulatory region via PmeI and Acc65I sites, and GFP in pMDC43 was exchanged with *tdTomato* via KpnI and AscI sites. The full-length *KEULE* gene was amplified from genomic DNA using primers GGGGACAAGTTTGTACAAAAAAGCAG GCTTGACCATGTCGTACTCTGACTCC and GGGGACCACTTTGTACAAGAAAGCTGGG TTTATTTGGAGATCGTCTA and then inserted into the modified pMDC43 vector using Gateway cloning technology (Invitrogen) as described previously[78]. The construct was introduced into plants heterozygous for the keulemm125 deletion mutation[79] via Agrobacterium-mediated transformation[80,81]. T2 plants were genotyped for the *KEULE* locus using PCR: primers CGACCTTATCAGAGAGCAAGG and Keu-as0 (ACAGGGATTTGATTTGAGATCTGG) amplified a 1661-bp fragment from the wild-type allele and a 1505-bp allele from the deletion allele. Additionally, the primer pair GTCTGTTAACGGCTGCAGAA and Keu-as0 was used to amplify a fragment (1103 bp) only present in the wild-type allele. A T2 plant homozygous for the deletion allele and hemizygous for the *pKEULE-tdTomato-KEULE* transgene was isolated. A doubly homozygous line was subsequently isolated in the T3 generation and used for experiments.

To generate double-labeled plants expressing markers for KEULE protein and cellulose synthase, homozygous *pKEULE-tdTomato-KEULE/ keulemm125* plants (ecotype Landsberg) were crossed with homozygous *pCESA6-Citrine-CESA6/cesa6prc1-1* plants (ecotype Columbia[53]. An F2 line was selected that was homozygous for both mutations and hemizygous for both markers. In the F3 generation, a quadruply homozygous plant was isolated together with a plant homozygous for both mutations, homozygous for *pCESA6-Citrine-CESA6* and hemizygous for *pKEULE-tdTomato-KEULE*. Seed from the latter plant thus show segregation for the *keule* mutant phenotype.

To generate double-labeled plants expressing markers for KEULE protein and microtubules, *p35S-EGFP-TUA5* was introduced into *pKEULE-tdTomato-KEULE/keulemm125* plants via Agrobacterium-mediated transformation. T2 seedlings expressing the microtubule label were used for experiments.

Growth

Seedlings were grown in the dark on vertical agar plates as described in Lindeboom et al.[69]. For hypocotyl elongation assays, seedlings were grown for 5 days. For live cell imaging experiments, seedlings were imaged 3 days after germination.

Drug treatments

Oryzalin, taxol and isoxaben were dissolved in DMSO for 100x stock solutions. Oryzalin was used at a final concentration of 20 µM, taxol (Sigma) at 20 µM and isoxaben (Sigma) at 100 nM in plant growth media (Hoagland's No. 2). Mannitol was dissolved in plant growth media

directly at a concentration of 200 mM. For control treatment (mock), DMSO was added to the
growth media to a final concentration of 0.01%. Incubation of the seedlings was performed in 6-
well plates (WVR, nontreated) with 2 mL of plant growth media in each well. The plates were
wrapped in aluminum foil to block light and incubated at 22 °C. The drug treatment durations
were 16 hours for oryzalin, and 2 hours for all other treatments. Likewise, the mock treatments
were 16 hours for comparison to oryzalin, and 2 hours for comparison to the remaining
treatments.
605
Confocal microscopy
Imaging was performed on an instrument featuring a CSU-X1 spinning disk head
(Yokogawa Electric), a DMI6000 B inverted microscope (Leica), a 100X/1.4 NA oil immersion
objective (Leica), an Evolve 512 EMCCD camera (Photometrics) and a 1.2X telescope between
the spinning disk unit and camera. EGFP/CITRINE and tdTomato were excited at 488 nm and
561 nm respectively, by solid-state lasers (Coherent Cube). Switching, shuttering and intensity of
these lasers was performed by direct electronic control. Emission filtering was accomplished
with bandpass filters, (525/50 nm for EGFP and CITRINE and 636/37 nm for tdTomato
(SemRock), together with a 405/488/568 nm dichroic filter (SemRock). Exposure times were
400 ms for tdTomato-KEULE, 200 ms for EGFP-TUA5 and 800 ms for Citrine-CESA6. Images
were acquired at 1 second intervals in experiments in which both tdTomato-KEULE and EGFP-
TUA5 were imaged and 2 second intervals in experiments featuring tdTomato-KEULE together
with Citrine-CESA6. All light energies were approximately 4 mW as measured at the end of the
optical fiber entering the confocal head. For high-speed acquisitions of tdTomato-KEULE,
image streaming was performed at an acquisition rate of 5 Hz (200 ms exposures). The focal
plane in all experiments was stabilized by hardware Adaptive Focus Control (Leica) to mitigate
changes in specimen intensity over time caused by thermal and mechanical drift in focus.
Photobleaching was performed by a Vector laser scanning unit (Intelligent Imaging
Innovations) integrated into the setup above at the right camera port. Focused laser light was
pulsed and scanned (pulse duration = 10 ms, raster block size = 20 units) on the image plane by a
pair of galvanometer-driven mirrors inside the Vector unit. Scanner positions were calibrated to
image coordinates using an automated procedure in Slidebook software (Intelligent Imaging
Innovations). Citrine photobleaching was performed using the 488 nm laser at the same power as
used for imaging but focused on the image plane. Areas of ~200 μm^2 were bleached in 20-30 s.
All imaging was performed on the upper hypocotyl cells (1-2 mm below the apical hook).
Specimen mounting was performed as described previously[69].
632
Image registration
To correct for lateral specimen drift in the movies, we performed image registration in
MATLAB. We first performed a running average on the image sequence using a three-frame
window (yielding an image series 2 frames shorter than the original series). We then performed
an intensity-based image registration algorithm using the MATLAB function imregtform, using
tformtype "translation" and conserving the pixel size. To minimize interpolation artifacts, we
first determined the transformation matrix that yielded the best intensity match for each
consecutive image pair, then calculated the cumulative transformation matrix for the whole
averaged image series and applied it to the original movie that was not frame averaged, minus
the first and last frame that were lost because of the frame averaging. Some of the KEULE-
tdTomato and Citrine-CESA colocalization movies (3 out of 8) still exhibited sample drift after

intensity-based registration. In those cases, we applied a feature-based method employing the
locations of detected KEULE spots (See Methods: Particle Tracking for details) as registration
features and using the MATLAB function estgeotform2d and a rigid transform.
KEULE particle tracking, streaming experiments, image series with labeled microtubules
Particle detection and tracking was performed on the registered movies. For streaming
experiments and experiments featuring labeled microtubules (1 second time intervals) we used
µTrack[49] to detect and estimate the positions of tdTomato-KEULE clusters in each frame and
subsequently link detected foci from frame to frame to track individual KEULE clusters over
time. For each experiment type, we used the following parameters:
653
654   Streaming image series.
655   For detection: initial psfSigma = 1.2 pixels, alpha for local max 0.05 and 10 max
656   iterations for psfSigma estimation.
657   For tracking: max detection gap = 2 frames, min track length = 4 frames, max search
658   radius = 2 pixels, gap penalty = 4.
659
660   1-second interval image series.
661   For detection: initial psfSigma = 1.2 pixels, alpha for local max 0.05 and 10 max
662   iterations for psfSigma estimation.
663   For tracking: max detection gap = 1 frame, min track length = 1 frames, max search
664   radius = 3 pixels, gap penalty = 4.
665
KEULE and CESA particle tracking in dual labeled movies
In experiments assessing KEULE localization and dynamics with respect to cellulose
synthase complex (CSC) trafficking and we used the Spots module in Imaris software (Bitplane)
to detect and track both tdTomato-KEULE clusters and Citrine-CESA6 particles. The Imaris
Spots detector and filters performed better for Citrine-CESA6 signals, in particular. CSCs as
labeled by Citrine-CESA6 were detected using a size value of 0.3 µm and quality value of 110.
Tracking was performed by applying the Autoregressive Motion algorithm (maximum distance =
0.35 µm, maximum gap size = 2 frames). Successful deliveries of CSC complexes to the cell
membrane were determined by dynamic criteria as described previously[7]. Briefly, a delivery
event was defined when puncta that were mobile in the streaming cytosol (1) stabilized
positionally in the optical plane of the plasma membrane, and (2) showed a transition to slow,
steady and linear movement in the velocity range (~200-450 nm/min) we previously have
determined be characteristic of CSC's actively synthesizing and extruding cellulose fibers into
the cell wall[53]. To make this assessment, we first filtered CSC tracks by duration (>30 seconds),
then used a custom GUI in MATLAB to manually inspect and verify each Spots track and
associated image data for these characteristics. As a part of this process, the frame in which a
tracked particle first showed evidence of stabilized lateral position was marked as the "tethering"
frame – the frame showing the time and location of first stable interaction with the cell cortex of
a presumed vesicle carrying one or more labeled CSC's.
Clusters of tdTomato-KEULE were detected with Spots module using a size parameter
value of 0.3 µm, a quality value of 100 and were tracked using the Connected Components
algorithm (frame to frame displacements of KEULE signal are small and not auto-correlated).
These settings adequately detected and tracked KEULE clusters based on visual inspection of the
results.

690
Maximum Likelihood Gaussian fitting for particles

We employed maximum likelihood Gaussian fitting as described in Mortensen et al.[50] to use photon counts in estimating the accuracy of particle position and shape. To determine the number of photons collected in each image pixel, we first needed to calibrate our camera gain (photoelectrons per grey value or ADU). Photon detection can be modeled as a Poisson process, which creates a distribution in which the mean equals the variance. Therefore, the camera gain is equal to the coefficient between the mean and variance of the gray values in the image. To calculate this coefficient, we first measured the noise floor of the camera by taking the average of the gray values (intensity) of images acquired with the camera capped. We then determined the grey value variance by subtracting one image from the other then calculating the variance among these pixel-by-pixel difference values. To obtain the final variance, we divided this number by 2 to account for the fact we are using two images for determining the differences. Finally, we calculated the mean intensity of the images, subtracted the noise floor, and divided by the variance to estimate the camera gain.

For each time point in every track, we cropped out a pixel grid with the estimated center of the detected KEULE spot in the middle pixel (9 x 9 pixels). Along with a cropped image, the fitting routine requires a set of initial parameters: xCoordinate, yCoordinate, point spread function standard deviation (PSF SD), background, and an estimate of number of photons that comprise the imaged PSF. The x- and y-coordinates we used are the estimates of position from the particle tracking described above. Based on the approximate emission wavelength and objective numerical aperture, we used 133 nm for the initial standard deviation of the PSF. We estimated the background by calculating the median intensity of the outer border of the pixel grid used for fitting after subtraction of the camera noise floor. If the center of the point source is in the middle of the pixel and the PSF SD is 133 nm, we expect 0.16 of the photoelectrons to be in the center pixel. Therefore, we estimated the number of photoelectrons in the imaged spot by first subtracting the noise floor and pixel background from the intensity in the center pixel and multiplying the result by 1/0.16 (6.25). Before performing fitting using these initial data, we needed to address noise that could have been introduced by bilinear interpolation during the image registration that was employed to mitigate specimen drift. To accomplish this, we used the transformation matrix from registration to map the locations of the tracked particles back to the locations in the original image sequence, we then performed MLE fitting on the non-interpolated images by employing the method of Mortenson et al. [50] in MATLAB.

Since we used free fit parameters and there is noise in the data from photon counting, free fluorescently tagged proteins, high local particle densities and unresolved distributions of labeled structures, we excluded distance measurement from fits that were outside the expected parameter ranges. In the case of the SD of the 2D Gaussian, we found that there was a distinct population where the SD fit parameter was < 50 nm. Since our pixels correspond to 133 nm, in these fits all the photoelectrons are contained in one pixel, which is not consistent for a diffraction-limited spot at the magnification onto our camera (roughly 4 pixels). Similarly, we found that fit SD values above 2x the estimated PSF SD in most cases represented a fit containing more than one particle. Therefore, SD fit results with SD values outside 50 to 266 nm were excluded from position measurements. Fit results with estimated numbers of photoelectrons > 1000 were also considered outliers and excluded. Finally, high local particle densities the distance in some cases caused the MLEwG fit to center on a nearby particle and not the particle marked by the starting position. We therefore also did not include position information for fits where the distance

736 between the original estimate and the MLEwG position differed more than 2 x 133 nm. See Fig
737 S6 for fit parameters for the tdTomato-KEULE clusters and CIT-CESA6 labeled CSCs.
738
<u>KEULE to microtubule distance measurements</u>
To determine the position of each KEULE spot to assess the spatial relationship of
KEULE accumulation with the microtubule cytoskeleton we first determined the single frame
with the highest number of measured photoelectrons for each KEULE track. For tracks that were
longer than 4 frames we determined the max number of photoelectrons using a running three
frames average, for shorter tracks we did not use frame averaging. The rational for this selection
was two-fold. First, we reasoned that maximum signal might best be associated with the site of
KEULE function, with initial signal rise being indicative of KEULE recruitment and/or assembly
into a membrane fusion structure, and signal loss indicating KEULE disassociation or
disassembly of the fusion machinery. Second, the precision of estimated position is proportional
to photon sampling. It is also relevant to note that frame-to-frame movement of KEULE tracks
were small (Fig. S7).
To locate positions of microtubule signal relative to each KEULE track location, we used
a rotating linear filter combined with analysis of signal peaks and a two-step fitting procedure.
At the coordinates of KEULE signal in each max value frame we drew a line 21 pixels long and
1 pixel wide extending 10.5 pixels to each side of the KEULE position and repeated this process
in increments of 9 degrees to cover 360 degrees (Fig 4C). We then projected this array of lines
on the microtubule channel and extracted the intensities (in photons) along each line at 1-pixel
(133 nm) intervals by bilinear interpolation. To locate potential positions of microtubule signal
along each sampling angle we used the MATLAB function Findpeaks with a
MinPeakProminence of 25 to identify signal peaks. The 10.5-pixel distance is typically long
enough to contain the peak intensity of the nearest microtubule. In fact, in many cases this length
intersects multiple microtubules, but it is also possible for the end of the lines to partially
intersect a microtubule with a peak outside the 10.5 pixels. For subsequent fitting steps, we
included pixels from the center until the furthest peak on each side plus 3 pixels to catch the
shoulder of these peaks, and at the same time, exclude signal from microtubules whose peaks are
just outside the 10.5 pixel range from influencing the fitting. We then estimated the center
position of each signal peak by modifying the 2D MLEwG[50] to a quasi-1D version, modeling the
microtubule as a Gaussian wall[82] that is one pixel wide, as shown in the formula below.
768
769   (Formula 1)
770

$$N \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right) \left(\frac{1}{a} \right) \exp \left(-\frac{(x - \mu_x)^2}{2\sigma^2} \right) + b^2$$

771
Where:
N = number of photoelectrons
σ = Standard deviation
a = nm per pixel
x = position on the line
μ_x = center position
b = background photoelectrons
779

780   We found up to four peaks on a single line and fitted all peaks simultaneously by adding
781 an additional term describing each additional peak as follows:

783   (Formula 2)

$$N_{\text{first}}\left(\frac{1}{\sqrt{2\pi\sigma_{\text{first}}^2}}\right)\left(\frac{1}{a}\right)\exp\left(-\frac{(x-\mu_{x\ \text{first}})^2}{2\sigma_{\text{first}}^2}\right)+$$

786   (Formula 3)

$$N_{\text{second}}\left(\frac{1}{\sqrt{2\pi\sigma_{\text{second}}^2}}\right)\left(\frac{1}{a}\right)\exp\left(-\frac{(x-\mu_{x\ \text{second}})^2}{2\sigma_{\text{second}}^2}\right)+b_{\text{global}}^2$$

For estimating the seeding parameters for the fit, we used the center of the maximum
value pixel for each peak identified by Findpeaks. We used 160 nm (median value of the peak
population) as the estimate for the standard deviation and we approximated the background by
taking the square root of the lowest 10 percentile of all the pixels for all angles after subtracting
the noise floor.
After fitting all microtubule peaks along all angular steps, we determined the distance
from the KEULE location to the center of the nearest microtubule signal. We did this in three
steps. First, we determined the nearest microtubule signal peak to the KEULE location from
among the initial angular samples. Second, we determined the angle of the microtubule identified
by the closest peak. Measuring the distance from the KEULE location to the center of the
microtubule signal is most accurate along a line perpendicular from the KEULE location to the
microtubule. By determining the local microtubule signal angle, we could then determine the
optimal perpendicular sample angle. To find the local orientation of the microtubule we rotated a
line of 21 pixels long around the center of the location where we initially found the closest
microtubule peak. We extracted the summed signal intensities along these lines and determined
the angle at which we found the maximum intensity, corresponding to the microtubule angle[69].
Finally, once we established the microtubule angle, we used the angle perpendicular to that
microtubule angle to extract the nearest microtubule fit for the representative KEULE position.
Distance values where the fit parameters for the microtubule positions were the number of
photoelectrons were < 50 or > 10000, as well as SD fit parameters < 50 or > 250 nm were
excluded from distance measurements. We required all the microtubule fits on the line to be
within these parameters to be included in the analysis. See Fig. S8 (A to C) for fit parameters for
KEULE (A) and microtubule signal (B) fits after selection as well as the standard error of the
distance measurement between them (C).

<u>Analysis of KEULE localization with CESA6 delivery sites</u>
KEULE and CESA6 spots were detected and tracked in Imaris as described above and
maximum likelihood 2D Gaussian fitting was applied to each set of tracks. CESA tracks tend to
be relatively long and can be confounded by a dynamic background produced by highly motile
CESA containing organelles including Golgi bodies and Trans-Golgi network compartments.

This background makes fully automated detection and quantitation of CSC delivery events
challenging. We therefore designed a graphical user interface in MATLAB that allowed us to
efficiently curate potential CESA6 delivery events identified in Imaris and verify whether
individual tracks were indeed depicting a CESA6 delivery. We have shown previously that
CESA delivery to the plasma membrane typically shows three distinct phases; (1) erratic
movement of CESA containing vesicles as they appear in the optical plane at the cell cortex, (2)
positional-stabilization and (3) slow and steady movement with typically linear trajectories. We
speculate that the second phase represents molecular tethering at the cell membrane and the
events that lead to vesicle fusion and cargo delivery. The third phase is behavior that we have
shown is characteristic of active CSC complexes in the plasma membrane as they synthesize
cellulose[53]. Thus, observation of this latter behavior indicates successful delivery of CESA
cargo to the cell membrane. For each CESA track that showed these three distinct phases, we
marked the first frame of CESA stabilization at the plasma membrane. From this position, we
determined the distance to the nearest KEULE spot in a 30-frame time range (60 seconds). Some
degree of coincidence between CESA deliveries and KEULE spots is expected by chance alone,
especially since the density of KEULE clusters at the plasma membrane is relatively high. To
evaluate whether the observed distribution of KEULE locations with sites of CESA stabilization
might be due to chance alone, we generated random positions in the same regions of interest
used for the experimental data and calculated the distances to the nearest KEULE spot. We used
these simulated data to model the expected distribution of distances and compared this
distribution to the distribution of distances found for confirmed CESA delivery events. First, we
divided both the experimental and simulated distance distributions up into 40 nm distance bins
and calculated the bin-by-bin count frequencies by the total number of simulated counts. We
then introduced a bias parameter δ that equates an observed frequency in each distance bin to the
expected value from the simulated distribution as shown in formula (4).

843
844   (Formula 4)

$$P_{inDistBin}(\delta) = \frac{\text{Expected}_{inDistBin}\,\delta}{\text{Expected}_{\text{inDistBin}}\,\delta + \text{Expected}_{\text{outsideDistBin}}(1-\delta)}$$

845
The value of δ may vary between 0 and 1. For a possible observed frequency that equals
the expected frequency, δ is equal 0.5. If δ is greater than 0.5, an observed frequency is higher
than expected. Likewise, if δ is less than 0.5 the observed frequency lower than expected under
the null hypothesis. Use of the parameter allows us to use a Bayesian approach to test multiple
hypotheses to ask what bias value best may best explain the actual observed frequencies. To do
this, we calculated the Likelihood Ratio of observing an experimental result given a range of
values for δ (in increments of 0.001, excluding 0) for each frequency bin:

853
854   (Formula 5)

$$\frac{L(\delta)}{L\frac{1}{2}} = \frac{P_{inDistBin}(\delta)^M (1 - P_{inDistBin}(\delta)^{N-M})}{Expected_{inDistBin}^N\, Expected_{outsideDistBin}^{N-M}}$$

Where:
$L(\delta)$ = likelihood of the data given the bias δ,
$L\frac{1}{2}$ = likelihood of the data in the neutral case (no positive or negative bias)
N = total number of measurements

M = number of measurements in this specific distance bin

We then found the maximum a posteriori probability (MAP) for δ and computed the
shortest range of δ values for the highest posterior density (HPD) that contains 95% of the
normalized probability density. Maximum likelihood estimates for δ below 0.5 indicate
underrepresentation at the distance of interest and above 0.5 indicate overrepresentation of
experimental data compared to the unbiased case.

<u>Analysis of KEULE localization with microtubules</u>

We used the same Bayesian approach describe above for analyzing the localization of
KEULE clusters with respect to the estimated center of the nearest microtubule. We changed the
distance bin size from 40 to 20 nm.

<u>Hot spot analysis</u>
To investigate if some KEULE events are independent of each other each or not (e.g. if
"hot spots" for KEULE accumulation may exist), we examined colocalization between
sequential KEULE spots in time series collected from cells expressing both labeled KEULE and
tubulin proteins. Within a region of interest in each time series, we selected all KEULE tracks
that were first detected between frames 2 and 31 and calculated the distance to all other KEULE
tracks that started subsequently up to and including frame 301. For position estimates we used
the frame in the track that showed the highest number of fitted photoelectrons. To compare the
degree of colocalization of the initial KEULE tracks with future KEULE tracks to that expected
from chance alone, we created a simulated initial dataset with random locations within the same
region of interest and calculated the distances to all KEULE tracks in the same time range.
Since KEULE foci are positionally stable we chose a stringent distance threshold of 133
nm (1 pixel) to declare two spots or locations to be co-localized. Because of the relatively high
density of KEULE spots at the plasma membrane, colocalization is expected at some point in
time, therefore the time frame of colocalization analysis is also important. Indeed, we observe a
large degree of colocalization in the whole dataset for both the experimental and simulated data,
but with very distinct waiting time distributions (Fig. S5D). Based on the differences in the
distribution of waiting times between random simulated positions and KEULE positions until the
next KEULE spot arrives, we determined that a 40 second time delay window captures most
biologically relevant colocalizations while excluding most random colocalizations due to
KEULE density and chance.

1. Noordstra, I., and Akhmanova, A. (2017). Linking cortical microtubule attachment and exocytosis. F1000Research *6*, 469. https://doi.org/10.12688/f1000research.10729.1.

2. Spiliotis, E.T., and Nelson, W.J. (2003). Spatial control of exocytosis. Curr. Opin. Cell Biol. *15*, 430–437. https://doi.org/10.1016/s0955-0674(03)00074-7.

3. Fourriere, L., Jimenez, A.J., Perez, F., and Boncompain, G. (2020). The role of microtubules in secretory protein transport. Journal of Cell Science *133*, jcs237016. https://doi.org/10.1242/jcs.237016.

4. Chang, F., and Martin, S.G. (2009). Shaping Fission Yeast with Microtubules. Cold Spring Harb. Perspect. Biol. *1*, a001347. https://doi.org/10.1101/cshperspect.a001347.

5. Bartolini, F., and Gundersen, G.G. (2006). Generation of noncentrosomal microtubule arrays. J. Cell Sci. *119*, 4155–4163. https://doi.org/10.1242/jcs.03227.

6. Shaw, S.L., Kamyar, R., and Ehrhardt, D.W. (2003). Sustained Microtubule Treadmilling in Arabidopsis Cortical Arrays. Science *300*, 1715–1718. https://doi.org/10.1126/science.1083529.

7. Gutierrez, R., Lindeboom, J.J., Paredez, A.R., Emons, A.M.C., and Ehrhardt, D.W. (2009). Arabidopsis cortical microtubules position cellulose synthase delivery to the plasma membrane and interact with cellulose synthase trafficking compartments. Nat. Cell Biol. *11*, 797–806. https://doi.org/10.1038/ncb1886.

8. Sampathkumar, A., Gutierrez, R., McFarlane, H.E., Bringmann, M., Lindeboom, J., Emons, A.-M., Samuels, L., Ketelaar, T., Ehrhardt, D.W., and Persson, S. (2013). Patterning and Lifetime of Plasma Membrane-Localized Cellulose Synthase Is Dependent on Actin Organization in Arabidopsis Interphase Cells. Plant Physiol. *162*, 675–688. https://doi.org/10.1104/pp.113.215277.

9. Vellosillo, T., Dinneny, J.R., Somerville, C.R., and Ehrhardt, D.W. (2021). TRANVIA (TVA) facilitates cellulose synthase trafficking and delivery to the plasma membrane. Proc. Natl. Acad. Sci. *118*, e2021790118. https://doi.org/10.1073/pnas.2021790118.

10. Gutierrez, R., Lindeboom, J.J., Paredez, A.R., Emons, A.M.C., and Ehrhardt, D.W. (2009). Arabidopsis cortical microtubules position cellulose synthase delivery to the plasma membrane and interact with cellulose synthase trafficking compartments. Nature Cell Biology *11*, 797–806. https://doi.org/10.1038/ncb1886.

11. Li, S., Lei, L., Somerville, C.R., and Gu, Y. (2012). Cellulose synthase interactive protein 1 (CSI1) links microtubules and cellulose synthase complexes. Proc. Natl. Acad. Sci. *109*, 185–190. https://doi.org/10.1073/pnas.1118560109.

12. Bringmann, M., Li, E., Sampathkumar, A., Kocabek, T., Hauser, M.-T., and Persson, S.
(2012). POM-POM2/CELLULOSE SYNTHASE INTERACTING1 Is Essential for the
Functional Association of Cellulose Synthase and Microtubules in Arabidopsis. Plant Cell *24*,
163–177. https://doi.org/10.1105/tpc.111.093575.

13. Zhu, X., Li, S., Pan, S., Xin, X., and Gu, Y. (2018). CSI1, PATROL1, and exocyst complex
cooperate in delivery of cellulose synthase complexes to the plasma membrane. Proc. Natl.
Acad. Sci. *115*, E3578–E3587. https://doi.org/10.1073/pnas.1800182115.

14. Lei, L., Singh, A., Bashline, L., Li, S., Yingling, Y.G., and Gu, Y. (2015). CELLULOSE
SYNTHASE INTERACTIVE1 Is Required for Fast Recycling of Cellulose Synthase Complexes
to the Plasma Membrane in Arabidopsis. The Plant Cell *27*, 2926–2940.
https://doi.org/10.1105/tpc.15.00442.

15. Paredez, A.R., Somerville, C.R., and Ehrhardt, D.W. (2006). Visualization of Cellulose
Synthase Demonstrates Functional Association with Microtubules. Science *312*, 1491–1495.
https://doi.org/10.1126/science.1126551.

16. Gutierrez, R., Lindeboom, J.J., Paredez, A.R., Emons, A.M.C., and Ehrhardt, D.W. (2009).
Arabidopsis cortical microtubules position cellulose synthase delivery to the plasma membrane
and interact with cellulose synthase trafficking compartments. Nature Cell Biology *11*, 797–806.
https://doi.org/10.1038/ncb1886.

17. DeBolt, S., Gutierrez, R., Ehrhardt, D.W., Melo, C.V., Ross, L., Cutler, S.R., Somerville, C.,
and Bonetta, D. (2007). Morlin, an inhibitor of cortical microtubule dynamics and cellulose
synthase movement. Proc. Natl. Acad. Sci. *104*, 5854–5859.
https://doi.org/10.1073/pnas.0700789104.

18. Emons, A.M.C., Mulder, B.M., Emons, A.M.C., and Mulder, B.M. (2000). How the
deposition of cellulose microfibrils builds cell wall architecture. Trends Plant Sci. *5*, 35–40.
https://doi.org/10.1016/s1360-1385(99)01507-1.

19. Hamant, O., Heisler, M.G., Jönsson, H., Krupinski, P., Uyttewaal, M., Bokov, P., Corson, F.,
Sahlin, P., Boudaoud, A., Meyerowitz, E.M., et al. (2008). Developmental Patterning by
Mechanical Signals in Arabidopsis. Science *322*, 1650–1655.
https://doi.org/10.1126/science.1165594.

20. Zhang, B., Karnik, R., Alvim, J., Donald, N., and Blatt, M.R. (2019). Dual Sites for SEC11
on the SNARE SYP121 Implicate a Binding Exchange during Secretory Traffic1. Plant Physiol
180, 228–239. https://doi.org/10.1104/pp.18.01315.

21. Honsbein, A., Sokolovski, S., Grefen, C., Campanoni, P., Pratelli, R., Paneque, M., Chen, Z.,
Johansson, I., and Blatt, M.R. (2009). A tripartite SNARE-K+ channel complex mediates in
channel-dependent K+ nutrition in Arabidopsis. Plant Cell *21*, 2859–2877.
https://doi.org/10.1105/tpc.109.066118.

22. Assaad, F.F., Qiu, J.-L., Youngs, H., Ehrhardt, D., Zimmerli, L., Kalde, M., Wanner, G.,
Peck, S.C., Edwards, H., Ramonell, K., et al. (2004). The PEN1 Syntaxin Defines a Novel
Cellular Compartment upon Fungal Attack and Is Required for the Timely Assembly of Papillae.
Mol. Biol. Cell *15*, 5118–5129. https://doi.org/10.1091/mbc.e04-02-0140.

23. Bloch, D., Pleskot, R., Pejchar, P., Potocký, M., Trpko¡ová, P., Cwiklik, L., Vuka¡inović, N.,
Sternberg, H., Yalovsky, S., and Žárský, V. (2016). Exocyst SEC3 and Phosphoinositides Define
Sites of Exocytosis in Pollen Tube Initiation and Growth. Plant Physiol. *172*, 980–1002.
https://doi.org/10.1104/pp.16.00690.

24. Fendrych, M., Synek, L., Pečenková, T., Drdová, E.J., Sekereš, J., Rycke, R. de, Nowack,
M.K., and Žárský, V. (2013). Visualization of the exocyst complex dynamics at the plasma
membrane of Arabidopsis thaliana. MBoC *24*, 510–520. https://doi.org/10.1091/mbc.e12-06-
0492.

25. Barg, S., Knowles, M.K., Chen, X., Midorikawa, M., and Almers, W. (2010). Syntaxin
clusters assemble reversibly at sites of secretory granules in live cells. Proceedings of the
National Academy of Sciences *107*, 20804–20809. https://doi.org/10.1073/pnas.1014823107.

26. Knowles, M.K., Barg, S., Wan, L., Midorikawa, M., Chen, X., and Almers, W. (2010).
Single secretory granules of live cells recruit syntaxin-1 and synaptosomal associated protein 25
(SNAP-25) in large copy numbers. Proceedings of the National Academy of Sciences *107*,
20810–20815. https://doi.org/10.1073/pnas.1014840107.

27. McNew, J.A., Weber, T., Parlati, F., Johnston, R.J., Melia, T.J., Söllner, T.H., and Rothman,
J.E. (2000). Close Is Not Enough. The Journal of Cell Biology *150*, 105–118.
https://doi.org/10.1083/jcb.150.1.105.

28. Südhof, T.C., and Rothman, J.E. (2009). Membrane Fusion: Grappling with SNARE and SM
Proteins. Science *323*, 474–477. https://doi.org/10.1126/science.1161748.

29. Burgoyne, R.D., Barclay, J.W., Ciufo, L.F., Graham, M.E., Handley, M.T.W., and Morgan,
A. (2009). The Functions of Munc18 1 in Regulated Exocytosis. Ann. N. York Acad. Sci. *1152*,
76–86. https://doi.org/10.1111/j.1749-6632.2008.03987.x.

30. Baker, R.W., Jeffrey, P.D., Zick, M., Phillips, B.P., Wickner, W.T., and Hughson, F.M.
(2015). A direct role for the Sec1/Munc18-family protein Vps33 as a template for SNARE
assembly. Science *349*, 1111–1114. https://doi.org/10.1126/science.aac7906.

31. Stepien, K.P., Prinslow, E.A., and Rizo, J. (2019). Munc18-1 is crucial to overcome the
inhibition of synaptic vesicle fusion by αSNAP. Nat. Commun. *10*, 4326.
https://doi.org/10.1038/s41467-019-12188-4.

32. Peng, R.-W., Guetg, C., Abellan, E., and Fussenegger, M. (2010). Munc18b regulates core
SNARE complex assembly and constitutive exocytosis by interacting with the N-peptide and the

closed-conformation C-terminus of syntaxin 3. Biochem. J. *431*, 353–361.
https://doi.org/10.1042/bj20100145.

33. Hashizume, K., Cheng, Y.-S., Hutton, J.L., Chiu, C., and Carr, C.M. (2009). Yeast Sec1p
Functions before and after Vesicle Docking. Mol. Biol. Cell *20*, 4673–4685.
https://doi.org/10.1091/mbc.e09-02-0172.

34. Mayer, U., Ruiz, R.A.T., Berleth, T., Miséra, S., and Jürgens, G. (1991). Mutations affecting
body organization in the Arabidopsis embryo. Nature *353*, 402–407.
https://doi.org/10.1038/353402a0.

35. Assaad, F.F., Mayer, U., Wanner, G., and Jürgens, G. (1996). The KEULE gene is involved
in cytokinesis in Arabidopsis. Mol. Gen. Genet. MGG *253*, 267–277.
https://doi.org/10.1007/pl00008594.

36. Karnik, R., Grefen, C., Bayne, R., Honsbein, A., Köhler, T., Kioumourtzoglou, D., Williams,
M., Bryant, N.J., and Blatt, M.R. (2013). Arabidopsis Sec1/Munc18 protein SEC11 is a
competitive and dynamic modulator of SNARE binding and SYP121-dependent vesicle traffic.
Plant Cell *25*, 1368–1382. https://doi.org/10.1105/tpc.112.108506.

37. Karnik, R., Zhang, B., Waghmare, S., Aderhold, C., Grefen, C., and Blatt, M.R. (2015).
Binding of SEC11 indicates its role in SNARE recycling after vesicle fusion and identifies two
pathways for vesicular traffic to the plasma membrane. Plant Cell *27*, 675–694.
https://doi.org/10.1105/tpc.114.134429.

38. Karnahl, M., Park, M., Krause, C., Hiller, U., Mayer, U., Stierhof, Y.-D., and Jürgens, G.
(2018). Functional diversification of Arabidopsis SEC1-related SM proteins in cytokinetic and
secretory membrane fusion. Proceedings of the National Academy of Sciences *115*, 6309–6314.
https://doi.org/10.1073/pnas.1722611115.

39. Assaad, F.F., Huet, Y., Mayer, U., and Jürgens, G. (2001). The Cytokinesis Gene KEULE
Encodes a Sec1 Protein That Binds the Syntaxin Knolle. The Journal of Cell Biology *152*, 531–
544. https://doi.org/10.1083/jcb.152.3.531.

40. Assaad, F.F., Mayer, U., Wanner, G., and Jürgens, G. (1996). The KEULE gene is involved
in cytokinesis in Arabidopsis. Molecular and General Genetics MGG *253*, 267–277.
https://doi.org/10.1007/pl00008594.

41. Carr, C.M., Grote, E., Munson, M., Hughson, F.M., and Novick, P.J. (1999). Sec1p Binds to
SNARE Complexes and Concentrates at Sites of Secretion. J. Cell Biol. *146*, 333–344.
https://doi.org/10.1083/jcb.146.2.333.

42. Scott, B.L., Komen, J.S.V., Irshad, H., Liu, S., Wilson, K.A., and McNew, J.A. (2004).
Sec1p directly stimulates SNARE-mediated membrane fusion in vitro. J. Cell Biol. *167*, 75–85.
https://doi.org/10.1083/jcb.200405018.

43. Zilly, F.E., Sørensen, J.B., Jahn, R., and Lang, T. (2006). Munc18-Bound Syntaxin Readily
Forms SNARE Complexes with Synaptobrevin in Native Plasma Membranes. PLoS Biol. *4*,
e330. https://doi.org/10.1371/journal.pbio.0040330.

44. Medine, C.N., Rickman, C., Chamberlain, L.H., and Duncan, R.R. (2007). Munc18-1
prevents the formation of ectopic SNARE complexes in living cells. J. Cell Sci. *120*, 4407–4415.
https://doi.org/10.1242/jcs.020230.

45. Rickman, C., and Duncan, R.R. (2010). Munc18/Syntaxin Interaction Kinetics Control
Secretory Vesicle Dynamics2. J. Biol. Chem. *285*, 3965–3972.
https://doi.org/10.1074/jbc.m109.040402.

46. Smyth, A.M., Rickman, C., and Duncan, R.R. (2010). Vesicle Fusion Probability Is
Determined by the Specific Interactions of Munc18*. J. Biol. Chem. *285*, 38141–38148.
https://doi.org/10.1074/jbc.m110.164038.

47. Trexler, A.J., Sochacki, K.A., and Taraska, J.W. (2016). Imaging the recruitment and loss of
proteins and lipids at single sites of calcium-triggered exocytosis. Mol. Biol. Cell *27*, 2423–2434.
https://doi.org/10.1091/mbc.e16-01-0057.

48. Somasundaram, A., and Taraska, J.W. (2018). Local protein dynamics during microvesicle
exocytosis in neuroendocrine cells. Mol. Biol. Cell *29*, 1891–1903.
https://doi.org/10.1091/mbc.e17-12-0716.

49. Jaqaman, K., Loerke, D., Mettlen, M., Kuwata, H., Grinstein, S., Schmid, S.L., and Danuser,
G. (2008). Robust single-particle tracking in live-cell time-lapse sequences. Nat. Methods *5*,
695–702. https://doi.org/10.1038/nmeth.1237.

50. Mortensen, K.I., Churchman, L.S., Spudich, J.A., and Flyvbjerg, H. (2010). Optimized
localization analysis for single-molecule tracking and super-resolution microscopy. Nat.
Methods *7*, 377–381. https://doi.org/10.1038/nmeth.1447.

51. Haigler, C.H., and Brown, R.M. (1986). Transport of rosettes from the golgi apparatus to the
plasma membrane in isolated mesophyll cells ofZinnia elegans during differentiation to tracheary
elements in suspension culture. Protoplasma *134*, 111–120. https://doi.org/10.1007/bf01275709.

52. Zhang, Y., Nikolovski, N., Sorieul, M., Vellosillo, T., McFarlane, H.E., Dupree, R., Kesten,
C., Schneider, R., Driemeier, C., Lathe, R., et al. (2016). Golgi-localized STELLO proteins
regulate the assembly and trafficking of cellulose synthase complexes in Arabidopsis. Nat.
Commun. *7*, 11656. https://doi.org/10.1038/ncomms11656.

53. Paredez, A.R., Somerville, C.R., and Ehrhardt, D.W. (2006). Visualization of Cellulose
Synthase Demonstrates Functional Association with Microtubules. Science *312*, 1491–1495.
https://doi.org/10.1126/science.1126551.

54. Scheible, W.-R., Eshed, R., Richmond, T., Delmer, D., and Somerville, C. (2001).
Modifications of cellulose synthase confer resistance to isoxaben and thiazolidinone herbicides
in Arabidopsis Ixr1 mutants. Proceedings of the National Academy of Sciences *98*, 10079–
10084. https://doi.org/10.1073/pnas.191361598.

55. Zhou, Q., Lai, Y., Bacaj, T., Zhao, M., Lyubimov, A.Y., Uervirojnangkoorn, M., Zeldin,
O.B., Brewster, A.S., Sauter, N.K., Cohen, A.E., et al. (2015). Architecture of the
synaptotagmin–SNARE machinery for neuronal exocytosis. Nature *525*, 62–67.
https://doi.org/10.1038/nature14975.

56. Brunger, A.T. (2005). Structure and function of SNARE and SNARE-interacting proteins. Q
Rev Biophys *38*, 1–47. https://doi.org/10.1017/s0033583505004051.

57. Rothman, J.E., Krishnakumar, S.S., Grushin, K., and Pincet, F. (2017). Hypothesis –
buttressed rings assemble, clamp, and release SNAREpins for synaptic transmission. FEBS Lett.
591, 3459–3480. https://doi.org/10.1002/1873-3468.12874.

58. Cho, W.J., Jeremic, A., and Jena, B.P. (2005). Size of Supramolecular SNARE Complex:
Membrane-Directed Self-Assembly. J. Am. Chem. Soc. *127*, 10156–10157.
https://doi.org/10.1021/ja052442m.

59. Cho, W.J., Lee, J., Zhang, L., Ren, G., Shin, L., Manke, C.W., Potoff, J., Kotaria, N.,
Zhvania, M.G., and Jena, B.P. (2011). Membrane   directed molecular assembly of the neuronal
SNARE complex. J. Cell. Mol. Med. *15*, 31–37. https://doi.org/10.1111/j.1582-
4934.2010.01152.x.

60. Giddings, T.H., and Staehelin, L.A. (1988). Spatial relationship between microtubules and
plasma-membrane rosettes during the deposition of primary wall microfibrils in Closterium sp.
Planta *173*, 22–30. https://doi.org/10.1007/bf00394482.

61. Lindeboom, J.J., Nakamura, M., Saltini, M., Hibbel, A., Walia, A., Ketelaar, T., Emons,
A.M.C., Sedbrook, J.C., Kirik, V., Mulder, B.M., et al. (2019). CLASP stabilization of plus ends
created by severing promotes microtubule creation and reorientation. J. Cell Biol. *218*, 190–205.
https://doi.org/10.1083/jcb.201805047.

62. Crowell, E.F., Bischoff, V., Desprez, T., Rolland, A., Stierhof, Y.-D., Schumacher, K.,
Gonneau, M., Höfte, H., and Vernhettes, S. (2009). Pausing of Golgi Bodies on Microtubules
Regulates Secretion of Cellulose Synthase Complexes in Arabidopsis. Plant Cell *21*, 1141–1154.
https://doi.org/10.1105/tpc.108.065334.

63. Martinière, A., Gayral, P., Hawes, C., and Runions, J. (2011). Building bridges: formin1 of
Arabidopsis forms a connection between the cell wall and the actin cytoskeleton. Plant J. *66*,
354–365. https://doi.org/10.1111/j.1365-313x.2011.04497.x.

64. Geelen, D., Leyman, B., Batoko, H., Sansebastiano, G.-P.D., Moore, I., and Blatt, M.R.
(2002). The abscisic acid-related SNARE homolog NtSyr1 contributes to secretion and growth:

evidence from competition with its cytosolic domain. Plant Cell *14*, 387–406.
https://doi.org/10.1105/tpc.010328.

65. Grefen, C., Chen, Z., Honsbein, A., Donald, N., Hills, A., and Blatt, M.R. (2010). A novel
motif essential for SNARE interaction with the K(+) channel KC1 and channel gating in
Arabidopsis. Plant Cell *22*, 3076–3092. https://doi.org/10.1105/tpc.110.077768.

66. Hoffmann, N., King, S., Samuels, A.L., and McFarlane, H.E. (2021). Subcellular
coordination of plant cell wall synthesis. Dev. Cell *56*, 933–948.
https://doi.org/10.1016/j.devcel.2021.03.004.

67. Cosgrove, D.J. (2015). Plant expansins: diversity and interactions with plant cell walls. Curr.
Opin. Plant Biol. *25*, 162–172. https://doi.org/10.1016/j.pbi.2015.05.014.

68. Baskin, T.I. (2005). ANISOTROPIC EXPANSION OF THE PLANT CELL WALL. Annu.
Rev. Cell Dev. Biol. *21*, 203–222. https://doi.org/10.1146/annurev.cellbio.20.082503.103053.

69. Lindeboom, J.J., Nakamura, M., Hibbel, A., Shundyak, K., Gutierrez, R., Ketelaar, T.,
Emons, A.M.C., Mulder, B.M., Kirik, V., and Ehrhardt, D.W. (2013). A Mechanism for
Reorientation of Cortical Microtubule Arrays Driven by Microtubule Severing. Science *342*,
1245533. https://doi.org/10.1126/science.1245533.

70. Baas, P.W., Rao, A.N., Matamoros, A.J., and Leo, L. (2016). Stability properties of neuronal
microtubules. Cytoskeleton *73*, 442–460. https://doi.org/10.1002/cm.21286.

71. Tojima, T., Itofusa, R., and Kamiguchi, H. (2014). Steering Neuronal Growth Cones by
Shifting the Imbalance between Exocytosis and Endocytosis. The Journal of Neuroscience *34*,
7165–7178. https://doi.org/10.1523/jneurosci.5261-13.2014.

72. Brown, A.-L., Wilkins, O.G., Keuss, M.J., Hill, S.E., Zanovello, M., Lee, W.C., Bampton,
A., Lee, F.C.Y., Masino, L., Qi, Y.A., et al. (2022). TDP-43 loss and ALS-risk SNPs drive mis-
splicing and depletion of UNC13A. Nature *603*, 131–137. https://doi.org/10.1038/s41586-022-
04436-3.

73. Ma, X.R., Prudencio, M., Koike, Y., Vatsavayai, S.C., Kim, G., Harbinski, F., Briner, A.,
Rodriguez, C.M., Guo, C., Akiyama, T., et al. (2022). TDP-43 represses cryptic exon inclusion
in the FTD–ALS gene UNC13A. Nature *603*, 124–130. https://doi.org/10.1038/s41586-022-
04424-7.

74. Liu, X., and Henty-Ridilla, J.L. (2022). Multiple roles for the cytoskeleton in ALS. Exp.
Neurol. *355*, 114143. https://doi.org/10.1016/j.expneurol.2022.114143.

75. Zhang, S., Cooper-Knock, J., Weimer, A.K., Shi, M., Moll, T., Marshall, J.N.G., Harvey, C.,
Nezhad, H.G., Franklin, J., Souza, C. dos S., et al. (2022). Genome-wide identification of the
genetic basis of amyotrophic lateral sclerosis. Neuron *110*, 992-1008.e11.
https://doi.org/10.1016/j.neuron.2021.12.019.

76. Curtis, M.D., and Grossniklaus, U. (2003). A Gateway Cloning Vector Set for High-
Throughput Functional Analysis of Genes in Planta. Plant Physiol. *133*, 462–469.
https://doi.org/10.1104/pp.103.027979.

77. Shaner, N.C., Campbell, R.E., Steinbach, P.A., Giepmans, B.N.G., Palmer, A.E., and Tsien,
R.Y. (2004). Improved monomeric red, orange and yellow fluorescent proteins derived from
Discosoma sp. red fluorescent protein. Nat. Biotechnol. *22*, 1567–1572.
https://doi.org/10.1038/nbt1037.

78. Kirik, V., Herrmann, U., Parupalli, C., Sedbrook, J.C., Ehrhardt, D.W., and Hülskamp, M.
(2007). CLASP localizes in two discrete patterns on cortical microtubules and is required for cell
morphogenesis and cell division in Arabidopsis. J. Cell Sci. *120*, 4416–4425.
https://doi.org/10.1242/jcs.024950.

79. Assaad, F.F., Huet, Y., Mayer, U., and Jürgens, G. (2001). The Cytokinesis Gene KEULE
Encodes a Sec1 Protein That Binds the Syntaxin Knolle. J. Cell Biol. *152*, 531–544.
https://doi.org/10.1083/jcb.152.3.531.

80. Clough, S.J., and Bent, A.F. (1998). Floral dip: a simplified method for Agrobacterium
mediated transformation of Arabidopsis thaliana. Plant J. *16*, 735–743.
https://doi.org/10.1046/j.1365-313x.1998.00343.x.

81. Logemann, E., Birkenbihl, R.P., Ülker, B., and Somssich, I.E. (2006). An improved method
for preparing Agrobacterium cells that simplifies the Arabidopsis transformation protocol. Plant
Methods *2*, 16. https://doi.org/10.1186/1746-4811-2-16.

82. Ruhnow, F., Zwicker, D., and Diez, S. (2011). Tracking Single Particles and Elongated
Filaments with Nanometer Precision. Biophys. J. *100*, 2820–2828.
https://doi.org/10.1016/j.bpj.2011.04.023.

1162

Acknowledgments: We thank Michael Davidson for the kind gift of the tdTomato construct. We
thank Zdeněk Lánský, Johnathan Cooper-Knock and Masayoshi Nakamura for advice
and insightful comments.

Funding:

National Institutes of Health grant R15GM137247-01 (VK)

National Institutes of Health grant R01GM123259-01 (DWE)

1169

Author contributions:

Conceptualization: JJL, RG, VK, DWE

Methodology: JJL, RG, VK, DWE

Investigation: JJL, RG, VK, DWE

Visualization: JJL, RG, DWE

Funding acquisition: VK, DWE

Writing – original draft: JJL, RG, DWE

Writing – review & editing: JJL, RG, VK, DWE

1178

1179   **Competing interests:** The authors declare no competing interests.

1180   **Data and materials availability:** Biological material, raw data, processed data and MATLAB
1181   scripts used for the analysis are available upon request.

 Figures

 Figure 1

Fig. 1. KEULE assembly dynamics at the plasma membrane. (**A**) Distribution of tdTomato-KEULE
signal at the plasma membrane of etiolated hypocotyl cells in *Arabidopsis*. Arrowheads point out
individual clusters of labeled KEULE protein. (**B**) Kymograph of the tdTomato-KEULE signal intensity
over time as sampled over the transect line shown in (**A**) (blue dashes). The yellow brackets highlight
several individual tdTomato-KEULE clusters. (**C**) Example of tdTomato-KEULE cluster intensity over
time. (**D**) Histogram of KEULE cluster lifetimes in 200 ms bins. (**E**) Average tdTomato-KEULE cluster
signal intensity with a lifetime of 4.2, 4.4 and 4.6 s. Note the initial rapid increase in signal over the first
acquisition step (darker green), the more gradual, near linear, ramp up (green) to a plateau (yellow),
followed by abrupt loss to non-detection (red). (**F**) Average tdTomato-KEULE cluster signal intensity
with lifetimes ranging from 0.8 to 6.8 s. (**G**) Average signal intensities of tdTomato-KEULE clusters in
three cumulative lifetime ranges, 0.8 to 1.6 s, 1.8 to 6.2 s and 6.4+ s. The inset shows the lifetime
distribution histogram of tdTomato-KEULE cluster lifetimes with the lifetime ranges color coded. Data
acquired from 12 cells on 12 seedlings. Error bars indicate S.E. Scale bars are 2 μm.

Figure 2

B Closest CESA-KEULE Distances

Observed n=194
Simulated n=45051

p<0.000001
Kolmogorov-Smirnov test

C Observed-Simulated Distances

D
CESA6
KEULE
Merge
MLE fit locations

E Likelihood ratios of biased models (∂) over unbiased models at 0 to 40 nm

MAP(δ) →
95% CI →

null hypothesis →

F MAP(δ) estimation for distance bias

G CESA6 delivery

****

1.59

0.09

WT   *keu* mutant

1199
1200

Fig. 2. KEULE is spatially associated with and required for CSC delivery to the plasma membrane.
(**A**) CESA exocytosis event. (Top panel) Time series (top panel) and kymograph (bottom panel) of
tdTomato-KEULE and CIT-CESA6 during a CESA exocytosis event. The kymograph shows evidence for
CSC-vesicle tethering at the cell membrane (start of static phase) and successful CSC secretion as an
active complex (steady movement phase, see text). (**B**) Distribution of distances between the positions of
CIT-CESA6 signals at start of static phase and the nearest tdTomato-KEULE spot, as estimated by MLE
PSF fitting. Data shown both for observed CIT-CESA6 locations and simulated random locations. (**C**)
Differences of the observed vs expected CESA6-KEULE distance frequencies in (**B**), portraying over-
and under-representation (green and red respectively). (**D**) Examples of colocalization and fitted positions
for events in (**B**). (**E**) Bayesian likelihood ratios of the observed frequency of distances between 0 and 40
nm given bias δ over $\delta(0.5)$. The position of the peak value indicates the most likely, or maximum a
posteriori probability (MAP), value for the bias δ for the observed frequency given the expected
frequency. (**F**) Bayesian analysis of the MAP(δ) for each distance bin, MAP(δ) = 0.5 indicates no
difference from expected. Error bars are 95% confidence intervals expressed as the highest posterior
density (HPD, the shortest interval containing 95% of the probability density). (**G**) CESA6 delivery rates
in wild type and *keule* mutant cells. Data acquired from 8 cells on 8 seedlings. (Scale bars = 1µm).

Figure 3

1218

Fig. 3. KEULE cluster appearance is temporally associated with CSC appearance at the plasma
membrane. (**A**) Comparison of the waiting time distribution from CIT-CESA6 signal stabilization until
the detection of the nearest tdTomato-KEULE cluster in observed and simulated data. (**B**) KEULE
dynamics in the context of CSC secretion. Histograms of waiting times (left pane) and lifetimes of
tdTomato-KEULE clusters co-localized with CIT-CESA6 delivery events (right panel (**B**)).

Figure 4

1225
Fig. 4. Microtubule dependent distribution of exocytosis events. (**A**) Images of GFP-TUA5 and
tdTomato-KEULE signal in epidermal cells. Arrowheads indicate examples of apparent co-localization of
signal. (**B**) Spatial patterning of tdTomato-KEULE clusters over the range of 36.8 seconds. Crosses show
positions of MLEwG fits. (**C**) Steps in the routine employed to calculation the distance from a KEULE
cluster to the center of the nearest microtubule distance. (**D**) Steps in quasi-1-dimensional MLEwG fitting
of microtubule signal along the most-orthogonal scan line of a rotating filter. (**E**) Overlay of a
microtubule channel image and pairs of markers for MLE fit locations of KEULE clusters and the center
of the nearest microtubule in observed data (left) and for simulated random KEULE locations (right).
KEULE cluster detects from -5 to + 5 frames from the microtubule image are shown. Scale bars are 2 μm.

Figure 5

1236

Fig. 5. Bayesian analysis of microtubule bias for exocytosis events. (**A**) Comparison of the
distributions of KEULE-nearest microtubule distances for observed and simulated data. 8-12 cells (**B**)
Differences of the observed vs expected KEULE-microtubule distance frequencies shown in (**A**),
portraying the pattern of over- and under-representation (green and red respectively). (**C**) Bayesian
likelihood ratios of the observed frequency of KEULE-microtubule distances between 0 and 20 nm given
bias δ over $\delta(0.5)$. (**D**) Bayesian analysis of the MAP(δ) of each distance bin. Data acquired from 14 cells
in 14 seedlings. Error bars are 95% confidence intervals expressed as HPD.

 Figure 6

A

- — Microtubule
- Golgi body
- ● Secretory vesicle
- Kinesin
- Targeted domain
- ● Focal adhesions
- ▐▌ Adherens junction

B

Membrane domain model   Vesicle-MT interaction model

400 nm
200 nm
100 nm
70 nm

- — Plasma membrane
- · Microtubule
- · Plasma membrane linker
- ● Vesicle linker (i.e. CSI1)
- ▪ Exocytosis complex
- ● KEULE/SNAREs
- ● Microtubule-directed vesicle

▬ KEULE-MT distance median CI

KEULE +/- bias zones

200 nm 500 nm

1400 nm   Mean MT-MT distance

Fig. 6. Patterning of secretion by microtubule arrays. (A) Schematic diagrams of microtubule-directed patterning of secretion. The three cells on the left show targeting of secretion by motor-driven transport of secretory vesicles along polarized microtubule tracks[3] . The positioning of microtubule plus ends determines the location of cargo delivery. Clockwise from upper left: migrating fibroblast, epithelial cell, neuronal cell. By contrast, the diagram at far right shows an epidermal cell in the plant axis, where a 2-D array of microtubules at the cell cortex acts as an template that directs secretion in a complex but ordered pattern. The organization of the array as a whole defines this pattern. **(B)** Models for spatial patterning of secretion by cortical microtubule arrays in plant cells. All structures are abstracted but depicted at approximate scale. Microtubules (grey) are shown spaced at the mean spacing that has been measured in the cortical arrays of growing cells in the hypocotyl epidermis of dark grown *Arabidopsis* seedings. The red to green color between microtubules is a red-green heat map scaled from the observed over-representation and under-representation of KEULE clusters compared to that expected from a null hypothesis of a random distribution (as depicted in Fig. 5B). The two sides of the diagram depict two alternative models to explain the measured spatial dimensions of the domains enriched in secretion. On the right is model where the extent of enriched domain is determined by secretory vesicle size and direct interaction of these vesicles with cortical microtubules. Secretory vesicles are depicted in a variety of sizes over a range that has been observed by SEM studies to carry cell wall secretory cargos. The diagram at the upper right relates the dimensions of these vesicles to roughly estimated locations for associated exocytotic machineries at the plasma membrane if the vesicles are targeted by direct association with microtubules by linking proteins such as CSI1. On the left, microtubules establish a membrane domain where proteins needed for vesicle capture are concentrated or secretion are concentrated. These models are not mutually exclusive. The grey scale bar at lower left depicts the mean 95% confidence interval for the measured distances of KEULE clusters to microtubule centers.