Distinct roles for the *Staphylococcus aureus* POT transporter DtpT in di/tripeptide uptake and glutathione utilisation inside human macrophages

Imran Khan[a], Sandy J MacDonald[b], Sigurbjörn Markusson[c], Callum Robson[a], Joanne L Parker[c,d], Simon Newstead[c,d], Dave Boucher[a,], Marjan Van Der Woude[e] and Gavin H Thomas[a]

[a] Department of Biology, University of York, York, United Kingdom
[b] Bioscience Technology Facility, University of York, York, United Kingdom
[c] Department of Biochemistry, University of Oxford, Oxford, United Kingdom
[d] Kavli Institute of Nanoscience Discovery, University of Oxford, Oxford, United Kingdom
[e] Hull York Medical School, University of York, York, United Kingdom

Abstract

Peptides available in biological niches inhabited by the human pathogen *Staphylococcus aureus* serve as a rich source of amino acids required for growth and successful host colonisation. Uptake of peptides by *S. aureus* involves at least two transport systems: the POT family di/tri-peptide permease DtpT and the oligopeptide ABC transporter Opp3. Here we study the individual and combined functions of DtpT and Opp3 in enabling utilisation of 282 di-/tri-peptides via a high-throughput phenotypic screen. We reveal that many peptides can be utilised via either transporter, though DtpT appears to be the primary route of uptake for dipeptides. Intriguingly, we demonstrate a preference for Asp/Glu-containing peptides among putative DtpT substrates. To measure DtpT-mediated peptide transport directly, the protein was purified and reconstituted into proteoliposomes and active transport of diverse di- and tri-peptides was demonstrated, supporting the conclusions of the phenotypic screen. During this *in vitro* analysis, we also found that DtpT could transport the biologically prevalent tripeptide reduced glutathione (GSH). In the absence of the known glutathione transporter Gis, *dtpT* is essential for growth under conditions where GSH is supplied as the sole sulphur source, identifying DtpT as the predicted second GSH uptake system of *S. aureus*. Site-directed mutagenesis of the predicted ligand binding site of DtpT suggests GSH binds the protein vertically in a manner dependent on interactions with Gln310. Finally, we demonstrate that GSH transport is required by *S. aureus* for complete fitness during *in vitro* macrophage infection experiments, and our data suggest that GSH uptake plays a role in modulating host-pathogen interactions in these infections. Together, these data reveal important new functions for DtpT, both in the utilisation of diverse peptides for growth but also providing the first clues toward a distinct role of DtpT during intracellular infection.

Introduction

Staphylococcus aureus is a Gram-positive human bacterial pathogen associated with a broad range of different disease manifestations including skin/soft tissue infections, systemic bacteraemia, endocarditis and other invasive disease states associated with significant mortality (Tong et al., 2015). Understanding the intrinsic factors contributing to bacterial fitness, colonisation and infection in these diverse niches is essential for informing the design of novel therapeutic approaches. In order to meet their nutritional requirements during infection, bacterial pathogens such as *S. aureus* possess diverse transport proteins which allow for the selective uptake of nutrients present within the host environment. While the importance of some individual transporters in successful colonisation is well documented – particularly those implicated in the acquisition of key carbon sources, amino acids and metals (Adolf et al., 2023; Handke et al., 2018; Lehman et al., 2023; Lensmire et al., 2020) – the distinct role of many transport systems and their contributions toward infection remains understudied.

Peptides serve as an abundant source of amino acids, carbon, nitrogen and sulphur in many biological contexts and, as such, most bacterial pathogens possess one or more routes of uptake for these molecules. Bacterial peptide transporters can be broadly divided into ATP-Binding Cassette (ABC) importers and secondary active transporters belonging to the Major Facilitator Superfamily (MFS), with the latter being further divided into a small number of functionally distinct groups including the proton-dependant oligopeptide transporters, or POTs (Garai et al., 2017). ABC peptide importers are unique to prokaryotes and are capable of recognising and transporting diverse oligopeptides with high affinity, with substrates ranging from di/tri-peptides up to 35-mers (Detmers et al., 2000; Doeven et al., 2004; Hughes et al., 2022; Smith et al., 1999). In addition to their canonical role in nutrient acquisition, there are examples of these transporters contributing to diverse physiological functions including cell-wall turnover (Hughes et al., 2022; Maqbool et al., 2011) and antimicrobial peptide resistance (Ackroyd et al., 2024; Wang et al., 2016), as well as acting as sensory units for intercellular and environmental signalling networks (Slamti & Lereclus, 2019). POTs, by comparison, are found across the Tree of Life and exclusively transport di- and tri-peptides or chemically related molecules of a similar size. The importance of bacterial POTs during

infection is poorly characterised, and little is known regarding their contribution to bacterial

physiology outside of their general role in nutrient uptake.

To date, there are two known promiscuous peptide transport systems which are widely

distributed in Staphylococci: the ABC importer-type system Opp3$_{ABCDF}$ (or Opp3) and the di-

/tri-peptide POT transporter DtpT (Yu et al., 2014). In the single elegant study comparing the

roles of these two systems in *S. aureus*, Opp3 was demonstrated to be required for the

transport of oligopeptides from 3 to 9 residues in length, while only DtpT was thought to be

responsible for dipeptide uptake (Hiron et al., 2007). *S. aureus* also possesses an ABC

importer system which is specific for the biologically prevalent tripeptide glutathione (Gis$_{ABCD}$,

or Gis) (Lensmire et al., 2023). In addition to the aforementioned systems, different *S. aureus*

strains can possess up to four additional Opp3-homologues, of which Opp1 and Opp2 are

likely nickel rather than peptide transporters (Hebbeln & Eitinger, 2004; Remy et al., 2013).

However, Opp4 plus one additional Opp system (ACME Opp) encoded as part of the widely

disseminated arginine catabolic mobile element (Yu et al., 2014) are mostly uncharacterised

and could function as additional peptide transporters.

The importance of peptide utilisation in *S. aureus* infection biology is a subject of ongoing

investigation. *S. aureus* secretes an arsenal of extracellular proteases with diverse functions

in host immune modulation and virulence, and these enzymes contribute to the generation

of peptides within the local environment which are likely to serve as a nutrient source for the

bacterium (Coulter et al., 1998; Shaw et al., 2004). Consistent with this, there is evidence that

S. aureus senses protein-rich environments via Opp3 and responds to this stimulus via

upregulation of extracellular proteases (Borezée-Durant et al., 2009; Lehman et al., 2019).

Recent work has demonstrated how the *S. aureus* protease aureolysin is capable of

contributing to the degradation of collagen and highlights how growth on collagen-

degradation products is decreased in an Opp3-deficient mutant (Lehman et al., 2019). These

observations point toward a model in which host- and bacterial-derived peptides support the

growth of *S. aureus* in an Opp3-dependent manner. Conversely, loss of DtpT – but seemingly

not Opp3 – leads to attenuation in several models of invasive *S. aureus* infection including

intravenous infection in mice and an endocarditis model in rabbits (Coulter et al., 1998). In

spite of this, the functional characterisation of DtpT to date is limited, and the molecular basis

of its role in bacterial survival during infection is unknown.

Recent work has aimed to understand the molecular basis of substrate promiscuity observed
in POTs and delineate the mechanistic basis of proton-coupled peptide transport in these
proteins (Kotov et al., 2023; Lichtinger et al., 2024; Parker et al., 2014). Despite this work,
little is known regarding the substrate preferences of these transporters and how this relates
to cellular lifestyle/environment. Here, we aim to address this gap in our knowledge. We apply
Phenotype Microarrays (BIOLOG) to compare the utilisation of varied di- and tri-peptides by
wild-type *S. aureus* JE2 against strains with disruptions in the known peptide transporters.
This approach provides novel insights into the diversity of peptides utilised via DtpT and
points toward distinct amino acid preferences among DtpT substrates. Further cell-based
and biochemical assays serve to both support these findings and further interrogate the
specificity of the system. Finally, we extend these assays to demonstrate how DtpT also
serves as a transporter for the biologically prevalent metabolite reduced glutathione (GSH)
and demonstrate how glutathione transport mediated by DtpT and Gis is a determinant of
bacterial survival inside human macrophages.

Results

Phenotypic screening defines the substrate landscapes of DtpT and Opp3A

In order to investigate the roles of the two known peptide transporters in *S. aureus*, we generated an isogenic series of strains in the USA300-derivative strain JE2 with disruptions of *dtpT* and *opp3A,* as well as a double-mutant strain. With these strains in hand, we exploited the power of phenotype microarrays (PMs) as a simple high-throughput approach for comparing the abilities of bacterial strains to metabolise diverse substrates as energy sources, including di- and tripeptides (Biolog PM Plates 6–8). (Bochner et al., 2001; Pletzer et al., 2014). Collectively, these plates allowed us to compare utilisation for 282 different peptides, comprising: 246 standard dipeptides (of a possible 400), 1 hydroxyproline-containing dipeptide, 7 β-bonded dipeptides, 2 γ-bonded dipeptides, 12 D-amino acid-containing dipeptides and 14 tripeptides. In this assay, a given peptide is supplied as the primary nitrogen source, and catabolism is measured via reduction of a redox-sensitive colourimetric dye, quantified over time to produce a signal curve for each strain/peptide combination. A positive metabolic signal is indicative of peptide utilisation enabled by a valid route of uptake and downstream catabolic pathways, while intermediate/weak signals are likely to result from poor uptake (or a complete lack thereof) or incomplete catabolism following internalisation.

Based on the signal curves produced from our PM analysis, we could observe a series of distinct utilisation patterns across our four test strains (Supplemental data SD1-4). To identify commonalities in the data, we subjected the raw signal curves for each peptide to unbiased cluster analysis and dimensional reduction using Uniform Manifold Approximation and Projection (UMAP) (Figure 1). These analytical methods provided complementary outputs, with peptides belonging to a common cluster demonstrating close spatial organisation in the final UMAP plot (Figure 1A-C).

Visual confirmation of the clusters suggested that each encompasses a distinct pattern which reflects differences in the routes of peptide utilisation (Figure 1B,D). Cluster 1 (n = 15) contains peptides for which a strong metabolic signal was observed for JE2 and all three mutant strains, suggesting – rather intriguingly – that neither of the two known peptide transporters in *S. aureus* is required for their uptake. In contrast, utilisation of peptides in cluster 2 (n = 41) was maintained in single mutant strains lacking either *dtpT* or *opp3A* but abolished in the double-mutant strain, suggesting that these peptides can be transported by either of our two

systems. Clusters 3 and 4 (n = 99) both present patterns wherein utilisation is abolished for
the *dtpT* mutant and the double mutant, but not the *opp3* mutant, suggesting DtpT serves as
the primary route of uptake for these peptides. Collectively, these two clusters account for
more than one-third of the tested peptides, with those in cluster 3 (n = 58) demonstrating a
more pronounced phenotype (that is, a greater difference in signal between "positive" and
"negative" utilisation) when compared to those in cluster 4 (n = 41). Such a distinction likely
results from either a higher rate of internalisation by DtpT or more productive downstream
catabolism following uptake for cluster 3 peptides.

The three remaining clusters contain peptides for which a weak metabolic signal was
observed in the wild-type JE2 strain. Cluster 5 (n = 16) encompasses peptides for which,
quite surprisingly, metabolism is permitted only in mutant strains lacking *dtpT*. While the
exact mechanistic basis for this pattern is unclear, we postulate that the DtpT-mediated
internalisation of these peptides may be cytotoxic under the conditions tested. However, a
strong signal in the *dtpT* mutant strain suggests an alternate route of internalisation for which
toxicity is not observed, perhaps due to a lower overall flux which prevents toxic over-
accumulation of the peptide. The remaining peptides each gave a weak/intermediate signal
or were not utilised at all or across the four strains tested here, as observed in cluster 6 (n =
58) and cluster 7 (n = 53) respectively. These phenotypic patterns suggest that the supplied
peptide alone cannot be effectively utilised for cellular metabolism. We do note that the mean
eAUC values recorded for cluster 6 are slightly higher than those in cluster 7 across the four
strains, perhaps indicating that some peptides in cluster 6 can be utilised to produce an
intermediate metabolic signal.

Notably, our analysis did not identify a cluster (nor even any individual peptides) for which the
patterns of metabolism suggested that utilisation was solely dependent on Opp3, which is
perhaps expected given that this system is primarily implicated in the utilisation of larger
oligopeptides (n ≥ 3) (Hiron et al., 2007; Lehman et al., 2019). Taken together, our findings
implicate DtpT in the utilisation of at least 140 of the peptides assessed here, with 41 of these
being recognised by both peptide transport systems.

Figure 1 – UMAP and cluster analysis reveal differential peptide utilisation patterns across *dtpT* and *opp3A* mutant strains. (**A**) PM signal curves were subjected to UMAP and k-means cluster analysis (k = 7). Peptides belonging to the same cluster are grouped by colour on the UMAP plot, as labelled. (**B**) Regions of the UMAP plot corresponding to distinct phenotypic patterns (as demonstrated by cluster analysis) are highlighted. (**C**) Pie-chart representation of the number of peptides in each of the seven clusters identified here. In each case, colours correspond to the seven clusters identified in this analysis, as depicted by the key. (**D**) (**Left**) Cluster analysis reveals seven strain-specific peptide utilisation patterns. All signal curves for each cluster are overlayed and separated by strain. Mean signal curves for each strain are indicated by the bold line. (**Right**) Empirical area under the curve (eAUC) was calculated for all signal curves, as shown on the right. Bars show the mean eAUC for each strain within a given cluster. Error bars indicate standard deviation in each case.

Cluster analysis demonstrates amino acid preference among putative DtpT substrates

Next, we aimed to determine whether specific amino acids were enriched within the peptides of each cluster. Given the limited panel of tripeptides assessed, we focussed our analysis on the 246 standard dipeptides present in the assay. First, we plotted a grid to identify potential cluster-specific trends in regard to the N- and C-terminal amino acid composition of dipeptides included in the phenotype microarray analysis (Figure 2). Then, we examined any statistically meaningful relationships by applying a Fisher exact test (Figure 3A-D). The strongest association was identified between glutamine-containing peptides and cluster 1, with the vast majority (9/13) of Gln-containing peptides assessed here belonging to this cluster (Figure 3A). Such an observation strongly suggests that an alternate route of uptake exists for Gln-containing peptides in *S. aureus,* which is independent of both DtpT and Opp3. Another clear relationship was observed between tyrosine-containing dipeptides and cluster 7 (Figure 3C), suggesting that tyrosine is either poorly transported or metabolised by *S. aureus* under the conditions tested here. Tantalisingly, we observe that aspartate and glutamate-containing peptides are both strongly enriched within the DtpT-specific cluster 3 (Figure 3D). This preference is seemingly independent of amino acid arrangement, with comparable frequencies for DtpT-specific utilisation in the N- or C-terminal positions (Figure 2). Our findings provide compelling evidence that Asp- and Glu-containing peptides are preferred transport targets for DtpT, in-turn suggesting that utilisation of these peptides is a distinct function of the transporter.

In addition, we found that arginine-containing peptides were highly enriched in cluster 2 (Figure 3B), implicating both peptide transport systems in the utilisation of these peptides. Arginine auxotrophy in *S. aureus* USA300 was previously demonstrated using chemically defined medium containing glucose (CDMG) and the same medium lacking arginine (CDMG-R) (Jeong et al., 2022; Reslane et al., 2022). Here, we exploit this auxotrophy to verify the DtpT/Opp3A-dependence patterns of three arginine-containing peptides as observed in our phenotypic screen. Specifically, we supplemented CDMG-R with two cluster 3 peptides (Arg-Asp and Arg-Phe) and one cluster 2 peptide (Arg-Ala) and separately inoculated these media with JE2 and the three mutant-derivative strains. As expected, robust growth for all strains was observed in CDMG, while none of the strains were able to grow in CDMG-R (Figure 3E; Figure S1A). Supplementing with Arg-Asp (cluster 3) restored growth in a completely DtpT-dependent manner, while Arg-Ala (cluster 2) restored growth in all but the double mutant

strain (Figure 3E; Figure S1B). These patterns entirely support the findings of our phenotypic

screen. Unexpectedly, supplementing with Arg-Phe (cluster 3) restored growth in all but the

double mutant strain similarly to Arg-Ala (cluster 2). Such deviation is likely indicative of

differences in the sensitivity of the two approaches utilised here; that is, Opp3-mediated flux

of Arg-Phe is sufficient to relieve Arg auxotrophy when supplied at 1 mM, but too limited to

support metabolism as a sole nitrogen source under PM conditions. Finally, to further verify

the role of *dtpT* in the utilisation of these three peptides, we cloned the gene into a low-copy

number vector and were able to observe genetic complementation of growth in each case

(Figure S1C). Hence, we confirm that DtpT serves as a route of entry for cluster 2 and cluster

3 peptides, supporting the findings of our PM assays.

Figure 2 – Phenotype microarrays reveal differential routes of uptake for diverse dipeptides in *S. aureus.* Grid representation of all possible dipeptide structures arranged with N-terminal amino acids on the Y axis and C-terminal amino acids on the X axis. Amino acids are displayed as single-letter symbols and colour-coded according to their chemical properties (purple = hydrophobic/non-polar, green = polar/unique, blue = basic, orange = acidic). Coloured cells correspond to peptides included in the PM analysis and their corresponding utilisation patterns, as indicated by the key.

Figure 3 – Specific amino acids are enriched in peptides belonging to a common utilisation pattern. (**A - D**) Enriched amino acids within each cluster were identified via a Fisher exact test with FDR q-value correction. Plots show the position of peptides containing strongly enriched amino acids ($q \leq$ 1E-03) overlayed on the original UMAP plot (as labelled). The value of q is given in each case. (**E**) Growth of *S. aureus* strain JE2 and mutant derivatives in CDMG – R supplemented with three dipeptides, as labelled. Arg-Asp (RD) and Arg-Phe (RF) are each cluster 3 peptides, while Arg-Ala (RA) belongs to cluster 2. Bars show the mean end-point OD_{600} value after 24 hours of growth ± standard deviation. End-point OD is reflective of the maximal cell density observed over this period. *** $p < 0.001$; unpaired t-test.

DtpT transports peptides with diverse chemical properties

Peptide utilisation (as determined from phenotype microarrays) produces a composite

phenotype representing the combined consequences of peptide internalisation and

catabolism, both of which are required to drive cellular metabolism and produce a positive

signal. As a result, peptides with limited uptake may produce a positive PM signal if efficiently

catabolised while other peptides may be rapidly internalised (by DtpT or otherwise) without

producing a positive PM signal due to incomplete catabolic pathways.

To measure transport of peptides directly, we expressed and purified *S. aureus* DtpT

recombinantly from *E. coli* cells (Figure S2) and reconstituted the pure protein into liposomes

(Figure S3A-B), using methods previously applied to study the *Staphylococcus hominis* DtpT

orthologue $PepT_{Sh}$ (Minhas et al., 2018). Transport of di/tri-peptide substrates was then

assessed by utilising the pH-sensitive dye pyranine, as described previously (Parker et al.,

2014). Briefly, peptide transport was assayed by monitoring the acidification of the liposome

lumen and quantified over time for each peptide after induction of a negative-inside

membrane potential ($\Delta\Psi$) (Figure 4A). To validate this methodology, we confirmed that

peptide transport is dependent on the presence of both a valid substrate and $\Delta\Psi$, as

evidenced by the lack of strong acidification when either of these components is absent

(Figure S3C) and were then able to measure transport for a range of selected di- and

tripeptides (Figure 3B-C).

By assessing transport of 20 peptides with diverse chemical properties, we were able to

confirm DtpT-mediated transport for 12 of the putative substrates identified by our phenotype

microarrays, including all Asp-/Glu-containing peptides tested (Figure 4C). We also

confirmed the transport of Arg-Phe, Arg-Ala and Arg-Asp by DtpT, as already demonstrated

in our Arg-auxotrophy experiments. Overall, there is good agreement between our liposome-

based transport assays and whole-cell peptide utilisation methods (Figure 4C). We do note

that a small number of peptides are transported by DtpT which did not sustain metabolism

of *S. aureus* in the phenotype microarray assay, namely; Ala-Ala, Ala-Ala-Ala and Cys-Gly.

Such an observation suggests that peptide internalisation is not the limiting factor in

utilisation of these peptides. Additionally, our transport assays shed new light on the

molecular factors contributing toward substrate preference in this transporter. For instance,

our data demonstrate how many favourable DtpT substrates incorporate acidic amino acids

(Arg-Asp and Gly-Glu) or amino acids with small, uncharged side chains (Ala-Ala, Gly-Gly

and Gly-Pro) (Figure 4C). In contrast, we demonstrate that dipeptides containing aromatic

residues in position 2 generally demonstrate poorer transport when compared to those with

less bulky side-chains in the equivalent position. Poor internalisation of these substrates (e.g.

Val-Tyr and Ala-Phe) by DtpT is consistent with their lack of utilisation in PM assays (Figure

4C). Overall, our transport assays demonstrate that DtpT is highly promiscuous and capable

of transporting substrates with a range of chemical structures and properties, which is

similarly noted for other POT transporters studied previously (Kotov et al., 2023; Parker et al.,

2014).

Figure 4 – DtpT transports structurally diverse di/tri-peptides. (**A**) DtpT was reconstituted into liposomes containing pyranine. Upon induction of negative-inside ΔΨ, DtpT transports peptides from the external environment into the liposome lumen. Peptide transport is proton-coupled, leading to acidification of the lumen and a decrease in the pyranine fluorescence (ex. 460 / emm.510). (**B**) Exemplary transport data for 5 peptides normalised immediately before the addition of valinomycin (red arrow) for ease of comparison. Curves show the mean of three replicates. (**C**) Transport of 20 peptides was compared by quantifying the change in pyranine fluorescence over the first 30 seconds of the assay. Bars indicate the mean ± standard deviation. Coloured bars indicate a change in fluorescence which differs significantly from the no peptide ("Water") control, as determined by a one-way ANOVA with Fisher LSD test (p ≤ 0.05). Peptides are labelled according to their single-letter amino acid code (dAdA = d-ala-d-ala). Corresponding phenotypic patterns identified by PM analysis for each peptide are shown below.

DtpT is a functional glutathione permease

Our phenotype microarray data points toward a distinct role for DtpT in the utilisation of Asp/Glu-containing peptides (Figure 3D), and we have demonstrated that dipeptides in this category are generally favourable transport targets (Figure 4C). Reduced glutathione (GSH) is a biologically important tripeptide metabolite with diverse roles in many organisms, primarily serving as an antioxidant in eukaryotes and some prokaryotes (Lushchak, 2012). GSH consists of a Cys-Gly dipeptide conjugated to glutamate via a γ-linked peptide bond between the glutamate side chain and amino terminus of the cysteine (Figure 5A). Given that DtpT recognises and transports Cys-Gly (Figure 4C), we predicted that DtpT may also transport GSH and that this activity may be relevant in biological environments where glutathione is enriched. To test this, we measured DtpT transport activity with either GSH or oxidised glutathione (GSSG) (Figure 5B). We demonstrate that GSH is transported by DtpT, with the acidification of the liposome lumen over the first 30 seconds comparable to that observed for Cys-Gly. We do not observe the transport of GSSG in this assay, likely due to the larger structure of GSSG compared to GSH.

The Gis system is an ABC-type importer which has recently been characterised as a route of glutathione acquisition in *S. aureus* and is the only known route of uptake for both GSH and GSSG identified in this bacterium to date (Lensmire et al., 2023). However, under conditions where GSH utilisation is essential, the growth of a *gis*-deficient mutant strain is restored to near wild-type levels when GSH is supplied at concentrations greater than 0.5 mM (Lensmire et al., 2023). This implies the existence of at least one additional route of GSH uptake. To determine whether DtpT fulfils this role, isogenic mutants of *dtpT* and *gisD,* as well as a double mutant, were grown under conditions where glutathione was supplied as the sole sulphur source. At 50 μM GSH, we found that the growth of *S. aureus* is limited and solely dependent on the Gis system (Figure S4A). However, both single mutant strains demonstrated growth defects relative to the wild-type at 1 mM GSH, while growth was abolished for the double mutant under this condition (Figure 5C). In comparison, we found that growth in the presence of GSSG was solely dependent on Gis (Figure S4B). We also found that *in trans* complementation with *dtpT* was sufficient to fully restore growth of the *dtpT* single-mutant and partially restore growth in the double mutant strain in the presence of 1 mM GSH (Figure 5D). Finally, given that GSH oxidises spontaneously over time in aerated environments, we repeated the growth experiment with 1 mM GSH under anaerobic conditions to limit changes in the oxidation state of GSH during growth (Figure S4C). Again, we observed phenotypes which were consistent with those observed during aerobic growth.

303   Taken together, these observations demonstrate that both Gis and DtpT contribute to GSH

304   uptake at millimolar concentrations.

Figure 5 – DtpT transports reduced glutathione (GSH). (**A**) Chemical structure of GSH. (**B**) (**Left**) Transport of GSH and GSSG was assessed via pyranine transport assays. Transport assay curves are normalised immediately before the addition of valinomycin (red arrow) for ease of comparison. Curves show the mean of three replicates. (**Right**) summarised transport data. Bars show the mean of three replicates ± standard deviation. (**C - D**) Growth of *S. aureus* strain JE2 and mutant derivatives was assessed in CDMG – *s* supplemented with 1 mM GSH. In each case, strains were grown over 24 hours and OD_{600} was measured every 30 minutes for three biological replicates. Curves indicate the mean values ± standard deviation. (**C**) Growth of single-mutants of *dtpT* and *gisD* as well as a *dtpT gisD* double-mutant strain in CDMG – *s* ± 1 mM GSH. (**D**) *In-trans* expression of DtpT restores growth of the *dtpT* single mutant to wild-type levels and partially restores growth in the *dtpT gisD* double mutant. (**E**) Transport data for DtpT mutant variants against Gly-Gly, Cys-Gly and GSH, expressed as a percentage of the activity for the wild-type protein. Bars indicate the mean ± standard deviation. *$p<0.05$, **$p<0.01$, ***$p<0.001$; One-way ANOVA with Tukey's test.

In order to better understand the structural determinants underpinning GSH transport in DtpT, mutant variants were generated with alterations in residues predicted to be involved in substrate binding and transport (Figure 5E, Figure S5-6). Glu33 and Tyr41 form part of the conserved $E_{33}xxERFxYY_{41}$ motif found ubiquitously in POT transporters and are known to contribute toward proton coupling during the transport cycle (Aduri et al., 2015; Lichtinger et al., 2024; Solcan et al., 2012) (Figure S5). Consistent with this, acidification of the liposome lumen was severely reduced in an E33A mutant to levels equivalent to that observed for wild-type DtpT in the absence of substrate (Figure S6), demonstrating that Glu33 is essential for proton-coupled transport in this protein. Acidification was similarly reduced for all substrates in the Y41A mutant protein, though to a lesser extent (Figure 5E). Similarly, Asn167 is conserved across diverse POT proteins and has been observed to facilitate substrate binding via anchoring of the peptide N-terminal amide group (Kotov et al., 2023; Lyons et al., 2014) (Figure S5). As expected, the transport of gly-gly and cys-gly is restricted in an N167Q mutant likely due to substrate occlusion. Transport of GSH is also restricted in this mutant, providing evidence that Asn167 plays a key role in transporting this novel substrate.

Based on co-crystal structures of the *Streptococcus thermophilus* POT protein PepT$_{St}$ solved in complex with Ala-Phe and Ala-Ala-Ala, a dual-binding model has been proposed in which POTs differentially accommodate dipeptides and tripeptides during transport (Lyons et al., 2014). Briefly, these structures suggest that dipeptides bind in a lateral conformation (relative to the membrane plane), while tripeptides can be accommodated in an alternative vertical conformation in which the peptide extends deeper into the binding groove and interacts with residues adjacent to the extracellular gate. In this conformation, the carboxyl terminus of Ala-Ala-Ala is coordinated by interactions with Glu299 and Glu300. Gln310 in the *Staphylococcus hominis* POT PepT$_{Sh}$ (conserved in the *S. aureus* DtpT studied here, Figure S5) is orthologous to Glu299 and was similarly found to form part of a deep pocket involved in accommodating the thioalcohol moiety of a vertically bound S-Cys-Gly-3M3SH ligand (Minhas et al., 2018). Here, we observe that the transport of GSH is significantly reduced in a Q310A variant of DtpT (Figure 5E). Such an observation, therefore, implicates Gln310 in the coordination of GSH, suggesting that GSH may be accommodated in a similar position to S-Cys-Gly-3M3SH in PepT$_{Sh}$, and may indicate a unique mechanism for recognising sulphur-containing peptides.

Glutathione transport-deficient strains demonstrate reduced fitness during macrophage infection

Although *S. aureus* was historically considered an extracellular pathogen, recent work has demonstrated that this bacterium survives inside phagocytic cells, highlighting how its intracellular lifestyle may contribute to dissemination and immune evasion during systemic infection (Flannagan et al., 2016; Kubica et al., 2008). To date, there is a poor understanding of the molecular factors governing fitness in these intracellular environments, particularly regarding nutrient availability and acquisition. Cytosolic concentrations of GSH in human cells range from 1 to 10 mM depending on the cell type, making it the most abundant intracellular non-protein thiol (Cui et al., 2023; Meister, 1988). Based on this, we hypothesised that glutathione utilisation may contribute toward the fitness of *S. aureus* in intracellular environments.

Differentiated THP1 macrophage-like cells and human monocyte-derived macrophages (hMDMs) were each infected with either JE2 or mutant derivatives of *dtpT* and/or *gisD*, and bacterial survival was quantified over the first 8 hours post-infection (Figure 6). To account for any changes in fitness resulting from the inclusion of the *bursa aurealis* transposon insertion sequence, we also assessed the survival of a mutant strain carrying the transposon in a presumed phenotypically-null locus corresponding to a predicted IS3-family transposase gene (SAUSA300_0060). In THP-1 cells, survival of both the *dtpT* and *gisD* single-mutant strains was reduced slightly on-average at 8 hours but did not differ significantly when compared to the wild-type. In comparison, survival of the *dtpT gisD* double mutant strain was significantly lower at this time-point, being reduced more than two-fold relative to the wild-type (Figure 6A). Similarly, survival of both the *gisD* single-mutant and *dtpT gisD* double-mutant strains was significantly reduced at this time-point in hMDMs derived from eight independent donors (Figure 6B). Overall, these data suggest that GSH transport is required for complete fitness of *S. aureus* during macrophage infection, though the presence of either *dtpT* or *gisD* is seemingly sufficient to restore survival to near wild-type levels.

In eukaryotic systems, glutathione acts as an important cellular antioxidant and contributes to varied physiological processes including metal homeostasis and alleviation of lipid oxidation stress, ultimately contributing to immune cell function (Dasgupta et al., 2010; Lushchak, 2012). We, therefore, wondered whether bacterial sequestration of glutathione may serve as a strategy to disrupt healthy phagocyte function and thereby promote bacterial

survival. On average, we found no differences in cytotoxicity between the tested strains at 4

hours post-infection (Figure S7B). In contrast, we observed that THP1 cells infected with the

GSH transporter-null double-mutant strain demonstrated increased cell death at 8 hours

relative to the wild-type (Figure S7B). To investigate this further, we quantified the production

of inflammatory cytokines at 6 hours post-infection, representing the mid-point between

these two states. At this time point, cells infected with the double-mutant strain produced

significantly lower levels of both IL-1β and IL6, though production of TNFα was seemingly

unaffected (Figure S7C). While preliminary, these observations suggest that glutathione

transporters allow *S. aureus* to modulate the macrophage pro-inflammatory status and

maintain the intracellular niche by ultimately prolonging host-cell survival.

Figure 6. A glutathione transport-null double-mutant demonstrates reduced survival inside macrophages. (A - B) Bacterial survival of JE2 and mutant derivatives after 8 hours post-infection in THP1 macrophage-like cells (A; $n = 9$) and hMDM cells (B; $n = 8$ donors, each infected in biological triplicate). In each case, survival is calculated as a percentage of the total bacterial uptake, as determined at 30 minutes post-infection. In each case *p<0.05, **p<0.01, ***p<0.001; as determined by paired-sample t-test.

Discussion

The promiscuous nature of peptide transport proteins and their functional redundancy in many organisms have made these systems historically challenging to characterise regarding substrate range and preference. Molecular dynamics simulations and similar computational methods have been combined with biochemical and structural methods in order predict substrate preferences in POTs, but experimental evidence for the applicability of these models to biological systems is lacking (Kotov et al., 2023; Li et al., 2022; Samsudin et al., 2016). Here, we present an extensive cell-based analysis of peptide transporter function in *S. aureus* and apply biochemical analysis of DtpT to both verify and expand our functional understanding of this transporter. Our findings demonstrate that DtpT is highly promiscuous, evidenced by the considerable heterogeneity observed among putative transport targets identified here. While earlier work has focussed on the contribution of Opp3 toward the utilisation of host-derived peptides (Lehman et al., 2019), our data instead highlights how DtpT serves as a major route of uptake for diverse dipeptides and tripeptides in *S. aureus*, including GSH. Indeed, this broad substrate range may underpin the essentiality of DtpT previously observed in various systemic animal infection models (Coulter et al., 1998).

A number of recent studies have demonstrated how exogenous amino acids play diverse functional roles during *S. aureus* infections, serving both as key nutrient sources and environmental signals with defined regulatory consequences (Urso et al., 2024). Despite this, many studies overlook the potential contribution of host-derived oligopeptides as reservoirs for these amino acids. For instance, glutamate is known to serve as a major gluconeogenic carbon source for *S. aureus* and strains lacking the glutamate dehydrogenase GudB are attenuated under glucose-limited conditions (Halsey et al., 2017). Glutamate restriction has pleiotropic effects on cellular physiology, ultimately resulting in increased biofilm formation *in vitro* and *in vivo* (Shibamura-Fujiogi et al., 2022). Taken together, these observations implicate exogenous glutamate as an important factor contributing to both growth and environmental sensing in *S. aureus*. Acquisition of exogenous glutamate in *S. aureus* is primarily facilitated by the amino acid permease GltS (Zeden et al., 2020). Here, we demonstrate that DtpT is responsible for the utilisation of diverse glutamate-containing dipeptides and show that these peptides are specifically enriched among DtpT substrates (Figure 3A). These findings suggest that DtpT-mediated peptide uptake may serve as an additional source of exogenous glutamate in peptide-rich growth environments.

Our PM data reinforces the fact that DtpT serves as the primary route of dipeptide uptake in

S. aureus, but also reveals some overlap in substrate specificity between this transporter and

Opp3. The identification of a subset of peptides with redundant routes of internalisation may

point toward these peptides being of high value to the cell, perhaps serving as reservoirs for

important amino acids in relevant biological contexts. For instance, our data demonstrates

how diverse arginine-containing dipeptides are utilised by *S. aureus*, being significantly

enriched within the cluster of peptides utilised via either DtpT or Opp3 in our PM data (Figure

2; 3B). Consistent with this, *S. aureus* USA300 rapidly internalises arginine during growth but

converts neither proline nor glutamate to arginine when grown *in vitro*, making exogenous

arginine essential for growth (Halsey et al., 2017; Reslane et al., 2022). Experimental evidence

has demonstrated how arginine restriction stimulates antibiotic tolerance in *S. aureus*

biofilms, implicating arginine availability as an important environmental stimulus modulating

S. aureus cell behaviour (Freiberg et al., 2024). We propose that the redundancy observed

here for uptake of Arg-containing peptides likely signifies an evolutionary strategy to increase

the diversity of potential arginine sources available to the cell, which is in agreement with the

apparent importance of exogenous arginine in *S. aureus* physiology. Furthermore, the

presence of two Arg-peptide transporters with varying substrate affinities in *S. aureus* may

facilitate the utilisation of these peptides in varied growth environments with variable Arg-

containing peptide availability.

In a similar vein, DtpT is capable of transporting the glutamine-containing dipeptide Ala-Gln

(Figure 4C), but our PM data points toward the existence of a yet-unidentified additional

utilisation system which is specific for glutamine-containing peptides (Figure 3A). The

existence of such a system may indicate a yet underappreciated role for exogenous

glutamine in *S. aureus* physiology. The Opp4 transport system is a yet-uncharacterised Opp-

homologue which is widespread in *S. aureus* and is likely to have been generated via

duplication of Opp3 (Yu et al., 2014). Given their close evolutionary relationship, we propose

that Opp4 is likely to be a peptide transporter like Opp3 and is, therefore, a strong candidate

to fulfil the role of the alternative Gln-peptide transport system in *S. aureus*. While ACME Opp

may also contribute toward peptide transport, this system is more closely related to Opp1/2

and is, therefore, more likely to play a role in metal ion transport.

In contrast to the examples listed above, we demonstrate that tyrosine-containing dipeptides

are scarcely utilised by *S. aureus* (Figure 3C). This is perhaps unsurprising, given that *S.*

aureus JE2 lacks a complete catabolic pathway for tyrosine utilisation and does not rapidly

consume exogenous tyrosine when grown in defined media (Halsey et al., 2017). Intriguingly,
no DtpT-mediated transport was observed for the sole tyrosine-containing peptide (Val-Tyr)
assessed here (Figure 4B-C), providing evidence that DtpT possesses a distinct biochemical
preference toward peptides which are readily catabolised over those which lack the potential
to serve as a nutrient source. That being said, we do observe the utilisation of some
dipeptides exclusively composed of the branched-chain amino acids (i.e. leucine, isoleucine
and valine) and demonstrate that several such peptides are utilised via DtpT, despite the lack
of a known catabolic pathway for these amino acids in JE2 (Halsey et al., 2017). For instance,
we demonstrate that Leu-Leu is utilised robustly by JE2 and belongs to cluster 2, implicating
both DtpT and Opp3 in the transport of this peptide (Figure 2). Our findings, therefore,
suggest that branched chain amino acids derived from short peptides are catabolised by
JE2, though the catabolic pathways involved remain to be elucidated.

Glutathione is a ubiquitous metabolite in eukaryotic systems and is the predominant thiol
present in intracellular niches, as well as being enriched in other environments associated
with *S. aureus* infection including the sputum of cystic fibrosis patients (Shull et al., 2024). In
most biological contexts, GSH is the most abundant form of glutathione by a significant
margin (Hansen et al., 2009). Here, we identify GSH as a substrate of DtpT and demonstrate
that either DtpT or Gis-mediated transport is sufficient to meet the nutrient sulphur
requirements of *S. aureus* at physiologically relevant GSH concentrations (Figure 5C). To the
best of our knowledge, our findings provide the first example of a bacterial POT contributing
toward glutathione acquisition; whether or not this function is conserved in the wider POT
superfamily remains unclear, especially given that Gln310 (or Glu in the equivalent position)
is required for strong transport of GSH but this feature is not conserved across bacterial POT
proteins (Figure 5E; Figure S5).

Generally, binding-protein dependant transport systems (including ABC transporters)
operate with high affinities, whilst POTs and similar MFS systems have been reported as
having substrate affinities in the high-micromolar or low millimolar range (Kotov et al., 2023;
Samsudin et al., 2016). We, therefore, predict that Gis may serve as the dominant system
involved in GSH scavenging under conditions where the metabolite is scarce, whilst both
systems likely contribute to transport in GSH-enriched environments. Indeed, the
phenotypes associated with our mutant strains support such a model (Figure 5C; Figure S4).

Finally, we identify *S. aureus* glutathione transporters as important fitness determinants

during intracellular growth inside macrophages, representing an environment in which GSH

is present at millimolar concentrations (Figure 5A-B). In *Mycobacterium tuberculosis* (Mtb),

glutathione transport has been implicated in regulating the activity of Mtb-infected

macrophages and thereby contributing toward intracellular survival and immune evasion

(Dasgupta et al., 2010). Our macrophage infection data similarly demonstrates how the

presence of glutathione transporters affects host-cell behaviour, seemingly allowing the

pathogen to modulate innate immune signalling responses and subvert premature host-cell

death (Figure S7). Hence, glutathione transport may represent a conserved strategy utilised

by intracellular bacterial pathogens in order to enable long-term intracellular survival, thereby

avoiding exposure to extracellular stresses and immune clearance. More work is needed to

better understand how the host-cell response is modified as a result of bacterial glutathione

transport and determine whether this response is conserved across different host-cell types.

Overall, this work has significantly expanded our knowledge of peptide utilisation in *S. aureus*

and shed light on a novel role for glutathione transport in host-pathogen interactions during

intracellular infection. However, with the diversity of roles ascribed to peptide transporters in

other organisms – including cell-wall turnover (Maqbool et al., 2011; Park et al., 1998;

Simpson et al., 2023), antimicrobial peptide resistance (Parra-Lopez et al., 1993; Rivera et

al., 2024) and cell signalling (Dubois et al., 2019; Nasher et al., 2018; Zheng et al., 2018) – it

is clear that this is an area of *S. aureus* biology where much remains to be discovered and

understood.

Methods

Bacterial strains and growth conditions

All *S. aureus* strains used in this study were routinely cultured in tryptic soy broth (TSB) with shaking or on solid tryptic soy agar (TSA) medium at 37°C. *E. coli* strains were routinely cultured in lysogeny broth (LB) with shaking or on LB agar at 37°C. Selective antibiotics were added during growth when appropriate at the following concentrations: 50 µg/ml kanamycin, 100 µg/ml ampicillin, 10 µg/ml erythromycin, 10 µg/ml chloramphenicol.

S. aureus JE2 transposon mutants were acquired from the Nebraska Transposon Mutant Library (NTML; (Fey et al., 2013) and transduced into wild-type JE2 via Φ85 bacteriophage in order minimise off-target genetic inconsistencies between strains. To generate double mutant strains, the Erm^R resistance cassette was exchanged with an unmarked sequence using the allelic exchange vector pTnT, as described previously (Bose et al., 2013). Plasmids were generated using the ClonExpress II One Step Cloning Kit (Vazyme) according to manufacturer's instructions. DtpT complementation in *S. aureus* mutant strains was achieved via introduction of the entire *dtpT* locus into the *E. coli* / *S. aureus* shuttle vector pSK5630 modified by the addition of the native *dtpT* promoter sequence (herein referred to as pSK56$_{pT}$) (Grkovic et al., 2003).

A full list of bacterial strains utilised in this work are provided in supplemental table ST1. Primers utilised for diagnostic PCR, cloning and sequencing of genetic elements are provided in supplemental table ST2.

Phenotype microarrays and analysis

Peptide utilisation by *S. aureus* strain JE2 and mutant derivatives was assessed on phenotype microarray plates PM 6-8 (Biolog) using a modified version of the manufacturer's protocol. Cells grown on TSA overnight were transferred from solid media into 3 ml IF-0α buffer using a sterile cotton swab and pelleted. Cells were washed twice and resuspended in IF-0α. This cell suspension was then diluted to a final OD_{600} of 0.004 in 9.975 ml IF-0α and supplemented with 110 µl redox dye H (Biolog) and 915 µl 12X PM additive solution (240 mM tricarballylic acid, pH 7.1; 24 mM $MgCl_2$; 12 mM $CaCl_2$; 150 µM L-cystine, pH 8.5; 0.06% w/v yeast extract; 0.06% v/v tween 80; 6 mM D-glucose; 12 mM pyruvate). PM plates were inoculated with 100 µl/well of the prepared cell suspension and incubated for 36 hours at

37°C with continuous slow shaking in an Epoch2 microplate spectrophotometer (Agilent).

OD_{590} and OD_{750} were measured every 15 minutes.

Signal curves (OD_{590} - OD_{750} against time) for all four strains were grouped for each peptide

and then analysed by Uniform Manifold Approximation and Projection (UMAP) using the

UMAP R library and a random seed of 888 (McInnes et al., 2018). K-means clustering was

used on the same grouped data with "kmeans" R function and k=7. Empirical area under the

curve (eAUC) was calculated using Growthcurver (Sprouffske & Wagner, 2016).

Peptide utilisation assays

Utilisation of specific peptides was assessed using glucose-supplemented chemically

defined medium (CDMG) based on a previously reported methodology (Hussain et al., 1991).

The complete composition of the CDMG used here is provided in supplemental table ST3.

For assessing the utilisation of arginine-containing peptides, L-arginine was omitted from the

medium (CDMG - *R*). For assessing the utilisation of glutathione, L-cysteine and L-methionine

were omitted from the medium and $(NH_4)_2SO_4 \cdot FeSO_4 \cdot 6H_2O$ was substituted for 15 µM iron

(III) citrate (CDMG - *s*). Reduced glutathione solutions were prepared under anaerobic

conditions to prevent changes in the oxidation state before commencing assays. Overnight

cultures of *S. aureus* strain JE2 and mutant derivatives were diluted 1:100 into fresh TSB and

grown to mid-exponential phase. Cells were then pelleted and washed twice in the assay

medium (CDMG - *R* or CDMG - *s*) before being resuspended and diluted to a final OD_{600} of

0.05 in peptide-supplemented medium. 200 µl of each cell suspension was transferred to a

Costar 96 well plate (Corning) and incubated for 24 hours at 37°C with shaking. OD_{600} was

measured every 30 minutes.

For anaerobic growth assays, cultures were prepared as before being transferred to a Whitley

A85 anaerobic workstation (Don Whitley Scientific) and diluted into assay medium which had

been equilibrated under anaerobic conditions for at least 3 hours and supplemented with 100

mM sodium nitrate. Growth was monitored as above in a Stratus 96 well plate reader (Cerillo)

for 18 hours at 37°C without shaking.

Protein purification

The DtpT open reading frame was amplified directly from the *S. aureus* JE2 genome and

cloned into the C-terminal octa-histidine/GFP-tagged fusion vector pWaldo (Drew et al.,

2006) before being transformed into *E. coli* C43(DE3) (Miroux & Walker, 1996). Mutant

variants were generated via site-directed PCR mutagenesis and blunt-end ligation with T4

DNA ligase (NEB) and all constructs were verified by sequencing.

Cells grown overnight in LB supplemented with kanamycin were diluted 1:100 into fresh

media. Cultures were incubated at 37°C until an OD_{600} of 0.5 was reached. Expression was

then induced by the addition of IPTG to a final concentration of 100 µM. Cultures were

incubated overnight at 25°C with shaking. Cells were then pelleted, resuspended in 50 ml

resuspension buffer (40 mM Tris, pH 7.5; 10% glycerol) and lysed by sonication. Crude

lysates were clarified via centrifugation at 27,000 x g for 30 minutes. Membranes were then

pelleted via ultracentrifugation at 164000 x g for 120 minutes. Membrane pellets were pooled

and resuspended in 40 ml protein solubilisation buffer (40 mM Tris, pH 7.5; 200 mM NaCl; 20

mM imidazole; 10% glycerol; 0.5% n-dodecyl-B-pyromaltoside (DDM)) using a glass

homogenizer before being incubated for 60 minutes at 4°C with rolling. The resultant

suspension was then ultracentrifuged as before for 60 minutes to pellet leftover membranes

and insoluble contaminants. Supernatant containing the solubilised membrane proteins was

pooled and applied to a His-TRAP nickel affinity column (GE Healthcare) using an ÄKTA

Protein Purification System (Cytiva). This column was washed with 50 ml column wash buffer

(40 mM Tris, pH 7.5; 200 mM NaCl; 20 mM imidazole; 10% glycerol; 0.04% DDM) and the

fusion protein was then eluted via application of column elution buffer (40 mM Tris, pH 7.5;

200 mM NaCl; 400 mM imidazole; 10% glycerol; 0.04% DDM).

Purified protein was exchanged into TEV reaction buffer (40 mM Tris, pH 7.5; 200 mM NaCl;

0.5 mM EDTA; 1 mM DTT; 10% glycerol; 0.04% DDM) using a HiTRAP desalting column

(SLS). Protein was then mixed with TEV protease to a final ratio of approximately 3:1. The

resultant reaction mixture was incubated overnight at 4°C. Pure DtpT was isolated by

applying the reaction mixture to a His-TRAP nickel affinity column as before and washing

with column wash buffer, this time collecting the flow-through fractions. Pure protein was

stored at -70°C.

Reconstitution into proteoliposomes and transport assays

Purified DtpT was exchanged into reconstitution buffer (20 mM Tris pH 7.5; 150 mM NaCl;

0.03% DDM) and diluted to a final concentration of 0.5 mg/ml. Liposomes (composed of

POPE and POPG in a 3:1 ratio) were prepared and extruded 11 times each through pre-

soaked 0.8 µm and 0.4 µm track-etched polycarbonate membranes (Whatman) before

diluting to a final concentration of 10 mg/ml in liposome buffer (50 mM KPi, pH 7).

10 mg lipid mixture was gradually added to 200 µg protein at room temperature before being

incubated on ice for 60 minutes. 100 µl SM2 Biobeads (Biorad) suspended in water were

added to the lipid-protein mixture, which was then incubated for 1 hour at 4°C while rotating.

A further 130 µl Biobeads were added and the mixture was incubated as before for a further

3 hours. Biobeads were pelleted and the lipid-protein mixture was transferred to a fresh tube.

160 µl Biobeads were added and the mixture was incubated as before for a further 16 hours.

Biobeads were removed as before and proteoliposomes were pelleted via ultracentrifugation

at 158 000 × g for 30 minutes. Proteoliposomes were then resuspended in liposome buffer

to a final protein concentration of 0.5 mg/ml (assuming 100% reconstitution efficiency) and

subjected to three freeze-thaw cycles before dialysing extensively against liposome buffer at

4°C. Proteoliposomes were pelleted as before and resuspended in liposome buffer before

being stored at -70°C.

For transport assays, proteoliposomes were pelleted as before and resuspended in inside

buffer (5 mM HEPES, pH 6.8; 120 mM KCl; 2 mM $MgSO_4$) spiked with 1 mM pyranine.

Proteoliposomes were subjected to seven freeze-thaw cycles before being extruded through

a pre-soaked 0.4 µm track-etched polycarbonate membrane. Proteoliposomes were then

pelleted at 20°C and resuspended in inside buffer. Excess pyranine was removed using a

microspin G25 desalting column (Cytiva) according to the manufacturer's protocol.

Liposomes were finally pelleted again at 20°C and resuspended to a final protein

concentration of 1 mg/ml. For each assay, 6 µg of proteoliposomes were diluted 1:100 into

outside buffer (5 mM HEPES, pH 6.8; 120 mM NaCl; 2 mM $MgSO_4$). Assays were performed

in 1 ml clear plastic microcuvettes using a Fluoromax-4 spectrophotometer (Horiba) with a

magnetic flee for stirring. Fluorescence of pyranine was measured via excitation at 460 nm

and emission at 510 nm. Substrate was added to the reaction at t = 25 s and transport was

induced by the addition of 1 µM valinomycin at t = 50 s.

Macrophage culture and differentiation

THP1 cells (ATCC) were cultured in macrophages media (RPMI1640 supplemented with 10%

Fetal Calf Serum and Glutamine). To obtain a macrophage-like differentiation state, THP1

were differentiated for 48 hrs in macrophage media supplemented with 50 ng/mL Phorbol

12-myristate 13-acetate (in DMSO, Sigma-Aldrich). PMA-containing media was replaced with

fresh macrophage media and cells were rested overnight prior to infection.

Human primary macrophages were differentiated from human peripheral blood mononuclear

cells (PBMCs) obtained from the NHS (University of York Biology ethic committee project

DB202111) and differentiated in complete macrophage media supplemented with 50 ng/mL

macrophage colony-stimulating factor (mCSF, Proteintech) over 7 days.

Macrophage infection

Infection of macrophages and macrophage-like cell lines was carried out based on previously

published methods (Alves et al., 2024). Overnight cultures of *S. aureus* strain JE2 and mutant

derivatives were diluted 1:100 into fresh TSB and grown to mid-exponential phase. Cells

were then pelleted and washed twice in ice-cold phosphate-buffered saline (PBS) before

being resuspended to a final cell density of 5×10^7 CFU/ml. For infection, 10 µl bacterial

suspension was added to 50,000 hMDM or THP-1 cells (D.O.I. = 10) in 90 µl Opti-MEM

reduced serum media (Thermo Fisher) and plates were centrifuged for 5 minutes at 500 x g

to promote contact. After 30 minutes of incubation at 37°C, the medium was exchanged for

Opti-MEM supplemented with 100 µg/ml gentamicin. The addition of gentamicin was

designated t = 0 h. After a further 30 minutes of incubation this medium was removed and

cells were then maintained in Opti-MEM supplemented with 5 µg/ml gentamicin (Gibc).

To compare bacterial internalisation and survival, cells at t = 0.5 / 8 h were incubated in

deionised water for 10 minutes before being lysed by repeated pipetting. Suspensions were

serially diluted in PBS, spotted onto TSA and incubated overnight at 37°C to determine viable

CFU. Cytotoxicity was measured by quantifying the activity of lactate dehydrogenase (LDH)

in the extracellular medium (Cytotox 96 non-radioactive cytotoxicity kit, Promega) and

normalised against cells lysed by 0.04% Triton X-100. Cytokine levels in cell culture medium

were determined at t = 6 h by utilising human IL 1-β, IL 6 and TNF-α ELISA assay kits

(invitrogen) according to the manufacturer's protocols.

Author contributions

Imran Khan: Conceptualization, Formal analysis, Investigation, Methodology, Data processing, Writing – original draft, review & editing. **Sandy MacDonald**: Formal analysis, Data Processing. **Sigurbjorn Markusson**: Investigation. **Callum Robson:** Investigation. **Joanne L Parker**: Conceptualization, Methodology. **Simon Newstead**: Conceptualization, Methodology. Writing – review & editing. **Dave Boucher**: Conceptualization, Investigation, Methodology, Writing – review & editing. **Marjan Van Der Woude**: Conceptualization, Methodology, Supervision, Project administration, Funding acquisition, Writing – review & editing. **Gavin H Thomas**: Conceptualization, Methodology, Supervision, Project administration, Funding acquisition, Writing – review & editing.

Acknowledgements

We are extremely grateful to the Burgess family for the award of a gift to the University of York that supported a PhD studentship to IK. We thank Dr Rebecca M. Corrigan (ORCiD: 0000-0002-6031-1148) for supplying the wild-type JE2 and NTML strains utilised in this study, and for providing valuable technical guidance. We acknowledge ongoing support of the GHT lab through BB/X003035/1.

References

1. Ackroyd, B. K., Dodson, E. J., Mehboob, J., Dowle, A. A., Thomas, G. H., & Wilkinson, A. J.
(2024). Structure and ligand binding in the putative anti-microbial peptide transporter
protein, YejA. *Microbiology (Reading, England)*, *170*(2).

2. Adolf, L. A., Müller-Jochim, A., Kricks, L., Puls, J.-S., Lopez, D., Grein, F., & Heilbronner, S.
(2023). Functional membrane microdomains and the hydroxamate siderophore transporter
ATPase FhuC govern Isd-dependent heme acquisition in Staphylococcus aureus. *Elife*, *12*.

3. Aduri, N. G., Prabhala, B. K., Ernst, H. A., Jørgensen, F. S., Olsen, L., & Mirza, O. (2015). Salt
bridge swapping in the EXXERFXYY motif of proton-coupled oligopeptide transporters. *The*
Journal of Biological Chemistry, *290*(50), 29931–29940.

4. Alves, J., Vrieling, M., Ring, N., Yebra, G., Pickering, A., Prajsnar, T. K., Renshaw, S. A., &
Fitzgerald, J. R. (2024). Experimental evolution of Staphylococcus aureus in macrophages:
dissection of a conditional adaptive trait promoting intracellular survival. *MBio*, *15*(6),
e0034624.

5. Bochner, B. R., Gadzinski, P., & Panomitros, E. (2001). Phenotype microarrays for high-
throughput phenotypic testing and assay of gene function. *Genome Research*, *11*(7), 1246–
1255.

6. Borezée-Durant, E., Hiron, A., Piard, J.-C., & Juillard, V. (2009). Dual role of the oligopeptide
permease Opp3 during growth of Staphylococcus aureus in milk. *Applied and Environmental*
Microbiology, *75*(10), 3355–3357.

7. Bose, J. L., Fey, P. D., & Bayles, K. W. (2013). Genetic tools to enhance the study of gene
function and regulation in Staphylococcus aureus. *Applied and Environmental Microbiology*,
79(7), 2218–2224.

8. Coulter, S. N., Schwan, W. R., Ng, E. Y., Langhorne, M. H., Ritchie, H. D., Westbrock-Wadman,
S., Hufnagle, W. O., Folger, K. R., Bayer, A. S., & Stover, C. K. (1998). Staphylococcus aureus
genetic loci impacting growth and survival in multiple infection environments. *Molecular*
Microbiology, *30*(2), 393–404.

9.   Cui, C.-Y., Li, B., & Su, X.-C. (2023). Real-time monitoring of the level and activity of

697   intracellular glutathione in live cells at atomic resolution by 19F-NMR. *ACS Central Science*,

698   *9*(8), 1623–1632.

699   10.   Dasgupta, A., Sureka, K., Mitra, D., Saha, B., Sanyal, S., Das, A. K., Chakrabarti, P., Jackson,

700   M., Gicquel, B., Kundu, M., & Basu, J. (2010). An oligopeptide transporter of Mycobacterium

701   tuberculosis regulates cytokine release and apoptosis of infected macrophages. *PloS One*,

702   *5*(8), e12225.

703   11.   Detmers, F. J., Lanfermeijer, F. C., Abele, R., Jack, R. W., Tampe, R., Konings, W. N., &

704   Poolman, B. (2000). Combinatorial peptide libraries reveal the ligand-binding mechanism of

705   the oligopeptide receptor OppA of Lactococcus lactis. *Proceedings of the National Academy*

706   *of Sciences of the United States of America*, *97*(23), 12487–12492.

707   12.   Doeven, M. K., Abele, R., Tampé, R., & Poolman, B. (2004). The binding specificity of OppA

708   determines the selectivity of the oligopeptide ATP-binding cassette transporter. *The Journal*

709   *of Biological Chemistry*, *279*(31), 32301–32307.

710   13.   Drew, D., Lerch, M., Kunji, E., Slotboom, D.-J., & de Gier, J.-W. (2006). Optimization of

711   membrane protein overexpression and purification using GFP fusions. *Nature Methods*, *3*(4),

712   303–313.

713   14.   Dubois, T., Lemy, C., Perchat, S., & Lereclus, D. (2019). The signaling peptide NprX controlling

714   sporulation and necrotrophism is imported into Bacillus thuringiensis by two oligopeptide

715   permease systems. *Molecular Microbiology*, *112*(1), 219–232.

716   15.   Fey, P. D., Endres, J. L., Yajjala, V. K., Widhelm, T. J., Boissy, R. J., Bose, J. L., & Bayles, K. W.

717   (2013). A genetic resource for rapid and comprehensive phenotype screening of

718   nonessential Staphylococcus aureus genes. *MBio*, *4*(1), e00537-12.

719   16.   Flannagan, R. S., Heit, B., & Heinrichs, D. E. (2016). Intracellular replication of Staphylococcus

720   aureus in mature phagolysosomes in macrophages precedes host cell death, and bacterial

721   escape and dissemination. *Cellular Microbiology*, *18*(4), 514–535.

722   17.   Freiberg, J. A., Reyes Ruiz, V. M., Gimza, B. D., Murdoch, C. C., Green, E. R., Curry, J. M.,

723   Cassat, J. E., & Skaar, E. P. (2024). Restriction of arginine induces antibiotic tolerance in

724   Staphylococcus aureus. *Nature Communications*, *15*(1), 6734.

725   18.   Garai, P., Chandra, K., & Chakravortty, D. (2017). Bacterial peptide transporters: Messengers

726   of nutrition to virulence. *Virulence*, *8*(3), 297–309.

727   19.   Grkovic, S., Brown, M. H., Hardie, K. M., Firth, N., & Skurray, R. A. (2003). Stable low-copy-

728   number Staphylococcus aureus shuttle vectors. *Microbiology*, *149*(Pt 3), 785–794.

729   20.   Halsey, C. R., Lei, S., Wax, J. K., Lehman, M. K., Nuxoll, A. S., Steinke, L., Sadykov, M., Powers,

730   R., & Fey, P. D. (2017). Amino acid catabolism in staphylococcus aureus and the function of

731   carbon catabolite repression. *MBio*, *8*(1).

732   21.   Handke, L. D., Gribenko, A. V., Timofeyeva, Y., Scully, I. L., & Anderson, A. S. (2018). MntC-

733   dependent manganese transport is essential for staphylococcus aureus oxidative stress

734   resistance and virulence. *MSphere*, *3*(4).

735   22.   Hansen, R. E., Roth, D., & Winther, J. R. (2009). Quantifying the global cellular thiol-disulfide

736   status. *Proceedings of the National Academy of Sciences of the United States of America*,

737   *106*(2), 422–427.

738   23.   Hebbeln, P., & Eitinger, T. (2004). Heterologous production and characterization of bacterial

739   nickel/cobalt permeases. *FEMS Microbiology Letters*, *230*(1), 129–135.

740   24.   Hiron, A., Borezée-Durant, E., Piard, J.-C., & Juillard, V. (2007). Only one of four oligopeptide

741   transport systems mediates nitrogen nutrition in Staphylococcus aureus. *Journal of*

742   *Bacteriology*, *189*(14), 5119–5129.

743   25.   Hughes, A. M., Darby, J. F., Dodson, E. J., Wilson, S. J., Turkenburg, J. P., Thomas, G. H., &

744   Wilkinson, A. J. (2022). Peptide transport in Bacillus subtilis - structure and specificity in the

745   extracellular solute binding proteins OppA and DppE. *Microbiology*, *168*(12).

746   26.   Hussain, M., Hastings, J. G., & White, P. J. (1991). A chemically defined medium for slime

747   production by coagulase-negative staphylococci. *Journal of Medical Microbiology*, *34*(3),

143–147.

27.   Jeong, B., Shah, M. A., Roh, E., Kim, K., Park, I., & Bae, T. (2022). Staphylococcus aureus does

not synthesize arginine from proline under physiological conditions. *Journal of Bacteriology*,

204(6), e0001822.

28.   Kotov, V., Killer, M., Jungnickel, K. E. J., Lei, J., Finocchio, G., Steinke, J., Bartels, K., Strauss, J.,

Dupeux, F., Humm, A.-S., Cornaciu, I., Márquez, J. A., Pardon, E., Steyaert, J., & Löw, C.

(2023). Plasticity of the binding pocket in peptide transporters underpins promiscuous

substrate recognition. *Cell Reports*, *42*(8), 112831.

29.   Kubica, M., Guzik, K., Koziel, J., Zarebski, M., Richter, W., Gajkowska, B., Golda, A., Maciag-

Gudowska, A., Brix, K., Shaw, L., Foster, T., & Potempa, J. (2008). A potential new pathway

for staphylococcus aureus dissemination: The silent survival of S. aureus phagocytosed by

human monocyte-derived macrophages. *PloS One*, *3*(1), e1409.

30.   Lehman, M. K., Nuxoll, A. S., Yamada, K. J., Kielian, T., Carson, S. D., & Fey, P. D. (2019).

Protease-mediated growth of Staphylococcus aureus on host proteins is opp3 dependent.

MBio, *10*(2).

31.   Lehman, M. K., Sturd, N. A., Razvi, F., Wellems, D. L., Carson, S. D., & Fey, P. D. (2023).

Proline transporters ProT and PutP are required for Staphylococcus aureus infection. *PLoS*

Pathogens, *19*(1), e1011098.

32.   Lensmire, J. M., Dodson, J. P., Hsueh, B. Y., Wischer, M. R., Delekta, P. C., Shook, J. C.,

Ottosen, E. N., Kies, P. J., Ravi, J., & Hammer, N. D. (2020). The staphylococcus aureus cystine

transporters TcyABC and TcyP facilitate nutrient sulfur acquisition during infection. *Infection*

and Immunity, *88*(3).

33.   Lensmire, J. M., Wischer, M. R., Kraemer-Zimpel, C., Kies, P. J., Sosinski, L., Ensink, E.,

Dodson, J. P., Shook, J. C., Delekta, P. C., Cooper, C. C., Havlichek, D. H., Jr, Mulks, M. H.,

Lunt, S. Y., Ravi, J., & Hammer, N. D. (2023). The glutathione import system satisfies the

Staphylococcus aureus nutrient sulfur requirement and promotes interspecies competition.

PLoS Genetics, *19*(7), e1010834.

34.   Li, C., Yue, Z., Newstead, S., & Voth, G. A. (2022). Proton coupling and the multiscale kinetic

mechanism of a peptide transporter. *Biophysical Journal*, *121*(12), 2266–2278.

35.   Lichtinger, S. M., Parker, J. L., Newstead, S., & Biggin, P. C. (2024). The mechanism of

mammalian proton-coupled peptide transporters. *ELife*, *13*.

36.   Lushchak, V. I. (2012). Glutathione homeostasis and functions: potential targets for medical

interventions. *Journal of Amino Acids*, *2012*, 736837.

37.   Lyons, J. A., Parker, J. L., Solcan, N., Brinth, A., Li, D., Shah, S. T. A., Caffrey, M., & Newstead,

S. (2014). Structural basis for polyspecificity in the POT family of proton-coupled

oligopeptide transporters. *EMBO Reports*, *15*(8), 886–893.

38.   Maqbool, A., Levdikov, V. M., Blagova, E. V., Hervé, M., Horler, R. S. P., Wilkinson, A. J., &

Thomas, G. H. (2011). Compensating stereochemical changes allow murein tripeptide to be

accommodated in a conventional peptide-binding protein. *The Journal of Biological*

Chemistry, *286*(36), 31512–31521.

39.   McInnes, L., Healy, J., & Melville, J. (2018). UMAP: Uniform Manifold Approximation and

Projection for Dimension Reduction. *ArXiv [Stat.ML]*. arXiv. http://arxiv.org/abs/1802.03426

40.   Meister, A. (1988). Glutathione metabolism and its selective modification. *The Journal of*

Biological Chemistry, *263*(33), 17205–17208.

41.   Minhas, G. S., Bawdon, D., Herman, R., Rudden, M., Stone, A. P., James, A. G., Thomas, G. H.,

& Newstead, S. (2018). Structural basis of malodour precursor transport in the human axilla.

ELife, *7*.

42.   Miroux, B., & Walker, J. E. (1996). Over-production of proteins in Escherichia coli: mutant

hosts that allow synthesis of some membrane proteins and globular proteins at high levels.

Journal of Molecular Biology, *260*(3), 289–298.

43.   Nasher, F., Heller, M., & Hathaway, L. J. (2018). Streptococcus pneumoniae Proteins AmiA,

AliA, and AliB Bind Peptides Found in Ribosomal Proteins of Other Bacterial Species.

Frontiers in Microbiology, *8*.

44.   Park, J. T., Raychaudhuri, D., Li, H., Normark, S., & Mengin-Lecreulx, D. (1998). MppA, a

802   periplasmic binding protein essential for import of the bacterial cell wall peptide L-alanyl-

803   gamma-D-glutamyl-meso-diaminopimelate. *Journal of Bacteriology*, *180*(5), 1215–1223.

804   45.   Parker, J. L., Mindell, J. A., & Newstead, S. (2014). Thermodynamic evidence for a dual

805   transport mechanism in a POT peptide transporter. *ELife*, *3*.

806   https://doi.org/10.7554/elife.04273

807   46.   Parra-Lopez, C., Baer, M. T., & Groisman, E. A. (1993). Molecular genetic analysis of a locus

808   required for resistance to antimicrobial peptides in Salmonella typhimurium. *EMBO J.*,

809   *12*(11), 4053–4062.

810   47.   Pletzer, D., Lafon, C., Braun, Y., Köhler, T., Page, M. G. P., Mourez, M., & Weingart, H. (2014).

811   High-throughput screening of dipeptide utilization mediated by the ABC transporter

812   DppBCDF and its substrate-binding proteins DppA1-A5 in Pseudomonas aeruginosa. *PloS*

813   *One*, *9*(10), e111311.

814   48.   Remy, L., Carrière, M., Derré-Bobillot, A., Martini, C., Sanguinetti, M., & Borezée-Durant, E.

815   (2013). The Staphylococcus aureus Opp1 ABC transporter imports nickel and cobalt in zinc-

816   depleted conditions and contributes to virulence. *Molecular Microbiology*, *87*(4), 730–743.

817   49.   Reslane, I., Halsey, C. R., Stastny, A., Cabrera, B. J., Ahn, J., Shinde, D., Galac, M. R., Sladek,

818   M. F., Razvi, F., Lehman, M. K., Bayles, K. W., Thomas, V. C., Handke, L. D., & Fey, P. D.

819   (2022). Catabolic ornithine carbamoyltransferase activity facilitates growth of

820   Staphylococcus aureus in defined medium lacking glucose and arginine. *MBio*, *13*(3),

821   e0039522.

822   50.   Rivera, K. G., Tanaka, K. J., Buechel, E. R., Origel, O., Jr, Harrison, A., Mason, K. M., & Pinkett,

823   H. W. (2024). Antimicrobial peptide recognition motif of the substrate binding protein SapA

824   from nontypeable Haemophilus influenzae. *Biochemistry*, *63*(3), 294–311.

825   51.   Samsudin, F., Parker, J. L., Sansom, M. S. P., Newstead, S., & Fowler, P. W. (2016). Accurate

826   prediction of ligand affinities for a proton-dependent oligopeptide transporter. *Cell Chemical*

827   *Biology*, *23*(2), 299–309.

52.   Shaw, L., Golonka, E., Potempa, J., & Foster, S. J. (2004). The role and regulation of the

extracellular proteases of Staphylococcus aureus. *Microbiology*, *150*(Pt 1), 217–228.

53.   Shibamura-Fujiogi, M., Wang, X., Maisat, W., Koutsogiannaki, S., Li, Y., Chen, Y., Lee, J. C., &

Yuki, K. (2022). GltS regulates biofilm formation in methicillin-resistant Staphylococcus

aureus. *Communications Biology*, *5*(1), 1284.

54.   Shull, L. M., Wolter, D. J., Kunkle, D. E., Legg, K. A., Giedroc, D. P., Skaar, E. P., Hoffman, L. R.,

& Reniere, M. L. (2024). Analysis of genetic requirements and nutrient availability for

Staphylococcus aureus growth in cystic fibrosis sputum. *BioRxiv.Org: The Preprint Server for*

Biology.

55.   Simpson, B. W., Gilmore, M. C., McLean, A. B., Cava, F., & Trent, M. S. (2023). Escherichia coli

utilizes multiple peptidoglycan recycling permeases with distinct strategies of recycling.

Proceedings of the National Academy of Sciences of the United States of America, *120*(44),

e2308940120.

56.   Slamti, L., & Lereclus, D. (2019). The oligopeptide ABC-importers are essential

communication channels in Gram-positive bacteria. *Research in Microbiology*, *170*(8), 338–

344.

57.   Smith, M. W., Tyreman, D. R., Payne, G. M., Marshall, N. J., & Payne, J. W. (1999). Substrate

specificity of the periplasmic dipeptide-binding protein from Escherichia coli: experimental

basis for the design of peptide prodrugs. *Microbiology*, *145 (Pt 10)*(10), 2891–2901.

58.   Solcan, N., Kwok, J., Fowler, P. W., Cameron, A. D., Drew, D., Iwata, S., & Newstead, S.

(2012). Alternating access mechanism in the POT family of oligopeptide transporters:

Alternating access mechanism in the POT family. *The EMBO Journal*, *31*(16), 3411–3421.

59.   Sprouffske, K., & Wagner, A. (2016). Growthcurver: an R package for obtaining interpretable

metrics from microbial growth curves. *BMC Bioinformatics*, *17*(1), 172.

60.　Tong, S. Y. C., Davis, J. S., Eichenberger, E., Holland, T. L., & Fowler, V. G., Jr. (2015).

Staphylococcus aureus infections: epidemiology, pathophysiology, clinical manifestations,

and management. *Clinical Microbiology Reviews*, *28*(3), 603–661.

61.　Urso, A., Monk, I. R., Cheng, Y.-T., Predella, C., Wong Fok Lung, T., Theiller, E. M., Boylan, J.,

Perelman, S., Baskota, S. U., Moustafa, A. M., Lohia, G., Lewis, I. A., Howden, B. P., Stinear, T.

P., Dorrello, N. V., Torres, V., & Prince, A. S. (2024). Staphylococcus aureus adapts to exploit

collagen-derived proline during chronic infection. *Nature Microbiology*, *9*(10), 2506–2521.

62.　Wang, Z., Bie, P., Cheng, J., Lu, L., Cui, B., & Wu, Q. (2016). The ABC transporter YejABEF is

required for resistance to antimicrobial peptides and the virulence of Brucella melitensis.

Scientific Reports, *6*, 31876.

63.　Yu, D., Pi, B., Yu, M., Wang, Y., Ruan, Z., Feng, Y., & Yu, Y. (2014). Diversity and evolution of

oligopeptide permease systems in staphylococcal species. *Genomics*, *104*(1), 8–13.

64.　Zeden, M. S., Kviatkovski, I., Schuster, C. F., Thomas, V. C., Fey, P. D., & Gründling, A. (2020).

Identification of the main glutamine and glutamate transporters in Staphylococcus aureus

and their impact on c-di-AMP production. *Molecular Microbiology*, *113*(6), 1085–1100.

65.　Zheng, F., Shao, Z.-Q., Hao, X., Wu, Q., Li, C., Hou, H., Hu, D., Wang, C., & Pan, X. (2018).

Identification of oligopeptide-binding protein (OppA) and its role in the virulence of

Streptococcus suis serotype 2. *Microbial Pathogenesis*, *118*, 322–329.

870