Aldh3a1-mediated detoxification of reactive aldehydes contributes to distinct muscle responses to Amyotrophic Lateral Sclerosis progression

Ang Li[1*], Li Dong[1], Xuejun Li[1], Jianxun Yi[1], Jianjie Ma[2], Jingsong Zhou[1*]

[1] Department of Kinesiology, College of Nursing and Health Innovation, University of Texas at Arlington, TX, 76019, USA

[2] Department of Surgery, Division of Surgical Sciences, University of Virginia, Charlottesville, VA, 22903, USA

* Corresponding author

Abstract

Different muscles exhibit varied susceptibility to degeneration in Amyotrophic Lateral Sclerosis (ALS), a fatal neuromuscular disorder. Extraocular muscles (EOMs) are particularly resistant to ALS progression, and exploring the underlying molecular nature may offer significant therapeutic value. Reactive aldehyde 4-hydroxynonenal (HNE) is implicated in ALS pathogenesis, and Aldh3a1 is an inactivation-resistant intracellular aldehyde dehydrogenase that detoxifies 4-HNE to protect eyes against UV-induced oxidative stress. We detected prominently higher levels of *Aldh3a1* in mouse EOMs compared to other muscles under normal physiological conditions. In an ALS mouse model (hSOD1^{G93A}) reaching end-stage, *Aldh3a1* expression was maintained high in EOMs, substantially elevated in soleus and diaphragm, but only moderately increased in extensor digitorum longus (EDL) muscle, which endured the most severe pathological remodeling, as demonstrated by unparalleled upregulation of a denervation marker *Ankrd1*. Importantly, sciatic nerve transection in wildtype mice further confirmed induced *Aldh3a1* and *Ankrd1* expression in an inverse manner across muscle types in response to denervation. Mechanistically, whole-muscle RNA-Seq and pharmacological tests indicate that higher basal levels of lipid oxidation in soleus and diaphragm muscles may predispose them to stronger Nrf2 antioxidant responses under pathological stress compared to EDL, leading to more prominent *Aldh3a1* upregulation. Additionally, the identification of the myoblast fusion marker *Mymk* as an EOM signature gene suggests that the spontaneous activation of satellite cells contributes to high levels of *Aldh3a1* in EOMs. Functionally, adeno-associated virus-mediated overexpression of *Aldh3a1* protected myotubes from 4-HNE-induced DNA fragmentation and plasma membrane leakage. It also restored MG53-mediated membrane repair, highlighting its potential for clinical applications.

Key words: Amyotrophic lateral sclerosis, skeletal muscle, extraocular muscle, reactive aldehydes, aldehyde dehydrogenase, sciatic nerve transection

Abbreviations: 4-hydroxy-2-nonenal (4-HNE); acetylcholine receptor (AChR); amyotrophic lateral sclerosis (ALS); extensor digitorum longus (EDL); soleus (SOL), diaphragm (DIA), extraocular muscle (EOM); enzyme-linked immunosorbent assay (ELISA); hindlimb (HL); malondialdehyde (MDA), neuromuscular junction (NMJ), satellite cell (SC), sciatic nerve transection (SNT), sulforaphane (SFN).

Introduction

Amyotrophic Lateral Sclerosis (ALS) is a devastating neuromuscular disorder characterized by progressive motor neuron death and skeletal muscle wasting. Remarkably, extraocular muscles (EOMs) exhibit superior preservation of structure, neuromuscular junction (NMJ) integrity and function in ALS patients and rodent models [1-5]. Beside EOMs, animal model studies also show that limb and body muscles respond to ALS progression differently, with fast-twitch muscles generally more susceptible to degeneration than in slow-twitch muscles [6-8]. Factors underlying this phenomenon can be multi-faceted and could hold clues for identifying novel therapeutic targets against muscle degeneration in ALS or other neuromuscular disorders.

Reactive aldehydes resulting from oxidative stress are involved in the pathological process of multiple neurodegenerative disorders including ALS [9]. One major route for intracellular production of reactive aldehydes is the peroxidation of polyunsaturated fatty acids (PUFA), which are major component of cellular membranous structures [10, 11]. Well-known examples include 4-hydroxynonenal (4-HNE), malondialdehyde (MDA) and acrolein [11], which can form adducts with protein through Michael addition or Schiff base formation [12-14]. These adducts contribute to protein crosslinking and aggregation [11], leading to broad pathological consequences including disrupted cell signaling, altered gene expression, inhibited enzyme activity, compromised mitochondrial function and disformed cytoskeleton [15]. Reactive aldehydes can also form adducts with DNA, resulting in inter-strand crosslinks, base substitution, mutation and fragmentation [16-22]. Furthermore, 4-HNE is known to form adducts with phosphatidylethanolamine and certain ion channel proteins at the plasma membrane, causing altered ionic homeostasis and cellular osmolarity [23, 24]. The deprivation of local antioxidants like glutathione by 4-HNE can promote further lipid peroxidation at the plasma membrane, creating a vicious cycle that leads to cell death [25, 26]. Elevated levels of lipid peroxidation markers, including 4-HNE adducts, have been detected in cells of central nervous system and body fluids from patients and/or models of amyotrophic lateral sclerosis (ALS), Alzheimer disease (AD), Parkinson disease (PD) and Huntington disease (HD) [9].

To neutralize reactive aldehydes, the human body has deployed a series of detoxification enzymes. Aldehyde dehydrogenases (ALDHs) aimed to oxidize the carbonyl group into corresponding acids, while aldo-keto reductases (AKRs) reduce aldehydes into corresponding alcohols [12]. Aldh2, Aldh1a1 and Aldh3a1 are the three most studied ALDHs [27]. Aldh2 is a mitochondrial ALDH abundantly present in liver, brain, heart and muscle [28]. Aldh1a1 and Aldh3a1 are primarily cytosolic and are extremely abundant in mammalian corneal and lens to protect against ultraviolet radiation (UVR) induced generation of reactive aldehydes and their pathological consequences [20, 29, 30]. Aldh3a1 knockout mice and Aldh1a1/Aldh3a1 double knockout mice develop cataracts by 1 month of age [30]. It is also worth noticing that a kinetics study of these three ALDHs reported that Aldh2 was irreversibly inactivated by 4-HNE and acrolein at above 10 μM. Aldh1a1 was inactivated by acrolein at concentrations higher than 1 mM, while no inactivation of Aldh3a1 was observed by either 4-HNE, acrolein or MDA even at 20 mM [27]. Thus, Aldh3a1 is the most inactivation-resistant isoform of the three and the focus of the current study.

It is unknown whether aldehyde dehydrogenases are involved in varied susceptibility of different muscles to degeneration under ALS. In this study, we examined the expression of *Aldh2*, *Aldh1a1* and *Aldh3a1* in EDL, soleus, diaphragm and EOMs of end-stage hSOD1^{G93A} (G93A) mice, a well-established ALS rodent model, as well as their wildtype (WT) littermates [31]. Only *Aldh3a1* exhibited dramatically higher expression in EOMs compared to other muscles in WT mice. Meanwhile it was prominently upregulated in G93A soleus and diaphragm compared to WT controls. However, the

upregulation was less pronounced in G93A EDL muscle, which suffers the most severe NMJ degeneration in these four muscle types in G93A mice [5]. Indeed, the commonly used denervation marker *Ankrd1* [32, 33] was induced most in G93A EDL muscle, suggesting that the expression level of *Aldh3a1* is inversely linked to the severity of muscle pathological remodeling. The distinct expression pattern of *Aldh3a1* gene was also confirmed at the protein level. In EDL and soleus muscles of WT mice with sciatic nerve transection (SNT), *Ankrd1* expression was quickly elevated post operation, especially in EDL muscles and gradually decreased over time. In contrast, the upregulation of *Aldh3a1* occurred with a multi-day delay in soleus, while no significant upregulation of *Aldh3a1* occurred in EDL muscles even after 14 days. Thus, *Ankrd1* and *Aldh3a1* exhibit inverse induction pattern over muscle type and time post denervation.

Whole muscle RNA-seq analysis and pharmacological tests revealed potential mechanisms underlying the differential regulations of *Aldh3a1* expression in various muscles involving Nrf2 signaling activation. Importantly, transduction of EDL and soleus-SC derived myotubes with adeno-associated virus (AAV) vector expressing human *Aldh3a1* markedly decreased 4-HNE-induced DNA fragmentation. Furthermore, MG53 is a muscle specific tripartite motif family protein nucleating the assembly of the repair machinery on injured plasma membrane [34]. We previously reported abnormal MG53 intracellular aggregation and compromised membrane repair in ALS muscles [35]. Here, we demonstrated that enforced expression of *Aldh3a1* in cultured myotubes protected against 4-HNE-induced plasma membrane damage and restored MG53 mediated membrane repair. Thus, the differential expression level of *Aldh3a1* in muscles in response to ALS progression likely reflects a varying protective role of Aldh3a1 against reactive aldehyde cytotoxicity in various muscles. Organ-targeted delivery of Aldh3a1 through gene therapy may add a new therapeutic avenue for ALS or diseases associated with elevated oxidative stress.

Results

Differential expression of *Aldh3a1* across muscles and ALS disease states

The transcription levels of three aldehyde dehydrogenases (*Aldh1a1*, *Aldh2* and *Aldh3a1*) in muscles from different anatomic origins, including hindlimb (EDL and soleus), diaphragm and EOMs were evaluated by qRT-PCR. These muscles were collected from WT mice and their end-stage G93A littermates (4 pairs of male mice and 4 pairs of female mice at 4-5 months of age). As shown in **Figure 1A** left panels and **Figure 1-figure supplement 1A**, *Aldh3a1* was expressed prominently higher in EOMs than other muscles in WT mice (59-fold that in EDL, 22-fold that in soleus and 91 folds that in diaphragm). Notable induction of *Aldh3a1* was detected for soleus (50-fold) and diaphragm (79-fold) from G93A mice compared to WT controls, while the induction was far less pronounced for EDL (7-fold). In G93A EOMs, *Aldh3a1* expression was maintained at comparable levels to WT EOMs. This pathological induction pattern is interesting as EDL is the most severely affected during ALS progression [4, 5]. Importantly, denervation-associated pathological remodeling marker *Ankrd1* (*CARP)* exhibited stronger induction in G93A EDL (69-fold) than diaphragm (35-fold), soleus (22-fold) and EOMs (3-fold) (**Figure 1A** right panels and **Figure 1-figure supplement 1B**). Data here suggests a potential relationship between *Aldh3a1* expression and muscle resistance to degeneration in ALS. In contrast to *Aldh3a1*, neither *Aldh1a1* nor *Aldh2* exhibited superior expression in EOMs, and none of the G93A muscles exhibited differential expression of *Aldh1a1* compared to WT controls, implying that it is unlikely involved in muscle dependent response to ALS

progression (**Figure 1-figure supplementary 1C,D** and **Figure 1-Source Data 1**). Thus, *Aldh3a1* is
the focus of this study*.*

The unique expression pattern of *Aldh3a1* is also present at the protein level. In WT mice,
Aldh3a1 protein was most abundant in EOMs (132-fold that in EDL, 24-fold that in soleus and 29-fold
that in diaphragm after normalization to housekeeping protein Gapdh). In line with the qPCR results,
the abundance of Aldh3a1 protein was maintained at high levels in EOMs in end-stage G93A mice.
Remarkably, significantly elevated levels of Aldh3a1 protein were detected in diaphragm (14-fold),
soleus (6-fold) and EDL muscle (5-fold) compared to WT counterparts (**Figure 1B, C, Figure 1-figure**
supplement 1A, 2 and **Figure 1-Source Data 2**). In comparison, Ankrd1 protein increased most
significantly in G93A EDL muscle compared to WT counterparts (594-fold), followed by diaphragm
(66-fold), soleus (51-fold) (**Figure 1B, C, Figure 1-figure supplement 1B, 2** and **Figure 1-Source**
Data 2). Consistently, in the transverse section of EDL muscles from G93A mice, most myofibers
were positively stained with anti-Ankrd1 antibodies (**Figure 2A**). Ankrd1 positive myofibers were
relatively sparse in G93A soleus and diaphragm muscles, while in G93A EOMs they were extremely
scarce. To further validate the association between Ankrd1 and denervation, we performed sciatic
nerve transection (SNT) in the right hindlimbs of WT mice 4-5 months of age. The majority of EDL
and soleus myofibers became Ankrd1 positive post SNT (**Figure 2B**). Thus, our data implied an
inverse relationship between *Aldh3a1* expression pattern and that of the denervation marker *Ankrd1*
in G93A muscles.

We did whole-mount immunostaining to investigate the distribution of Aldh3a1 protein in
myofibers (**Figure 3A** and **Figure 3-figure supplement 1A**). In WT mice, EOM exhibited higher
levels of cytosolic Aldh3a1 than other muscles, confirming the Western blot and qPCR data in **Figure**
1. In muscles from end-stage G93A mice, soleus and diaphragm show increased Aldh3a1 in the
cytosol and myonuclei, while Aldh3a1 positive myofibers were scarce in EDL (**Figure 3A** and **Figure**
3-figure supplement 1A, yellow arrows). Additionally, axon terminals (labelled by Sg2 antibody)
were partially or fully absent from AChR positive area (labelled by BTX), confirming partial or full
denervation of G93A muscles. These data further support that the induction of *Aldh3a1* expression is
less robust in G93A EDL than in soleus and diaphragm.

It is worth noticing that the nuclear Aldh3a1 signal dips in the DAPI dense foci, where genomic
DNA is highly compacted (**Figure 3B** and **Figure 3-figure supplement 1B**) [36], implying that
nuclear Aldh3a1 preferentially localizes to the euchromatin regions, a pattern commonly seen for
histone modifications associated with actively transcribing genes [37]. We speculate that the elevation
of Aldh3a1 levels in G93A muscles is a self-defense mechanism protecting cytosolic protein and
vulnerable genomic DNA regions against lipid peroxidation stress under ALS.

We also examined whether there is an inverse correlation between aldh3a1 and Ankrd1
distribution on the level of individual myofibers. Most myofibers in G93A EDL were positive for Ankrd1,
with few of them also positive for Aldh3a1 (**Figure 4A**, yellow arrows). Ankrd1 positive myofibers
were sparser in G93A soleus and diaphragm, whereas myofibers with elevated cytosolic and nuclear
Aldh3a1 were more frequently seen (**Figure 4B, C**, yellow arrows). These Aldh3a1 positive myofibers
could either be Ankrd1 positive (yellow arrows) or negative (blue arrows) (**Figure 4B, C**). In G93A
EOMs, Ankrd1 positive myofibers were extremely scarce, whereas most G93A EOM myofibers
exhibited cytosolic distribution of Aldh3a1 (**Figure 4D**), a feature shared by their WT peers (**Figure 4-**
figure supplement 1). In sum, although Aldh3a1 and Ankrd1 are not mutually exclusive in individual

myofibers, their abundance does exhibit an inverse relationship on the whole-muscle scale across
different muscles in end-stage G93A mice.

Sciatic nerve transection confirms the differential *Aldh3a1* and *Ankrd1* expression in EDL and
soleus post denervation

Previous studies from us and others identified increased production of reactive oxygen species
(ROS) in mitochondria as a common pathological feature in denervated muscles and muscle from
ALS mouse models [38-40]. To better understand the relationship between *Aldh3a1* expression
changes and muscle denervation, we performed sciatic nerve transection (SNT) in the right hindlimbs
of WT mice 4-5 months of age. The left hindlimbs were sham operated as controls. The changes of
gene expression in EDL and soleus with SNT (compared to their sham-operated controls) were
examined at Day 3, 7 and 14 post operation by qRT-PCR (**Figure 5A** and **Figure 5-Source Data 1**).
In EDL with SNT, there is no notable increase of *Aldh3a1* expression throughout the two weeks. In
soleus with SNT, the elevated *Aldh3a1* expression was observed at Day 7 and 14. This time-
dependent change of *Aldh3a1* expression was also confirmed at the protein level, with significant
elevation detected in soleus but not in EDL post SNT (**Figure 5B**, **Figure 5-figure supplement 1** and
Figure 5-Source Data 2). The *Ankrd1* expression was elevated in both EDL and soleus and peaked
at Day 3 post SNT and decreased afterwards (**Figure 5**, **Figure 5-figure supplement 1** and **Figure**
5-Source Data 3). It is worth noticing that the increase of Ankrd1 protein was more pronounced in
EDL (333-fold) compared to soleus (45-fold) at Day 3 post SNT, while the increase of Aldh3a1 protein
was significant in soleus (7.6-fold) but not in EDL (1.6-fold) at Day 14 post SNT.

The whole-mount immunostaining confirmed the complete denervation of myofibers in both EDL
and soleus after SNT by the absence of axonal terminals in NMJs (**Figure 6A**). Myofibers with
nuclear enriched Aldh3a1 was spotted in soleus at Day 7 post SNT. At Day 14, more myofibers
showed increased Aldh3a1, not only in myonuclei, but also in cytosol (**Figure 6A**, insets). In line with
the qPCR and Western Blot results, no noticeable increase of Aldh3a1 was observed in EDL
myofibers after SNT compared to sham controls. Consistently, immunostaining revealed presence of
Aldh3a1 positive myofibers in transverse sections of soleus but not EDL at Day 14 post SNT,
whereas higher proportions of Ankrd1 positive myofibers were observed in these EDL muscles than
soleus (**Figure 6B**). These data further demonstrate that *Aldh3a1* induction varies significantly
between muscle types following denervation. Muscles with high proportions of mitochondria-rich
oxidative fibers, like soleus and diaphragm, seem to be more prone to Aldh3a1 induction under
elevated oxidative stress, likely as a self-protection mechanism.

RNA-seq and pharmacological tests reveal potential mechanisms underlying muscle type
dependent regulation of *Aldh3a1* expression

To explore the molecular mechanism underlying muscle type dependent regulation of *Aldh3a1*
expression, whole muscle RNA-Seq were performed to compare between EDL, soleus, diaphragm
and EOMs derived from end-stage G93A mice and age-matched WT controls (2 male and 2 female
mice per group). In principle component analysis, EOM is the only muscle group with WT and G93A
samples clustered together, whereas other muscle groups have more distinct expression profiles
between WT and G93A (**Figure 7A**). This confirms that EOMs are less affected by ALS progression.
To identify EOM signature genes, we generated lists of differentially expressed genes (DEGs, log2
fold change > 0.5, p.adjust < 0.05) comparing EOMs to EDL, soleus and diaphragm in WT mice. A
total of 866 DEGs, including *Aldh3a1*, were in common between the three lists (**Figure 7-Source**

data 1). Gene ontology analysis of these genes reveals that the most significant biological processes
are related to eye development (**Figure 7-figure supplement 1A**). In addition, a good proportion of
these genes are involved in axon guidance/neuron migration (**Figure 7-figure supplement 1B, C**),
which is consistent with our previous discovery in EOM SC-derived myoblasts and myotubes [5] and
may explain why neuromuscular junctions are preserved better in EOMs than other muscles in ALS.

Since eye development is one of the top biological processes highlighted by gene ontology
analysis, we suspected that the superior expression of *Aldh3a1* in EOMs was due to *Pax6*, one of the
EOM signature genes (**Figure 7C**) that is a master regulator of eye formation and has been reported
to activate *Aldh3a1* expression in mouse cornea [41]. However, upon closer inspection, we found that
the transcripts per million (TPM) of *Pax6* is extremely low even for EOMs (the average is 1.47, not
detected in one sample). Meanwhile *Pax6* transcripts were not detected in EOM SC-derived
myoblasts or myotubes in our previous RNA-Seq results (GSE249484), which argues against our
hypothesis. Another EOM signature gene that catches our attention is *Mymk* (**Figure 7C**), whose
elevated expression in EOMs are also confirmed by qPCR (**Figure 1-figure supplement 1E**). *Mymk*
mediates membrane fusion of myoblasts to existing myofibers during muscle development and
regeneration [42]. Distinct from other muscles, EOMs are known for spontaneous activation of SCs to
fuse with existing myofibers even without injury [5, 43]. That is why *Mymk* expression, which is
supposed to be very low in uninjured WT muscles, is high in EOMs. Furthermore, our previous RNA-
Seq data revealed notably higher expression of *Aldh3A1* in myoblasts than differentiated myotubes
(GSE249484), implying that the spontaneous activation of SCs could contribute to the maintenance of
high *Aldh3a1* levels in EOMs.

As to the differential regulation of *Aldh3a1* expression in other muscles, gene ontology analysis
reveals that the most significantly upregulated biological processes in soleus and diaphragm
compared to EDL in WT mice are fatty acid metabolism and oxidation (**Figure 7B**). Representative
genes include *Acadl*, *Acadm*, *Acat1, Adipoq*, *Alox12* and *Alox15* (**Figure 7C** and **Figure 7-figure
supplement 1D**). High basal level of lipid oxidation could predispose these muscles to stronger Nrf2
antioxidant response to additional oxidative stress caused by denervation or ALS [44]. Indeed, soleus
and diaphragm showed a significant elevation of *Hmox1,* a classic target gene of Nrf2 signaling in
G93A mice compared to EDL (**Figure 7C** and **Figure 1-figure supplement 1F**). Meanwhile more
Aldh3a1 and less *Ankrd1* transcripts were detected in soleus and diaphragm than EDL of G93A mice
(**Figure 7C**), confirming the previous qPCR results.

Nrf2 signaling has been reported to activate *Aldh3a1* expression through the antioxidant response
element in its promoter in human cancer cells [45, 46]. To examine whether Nrf2 dependent
activation of *Aldh3a1* expression also occurs in skeletal muscle, we conducted pharmacological tests
using Nrf2 signaling activator sulforaphane (SFN). The myotubes derived from EDL and soleus SCs
(SCs were isolated from corresponding muscles in WT mice 4-5 months of age) were treated with 10
μM SFN for 48 hrs. As shown in **Figure 7D** and **Figure 7-Source Data 2**, more prominent
upregulation of *Hmox1* and *Aldh3a1* expression were observed in soleus-SC derived myotubes than
EDL-SC derived ones. These results support the differential predisposition to Nrf2 signaling activation
as a contributor to differential regulation of *Aldh3a1* between muscles.

**Transducing myotubes with AAV-Aldh3a1 protects against 4-HNE induced apoptosis and
plasma membrane defects**

4-HNE is the most toxic reactive aldehyde generated during lipid peroxidation [47]. We evaluated
4-HNE levels in EDL and soleus muscles from end-stage G93A mice and age-matched WT controls
through ELISA (**Figure 8A** and **Figure 8-Source Data 1**). 4-HNE levels were higher in soleus than
EDL muscles from WT mice, potentially due to more active fatty acid oxidation in soleus as indicated
in **Figure 7B**. However, 4-HNE level significantly increased in EDL but not in soleus muscles from
end-stage G93A mice compared to WT controls, implicating the more pronounced induction of
Aldh3a1 expression in G93A soleus in preventing further elevation of 4-HNE levels under ALS.

To assess 4-HNE cytotoxicity in skeletal muscles, the SCs-derived myotubes from WT EDL and
soleus muscles were first treated with 0, 7.5, 15 or 30 μM of 4-HNE for 2 hours. Afterwards the
myotubes were cultured for another 16 hours in differentiation medium before TUNEL staining, as
apoptosis can take 12-24 hours to occur [48]. TUNEL positive nuclei (indicative of apoptotic cell death
hallmarked by DNA fragmentation) were scarce in myotubes treated with 0, 7.5 or 15 μM 4-HNE but
dramatically increased in myotubes treated with 30 μM 4-HNE (53% for myotubes derived from EDL
myoblasts, 20% for myotubes derived from soleus myoblasts), accompanied by notable detachment
of myotubes from the culture chamber (**Figure 8B, C** and **Figure 8-Source Data 2**). Due to the
extensive occurrence of myotube detachment, calculating the percentage of TUNEL positive nuclei
may not accurately reflect 4-HNE cytotoxicity. Since plasma membrane is commonly the primary site
of 4-HNE production, we assessed the plasma membrane integrity by measuring the leakage of the
cytosolic housekeeping enzyme lactate dehydrogenase (LDH) to the culture medium [49]. As shown
in **Figure 8D** and **Figure 8-Source Data 3,** LDH leakage assay revealed a notable uptick of plasma
membrane damage at 30 μM 4-HNE (65.6% for EDL-derived myotubes, 78.4% for soleus -derived
myotubes).

Intriguingly, AAV-Aldh3a1 transduced myotubes, which had 2 to 6-fold higher levels of Aldh3a1
protein (**Figure 9A** and **Figure 9-Source Data 1**), exhibited dramatically lower percentage of
apoptotic nuclei (3.5% for EDL-derived myotubes, 0.83% for soleus-dervied myotubes (**Figure 9B, C**
and **Figure 9-Source Data 2**) and LDH leakage (30.4% for EDL-derived myotubes, 38.4% for soleus-
derived myotubes (**Figure 9D** and **Figure 9-Source Data 3**). The transduction of AAV-Aldh3a1 alone
did not change myotube viability compared to non-transduced controls (**Figure 9D**), indicating that
Aldh3a1 protein is not toxic to cultured muscle cells.

Increased LDH leakage and myotube detachment following 4-HNE treatment indicate
compromised plasma membrane integrity. We previously reported defects in MG53 mediated
membrane repair mechanism G93A mice [35, 50]. Here we examined the impact of 4-HNE treatment
on the formation of stable MG53 repair patches in myotubes with overexpression of GFP-MG53 [34].
Without 4-HNE treatment, GFP-MG53 was predominantly cytosolic and quickly translocated to the
plasma membrane in response to cell membrane injury induced by saponin, a compound
permeabilizing plasma membrane. About 86% recorded myotubes maintained elevated membranous
GFP-MG53 signal at the end of recording, implying the formation of stable MG53 repair patches
(**Video 1, Figure 10** and **Figure 10-Source Data 1**), while the proportion dropped to 8.6% for
myotubes treated with 30 μM 4-HNE for 2 hours before recording (**Figure 10B, D** and **Figure 10-
Source Data 1**). A good portion of 4-HNE treated myotubes already had blebs on the plasma
membrane before saponin administration. Additionally, the peak intensity of membranous GFP-MG53
signals were also lower than those not treated with 4-HNE (**Figure 10C**). These observations indicate
that plasma membrane damage has already occurred during 4-HNE treatment (**Video 2** and **Figure
10**). Intriguingly, pretreatment of myotubes with AAV-Aldh3a1 before 4-HNE treatment not only

prevented 4-HNE induced plasma membrane blebbing, restored the peak intensity of membranous
GFP-MG53 signal, but also increased the proportion of myotubes maintaining elevated membranous
GFP-MG53 signal to 77% (**Video 3**, **Figure 10** and **Figure 10-Source Data 1**).

Discussion

In this study we discovered that Aldh3a1 protein is extremely abundant in mouse EOMs and can
be upregulated in other muscles under pathological conditions such as denervation and ALS
progression. The extent of *Aldh3a1* upregulation in the ALS mouse model (G93A) varies over muscle
type, more prominent in soleus, diaphragm than EDL. In contrast, the muscle pathological remodeling
marker *Ankrd1* is more substantially elevated in EDL than soleus and diaphragm of G93A mice. The
induction pattern of *Aldh3a1* is also inverse to that of *Ankrd1* over muscle type and time in WT mice
with sciatic nerve transection. In addition, the protective effects of Aldh3a1 against 4-HNE in muscle
are multi-faceted, from DNA fragmentation in the nuclei to plasma membrane damage and repair
defects involving MG53.

Aldh3a1 has been reported to be an inactivation-resistant detoxifier of reactive aldehydes such
as 4-HNE in an enzyme kinetics study [27]. Here we detected increased 4-NHE levels in G93A
muscles without pronounced increase of Aldh3a1. Meanwhile 4-HNE treated myotubes showed an
increase in nuclear DNA fragmentation, cell membrane leakage and loss of MG53 membrane repair
function, implying that elevated 4-HNE level overwhelmed the physiological detoxification capacity of
muscle cells. Remarkably, pretreating myotubes with AAV-Aldh3a1 potently protected myotubes
against all the above cytotoxic effects caused by 4-HNE. In addition, AAV-Aldh3a1 transduction alone
did not alter myotube viability, which increases the likelihood for its therapeutic application in the
future.

Nuclear enrichment of Aldh3a1 has been observed previously in cultured human and rabbit
corneal epithelial cells overexpressing *Aldh3a1* [29, 51, 52]. Since reactive aldehydes such as 4-HNE
can directly modify genomic DNA, leading to mutagenesis and fragmentation in multiple non-muscle
cell types [16, 21, 22], the DNA protection role of Aldh3a1 has been investigated. Indeed, it was
reported to prevent DNA damage and apoptosis induced by 4-HNE, hydrogen peroxide, or UVR in
cultured corneal epithelial (HCE) cells and stromal fibroblasts [20, 51-54]. Aside from its role to
directly detoxify reactive aldehydes, Aldh3a1 was also reported to promote a series of DNA damage
detection/repair related processes in HCE cells including prolonging cell cycle, increasing levels of
total and phospho (Ser15) p53 and activating ATM/ATR signaling pathway, which is central to the
maintenance of genome integrity [51, 52, 54-56]. These effects could be a unique advantage of
applying Aldh3a1 in therapy. Interestingly, in HCE cells overexpressing *Aldh3a1,* the expression of
GADD45A, another muscle denervation marker [57], was found to be downregulated [54], indicating
the inverse correlation between the expression levels of *Aldh3a1* and muscle denervation marker
Ankrd1 may not be a coincidence, but a phenomenon broadly presents in different tissues. In this
study, we did kymographic profiling of Aldh3a1 inside myonuclei and unveiled its preferential
distribution in the euchromatin regions, where genomic DNA is less compacted and could be more
vulnerable to attacks by reactive aldehydes. It is reasonable to hypothesize that *Aldh3a1* upregulation
in muscle is a self-defense mechanism against pathologically elevated lipid peroxidation stress not
only to protect cytosolic protein but also to maintain genome DNA integrity.

An alternative scenario would be Aldh3a1 self-regulates its own expression via a positive
feedback loop. Yet this is unlikely as Aldh3a1 is not known to directly bind DNA like a transcription

factor. It is also worth noticing that the high expression of *Aldh3a1* gene does not guarantee the
nuclear enrichment of Aldh3a1 protein. In EOMs and myotubes transduced with AAV-Aldh3a1, we did
not observe notable nuclear enrichment of Aldh3a1 (**Figure 3A**, **Figure 3-figure supplement 1A**,
Figure 4D, **Figure 4-figure supplement 1** and **Figure 9B**). The mechanism driving Aldh3a1 nuclear
translocation is still under investigation, as the deactivation of its predicted nuclear localization
sequence did not achieve the expected outcome [52]. It is possible that the extent of Aldh3a1 unclear
accumulation depends on the status of the oxidative stress inside the nuclei.

Aldh3a1 expression has been reported to be activated by Nrf2 signaling through antioxidant
response element (ARE) in its promoter in human cancer cells [45]. The activation of Nrf2 signaling
has been reported to enhance exercise endurance capacity, augment skeletal muscle regeneration
after ischemia-reperfusion injury and ameliorate muscle mass/contractility decline during aging [58-
63]. In addition, Nrf2 signaling activator edaravone (Radicava) has been approved by FDA to combat
ALS [64, 65].

Our whole muscle RNA-Seq detected higher expression levels of lipid oxidation related genes in
soleus and diaphragm than EDL muscles. This could be due to higher content of oxidative myofibers
in these two muscles [66], whose mitochondria have a predisposition toward using fatty acids as fuel
[67]. This is also supported by higher 4-HNE levels in soleus than EDL muscles from WT mice
indicated by ELISA. The high basal lipid oxidation levels could predispose soleus and diaphragm to
Nrf2 signaling activation [44], potentially through making the promoter regions of antioxidant genes
more accessible to Nrf2. In addition, the higher proportion of mitochondria-rich oxidative fibers in
soleus and diaphragm could also accelerate the activation of Nrf2 signaling as a self-defense
mechanism due to faster accumulation of mitochondria-produced ROS under ALS or denervation [38-
40]. Indeed, we detected more transcripts of Nrf2 target gene *Hmox1* in soleus and diaphragm than
EDL of G93A mice. In addition, treatment of Nrf2 activator SFN induced more prominent upregulation
of *Hmox1* and *Aldh3a1* in soleus-SC derived myotubes than EDL-SC derived ones. Thus, the
differential predisposition to Nrf2 signaling activation is likely a contributing factor to the differential
regulation of *Aldh3a1* expression between soleus, diaphragm and EDL muscles.

However, Nrf2 signaling alone cannot explain the superior expression of *Aldh3a1* in EOMs as
none of its classic target genes, such as *Hmox1*, *Nqo1*, *gclm* or *gsta4* [64], shows up in the EOM
signature gene list. Based on the results of gene ontology analysis, we suspected that eye
development-related genes, such as *Pax6*, were involved in activating *Aldh3a1* expression in EOMs
[41]. However, the extremely low levels of *Pax6* transcripts in EOM samples, combined with their
absence in EOM-SC derived myoblasts and myotubes in our previous RNA-Seq data (GSE249484),
argues that they are more likely due to residual eye tissue left during dissection, rather than
endogenous expression in the muscles. On the other hand, the identification of *Mymk* as an EOM
signature gene provides an alternative scenario to explain high *Aldh3a1* levels in EOMs. *Mymk* is
known for mediating the fusion of myoblasts with existing myofibers during muscle regeneration [42].
Its expression should be low in uninjured muscles, but EOM is an exception because of spontaneous
activation of SCs. This unique property was first documented due to the observation of BrdU positive
nuclei (from proliferating myoblasts) in the center of uninjured EOM myofibers [43]. Our recent
fluorescence-activated cell sorting analysis also revealed significantly more heterogenous levels of
VCAM (quiescent satellite cell marker) in EOM SCs than those from other muscles of WT mice,
confirming spontaneous activation of EOM SCs without pathological stimuli [5]. Furthermore, we
detected notably more *Aldh3a1* transcripts in myoblasts than myotubes (either originated from EOM,

diaphragm or hindlimb muscle SCs) in previous RNA-Seq results (GSE249484). This is not surprising
as many types of stem cells are characterized by upregulated expression of specific ALDH
isoenzymes, like *Aldh1a1*, *Aldh1a3* and *Aldh3a1*, which promote their survival [68-70]. High *Aldh3a1*
expression has been correlated with cell proliferation and resistance to reactive aldehyde cytotoxicity
[70]. Therefore, the spontaneous activation of abundant SCs in EOMs and their fusion with existing
myofibers could contribute to high *Aldh3a1* levels.

As to 4-HNE targets in ALS patients, a few key proteins in the ventral horn motor neurons, such
as the astrocytic glutamate transporter EAAT2, have been found to be modified by 4-HNE [71]. In
ALS rodent models, 4-HNE has been reported to form adducts with Drp2, Hsp70, glutathione and
carnosine [72, 73]. Here, we demonstrated functional disruption of MG53 by 4-HNE. Previous studies
implied cysteine 242 of mouse MG53 as a redox sensor to oligomerize or crosslink [34, 74].
Meanwhile cysteine is the residue most prone to 4-HNE attack through Michael addition [75, 76].
Interestingly, similar to the time-lapse imaging results of 4-HNE treated myotubes expressing GFP-
MG53 we reported here, cardiomyocytes expressing GFP-MG53(C242A), which mutates cysteine
242 to alanine, also demonstrated failed maintenance of membranous signal after the application of
saponin [74]. Thus, whether C242 of MG53 is one of the residues forming Michael adducts with 4-
HNE in muscles of G93A mice is worth further investigation in future studies. The result may explain
the abnormal formation of cytosolic MG53 aggregates in these muscles [35].

In summary, we identified multi-faceted benefits of Aldh3a1 in protecting against reactive
aldehyde cytotoxicity, which likely contribute to the distinct muscle response observed in ALS
progression. The spectrum of vulnerability among motor neurons innervating different muscles
remains a mystery in ALS. Specifically, the ocular motor neurons are notably resistant to ALS,
persisting throughout the disease [77]. While our results do not exclude a primary effect of those
neurons on EOMs, we can only speculate that ocular motor neurons, like EOMs, may possess a
superior capacity to mitigate lipid peroxidation cytotoxicity due to their close developmental
relationship with the eye. Future studies are warranted to experimentally validate this hypothesis and
test the therapeutic potential of motor neuron targeted expression of Aldh3a1 for ALS.

Materials and methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Strain, strain background (*Mus musculus*)	B6SJLF1/J	Jackson Laboratory	Stock # 100012	WT mice Both male and female Age up to 4 months
Strain, strain background (*Mus*	B6SJL-Tg (SOD1*G93A)	Jackson Laboratory	Stock # 002726	G93A mice Both male and female Age up to 4 months

musculus)				
Cell line (*Mus musculus*)	Primary cultured mouse satellite cells	This paper		Isolated from EDL or soleus muscles dissected from 4-month-old WT female mice, enriched by 3 rounds of preplating during culture.
Cell line (*Rattus norvegicus*)	Primary cultured rat spinal motor neurons (RSCMNs)	iXCells	SKU # 10RA-033	Isolated from D16 rat embryo spinal cord; Negative for mycoplasma, bacteria, yeast, and fungi.
Transfected construct (*Mus musculus*)	AAVMYO(9P1)-tMCKp-huAldh3a1.HA-WPRE	Vector Biolabs	SKU # AAV-200734	Adeno-associated virus for muscle-specific expression of human ALHD3A1.
Peptide, recombinant protein	Collagenase II	Worthington	Cat # LS004176	Final concentration 0.26%
Peptide, recombinant protein	Dispase II	Sigma Aldrich	Cat # D46931G	Final concentration 0.24%
Peptide, recombinant protein	Hyaluronidase	Worthington	Cat # LS002592	Final concentration 0.16%
Peptide, recombinant protein	DNase I	Worthington	Cat # LS002139	Final concentration 0.04%
Antibody	Anti-human Aldh3a1 (Rabbit polyclonal IgG)	Proteintech	Cat # 15578-1-AP	IF (1:150) WB (1: 2000)
Antibody	Anti-chicken MHC (Mouse monoclonal IgG2b)	Developmental Studies Hybridoma Bank	Cat # MF20	IF (1:200)
Antibody	Anti-rat Ankrd1 (Mouse	Sigma Aldrich	Cat # MABS1228	IF (1:300) WB (1: 2000)

	monoclonal IgG1)			
Antibody	Anti-human Gapdh (Mouse monoclonal IgG2b)	Proteintech	Cat # 60004-1-Ig	WB (1:5000)
Antibody	Anti-human Gapdh (Rabbit polyclonal IgG)	Proteintech	Cat # 10494-1-AP	WB (1:5000)
Antibody	Anti-human β-actin (Rabbit polyclonal IgG)	Proteintech	Cat # 20536-1-AP	WB (1:5000)
Antibody	Anti-human Sod1 (Rabbit polyclonal IgG)	Proteintech	Cat # 10269-1-AP	WB (1:5000)
Antibody	HRP-conjugated affinipure goat anti-mouse IgG(H+L)	Proteintech	Cat # SA00001-1	WB (1:5000)
Antibody	HRP-conjugated affinipure goat anti-rabbit IgG(H+L)	Proteintech	Cat # SA00001-2	WB (1:5000)
Antibody	Anti-zebrafish Synaptotagmin 2 (Mouse monoclonal IgG2a)	Developmental Studies Hybridoma Bank	Cat # znp-1	IF (1:150)
Recombinant DNA reagent	GFP-MG53	Laboratory of Dr. Jianjie Ma		
Sequence-based reagent	qPCR primers for mouse *Aldh1a1*	Sigma Aldrich		Fwd: GGAATACCGTGGTTGTCAAGCC Rev: CCAGGGACAATGTTTACCACGC
Sequence-based reagent	qPCR primers for mouse *Aldh3a1*	Sigma Aldrich		Fwd: GGTCCTTGTCATAGGTGCTTGG Rev: GAAAGCAGGTCTG

				CCATGTGATC
Sequence-based reagent	qPCR primers for mouse *Ankrd1*	Sigma Aldrich		Fwd: GCTTAGAAGGACAC TTGGCGATC Rev: GACATCTGCGTTTC CTCCACGA
Commercial assay or kit	CF640R TUNEL Assay Apoptosis Detection Kit	Biotium	Cat # 30074	1.5 hours incubation with the reaction mix containing TdT
Commercial assay or kit	Direct-zol RNA MiniPrep Plus	Zymo Research	Cat # R2070	
Commercial assay or kit	GoScript Reverse Transcriptase Mix + Oligo(dT)	Promega	Cat # A2791	
Commercial assay or kit	Universal SYBR Green Fast qPCR Mix	Abclonal	Cat # RK21203	
Commercial assay or kit	CytoTox 96 Non-Radioactive Cytotoxicity Assay	Promega	Cat # G1780	
Commercial assay or kit	Pierce BCA Protein Assay Kits	ThermoFisher Scientific	Cat # 23225	
Commercial assay or kit	4-Hydroxynonenal ELISA Kit	Abcam	Cat # ab287803	
Chemical compound, drug	AF 488 conjugated α-Bungarotoxin	ThermoFisher Scientific	Cat # B13422	IF (1:1000)
Chemical compound, drug	DAPI in water	Biotium	Cat # 40043	IF (1:10000)
Chemical compound, drug	TransfeX transfection reagent	ATCC	Cat #ACS-4005	
Chemical	4-HNE	Cayman	Cat #32100	

compound, drug		Chemical		
Chemical compound, drug	Saponin	ThermoFisher Scientific	Cat #A18820.14	Final concentration 0.00375%
Chemical compound, drug	Sulforaphane	Cayman Chemical	Cat #4478-93-7	
Chemical compound, drug	Sodium butyrate	SAFC	Product #ARK2161	
Chemical compound, drug	EGTA	Sigma Aldrich	Cat #E4378	
Chemical compound, drug	BTS	Sigma Aldrich	Cat #203895	
Chemical compound, drug	Protease and Phosphatase Inhibitor Cocktail	ThermoFisher Scientific	Cat #1861282	
Chemical compound, drug	EDTA	ThermoFisher Scientific	Cat #1861276	

Animals

All animal experiments were carried out in accordance with the recommendations in the *Guide for*
the Care and Use of Laboratory Animals of the National Institutes of Health. The protocol on the
usage of mice was approved by the Institutional Animal Care and Use Committee of the University of
Texas at Arlington (A19.001, approval date: 09/20/2018). WT mice used in this study were of B6SJL
background. The ALS transgenic mouse model (hSOD1G93A) with genetic background of B6SJL
was originally generated by Drs. Deng and Siddique's group at Northwestern University and
deposited to the Jackson Lab as B6SJL-Tg (SOD1*G93A) [31]. G93A mice of both genders were
euthanized for sample collection at the endpoint when they were unable to right themselves to a
sternal position within 30 seconds (4-5 months of age). WT mice of the corresponding gender were
euthanized for sample collection at the same day as their G93A littermates.

Sciatic nerve transection

Surgical instruments were sterilized by autoclave before operation. Mice were anesthetized with
constant flow isoflurane inhalation. Hairs in the two posterior thighs and lower back were shaved as
much as possible with electric clipper. The skin region to be operated on was aseptically prepared
using surgical scrub with betadine or equivalent surgical soap and rinse with 70% alcohol. Incision
through the skin and superficial muscles was made parallel and just inferior to the femur of the right

hindlimb. Curved-end forceps were used to divide the muscles and expose the sciatic nerve. 5 mm of
the sciatic nerve was removed with fine surgical scissors. Both ends of the nerve were sutured to
prevent regeneration. The incision on the skin was closed with stainless steel wound clips. A sham
procedure following the same steps without severing the sciatic nerve was performed for the left
hindlimb as control.

Immunofluorescence (IF) and imaging of whole mount and muscle samples

For whole mount immunofluorescence, EDL, soleus, diaphragm and EOM samples were fixed
and permeabilized in precooled methanol at -20 °C for 15 min. The samples were rehydrated by three
changes of PBS and incubated with Alexa Fluor 488 conjugated α-Bungarotoxin (ThermoFisher
B13422, 1:1000) in blocking buffer (PBS containing 2% BSA, 2% horse serum, 0.1% Tween-20, 0.1%
Triton X-100 and 0.05% sodium azide) at 4 °C for 1 day. The samples were then washed with PBS
and further incubated with the primary antibodies against Aldh3a1 (Proteintech 15578-1-AP 1:150)
and Synaptotagmin-2 (Sg2 for short, DSHB ZNP-1 concentrate + 50% glycerol 1:150) in blocking
buffer at 4 °C for 1 day. On the third day, the samples were washed with PBS for 3 times and
incubated with corresponding secondary antibodies labelled with Alexa Fluor.

For section immunofluorescence, EDL, soleus, diaphragm and EOM were fixed in either 4%
paraformaldehyde or 3% glyoxal fixative (pH 4.5, containing 20% ethanol) for 8-12 hours at 4 °C.
Glyoxal based fixative can penetrate tissue faster due to the presence of ethanol [78-80]. Antigen
retrieval was conducted in Tris-EDTA buffer (pH 9.0) for 30 min at 95 °C. Afterwards the samples
were washed with PBS containing 1% glycine once and two more times with PBS. Then the samples
were immersed in blocking buffer for 45 min at room temperature followed by primary antibody
incubation at 4 °C overnight. Next day, after washing with PBS for three times, the samples were
incubated with Alexa Fluor labelled secondary antibodies (ThermoFisher Scientific 1:800) for 2 hours
at room temperature. The samples were then washed with PBS, counterstained with DAPI and
mounted in antifade mounting media (20 mM Tris, 0.5% N-propyl gallate, 80% glycerol) for imaging.
Primary antibodies used: Aldh3a1, Ankrd1 (Sigma MABS1228 1:200)

RNA extraction and qRT-PCR

Homogenization was performed in FastPrep-24 Classic bead beating grinder (MP Biomedicals
116004500). The muscle homogenate was transferred to centrifuge tubes containing phase lock gel
(QuantaBio 2302830) and 1/10 volume of 1- bromo-3-chloropropane was added. The tubes were
hand shaken for 12 seconds, left at bench top for 2 minutes and then centrifuged at 16000 x g at 4 °C
for 15 minutes. The upper phase was transferred to a new centrifuge tube and mixed with equal
volume of ethanol. The following steps of RNA purification were performed with Direct-zol RNA
Miniprep Plus kit (Zymo Research R2070). RNA concentration was measured with Quantus
Fluorometer (Promega E6150). GoScript Reverse Transcription Mix, oligo(dT) was used for the
reverse transcription reaction (Promega A2791). First-strand cDNAs were diluted and mixed with 2X
Universal SYBR Green Fast qPCR Mix (Abclonal RK21203), as well as corresponding primers for
qPCR using StepOnePlus Real-Time PCR system (ThermoFisher Scientific 4376600). Relative
quantification (RQ) of gene expression was generated by ΔΔCt method. The sequences of primers
used for qPCR were listed in Key Resource Table.

Bulk RNA-Seq and analysis

475 RNAs were extracted from EDL, soleus, diaphragm and extraocular muscles from end-stage
476 G93A mice and age-matched WT controls (2 male and 2 female mice per group) as described above.
477 cDNA library preparation and sequencing were carried out by Innomics.inc. The average number of

clean reads per sample was 23 million. The clean reads were mapped to Genome Reference Consortium Mouse Build 39 (GRCm39) using Rsubread package of Bioconductor [81] and the average percentage of aligned reads was 87%. DESeq2 package [82] was used to identify differentially expressed genes (DEGs) between different samples. The data were prefiltered to keep only genes with counts ≥ 10 in at least 4 samples. Only genes with log2FC > 0.5 and p.adjust < 0.05 were considered differentially expressed in volcano plots and gene ontology analysis. Principle component analysis (PCA) was performed with pcaExplorer package [83, 84]. Volcano plots were created using EnhancedVolcano package. Gene ontology analysis was performed with clusterProfiler package [85]. Fastq files of clean reads and the excel file containing TPM values will be uploaded to GEO database upon acceptance of the manuscript.

Western blot

Proteins were extracted by grinding corresponding muscles in 10 volumes of RIPA buffer containing protease inhibitors (ThermoFisher Scientific), resolved by SDS-PAGE, and transferred to PVDF membrane with Bio-Rad semidry transfer cell. The primary antibodies were Aldh3a1 (Proteintech 15578-1-AP), Ankrd1 (Sigma MABS1228) and Gapdh (Proteintech 60004-1-Ig). Protein bands were detected with Bio-Rad Clarity ECL kit and ChemiDoc Imaging system. Signal strengths and backgrounds were measured using ImageJ software.

BCA protein assay and 4-HNE enzyme-linked immunosorbent assay (ELISA)

To evaluate 4-HNE levels in EDL and soleus muscles from end-stage G93A mice and age-matched WT controls. We homogenized corresponding muscles in lysis buffer (50mM Tris + 0.9% NaCl + 0.1%SDS + Protease and Phosphatase Inhibitor Cocktail + 1 mM EDTA) using FastPrep-24 Classic bead beating grinder (MP Biomedicals 116004500). The homogenates were briefly centrifuged and transferred to new 1.5 ml microcentrifuge tubes. The tubes were centrifuged again for 15 min at 12000 x g and the supernatants were collected for concentration measurement using Pierce BCA Protein Assay Kit (ThermoFisher Scientific 23225) according to their protocol on a plate reader (SpectraMax i3x, Molecular Devices).

The homogenates were then diluted to ~1 mg/ml with the sample/standard dilution buffer of 4-HNE ELISA kit (Abcam ab287803). Each well in the Micro ELISA Plate of the kit was washed twice with 1x wash solution. 50 µl of diluted samples or 4-HNE standards were loaded to each well of the Micro ELISA Plate and incubated with 50 µl of diluted biotin-detection antibodies at 37 °C for 45 min. The plate was washed again with 1x wash solution for 3 times and incubated with 0.1 ml SABC working solution/well at 37 °C for 30 min. The plate was then washed again with 1x wash solution for 5 times and loaded with 90 µl TMB substrate solution/well at 37 °C for 15 min. 50 µl stop solution was added per well and the absorbance of each well at 450 nm and 620 nm were immediately measured by a plate reader. 4-HNE concentration derived from the standard curve was normalized to the protein concentration of the corresponding well.

Primary culture of myoblasts

EDL or soleus muscles were dissected from two 4-month-old WT female mice and placed in 0 mM Ca^{2+} Ringer's solution. Excessive connective tissue, tendons, fat, blood clots and vessels were removed as much as possible. After cleanup, the muscles were transferred into the digestion media (DMEM containing 5% FBS, 1% Pen/Strep and 10 mM HEPES) and minced into small blocks using scissors. Collagenase II (Worthington LS004176, ≥125 units/mg), dispase II (Sigma D46931G, ≥0.5

units/mg), hyaluronidase (Worthington LS002592, ≥300 USP/NF units/mg) and Dnase I (Worthington
LS002139, ≥2,000 Kunitz units/mg) were added at the final concentration of 0.26%, 0.24%, 0.16%
and 0.04% (weight/volume), respectively. The digestion system was placed in an orbital shaker
running at 70 rpm at 37 °C for 45 min. The digested muscles were triturated 10 times with a 20-gauge
needle attached to a 5 ml syringe. Afterwards the triturated mixture was pipetted onto a pre-wetted
100 μm strainer and centrifuged at 100 x g for 2.5 min to precipitate the bulky myofiber debris. The
supernatant was transferred onto a pre-wetted 40 μm strainer, and cells were collected by centrifuged
at 1200 x g for 6.5 min. After the removal of supernatant, cells were resuspended with 5 ml growth
medium (Ham's F-10 medium + 20% FBS + 1%Pen/Strep + 5ng/ml bFGF + 10 μg/ml Plasmocin and
pre-plated in non-coated T25 flasks at 37 °C for 30 min. The unattached cells were transferred
together with medium to Matrigel coated T25 flasks and cultured till reaching about 50% confluence.
Pre-plating continued for three more passages to diminish non-myoblast cells.

Chemical and AAV treatment of cultured myotubes

Cultured myoblasts derived from soleus or EDL muscles of 4-month-old WT mice (seeded in
laminin coated chambered cover glass at about 2.5x10^4/compartment) were induced to differentiate
into myotubes in low serum medium (DMEM+2% horse serum) for four days, with medium changed
every other day. 4-HNE treatment was performed at 0, 7.5, 15, 30 μM for 2 hours. The treatment time
was determined based on a previous study showing 4-HNE notably induced oxidative stress as early
as 2 hours in Swiss 3T3 fibroblasts [22].

AAVMYO(9P1)-tMCKp-huAldh3a1.HA-WPRE (AAV-Aldh3a1 for short) was produced by Vector
Biolabs. AAVMYO(9P1) is a mutant of AAV9 that shows superior efficiency and specificity in skeletal
and cardiac muscles [86]. The triple muscle creatine kinase promoter (tMCKp) specifically drives
gene expression in differentiated muscle tissue [87]. To carry out AAV transduction, cultured
myoblasts derived from soleus or EDL muscles of WT mice 4-5 months of age (seeded in laminin
coated 48-well plate or chambered cover glass at about 2.5x10^4/compartment) were induced to
differentiate into myotubes with low serum medium containing AAV-Aldh3a1 (1x10^11 GC/ml) for 2
days and regular differentiation medium for another 2 days. 4-HNE treatment (30 μM for 2 hours) was
performed at the end of the second day. After the transient exposure to 4-HNE, myotubes were
cultured for another 16 hours in regular low serum medium before TUNEL assay (Biotium 30074) or
LDH leakage-based cytotoxicity assay (Promega G1780). This is because apoptosis can take 12-24
hours to occur [48].

TUNEL assay

Triplicate cultures of myotubes in chambered cover glass (Cellvis C8-1.5H-N) were fixed and
permeabilized with precooled methanol at -20 °C for 15 min. Afterwards, the samples were
rehydrated by washing with PBS for 3 times (5 min each). TUNEL staining was performed following
the manual of Biotum CF640R TUNEL Assay Apoptosis Detection Kit. The incubation time with the
reaction mix (containing TdT enzyme) was 1.5 hours at 37 °C. Afterwards, the samples were washed
twice with PBS and incubated with blocking buffer for 45 min at room temperature, followed by
immunostaining with Aldh3a1 and myosin heavy chain (DSHB, MF-20 1:100) antibodies.
Counterstaining with DAPI was done to calculate the ratio of nuclei with fragmented genomic DNA.

Non-radioactive cytotoxicity (LDH leakage) assay

We generally followed the protocol of the assay kit (Promega, G1780). In brief, 5-6 replicates of
myotube cultures with or without AAV transduction, 4-HNE treatment, were cultured in 300 µl regular
low serum medium for 16 hours, along with wells containing only 300 µl medium but no cells (to serve
as medium only controls and volume correction controls). Next morning, 50 µl of culture supernatant
was collected and 50 µl of lysis solution (contains Triton X-100) was added to lyse all the remaining
myotubes (37 °C, 45 min). The supernatant, the medium only control, the all-cell-lysed solution (12.5
µl diluted to 50 µl with PBS+1% BSA, 1:4 dilution) and the volume correction controls (medium only
wells with 50 µl replaced with lysis solution and diluted 1:4 with PBS+1% BSA) were combined with
50 µl of chromogenic substrates of lactate dehydrogenase (LDH, a stable cytosolic enzyme that is
released upon cell death), respectively, for absorbance measurement by a plate reader (SpectraMax
i3x, Molecular Devices). Absorbance at 490 nm was recorded every 5 min for 35 min for each sample.
The largest rate of absorbance increase (Vmax) calculated from 3 continuous points out of the 7 time
points measured reflects the concentration of LDH. To compare the percentages of dead myotubes
under different treatment conditions, we calculated the percentage of LDH in the culture supernatant
to total LDH as follows: 100*(Vmax(supernatant)-Vmax(medium only control))/(4*(Vmax(all-cell-
lysed)-Vmax(volume correction control))+(Vmax(supernatant)-Vmax(medium only control))/6).

Transfection of GFP-MG53 and recording of saponin induced MG53 membrane translocation

4.5×10^4 of myoblasts derived from soleus muscles of 4-month-old WT mice were seeded into glass
bottom dishes (Matsunami D35-14-1.5-U) and cultured in growth medium overnight before
transfection of 3 µg GFP-MG53 plasmids with TransfeX transfection reagent (ATCC ACS-4005). We
selected soleus derived myoblasts because they form multi-nuclei, lengthy myotubes more robustly
than EDL derived ones, facilitating the measurement of membranous GFP-MG53 signals.GFP-MG53
was a gift from Dr. Jianjie Ma's lab [34]. One day after transfection, the culture medium was changed
to low serum medium to induce differentiation for 4 days, with medium changed every two days. AAV-
Aldh3a1 treatment group was incubated with the AAV vector at the first two days of differentiation. 4-
HNE treatment group was incubated with 30 µM 4-HNE for 2 hours at the end of the 4-day
differentiation. The myotubes were washed three times with 0 mM Ca^{2+} Ringer's solution containing
0.5 mM EGTA (Sigma Aldrich E4378) and 90 µM BTS (Sigma Aldrich 203895) to prevent contraction.
Time lapse imaging of myotubes were performed with Leica TCS SP8 confocal microscope at an
interval of 2 sec for 3 min. 0.00375% saponin (Thermo Scientific A18820.14) was applied 10 sec after
recording started. Regions of interest were generated at plasma membrane regions and background
signals from myotube free area was deduced to acquire GFP-MG53 intensity profile (*F*) over time. F_0
was the averaged intensity of the first 4 frames.

Data analysis and statistics

Box-and-dot plots were generated with ggplot2 package of R (version 4.2.3). The lower hinge,
median line and upper hinge correspond to the first, second and third quartiles. The lower and upper
whiskers extend from the hinges to the largest value no further than 1.5 times of inter-quartile range
(distance between the first and third quartiles). Data beyond the end of the whiskers are outlying
points. The stat_compare_means function of ggplot 2 was used to compare the means of two groups.
Wilcoxon rank-sum test was used unless there were tie values between two groups (**Figure 8C**) or
the group had less than 4 samples (**Figure 9A**). In those cases, t-test was used instead. For multi-
group data, one-way ANOVA P values were generated by aov function of R.
603

Acknowledgement: We thank Dr. Yongfu Wang at and Dr. Ji Pang, who previously worked at Stowers Institute for Medical Research, for the help in trouble-shooting the glyoxal fixation protocol for section immunofluorescence. This study has been supported by grants from NIH (R01NS105621 to JZ, R01NS129219 to JZ and JM) and the Department of Defense AL170061(W81XWH1810684) to JZ.

Author contributions

Ang Li, Conceptualization, Data curation, Formal analysis, Validation, Investigation, Visualization, Methodology, Writing – original draft, Project administration, Writing – review and editing; Li Dong, Xuejun Li, Jianxun Yi, Data curation, Validation, Investigation, Methodology; Jianjie Ma, Conceptualization, Funding acquisition, Writing – review and editing; Jingsong Zhou, Conceptualization, Resources, Supervision, Funding acquisition, Validation, Investigation, Visualization, Methodology, Writing – original draft, Project administration, Writing – review and editing

References

1. Domellöf, F.P. (2019). The Extraocular Muscles Are Selectively Spared in ALS. In Amyotrophic Lateral Sclerosis-Recent Advances and Therapeutic Challenges. (IntechOpen).

2. Tjust, A.E., Brännstrom, T., and Pedrosa Domellöf, F. (2012). Unaffected motor endplate occupancy in eye muscles of ALS G93A mouse model. Frontiers in bioscience (Scholar edition) *4*, 1547-1555.

3. Ahmadi, M., Liu, J.-X., Brännström, T., Andersen, P.M., Stål, P., and Pedrosa-Domellöf, F. (2010). Human extraocular muscles in ALS. Investigative ophthalmology & visual science *51*, 3494-3501.

4. Valdez, G., Tapia, J.C., Lichtman, J.W., Fox, M.A., and Sanes, J.R. (2012). Shared resistance to aging and ALS in neuromuscular junctions of specific muscles. PloS one *7*, e34640.

5. Li, A., Yi, J., Li, X., Dong, L., Ostrow, L.W., Ma, J., and Zhou, J. (2024). Distinct transcriptomic profile of satellite cells contributes to preservation of neuromuscular junctions in extraocular muscles of ALS mice. Elife *12*, RP92644.

6. Derave, W., Van Den Bosch, L., Lemmens, G., Eijnde, B.O., Robberecht, W., and Hespel, P. (2003). Skeletal muscle properties in a transgenic mouse model for amyotrophic lateral sclerosis: effects of creatine treatment. Neurobiology of disease *13*, 264-272.

7. Chiu, A.Y., Zhai, P., Dal Canto, M.C., Peters, T.M., Kwon, Y.W., Prattis, S.M., and Gurney, M.E. (1995). Age-dependent penetrance of disease in a transgenic mouse model of familial amyotrophic lateral sclerosis. Molecular and Cellular Neuroscience *6*, 349-362.

8. Frey, D., Schneider, C., Xu, L., Borg, J., Spooren, W., and Caroni, P. (2000). Early and selective loss of neuromuscular synapse subtypes with low sprouting competence in motoneuron diseases. J Neurosci *20*, 2534-2542.

9. Di Domenico, F., Tramutola, A., and Butterfield, D.A. (2017). Role of 4-hydroxy-2-nonenal (HNE) in the pathogenesis of alzheimer disease and other selected age-related neurodegenerative disorders. Free Radical Biology and Medicine *111*, 253-261.

10. Li, Y., Zhao, T., Li, J., Xia, M., Li, Y., Wang, X., Liu, C., Zheng, T., Chen, R., and Kan, D. (2022). Oxidative Stress and 4-hydroxy-2-nonenal (4-HNE): Implications in the Pathogenesis and Treatment of Aging-related Diseases. Journal of Immunology Research *2022*.

11. Guéraud, F., Atalay, M., Bresgen, N., Cipak, A., Eckl, P., Huc, L., Jouanin, I., Siems, W., and Uchida, K. (2010). Chemistry and biochemistry of lipid peroxidation products. Free radical research *44*, 1098-1124.

649 12. Mol, M., Regazzoni, L., Altomare, A., Degani, G., Carini, M., Vistoli, G., and Aldini, G. (2017). Enzymatic
650   and non-enzymatic detoxification of 4-hydroxynonenal: Methodological aspects and biological
651   consequences. Free Radical Biology and Medicine *111*, 328-344.
652 13. Cai, J., Bhatnagar, A., and Pierce Jr, W.M. (2009). Protein modification by acrolein: formation and
653   stability of cysteine adducts. Chemical research in toxicology *22*, 708-716.
654 14. Aubourg, S.P. (1993). Interaction of malondialdehyde with biological molecules—new trends about
655   reactivity and significance. International journal of food science & technology *28*, 323-335.
656 15. Esterbauer, H. (1993). Cytotoxicity and genotoxicity of lipid-oxidation products. The American journal
657   of clinical nutrition *57*, S779-S786.
658 16. Voulgaridou, G.-P., Anestopoulos, I., Franco, R., Panayiotidis, M.I., and Pappa, A. (2011). DNA damage
659   induced by endogenous aldehydes: current state of knowledge. Mutation Research/Fundamental and
660   Molecular Mechanisms of Mutagenesis *711*, 13-27.
661 17. Falletti, O., and Douki, T. (2008). Low glutathione level favors formation of DNA adducts to 4-hydroxy-
662   2 (E)-nonenal, a major lipid peroxidation product. Chemical research in toxicology *21*, 2097-2105.
663 18. Wang, H.-T., Zhang, S., Hu, Y., and Tang, M.-s. (2009). Mutagenicity and sequence specificity of
664   acrolein-DNA adducts. Chemical research in toxicology *22*, 511-517.
665 19. Liu, X.-y., Zhu, M.-x., and Xie, J.-p. (2010). Mutagenicity of acrolein and acrolein-induced DNA adducts.
666   Toxicology mechanisms and methods *20*, 36-44.
667 20. Pappa, A., Chen, C., Koutalos, Y., Townsend, A.J., and Vasiliou, V. (2003). Aldh3a1 protects human
668   corneal epithelial cells from ultraviolet-and 4-hydroxy-2-nonenal-induced oxidative damage. Free
669   Radical Biology and Medicine *34*, 1178-1189.
670 21. Brambilla, G., Sciaba, L., Faggin, P., Maura, A., Marinari, U., Ferro, M., and Esterbauer, H. (1986).
671   Cytotoxicity, DNA fragmentation and sister-chromatid exchange in Chinese hamster ovary cells
672   exposed to the lipid peroxidation product 4-hydroxynonenal and homologous aldehydes. Mutation
673   Research/Genetic Toxicology *171*, 169-176.
674 22. Kutuk, O., Adli, M., Poli, G., and Basaga, H. (2004). Resveratrol protects against 4‐HNE induced
675   oxidative stress and apoptosis in Swiss 3T3 fibroblasts. Biofactors *20*, 1-10.
676 23. Jovanović, O., Škulj, S., Pohl, E.E., and Vazdar, M. (2019). Covalent modification of
677   phosphatidylethanolamine by 4-hydroxy-2-nonenal increases sodium permeability across
678   phospholipid bilayer membranes. Free Radical Biology and Medicine *143*, 433-440.
679 24. Bhatnagar, A. (1995). Electrophysiological effects of 4-hydroxynonenal, an aldehydic product of lipid
680   peroxidation, on isolated rat ventricular myocytes. Circulation research *76*, 293-304.
681 25. Mustafa, A.G., Alfaqih, M.A., and Al-Shboul, O. (2018). The 4-hydroxynonenal mediated oxidative
682   damage of blood proteins and lipids involves secondary lipid peroxidation reactions. Experimental and
683   Therapeutic Medicine *16*, 2132-2137.
684 26. Yamashima, T. (2024). 4-Hydroxynonenal from Mitochondrial and Dietary Sources Causes Lysosomal
685   Cell Death for Lifestyle-Related Diseases. Nutrients *16*, 4171.
686 27. Yoval-Sánchez, B., and Rodríguez-Zavala, J.S. (2012). Differences in susceptibility to inactivation of
687   human aldehyde dehydrogenases by lipid peroxidation byproducts. Chemical research in toxicology *25*,
688   722-729.
689 28. Stewart, M., Malek, K., and Crabb, D.W. (1996). Distribution of messenger RNAs for aldehyde
690   dehydrogenase 1, aldehyde dehydrogenase 2, and aldehyde dehydrogenase 5 in human tissues.
691   Journal of investigative medicine: the official publication of the American Federation for Clinical
692   Research *44*, 42-46.
693 29. Stagos, D., Chen, Y., Cantore, M., Jester, J.V., and Vasiliou, V. (2010). Corneal aldehyde
694   dehydrogenases: multiple functions and novel nuclear localization. Brain research bulletin *81*, 211-218.
695 30. Lassen, N., Bateman, J.B., Estey, T., Kuszak, J.R., Nees, D.W., Piatigorsky, J., Duester, G., Day, B.J.,
696   Huang, J., and Hines, L.M. (2007). Multiple and additive functions of ALDH3A1 and ALDH1A1: cataract

phenotype and ocular oxidative damage in Aldh3a1 (-/-)/Aldh1a1 (-/-) knock-out mice. Journal of
Biological Chemistry *282*, 25668-25676.

31.   Gurney, M.E., Pu, H., Chiu, A.Y., Dal Canto, M.C., Polchow, C.Y., Alexander, D.D., Caliendo, J., Hentati,
A., Kwon, Y.W., and Deng, H.-X. (1994). Motor neuron degeneration in mice that express a human Cu,
Zn superoxide dismutase mutation. Science *264*, 1772-1775.

32.   Laure, L., Suel, L., Roudaut, C., Bourg, N., Ouali, A., Bartoli, M., Richard, I., and Danièle, N. (2009).
Cardiac ankyrin repeat protein is a marker of skeletal muscle pathological remodelling. The FEBS
journal *276*, 669-684.

33.   Nakamura, K., Nakada, C., Takeuchi, K., Osaki, M., Shomori, K., Kato, S., Ohama, E., Sato, K., Fukayama,
M., and Mori, S. (2003). Altered expression of cardiac ankyrin repeat protein and its homologue,
ankyrin repeat protein with PEST and proline-rich region, in atrophic muscles in amyotrophic lateral
sclerosis. Pathobiology *70*, 197-203.

34.   Cai, C., Masumiya, H., Weisleder, N., Matsuda, N., Nishi, M., Hwang, M., Ko, J.-K., Lin, P., Thornton, A.,
and Zhao, X. (2009). MG53 nucleates assembly of cell membrane repair machinery. Nature cell biology
11, 56-64.

35.   Yi, J., Li, A., Li, X., Park, K., Zhou, X., Yi, F., Xiao, Y., Yoon, D., Tan, T., and Ostrow, L.W. (2021). MG53
preserves neuromuscular junction integrity and alleviates ALS disease progression. Antioxidants *10*,
1522.

36.   Imai, R., Nozaki, T., Tani, T., Kaizu, K., Hibino, K., Ide, S., Tamura, S., Takahashi, K., Shribak, M., and
Maeshima, K. (2017). Density imaging of heterochromatin in live cells using orientation-independent-
DIC microscopy. Molecular biology of the cell *28*, 3349-3359.

37.   Alvarenga, E.M., Rodrigues, V.L., Moraes, A.S., Naves, L.S., Mondin, M., Felisbino, M.B., and Mello,
M.L.S. (2016). Histone epigenetic marks in heterochromatin and euchromatin of the Chagas' disease
vector, Triatoma infestans. Acta histochemica *118*, 401-412.

38.   Xiao, Y., Karam, C., Yi, J., Zhang, L., Li, X., Yoon, D., Wang, H., Dhakal, K., Ramlow, P., and Yu, T. (2018).
ROS-related mitochondrial dysfunction in skeletal muscle of an ALS mouse model during the disease
progression. Pharmacological research *138*, 25-36.

39.   Karam, C., Yi, J., Xiao, Y., Dhakal, K., Zhang, L., Li, X., Manno, C., Xu, J., Li, K., and Cheng, H. (2017).
Absence of physiological Ca 2+ transients is an initial trigger for mitochondrial dysfunction in skeletal
muscle following denervation. Skeletal muscle *7*, 6.

40.   Muller, F.L., Song, W., Jang, Y.C., Liu, Y., Sabia, M., Richardson, A., and Van Remmen, H. (2007).
Denervation-induced skeletal muscle atrophy is associated with increased mitochondrial ROS
production. American journal of physiology-Regulatory, integrative and comparative physiology *293*,
R1159-R1168.

41.   Davis, J., Davis, D., Norman, B., and Piatigorsky, J. (2008). Gene expression of the mouse corneal
crystallin Aldh3a1: activation by Pax6, Oct1, and p300. Investigative ophthalmology & visual science *49*,
1814-1826.

42.   Millay, D.P., O'Rourke, J.R., Sutherland, L.B., Bezprozvannaya, S., Shelton, J.M., Bassel-Duby, R., and
Olson, E.N. (2013). Myomaker is a membrane activator of myoblast fusion and muscle formation.
Nature *499*, 301-305.

43.   McLoon, L.K., and Wirtschafter, J. (2002). Activated satellite cells are present in uninjured extraocular
muscles of mature mice. Transactions of the American Ophthalmological Society *100*, 119.

44.   Hayashi, T., Kudo, T., Fujita, R., Fujita, S.-i., Tsubouchi, H., Fuseya, S., Suzuki, R., Hamada, M., Okada, R.,
and Muratani, M. (2021). Nuclear factor E2-related factor 2 (NRF2) deficiency accelerates fast fibre
type transition in soleus muscle during space flight. Communications Biology *4*, 787.

45.   Sreerama, L., and Sládek, N.E. (2001). Three different stable human breast adenocarcinoma sublines
that overexpress ALDH3A1 and certain other enzymes, apparently as a consequence of constitutively

upregulated gene transcription mediated by transactivated EpREs (electrophile responsive elements)
present in the 5′-upstream regions of these genes. Chemico-Biological Interactions *130*, 247-260.

46. Baird, L., and Yamamoto, M. (2020). The Molecular Mechanisms Regulating the KEAP1-NRF2 Pathway.
Mol Cell Biol *40*.

47. Esterbauer, H., Schaur, R.J., and Zollner, H. (1991). Chemistry and biochemistry of 4-hydroxynonenal,
malonaldehyde and related aldehydes. Free radical Biology and medicine *11*, 81-128.

48. Saraste, A. (1999). Morphologic criteria and detection of apoptosis. Herz *24*, 189.

49. Korzeniewski, C., and Callewaert, D.M. (1983). An enzyme-release assay for natural cytotoxicity.
Journal of immunological methods *64*, 313-320.

50. Li, A., Yi, J., Li, X., Dong, L., Ostrow, L.W., Ma, J., and Zhou, J. (2022). Deficient Sarcolemma Repair in
ALS: A Novel Mechanism with Therapeutic Potential. Cells *11*, 3263.

51. Pappa, A., Brown, D., Koutalos, Y., DeGregori, J., White, C., and Vasiliou, V. (2005). Human aldehyde
dehydrogenase 3A1 inhibits proliferation and promotes survival of human corneal epithelial cells.
Journal of Biological Chemistry *280*, 27998-28006.

52. Voulgaridou, G.-P., Theologidis, V., Venetikidou, M., Tsochantaridis, I., Tsolou, A., Koffa, M.,
Panayiotidis, M.I., and Pappa, A. (2023). Investigating the functional roles of aldehyde dehydrogenase
3A1 in human corneal epithelial cells. International Journal of Molecular Sciences *24*, 5845.

53. Lassen, N., Pappa, A., Black, W.J., Jester, J.V., Day, B.J., Min, E., and Vasiliou, V. (2006). Antioxidant
function of corneal ALDH3A1 in cultured stromal fibroblasts. Free Radical Biology and Medicine *41*,
1459-1469.

54. Voulgaridou, G.-P., Tsochantaridis, I., Tolkas, C., Franco, R., Giatromanolaki, A., Panayiotidis, M.I., and
Pappa, A. (2020). Aldehyde dehydrogenase 3A1 confers oxidative stress resistance accompanied by
altered DNA damage response in human corneal epithelial cells. Free Radical Biology and Medicine
150, 66-74.

55. Canman, C.E., Lim, D.-S., Cimprich, K.A., Taya, Y., Tamai, K., Sakaguchi, K., Appella, E., Kastan, M.B., and
Siliciano, J.D. (1998). Activation of the ATM kinase by ionizing radiation and phosphorylation of p53.
Science *281*, 1677-1679.

56. Awasthi, P., Foiani, M., and Kumar, A. (2015). ATM and ATR signaling at a glance. Journal of cell
science *128*, 4255-4262.

57. Bongers, K.S., Fox, D.K., Ebert, S.M., Kunkel, S.D., Dyle, M.C., Bullard, S.A., Dierdorff, J.M., and Adams,
C.M. (2013). Skeletal muscle denervation causes skeletal muscle atrophy through a pathway that
involves both Gadd45a and HDAC4. Am J Physiol Endocrinol Metab *305*, E907-915.

58. Onoki, T., Izumi, Y., Takahashi, M., Murakami, S., Matsumaru, D., Ohta, N., Wati, S.M., Hatanaka, N.,
Katsuoka, F., and Okutsu, M. (2021). Skeletal muscle-specific Keap1 disruption modulates fatty acid
utilization and enhances exercise capacity in female mice. Redox Biology *43*, 101966.

59. Wang, L., Yang, S., Yan, L., Wei, H., Wang, J., Yu, S., Kong, A.-N.T., and Zhang, Y. (2019). Hypoxia
preconditioning promotes endurance exercise capacity of mice by activating skeletal muscle Nrf2.
Journal of Applied Physiology *127*, 1267-1277.

60. Al‐Sawaf, O., Fragoulis, A., Rosen, C., Keimes, N., Liehn, E.A., Hölzle, F., Kan, Y.W., Pufe, T., Sönmez,
T.T., and Wruck, C.J. (2014). Nrf2 augments skeletal muscle regeneration after ischaemia–reperfusion
injury. The Journal of pathology *234*, 538-547.

61. Shelar, S.B., Narasimhan, M., Shanmugam, G., Litovsky, S.H., Gounder, S.S., Karan, G., Arulvasu, C.,
Kensler, T.W., Hoidal, J.R., and Darley-Usmar, V.M. (2016). Disruption of nuclear factor (erythroid-
derived-2)–like 2 antioxidant signaling: a mechanism for impaired activation of stem cells and delayed
regeneration of skeletal muscle. The FASEB Journal *30*, 1865.

62. Huang, D.-D., Yan, X.-L., Fan, S.-D., Chen, X.-Y., Yan, J.-Y., Dong, Q.-T., Chen, W.-Z., Liu, N.-X., Chen, X.-L.,
and Yu, Z. (2020). Nrf2 deficiency promotes the increasing trend of autophagy during aging in skeletal
muscle: a potential mechanism for the development of sarcopenia. Aging (Albany NY) *12*, 5977.

63. Ahn, B., Pharaoh, G., Premkumar, P., Huseman, K., Ranjit, R., Kinter, M., Szweda, L., Kiss, T., Fulop, G.,
and Tarantini, S. (2018). Nrf2 deficiency exacerbates age-related contractile dysfunction and loss of
skeletal muscle mass. Redox biology *17*, 47-58.
64. Jiménez-Villegas, J., Ferraiuolo, L., Mead, R., Shaw, P., Cuadrado, A., and Rojo, A.I. (2021). NRF2 as a
therapeutic opportunity to impact in the molecular roadmap of ALS. Free Radical Biology and
Medicine *173*, 125-141.
65. Nightingale, B. (2020). The effect of edaravone on amyotrophic lateral sclerosis. Journal of Neurology
Research *10*, 150-159.
66. Augusto, V., Padovani, C.R., and Campos, G.E.R. (2017). Skeletal muscle fiber types in C57BL6J mice.
Journal of Morphological Sciences *21*, 0-0.
67. Leary, S., Lyons, C., Rosenberger, A., Ballantyne, J., Stillman, J., and Moyes, C. (2003). Fiber-type
differences in muscle mitochondrial profiles. American Journal of Physiology-Regulatory, Integrative
and Comparative Physiology *285*, R817-R826.
68. Vassalli, G. (2019). Aldehyde dehydrogenases: not just markers, but functional regulators of stem cells.
Stem cells international *2019*, 3904645.
69. Moreb, J.S. (2008). Aldehyde dehydrogenase as a marker for stem cells. Current stem cell research &
therapy *3*, 237-246.
70. Muzio, G., Maggiora, M., Paiuzzi, E., Oraldi, M., and Canuto, R.A. (2012). Aldehyde dehydrogenases
and cell proliferation. Free Radical Biology and Medicine *52*, 735-746.
71. Pedersen, W.A., Fu, W., Keller, J.N., Markesbery, W.R., Appel, S., Smith, R.G., Kasarskis, E., and
Mattson, M.P. (1998). Protein modification by the lipid peroxidation product 4‐hydroxynonenal in
the spinal cords of amyotrophic lateral sclerosis patients. Annals of neurology *44*, 819-824.
72. Reis, P.V., Vargas, B.S., Rebelo, R.A., Massafera, M.P., Prado, F.M., Oreliana, H., de Oliveira, H.V.,
Freitas, F.P., Ronsein, G.E., and Miyamoto, S. (2024). Quantitative Analysis of Glutathione and
Carnosine Adducts with 4-Hydroxy-2-nonenal in Muscle in a hSOD1G93A ALS Rat Model. Chemical
Research in Toxicology *37*, 1306-1314.
73. Perluigi, M., Poon, H.F., Hensley, K., Pierce, W.M., Klein, J.B., Calabrese, V., De Marco, C., and
Butterfield, D.A. (2005). Proteomic analysis of 4-hydroxy-2-nonenal-modified proteins in G93A-SOD1
transgenic mice-A model of familial amyotrophic lateral sclerosis. Free Radical Biology and Medicine
38, 960-968.
74. Wang, X., Xie, W., Zhang, Y., Lin, P., Han, L., Han, P., Wang, Y., Chen, Z., Ji, G., and Zheng, M. (2010).
Cardioprotection of ischemia/reperfusion injury by cholesterol-dependent MG53-mediated
membrane repair. Circulation research *107*, 76-83.
75. Milkovic, L., Zarkovic, N., Marusic, Z., Zarkovic, K., and Jaganjac, M. (2023). The 4-Hydroxynonenal–
Protein Adducts and Their Biological Relevance: Are Some Proteins Preferred Targets? Antioxidants *12*,
856.
76. Dalleau, S., Baradat, M., Guéraud, F., and Huc, L. (2013). Cell death and diseases related to oxidative
stress: 4-hydroxynonenal (HNE) in the balance. Cell Death & Differentiation *20*, 1615-1630.
77. Nijssen, J., Comley, L.H., and Hedlund, E. (2017). Motor neuron vulnerability and resistance in
amyotrophic lateral sclerosis. Acta neuropathologica *133*, 863-885.
78. Richter, K.N., Revelo, N.H., Seitz, K.J., Helm, M.S., Sarkar, D., Saleeb, R.S., d'Este, E., Eberle, J., Wagner,
E., and Vogl, C. (2018). Glyoxal as an alternative fixative to formaldehyde in immunostaining and super
‐resolution microscopy. The EMBO journal *37*, 139-159.
79. Dapson, R. (2007). Glyoxal fixation: how it works and why it only occasionally needs antigen retrieval.
Biotechnic & Histochemistry *82*, 161-166.
80. Pang, J., Zhao, X., Deng, F., Tsuchiya, D., Malloy, S., Parmely, T., Xie, T., and Wang, Y. (2022). A freeze-
substitution approach with solvent-based glyoxal fixative to prevent distortion of ocular structures.
Journal of Histotechnology *45*, 172-181.

81. Liao, Y., Smyth, G.K., and Shi, W. (2019). The R package Rsubread is easier, faster, cheaper and better
for alignment and quantification of RNA sequencing reads. Nucleic acids research *47*, e47-e47.
82. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol *15*, 550.
83. Marini, F., and Binder, H. (2019). pcaExplorer: an R/Bioconductor package for interacting with RNA-seq
principal components. BMC bioinformatics *20*, 1-8.
84. Ludt, A., Ustjanzew, A., Binder, H., Strauch, K., and Marini, F. (2022). Interactive and Reproducible
Workflows for Exploring and Modeling RNA‐seq Data with pcaExplorer, Ideal, and GeneTonic.
Current protocols *2*, e411.
85. Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R package for comparing biological
themes among gene clusters. Omics: a journal of integrative biology *16*, 284-287.
86. Weinmann, J., Weis, S., Sippel, J., Tulalamba, W., Remes, A., El Andari, J., Herrmann, A.-K., Pham, Q.H.,
Borowski, C., and Hille, S. (2020). Identification of a myotropic AAV by massively parallel in vivo
evaluation of barcoded capsid variants. Nature communications *11*, 5432.
87. Wang, B., Li, J., Fu, F., Chen, C., Zhu, X., Zhou, L., Jiang, X., and Xiao, X. (2008). Construction and
analysis of compact muscle-specific promoters for AAV vectors. Gene therapy *15*, 1489-1499.

882   **Figure and figure legends**

Figure 1

Figure 1. Differential expression of *Aldh3a1* and *Ankrd1* in different muscles from end-stage
G93A mice and WT littermates. (**A**) Comparing RNA levels of *Aldh3a1* and *Ankrd1* in Extensor
Digitorum Longus (EDL), soleus (SOL), diaphragm (DIA) to those in extraocular muscles (EOMs) by
qRT-PCR. N = 8 (4 pairs of male and 4 pairs of female). RQ, relative quantification (WT EDL samples

served as the reference group). *** P < 0.001; * P < 0.05; ns, not significant (Wilcoxon rank-sum test).
ANOVA P values are also shown. Please also see **Figure 1-figure supplement 1** and **Figure 1-**
Source Data 1. (**B**) Western blots of Aldh3a1, Ankrd1 and housekeeping protein Gapdh in different
muscles from end-stage G93A mice and WT littermates. The quantification results used WT EDL
samples as the reference group. N = 6 (3 pairs of male and 3 pairs of female). * P < 0.05; ** P < 0.01;
ns, not significant (Wilcoxon rank-sum test). ANOVA P values are also provided. Please also see
Figure 1-figure supplement 2 and **Figure 1-Source Data 2**. (**C**) Western blots of Aldh3a1, Ankrd1
and housekeeping protein Gapdh in the same muscle from end-stage G93A mice and WT littermates.
F: female, M: male.
897

Figure 2

A

Female | Male

EDL
WT | G93A | WT | G93A

SOL
WT | G93A | WT | G93A

DIA
WT | G93A | WT | G93A

EOM
WT | G93A | WT | G93A

B

Sham | SNT Day3 | SNT Day7 | SNT Day14

Ankrd1
DAPI

EDL

SOL

Figure 2. Global view of Ankrd1 immunostaining patterns in transverse sections of different
muscles. (**A**) Transverse sections of different muscles from end-stage G93A mice and WT controls
stained with anti-Ankrd1 antibodies and counterstained with DAPI. N = 6 (3 pairs of male and 3 pairs
of female). Scale bars, 200 μm. (**B**) Transverse sections of EDL and soleus muscles dissected from
WT mice (4-5 months of age) at Day 3, Day 7 and Day 14 post sciatic nerve transection and sham
operated controls stained with Ankrd1 antibodies and counterstained with DAPI. N = 6 (3 male and 3
female). Scale bars, 200 μm.

Figure 3

A

WT (female)   G93A (female)

Figure 3

A

WT (female)   G93A (female)

	Aldh3a1 (green), Sg2 (red), BTX (white)	DAPI		DAPI

EDL

SOL

DIA

EOM

B

G93A SOL (enlarged)

Aldh3a1

DAPI

Relative intensity — Aldh3a1, DAPI

Distance (µm)

0   6.25   12.5

1.2
0.9
0.6
0.3
0

G93A DIA (enlarged)

Aldh3a1

DAPI

Relative intensity — Aldh3a1, DAPI

Distance (µm)

0   6.25

1.2
0.9
0.6
0.3
0

Figure 3. Whole-mount immunostaining of Aldh3a1 in different muscles from end-stage G93A mice and WT littermates. (**A**) Representative compacted z-stack scan images of whole-mount EDL, soleus diaphragm extraocular muscles stained with antibodies against Aldh3a1, Sg2 (labeling axon terminals), Alexa Fluor conjugated α-Bungarotoxin (BTX, labeling AChRs on muscle membrane) and DAPI (labeling nuclei). Yellow arrows highlight nuclei with Aldh3a1 enrichment. Dashed yellow boxes denote regions enlarged in Panel B for kymographic measurement. N = 6 (3 pairs of male and 3 pairs of female). Scale bars, 50 µm. Please also see **Figure 3-figure supplement 1A**. (**B**) Profiling the relative intensity of Aldh3a1 and DAPI fluorescent signals along the strips denoted by dashed red boxes. Relative intensities are calculated as $(F - F_{min})/(F_{max} - F_{min})$. Also see **Figure 3-figure supplement 1B**.

Figure 4

Figure 4. Section immunostaining results of Aldh3a1 and Ankrd1 in different muscles from end-stage G93A mice. (**A-D**) Transverse sections of EDL, soleus, diaphragm and EOMs from end-stage G93A mice stained with antibodies recognizing Aldh3a1 and Ankrd1. Yellow arrows denote nuclear Aldh3a1 positive myofibers that are also positive for Ankrd1. Blue arrows denote nuclear Aldh3a1 positive myofibers negative for Ankrd1. N = 6 (3 male and 3 female). Scale bars: 50 μm.

925   Please also see **Figure 4-figure supplement 1** for section immunostaining results of Aldh3a1 in WT
926   muscles.

Figure 5

Figure 5. Time and muscle-dependent change of Aldh3a1 abundance after sciatic nerve
transection. (**A**) Comparing the changes of *Aldh3a1* and *Ankrd1* expression levels in EDL and SOL
from WT mice (4-5 months of age) collected 3 days, 7 days and 14 days after sciatic nerve
transection (SNT). Fold of change refers to expression level differences between muscles with SNT
and their sham-operated controls examined by qRT-PCR. N = 6 (3 male and 3 female). * P < 0.05; **
P < 0.01; ns, not significant (Wilcoxon rank-sum test). ANOVA P values are also shown. Please also
see **Figure 5-Source Data 1**. (**B, C**) Comparing relative changes of Aldh3a1 and Ankrd1 protein
(normalized by housekeeping protein Gapdh) in EDL and soleus (SOL) at 3 days, 7 days and 14 days
after SNT to corresponding sham-operated controls by Western blot. N = 6 (3 male and 3 female). * P
< 0.05; ** P < 0.01; ns, not significant (Wilcoxon rank-sum test). ANOVA P values are also shown.
Please also see **Figure 5-figure supplement 1** and **Figure 5-Source Data 2, 3**.

Figure 6

Figure 6. Examining subcellular distribution of Aldh3a1 protein in EDL and soleus muscles from WT mice with sciatic nerve transection. (**A**) Examining subcellular distribution of Aldh3a1 in EDL and soleus (SOL) muscles collected 3 days, 7 days and 14 days after SNT and sham-operated controls by whole-mount immunostaining. Antibodies against Sg2 labels axon terminals at NMJ and Alexa Fluor tagged BTX labels AChRs on the sarcolemma at NMJ. Regions denoted by dashed yellow boxes are enlarged in insets. N = 6 (3 male and 3 female). Scale bars: 50 µm. (**B**) Transverse sections of EDL and soleus muscles collected 14 days after SNT stained with Ankrd1, Aldh3a1 antibodies and DAPI. Yellow arrows denote nuclear Aldh3a1 positive myofibers that are also positive for Ankrd1. Blue arrows denote nuclear Aldh3a1 positive myofibers negative for Ankrd1. Scale bars: 50 µm.

Figure 7

Figure 7. Whole muscle RNA-Seq reveals potential mechanisms underlying differential regulation of *Aldh3a1* expression between muscles. (**A**) Principle component analysis of different muscle samples from end-stage G93A mice and age-matched WT controls (2 male and 2 female mice per group). Dashed circles highlight samples with similar expression profiles. (**B**) Top 20 most significant gene ontology biological processes for differentially expressed genes (log2 fold change > 0.5, p.adjust < 0.05) comparing soleus and EDL muscles, diaphragm and EDL muscles in WT mice, respectively. Arrows highlight processes related to fatty acid metabolism and oxidation. (**C**) Volcano plots of genes differentially expressed between different muscle groups. The three plots in the left-most panel highlight three EOM signature genes *Aldh3a1*, *Mymk* and *Pax6* by arrowheads. In the upper two panels on the right, genes related to fatty acid/lipid oxidation in the gene ontology analysis above are denoted by lines. The lower two panels demonstrate the higher expression of *Aldh3a1*, *Hmox1* and the lower expression of *Ankrd1* in soleus and diaphragm compared to the EDL muscles from G93A mice (arrowheads). (D) qRT-PCR results of *Hmox1* and *Aldh3a1* in EDL and soleus SC-derived myotubes treated with DMSO (Ctrl) or 10 µM sulforaphane (SFN) for 48 hrs (N = 5 culture replicate). ** P < 0.01; * P < 0.05; ns, not significant (Wilcoxon rank-sum test). ANOVA P values are also shown. Please also see **Figure 7-source data 2**.

Figure 8

Figure 8. Characterizing 4-HNE cytotoxicity to myotubes derived from EDL and soleus satellite cells. (**A**) 4-HNE levels (normalized to protein amount) in EDL and soleus (SOL) muscles from end-stage G93A mice and age-matched WT controls (3 male and 3 female mice/group) measured by ELISA. ** P < 0.01; ns, not significant (Wilcoxon rank-sum test). ANOVA P value is also shown. Please also see **Figure 8-Source Data 1.** (**B**) Myotubes differentiated from EDL or soleus SCs isolated from 4-month-old WT mice for 5 days were treated with 0, 7.5, 15 or 30 µM 4-HNE for 2 hrs, respectively. Afterwards the myotubes were cultured for another 16 hrs in regular differentiation medium (as apoptosis takes time to occur) before fixation, TUNEL staining and immunostaining against MF-20 (labeling the cytosolic part of myotubes). DAPI counterstaining labelled the nuclei. N =

3 culture replicates. Regions denoted by dashed yellow boxes are enlarged in insets. Scale bars: 100
µm. (**C**) Averaged TUNEL positive percentage of nuclei in myotubes treated with different
concentrations of 4-HNE (N = 5 culture replicates, 4 images analyzed and averaged for each culture
replicate). ** $P < 0.01$; ns, not significant (t.test). ANOVA P values are also shown. Please also see
Figure 8-Source Data 2. (**D**) Measuring 4-HNE cytotoxicity to myotubes using LDH leakage-based
assay (N = 5 culture replicates). ** $P < 0.01$; * $P < 0.05$; ns, not significant (Wilcoxon rank-sum test).
ANOVA P values are also shown. Please also see **Figure 8-Source Data 3**.

Figure 9

Figure 9. Transducing myotubes with AAV-Aldh3a1 protects against 4-HNE cytotoxicity. (**A**) Myotubes differentiated from EDL or soleus SCs for 1 day were incubated with or without AAV-Aldh3a1 for 2 days, followed by another 2-day culture in regular differentiation medium before protein extraction and Western blot. N = 3 culture replicates. ** P < 0.01; * P < 0.05 (t-test). Please also see **Figure 9-Source Data 1**. (**B**) Myotubes differentiated from EDL or soleus satellite cells isolated from

4-month-old female WT mice with or without AAV-Aldh3a1 transduction were treated with 30 µM 4-
HNE for 2 hrs. Afterwards the myotubes were cultured for another 16 hrs in regular differentiation
medium before fixation, TUNEL staining, immunostaining against MF-20 (labeling the cytosolic part of
myotubes) and Aldh3a1. DAPI counterstaining labelled the nuclei. N = 3 culture replicates. Regions
denoted by dashed yellow boxes are enlarged in insets. Scale bars: 100 µm. (**C**) Quantification
results of averaged TUNEL positive percentage of nuclei in myotubes (N = 3 culture replicates, 4
images analyzed and averaged for each culture replicate). ANOVA P values are also shown. * P <
0.05 (t-test). Please also see **Figure 9-Source Data 2**. (**D**) Measuring the protective effect of AAV-
Aldh3a1 against 4-HNE cytotoxicity to myotubes using LDH leakage-based assay (N = 6 culture
replicates). ** P < 0.01; ns, not significant (Wilcoxon rank-sum test). ANOVA P values are also shown.
Please also see **Figure 9-Source Data 3**.

Figure 10

A

GFP-MG53

B

① ② ③

C

ANOVA P=2.45e-08

Max F/F_0 on membrane

AAV-Aldh3a1 − − +
4-HNE − + +

D

Percentage

AAV-Aldh3a1 − − +
4-HNE − + +

■ F_{180} higher than F_0 ■ F_{180} lower than F_0

1003

Figure 10. AAV-Aldh3a1 protects against 4-HNE compromised stabilization of MG53 repair patches in partially permeabilized myotubes. (**A**) Time lapse imaging of saponin-induced formation of GFP-MG53 repair patches on the plasma membrane in soleus SC-derived myotubes with or without AAV-Aldh3a1 transduction and/or 4-HNE treatment (30 µM for 2 hrs). Regions denoted by dashed yellow boxes are enlarged in insets. Arrows highlight myotubes whose GFP-MG53 intensity on the plasma membrane were profiled in panel B. Scale bars: 50 µm. Please also see Videos 1-3. (**B**) Plotting background-corrected, initial timepoint intensity normalized signal of GFP-MG53 on plasma membrane (F/F_0) over time for the three myotubes indicated in Panel A. Arrowheads denote the time point when saponin was applied. (**C**) Maximum F/F_0 of recorded myotubes under different treatment conditions (N = 35 for each group). **** P < 0.0001; ns, not significant (Wilcoxon rank-sum test). ANOVA P values are also shown. Please also see **Figure 10-Source Data 1**. (**D**) Percentage stacked column chart highlights the proportion of recorded myotubes with F_{180} (end of recording) higher than F_0 (blue) and the proportion of those with F_{180} lower than F_0 (orange) in different treatment groups. Please also see **Figure 10-source data 1**.

Video 1. Time lapse imaging of GFP-MG53 transfected myotubes without 4-HNE or AAV-Aldh3a1 treatment.

Video 2. Time lapse imaging of GFP-MG53 transfected myotubes treated with 30 µM 4-HNE for 2 hours before recording.

Video 3. Time lapse imaging of GFP-MG53 transfected, AAV-Aldh3a1 transduced myotubes treated with 30 µM 4-HNE for 2 hours before recording.

Figure 1-figure supplement 1

Figure 1-figure supplement 1. Additional qRT-PCR and Western blot quantification results for
different muscles from end-stage G93A mice and WT littermates. (**A**) qRT-PCR (upper) and
Western blot (lower) relative quantification results of *Aldh3a1* (against housekeeping gene *Gapdh*) in
EDL, soleus (SOL), diaphragm (DIA) and extraocular muscles (EOM) from end-stage G93A mice and
WT littermates. (**B**) qRT-PCR (upper) and Western blot (lower) relative quantification results of
Ankrd1 in different muscles from end-stage G93A mice and WT littermates. (**C-F**) qRT-PCR relative
quantification results of *Aldh1a1*, *Aldh2*, *Mymk* and *Hmox1*. *** P < 0.001; ** P < 0.01; * P < 0.05; ns,
not significant (Wilcoxon rank-sum test). ANOVA P values are also shown. N = 8 (4 pairs of male
mice and 4 pairs of female mice). RQ, relative quantification. Please also see **Figure 1-Source Data**
1.

Figure 1-figure supplement 2

1038
1039 **Figure 1-figure supplement 2. Original images for Western blots in Figure 1.** (**A**) Original images
1040 for Western blots in Figure 1B. Red arrows indicate Aldh3a1 (51 kDa) or Ankrd1 (36 kDa) bands. Blue
1041 arrows indicate Gapdh (36 kDa) bands. Green arrows indicate β-actin (42 kDa) bands. Orange

1042   arrows indicate hSod1 (16-20 kDa) bands (the upper band is human Sod1^{G93A}, the lower band is
1043   mouse endogenous Sod1). (**B**) Original images for Western blots in Figure 1C.
1044
1045

Figure 3-figure supplement 1

A

WT (male)   G93A (male)

B   G93A SOL (enlarged)   G93A DIA (enlarged)

Figure 3-figure supplement 1. Additional images of Aldh3a1 whole-mount immunostaining in
different muscles from end-stage G93A mice and WT littermates. (**A**) Representative compacted
z-stack scan images of whole-mount EDL, soleus diaphragm extraocular muscles stained with
antibodies against Aldh3a1, Sg2 (labeling axon terminals), Alexa Fluor conjugated α-Bungarotoxin
(BTX, labeling AChRs on muscle membrane) and DAPI (labeling nuclei). N = 6 (3 pairs of male and 3
pairs of female). Scale bars, 50 μm. Yellow arrows highlight nuclei with Aldh3a1 enrichment. Dashed
yellow boxes denote regions enlarged in Panel B for kymographic measurement. (**B**) Profiling the
relative intensity of Aldh3a1 and DAPI fluorescent signals along the strips denoted by the red boxes.
Relative intensities are calculated as $(F - F_{min})/(F_{max} - F_{min})$.

1056

Figure 4-figure supplement 1

 Figure 4-figure supplement 1. Section immunostaining results of Aldh3a1 in different muscles
 from WT mice 4-5 months of age. (**A**) Transverse sections of EDL, soleus, diaphragm and EOMs
 from WT mice stained with antibodies recognizing Aldh3a1. N = 6 (3 male and 3 female). Scale bars:
 50 μm.

Figure 5-figure supplement 1

1063 **Figure 5-figure supplement 1. Original images of Western blots in Figure 5.** Red arrows indicate
1064 Aldh3a1 (51 KDa) or Ankrd1 (36 kDa) bands. Blue arrows indicate Gapdh (36 kDa) bands. Green
1065 arrows indicate β-actin (42 kDa) bands.

Figure 7-figure supplement 1

Figure 7-figure supplement 1. Gene ontology analysis and heatmaps of genes differentially expressed between different groups of muscles in WT mice. (**A**) Top 20 biological processes in gene ontology analysis for genes expressed higher in EOMs than other muscles from WT mice. Processes related to eye development are denoted. (**B**) Axon guidance/neuron migration related biological processes highlighted by gene ontology analysis for genes expressed higher in EOMs than other muscles from WT mice. (**C**) Heatmaps of axon guidance related DEGs comparing EOMs to other muscles from WT mice. (**D**) Heatmaps of lipid oxidation related DEGs comparing soleus and diaphragm to EDL muscles from WT mice.

Figure 1-Source Data 1. qRT-PCR results for *Aldh1a1*, *Aldh2*, *Aldh3a1*, *Ankrd1*, *Mymk* and *Hmox1* expression levels in whole muscles of different origins.

Figure 1-Source Data 2. Western blot quantification results for Aldh3a1 and Ankrd1 protein levels in whole muscles of different origins.

Figure 5-Source Data 1. qRT-PCR results of the fold of change of *Aldh3a1* and *Ankrd1* expression levels in whole muscles with SNT compared to the sham operated controls.

Figure 5-Source Data 2. Western blot quantification results of the fold of change of Aldh3a1 protein levels in whole muscles with SNT compared to the sham operated controls.

Figure 5-Source Data 3. Western blot quantification results of the fold of change of Ankrd1 protein levels in whole muscles with SNT compared to the sham operated controls.

Figure 7-Source Data 1. Entrez IDs, symbols, log2 fold changes and adjusted p values of the 866 EOM signature genes.

Figure 7-Source Data 2. qRT-PCR results for *Hmox1* and *Aldh3a1* expression levels in EDL and soleus-SC derived myotubes with and without SFN treatment.

Figure 8-Source Data 1. 4-HNE levels in EDL and soleus muscles from end-stage G93A mice and age-matched WT controls measured by ELISA. .

1099

Figure 8-Source Data 2. Quantification results of the percentages of TUNEL positive nuclei in myotubes treated with different concentrations of 4-HNE.

1102

Figure 8-Source Data 3. Quantification results of LDH leakage percentages of myotubes treated with different concentrations of 4-HNE.

1105

Figure 9-Source Data 1. Western blot quantification results for Aldh3a1 protein levels in myotubes with or without transduction of AAV-Aldh3a1.

1108

Figure 9-Source Data 2. Quantification results of the percentages of TUNEL positive nuclei in myotubes with or without transduction of AAV-Aldh3a1 before 4-HNE treatment.

1111

Figure 9-Source Data 3. Quantification results of LDH leakage percentages of myotubes with or without transduction of AAV-Aldh3a1 before 4-HNE treatment.

1114

Figure 10-Source Data 1. Ratios between maximum F and F_0 in each recorded myotubes transfected with GFP-MG53 and whether the F_{180} is higher or lower than F_0.

1117