The complete genome sequence of the crayfish pathogen Candidatus

***Paracoxiella cheracis* n.g. n.sp. provides insight into pathogenesis and the**

phylogeny of the Coxiellaceae family.

Danielle J. Ingle[1#], Calum J. Walsh[1,2#], Genevieve R. Samuel[3], Ryan R. Wick[1,2],

Nadav Davidovich[4], Eleonora Fiocchi[5], Louise M. Judd[2], Jennifer Elliman[6], Leigh

Owens[6], Timothy P. Stinear[1,2], Andrea Basso[6], Tobia Pretto[6], Hayley J. Newton[3]

[1] Department of Microbiology and Immunology, The University of Melbourne at The

Peter Doherty Institute for Infection and Immunity, Melbourne, Australia

[2] Centre for Pathogen Genomics, University of Melbourne, Parkville, Victoria

Australia

[3] Infection Program, Monash Biomedicine Discovery Institute and Department of

Microbiology, Monash University, Clayton, VIC 3800, Australia

[4] Israeli Veterinary Services, Bet Dagan, Israel

[5] Istituto Zooprofilattico Sperimentale delle Venezie, Legnaro, Italy

[6] College of Public Health, Medical and Veterinary Sciences, James Cook University,

1 Solander Road, Townsville, QLD, 4811, Australia

[#] authors contributed equally

Corresponding author:

Hayley J. Newton

Ph: +619902 9146

Email: hayley.newton@monash.edu

Running Title: Analysis of the *Paracoxiella cheracis* genome

Keywords: Candidatus *Paracoxiella cheracis*, Nanopore long-read sequencing,

Coxiellaceae, Dot/Icm type 4 secretion system, effector proteins

ABSTRACT

The Coxiellaceae bacterial family, within the order Legionellales, is defined by a collection of poorly characterized obligate intracellular bacteria. The zoonotic pathogen and causative agent of human Q fever, *Coxiella burnetii*, represents the best characterized member of this family. Coxiellaceae establish replicative niches within diverse host cells and rely on their host for survival, making them challenging to isolate and cultivate within a laboratory setting. Here we describe a new genus within the Coxiellaceae family that has been previously shown to infect economically significant freshwater crayfish. Using culture-independent long-read metagenomics, we reconstructed the complete genome of this novel organism and demonstrate that the previously referred to as Candidatus *Coxiella cheraxi* represents a novel genus within this family, herein denoted Candidatus *Paracoxiella cheracis*. Interestingly, we demonstrate that Candidatus *P. cheracis* encodes for a complete, putatively functional Dot/Icm type 4 secretion system that likely mediates the intracellular success of this pathogen. In silico analysis defined a unique repertoire of Dot/Icm effector proteins and highlighted homologues of several important *C. burnetii* effectors including a homologue of CpeB that was demonstrated to be a Dot/Icm substrate in *C. burnetii*.

IMPORTANCE

Using long-read sequencing technology we have uncovered the full genome sequence of Candidatus *Paracoxiella cheracis*, a pathogen of economic importance in aquaculture. Analysis of this sequence has revealed new insight into this novel member of the Coxiellaceae family, demonstrating that it represents a new genus within this poorly characterized family of intracellular organisms. Importantly, the genome sequence reveals invaluable information that will support diagnostics and potentially both preventative and treatment strategies within crayfish breeding facilities. Candidatus *P. cheracis* also represents a new member of Dot/Icm pathogens that rely on this system to establish an intracellular niche. Candidatus *P. cheracis* possesses a unique cohort of putative Dot/Icm substrates that constitute a collection of new eukaryotic cell biology manipulating effector proteins.

INTRODUCTION

Australian native freshwater redclaw crayfish, *Cherax quadricarinatus*, are bred for commercial market. Aquaculture of these animals has led to the observation of infectious diseases that impact both wild and farmed *C. quadricarinatus* (reviewed in (1)). In particular, a Rickettsial-like intracellular bacterium, designated TO-98, isolated from infected crayfish was recognized as the likely causative agent of mass mortality in farmed redclaw crayfish in the early 1990s (2, 3). Subsequent investigations were able to recapitulate disease in experimentally infected crayfish and demonstrate that TO-98 causes a lethal infection in redclaw crayfish (4). This study conducted 16s rRNA sequence analysis and revealed that TO-98 was closely related to the zoonotic human pathogen *Coxiella burnetii*, coining the name Candidatus *Coxiella cheraxi* (4).

The initial observation that this pathogen is a member of the Coxiellaceae family has been confirmed by examination of additional DNA sequences including comparison of specific gene sequences (5) and MinION sequencing data showing fragments of DNA with closest similarity to *C. burnetii* and *Coxiella*-like endosymbionts (C-LEs) (6). However, this fragmented genomic data does not allow complete and comprehensive understanding of the relationship between *C. burnetii* and Candidatus *C. cheraxi*.

C. burnetii, the causative agent of human Q fever, has a large reservoir in domesticated ruminants (7). *C. burnetii* is a Gammaproteobacteria within the order Legionellales and family Coxiellaceae. While the order Legionellales is best known for the two major human pathogens, *Legionella pneumophila* and *C. burnetii*, these bacteria are distantly related and the order is poorly characterized, including over 450 uncultured genera (8). The diversity of Legionellales has been revealed by increased sampling of different environments and ecological surveys employing metagenomic techniques that have demonstrated the presence of Legionellales within arthropods, amoeba and aquatic environments (reviewed in (9)). Most of these newly discovered Legionellales are likely nonpathogenic bacteria that have evolved mutualistic relationships with non-vertebrate hosts. However, recent data suggests that some poorly characterized Legionellales harbor the potential to be zoonotic

pathogens and/or pathogenic to important animal species. This includes a recently

identified *Coxiella* species found in the placenta of fur seals (10).

Essential to the ability of *C. burnetii* and *L. pneumophila* to replicate intracellularly,

and cause disease, is the Dot/Icm type IVB secretion system (T4SS) (11-14). This

multiprotein apparatus is responsible for the delivery of a large cohort of effector

proteins into the host cell where they modulate a range of cellular functions and

pathways (reviewed in (15)). Dot/Icm-deficient *L. pneumophila* are rapidly destroyed

via lysosomal degradation as many of the effectors act to remodel the phagosome

and block endocytic maturation (reviewed in (16)). In contrast, internalized *C. burnetii*

is trafficked through the endocytic pathway before the lysosomal environment

triggers activation of the Dot/Icm system (17, 18). Dot/Icm-deficient *C. burnetii* are

not destroyed by the lysosome but they are also incapable of intracellular replication

(11, 12). Interestingly, comparative genomic studies show that the Dot/Icm system is

lost or pseudogenized in C-LE, indicating its association with pathogenesis (19).

Despite *L. pneumophila* and *C. burnetii* relying on the same key virulence factor, the

cohort of effector proteins translocated by this secretion system are pathogen

specific. This genetic divergence is represented by phenotypic divergence through

the establishment of distinct intracellular niches. The biochemical and functional

characterization of these effector proteins is an active area of research as these

novel proteins facilitate insight into both mechanisms of pathogenesis and novel

strategies to manipulate eukaryotic cell biology.

Here we report the complete genome sequence of Candidatus *C. cheraxi* revealing a

more distant relationship to *C. burnetii* than expected, notably that the Candidatus *C.*

cheraxi represents a novel genus within Coxiellaceae. We propose to rename this

crayfish pathogen Candidatus *Paracoxiella cheracis.* Interestingly, we report that this

novel Coxiellaceae family member possesses a Dot/Icm T4SS most closely

associated with *C. burnetii. In silico* analysis allowed us to identify a large cohort of

putative effectors of this secretion system with only 12 out of 238 predicted effectors

sharing identity to known *C. burnetii* T4SS effectors. We have been able to

demonstrate translocation of the Candidatus *P. cheracis* CpeB homologue via the *C.*

burnetii Dot/Icm system. This study provides additional insight into the poorly

characterized Coxiellaceae family and demonstrates that the Dot/Icm secretion

system is a common tool used by these bacteria to communicate with their host. It

also demonstrates the utility of long-read sequencing approaches for generating a

complete metagenome-assembled genome for an obligate intracellular pathogen.

METHODS

Collection and preparation of genetic material.

Samples of hepatopancreas from infected redclaw crayfish were collected during an outbreak of crayfish rickettsiosis, which occurred after the crayfish were imported from Australia to Israel in 2019 (20). Briefly, 10,000 juvenile Redclaw crayfish (2–3 weeks old) were imported to Israel and quarantined in a facility. From April to July, juvenile mortality sharply increased despite the absence of clinical signs or gross lesions. Specimens collected for histological and molecular evaluations were sent to the Istituto Zooprofilattico Sperimentale delle Venezie in Legnaro, Italy, and revealed a severe infection by Candidatus *C. cheraxi.* DNA was extracted from these samples, as previously described (20), and sent to Monash University, Australia, for further analysis.

Long-read sequencing and genome assembly

Metagenomic sequencing of extracted DNA was performed on two separate sequencing runs. The first run was performed on an Oxford Nanopore Technologies (ONT) MinION while the second employed a PromethION. Both runs were carried out on R10.4.1 flow cells. Basecalling and quality filtering was performed by Dorado (v. 0.5.0) using models dna_r10.4.1_e8.2_400bps_sup@v4.1.0 and dna_r10.4.1_e8.2_400bps_sup@v4.3.0 respectively. Reads shorter than 1kbp were discarded. Kraken2 (21) was used for initial read-level classification of metagenomic data.

Initial assemblies were generated using Flye (22), MetaMDBG (23), Miniasm (24), and Raven (25), each of which produced a single, continuous sequence representing the bacterial chromosome, and most of which contained two medium-sized plasmids at a similar read depth to the chromosome. For each assembly, the length-filtered reads were mapped to the chromosome and plasmids Minimap2 (26) and mapped reads were retained for downstream analysis. The chromosomal and plasmid reads were then downsampled to 500× coverage using Filtlong (github.com/rrwick/Filtlong) and combined into a single high-quality assembly with Trycycler (27) incorporating 24 separate assemblies from six assemblers: Canu (28), Flye (22), Miniasm (24), NECAT (29), NextDenovo (30), and Raven (25). The final complete genome assembly was annotated with Bakta v1.9.1 (31). Full details of the individual steps

for quality controls, kraken assignment and genome assembly are available in

Supplementary File 1.

Phylogeny of Coxiellaceae Family

The Genome Taxonomy Database Toolkit (GTDB-Tk) (32) classified this genome as

Coxiellaceae. Representative isolates from the order Coxiellales

(gtdb.ecogenomic.org/tree?r=o__Coxiellales) were used in comparative analyses to

infer the relative location of Candidatus *P. cheracis* in the order. Given the large

number of Coxiellales genomes in the GTDB database, the majority of which belong

to the clinically important species *Coxiella burnetii*, the dataset was first dereplicated

by Assembly Dereplicator (github.com/rrwick/Assembly-Dereplicator) to a Mash

distance of 0.001, resulting in a representative set of 60 genomes capturing the

available diversity of the Order. A phylogeny was inferred using GTDB-Tk (33), a

marker gene-based tool for taxonomic classification and phylogenetic placement of

prokaryotic genomes based on the Genome Taxonomy Database (GTDB) taxonomy

system (34) was used to infer a phylogeny for the representative 60 genomes and to

taxonomically classify our long-read assembly.

Investigation of the complete genome for features of interest.

The genome content of Candidatus *P. cheracis* was explored using publicly available

tools. First, known antimicrobial resistance (AMR) determinants were screened with

abritAMR (v1.0.14) with the AMRfinderPlus database (v2022-08-09.1) with no

species flag and default parameters (35). Mob-typer, which is part of Mob-Suite, was

used to identify plasmid replicons and mob genes with default parameters (36, 37).

Phage defense mechanisms were identified with PADLOC v2 using the annotated gff

file as input and default parameters (38).

Putative Dot/Icm effectors were screened in the complete genome with Bastion4 (39)

and T4Sepp (40) tools. The predictions were compared, with hits for putative IS

elements not included in further analysis due to the likelihood of being false positive

results.

Components of the T4SS were identified with SECRET4 (41). T4SSs were also

identified in selected, publicly available genomes that have been established to have

functional T4SSs. These included two *C. burnetii* genomes as the type strains for the

species (RSA493 accession: AE016828.3, and Dugway accession: CP000733.1),

two genomes from the *Legionella* genus, *Legionella pneumophila* subsp*.*

pneumophila str. Philadelphia 1 (accession: AE017354.1) and *Legionella*

longbeachae NSW150 (accession:NC_013861.1)(42) and the Candidatus

Rickettsiella viridis Ap-RA04 genome (accession: AP018005.1). We also screened

representative isolates with complete genomes from *Coxiella* like-endosymbionts (C-

LEs) (accessions: CP021379.1, CP011126.1, CP033868.1 and CP064834.1) with

SECRET4 confirming the reported lack of a complete T4SS in these genomes (43).

Protein identity between the six genomes was undertaken with *blastp* using the

Dot/Icm proteins from *C. burnetii* RSA493 (accession: AE016828.3) as a query

sequence. For the three *Coxiella* genomes, the output was filtered to the top hit for

each with a bitscore >100. A modified approach was undertaken for the two

Legionella and Candidatus *Rickettsiella viridis* genomes due to the lower sequence

homology. For these three genomes, the results were filtered to the loci that encode

the T4SS and the hits with highest bitscore for each protein was included.

Visualization of the Candidatus *P. cheracis* genome with features of interest

including phage defense, T4SS and putative Dot/Icm effectors was done in BRICK

(github.com/esteinig/brick). The phylogenetic tree was visualized in R using ape (44)

and ggtree (45). The R package gggenomes was used to plot the T4SS

(github.com/thackl/gggenomes) and pheatmap was used to visualize the protein

identity (github.com/raivokolde/pheatmap).

Cultivation of bacterial and tissue culture cells

C. burnetii Nine Mile phase II (NMII) strain RSA439 clone 4 was axenically cultivated

in liquid ACCM-2 or ACCM-2 agarose at 37°C in 5% CO_2 and 2.5% O_2 as previously

described (46). When required chloramphenicol was added to *C. burnetii* cultures at

3 μg/ml. HeLa CCL-2 cells (ATCC, VA) were maintained in Dulbecco's Modified

Eagle's Media (DMEM) supplemented with 10% heat inactivated fetal bovine serum
at 37°C in 5% CO_2.

Construction of *C. burnetii* translocation reporter strains

The open reading frame encoding for the Candidatus *P. cheracis* CpeB homologue
(*Pc*CpeB) was synthesized by IDT and cloned into pJB-CAT:BlaM (47) using BamHI
and XhoI restriction enzymes. The resulting pBlaM-*Pc*CpeB was introduced into
C. burnetii strains using a standard electroporation protocol (11). Following selection
of transformants, expression of BlaM fusion proteins was confirmed using an anti-
BlaM (1:1500, QED Biosciences). *C. burnetii* strains carrying either pJB-CAT:BlaM
or pJB-CAT:BlaM:CBU0021 were used as negative and positive translocation
controls (48).

BlaM reporter translocation assay

Translocation assays were conducted as previously reported (17, 49). HeLa CCL2
cells were seeded in 96-well black, clear bottom tissue culture plates at a density of
10^4 per well. Approximately 24 h later, cell monolayers were infected with *C. burnetii*
strains at an MOI of 300. Following a 22 h incubation, wells were loaded with CCF2-
AM substrate from the LiveBLAzer FRET loading kit (Thermo Fisher Scientific) and
incubated at room temperature for 2 h. Wells were excited at 415 nm and the
emission at 450 nm (blue) and 520 nm (green) was collected using a Clariostar
fluorescence plate reader. Representative images were acquired using an Invitrogen
EVOS™ FL Imaging System, images taken at 10✕ objective.

Data availability

The complete genome and the long-read data are available at BioProject
PRJNA1189271.

RESULTS

A putative new genus in the Coxiellaceae family

Here we describe the complete genome of a putative new genus within
Coxiellaceae, Candidatus *C. cheraxi,* that is the likely causative agent of mass
mortality events in *C. quadricarinatus*. The Candidatus *C. cheraxi* genome consisted
of one chromosome (2,233,862 bp) and two smaller plasmids (59,783 and 34,021
bp). The chromosome is similar in size to previously sequenced *C. burnetii* isolates
ranging from approximately 1.9 Mb to 2.2 Mb (50). To confirm that this organism is
related to the original Candidatus *C. cheraxi* TO-98 organism, the 21 available
sequence reads from TO-98 (6) were mapped to this genome with an average
identify of 89%. This is a high level of similarity, even with the TO-98 dataset
produced using older minion chemistry (R9.4.1), provided confidence that this
genome is representative of the same species as TO-98.

The Candidatus *C. cheraxi* genome was classified as Coxiellaceae by GTDB-Tk
(32), meaning that the novel genome could be classified to the family level but no
further, suggesting that this genome represents a novel genus, supported by a
Relative Evolutionary Divergence (RED) value of 0.79373 (32). Based on this data
we propose to consider this Coxiellaceae as a novel genus: Candidatus *Paracoxiella*
cheracis, also adopting the adjusted genus nomenclature (51).

A phylogenetic tree inferred from bacteria-specific marker genes placed Candidatus
P. cheracis near genomes of other Coxiellaceae bacteria (**Figure 1**) isolated from
similar aquatic ecological niches and distant to *C. burnetii* (largely associated with
mammalian hosts) and the *Coxiella*-like endosymbionts (C-LEs) that are associated
with ticks as hosts (**Figure 1**). Interestingly, Kraken2 classification of the
chromosomal reads using a GTDB-based database identified multiple bacterial
families in the read set. These include 10.6% Alteromonadaceae, 4.1%
Vibrionaceae, 3.7% Enterobacteriaceae, 2.9% Burkholderiaceae, 2.5%
Legionellaceae, 2.3% Moraxellaceae and only 8.5% Coxiellaceae (**Supplementary
File 2**), which are all members of the Gammaproteobacteria class. This
demonstrates that Kraken2 results for an isolate in a novel taxon not represented in
the database can be misleading and may falsely suggest a mixed sample. However,

assembly of the long-read data successfully recovered the bacterial chromosome
and two plasmids.

The complete genome of Candidatus *P. cheracis* was explored for antimicrobial
resistance, phage defense mechanisms, the Dot/Icm T4SS and putative effectors of
this system (**Figure 2**). No known AMR mechanisms were identified, consistent with
the AMR profiles of *C. burnetii* genomes. No replicon or mob genes were detected in
the two plasmids. Further, these two plasmid sequences had no homology to publicly
available data on NCBI. Several phage defense systems were identified on the
chromosome in the novel genome. Two SoFic defense systems and the AbiE
system, comprised of AbiEi and AbiEii components, were identified in the genome.
The remaining ten defense systems identified were from the curated PADLOC
database. Multiple putative effectors were identified (expanded upon in Predicted
substrates of the *C. cheraxi* Dot/Icm secretion system), in addition to many copies
and variants of a transposase that shares identity with a transposase from a
symbiont of *Solemya velum*, the Atlantic awning clam [ref, accession:
JRAA00000000] (52).

Detection and analysis of the Dot/Icm locus encoded by Candidatus *P.
cheracis*
The Dot/Icm system was detected in the Candidatus *P. cheracis* genome,
suggesting that the isolate has a functional T4SS. The site of integration into the
chromosome and gene arrangement were similar to the two isolates of *C. burnetii*
examined here (**Figure 3A**). The two *C. burnetii* genomes have additional genes
between *icmT* and *icmQ,* some of which have been shown to be effector proteins
(53), which are lacking in the Candidatus *C. cheraxi* genome. Additionally, *icmT* and
icmS were encoded in the reverse orientation in the Candidatus *P. cheracis* genome
compared to the two *C. burnetii* genomes. These two genes were in the same
orientation as the two *Legionella* and Candidatus *R. viridis* genomes despite the
Dot/Icm system being fragmented into multiple regions in these three genomes. Of
note, the protein homology between 21 genes of the Dot/Icm system differed within
the six genomes by species and genus (**Figure 3B**). The proteins from the *C.*
burnetii Dugway isolate were homologous to the Dot/Icm proteins of *C. burnetii*

RSA493 (>99% protein identity for all). In contrast, the protein identity for the Dot/Icm
proteins in Candidatus *P. cheracis* varied between 52% (IcmG) and 86% (IcmB)
identity relative to the *C. burnetii* RSA493 proteins. This was higher than the protein
similarity of the Dot/Icm proteins from the two *Legionella* and the Candidatus *R.*
viridis genomes which ranged between 24% (IcmX in *L. longbeachae*) to 67% (DotB
for Candidatus *R. viridis*). As previously reported, the Dot/Icm system was not
detected in the three C-LEs genomes.
343

Predicted substrates of the Candidatus *P. cheracis* Dot/Icm secretion system
Given the conservation of genes encoding for the Dot/Icm apparatus we
hypothesized that Candidatus *P. cheracis* possesses a unique repertoire of putative
T4SS substrates. To identify these proteins, two effector prediction pipelines,
Bastion4 and T4SEpp, were employed. Bastion4, bacterial secretion effector
predictor for T4SS, is an ensemble effector predictor based on six distinct machine
learning models (39). More recently, T4SEpp was developed using an integrated
pipeline that incorporates homology-based predictions with machine learning models
to generate a prediction score for the likelihood of a protein being a T4SS substrate
(40). Combined, these tools predicted 355 putative Dot/Icm substrates within the
Candidatus *P. cheracis* genome. Bastion4 predicted 296 effectors and T4SEpp
predicted 114 with 55 putative effectors predicted by both approaches
(**Supplementary File 3**). Bastion4 predicted 117 instances of IS elements as
putative effectors which are likely false positives bringing the pool of potential
Dot/Icm effectors to 238. Of particular interest, 66 putative effectors (25 predicted by
both tools) have no significant similarity to protein sequences found in NCBI.
360

Candidatus *P. cheracis* CpeB, *Pc*CpeB, can be translocated by the *C. burnetii*
Dot/Icm system.
Each of the predicted Dot/Icm effectors were examined for similarity to known
Dot/Icm effectors of *L. pneumophila* and *C. burnetii*. No putative effectors showed
similarity within the *L. pneumophila* effector cohort however 12 putative effectors
have similarity to putative and experimentally characterized *C. burnetii* effectors
(**Table 1**).
368

To test the feasibility of using *C. burnetii* as a surrogate bacterium to demonstrate

Dot/Icm-dependent translocation of putative Candidatus *P. cheracis* effectors, the

BlaM translocation reporter assay was employed. This assay is routinely used to

demonstrate Dot/Icm substrate translocation into host cells and relies on bacterial

delivery of a β-lactamase (BlaM) enzyme into the host cytosol where its activity can

be measured by cleavage to the BlaM substrate CCF2-AM (11). Dot/Icm-dependent

translocation of the Candidatus *P. cheracis* homologue of *C. burnetii* effector CpeB

(*Pc*CpeB) was confirmed (**Figure 4A and B**). *C. burnetii* WT and Dot/Icm-deficient

(*icmL::*Tn) strains were engineered to express BlaM alone (negative control), BlaM-

CvpB, a well characterized *C. burnetii* Dot/Icm effector (positive control) (48, 54-56)

or BlaM-*Pc*CpeB (**Fig 4C**). HeLa cells were infected with these *C. burnetii* strains for

22 h before addition of CCF2-AM and subsequent detection of fluorescence

emission via both a fluorescence plate reader (**Fig 4A**) and microscopy (**Fig 4B**).

BlaM-*Pc*CpeB was translocated in a Dot/Icm-dependent manner. Interestingly,

comparison of the amino acid sequences of *C. burnetii* CpeB (*Cb*CpeB) and *Pc*CpeB

showed only 29% identity however the region determined to mediate interaction

between *Cb*CpeB and Rab11 shows greater conservation potentially indicating

functional similarity (purple box, **Fig 4D**) (57).

DISCUSSION

In the absence of live samples of Candidatus *P. cheracis* n.g. n. sp.*,* progressing our understanding of this novel crayfish pathogen is limited to the exploration of genetic material. This report of the complete genome sequence of this pathogen provides both an opportunity to expand our understanding of the evolutionary relationships between members of the family Coxiellaceae and formulate hypotheses regarding the host-pathogen interactions mediated by this organism.

The phylogenetic placement, ANI to reference genomes, and Relative Evolutionary Divergence of Candidatus *P. cheracis* provides evidence that this novel genome represents a new genus within the Coxiellaceae family and is more distantly related to both *C. burnetii* and C-LEs than previously appreciated. Of note, the Dot/Icm T4SS detected in Candidatus *P. cheracis* was found to have inserted into the chromosome near the same genes as in *C. burnetii* genomes. From this, we hypothesize that the Dot/Icm T4SS may have been acquired in the most recent common ancestor of both Candidatus *P. cheracis, C. burnetii* and C-LE. Moreover, the loss of the Dot/Icm T4SS in the C-LEs which are associated with invertebrate hosts suggests that the Dot/Icm T4SS did not confer a fitness advantage to warrant being maintained in C-LEs, which are maternally inherited in ticks (19).

The Dot/Icm T4SS has co-evolved in these different genetic backgrounds, resulting in the observed differences in protein identity between Candidatus *P. cheracis* and the two *C. burnetii* reference genomes, with adaptation of the effectors to the different host niches. The observed differences in Dot/Icm protein identity between these genomes is greater than the average values recently reported for 58 *Legionella* species (58). However, the Dot/Icm components of *Legionella* with the greatest variation in protein identity were the same in the Candidatus *P. cheracis* (relative to *C. burnetii*), specifically IcmX, DotA and IcmG/DotF, consistent with exposure of these protein to hosts (58). In contrast, IcmB and DotB had the highest protein identity, which is consistent with previous work in *Legionella* species suggestive that these proteins are under functional selection (58).

Key challenges in the detection of obligate intracellular pathogens is the difficulty in culturing the organisms, and the lack of diagnostic targets that can detect these

bacteria at low concentrations in often complex metagenomic samples. Here we identified that Candidatus *P. cheracis* possesses a high copy transposase that had homology to a symbiont of *Solemya velum*, the Atlantic awning clam. Future efforts to screen for the presence of Candidatus *P. cheracis* could be facilitated by PCR amplification of the transposase in aquatic samples. This would be akin to diagnostic efforts of members of the *Bordetella* genus (59). *Bordetella pertussis,* the causative agent of whooping cough, is commonly identified through the PCR of high copy number insertion sequences (IS) with the primary target IS481. This would provide a potentially non-invasive, sensitive screening strategy for ongoing surveillance of the crayfish farms that would facilitate the prevention and absence of this pathogen in aquaculture environments. Additionally, if proven as a sensitive strategy for detection of Candidatus *P. cheracis*, this PCR approach could also be employed to test different aquatic environments for the presence of this Coxiellaceae genus.

Comparative genomics of 65 *Legionella* species has reported an expansive repertoire of putative Dot/Icm effector proteins (60). While many species possess putative effectors with specific eukaryotic domains, such as GTPase, F-box and SET domains, these motifs are encoded within different proteins that are not orthologous (60). Interestingly, only 8 Dot/Icm substrates are conserved among these 65 species of *Legionella* (60). Using two distinct *in silico* tools, we have developed a list of 238 putative Dot/Icm effectors of Candidatus *P. cheracis.* Unlike *Legionella* species, very few putative Dot/Icm effectors encoded by Candidatus *P. cheracis* possess similarity to eukaryotic domains and a large proportion, ~28%, are novel. We observed 12 predicted effectors with similarity to known or putative *C. burnetii* Dot/Icm effectors, including CpeB and CvpE which have been shown to make important contributions to the intracellular success of *C. burnetii* (48, 54, 56, 57)*.* Further expanding the collection of sequenced Coxiellaceae genomes will facilitate greater comparison of the effector repertoire in this family, allowing us to determine whether some of these 12 common effectors act as "core" Coxiellaceae effectors that facilitate their characteristic interaction with eukaryotic cells. We were able to demonstrate Dot/Icm-dependent translocation of Candidatus *P. cheracis* CpeB (*Pc*CpeB) using *C. burnetii* as a surrogate host. In the absence of methodology for the isolation and cultivation of Candidatus *P. cheracis,* the success of this approach provides an

alternative strategy for developing our understanding of the host-pathogen

interactions mediated by Candidatus *P. cheracis*.

The full genome sequence of Candidatus *P. cheracis* has highlighted potential

screening or diagnostic targets and unveiled likely pathogenesis strategies. In

addition, the genome sequence shows that Candidatus *P. cheracis* encodes for Mip,

a peptidyl-prolyl cis-trans isomerase that has been shown to be essential for

intracellular replication and pathogenesis of *C. burnetii* (61). Candidatus *P. cheracis*

Mip shares 66% identity with *C. burnetii* Mip and these proteins are conserved within

the PPIase catalytic domain which suggests that Mip inhibitors developed by

Debowski et al., may inhibit Candidatus *P. cheracis* Mip (61). Treatment with

rationally designed small molecule inhibitors of Mip may represent a plausible

intervention measure for future outbreaks of Candidatus *P. cheracis* in aquaculture

facilities.

Here, we have further demonstrated the effective use of ONT sequencing to

reconstruct the genome of a novel intracellular pathogen - a member of a family that

is difficult to culture in standard laboratory conditions - directly from its host. The

complete genome of a chromosome and two plasmids enabled the exploration of a

previously unexplored pathogen that is important for Australian aquaculture. ONT

was advantageous here for the capture of the two plasmids, elements that are often

missed by short-read metagenomic binning approaches but can be critical for

virulence (62, 63) and for dealing with the significant number of IS elements

throughout this genome (64). The detection of a T4SS and cohort of novel effector

proteins that likely facilitate intracellular success of the pathogen provides the first

insights into the host pathogen interactions of this novel pathogen. ONT sequencing

has shown promise in recent years for outbreak investigations due to its low-cost,

portability, minimal sample preparation, and rapid, real-time data output (65). If such

an outbreak was to occur today, in addition to traditional histopathology, ONT

metagenomics would enhance efforts to detect and characterize unknown emerging

pathogens.

ACKNOWLEDGEMENTS

DJI is supported by a National Health and Medical Research Council (NHMRC)

Investigator Grant (GNT1195210). CJW and RRW are funded through the NHMRC

(GNT1194325). GRS is the recipient of an Australian Government Research Training

Program Scholarship and Monash Graduate Excellence Scholarship. Research in

HJNs laboratory was funded by the NHMRC (GNT2010841). The funders had no

role in study design, data collection and analysis, decision to publish or preparation

of the manuscript.

REFERENCES

1. Dragičević P, Bielen A, Petrić I, Hudina S. 2021. Microbial pathogens of freshwater crayfish: A critical review and systematization of the existing data with directions for future research. J Fish Dis 44:221-247.
2. Ketterer PJ, Taylor DJ, Prior HC. 1992. Systemic rickettsia-like infection in farmed freshwater crayfish, *Cherax quadricarinatus*, p 173-179. *In* Shariff M, Subasinghe RP, Arthur JR (ed), Diseases in Asian Aquaculture. Asian Fisheries Society.
3. Owens L, Muir P, Sutton D, Wingfield M. 1992. The pathology of microbial diseases in tropical Australian Crustacea, p 165-172. *In* Shariff M, Subasinghe RP, Arthur JR (ed), Diseases in Asian Aquaculture. Asian Fisheries Society.
4. Tan CK, Owens L. 2000. Infectivity, transmission and 16S rRNA sequencing of a rickettsia, *Coxiella cheraxi* sp. nov., from the freshwater crayfish *Cherax quadricarinatus*. Dis Aquat Organ 41:115-22.
5. Cooper A, Layton R, Owens L, Ketheesan N, Govan B. 2007. Evidence for the classification of a crayfish pathogen as a member of the genus *Coxiella*. Lett Appl Microbiol 45:558-63.
6. Ruth Elliman J, Owens L. 2020. Confirmation that candidatus *Coxiella cheraxi* from redclaw crayfish (*Cherax quadricarinatus*) is a close relative of Coxiella burnetii, the agent of Q-fever. Lett Appl Microbiol 71:320-326.
7. Luhrmann A, Newton HJ, Bonazzi M. 2017. Beginning to Understand the Role of the Type IV Secretion System Effector Proteins in *Coxiella burnetii* Pathogenesis. Curr Top Microbiol Immunol 413:243-268.
8. Graells T, Ishak H, Larsson M, Guy L. 2018. The all-intracellular order Legionellales is unexpectedly diverse, globally distributed and lowly abundant. FEMS Microbiol Ecol 94.
9. Duron O, Doublet P, Vavre F, Bouchon D. 2018. The Importance of Revisiting Legionellales Diversity. Trends Parasitol 34:1027-1037.
10. Gardner BR, Bachmann NL, Polkinghorne A, Hufschmid J, Tadepalli M, Marenda M, Graves S, Arnould JPY, Stenos J. 2023. A Novel Marine Mammal *Coxiella burnetii*-Genome Sequencing Identifies a New Genotype with Potential Virulence. Pathogens 12.
11. Carey KL, Newton HJ, Luhrmann A, Roy CR. 2011. The *Coxiella burnetii* Dot/Icm system delivers a unique repertoire of type IV effectors into host cells and is required for intracellular replication. PLoS Pathog 7:e1002056.
12. Beare PA, Gilk SD, Larson CL, Hill J, Stead CM, Omsland A, Cockrell DC, Howe D, Voth DE, Heinzen RA. 2011. Dot/Icm type IVB secretion system requirements for *Coxiella burnetii* growth in human macrophages. MBio 2:e00175-11.
13. Vogel JP, Andrews HL, Wong SK, Isberg RR. 1998. Conjugative transfer by the virulence system of *Legionella pneumophila*. Science 279:873-6.
14. Segal G, Purcell M, Shuman HA. 1998. Host cell killing and bacterial conjugation require overlapping sets of genes within a 22-kb region of the *Legionella pneumophila* genome. Proc Natl Acad Sci U S A 95:1669-74.
15. Qiu J, Luo ZQ. 2017. *Legionella* and *Coxiella* effectors: strength in diversity and activity. Nat Rev Microbiol 15:591-605.
16. Hubber A, Roy CR. 2010. Modulation of host cell function by *Legionella pneumophila* type IV effectors. Annu Rev Cell Dev Biol 26:261-83.

17. Newton HJ, McDonough JA, Roy CR. 2013. Effector protein translocation by the
Coxiella burnetii Dot/Icm type IV secretion system requires endocytic
maturation of the pathogen-occupied vacuole. PLoS One 8:e54566.
18. Newton P, Thomas DR, Reed SCO, Lau N, Xu B, Ong SY, Pasricha S,
Madhamshettiwar PB, Edgington-Mitchell LE, Simpson KJ, Roy CR, Newton HJ.
2020. Lysosomal degradation products induce *Coxiella burnetii* virulence. Proc
Natl Acad Sci U S A 117:6801-6810.
19. Santos-Garcia D, Morel O, Henri H, El Filali A, Buysse M, Noël V, McCoy KD,
Gottlieb Y, Klasson L, Zenner L, Duron O, Vavre F. 2023. Genomic changes
during the evolution of the *Coxiella* genus along the parasitism-mutualism
continuum. Peer Community Journal 3.
20. Davidovich N, Fiocchi E, Basso A, Budai J, Montesi F, Pretto T. 2022. An outbreak
of crayfish rickettsiosis caused by *Coxiella cheraxi* in redclaw crayfish (*Cherax*
quadricarinatus) imported to Israel from Australia. Transbound Emerg Dis
69:204-212.
21. Wood DE, Lu J, Langmead B. 2019. Improved metagenomic analysis with Kraken
2. Genome Biol 20:257.
22. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone
reads using repeat graphs. Nat Biotechnol 37:540-546.
23. Benoit G, Raguideau S, James R, Phillippy AM, Chikhi R, Quince C. 2024. High-
quality metagenome assembly from long accurate reads with metaMDBG. Nat
Biotechnol 42:1378-1383.
24. Li H. 2016. Minimap and miniasm: fast mapping and de novo assembly for noisy
long sequences. Bioinformatics 32:2103-10.
25. Vaser R, Šikić M. 2021. Time- and memory-efficient genome assembly with
Raven. Nat Comput Sci 1:332-336.
26. Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences.
Bioinformatics 34:3094-3100.
27. Wick RR, Judd LM, Cerdeira LT, Hawkey J, Méric G, Vezina B, Wyres KL, Holt KE.
2021. Trycycler: consensus long-read assemblies for bacterial genomes.
Genome Biol 22:266.
28. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu:
scalable and accurate long-read assembly via adaptive k-mer weighting and
repeat separation. Genome Res 27:722-736.
29. Chen Y, Nie F, Xie SQ, Zheng YF, Dai Q, Bray T, Wang YX, Xing JF, Huang ZJ, Wang
DP, He LJ, Luo F, Wang JX, Liu YZ, Xiao CL. 2021. Efficient assembly of nanopore
reads via highly accurate and intact error correction. Nat Commun 12:60.
30. Hu J, Wang Z, Sun Z, Hu B, Ayoola AO, Liang F, Li J, Sandoval JR, Cooper DN, Ye
K, Ruan J, Xiao CL, Wang D, Wu DD, Wang S. 2024. NextDenovo: an efficient
error correction and accurate assembly tool for noisy long reads. Genome Biol
25:107.
31. Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J, Goesmann A. 2021.
Bakta: rapid and standardized annotation of bacterial genomes via alignment-
free sequence identification. Microb Genom 7.
32. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2019. GTDB-Tk: a toolkit to
classify genomes with the Genome Taxonomy Database. Bioinformatics
36:1925-7.

33.   Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2022. GTDB-Tk v2: memory
friendly classification with the genome taxonomy database. Bioinformatics
38:5315-5316.

34.   Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil PA, Hugenholtz P. 2022.
GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based
taxonomy. Nucleic Acids Res 50:D785-d794.

35.   Sherry NL, Horan KA, Ballard SA, Gonçalves da Silva A, Gorrie CL, Schultz MB,
Stevens K, Valcanis M, Sait ML, Stinear TP, Howden BP, Seemann T. 2023. An
ISO-certified genomics workflow for identification and surveillance of
antimicrobial resistance. Nat Commun 14:60.

36.   Robertson J, Nash JHE. 2018. MOB-suite: software tools for clustering,
reconstruction and typing of plasmids from draft assemblies. Microb Genom 4.

37.   Robertson J, Bessonov K, Schonfeld J, Nash JHE. 2020. Universal whole-
sequence-based plasmid typing and its utility to prediction of host range and
epidemiological surveillance. Microb Genom 6.

38.   Payne LJ, Todeschini TC, Wu Y, Perry BJ, Ronson CW, Fineran PC, Nobrega FL,
Jackson SA. 2021. Identification and classification of antiviral defence systems
in bacteria and archaea with PADLOC reveals new system types. Nucleic Acids
Res 49:10868-10878.

39.   Wang J, Yang B, An Y, Marquez-Lago T, Leier A, Wilksch J, Hong Q, Zhang Y,
Hayashida M, Akutsu T, Webb GI, Strugnell RA, Song J, Lithgow T. 2019.
Systematic analysis and prediction of type IV secreted effector proteins by
machine learning approaches. Brief Bioinform 20:931-951.

40.   Hu Y, Wang Y, Hu X, Chao H, Li S, Ni Q, Zhu Y, Hu Y, Zhao Z, Chen M. 2024.
T4SEpp: A pipeline integrating protein language models to predict bacterial type
IV secreted effectors. Comput Struct Biotechnol J 23:801-812.

41.   Bi D, Liu L, Tai C, Deng Z, Rajakumar K, Ou HY. 2013. SecReT4: a web-based
bacterial type IV secretion system resource. Nucleic Acids Res 41:D660-5.

42.   Cazalet C, Gomez-Valero L, Rusniok C, Lomma M, Dervins-Ravault D, Newton
HJ, Sansom FM, Jarraud S, Zidane N, Ma L, Bouchier C, Etienne J, Hartland EL,
Buchrieser C. 2010. Analysis of the *Legionella longbeachae* genome and
transcriptome uncovers unique strategies to cause Legionnaires' disease. PLoS
Genet 6:e1000851.

43.   Brenner AE, Muñoz-Leal S, Sachan M, Labruna MB, Raghavan R. 2021. *Coxiella*
burnetii and Related Tick Endosymbionts Evolved from Pathogenic Ancestors.
Genome Biol Evol 13.

44.   Paradis E, Claude J, Strimmer K. 2004. APE: Analyses of Phylogenetics and
Evolution in R language. Bioinformatics 20:289-90.

45.   Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. GGTREE: an R package for
visualization and annotation of phylogenetic trees with their covariates and
other associated data. Methods in Ecology and Evolution 8:28-36.

46.   Omsland A, Beare PA, Hill J, Cockrell DC, Howe D, Hansen B, Samuel JE,
Heinzen RA. 2011. Isolation from animal tissue and genetic transformation of
Coxiella burnetii are facilitated by an improved axenic growth medium. Appl
Environ Microbiol 77:3720-5.

47. Voth DE, Beare PA, Howe D, Sharma UM, Samoilis G, Cockrell DC, Omsland A,
Heinzen RA. 2011. The *Coxiella burnetii* cryptic plasmid is enriched in genes
encoding type IV secretion system substrates. J Bacteriol 193:1493-503.

48. Newton HJ, Kohler LJ, McDonough JA, Temoche-Diaz M, Crabill E, Hartland EL,
Roy CR. 2014. A screen of *Coxiella burnetii* mutants reveals important roles for
Dot/Icm effectors and host autophagy in vacuole biogenesis. PLoS Pathog
10:e1004286.

49. Thomas DR, Garnish SE, Khoo CA, Padmanabhan B, Scott NE, Newton HJ. 2024.
Coxiella burnetii protein CBU2016 supports CCV expansion. Pathog Dis 82.

50. Abou Abdallah R, Million M, Delerce J, Anani H, Diop A, Caputo A, Zgheib R,
Rousset E, Sidi Boumedine K, Raoult D, Fournier PE. 2022. Pangenomic analysis
of *Coxiella burnetii* unveils new traits in genome architecture. Front Microbiol
13:1022356.

51. Oren A, Garrity GM. 2022. CANDIDATUS LIST No. 3. Lists of names of prokaryotic
Candidatus taxa. Int J Syst Evol Microbiol 72.

52. Dmytrenko O, Russell SL, Loo WT, Fontanez KM, Liao L, Roeselers G, Sharma R,
Stewart FJ, Newton IL, Woyke T, Wu D, Lang JM, Eisen JA, Cavanaugh CM. 2014.
The genome of the intracellular bacterium of the coastal bivalve, *Solemya*
velum: a blueprint for thriving in and out of symbiosis. BMC Genomics 15:924.

53. Weber MM, Chen C, Rowin K, Mertens K, Galvan G, Zhi H, Dealing CM, Roman
VA, Banga S, Tan Y, Luo ZQ, Samuel JE. 2013. Identification of *Coxiella burnetii*
type IV secretion substrates required for intracellular replication and *Coxiella*-
containing vacuole formation. J Bacteriol 195:3914-24.

54. Martinez E, Allombert J, Cantet F, Lakhani A, Yandrapalli N, Neyret A, Norville IH,
Favard C, Muriaux D, Bonazzi M. 2016. *Coxiella burnetii* effector CvpB modulates
phosphoinositide metabolism for optimal vacuole development. Proc Natl Acad
Sci U S A 113:E3260-9.

55. Kohler LJ, Reed Sh C, Sarraf SA, Arteaga DD, Newton HJ, Roy CR. 2016. Effector
Protein Cig2 Decreases Host Tolerance of Infection by Directing Constitutive
Fusion of Autophagosomes with the *Coxiella*-Containing Vacuole. MBio 7.

56. Larson CL, Beare PA, Voth DE, Howe D, Cockrell DC, Bastidas RJ, Valdivia RH,
Heinzen RA. 2015. *Coxiella burnetii* effector proteins that localize to the
parasitophorous vacuole membrane promote intracellular replication. Infect
Immun 83:661-70.

57. Fu M, Zhang J, Zhao M, Zhang S, Dai L, Ouyang X, Yu Y, Wen B, Zhou D, Sun Y,
Jiao J, Xiong X. 2022. *Coxiella burnetii* Plasmid Effector B Promotes LC3-II
Accumulation and Contributes To Bacterial Virulence in a SCID Mouse Model.
Infect Immun 90:e0001622.

58. Gomez-Valero L, Chiner-Oms A, Comas I, Buchrieser C. 2019. Evolutionary
Dissection of the Dot/Icm System Based on Comparative Genomics of 58
Legionella Species. Genome Biol Evol 11:2619-2632.

59. Martini H, Detemmerman L, Soetens O, Yusuf E, Piérard D. 2017. Improving
specificity of *Bordetella pertussis* detection using a four target real-time PCR.
PLoS One 12:e0175587.

60. Gomez-Valero L, Rusniok C, Carson D, Mondino S, Pérez-Cobas AE, Rolando M,
Pasricha S, Reuter S, Demirtas J, Crumbach J, Descorps-Declere S, Hartland EL,
Jarraud S, Dougan G, Schroeder GN, Frankel G, Buchrieser C. 2019. More than

18,000 effectors in the *Legionella* genus genome provide multiple, independent
combinations for replication in human cells. Proc Natl Acad Sci U S A 116:2265-
2273.
61.   Debowski AW, Bzdyl NM, Thomas DR, Scott NE, Jenkins CH, Iwasaki J, Kibble EA,
Khoo CA, Scheuplein NJ, Seibel PM, Lohr T, Metters G, Bond CS, Norville IH,
Stubbs KA, Harmer NJ, Holzgrabe U, Newton HJ, Sarkar-Tyson M. 2023.
Macrophage infectivity potentiator protein, a peptidyl prolyl cis-trans isomerase,
essential for *Coxiella burnetii* growth and pathogenesis. PLoS Pathog
19:e1011491.
62.   Luo S, Lu S, Fan H, Sun Z, Hu Y, Li R, An X, Uversky VN, Chen Z, Tong Y, Song L.
2021. The *Coxiella burnetii* QpH1 plasmid is a virulence factor for colonizing
bone marrow-derived murine macrophages. J Bacteriol 203.
63.   Wachter S, Cockrell DC, Miller HE, Virtaneva K, Kanakabandi K, Darwitz B,
Heinzen RA, Beare PA. 2022. The endogenous *Coxiella burnetii* plasmid encodes
a functional toxin-antitoxin system. Mol Microbiol 118:744-764.
64.   Maguire F, Jia B, Gray KL, Lau WYV, Beiko RG, Brinkman FSL. 2020.
Metagenome-assembled genome binning methods with short reads
disproportionately fail for plasmids and genomic Islands. Microb Genom 6.
65.   Buytaers FE, Saltykova A, Denayer S, Verhaegen B, Vanneste K, Roosens NHC,
Piérard D, Marchal K, De Keersmaecker SCJ. 2021. Towards Real-Time and
Affordable Strain-Level Metagenomics-Based Foodborne Outbreak
Investigations Using Oxford Nanopore Sequencing Technologies. Front Microbiol
12:738284.
66.   Pan X, Lührmann A, Satoh A, Laskowski-Arce MA, Roy CR. 2008. Ankyrin repeat
proteins comprise a diverse family of bacterial type IV effectors. Science
320:1651-4.
67.   Voth DE, Howe D, Beare PA, Vogel JP, Unsworth N, Samuel JE, Heinzen RA. 2009.
The *Coxiella burnetii* ankyrin repeat domain-containing protein family is
heterogeneous, with C-terminal truncations that influence Dot/Icm-mediated
secretion. J Bacteriol 191:4232-42.
68.   Larson CL, Pullman W, Beare PA, Heinzen RA. 2023. Identification of Type 4B
Secretion System Substrates That Are Conserved among *Coxiella burnetii*
Genomes and Promote Intracellular Growth. Microbiol Spectr 11:e0069623.
69.   Burette M, Allombert J, Lambou K, Maarifi G, Nisole S, Di Russo Case E, Blanchet
FP, Hassen-Khodja C, Cabantous S, Samuel J, Martinez E, Bonazzi M. 2020.
Modulation of innate immune signaling by a *Coxiella burnetii* eukaryotic-like
effector protein. Proc Natl Acad Sci U S A 117:13708-13718.
70.   Chen C, Banga S, Mertens K, Weber MM, Gorbaslieva I, Tan Y, Luo ZQ, Samuel
JE. 2010. Large-scale identification and translocation of type IV secretion
substrates by *Coxiella burnetii*. Proc Natl Acad Sci U S A 107:21755-60.
71.   Crabill E, Schofield WB, Newton HJ, Goodman AL, Roy CR. 2018. Dot/Icm-
translocated proteins important for biogenesis of the *Coxiella burnetii-*
containing vacuole identified by screening of an effector mutant sub-library.
Infect Immun doi:10.1128/iai.00758-17.
72.   Fielden LF, Scott NE, Palmer CS, Khoo CA, Newton HJ, Stojanovski D. 2020.
Proteomic identification of *Coxiella burnetii* effector proteins targeted to the

host cell mitochondria during infection. Mol Cell Proteomics
doi:10.1074/mcp.RA120.002370.

73. Zhang Y, Fu J, Liu S, Wang L, Qiu J, van Schaik EJ, Samuel JE, Song L, Luo ZQ. 2022. *Coxiella burnetii* inhibits host immunity by a protein phosphatase adapted from glycolysis. Proc Natl Acad Sci U S A 119.

74. Martinez E, Huc-Brandt S, Brelle S, Allombert J, Cantet F, Gannoun-Zaki L, Burette M, Martin M, Letourneur F, Bonazzi M, Molle V. 2020. The secreted protein kinase CstK from *Coxiella burnetii* influences vacuole development and interacts with the GTPase-activating host protein TBC1D5. J Biol Chem 295:7391-7403.

Table 1. Candidatus *P. cheracis* proteins with similarity to *C. burnetii* Dot/Icm
substrates

Candidatus P. *cheracis* gene	*C. burnetii* gene	Identity (%)	Coverage (%)	Role of *C. burnetii* effector
PBJCNB_00280	CBUD_0487	26	22	Experimentally shown to be a Dot/Icm substrate (56).
PBJCNB_01005	CBU_1863 (CvpE)	40	78	*Coxiella* vacuolar protein E localizes to the *Coxiella*-containing vacuole (CCV). Characterization of a deletion mutant demonstrated that CvpE contributes to intracellular replication of *C. burnetii* (47, 56).
PBJCNB_01155	CBU_0201 (AnkC)	84	95	Early studies using *L. pneumophila* as a surrogate system did not demonstrate translocation of AnkC (53, 66, 67). Similarly, no translocation was observed in *C. burnetii* using the CyaA reporter assay (68). However, using a BlaM reporter assay in *C. burnetii* low level of translocation were observed at a later time point, 72 h post-infection (69).
PBJCNB_01790	CBUA_0013 (CpeB)	29	90	Plasmid encoded *C. burnetii* effector that contributes to LC3-II accumulation in a Rab11-dependent manner (57).
PBJCNB_05220	CBU_0937 (CirC/MceB)	57	100	Identified as a DotF binding protein and demonstrated to be translocated by *L. pneumophila* (70). Two independent *cbu0937* transposon mutants

				were reported to have an intracellular replication defect (53, 71). Detected in mitochondria isolated from *C. burnetii* infected cells (72). However, recent translocation reporter assays conducted in *C. burnetii* could not confirm translocation of this protein (68).
PBJCNB_05510	CBU_1079	76	97	5% translocation observed using the BlaM reporter in *L. pneumophila* (53). Not validated in *C. burnetii* using the CyaA reporter (68).
PBJCNB_08410	CBU_0513 (CinF)	83	99	Shown to be translocated as a BlaM-fusion protein by *L. pneumophila* with Dot/Icm-dependent translocation confirmed in *C. burnetii* (53, 73).Loss of CinF, either through transposon disruption of silencing expression, inhibits intracellular replication of *C. burnetii* (53, 73). CinF is a protein phosphatase that dephosphorylates IκBα (73).
PBJCNB_09120	CBU_1425 (MceC)	74	100	Identified as a DotF binding protein and demonstrated to be translocated by *L. pneumophila* (70). Identified as being enriched in the mitochondria of *C. burnetii* infected cells with the capacity to

				localise to the inner mitochondrial membrane and interact with components of the mitochondrial quality control machinery (72). Translocation by C. burnetii not confirmed using the BlaM reporter assay (68).
PBJCNB_09430	CBU_0469	61	95	Low level (1%) translocation of BlaM fusion protein in L. pneumophila (53). Translocation has not been confirmed in *C. burnetii*.
PBJCNB_10235	CBU_1566	79	99	BlaM fusion protein was shown to be translocated by *L. pneumophila* (53) but no translocation was observed for a Cya fusion in *C. burnetii* (68).
PBJCNB_10940	CBU_0175 (CoxK1)	82	100	BlaM fusion protein translocated by *C. burnetii* in a Dot/Icm-dependent manner (74). Shown to interact with the Rab7 GTPase-activating protein TBC1D5 (74).
PBJCNB_10315	CBU_1594 (MceD)	71	97	BlaM fusion protein was shown to be translocated by *L. pneumophila* (53) but no translocation was observed for a Cya fusion in *C. burnetii* (68). Identified as being enriched in the mitochondria of *C. burnetii* infected cells (72).

 FIGURE LEGENDS

Figure 1. Phylogeny of Coxiellales Order demonstrating relationship to

Candidatus *P. cheracis* genome

The inferred evolution of the Candidatus *P. cheracis* genome in representative

isolates from Coxiellales Order (gtdb.ecogenomic.org/tree?r=o__Coxiellales).

The phylogeny was inferred using GTDB-Tk, a marker gene-based method of 120

domain-specific marker genes for bacteria (33). The tips of the tree are coloured by

the NCBI organism for the Coxiellales lineages, and additional GTDB lineages at the

genus level. The associated hosts or source from where the samples were collected

are shown to the right of the tree. The Candidatus *P. cheracis* genome is indicated in

green and by the crayfish icon (sourced from www.phylopic.org/).

Figure 2. Characterization of the Candidatus *P. cheracis* genome

The complete genome showing A) the chromosome and B) two plasmids. The

middle ring shows the coding sequence (CDS) on the chromosome and plasmids.

Genes of interest are shown in the outer ring with colours indicating broad function

(T4SS, multicopy transposase (TS), phage defense and putative effector). The total

number of these four functional classes on each genetic element are shown on panel

C.

Figure 3. Structure and protein identity of the T4SS in the Candidatus *P.*

***cheracis* genome**

A) Characterization of the T4SS identified in Candidatus *P. cheracis, Coxiella*

burnetii, *Legionella pneumophila, Legionella longbeachae* and Candidatus

Rickettsiella viridis. Genes encoding the T4SS are colored by function within the

system. B) Visualization of Dot/Icm protein identity relative to *Coxiella burnetii* RSA

493.

Figure 4. Candidatus *P. cheracis Pc*CpeB is a Dot/Icm substrate.

HeLa cells were infected for 24 h with either wildtype (NMII) or Dot/Icm-deficient

(*icmL*::Tn) *C. burnetii* expressing the indicated BlaM fusion proteins. **(A)** Change in

the fluorescence emission ratio 450nm::520nm was measured as a readout of BlaM

cleaving the CCF2-AM substrate (y-axis). The graph depicts this ratio from 4

independent experiments (circles) with the bar representing the mean and SEM

ratio. The threshold for translocation positive was set to 2.5× the BlaM alone

negative control (dotted line) and Šidák's multiple comparisons test was used to

determine P-values compared to the BlaM and between NMII and *icmL*::Tn strains

expressing the same fusion proteins. **(B)** Representative fluorescent images of

infected HeLa cells demonstrate the translocation of CvpB (*C. burnetii* effector

positive control) and *Pc*CpeB by *C. burnetii* NMII but not *C. burnetii icmL*::Tn.

Fluorescence intensity at 520 nm of uncleaved CCF2-AM is shown in green and

cleaved CCF2-AM, at 450 nm is blue, scale bars represent 100 μm. **(C)** Immunoblot

analysis of the indicated *C. burnetii* strains confirmed expression of BlaM (29 kDa),

BlaM-CvpB (117.6 kDa) and BlaM-*Pc*CpeB (54.5 kDa). **(D)** ClustalOmega alignment

showing the similarity between *C. burnetii* CpeB (*Cb*CpeB) and Candidatus *P.*

cheracis CpeB (*Pc*CpeB). The two proteins share 26% identity (*) with additional

conserved (:) and semi-conserved (.) substitutions. The purple boxed area

represents amino acids 164 to 204 that were previously shown to mediate interaction

of *Cb*CpeB with Rab11a (57).

SUPPLEMENTARY FILES

Supplementary File 1. Workflow for quality controls, kraken assignment and

genome assembly.

Supplementary File 2. The Kraken2 classification of the chromosomal reads using

a GTDB-based database.

Supplementary File 3. Predicted Dot/Icm effector proteins based on Bastion4 and

T4Sepp tools.

marine water and coral reef sponge

activated sludge

ground water, soil and subsurface

estauarine and marine water

lake water
drinking water
subsurface
crayfish
coral reef sponge

tick

tick

human, goat, dog, mouse, tick

NCBI Organism

- Candidatus *Paracoxiella cheracis*
- Candidatus *Coxiella mudrowiae*
- *Coxiella burnetii*
- *Coxiella* species
- *Coxiella* endosymbiont
- *Coxiella*–like endosymbiont
- Coxiellaceae bacterium
- uncultured *Coxiella* species
- uncultured Coxiellaceae bacterium
- gtdb JAFEDN01 genus
- gtdb JAHFRQ01, 2-02-FULL-42-43, JAHZIV01, JAHZKQ01, JAJFJO01 genus
- gtdb CAIQCE01 and CACHNP01 genus

A

Phage defense
Phage defense
(SoFic)
Phage defense
Phage defense
Phage defense
Phage defense
Phage defense (SoFic)
Phage defense
Phage defense

Paracoxiella cheracis
chromosome

2,233,862 bp

Phage defense
(AbiEi,AbiEii)
Phage defense
Dot/Icm
T4SS

Phage defense

Phage defense

Putative function

- Effector
- Phage defense
- Type IV Secretion System (T4SS)
- Transposase (TS)

B

Paracoxiella cheracis
plasmid 1

59,783 bp

Paracoxiella cheracis
plasmid 2

34,021 bp

C

Chromosome | Plasmid 1

Number detected

Plasmid 2

Putative function

Effector | Phage | T4SS | TS

A

CCF2-AM 450::520 ratio

P=0.0037
P<0.0001
P<0.0001
P=0.0046

Uninfected
BlaM (NMII)
CvpB (NMII)
CvpB (*icmL*::Tn)
*Pc*CpeB (NMII)
Pc CpeB (*icmL*::Tn)

B

NMII   *icmL*::Tn

BlaM-CvpB

BlaM-*Pc*CpeB

C

117.6 kDa
54.5 kDa
29 kDa

D

CbCpeB MPYFFTLPKIDIKE--------KSASQTPEKVLREWLRGASNQEDASLFIGNSPIF----
PcCpeB MSAFFTVPKPQYGGESELLEREKKDDTSGKITLKDWVRNASNACEARVFVMSGTSPNTLT 60
        * ***:** :   *. . : : .*::*:*.*** :* :*: ..

        ---------GEHQDTSGSSLPKVEIPEFINSIKAVEAVFKKEGLKDEEILNYILHFGGIN 99
        TAINESDLNAYMRDDDGE-----RIPLLSDADQYIKQLLEKHGLENHRIYKYIRMFANIN 115
          . :* .*.   .** : :: : :: :::*.**::..* :** *..**

        GYMTLGQVVFEEISMNKG-KSSRMENRRYSYRVKNKGEVEYIEQFDFYKKNTCDRNEIGD 158
        GFLPAGQIVMSQVKNSGEQSPIHTENPKYTFFVKDKTTVQYTEESDIVLNPTH-EYGKGE 174
        *:: **:*:.:: .   .   : ** *:: **:* *:* *: *: : * .   *:

        FIGQV⎡KIVSIISKDGD-IQHACKKVEI-AVPGEKLNSFRKHFGDQLNF⎤VNRMKLRLVELY 216
        TIAKL⎣KVVSIISSNDQKIHHILDSASLTAYPGK-EPQCIKCFGDNRSV⎦MVKLIHKLKEMF 233
         .:::*****:.:: *:* ....: * **: . * ***: .: :: :* *::

        NSLVERFRAAINKADRTN-----NNCFPFCSR 243
        MAFVALFTKRSSCEQDYNGFNDVASNAASCRR 265
         ::* *   . : *   .   * *