Sex-specific thermoregulatory effects of estrogen signaling in *Reprimo* lineage cells

Authors:
Jae W. Park[1*], Laura R. Cortes[1*#], Norma P. Sandoval[1], Adriana R. Vree[1], Alejandra G. Baron[1], Kelly Vranich, Higor J. Fideles[1], Rosalizbeth M. Martinez, Elizabeth A. Dilday[1], Mia R. Hansen[1], Weronika Budek, Julissa I. Lopez[1,2], Laura G. Kammel[1], J. Edward van Veen[1], Stephanie M. Correa[1#]

Affiliation:
[1]Department of Integrative Biology and Physiology, University of California, Los Angeles, California, USA
[2] Cypress College, Cypress, CA, USA

*co-first authors
#authors for correspondence: lauracortes@ucla.edu and stephaniecorrea@ucla.edu

The authors have nothing to disclose.

Corresponding authors ORCiD IDs:
 Stephanie M Correa: https://orcid.org/0000-0002-6221-689X
 Laura R Cortes https://orcid.org/0000-0002-8202-7949

Abstract

Estrogens have considerable impact on energy homeostasis and metabolic health. In mice, signaling through estrogen receptor alpha (ERα) alters energy intake and multiple aspects of energy expenditure, effects that may be mediated by specific regions or neuronal sub-populations of the hypothalamus. This study investigates the function of ERα in neurons of the lineage that expresses *Rprm* (Reprimo), a gene we previously linked to thermoregulation in females. Here, we engineered a novel *Reprimo^{Cre}* mouse to selectively knock out ERα in *Rprm* lineage cells (Reprimo-specific estrogen receptor α KO; RERKO) and report changes in core temperature in female mice, with no changes in movement or food intake. RERKO females have elevated brown adipose tissue (BAT) temperature and lower tail temperature relative to controls, suggesting increased heat production and reduced heat dissipation, respectively. Developmental expression of *Rprm* was detected in the brain, but not in BAT or white adipose tissue suggesting temperature changes may be mediated by the nervous system. To confirm centrally mediated effects on temperature, we ablated *Rprm* expressing cells in the mediobasal hypothalamus and observed a reduction in core temperature relative to controls. Taken together, these results indicate that estrogen signaling in the *Rprm* lineage is critical for thermoregulation, mainly through the modulation of brown adipose tissue thermogenesis in female, but not male, mice.

Keywords: Estrogen receptor alpha, thermogenesis**,** energy expenditure, *Reprimo*, metabolism, hypothalamus

Introduction

Estrogens are potent modulators of energy homeostasis [1–3], and thus, plummeting levels of estrogen during menopause dysregulate metabolism [4]. Menopausal people experience hot flushes and reduced energy expenditure, which leads to greater adiposity, body weight, and risk of metabolic disorders [5–8]. Energy expenditure is regulated by estrogen signaling in the hypothalamus [9–13]. Recent work suggests that distinct groups of hypothalamic estrogen receptor alpha expressing (ERα+) neurons may regulate specific aspects of energy homeostasis [10,11,13]. Pinpointing these neural subsets and their precise functions can lead to a mechanistic understanding of how energy expenditure is regulated and inform the design of cell-based therapeutics tailored to specific symptoms in menopausal or obese patients.

Rodents are suitable models to study the hormonal regulation of energy expenditure. Like menopausal people, mice lacking ovaries have lower energy expenditure and increased body weight and adiposity [14], and estradiol administration blunts these changes [15]. Although ERα (*Esr1*) is found throughout the body, its expression in the central nervous system (CNS) is necessary for energy homeostasis. Mice with *Esr1* knocked out (KO) in the *Nestin-Cre* cell lineage that labels the nervous system have greater body mass, and females, in particular, eat more and show reduced energy expenditure relative to controls [9]. The neural circuitry involved in energy homeostasis includes the ventrolateral area of the ventromedial nucleus (VMHvl), the medial preoptic area (MPO), and the arcuate nucleus (ARH). These regions contain abundant expression of ERα. Female mice and rats with *Esr1* knockdown in the VMHvl have lower activity levels, basal metabolic rate, and energy expenditure, resulting in higher body weight relative to controls [16,17]. Similarly, mice with *Esr1* KO in the steroidogenic factor 1 (*Sf1/Nr5a1*)-lineage have lower metabolic rates and thermogenesis [9]. In contrast, activating *Esr1+* neurons in the VMHvl (VMHvl^{Esr1}) increases heat production and movement [11]. Thus, the VMHvl is essential for the regulation of energy expenditure. The effects of estrogens on food intake are regulated by other regions, including the nearby ARH. Knocking out *Esr1* in pro-opiomelanocortin (POMC) lineage cells increases food intake and body weight in females relative to controls [9]. Similarly, silencing steroid and metabolic hormone-responsive ERα/Kisspeptin+ neurons in the ARH causes obesity in female mice and alters circadian feeding patterns [18–21]. Lastly, activation of ERα+ cells in the MPO leads to a hypometabolic state akin to torpor; temperature, energy expenditure, metabolic rate, and movement plummet [12]. In

summary, ERα+ cells or expression in the VMHvl and ARH promote energy expenditure, while

Esr1 expression in ARH cells reduce energy intake, and ERα+ MPO cells trigger a profound

energy conservation state.

Intermingled within the same nuclei, ERα+ cells may differ in their function,

connections, and neurochemical identity (*i.e.*, their gene expression pattern) [22,23,11,24]. For

example, a study using preoptic area slices from rats showed that a portion of estrogen-

responsive MPO neurons (likely expressing ERα) are temperature-responsive, with some

responding to warmth and others to cold [25]. In a similar fashion, VMHvl^{Esr1} cells differ in their

response to glucose and their neural connectivity [26,24]. About half are excited by an increase

in glucose levels and project to the dorsal raphe nucleus, whereas the other half are excited by a

depletion of glucose levels and project to the arcuate nucleus [24]. Functional specialization

among ERα+ cells suggests that they may differ in the expression of peptides and proteins that

shape their roles and responses. Indeed, our lab found that ERα+ VMHvl cells are

neurochemically heterogenous. Using single-cell sequencing, we found sub-populations of ERα+

VMHvl neurons marked by the expression of substance P (Tachykinin 1; *Tac1*), reprimo (*Rprm*),

or prodynorphin (*Pdyn*). Intriguingly, females have many more ERα neurons that express *Tac1*

and *Rprm* than males, and manipulating these cells differentially alters energy expenditure. Mice

that are missing a subset of ERα+ VMHvl cells that co-express *Tac1* and *Nkx2-1* or have

inactivation of the melanocortin receptor 4 (*Mc4r*) subpopulation move less [10,13]. With

Tac1/Nkx2-1 cell loss, this also leads to increased adiposity and body weight [10]. Interestingly,

Rprm knockdown in the VMHvl leaves movement intact, but increases core temperature and

thermogenesis. These findings suggest specialization of cell populations or gene function within

estrogen sensitive hypothalamic regions.

The function of *Rprm*, a tumor suppressor gene, and *Rprm*+ neurons has been largely

unexplored outside of pathological contexts [27–29], with the exception of one developmental

gene expression study [30]. In the present work, we test the function of *Rprm*+ cells on energy

expenditure. Specifically, we hypothesize that *Esr1* expression in *Rprm*+ cells modulates body

temperature via thermogenic brown adipose tissue (BAT) [31]. We generated a knock-in mouse

that expresses Cre recombinase under the control of *Rprm* to selectively knock out *Esr1* in *Rprm*

lineage cells. Despite the extensive effects of global and CNS-specific *Esr1* knockout on weight,

food intake and movement [9,32,33], *Esr1* KO in *Rprm* lineage cells produces a more restricted

phenotype. Core temperature is altered between controls and KO, while body weight, body

composition, and food intake are largely unaffected. Furthermore, BAT mass is heavier and

produces more heat, indicating increased thermogenesis. Because *Esr1* and *Rprm* are co-

expressed in several brain areas and peripheral organs, we next determined the specific role of

Rprm cells in the mediobasal hypothalamus (MBH) using a Cre-dependent caspase. Loss of

Rprm cells led to lower core temperature and higher BAT mass compared with controls. In

summary, our data suggest that *Esr1* expression in *Rprm* lineage cells and MBH^{Rprm} neurons

seem to largely modulate core temperature, while leaving other aspects of energy balance intact.

Methods

Generation and validation of a Rprm^{Cre} knock-in mouse

The donor construct consisted of a double stranded DNA repair cassette that included the coding

sequence for *Rprm* linked to the codon-improved cre recombinase (*iCre*) sequence via a P2A

peptide (Fig. 1A). The P2A peptide leads to ribosomal skipping and the generation of distinct

peptide products, in roughly equal amounts, from a single multi-cistronic construct [34]. A 21-bp

nuclear localization sequence (NLS) is located at the 5' end of the *iCre* exon and helps direct

proteins into the nucleus. To induce proper DNA insertion, we also included homology arms

(approximately 2,000 bp) both upstream and downstream of the *Rprm* and *iCre* coding

sequences. Finally, 34-bp long flippase recognition target (FRT) sites were inserted in the

upstream and downstream homology arms within 250 bp of the 5' and 3' exons. The total length

of the construct is approximately 8.3 Kb. The engineered DNA sequences were based on the

C57BL/6J mm8 reference genome and introduced into albino C57BL/6 blastocysts by CRISPR

injection by the University of California San Diego Transgenic & CRISPR Mouse Core. Two

founder lines were generated but only one was confirmed to contain the entire donor sequence by

PCR validation and sequencing. The founder was crossed to C57BL/6J females and

heterozygous offspring were used to generate a homozygous line using in-crosses.

Esr1$^{f/f}$ female mice [35] were paired with *Rprm*^{Cre};*Esr1*$^{f/f}$ studs to generate **R**eprimo-

specific **E**strogen **R**eceptor α **K**nock **O**ut (RERKO) mice on a mixed B6;129P background. In

this RERKO line, mice had *Esr1* knocked out (KO) in *Rprm* lineage cells. Mice were bred and

maintained in approved conditions at the University of California Los Angeles. Control

littermates were *Esr1*$^{f/f}$ and denoted WT for the *Rprm*^{Cre} mutation. For lineage experiments,

Rprm^{Cre} mice on a C57BL/6J background were paired with *Ai14*$^{f/f}$ reporter mice (Jax #007914;

[36]) to generate offspring that fluoresce tdTomato in tissues expressing *Rprm* (*Rprm^Cre^*; *Ai14^f/+^*).

Mice were kept in 12:12 light cycle conditions (zeitgeber time (ZT) 0 = 7 am) at ~23 °C and

~30% humidity with food and water provided *ad libitum*. All animal studies were approved by

the UCLA Institutional Animal Care and Use Committee (IACUC) and were carried out

following the recommendations of the Guide for the Care and Use of Laboratory Animals of the

National Institutes of Health.

Surgeries

For all surgeries, mice were anesthetized with isoflurane and administered buprenorphine (0.01

mg/mL) and carprofen (0.58 mg/mL) on the day of surgery and the following day to manage pain

and inflammation.

Probe implants

G2 eMitters (Starr Life Sciences; Oakmont, PA, USA) were placed into the abdominal cavity to

record core temperature and locomotion. Cubisens TS100/TS110 probes (Cubeworks; Ann

Arbor, MI, USA) or Nano-T loggers (Star-Oddi, Garðabær, Iceland) were secured

subcutaneously above the BAT depots or over the base of the tail to gauge heat generation and

dissipation, respectively. Custom sleeves were designed to secure probes above BAT and tail

using Autodesk Fusion 360 (San Francisco, CA, USA) and UltiMakerCura software (New York,

NY, USA). Sleeves were printed on a Creality Ender-2 V2 3D printer using PLA filament. Files

are publicly available on NIH 3D (Reference model: 3DPX-020322;

https://3d.nih.gov/entries/3DPX-020322).

Stereotaxic surgery

Control (*pAAV8-Syn-FLEX-Mac-GFP*) or caspase virus (*pAAV2-flex-taCasp3-TEVp*) was

delivered into the VMHvl (AP 1.58, ML ± 0.65, DV 5.7) in 8-week-old C57BL/6J wildtype or

*Rprm^Cre^* mice [37,38]. Each hemisphere received 100 nL at 5-10 nL/sec using a 10 µL Hamilton

syringe attached to a pulled glass needle. Mice recovered for two weeks before starting three

days of temperature and locomotion recordings which were averaged hourly. Targeting of the

VMHvl was validated by immunohistochemical labeling of ERα (described below), since there

is no suitable reprimo antibody and brains were not collected RNAse-free. Because it could not

be confirmed that *Rprm+* cells were unaffected in nearby nuclei, such as the ARH, this

manipulation is termed MBH-specific.

 Collection

Mice were perfused with 4% paraformaldehyde (PFA; Electron Microscopy Sciences; Hatfield,

PA, USA) in 0.01M phosphate-buffered saline (1x PBS). Brains were stored in 4% PFA

overnight at 4°C and transferred to 30% sucrose in 1x PBS until brains sank. Brains were

embedded in Optimal Cutting Temperature compound (OCT; Fisher Scientific, Fair Lawn, NJ,

USA) and stored at -80°C until cutting. BAT, inguinal white adipose tissue (iWAT), and gonadal

white adipose tissue (gWAT) were dissected and stored in tissue cassettes submerged in 70%

ethanol until staining.

Metabolic and reproductive phenotyping

Mice were group housed according to sex, and body weight was recorded weekly from 5 to 16

weeks of age. A separate cohort of mice were weighed and placed into an EchoMRI^{tm} (Houston,

TX, USA) to record fat and lean mass at 16 weeks of age. To record food intake, a pre-weighed

allotment was provided to singly housed mice daily for 7 consecutive days. Every 24 hours, the

remaining food and any crumbs in the cage were weighed and subtracted from the amount

initially provided to determine daily food intake, which was averaged over seven days to provide

a daily average of food consumption. To record core temperature and locomotion, mice were

placed on ER4000 energizer/receiver pads in their home cages (version 5, VitalView software,

Starr Life Sciences). All temperature and activity measurements were collected every five

minutes across three to five days. For temperature and activity recordings, an average for each

hour of the day (ZT 0 – 23) across recording days was calculated. Raw data from G2 emitters,

TS100/TS110 probes, and Nano-T loggers were exported in CSV format using VitalView (Starr

Life Science), TS GUI (Cubeworks), and Mercury (Star-Oddi), respectively. Data were prepared

for analysis in Microsoft Excel (Microsoft Corporation; Redmond, Washington, USA). R Studio

(R version 4.3.1; Boston, MA, USA) was used to process the data and generate hourly averages

(code for processing and plotting provided at https://github.com/kodori00/Rscript). To track the

estrous cycle, mice underwent vaginal lavage using 0.9% saline every morning between ZT 1-2.

Histology
Cytology
For estrous cycling, vaginal lavages were deposited onto slides, stained with Giemsa (0.6% in 1x

PBS;) to visualize nucleic acid and distinguish cell morphology, and imaged using light

microscopy as described in Massa et al., 2023. The estrous stage was determined by the relative

amounts of leukocytes and cornified and nucleated epithelial [39]. BAT and iWAT were

sectioned into 4 µm slices and stained with hematoxylin and eosin by the UCLA Translational

Pathology Core Laboratory.

Immunohistochemistry

Brains were coronally cryo-sectioned into four or six 20-30 µm series and stored on slides or in

cryoprotectant (30% sucrose, 30% ethylene glycol, and 1% polyvinylpyrrolidone in 1x PBS) at -

80°C until use. One series of 30 µm thick sections underwent immunohistochemistry to validate

the loss of ERα expression in RERKO mice (rabbit anti-ERα, 1:1000, ThermoFisher Scientific,

Waltham, MA; Cat# PA1-309, RRID:AB_325814) and one series of 20 µm sections was used to

validate the loss of ERα cells in caspase-treated mice (rabbit anti-ERα, 1:500, Millipore Sigma,

Burlington, MA; Cat# 06-935, RRID:AB_310305). Tissue was rinsed in 1x PBS and blocked for

one hour at room temperature (RT) in either 10% bovine serum albumin (BSA) and 2% Normal

Goat Serum (NGS) or 10% NGS in 0.3% PBS-Triton X depending on antibody. After blocking,

tissue was incubated in primary antibody overnight at 4°C or RT. The following day, the tissue

was incubated with either Alexa Fluor 546 (1:500; Thermo Fisher Scientific Cat# A-11035,

RRID:AB_2534093) or 647 (1:1000; Jackson ImmunoResearch Labs, West Grove, PA; Cat#

111-605-144, RRID:AB_2338078) anti-rabbit secondary antibody for 1.5-2 hours at RT. Nuclei

were counterstained with 4′,6-diamidino-2-phenylindole (DAPI, 1:500-1000, ThermoFisher

Invitrogen), and Fluoromount-G was used to coverslip (Southernbiotech, Birmingham, AL,

USA).

Fluorescent In-Situ Hybridization

Perfused brains (4% PFA) from female mice were coronally cryo-sectioned into six series at 20

µm. Sections containing the MPO, VMH, and ARH were anatomically-matched across mice.

Fixed tissue underwent single molecule *in-situ* hybridization according to manufacturer

instructions in the RNAscope Multiplex Fluorescent Detection Kit version 2 (Advanced Cell

Diagnostics, Newark, CA, USA). Tissue was fixed in 10% neutral buffered formalin for 20

minutes and pre-treated using serial ethanol dilutions, hydrogen peroxide, and protease III

solution (incubation time shortened to 17 minutes to prevent tissue degradation). Tissue was

incubated with probes targeting *Rprm* (Cat# 466071), *Cre* (Cat# 1058921)*, and *Esr1* (Cat #

478201) at 40°C for 2 hours. Afterward, the probe signal was amplified and fluorescently labeled

with Opal 520 (1:750), 570 (1:500), and 690 (1:500; Akoya Biosciences, MA) for *Esr1*, *Rprm*

and *Cre*, respectively. Signal was developed using horseradish peroxidase and counterstained

with reagents provided in the kit. Slides were cover-slipped using Prolong Gold Antifade

(ThermoFisher Scientific).

Imaging & analysis

Brightfield images of BAT and iWAT histological sections were taken by a Leica DM1000

microscope (Leica Microsystems Inc, Deerfield, IL, USA). Brain, uterine and ovarian sections

were imaged using a Nikon Eclipse Ti2 inverted microscope. Peripheral tissues were illuminated

using a Leica MZ10F and imaged using a dual MP camera system on an iPhone13. ERα cells

were counted in the anteroventral periventricular area (AVPV), medial preoptic area (MPO),

ventromedial ventrolateral area (VMHvl), and arcuate nucleus (ARH) of the hypothalamus. The

machine learning software Ilastik (Berg et al., 2019; www.ilastik.org/about) segmented ERα+

cells from the background. Positive object outlines were loaded to CellProfiler 4.2.5 (Stirling et

al., 2021; www.cellprofiler.org), overlaid over the fluorescent images, manually edited if

necessary, and counted. Cell counts were averaged across hemispheres; if one hemisphere was

damaged, the count on the intact side was used.

For *in situ* hybridization, cell counts in the MPO, VMH, and ARH were done in one hemisphere

using ImageJ Version 2.1.0/1.53c. A cell was considered positive for a given gene if it contained

more than approximately 7 puncta and co-localized with DAPI. For studies interrogating the co-

localization of *Cre*/*Rprm* or *Cre*/*Esr1*, each channel was analyzed independently and channels

were then merged to identify co-expressing cells. For all analyses, regions of interest were

determined by examining the shape of the anterior commissure, median eminence, and third

ventricle referencing the Allen Brain Mouse (v2, 2011) and Paxinos and Franklin (v3, 2008)

atlases. Area-proportional Venn diagrams, generated using BioVenn [40], show the mean number

of cells expressing *Esr1* only, *Cre* only, or *Esr1* and *Cre* in each brain region. To estimate the

proportion of the *Esr1*+ population that expresses *Rprm-Cre*, the number of *Esr1+/Cre+* cells

was divided by the total number of *Esr1*+ cells (*Esr1+/Cre+* and *Esr1+/Cre-*) and multiplied by

100 to get a percentage. This proportion was calculated per animal, averaged across the group,

and described in text as an average ± SEM.

Statistics

Statistical analyses and plots were made in Prism (version 10.2.3 for Mac; GraphPad Software,

Boston, MA, USA). When applicable, two-way ANOVAs or mixed-effects models were run with

repeated (age or ZT) and independent (genotype or sex) factors. Post hoc comparisons between

control and RERKO mice at each time point were performed using Šídák's multiple comparisons

test. Comparisons between two samples were tested with t-tests, as indicated in the figure

legends.

Results

3.1 Validation of *Rprm^Cre* mouse

We bred *Rprm^Cre* mice with *Ai14* reporter mice to identify the tissues expressing *Rprm* at any

point through development. We report *Cre*-mediated expression of TdTomato in the brain,

uterus, ovary, pancreas and pituitary (Supp. Fig. 1A,B). Conversely, TdTomato expression was

absent from BAT, iWAT, and gWAT (Supp. Fig. 1C). We note that the brain exhibited

widespread TdTomato expression that is most striking in the cortex. This was somewhat

expected as *Rprm* expression is pronounced in the cortex and extends to many regions in

development. Notably, *Rprm* expression has an interesting developmental pattern, peaking just

before birth (E18.5) and dampening down by weaning (Allen Institute for Brain Science

developing mouse brain, ISH data, entrez ID: 67874).

Therefore, to validate the fidelity of the transgenic allele, we assessed the expression of *Cre* and

Rprm transcripts in adult mice. We analyzed three regions that abundantly express *Esr1*: the

MPO, VMHvl, and ARH (Fig. 1B-F). In the VMHvl and ARH, *Cre* expression was exclusively

restricted to *Rprm+* cells (Fig. 1E,F). This was also the case in the MPO, except for a small

number of *Cre+/Rprm-* and *Rprm+/Cre-* cells (Fig. 1D). Next, we quantified the number of

Esr1+, *Cre+* and co-expressing *Esr1+/Cre+* cells (Fig. 1G-I) and calculated the proportion of

Esr1+ cells that co-express *Cre+* (and by extension, *Rprm*), reporting 58.9 ± 6.0 % in the MPO,

39.5 ± 3.4 % in the VMHvl, and 11.3 ± 0.9 % in the ARH, consistent with previous work [11].

Finally, we confirmed that the *Cre* knock-in allele does not exhibit ectopic *Rprm* expression in

the MPO and VMHvl by finding no difference in *Rprm+* cell counts between *Rprm^Cre* and WT

C57BL/6J controls (Supp. Fig 2). Together, these results validate the fidelity of the expression of

the *Rprm^Cre* allele in adults. Therefore, the widespread expression observed with lineage tracing

likely reflects transient expression of *Rprm* throughout the brain and other tissues that

extinguishes prior to adulthood.

3.2 *Esr1* knockout in *Rprm* lineage cells (RERKO) leads to an increase in brown adipose tissue

mass, specifically in females

Next, we knocked out *Esr1* in *Rprm*-lineage cells which, based on TdTomato and expected *Esr1*

expression, is expected to occur in the hypothalamus, uterus, ovary, pancreas, and pituitary

[41,42]. While previous studies report that global ERα knockout or VMHvl-specific ERα knock-

down increases body weight, feeding, and fat mass in mice [16,32], we find that body weight in

RERKO mice is largely unchanged, relative to controls (Supp. Fig. 3). Male RERKO mice

became subtly heavier as they aged (genotype x age interaction: $F_{11,318} = 2.398$, $p < 0.0001$),

while females did not differ from controls. However, we observed a sex-specific effect of

RERKO on BAT mass (genotype x sex interaction: $F_{1,40} = 24.5$, $p < 0.0001$; Fig. 2A) and no

effect on iWAT (Fig. 2B). Female RERKO mice had larger BAT depots than controls ($p <$

0.0001), while BAT size in males did not differ (Fig. 2A). BAT tissue in RERKO mice appeared

lighter in color and typically had larger adipose droplets relative to controls (Fig. 2C, D) –

although an analysis comparing nuclei and adiposity in BAT did not reach significance (Supp.

Fig. 4). In contrast to global *Esr1* KO [16,32], RERKO mice have no changes in adiposity: body

mass and percent fat and lean mass as recorded via EchoMRI were similar between RERKO and

control littermates in a separate cohort of animals (Fig. 2E-G). In addition, RERKO mice did not

consume more food relative to controls (Fig. 2H) as reported in CNS-restricted *Esr1* KO mice

[9]. Together, these data suggest that *Esr1* in *Rprm* lineage cells may regulate discrete facets of

metabolism (*i.e.*, BAT), while sparing energy intake and storage.

3.3 *Esr1* knockout in *Rprm* lineage cells sex-specifically impacts energy expenditure

Because BAT contributes to thermoregulation, we next tested if RERKO mice had

changes in body temperature by implanting telemetry probes that simultaneously and passively

record core, BAT, and tail temperature (Fig. 3A). In males, core temperature and locomotion

were unaffected by genotype (Fig. 3B, C), consistent with no change in BAT mass. In contrast,

both temperature (genotype x ZT: $F_{23,437} = 6.0$, $p < 0.0001$; Fig 3D) and locomotion (ZT x

genotype: $F_{23,391} = 1.7$, $p = 0.02$; Fig. 3E) differed between WT and RERKO female mice.

Probing further, we compared BAT and tail temperature as proxies for heat generation and

dissipation, respectively. Female RERKO mice had higher BAT temperature than WT mice
(main effect of genotype: $F_{1,14} = 4.7$, $p < 0.05$; Fig. 3F) and lower tail temperature than WT mice
(main effect of genotype: $F_{1,6} = 8.6$, $p < 0.05$; Fig. 3G). Hence, female RERKO mice appear to
produce more heat and dissipate less heat, contributing to their altered core temperature
phenotype. In line with the morphological findings, ERα in *Rprm+* cells appears to specifically
regulate thermogenesis in a sex-specific manner.

In addition to altered thermogenesis, RERKO female mice appeared to have reproductive
deficits. RERKO mice did not cycle well, mostly remaining in diestrus or metestrus (Supp. Fig.
4A, B). RERKO mice had heavier ovarian ($t_8 = 2.5$, $p < 0.05$) and decreased uterine masses ($t_{23} =$
5.9, $p < 0.0001$; Supp. Fig. 5D), and histological examination revealed abnormal ovarian and
uterine anatomy (Supp. Fig. 5C), including the presence of follicular cysts and the absence of
corpora lutea. Sex steroid hormones influence thermoregulation [43,44]; thus, it is possible that
differing hormonal milieu between WT and RERKO underlie the temperature phenotype. To
address this, females were ovariectomized, and temperature was compared between WT
littermates and RERKO mice seven days later. There was a main effect of genotype ($F_{1,12} = 5.5$,
$p < 0.05$) and an interaction between genotype and hour of day ($F_{23,270} = 1.7$, $p < 0.05$; Supp. Fig.
6). Ovariectomized RERKO mice had elevated temperature relative to ovariectomized controls,
particularly during the light phase (Supp. Fig. 6). Thus, RERKO mice have altered
thermoregulation, that is at least partially, independent from changes in ovarian hormones.

3.4 *Esr1* knockout in *Rprm* cells reduces ERα immunoreactivity in thermoregulatory hypothalamic
regions.
In the brain, ERα is expressed in brain regions involved in thermoregulation and energy intake
and expenditure (*e.g.*, MPO, VMHvl, and the ARH) [12,45], and *Rprm* is co-expressed in all of
these regions (Fig. 1). RERKO mice had fewer ERα-expressing cells in the AVPV, VMHvl, and
ARH (Fig. 4D, F, G), but not in the MPO (Fig. 4E). We note that the sex difference in ERα in the
VMHvl was completely abolished in RERKO mice, due to a reduction in both sexes (Figure 4F).
We did not have sufficient power to confirm the expected sex difference in ERα cell counts in
the AVPV, but nevertheless, observed a decrease in RERKO mice relative to controls in males
and females (Fig. 4D).

3.5 Ablating *Rprm+* cells in the MBH decreases core temperature and increases fat depots.

We hypothesized that the temperature phenotype in RERKO mice may involve the VMHvl given

our previous work [11]. To test this, we delivered an AAV expressing a Cre-dependent and self-

activating caspase into the VMHvl of $Rprm^{Cre}$ mice to ablate $Rprm+$ cells in this brain region

[38]. We labeled ERα to validate caspase-mediated cell death of $Esr1/Rprm$ cells in the VMHvl

and ARH and observed decrease in ERα+ immunoreactivity between controls and caspase-

treated mice in the VMHvl, but not ARH (Figure 5C). Although we did not detect a decrease in

ERα+ cell counts in the ARH, we note that this does not rule out the possible loss of other

$Rprm+$ cells in a broader area of the MBH. Caspase-treated mice had decreased core temperature

(main effect of group: $F_{1,10}= 11.20$, $p < 0.01$; Figure 5D) and heavier BAT ($t_{10} = 4.02$, $p < 0.01$,

Figure 5G), iWAT ($t_{10} = 2.23$, $p = 0.05$; Figure 5H) and gWAT ($t_{12} = 4.03$, $p < 0.01$; Figure 5I).

Statistical differences in activity levels (Figure 5E), total body mass (Figure 5F), or uterine mass

(Figure 5J) were not detected. This provides additional evidence that $Rprm+$ neurons in the MBH

regulate body temperature [11]. Importantly, we observe these effects without an effect on the

uterus, a bioassay for circulating estrogen levels [46], suggesting that this effect is not secondary

to any potential changes in circulating estrogens.

Discussion

We investigated the role of $Esr1$ in $Rprm+$ cells and the specific contribution of MBH^{Rprm}

neurons on energy homeostasis. First, we characterized the $Rprm$ lineage using a novel $Rprm^{Cre}$

mouse and an $Ai14$ reporter. We observed $Rprm$-driven tdTomato expression in the pancreas,

uterus, ovary, pituitary, and brain. Interestingly, we did not observe evidence of $Rprm$ expression

in fat depots associated with heat generation or retention, including BAT, iWAT, or gWAT. It is

intriguing that $Rprm$ appears predominantly expressed in hormone-producing or hormone-

transporting tissues (*i.e.*, ovaries, pituitary, pancreas, and brain). In particular, these tissues are

estrogen sensitive [47–49], raising the possibility that $Rprm$ may play a role in the development

or function of hormone-sensitive tissues. In contrast to the widespread expression of tdTomato,

which labels the the $Rprm$ lineage, we found that Cre expression in adults is more restricted and

a reliable indicator of endogenous $Rprm$ expression. The findings in adults validate the fidelity of

the $Rprm^{Cre}$ mouse line and suggest that tdTomato expression is a result of previous, transient

$Rprm$ expression in $Ai14^{fl/+};Rprm^{Cre}$ mice. Future studies will use tools that are spatially or

temporally restricted, such as AAVs injected to specific tissues in adulthood, in $Rprm^{Cre}$ mice to

understand the role of cells or tissues that actively express $Rprm$ at the time of manipulation.

Here we tested the role of *Esr1* in *Rprm*-lineage cells using *Rprm*^{Cre}; *Esr1*$^{f/f}$ (RERKO) mice.

In the adult brain, *Esr1* and *Rprm* are co-expressed in at least four hypothalamic nuclei: the

AVPV, the MPO, the VMHvl, and the ARH. RERKO mice had reduced ERα+ cell counts in all

of these regions, except the MPO. These regions are associated with energy homeostasis and

reproduction prompting us to compare these phenotypes in wildtype and RERKO mice.

As mice aged to four months, there was a subtle increase in body weight in males, but not

females. However, there were no significant differences in fat mass, lean mass, or food

consumption between RERKO mice and their WT siblings in male or female mice. We observed

a reduction in ERα+ cells in the ARH but did not detect an effect on food intake, suggesting that

the *Rprm* lineage does not substantially contribute to ERα cells in the ARH that modulate food

intake. The most notable phenotype was a doubling of BAT mass that was specific to females.

This was surprising since previous studies on global *Esr1* KO mice report significant weight gain

and adiposity, but no change in BAT mass [32,33]. Future studies could assess if there are

differences in thermogenic capabilities of BAT tissue. In summary, while *Esr1* was knocked out

in the *Rprm* lineage, which extends to multiple tissues, RERKO mice exhibited a more restricted

phenotype than global *Esr1* KO.

Because we found the strongest effect of RERKO on the mass of thermogenic BAT, we

compared the temperature of the core, BAT and tail skin. Interestingly, we found effects on core

temperature only in females, not males, suggesting that estrogen signaling in the *Rprm* lineage is

particularly important for maintaining normal temperatures in females. In females, there was no

clear directionally of the temperature change; instead RERKO females appear to show reduced

differences in temperature between day and night. A similar pattern is observed in BAT and tail

temperature, although we only detected a main effect of genotype and no effect of the interaction

between genotype and time. Thus, overall energy expenditure may be unchanged, as suggested

by no change in body mass. Nonetheless, our results suggest that estrogen signaling in *Rprm*

lineage cells may help maintain day-night patterns of core temperature by modulating both heat

generation and dissipation processes.

In addition to differences in thermoregulatory patterns, we observed reproductive deficits in

female RERKO mice: they did not cycle well and had abnormal reproductive anatomy. Thus, a

limitation of this mouse model is potential differences in hormone levels, which may indirectly

alter physiology and metabolism. To address this, we ovariectomized RERKO and WT mice and

found persistent differences in core temperature, albeit with a different pattern than intact

RERKO mice suggesting some hormonal effects.

Because we did not detect *Rprm* expression in BAT nor iWAT, we hypothesized that RERKO

mice had altered central regulation of energy expenditure. Movement and BAT-mediated

thermogenesis are associated with the VMHvl [10,11], making it the most promising mediator of

a temperature phenotype. To conclusively test that *Rprm* cells in the brain, and not peripheral

organs, regulate thermogenesis, we ablated *Rprm*+ cells in the MBH by stereotaxic delivery of a

Cre-dependent caspase. We observed a ~40% decrease in the number of ERα cells in the

VMHvl, consistent with our prior observation and confirmation in this paper that 41% of ERα

cells co-express *Rprm* [11]. Caspase-treated mice had lower core temperature than controls with

no changes in movement, similar to previous effects of knocking down *Rprm* in the same region

[11]. We also observed heavier BAT mass in caspase-ablated mice, suggesting an effect on BAT-

mediated thermogenesis. Together, these studies suggest that MBH^{Rprm} cells regulate core body

temperature. Taken together with the effects of ERα KO in the *Rprm* lineage, these findings

suggest that MBH^{Rprm} cells may mediate some of the effects of estrogens on core temperature

and thermogenesis. However, future studies will be required to definitively test this hypothesis

using intersectional genetic approaches to specifically knock out *Rprm* in *Esr*1+ cells in the

VMHvl or MBH.

Conclusion

In summary, we use a new genetically engineered mouse model to show that estrogen-

dependent regulation of cells in the *Rprm* lineage alters temperature, partly by modulating BAT

thermogenesis and heat dissipating processes. Evidence of *Rprm* expression is not found in

brown or white adipose tissue, suggesting that *Esr1* in *Rprm* cells within the brain (as opposed to

in adipose tissue) is critical for proper thermoregulation. Specifically, it appears that *Rprm*+ cells

in the MBH drive patterns of heat generation via BAT. This work adds to a body of literature

showing specialization of sub-populations of ERα neurons [10,13,22,24,50]. By identifying the

cells driving regulation of thermogenesis, our research opens potential avenues for new precision

treatments in metabolic disorders, such as obesity and menopause.

Acknowledgements

The authors thank members of the Correa lab, particularly Paul Vander and Rachel Scott, for their feedback on the manuscript, as well as, Sakina Rashid and Hannah Azadi for technical assistance.

Author contributions

JWP, LRC, and SMC conceptualized the present studies. JEV and LGK designed the *Rprm-Cre-FRT* mouse, and JWP designed the telemetry probe sleeves. JWP, LRC, NPS, ARV, AGB, HJF, MRH, EAD, WB, and KV contributed to experimental research and data collection. JWP, LRC, NPS, ARV, HJF, KV, and JIL provided data analysis and validation. SMC, LRC, and LGK provided funding and/or resources to complete the studies, and SMC provided mentoring and laboratory space. LRC and SMC wrote the manuscript with input from JWP, NPS, and JEV.

Funding Sources

These studies were supported by the National Institute on Aging (R01AG066821) and National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK; R01DK136073) to SMC and by the Eunice Kennedy Shriver National Institute of Child Health and Development (K00HD109205), the Burroughs Wellcome Fund Postdoctoral Enrichment Fellowship, the Iris Cantor-UCLA Women's Health Center Executive Advisory Board (NCATS UCLA Clinical and Translational Science Institute; UL1TR001881), and the NIDDK UCLA LIFT-UP (Leveraging Institutional support for Talented, Upcoming Physicians and/ or Scientists; National Institutes of Health Office of Disease Prevention, ODP; U24DK132746) to LRC. The generation of the *Rprm-Cre-FRT* mouse was funded by a UCLA Brain Research Institute Predoctoral Research Grant to LGK.

Disclosures

Data available

Data published in this article may be shared by emailing SMC.

References

[1] Vigil, P., Meléndez, J., Petkovic, G., Del Río, J.P., 2022. The importance of estradiol for body weight regulation in women. Frontiers in Endocrinology 13, Doi: 10.3389/fendo.2022.951186.

[2] Mahboobifard, F., Pourgholami, M.H., Jorjani, M., Dargahi, L., Amiri, M., Sadeghi, S., et al., 2022. Estrogen as a key regulator of energy homeostasis and metabolic health. Biomedicine & Pharmacotherapy 156: 113808, Doi: 10.1016/j.biopha.2022.113808.

[3] Weidlinger, S., Winterberger, K., Pape, J., Weidlinger, M., Janka, H., von Wolff, M., et al., 2023. Impact of estrogens on resting energy expenditure: A systematic review. Obesity Reviews: An Official Journal of the International Association for the Study of Obesity 24(10): e13605, Doi: 10.1111/obr.13605.

[4] Rehman, A., Lathief, S., Charoenngam, N., Pal, L., 2024. Aging and Adiposity—Focus on Biological Females at Midlife and Beyond. International Journal of Molecular Sciences 25(5): 2972, Doi: 10.3390/ijms25052972.

[5] Lovejoy, J.C., Champagne, C.M., de Jonge, L., Xie, H., Smith, S.R., 2008. Increased visceral fat and decreased energy expenditure during the menopausal transition. International Journal of Obesity (2005) 32(6): 949–58, Doi: 10.1038/ijo.2008.25.

[6] Duval, K., Prud'homme, D., Rabasa-Lhoret, R., Strychar, I., Brochu, M., Lavoie, J.-M., et al., 2013. Effects of the menopausal transition on energy expenditure: a MONET Group Study. European Journal of Clinical Nutrition 67(4): 407–11, Doi: 10.1038/ejcn.2013.33.

[7] Ward, E., Gold, E.B., Johnson, W.O., Ding, F., Chang, P.-Y., Song, P., et al., 2019. Patterns of Cardiometabolic Health as Midlife Women Transition to Menopause: A Prospective Multiethnic Study. The Journal of Clinical Endocrinology & Metabolism 104(5): 1404–12, Doi: 10.1210/jc.2018-00941.

[8] Ou, Y.-J., Lee, J.-I., Huang, S.-P., Chen, S.-C., Geng, J.-H., Su, C.-H., 2023. Association between Menopause, Postmenopausal Hormone Therapy and Metabolic Syndrome. Journal of Clinical Medicine 12(13): 4435, Doi: 10.3390/jcm12134435.

[9] Xu, Y., Nedungadi, T.P., Zhu, L., Sobhani, N., Irani, B.G., Davis, K.E., et al., 2011. Distinct hypothalamic neurons mediate estrogenic effects on energy homeostasis and reproduction. Cell Metabolism 14(4): 453–65, Doi: 10.1016/j.cmet.2011.08.009.

[10] Correa, S.M., Newstrom, D.W., Warne, J.P., Flandin, P., Cheung, C.C., Lin-Moore, A.T., et al., 2015. An estrogen-responsive module in the ventromedial hypothalamus selectively drives sex-specific activity in females. Cell Reports 10(1): 62–74, Doi: 10.1016/j.celrep.2014.12.011.

[11] van Veen, J.E., Kammel, L.G., Bunda, P.C., Shum, M., Reid, M.S., Massa, M.G., et al., 2020. Hypothalamic estrogen receptor alpha establishes a sexually dimorphic regulatory node of energy expenditure. Nature Metabolism 2(4): 351–63, Doi: 10.1038/s42255-020-0189-6.

[12] Zhang, Z., Reis, F.M.C.V., He, Y., Park, J.W., DiVittorio, J.R., Sivakumar, N., et al., 2020. Estrogen-sensitive medial preoptic area neurons coordinate torpor in mice. Nature Communications 11(1): 6378, Doi: 10.1038/s41467-020-20050-1.

[13] Krause, W.C., Rodriguez, R., Gegenhuber, B., Matharu, N., Rodriguez, A.N., Padilla-Roger, A.M., et al., 2021. Oestrogen engages brain MC4R signalling to drive physical activity in female mice. Nature 599(7883): 131–5, Doi: 10.1038/s41586-021-04010-3.

[14] Rogers, N.H., Perfield, J.W., Strissel, K.J., Obin, M.S., Greenberg, A.S., 2009. Reduced
energy expenditure and increased inflammation are early events in the development of
ovariectomy-induced obesity. Endocrinology 150(5): 2161–8, Doi: 10.1210/en.2008-1405.
[15] Camporez, J.P.G., Jornayvaz, F.R., Lee, H.-Y., Kanda, S., Guigni, B.A., Kahn, M., et al.,
2013. Cellular Mechanism by Which Estradiol Protects Female Ovariectomized Mice From
High-Fat Diet-Induced Hepatic and Muscle Insulin Resistance. Endocrinology 154(3):
1021–8, Doi: 10.1210/en.2012-1989.
[16] Musatov, S., Chen, W., Pfaff, D.W., Kaplitt, M.G., Ogawa, S., 2006. RNAi-mediated
silencing of estrogen receptor α in the ventromedial nucleus of hypothalamus abolishes
female sexual behaviors. Proceedings of the National Academy of Sciences 103(27):
10456–60, Doi: 10.1073/pnas.0603045103.
[17] Sano, K., Tsuda, M.C., Musatov, S., Sakamoto, T., Ogawa, S., 2013. Differential effects of
site-specific knockdown of estrogen receptor α in the medial amygdala, medial pre-optic
area, and ventromedial nucleus of the hypothalamus on sexual and aggressive behavior of
male mice. European Journal of Neuroscience 37(8): 1308–19, Doi: 10.1111/ejn.12131.
[18] Frazao, R., Lemko, H.M.D., da Silva, R.P., Ratra, D.V., Lee, C.E., Williams, K.W., et al.,
2014. Estradiol modulates Kiss1 neuronal response to ghrelin. American Journal of
Physiology - Endocrinology and Metabolism 306(6): E606–14, Doi:
10.1152/ajpendo.00211.2013.
[19] Qiu, J., Zhang, C., Borgquist, A., Nestor, C.C., Smith, A.W., Bosch, M.A., et al., 2014.
Insulin excites anorexigenic proopiomelanocortin neurons via activation of canonical
transient receptor potential channels. Cell Metabolism 19(4): 682–93, Doi:
10.1016/j.cmet.2014.03.004.
[20] Padilla, S.L., Perez, J.G., Ben-Hamo, M., Johnson, C.W., Sanchez, R.E.A., Bussi, I.L., et
al., 2019. Kisspeptin Neurons in the Arcuate Nucleus of the Hypothalamus Orchestrate
Circadian Rhythms and Metabolism. Current Biology 29(4): 592-604.e4, Doi:
10.1016/j.cub.2019.01.022.
[21] Rønnekleiv, O.K., Qiu, J., Kelly, M.J., 2019. Arcuate Kisspeptin Neurons Coordinate
Reproductive Activities with Metabolism. Seminars in Reproductive Medicine 37(3): 131–
40, Doi: 10.1055/s-0039-3400251.
[22] Hashikawa, K., Hashikawa, Y., Tremblay, R., Zhang, J., Feng, J.E., Sabol, A., et al., 2017.
Esr1+ cells in the ventromedial hypothalamus control female aggression. Nature
Neuroscience 20(11): 1580–90, Doi: 10.1038/nn.4644.
[23] Lo, L., Yao, S., Kim, D.-W., Cetin, A., Harris, J., Zeng, H., et al., 2019. Connectional
architecture of a mouse hypothalamic circuit node controlling social behavior. Proceedings
of the National Academy of Sciences 116(15): 7503–12, Doi: 10.1073/pnas.1817503116.
[24] He, Y., Xu, P., Wang, C., Xia, Y., Yu, M., Yang, Y., et al., 2020. Estrogen receptor-α
expressing neurons in the ventrolateral VMH regulate glucose balance. Nature
Communications 11(1): 2165, Doi: 10.1038/s41467-020-15982-7.
[25] Silva, N.L., Boulant, J.A., 1986. Effects of testosterone, estradiol, and temperature on
neurons in preoptic tissue slices. The American Journal of Physiology 250(4 Pt 2): R625-
632, Doi: 10.1152/ajpregu.1986.250.4.R625.
[26] Song, Z., Levin, B.E., McArdle, J.J., Bakhos, N., Routh, V.H., 2001. Convergence of pre-
and postsynaptic influences on glucosensing neurons in the ventromedial hypothalamic
nucleus. Diabetes 50(12): 2673–81, Doi: 10.2337/diabetes.50.12.2673.

[27] Ohki, R., Nemoto, J., Murasawa, H., Oda, E., Inazawa, J., Tanaka, N., et al., 2000.
Reprimo, a new candidate mediator of the p53-mediated cell cycle arrest at the G2 phase.
The Journal of Biological Chemistry 275(30): 22627–30, Doi: 10.1074/jbc.C000235200.
[28] Xu, M., Knox, A.J., Michaelis, K.A., Kiseljak-Vassiliades, K., Kleinschmidt-DeMasters,
B.K., Lillehei, K.O., et al., 2012. Reprimo (RPRM) Is a Novel Tumor Suppressor in
Pituitary Tumors and Regulates Survival, Proliferation, and Tumorigenicity. Endocrinology
153(7): 2963–73, Doi: 10.1210/en.2011-2021.
[29] Ye, Z., Wang, J., Shi, W., Zhou, Z., Zhang, Y., Wang, J., et al., 2023. Reprimo (RPRM) as a
Potential Preventive and Therapeutic Target for Radiation-Induced Brain Injury via
Multiple Mechanisms. International Journal of Molecular Sciences 24(23): 17055, Doi:
10.3390/ijms242317055.
[30] Figueroa, R.J., Carrasco-Avino, G., Wichmann, I.A., Lange, M., Owen, G.I., Siekmann,
A.F., et al., 2017. Reprimo tissue-specific expression pattern is conserved between zebrafish
and human. PloS One 12(5): e0178274, Doi: 10.1371/journal.pone.0178274.
[31] Nedergaard, J., Cannon, B., 2018. Brown adipose tissue as a heat-producing thermoeffector.
Handbook of Clinical Neurology 156: 137–52, Doi: 10.1016/B978-0-444-63912-7.00009-6.
[32] Heine, P.A., Taylor, J.A., Iwamoto, G.A., Lubahn, D.B., Cooke, P.S., 2000. Increased
adipose tissue in male and female estrogen receptor-α knockout mice. Proceedings of the
National Academy of Sciences 97(23): 12729–34, Doi: 10.1073/pnas.97.23.12729.
[33] Ohlsson, C., Hellberg, N., Parini, P., Vidal, O., Bohlooly-Y, M., Rudling, M., et al., 2000.
Obesity and disturbed lipoprotein profile in estrogen receptor-alpha-deficient male mice.
Biochemical and Biophysical Research Communications 278(3): 640–5, Doi:
10.1006/bbrc.2000.3827.
[34] Kim, J.H., Lee, S.-R., Li, L.-H., Park, H.-J., Park, J.-H., Lee, K.Y., et al., 2011. High
cleavage efficiency of a 2A peptide derived from porcine teschovirus-1 in human cell lines,
zebrafish and mice. PloS One 6(4): e18556, Doi: 10.1371/journal.pone.0018556.
[35] Feng, Y., Manka, D., Wagner, K.-U., Khan, S.A., 2007. Estrogen receptor-α expression in
the mammary epithelium is required for ductal and alveolar morphogenesis in mice.
Proceedings of the National Academy of Sciences of the United States of America 104(37):
14718–23, Doi: 10.1073/pnas.0706933104.
[36] Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., et al., 2010.
A robust and high-throughput Cre reporting and characterization system for the whole
mouse brain. Nature Neuroscience 13(1): 133–40, Doi: 10.1038/nn.2467.
[37] Chow, B.Y., Han, X., Dobry, A.S., Qian, X., Chuong, A.S., Li, M., et al., 2010. High-
performance genetically targetable optical neural silencing by light-driven proton pumps.
Nature 463(7277): 98–102, Doi: 10.1038/nature08652.
[38] Yang, C.F., Chiang, M.C., Gray, D.C., Prabhakaran, M., Alvarado, M., Juntti, S.A., et al.,
2013. Sexually Dimorphic Neurons in the Ventromedial Hypothalamus Govern Mating in
Both Sexes and Aggression in Males. Cell 153(4): 896–909, Doi:
10.1016/j.cell.2013.04.017.
[39] Byers, S.L., Wiles, M.V., Dunn, S.L., Taft, R.A., 2012. Mouse Estrous Cycle Identification
Tool and Images. PLoS ONE 7(4): e35538, Doi: 10.1371/journal.pone.0035538.
[40] Hulsen, T., de Vlieg, J., Alkema, W., 2008. BioVenn – a web application for the comparison
and visualization of biological lists using area-proportional Venn diagrams. BMC Genomics
9(1): 488, Doi: 10.1186/1471-2164-9-488.

[41] Couse, J.F., Lindzey, J., Grandien, K., Gustafsson, J.-Å., Korach, K.S., 1997. Tissue Distribution and Quantitative Analysis of Estrogen Receptor-α (ERα) and Estrogen Receptor-β (ERβ) Messenger Ribonucleic Acid in the Wild-Type and ERα-Knockout Mouse. Endocrinology 138(11): 4613–21, Doi: 10.1210/endo.138.11.5496.

[42] Le May, C., Chu, K., Hu, M., Ortega, C.S., Simpson, E.R., Korach, K.S., et al., 2006. Estrogens protect pancreatic β-cells from apoptosis and prevent insulin-deficient diabetes mellitus in mice. Proceedings of the National Academy of Sciences 103(24): 9232–7, Doi: 10.1073/pnas.0602956103.

[43] Sanchez-Alavez, M., Alboni, S., Conti, B., 2011. Sex- and age-specific differences in core body temperature of C57Bl/6 mice. Age (Dordrecht, Netherlands) 33(1): 89–99, Doi: 10.1007/s11357-010-9164-6.

[44] Krajewski-Hall, S.J., Blackmore, E.M., McMinn, J.R., Rance, N.E., 2018. Estradiol alters body temperature regulation in the female mouse. Temperature (Austin, Tex.) 5(1): 56–69, Doi: 10.1080/23328940.2017.1384090.

[45] Frank, A., Brown, L.M., Clegg, D.J., 2014. The role of hypothalamic estrogen receptors in metabolic regulation. Frontiers in Neuroendocrinology 35(4): 550–7, Doi: 10.1016/j.yfrne.2014.05.002.

[46] Ingberg, E., Theodorsson, A., Theodorsson, E., Strom, J.O., 2012. Methods for long-term 17β-estradiol administration to mice. General and Comparative Endocrinology 175(1): 188–93, Doi: 10.1016/j.ygcen.2011.11.014.

[47] Rubanyi, G.M., Kauser, K., Johns, A., 2002. Role of estrogen receptors in the vascular system. Vascular Pharmacology 38(2): 81–8, Doi: 10.1016/S0306-3623(02)00130-1.

[48] Gieske, M.C., Kim, H.J., Legan, S.J., Koo, Y., Krust, A., Chambon, P., et al., 2007. Pituitary Gonadotroph Estrogen Receptor-α Is Necessary for Fertility in Females. Endocrinology 149(1): 20, Doi: 10.1210/en.2007-1084.

[49] Nadal, A., Alonso-Magdalena, P., Soriano, S., Quesada, I., Ropero, A.B., 2009. The pancreatic beta-cell as a target of estrogens and xenoestrogens: Implications for blood glucose homeostasis and diabetes. Molecular and Cellular Endocrinology 304(1–2): 63–8, Doi: 10.1016/j.mce.2009.02.016.

[50] Hirschberg, P.R., Sarkar, P., Teegala, S.B., Routh, V.H., 2020. Ventromedial hypothalamus glucose-inhibited neurones: A role in glucose and energy homeostasis? Journal of Neuroendocrinology 32(1): e12773, Doi: 10.1111/jne.12773.

Figure 1: Characterization of a novel *Rprm^Cre* knock-in mouse model. (A) Schematic of the knock-in construct. FRT = flippase recognition target sites, UTR = untranslated region, NLS = nuclear localization sequence. (B, C) Single-molecule *in situ* hybridization labeling *Rprm* (magenta), *Cre* (Cyan) and *Esr1^+* (yellow) cells in coronal sections of the brain containing the (B) MPO and (C) VMHvl & ARH in female mice. (D-F) Quantification of cells expressing *Rprm* alone (magenta), *Cre* alone (*Cyan*), and both *Rprm* and *Cre* (purple) in the MPO (D), VMHvl (E), and ARH (F) show that more than 91% of *Rprm^+* cells express *Cre* (n = 6-9 per region). (G-I) Average number of cells expressing *Esr1* alone (yellow), *Cre* alone (Cyan), and both *Esr1* and *Cre* (green) are depicted in the MPO (G), VMHvl (H), and ARH (I; n = 6-9 per group). The mean ± SEM are depicted in D-I, as well as individual data points in D-F. Scale bar = 100 μm. MPO = medial preoptic area, MPN = medial preoptic nucleus, VMHvl = ventrolateral area of the ventromedial nucleus of the hypothalamus, ARH = arcuate nucleus, 3V = third ventricle.

Figure 2. *Esr1* **KO in** *Rprm* **lineage cells (RERKO) sex-specifically alters brown adipose tissue.** (A,B) BAT (A) and iWAT (B) mass from adult male and female control littermates (black dots) and RERKO mice (gray dots). N = 10 WT males, 11 WT females, 9 RERKO males, and 14 RERKO females. (C) Representative images of iWAT (top) and BAT (bottom) in WT (left) and RERKO (right) female mice. Ruler with centimeters added for scale. (D) Representative images of hematoxylin and eosin staining of WT (top) and RERKO (bottom) in 8-week-old mice. Scale = 20 μm. (E-G) Body composition analysis of male and female WT and RERKO mice at 16 weeks of age via EchoMRI: body mass (E), percent fat mass (F), and percent lean mass (G). N = 5 WT males, 5 WT females, 5 RERKO males, and 6 RERKO females. H) Daily food consumption, accounting for spillage, averaged over seven days in WT and RERKO mice. N = 8 WT males, 6 WT females, 9 RERKO males, and 10 RERKO females. The mean ± SEM are depicted on all graphs, as well as individual data points. Two-way ANOVAs used for statistical analysis; differences between WT and RERKO tested within each sex using Šídák's multiple comparisons test following a significant interaction. ** $p < .01$, *** $p < 0.001$, **** $p < 0.0001$.

Figure 3. KO of *Esr1* in *Rprm* lineage cells sex-specifically impacts thermoregulation.
(A) Illustration of a mouse implanted with an intraperitoneal probe to record temperature and locomotion. In addition, a probe is placed above the BAT and on the base of the tail to measure heat generation and heat dissipation, respectively. Custom 3d printed sleeves are used to secure the probes. (B-E) Temperature in Celsius or activity counts of WT siblings (black) and RERKO mice (males: blue, females: pink) averaged hourly across 24 hours. N = 8 WT males, 8 RERKO males, 11 WT females, 10 RERKO females. (F, G) Temperature averaged hourly above the BAT or at base of the tail in female control (black) and RERKO (pink) across 24 hours. N for BAT = 8 WT, 8 RERKO; N for tail = 4 WT, 4 RERKO. Mean ± SEM plotted at each time point. Two-way mixed ANOVAs used for statistical analyses. Significant interaction followed by Šídák's multiple comparisons at each time point. ZT = Zeitgeber time. C = Celsius. Shaded box indicates the dark (active) phase of the day.

Figure 4. RERKO mice have reduced ERα immunoreactivity in hypothalamic thermoregulatory hubs.
(A-C) Images of ERα immunoreactivity (red) in coronal sections of the brain containing the anteroventral periventricular area (A; AVPV), medial preoptic area (B; MPO), ventrolateral area of the ventromedial hypothalamus (C; VMHvl), and arcuate nucleus (C; ARH). Tissue is counterstained with DAPI (blue). (D-G) Quantification of ERα+ cells in male and female WT (black dots) and RERKO (gray dots) mice in the AVPV (D), MPO (E), VMHvl (F) and ARH (G) (N = 2 - 8 per group). Statistical analysis performed using Two-way ANOVA followed by Šídák's multiple comparisons test, if applicable. The mean ± SEM are depicted. Individual data points are shown in figures. ** $p < .01$, *** $p < 0.001$, **** $p < 0.0001$. Scale bars = 100 μm. Boundaries of nuclei and landmarks outlined to aid in visualization. LPO = lateral preoptic area, MPN = medial preoptic nucleus, VLPO = ventrolateral preoptic area. dm = dorsomedial, c = caudal, 3V = third ventricle.

Figure 5. Ablating *Reprimo+* cells in the MBH reduces core temperature in female mice. (A) Illustration depicting the stereotaxic delivery of a cre-dependent caspase or control virus targeting the VMHvl of female mice, with possible spread to other areas of the MBH. (B) Image of ERα immunoreactivity (yellow) in a control (GFP; left) and caspase-treated (casp; right) mouse. VMHvl delineated to aid in visualization. (C) Quantification of ERα+ cells in the VMHvl (left) and ARH (right) in control (gray bar) and caspase-treated (purple bar) female mice. (E, F) Core temperature and activity counts averaged hourly across 24 hours in control (black) and caspase-treated (purple) mice. (G) Body mass of control (GFP) and caspase-treated mice at 8 weeks of age. (H-K) Mass of dissected BAT (H), iWAT (I), gonadal WAT (J), and uterus (K) from GFP and caspase mice. N = 5 GFP and N = 7 caspase for all plots. Statistical analysis performed using mixed model Two-way ANOVAs (D-E) or t-tests (F-J). The mean ± SEM are depicted. Individual data points are shown in figures C-D and G-K.* $p < 0.05$, ** $p < .01$, **** $p < 0.0001$. Scale bar = 100 μm. VMH = ventromedial hypothalamus (c = caudal, dm = dorsomedial, and vl = ventrolateral areas), ARH = arcuate nucleus, 3V = third ventricle.