Title

High-throughput synapse profiling reveals cell-type-specific spatial configurations in the

fly brain

Authors

Hongyang Wu[1], Yoh Maekawa[2], Sayaka Eno[1], Shu Kondo[3], Nobuhiro Yamagata[1,4],

Hiromu Tanimoto[1,2]*

Affiliations

[1] Graduate School of Life Sciences, Tohoku University, Sendai, Japan.

[2] Department of Biology, Faculty of Science, Tohoku University, Sendai, Japan.

[3] Department of Biological Science and Technology, Tokyo University of Science, Tokyo,

Japan.

[4] Department of Life Science, Graduate School of Engineering Science, Akita, Akita

University.

* Correspondence: Dr. Hiromu Tanimoto (hiromut@m.tohoku.ac.jp)

Abstract

Characterization of intracellular synapse heterogeneity aides to understand the intricate computational logic of neuronal circuits. Despite recent advances in connectomics, the spatial patterns of synapses and their inter-individual variability remain largely unknown. Using directed split-GFP reconstitution, we achieved visualization of endogenous Bruchpilot (Brp), a presynaptic active zone (AZ) scaffold protein, in a cell-type-specific manner. By developing a high-throughput quantification pipeline, we profiled AZ structures in identified neurons of the mushroom body circuit, where intracellular synaptic patterns are crucial due to compartmentalized connectivity. Quantitative characterization of the pattern of Brp clusters across multiple individuals revealed cell-type-dependent synapse heterogeneity and stereotypy. Furthermore, we discovered previously unidentified sub-compartmental synapse configuration and its transient structural plasticity triggered by associative learning. Synapse profiling thus uncovered multi-layered organizations of AZs, ranging from neighboring synapses to consistent patterns across individuals.

Keywords:

Introduction

The brain is a complex network of neurons, and understanding their wiring strategies

provides crucial insights into their roles in neuronal computation. Release sites in the

presynaptic terminal of *Drosophila* neurons are decorated with electron-dense projections

composed of the active-zone (AZ) scaffold protein Brp [1–7]. These Brp projections serve as a

synapse marker enabling the whole-brain connectomes, comprehensive mapping of

synaptic connections at the electron microscopic level [8–12]. While connectome-based

approaches catalyzed the exploration of cell type distinctions, circuit motifs, and

neurotransmitters, their application to comparisons across different conditions is still

challenging due to the limited throughput of data acquisition and dense reconstruction.

Since the synaptic structure and connectivity reflect substantial variability of development

and experiences [11,13–18], cross-individual comparisons are necessary. Fluorescence labeling

offers high throughputs and contents, especially when comparing synaptic organizations

across cell-types [19] or brain regions [20]. Nevertheless, characterizing endogenous synapses in

specific cells using fluorescence-labeling remains challenging in the central nervous system

(CNS) due to the dense synaptic connections from different neurons. This has driven

growing interest in designing cell-type-specific fluorescence-tagging strategies for

endogenous synaptic protein [21].

The circuit of *Drosophila* mushroom bodies (MBs) plays a central role in olfactory

associative learning [22]. The major intrinsic MB neuron, known as Kenyon cells (KCs)

respond to odor stimulation and synapse onto the spatially segregated dendrites of MB

output neurons (MBONs) that divide the MB into compartments. Distinct types of

dopaminergic neurons (DANs) synapse onto specific compartments and thereby modulate

KC-MBON synapses in the corresponding compartments [23]. Presynaptic calcium levels of these DANs undergo sub-compartmental GABAergic modulation and inform memory specificity, postulating the distinct synapse structures at individual release sites [24] . Two types of giant interneurons, the anterior paired lateral neuron (APL) and the dorsal paired medial (DPM) neuron extensively ramify across the MB lobes and provide recurrent modulations to KC synapses [25–32]. Considering their comprehensive projections throughout the MB lobes, the modulation of local circuits requires intracellular synaptic tuning. These studies together underscore the importance to spatially distinguish individual synapses within a cell. However, the high AZ density in the MB lobes precluded conventional approaches of fluorescent microscopy from profiling synaptic structures of specific neurons [9].

Here, we present a spatial analysis of individual AZs within the MB circuit at the single-cell resolution, achieved by cell-type-specific visualization of endogenous Brp using the CRISPR/Cas9-mediated split-GFP tagging system. We developed a high-throughput quantification pipeline to systematically profile Brp clusters of individual AZs in different MB-innervating neurons. Characterizing multiple structural parameters revealed a significant synaptic heterogeneity within single neurons and AZ distribution stereotypy across individuals. These cell-type-specific synapse profiles suggest that AZs are organized at multiple scales, ranging from neighboring synapses to across individuals.

Results

Establishing an experimental system for cell-type-specific profiling of active zone structures

As the molecular assembly of synapses is reported to be sensitive to the dosage of Brp, we visualized endogenous Brp instead of transgenic expression of tagged Brp [33]. To label endogenous Brp specifically in designated cell types, we employed the split-GFP tagging system [34,35]. Using CRISPR/Cas9, we inserted the GFP_{11} fragment (the eleventh β-strand of the super-folder GFP) just prior to the stop codon of *brp* gene. The GFP_{1-10} fragment is expressed in cells of interest by the GAL4/UAS system. The self-assembly of the split-GFP fragments is therefore directed only in GAL4-expressing cells and visualizes endogenous Brp through reconstituted GFP fluorescence (Fig. 1A). As a proof of principle, we directed GFP reconstitution and expressed a plasma membrane marker in the single serotonergic DPM neuron using *VT64246-GAL4* (Fig. 1B). Confocal microscopy revealed Brp::reconstituted GFP (Brp::rGFP) signals only to the MB lobes, aligning with DPM terminals [32]. These signals were distributed as discrete puncta within plasma membrane varicosities resembling presynaptic boutons (Fig. 1B), suggesting that Brp::rGFP clusters labeled individual AZs in the DPM neuron.

To systematically profile Brp::rGFP clusters, we developed an image processing pipeline (Fig. 1A). In brief, image deconvolution was applied to reduce out-of-focus light and enhance signal sharpness. Individual Brp::rGFP clusters were then segmented using the 3D suite plugin in Fiji [36], generating the 3D region-of-interests (3D ROIs) that encircle each cluster. Brp::rGFP signal intensity, volume and location of individual clusters were measured using the 3D ROIs. To optimize the segmentation, we systematically varied the

parameters and calculated the F-score to measure detection accuracy against manually

defined ground truth (see Materials and Methods). The F-score is near 1.0 in segmenting

Brp::rGFP clusters within DPM and APL neurons and achieved > 0.90 in KCs with certain

combinations of segmentation parameters (Fig. 1B).

To confirm that Brp puncta represent AZs, we quantified the numbers of Brp::rGFP

clusters in DPM and APL neurons specifically and compared them to the connectome data.

We found that cluster counts closely matched the number of AZs annotated by the

hemibrain connectome (Fig. 1C) [9]. This result validated our method that accurately isolated

Brp::rGFP clusters corresponding to single AZs.

AZ profiles within KCs are compartmentally heterogeneous

Brp accumulation at individual AZs is known to be heterogeneous even within a single

motor neuron [5,6,37,38]. To study such AZ heterogeneity in CNS neurons, we quantified the

signal intensity of individual Brp::rGFP clusters in MB lobes. Using KC subtype-specific

drivers to direct split-GFP reconstitution in γ KC (*MB009B-GAL4*), α'/β' KCs (*MB370B-*

GAL4) and α/β KCs (*MB008B-GAL4*). we found that within-cluster Brp::rGFP intensities

are highly diverse in all KC subtypes (Fig. 2A).

To visualize spatial patterns of Brp::rGFP intensity in each KC subtype, we

reconstructed 3D distribution of Brp::rGFP clusters and color-coded them by signal

intensities (Fig. 2B). We found that clusters showed noticeable intensity differences

between compartments in all subtypes. Even adjacent compartments such as γ1 and γ2, β'2

and β'1, α2 and α3, showed drastic differences (Fig. 2B). Importantly, these compartmental

patterns were stable across individuals (Fig. 2C) [39]. This result suggests that AZs in

different compartments have distinct structures. Since individual KCs of each subtype

synapse onto all compartments [8–10,40], this Brp compartmental heterogeneity is likely

formed intracellularly.

Intracellular heterogeneity of Brp concentration at individual AZs and cell-type

dependent diversity

Super-resolution microscopy has shown that the Brp molecular density at individual AZs is

dynamically adjusted in motor neurons [41,42]. We hypothesized that the amount of Brp does

not necessarily correlate with AZ cluster size. To verify this, we calculated the Brp

concentration by quantifying both the volume (number of voxels within a 3D ROI) and the

Brp::rGFP intensity of each cluster. We found that clusters with similar sizes can exhibit

vastly different signal intensities in the DPM neuron, suggesting the heterogeneity of Brp

concentration at individual AZs (Fig. 3A). Correlation analysis between the Brp::rGFP

intensity and cluster volume revealed a substantial variability among cell types (Fig. 3B-E).

There were even significant differences among KC subtypes. These results suggest cell-

type dependent variability of the Brp concentration of individual clusters.

Stereotypy of intracellular distribution of AZs

To study the spatial distribution of Brp clusters within single neurons, we focused on

DANs. *MB504B-GAL4* labels four types of single PPL1 DANs projecting to distinct

compartments [43]. Using this driver, we analyzed Brp::rGFP distribution in each

compartment and found it uneven in terminals of both PPL1-α2α'2 and α3 (Fig. 4A-B). To

better characterize cluster distribution, we calculated the "AZ density" defined as the

number of Brp::rGFP clusters surrounding a particular cluster within a specified radius (see

Fig. 1A and Materials and Methods for detail). By color-coding Brp::rGFP clusters

according to their AZ density, we found that AZs in PPL1-α3 were more localized to the

core of the α3 compartment. This pattern was consistent across individuals (Fig. 4C-D),

suggesting a stereotyped AZ distribution within PPL1-α3 neurons.

We extended the analysis to two other single pairs of neurons innervating the MBs,

the DPM and APL neurons. Visualizing AZ density revealed a stereotyped and

compartment specific distribution in the DPM neuron (Fig. 5A). Specifically, the AZ

density was constantly high in the α'/β' lobes across all individuals, supporting the known

functional importance of DPM branches in the α'/β' lobes [30]. In contrast, this compartmental

pattern was less pronounced in the APL neuron. Notably, the ratio of AZ density between

α3 and α'3 is strikingly different among individuals (Fig. 5B). Taken together, these data

suggest that the individual variability in AZ distribution depends on the cell type.

Regulation of spatial configuration of AZs

Characteristics of Brp clusters and AZ distribution in KCs revealed spatial

organization at the level of compartments. Recent studies further showed intra-

compartmental variability of synaptic plasticity upon associative learning, prompting us to

analyze sub-compartmental organization of AZs [44,45]. We thus examined potential local

patterns among AZs in proximity by calculating the correlation between nearest neighbor

clusters in terms of their Brp::rGFP intensities. This analysis revealed strong correlations in

KCs and APL but not in DPM (Fig. 6A).

We further analyzed the sub-compartmental AZ patterns by examining the spatial

distribution of Brp::rGFP intensity. To reduce the high frequency variability of Brp::rGFP

intensities (Fig. 2B), we calculated the "local intensity" for each cluster by applying mean

filtering (the average Brp::rGFP intensity of all surrounding clusters within a specified

radius r, see Materials and Methods for detail). By visualizing the local intensity in γ KCs,

we identified AZ "hot spots", sub-compartment-sized small groups of AZs with high local

intensities, on top of the compartmental differences (Fig. 6B). Since Brp molecules are

clustered more in AZ-dense boutons of a single motor neuron [5], we hypothesized that these

hot spots correspond to regions with higher AZ densities. Indeed, we found that hot spots

appear in high AZ densities regions in γ KCs (Fig. 6B). Consistently, correlation between

the local intensity and the AZ density was high in all KC subtypes. In contrast, for DPM

and APL neurons, the local intensity of Brp::rGFP was not associated with AZ density (Fig.

6C-D). Taken together, these results suggest previously unidentified sub-compartmental

synaptic configuration organized across individual KCs.

Associative learning re-organizes sub-compartmental synaptic configuration

Sub-compartmental synaptic configuration may undergo experience-dependent

changes, such as through associative learning [17,18]. To examine such structural plasticity,

we visualized Brp::rGFP specifically in KCs using *R13F02-GAL4* (Fig. 7A) and presented

flies with odor (4-methylcyclohexanol) with a concomitant (paired) or shifted (unpaired)

electric shock (Fig. 7B). The difference between the paired and unpaired groups therefore

represents the effect of associative learning. We prepared the brains at different time points

(ranging from 3 min to 1 day) after conditioning and measured the correlation between the

AZ density and local intensity of individual Brp::rGFP clusters. Strikingly, we found that

associative learning induced a transient modification to the KC synaptic configurations in a

compartment-specific manner (Fig. 7C). This learning-induced structural plasticity was

specific at 90 min after conditioning and disappeared within one day (Fig. 7D), consistent

with previous studies reporting short-lived AZ remodeling in the MBs by conditioning [17,18].

These results showcase the advantage of split-GFP system in reporting the dynamics of

local synapse subsets with high throughputs and contents. Altogether, we suggest that

associative learning induces transient memory traces of local structural plasticity in AZ

configuration.

Discussion

By leveraging the CRISPR/Cas9 genome editing and split-GFP technique, we

systematically profiled presynaptic structures of MB neurons in a cell-type-specific

manner. This approach enabled high-throughput analysis of multiple brain samples and

detailed characterizations, revealing previously inaccessible structural features of AZs in

CNS neurons.

Single-cell level analysis of AZ profiles revealed cell-type-specific stereotypy in

spatial regulation (Fig. 4-6). We found that AZ density in the DPM neurons is consistently

high in the α'/β' lobes across individuals (Fig. 5). This AZ stereotypy of DPM explains the

functional significance of the branches in the α'/β' lobes. Overexpression of a Down

syndrome cell adhesion molecule (DScam) variant in the DPM neurons was shown to

disrupt their innervation except in the α'/β' lobe, while memory formation remained

unaffected [30]. Similarly, our analysis identified the intracellular stereotypy in AZ

distribution of the PPL1-α3 DAN within the α3 compartment (Fig. 4). This sub-

compartmental organization of the single DAN and layered projections of α/β KCs may

underlie the differential dopaminergic modulation of KC subtypes in associative learning [46].

While such stereotypy is less pronounced in APL, we found strong correlations between

Brp::rGFP intensities of neighboring AZs in contrast to DPM (Fig. 6D). This suggests that

AZs in APL neurons are organized more locally, allowing them to modulate microcircuits

composed of KCs, DANs and MBONs [47]. Collectively, the spatial scale of AZ

organizations depends on cell types, reflecting the function of individual MB neurons.

We observed that Brp::rGFP intensity correlates strongly with the AZ size in KC

subtypes, in contrast to DPM and APL neurons (Fig. 3). Previous studies using localization

microscopy showed that the molecular density of synaptic proteins at AZs can be

dynamically adjusted in motor neurons, potentially modulating the synaptic vesicle release

[41,42,48]. The variable Brp concentrations (i.e. the ratio of the Brp::rGFP intensity to the AZ

size) in DPM and APL neurons may represent AZs with distinct Brp densities and therefore

different functional modes (Fig. 3) [41]. Notably, a recent study successfully predicted the

neurotransmitter type of neurons using the ultrastructure of dense projections as one of the

parameters [49], suggesting neurotransmitter-specific AZ structures. Interestingly, both the

DPM and APL neurons are reported to release multiple neurotransmitters [27,31,50–53]. Distinct

Brp concentrations at AZs might thus support particular release modes or neurotransmitter

types.

In KCs, the spatial analysis of local AZ configuration revealed strong correlations

between Brp::rGFP intensities at neighboring AZs as well as AZ "hot spots", regions with

high Brp localization and AZ densities (Fig. 6). These results suggest previously

unidentified, highly localized sub-compartmental AZ structures, which are likely organized

across individual KCs. Presynaptic terminals of a single KC are found to be highly

heterogeneous in terms of their plasticity upon dopamine modulation and associative

learning [44,45]. A recent study found the sub-compartmentally heterogeneous activities of

dopamine terminals explain memory specificity [24]. Such locally distinct dopamine input

may be a source of the AZ hot spot formation in KC terminals. This model is consistent

with the localization of two opposing dopamine receptors Dop1R1 and Dop2R to AZs [54].

Furthermore, we showed that the local configuration undergoes structural plasticity upon

associative learning that involves presynaptic dopaminergic modulation [55–60] (Fig. 7). Since

cAMP signaling plays a crucial role in the AZ structural plasticity especially during

associative memory formation [39,48,61], it may underlie learning-induced plasticity in sub-

compartmental AZ configuration [17,18].

Materials and Methods

Animals

Flies were maintained on standard cornmeal food at 25 °C under a 12:12 h light-dark cycle

for all experiments. All flies used for experiments are 3-7 days old adult males. Flies were

transferred to fresh food vials after hatching and flipped every two days before

experiments. The GAL4-UAS system was used to express transgenes, and balancers were

removed for animals used in all experiments. Fly strains and resources used in this study

are as follow: *MB008B-GAL4* (BDSC 68291) [23], *MB009B-GAL4* (BDSC 68292) [23];

MB370B-GAL4 (BDSC 68319) [23], *MB504B-GAL4* (BDSC 68329) [23] , *VT64246-GAL4*

(VDRC 204311) [62], *GH146-GAL4* (BDSC 30026) [63], *R58E02-GAL4* (BDSC 41347) [64],

R13F02-GAL4 (BDSC 48571) [64] , *UAS-CD4::tdTomato* (BDSC 35841, by Dr. Yhu Nung

Jan & Chun Han), [59]*brp::GFP$_{11}$* [39], *UAS-GFP$_{1-10}$* [35].

Single-odor aversive conditioning

Adult flies at 3-7 days after eclosion were kept in a 25 °C incubator before and after

conditioning. Approximately 50 flies were put in a training tube that is set inside the

climate box for 2 minutes before the conditioning. Temperature in the climate box was set

at 24 °C and the relative humidity was set to 70%. For the paired group, 2% 4-

methycyclohexanol (4-MCH, Sigma-Aldrich, diluted in paraffin oil, Sigma-Aldrich) was

presented in a cup with the diameter of 5 mm, together with 12 pulses of electric shocks at

90 V for 1 minute inside the training tube. The unpaired group received the same 1-minute

electric shock first, followed by 1-minute rest, and then the same 1-minute 4-MCH

presentation. Flies were dissected at 3 min, 20 min, 90 min, 270 min and 1 day after

conditioning. Experiments were performed multiple times on different days for a single

group.

Sample preparation

Animals used in experiments involving comparisons between control and experimental

groups were dissected on the same day. Data were collected from multiple batches of

experiments performed on different days. Flies were anesthetized by ice and placed on ice

before dissection. The dissection was performed in cold PBS to collect the brain without

the ventral nerve cord. The brains were subsequently fixed by 2% paraformaldehyde for 1 h

and then washed 3×20 min by 0.1% PBT (0.1% Triton X-100 in PBS) in a PCR tube,

typically 5-6 brains in one tube. Samples were mounted on microscope slides using

mounting medium SeeDB2S [65].

Confocal microscopy image acquisition

The image acquisition was performed using an Olympus FV1200 confocal microscope

platform equipped with both PMT and GaAsP high sensitivity detectors and a 60×/1.42 NA

oil immersion objective (PLAPON60XO, Olympus). The Kalman filter was turned on to 2×

averaging while scanning. For Brp::rGFP samples in all GAL4 types, scanning speed was

set at 2 μs/pixel with a voxel size of lateral 0.079 × 0.079 × axial 0.370 μm. For

experimental point spread function (PSF) imaging, SeeDB2S immersed beads were

scanned in a setting that is 60×/1.42 NA oil immersion objective; 473 nm laser power:

2.0%; 559 nm laser power: 1.5%; voxel size: 0.079 μm × 0.079 μm × 0.370 μm; scanning

speed: 4 μs/pixel, to produce multiple bead images.

PSF acquisition and image deconvolution

Sub-resolution fluorescent beads (Tetraspeck Microspheres 0.1 μm, Thermo Fisher

Scientific, T7279) were imaged to generate the PSF. The bead solution was diluted with

distilled water and sonicated several times to eliminate aggregation. PSF images acquired

were processed using Amira software (Thermo Fisher Scientific) with the Extract Point

Spread Function module. The PSFs extracted from each image were averaged to create a

single PSF for later image deconvolution (Resizing voxel size: 0.079 μm × 0.079 μm ×

0.370 μm; 32 pixels × 32 pixels × 21 slices of image size). Image deconvolution was

performed by using the Richardson-Lucy iterative non-blind algorithm in a Fiji plugin

DeconvolutionLab2 [66], or by CLIJ2 GPU-based Richardson-Lucy deconvolution [67]. Images

were deconvolved 20 times iteratively to improve image quality.

Brp clusters segmentation and data analysis

For Brp::rGFP cluster detection, the image stack size was adjusted to double the XY plate

pixel number using Fiji to increase the detection precision. The images were subsequently

processed using the 3D maxima finder and the 3D spot segmentation function of the 3D

suite package in Fiji. 3D maxima were identified for each cluster and used as the starting

point for pixel clustering [36]. An intensity threshold (threshold) and the noise tolerance were

set to reduce noise and adjust the sensitivity of 3D maxima detection respectively. The

threshold set the value of all pixels that have values below the threshold to NaN (not a

number). The noise tolerance is set in the 3D maxima finder. To be noticed, the threshold is

only applied to the image used to find 3D maxima. The 3D maxima finder generates the 3D

maxima. The 3D segmentation function then requires the 3D maxima and the raw image, so

that there is no intensity cut affecting the measurement of cluster volume (used

deconvolved image). The local thresholding method for the 3D segmentation was set to

Gaussian fit, with the "SD value" set to 2. Cluster volume was measured as the voxel count

within the 3D ROI.

The above procedures may include a tiling and stitching step for images and data

because of the limited computer power. Typically, an image was tiled into 25 sub-stacks

with overlapped margins. After being processed separately, the data acquired from all tiled

sub-stacks was stitched back with a deletion of overlapping data points to generate the

complete detection result. Nearest neighbors were identified for each particle and spatial

distances in between (NND) were calculated using Python. For each sample, mean NND

was calculated separately. For a particular Brp::rGFP cluster, the number of surrounding

clusters within r ($r = 3 \times$ mean NND) spherically including itself is referred as the "AZ

density". The "local intensity" of a particular Brp::rGFP cluster is the average of Brp::rGFP

intensity of all clusters surrounding within r ($r = 3 \times$ mean NND) including itself. The plot

function of Python was used to plot the color-coded 3D reconstructed maps.

F-score calculation

The 3D detection parameters were optimized by matching the 3D maxima detection results

to the ground truths. Ground truths were defined by annotating AZs manually, according to

the size, position and signal pattern of Brp puncta in the images. The F-score was used to

assess detection precision. F-score is the harmonic mean of a system's precision and recall

value, calculated by the following formula: $2 \times [(\text{Precision} \times \text{Recall}) / (\text{Precision} + \text{Recall})]$.

The precision is the number of true positives divided by the total number of identified

positives, including incorrect identification. The recall is the number of true positives

divided by the total number of true data points defined as the ground truth. F-scores shown

in Fig. 1 were automatically calculated using a customized Fiji Marco that aligned 3D

maxima with ground truth. Different combinations of noise threshold and noise tolerance

were tested for the test images cropped from raw images.

Statistical analysis

Statistical analysis was performed using GraphPad Prism version 8, 9 and 10 or Python.

For statistical comparisons, Kruskal-Wallis test with original False Discovery Rate method

of Benjamini and Hochberg correction was applied expect as explicitly mentioned. A

desired false discovery rate of 0.05. Pearson's correlation coefficient (R) was calculated

using Python.

References

1. Kittel, R.J., Wichmann, C., Rasse, T.M., Fouquet, W., Schmidt, M., Schmid, A., Wagh,
D.A., Pawlu, C., Kellner, R.R., Willig, K.I., et al. (2006). Bruchpilot Promotes Active Zone

Assembly, Ca^{2+} Channel Clustering, and Vesicle Release. Science *312*, 1051–1054. https://doi.org/10.1126/science.1126308.

2. Wagh, D.A., Rasse, T.M., Asan, E., Hofbauer, A., Schwenkert, I., Dürrbeck, H., Buchner, S., Dabauvalle, M.-C., Schmidt, M., Qin, G., et al. (2006). Bruchpilot, a Protein with Homology to ELKS/CAST, Is Required for Structural Integrity and Function of Synaptic Active Zones in Drosophila. Neuron *49*, 833–844. https://doi.org/10.1016/j.neuron.2006.02.008.

3. Fouquet, W., Owald, D., Wichmann, C., Mertel, S., Depner, H., Dyba, M., Hallermann, S., Kittel, R.J., Eimer, S., and Sigrist, S.J. (2009). Maturation of active zone assembly by Drosophila Bruchpilot. J Cell Biol *186*, 129–145. https://doi.org/10.1083/jcb.200812150.

4. Matkovic, T., Siebert, M., Knoche, E., Depner, H., Mertel, S., Owald, D., Schmidt, M., Thomas, U., Sickmann, A., Kamin, D., et al. (2013). The Bruchpilot cytomatrix determines the size of the readily releasable pool of synaptic vesicles. J Cell Biol *202*, 667–683. https://doi.org/10.1083/jcb.201301072.

5. Paul, M.M., Pauli, M., Ehmann, N., Hallermann, S., Sauer, M., Kittel, R.J., and Heckmann, M. (2015). Bruchpilot and Synaptotagmin collaborate to drive rapid glutamate release and active zone differentiation. Front Cell Neurosci *9*, 29. https://doi.org/10.3389/fncel.2015.00029.

6. Akbergenova, Y., Cunningham, K.L., Zhang, Y.V., Weiss, S., and Littleton, J.T. (2018). Characterization of developmental and molecular factors underlying release heterogeneity at Drosophila synapses. Elife *7*, e38268. https://doi.org/10.7554/elife.38268.

7. Newman, Z.L., Bakshinskaya, D., Schultz, R., Kenny, S.J., Moon, S., Aghi, K., Stanley, C., Marnani, N., Li, R., Bleier, J., et al. (2022). Determinants of synapse diversity revealed by super-resolution quantal transmission and active zone imaging. Nat Commun *13*, 229. https://doi.org/10.1038/s41467-021-27815-2.

8. Zheng, Z., Lauritzen, J.S., Perlman, E., Robinson, C.G., Nichols, M., Milkie, D., Torrens, O., Price, J., Fisher, C.B., Sharifi, N., et al. (2018). A Complete Electron Microscopy Volume of the Brain of Adult Drosophila melanogaster. Cell *174*, 730-743.e22. https://doi.org/10.1016/j.cell.2018.06.019.

9. Scheffer, L.K., Xu, C.S., Januszewski, M., Lu, Z., Takemura, S., Hayworth, K.J., Huang, G.B., Shinomiya, K., Maitlin-Shepard, J., Berg, S., et al. (2020). A connectome and analysis of the adult Drosophila central brain. Elife *9*, e57443. https://doi.org/10.7554/elife.57443.

10. Takemura, S., Hayworth, K.J., Huang, G.B., Januszewski, M., Lu, Z., Marin, E.C.,
Preibisch, S., Xu, C.S., Bogovic, J., Champion, A.S., et al. (2024). A Connectome of the
Male Drosophila Ventral Nerve Cord. https://doi.org/10.7554/elife.97769.1.

11. Schlegel, P., Yin, Y., Bates, A.S., Dorkenwald, S., Eichler, K., Brooks, P., Han, D.S.,
Gkantia, M., Santos, M. dos, Munnelly, E.J., et al. (2024). Whole-brain annotation and
multi-connectome cell typing of Drosophila. Nature *634*, 139–152.
https://doi.org/10.1038/s41586-024-07686-5.

12. Dorkenwald, S., Matsliah, A., Sterling, A.R., Schlegel, P., Yu, S., McKellar, C.E., Lin,
A., Costa, M., Eichler, K., Yin, Y., et al. (2024). Neuronal wiring diagram of an adult brain.
Nature *634*, 124–138. https://doi.org/10.1038/s41586-024-07558-y.

13. Sachse, S., Rueckert, E., Keller, A., Okada, R., Tanaka, N.K., Ito, K., and Vosshall,
L.B. (2007). Activity-Dependent Plasticity in an Olfactory Circuit. Neuron *56*, 838–850.
https://doi.org/10.1016/j.neuron.2007.10.035.

14. Gilestro, G.F., Tononi, G., and Cirelli, C. (2009). Widespread Changes in Synaptic
Markers as a Function of Sleep and Wakefulness in Drosophila. Science *324*, 109–112.
https://doi.org/10.1126/science.1166673.

15. Fernández, M.P., Berni, J., and Ceriani, M.F. (2008). Circadian Remodeling of
Neuronal Circuits Involved in Rhythmic Behavior. PLoS Biol. *6*, e69.
https://doi.org/10.1371/journal.pbio.0060069.

16. Kremer, M.C., Christiansen, F., Leiss, F., Paehler, M., Knapek, S., Andlauer, T.F.M.,
Förstner, F., Kloppenburg, P., Sigrist, S.J., and Tavosanis, G. (2010). Structural Long-Term
Changes at Mushroom Body Input Synapses. Curr Biol *20*, 1938–1944.
https://doi.org/10.1016/j.cub.2010.09.060.

17. Zhang, X., Li, Q., Wang, L., Liu, Z.-J., and Zhong, Y. (2018). Active Protection:
Learning-Activated Raf/MAPK Activity Protects Labile Memory from Rac1-Independent
Forgetting. Neuron *98*, 142-155.e4. https://doi.org/10.1016/j.neuron.2018.02.025.

18. Turrel, O., Ramesh, N., Escher, M.J.F., Pooryasin, A., and Sigrist, S.J. (2022).
Transient active zone remodeling in the Drosophila mushroom body supports memory.
Curr Biol. https://doi.org/10.1016/j.cub.2022.10.017.

19. Mosca, T.J., and Luo, L. (2014). Synaptic organization of the Drosophila antennal lobe
and its regulation by the Teneurins. eLife *3*, e03726. https://doi.org/10.7554/elife.03726.

20. Gao, R., Asano, S.M., Upadhyayula, S., Pisarev, I., Milkie, D.E., Liu, T.-L., Singh, V.,
Graves, A., Huynh, G.H., Zhao, Y., et al. (2019). Cortical column and whole-brain imaging

with molecular contrast and nanoscale resolution. Science *363*.
https://doi.org/10.1126/science.aau8302.

21. Chen, Y., Akin, O., Nern, A., Tsui, C.Y.K., Pecot, M.Y., and Zipursky, S.L. (2014).
Cell-type-Specific Labeling of Synapses In Vivo through Synaptic Tagging with
Recombination. Neuron *81*, 280–293. https://doi.org/10.1016/j.neuron.2013.12.021.

22. Davis, R.L. (2023). Learning and memory using Drosophila melanogaster : a focus on
advances made in the fifth decade of research. GENETICS.
https://doi.org/10.1093/genetics/iyad085.

23. Aso, Y., Hattori, D., Yu, Y., Johnston, R.M., Iyer, N.A., Ngo, T.-T., Dionne, H.,
Abbott, L., Axel, R., Tanimoto, H., et al. (2014). The neuronal architecture of the
mushroom body provides a logic for associative learning. Elife *3*, e04577.
https://doi.org/10.7554/elife.04577.

24. Yamagata, N., Ezaki, T., Takahashi, T., Wu, H., and Tanimoto, H. (2021). Presynaptic
inhibition of dopamine neurons controls optimistic bias. Elife *10*, e64907.
https://doi.org/10.7554/elife.64907.

25. Pitman, J.L., Huetteroth, W., Burke, C.J., Krashes, M.J., Lai, S.-L., Lee, T., and
Waddell, S. (2011). A Pair of Inhibitory Neurons Are Required to Sustain Labile Memory
in the Drosophila Mushroom Body. Curr Biol *21*, 855–861.
https://doi.org/10.1016/j.cub.2011.03.069.

26. Lin, A.C., Bygrave, A.M., Calignon, A. de, Lee, T., and Miesenböck, G. (2014).
Sparse, decorrelated odor coding in the mushroom body enhances learned odor
discrimination. Nat Neurosci *17*, 559–568. https://doi.org/10.1038/nn.3660.

27. Liu, X., and Davis, R.L. (2009). The GABAergic anterior paired lateral neuron
suppresses and is suppressed by olfactory learning. Nat. Neurosci. *12*, 53–59.
https://doi.org/10.1038/nn.2235.

28. Yu, D., Keene, A.C., Srivatsan, A., Waddell, S., and Davis, R.L. (2005). Drosophila
DPM Neurons Form a Delayed and Branch-Specific Memory Trace after Olfactory
Classical Conditioning. Cell *123*, 945–957. https://doi.org/10.1016/j.cell.2005.09.037.

29. Keene, A.C., Stratmann, M., Keller, A., Perrat, P.N., Vosshall, L.B., and Waddell, S.
(2004). Diverse Odor-Conditioned Memories Require Uniquely Timed Dorsal Paired
Medial Neuron Output. Neuron *44*, 521–533. https://doi.org/10.1016/j.neuron.2004.10.006.

30. Keene, A.C., Krashes, M.J., Leung, B., Bernard, J.A., and Waddell, S. (2006).
Drosophila Dorsal Paired Medial Neurons Provide a General Mechanism for Memory
Consolidation. Curr Biol *16*, 1524–1530. https://doi.org/10.1016/j.cub.2006.06.022.

31. Haynes, P.R., Christmann, B.L., and Griffith, L.C. (2015). A single pair of neurons
links sleep to memory consolidation in Drosophila melanogaster. eLife *4*, e03868.
https://doi.org/10.7554/elife.03868.

32. Waddell, S., Armstrong, J.D., Kitamoto, T., Kaiser, K., and Quinn, W.G. (2000). The
amnesiac Gene Product Is Expressed in Two Neurons in the Drosophila Brain that Are
Critical for Memory. Cell *103*, 805–813. https://doi.org/10.1016/s0092-8674(00)00183-5.

33. Huang, S., Piao, C., Beuschel, C.B., Götz, T., and Sigrist, S.J. (2020). Presynaptic
Active Zone Plasticity Encodes Sleep Need in Drosophila. Curr Biol *30*, 1077-1091.e5.
https://doi.org/10.1016/j.cub.2020.01.019.

34. Kamiyama, D., Sekine, S., Barsi-Rhyne, B., Hu, J., Chen, B., Gilbert, L.A., Ishikawa,
H., Leonetti, M.D., Marshall, W.F., Weissman, J.S., et al. (2016). Versatile protein tagging
in cells with split fluorescent protein. Nat. Commun. *7*, 11046.
https://doi.org/10.1038/ncomms11046.

35. Kondo, S., Takahashi, T., Yamagata, N., Imanishi, Y., Katow, H., Hiramatsu, S., Lynn,
K., Abe, A., Kumaraswamy, A., and Tanimoto, H. (2020). Neurochemical Organization of
the Drosophila Brain Visualized by Endogenously Tagged Neurotransmitter Receptors.
Cell Reports *30*, 284-297.e5. https://doi.org/10.1016/j.celrep.2019.12.018.

36. Ollion, J., Cochennec, J., Loll, F., Escudé, C., and Boudier, T. (2013). TANGO: a
generic tool for high-throughput 3D image analysis for studying nuclear organization.
Bioinformatics *29*, 1840–1841. https://doi.org/10.1093/bioinformatics/btt276.

37. Ehmann, N., Linde, S. van de, Alon, A., Ljaschenko, D., Keung, X.Z., Holm, T., Rings,
A., DiAntonio, A., Hallermann, S., Ashery, U., et al. (2014). Quantitative super-resolution
imaging of Bruchpilot distinguishes active zone states. Nat Commun *5*, 4650.
https://doi.org/10.1038/ncomms5650.

38. Gratz, S.J., Goel, P., Bruckner, J.J., Hernandez, R.X., Khateeb, K., Macleod, G.T.,
Dickman, D., and O'Connor-Giles, K.M. (2019). Endogenous tagging reveals differential
regulation of Ca2+ channels at single AZs during presynaptic homeostatic potentiation and
depression. J Neurosci *39*, 3068–18. https://doi.org/10.1523/jneurosci.3068-18.2019.

39. Wu, H., Eno, S., Jinnai, K., Maekawa, Y., Saito, K., Abe, A., Williams, D.W.,
Yamagata, N., Kondo, S., and Tanimoto, H. (2024). Octopamine signals coordinate the

spatial pattern of presynaptic machineries within individual Drosophila mushroom body neurons. bioRxiv, 2024.12.02.626512. https://doi.org/10.1101/2024.12.02.626512.

40. Schlegel, P., Yin, Y., Bates, A.S., Dorkenwald, S., Eichler, K., Brooks, P., Han, D.S., Gkantia, M., Santos, M. dos, Munnelly, E.J., et al. (2023). A consensus cell type atlas from multiple connectomes reveals principles of circuit stereotypy and variation. bioRxiv, 2023.06.27.546055. https://doi.org/10.1101/2023.06.27.546055.

41. Mrestani, A., Pauli, M., Kollmannsberger, P., Repp, F., Kittel, R.J., Eilers, J., Doose, S., Sauer, M., Sirén, A.-L., Heckmann, M., et al. (2021). Active zone compaction correlates with presynaptic homeostatic potentiation. Cell Rep. *37*, 109770. https://doi.org/10.1016/j.celrep.2021.109770.

42. Ghelani, T., Escher, M., Thomas, U., Esch, K., Lützkendorf, J., Depner, H., Maglione, M., Parutto, P., Gratz, S., Matkovic-Rachid, T., et al. (2023). Interactive nanocluster compaction of the ELKS scaffold and Cacophony Ca2+ channels drives sustained active zone potentiation. Sci. Adv. *9*, eade7804. https://doi.org/10.1126/sciadv.ade7804.

43. Vogt, K., Schnaitmann, C., Dylla, K.V., Knapek, S., Aso, Y., Rubin, G.M., and Tanimoto, H. (2014). Shared mushroom body circuits underlie visual and olfactory memories in Drosophila. Elife *3*, e02395. https://doi.org/10.7554/elife.02395.

44. Davidson, A.M., Kaushik, S., and Hige, T. (2023). Dopamine-Dependent Plasticity Is Heterogeneously Expressed by Presynaptic Calcium Activity across Individual Boutons of the Drosophila Mushroom Body. eNeuro *10*, ENEURO.0275-23.2023. https://doi.org/10.1523/eneuro.0275-23.2023.

45. Bilz, F., Geurten, B.R.H., Hancock, C.E., Widmann, A., and Fiala, A. (2020). Visualization of a Distributed Synaptic Memory Code in the Drosophila Brain. Neuron *106*, 963-976.e4. https://doi.org/10.1016/j.neuron.2020.03.010.

46. Perisse, E., Yin, Y., Lin, A.C., Lin, S., Huetteroth, W., and Waddell, S. (2013). Different Kenyon Cell Populations Drive Learned Approach and Avoidance in Drosophila. Neuron *79*, 945–956. https://doi.org/10.1016/j.neuron.2013.07.045.

47. Amin, H., Apostolopoulou, A.A., Suárez-Grimalt, R., Vrontou, E., and Lin, A.C. (2020). Localized inhibition in the Drosophila mushroom body. Elife *9*, e56954. https://doi.org/10.7554/elife.56954.

48. Sachidanandan, D., Aravamudhan, A., Mrestani, A., Nerlich, J., Lamberty, M., Hasenauer, N., Ehmann, N., Pauls, D., Seubert, T., Maiellaro, I., et al. (2023). Rab3 mediates cyclic AMP-dependent presynaptic plasticity and olfactory learning. bioRxiv, 2023.12.21.572589. https://doi.org/10.1101/2023.12.21.572589.

49. Eckstein, N., Bates, A.S., Champion, A., Du, M., Yin, Y., Schlegel, P., Lu, A.K.-Y., Rymer, T., Finley-May, S., Paterson, T., et al. (2024). Neurotransmitter classification from electron microscopy images at synaptic sites in Drosophila melanogaster. Cell *187*, 2574-2594.e23. https://doi.org/10.1016/j.cell.2024.03.016.

50. Wu, C.-L., Shih, M.-F.M., Lee, P.-T., and Chiang, A.-S. (2013). An Octopamine-Mushroom Body Circuit Modulates the Formation of Anesthesia-Resistant Memory in Drosophila. Curr Biol *23*, 2346–2354. https://doi.org/10.1016/j.cub.2013.09.056.

51. Lee, P.-T., Lin, H.-W., Chang, Y.-H., Fu, T.-F., Dubnau, J., Hirsh, J., Lee, T., and Chiang, A.-S. (2011). Serotonin–mushroom body circuit modulating the formation of anesthesia-resistant memory in Drosophila. Proc. Natl. Acad. Sci. *108*, 13794–13799. https://doi.org/10.1073/pnas.1019483108.

52. Davie, K., Janssens, J., Koldere, D., Waegeneer, M.D., Pech, U., Kreft, Ł., Aibar, S., Makhzami, S., Christiaens, V., González-Blas, C.B., et al. (2018). A Single-Cell Transcriptome Atlas of the Aging Drosophila Brain. Cell *174*, 982-998.e20. https://doi.org/10.1016/j.cell.2018.05.057.

53. Zeng, J., Li, X., Zhang, R., Lv, M., Wang, Y., Tan, K., Xia, X., Wan, J., Jing, M., Zhang, X., et al. (2023). Local 5-HT signaling bi-directionally regulates the coincidence time window for associative learning. Neuron *111*, 1118-1135.e5. https://doi.org/10.1016/j.neuron.2022.12.034.

54. Hiramatsu, S., Saito, K., Kondo, S., Katow, H., Yamagata, N., Wu, C.-F., and Tanimoto, H. (2024). Cell-type-specific fluorescent tagging of endogenous target proteins reveals synaptic enrichment and dynamic regulations of dopamine receptors. https://doi.org/10.7554/elife.98358.1.

55. Aso, Y., Siwanowicz, I., Bräcker, L., Ito, K., Kitamoto, T., and Tanimoto, H. (2010). Specific Dopaminergic Neurons for the Formation of Labile Aversive Memory. Curr Biol *20*, 1445–1451. https://doi.org/10.1016/j.cub.2010.06.048.

56. Liu, C., Plaçais, P.-Y., Yamagata, N., Pfeiffer, B.D., Aso, Y., Friedrich, A.B., Siwanowicz, I., Rubin, G.M., Preat, T., and Tanimoto, H. (2012). A subset of dopamine neurons signals reward for odour memory in Drosophila. Nature *488*, 512–516. https://doi.org/10.1038/nature11304.

57. Aso, Y., and Rubin, G.M. (2016). Dopaminergic neurons write and update memories with cell-type-specific rules. Elife *5*, e16135. https://doi.org/10.7554/elife.16135.

58. Yamagata, N., Ichinose, T., Aso, Y., Plaçais, P.-Y., Friedrich, A.B., Sima, R.J., Preat, T., Rubin, G.M., and Tanimoto, H. (2015). Distinct dopamine neurons mediate reward

signals for short- and long-term memories. Proc National Acad Sci *112*, 578–583. https://doi.org/10.1073/pnas.1421930112.

59. Aso, Y., Herb, A., Ogueta, M., Siwanowicz, I., Templier, T., Friedrich, A.B., Ito, K., Scholz, H., and Tanimoto, H. (2012). Three Dopamine Pathways Induce Aversive Odor Memories with Different Stability. PLoS Genet. *8*, e1002768. https://doi.org/10.1371/journal.pgen.1002768.

60. Burke, C.J., Huetteroth, W., Owald, D., Perisse, E., Krashes, M.J., Das, G., Gohl, D., Silies, M., Certel, S., and Waddell, S. (2012). Layered reward signalling through octopamine and dopamine in Drosophila. Nature *492*, 433–437. https://doi.org/10.1038/nature11614.

61. Baltruschat, L., Prisco, L., Ranft, P., Lauritzen, J.S., Fiala, A., Bock, D.D., and Tavosanis, G. (2021). Circuit reorganization in the Drosophila mushroom body calyx accompanies memory consolidation. Cell Reports *34*, 108871. https://doi.org/10.1016/j.celrep.2021.108871.

62. Tirian, L., and Dickson, B.J. (2017). The VT GAL4, LexA, and split-GAL4 driver line collections for targeted expression in the Drosophila nervous system. bioRxiv, 198648. https://doi.org/10.1101/198648.

63. Stocker, R.F., Heimbeck, G., Gendre, N., and Belle, J.S. de (1997). Neuroblast ablation in Drosophila P[GAL4] lines reveals origins of olfactory interneurons. J. Neurobiol. *32*, 443–456. https://doi.org/10.1002/(sici)1097-4695(199705)32:5<443::aid-neu1>3.0.co;2-5.

64. Pfeiffer, B.D., Jenett, A., Hammonds, A.S., Ngo, T.-T.B., Misra, S., Murphy, C., Scully, A., Carlson, J.W., Wan, K.H., Laverty, T.R., et al. (2008). Tools for neuroanatomy and neurogenetics in Drosophila. Proc. Natl. Acad. Sci. *105*, 9715–9720. https://doi.org/10.1073/pnas.0803697105.

65. Ke, M.-T., Nakai, Y., Fujimoto, S., Takayama, R., Yoshida, S., Kitajima, T.S., Sato, M., and Imai, T. (2016). Super-Resolution Mapping of Neuronal Circuitry With an Index-Optimized Clearing Agent. Cell Reports *14*, 2718–2732. https://doi.org/10.1016/j.celrep.2016.02.057.

66. Sage, D., Donati, L., Soulez, F., Fortun, D., Schmit, G., Seitz, A., Guiet, R., Vonesch, C., and Unser, M. (2017). DeconvolutionLab2: An open-source software for deconvolution microscopy. Methods *115*, 28–41. https://doi.org/10.1016/j.ymeth.2016.12.015.

67. Haase, R., Royer, L.A., Steinbach, P., Schmidt, D., Dibrov, A., Schmidt, U., Weigert, M., Maghelli, N., Tomancak, P., Jug, F., et al. (2020). CLIJ: GPU-accelerated image

processing for everyone. Nat. Methods *17*, 5–6. https://doi.org/10.1038/s41592-019-0650-
1.

Acknowledgments

We thank all lab members of Tanimoto Lab at Tohoku University for valuable discussions.

We thank Dr. Yuki Tsukada at Keio University for crucial advice on Fiji-based image

analysis.

We thank Vienna Drosophila Resource Center and Bloomington Drosophila Stock Center

for transgenic flies.

Funding:

Ministry of Education, Culture, Sports, Science and Technology 22H05481 (HT)

Ministry of Education, Culture, Sports, Science and Technology 22KK0106 (HT)

Ministry of Education, Culture, Sports, Science and Technology 20H00519 (HT)

Japan Science and Technology Agency JPMJSP2114 (HW)

Tohoku University Research Program 'Frontier Research in Duo' (HT).

Author contributions:

Conceptualization: HW, HT

Methodology: HW, YM, NY, SE, SK

Investigation: HW, YM, SE

Visualization: HW, SE

Resources: SK

Supervision: HW, NY, HT

Writing—original draft: HY, HT

Writing—review & editing: HW, SE, HT

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

Further information and requests on resources and reagents should be directed to and will

be fulfilled by the corresponding author, Dr. Hiromu Tanimoto (hiromut@m.tohoku.ac.jp).

This study did not generate any unique code. All data needed to evaluate the conclusions in

the paper are present in the paper.

Figures and Tables

Fig. 1. Spatial characterization of Brp::rGFP clusters at single-AZ level.

(**A**) Schematic of the split-GFP tagging strategy. The GFP₁₁ fragment is inserted by

CRISPR/Cas9 just prior to the stop codon of *brp*, while GFP1-10 is expressed in specific

cell types via GAL4-UAS system.

(**B**) Brp::rGFP signals in DPM. The entire MB lobe structure is shown. The black box in

the left panel indicates the zoom-in area (right panel). GFP1-10 and CD4::tdTomato

(magenta) were co-expressed using *VT64246-GAL4*. Scale bar, 20 μm (left); 5 μm (right).

(**C**) Image processing pipeline. Raw images were processed with image deconvolution to

improve the signal quality. 3D maxima (green pixels) are detected for each Brp::rGFP

cluster. Pixels around the maxima were clustered to created 3D ROIs, indicated by yellow

circles. The "AZ density" is the number of Brp::rGFP clusters within r (3 × mean NND).

Scale bar, 1 μm.

(**D**) Heatmaps of F-scores showing the detection performance across in different cell types.

An intensity threshold (threshold) is applied to reduce the background noise when detecting

3D maxima. The noise tolerance is the parameter that adjusts the sensitivity of 3D maxima

detection. The color represents the F-score value.

(**E**) Number of Brp::rGFP clusters detected in DPM and APL. The dashed lines indicate the

number of electron dense projections identified in the hemibrain connectome for DPM

(17530) and APL (10789) on the MB lobes and peduncle (cite the resource here). For both

DPM and APL, n = 5. Lower panels show the representative 3D reconstructions of

Brp::rGFP clusters in DPM and APL. Scale bars, 20 μm. Data were presented as mean ±

s.e.m.

Fig. 2. Compartmentalized AZ structures of KCs.

(**A**) Histograms of the signal intensity of individual Brp::rGFP clusters from three KC

subtypes. Values are normalized by the mean intensity in each data set (each sample). Each

line represents the histogram of one independent sample. γ KCs (*MB009B-GAL4*), n = 6;

α/β KCs (*MB370B-GAL4*), n = 5; α'/β' KCs (*MB008B-GAL4*), n = 5.

(**B**) 3D reconstructions of Brp::rGFP clusters in three KC subtypes, colored by the

Brp::rGFP intensity. The approximate locations of compartments are as indicated. Min. and

Max. were set to represent the lowest and highest 5% of Brp::rGFP intensity value in the

dataset respectively. Scale bars, 20 μm.

664   (C) Signal intensity of Brp:rGFP clusters in each compartment. Brp::rGFP puncta were

665   quantified compartmentally. Medians in different compartments are showed as the ratio

666   against the average of five compartments in the corresponding KC subtype. Each line

667   represents one independent sample.

Fig. 3. The variability of Brp::rGFP concentrations depends on cell type.

(**A**) Brp::rGFP clusters with distinct intensities but similar size in DPM. The left panel

shows a cropped image of Brp::rGFP in DPM. White boxes indicate areas zoomed-in in

panels 1 and 2. Transparent white lines in panels 1 and 2 show the lines on which the

intensity profiles were plotted. Intensity profiles were plotted for AZ1 (magenta) and AZ2

(blue) respectively.

(**B-D**) Scatter plots showing the correlation between the Brp::rGFP intensity and cluster

volume in DPM, APL and α/β KCs. Data of three representative samples were shown.

Pearson correlation coefficient R were calculated for each sample.

(**E**) Correlations between the intensity and volume of Brp::rGFP clusters in different cell

types. DPM (n = 6), APL (n = 5), γ KCs (n = 6, *MB009B-GAL4*), α'/β' KCs (n = 5,

MB370B-GAL4), α/β KCs (n = 5, *MB008B-GAL4*). DPM vs. α'/β' KCs: P = 0.0253; DPM

681   vs. α/β KCs: P = 0.0009; APL vs. α/β KCs: P = 0.0035; γ KCs vs. α/β KCs: P = 0.0418.

682   Values marked with different lowercase letters represent significant difference (P < 0.05);

683   Data were presented as box plots showing center (median), whiskers (Min. to Max.).

Fig. 4. Stereotyped AZ distribution of PPL1-α3 DAN.

(**A**) Brp::rGFP in PPL1 DANs. UAS-GFP$_{1-10}$ was expressed using *MB504B-GAL4*. Dashed line marks the rough boundary between α3 and α2α'2 compartments. Scale bar, 20 μm.

(**B**) Brp::rGFP intensity profiles of α3 and α2α'2 compartments. Left panels show the max-projection images of α3 and α2α'2 optical coronal sections. Transparent gray stripes indicate areas where intensity profiles are plotted in the right panels. Scale bars, 10 μm.

(**C**) 3D reconstruction colored by the AZ density in PPL-1 DANs. Color scale: Min. = 0, Max. = 40. Dashed line indicates the rough boundary between α3 and α2α'2 compartments. Scale bar, 20 μm.

(**D**) Stereotyped AZ distribution pattern in PPL1-α3 across individuals. 3D reconstructions show the AZ density across different brain samples. Scale bar, 20 μm.

Fig. 5. Cell-type-specific stereotypy of AZ spatial distributions.

(**A**) 3D reconstructions of Brp::rGFP clusters in DPMs, colored by AZ density. Color scale:

Min. = 0, Max. = 35. Black arrows indicate consistently high AZ density regions across

brain samples. Scale bars, 20 μm.

(**B**) 3D reconstructions of Brp::rGFP clusters in APLs, colored by AZ density. Color scale:

Min. = 0, Max. = 30. Black dashed square indicates the area zoomed-in. APL

reconstructions are arranged from left to right according to the overall AZ density in α3.

Fig. 6. Local intensity analysis revealed sub-compartmental AZ structures.

(**A**) Correlation of Brp::rGFP intensities between nearest neighbor AZs in different cell types. Brp::rGFP intensities were log transformed. DPM vs. γ KCs: P = 0.0066; DPM vs. α'/β' KCs: P = 0.0027; DPM vs. α/β KCs: P = 0.0066; Scale bar: 1 μm.

(**B**) 3D reconstructions of Brp::rGFP clusters in a γ KC sample, colored by local intensity and AZ density. Black arrows indicate areas with both high local intensities and high AZ densities. Min. and Max. were set to represent the lowest and highest 5% in local intensity value respectively. Color scale of AZ density: Min. = 0 and Max = 35.

(**C**) 3D reconstructions of Brp::rGFP clusters in a DPM neuron, colored by local intensity and AZ density. Min. and Max. are set to represent the lowest and highest 5% in local intensity value respectively. Color scale of AZ density: Min. = 0, Max. = 35.

716　**(D)** Correlations between AZ density and local intensity in different cell types. DPM (n =

717　6), APL (n = 5), γ KCs (n = 6, *MB009B-GAL4*), α'/β' KCs (n = 5, *MB370B-GAL4*), α/β KCs

718　(n = 5, *MB008B-GAL4*). DPM vs. γ KCs: P = 0.0171; DPM vs. α'/β' KCs: P = 0.0171; APL

719　vs. γ KCs: P = 0.0014; APL vs. α'/β' KCs: P = 0.0014; APL vs. α/β KCs: P = 0.0171.

Fig. 7. Associative conditioning re-organizes sub-compartmental active zone clusters.

(A) Brp::rGFP (green) and CD4::tdTtomato cytoplasm membrane marker (magenta) in

KCs, visualized by using *R13F02-GAL4*.

(B) The experimental design of the aversive single odor conditioning. Files in the pair

group receive concurrent presentation of 2% 4-MCH and 90 V electric shock. Flies in the

unpair group first receive the electric shock first and then 4-MCH 1 min later. Flies were

dissected at different time points after the conditioning.

(C) Correlation coefficient (AZ density vs. local intensity of individual Brp::rGFP puncta)

of each compartment at 90 min after conditioning. Pair (n = 11) vs. Unpair (n = 12), γ2 (P =

0.0261), β2 (P = 0.0060), α1 (P = 0.0026), α'3 (P = 0.0026).

(D) Heatmap showing the difference of correlation coefficient (AZ density vs. local

intensity of Brp::rGFP puncta) between the pair and unpair group at different time points

after conditioning. The color indicates the difference and the asterisks in the compartments

indicate the significant difference. For 3 min, Pair (n = 12), Unpair (n = 11); For 20 min,

Pair (n = 12), Unpair (n = 11); For 90 min, Pair (n = 11) vs. Unpair (n = 12), P = 0.0247;

For 270 min, Pair (n = 11), Unpair (n = 11), ; For 1 D, Pair (n = 12), Unpair (n = 9);

For all results involved statistical comparison, only significant results are shown. Scale

bars, 20 μm. Mann-Whitney test with original False Discovery Rate (FDR) method of

Benjamini and Hochberg correction. *P<0.05. Data were presented as box plots showing

centre (median), whiskers (Min. to Max.).