Title

Profiling presynaptic scaffolds using split-GFP reconstitution reveals cell-type-specific

spatial configurations in the fly brain

Authors

Hongyang Wu[1], Yoh Maekawa[2], Sayaka Eno[1], Shu Kondo[3], Nobuhiro Yamagata[1,4],

Hiromu Tanimoto[1,2]*

Affiliations

[1] Graduate School of Life Sciences, Tohoku University, Sendai, Japan.

[2] Department of Biology, Faculty of Science, Tohoku University, Sendai, Japan.

[3] Department of Biological Science and Technology, Tokyo University of Science, Tokyo,

Japan.

[4] Department of Life Science, Graduate School of Engineering Science, Akita, Akita

University.

* Correspondence: Dr. Hiromu Tanimoto (hiromut@m.tohoku.ac.jp)

Abstract

Characterization of intracellular synapse heterogeneity aides to understand the intricate computational logic of neuronal circuits. Despite recent advances in connectomics, the spatial patterns of synapses and their inter-individual variability remain largely unknown. Using directed split-GFP reconstitution, we achieved visualization of endogenous Bruchpilot (Brp), a presynaptic active zone (AZ) scaffold protein, in a cell-type-specific manner. By developing a high-throughput quantification pipeline, we profiled AZ structures in identified neurons of the mushroom body circuit, where intracellular synaptic patterns are crucial due to compartmentalized connectivity. Quantitative characterization of the pattern of Brp clusters across multiple individuals revealed cell-type-dependent synapse heterogeneity and stereotypy. Furthermore, we discovered previously unidentified sub-compartmental synapse configuration and its transient structural plasticity triggered by associative learning. These profiles thus uncovered multi-layered spatial configurations of AZs, from stereotyped overall AZ distribution patterns, to local arrangements of neighboring synapses.

Keywords:

Intracellular synaptic configuration; synapse profiling; active zones; Bruchpilot; mushroom body circuit; split-GFP; synapse heterogeneity; synapse stereotypy; structural plasticity; associative learning

Introduction

The brain is a complex network of neurons, and understanding their wiring strategies provides crucial insights into their roles in neuronal computation. Release sites in the presynaptic terminal of *Drosophila* neurons are decorated with electron-dense projections composed of active-zone (AZ) scaffolds that are essential for efficient synaptic transmission (Akbergenova et al., 2018; Fouquet et al., 2009; Kittel et al., 2006; Matkovic et al., 2013; Newman et al., 2022; Paul et al., 2015; Wagh et al., 2006). These dense projections serve as a synapse marker enabling the whole-brain connectomes,

comprehensive mapping of synaptic connections at the electron microscopic level (Dorkenwald et al., 2024; Scheffer et al., 2020; Schlegel et al., 2024; Takemura et al., 2024; Zheng et al., 2018). While connectome-based approaches catalyzed the exploration of cell type distinctions, circuit motifs, and neurotransmitters, their application to comparisons across different conditions is still challenging due to the limited throughput of data acquisition and dense reconstruction. Since the synaptic structure and connectivity reflect substantial variability of development and experiences (Fernández et al., 2008; Gilestro et al., 2009; Kremer et al., 2010; Sachse et al., 2007; Schlegel et al., 2024; Turrel et al., 2022; Zhang et al., 2018), cross-individual comparisons are necessary. Fluorescence labeling offers high throughputs and contents, especially when comparing synaptic organizations across cell-types (Mosca and Luo, 2014) or brain regions (Gao et al., 2019). Nevertheless, characterizing endogenous synaptic proteins in specific cells using fluorescence-labeling remains challenging in the central nervous system (CNS). This has driven growing interest in designing cell-type-specific fluorescence-tagging strategies for endogenous synaptic protein (Chen et al., 2014).

The circuit of *Drosophila* mushroom bodies (MBs) plays a central role in olfactory associative learning (Davis, 2023). The major intrinsic MB neuron, known as Kenyon cells (KCs) respond to odor stimulation and synapse onto the spatially segregated dendrites of MB output neurons (MBONs) that divide the MB into compartments. Distinct types of dopaminergic neurons (DANs) synapse onto specific compartments and thereby modulate KC-MBON synapses in the corresponding compartments (Aso et al., 2014). Presynaptic calcium levels of these DANs undergo sub-compartmental GABAergic modulation and inform memory specificity, postulating the distinct synapse structures at individual release sites (Yamagata et al., 2021) . Two types of giant interneurons, the anterior paired lateral neuron (APL) and the dorsal paired medial (DPM) neuron extensively ramify across the MB lobes and provide recurrent modulations to KC synapses (Haynes et al., 2015; Keene et al., 2006, 2004; Lin et al., 2014; Liu and Davis, 2009; Pitman et al., 2011; Waddell et al., 2000; Yu et al., 2005). Considering their comprehensive projections throughout the MB lobes, the modulation of local circuits requires intracellular synaptic tuning. These studies

together underscore the importance to spatially distinguish individual synapses within a cell. However, the high AZ density in the MB lobes precluded conventional approaches of fluorescent microscopy from profiling synaptic structures of specific neurons (Scheffer et al., 2020).

Here, we present a spatial analysis of individual AZs within the MB circuit at the single-cell resolution, achieved by cell-type-specific visualization of the endogenous active zone protein Bruchpilot (Brp) using the CRISPR/Cas9-mediated split-GFP tagging system. *Drosophila* ELKS/CAST/ERC family member Brp plays a central role in molecular assemblies at AZs by accumulating calcium channels and synaptic vesicles (Fouquet et al., 2009; Hallermann et al., 2010; Kittel et al., 2006; Matkovic et al., 2013; Wagh et al., 2006). Therefore, Brp enrichment serve as a structural proxy suited for estimating synapse function, such as release probability at single AZs (Akbergenova et al., 2018; Newman et al., 2022). We developed a high-throughput quantification pipeline to systematically profile Brp clusters of individual AZs in different MB-innervating neurons. Characterizing the distinct parameters and localization of Brp clusters revealed AZ distribution stereotypy across individuals and significant synaptic heterogeneity within single neurons. These cell-type-specific synapse profiles suggest that AZs are spatially organized at multiple scales, ranging from interindividual stereotypy to neighboring synapses in the same cells.

Results

Establishing an experimental system for cell-type-specific profiling of active zone structures

As the molecular assembly of synapses is reported to be sensitive to the dosage of Brp, we visualized endogenous Brp instead of transgenic expression of tagged Brp (Huang et al., 2020). To label endogenous Brp specifically in designated cell types, we employed the split-GFP tagging system (Kamiyama et al., 2016; Kondo et al., 2020) . Using CRISPR/Cas9, we inserted the GFP_{11} fragment (the eleventh β-strand of the super-folder GFP) just prior to the stop codon of *brp* gene. The GFP_{1-10} fragment is expressed in cells of interest by the GAL4/UAS system. The self-assembly of the split-GFP fragments is

therefore directed only in GAL4-expressing cells and visualizes endogenous Brp through

reconstituted GFP fluorescence (Figure 1A).

As a proof of principle, we first validated the localization and plasticity of split-GFP

tagged Brp (Figure 1B). To verify tagged Brp localization in an individual neuron, we

directed GFP reconstitution and expressed a plasma membrane marker in the serotonergic

DPM neuron using *VT64246-GAL4* (Figure 1C). Confocal microscopy revealed

Brp::reconstituted GFP (Brp::rGFP) signals only to the MB lobes, aligning with DPM

terminals (Waddell et al., 2000). These signals were distributed as discrete clusters within

plasma membrane varicosities resembling presynaptic boutons (Figure 1C), suggesting that

Brp::rGFP clusters labeled individual AZs in the DPM neuron. AZ localization of

Brp::rGFP was further validated by co-localization with anti-Brp immunostaining in the

calyx and the voltage-gated calcium channel subunit Cacophony (Cac) in the PAM

dopaminergic neurons (Figure 1D-E) (Kawasaki et al., 2004; Kittel et al., 2006). We found

that GFP_{11} tagging did not affect *brp* mRNA levels, but slightly decreased protein

expression without altering overall expression pattern in the brain (Figure 1 – figure

supplement 1-2). Brp::rGFP successfully visualized previously reported light-induced AZ

plasticity in the photoreceptors (Sugie et al., 2015). Constant light exposure reduced the

numbers of Brp::rGFP clusters in the R8 terminals in the medulla, compared to constant

darkness (DD) (Figure 1F). This was further substantiated by increased Brp::rGFP cluster

intensities upon knocking-down *rab3* expression (Figure 1 – figure supplement 3),

recapitulating Rab3-dependent regulation of Brp allocation in the presynaptic terminals of

the motor neurons (Graf et al., 2009).

To systematically profile Brp::rGFP clusters, we developed an image processing

pipeline (Figure 1A). In brief, image deconvolution was applied to reduce out-of-focus light

and enhance signal sharpness (Figure 1 – figure supplement 4). Individual Brp::rGFP

clusters were then segmented using the 3D suite plugin in Fiji (Ollion et al., 2013),

generating the 3D region-of-interests (3D ROIs) that encircle each cluster. Brp::rGFP signal

intensity, volume and location of individual clusters were measured using the 3D ROIs. To

optimize the segmentation, we systematically varied the parameters and calculated the F-

score to measure detection accuracy against manually defined ground truth (see Materials and Methods). The F-score is near 1.0 in segmenting Brp::rGFP clusters within DPM and APL neurons and achieved > 0.90 in KCs with certain combinations of segmentation parameters (Figure 1 – figure supplement 5).

To confirm that Brp clusters represent AZs, we quantified the numbers of Brp::rGFP clusters in DPM and APL neurons specifically and compared them to the connectome data. We found that cluster counts closely matched the number of AZs annotated by the hemibrain connectome (Figure 1C) (Scheffer et al., 2020). This result validated our method that accurately isolated Brp::rGFP clusters corresponding to single AZs.

AZ profiles within KCs are compartmentally heterogeneous

Brp accumulation at individual AZs is known to be heterogeneous even within a single motor neuron (Akbergenova et al., 2018; Ehmann et al., 2014; Gratz et al., 2019; Paul et al., 2015). To study such AZ heterogeneity in CNS neurons, we quantified the signal intensity of individual Brp::rGFP clusters in MB lobes. Using KC subtype-specific drivers to direct split-GFP reconstitution in γ KC (*MB009B-GAL4*), α'/β' KCs (*MB370B-GAL4*) and α/β KCs (*MB008B-GAL4*). we found that within-cluster Brp::rGFP intensities are highly diverse in all KC subtypes (Figure 2A).

To visualize spatial patterns of Brp::rGFP intensity in each KC subtype, we reconstructed 3D distribution of Brp::rGFP clusters and color-coded them by signal intensities (Figure 2B). We found that clusters showed noticeable intensity differences between compartments in all subtypes. Even adjacent compartments such as γ1 and γ2, β'2 and β'1, α2 and α3, showed drastic differences (Figure 2B). Importantly, these compartmental patterns were stable across individuals (Figure 2C) (Wu et al., 2025). This result suggests that AZs in different compartments have distinct structures. Since individual KCs of each subtype synapse onto all compartments (Scheffer et al., 2020; Schlegel et al., 2023; Takemura et al., 2024; Zheng et al., 2018), this Brp compartmental heterogeneity is likely formed intracellularly.

Intracellular heterogeneity of Brp concentration at individual AZs and cell-type

dependent diversity

Super-resolution microscopy has shown that the Brp molecular density at individual AZs is

dynamically adjusted in motor neurons (Ghelani et al., 2023; Mrestani et al., 2021). We

hypothesized that the amount of Brp does not necessarily correlate with AZ cluster size. To

verify this, we calculated the Brp concentration by quantifying both the volume (number of

voxels within a 3D ROI) and the Brp::rGFP intensity of each cluster. We found that

clusters with similar sizes can exhibit vastly different signal intensities in the DPM neuron,

suggesting the heterogeneity of Brp concentration at individual AZs (Figure 3A).

Correlation analysis between the Brp::rGFP intensity and cluster volume revealed a

substantial variability among cell types (Figure 3B-E). There were even significant

differences among KC subtypes. These results suggest cell-type dependent variability of the

Brp concentration of individual clusters.

Stereotypy of intracellular distribution of AZs

To study the spatial distribution of Brp clusters within single neurons, we focused on

DANs. *MB504B-GAL4* labels four types of single PPL1 DANs projecting to distinct

compartments (Vogt et al., 2014). Using this driver, we analyzed Brp::rGFP distribution in

each compartment and found it uneven in terminals of both PPL1-α2α'2 and α3 (Figure 4A-

B). To better characterize cluster distribution, we calculated the "AZ density" defined as

the number of Brp::rGFP clusters surrounding a particular cluster within a specified radius

(see Figure 1A and Materials and Methods for detail). By color-coding Brp::rGFP clusters

according to their AZ density, we found that AZs in PPL1-α3 were more localized to the

core of the α3 compartment. This pattern was consistent across individuals (Figure 4C-D),

suggesting a stereotyped AZ distribution within PPL1-α3 neurons.

We extended the analysis to two other single pairs of neurons innervating the MBs,

the DPM and APL neurons. Visualizing AZ density revealed a stereotyped and

compartment specific distribution in the DPM neuron (Figure 5A). Specifically, the AZ

density was constantly high in the α'/β' lobes across all individuals, supporting the known

functional importance of DPM branches in the α'/β' lobes (Keene et al., 2006). In contrast,

this compartmental pattern was less pronounced in the APL neuron. Notably, the ratio of

AZ density between α3 and α'3 is strikingly different among individuals (Figure 5B). Taken

together, these data suggest that the individual variability in AZ distribution depends on the

cell type.

Regulation of spatial configuration of AZs

Characteristics of Brp clusters and AZ distribution in KCs revealed spatial organization at

the level of compartments. Recent studies further showed intra-compartmental variability

of synaptic plasticity upon associative learning, prompting us to analyze sub-

compartmental organization of AZs (Bilz et al., 2020; Davidson et al., 2023). We thus

examined potential local patterns among AZs in proximity by calculating the correlation

between nearest neighbor clusters in terms of their Brp::rGFP intensities (Figure 6A-B).

This analysis revealed strong correlations in KCs and APL but not in DPM (Figure 6C).

We further analyzed the sub-compartmental AZ patterns by examining the spatial

distribution of Brp::rGFP intensity. To reduce the high frequency variability of Brp::rGFP

intensities (Figure 2B), we calculated the "local intensity" for each cluster by applying

mean filtering (the average Brp::rGFP intensity of all surrounding clusters within a

specified radius r, see Materials and Methods for detail). By visualizing the local intensity

in γ KCs, we identified AZ "hot spots", sub-compartment-sized small groups of AZs with

high local intensities, on top of the compartmental differences (Figure 6D). Since Brp

molecules are clustered more in AZ-dense boutons of a single motor neuron (Paul et al.,

2015), we hypothesized that these hot spots correspond to regions with higher AZ densities.

Indeed, we found that hot spots appear in high AZ densities regions in γ KCs (Figure 6D).

Consistently, correlation between the local intensity and the AZ density was high in all KC

subtypes. In contrast, for DPM and APL neurons, the local intensity of Brp::rGFP was not

associated with AZ density (Figure 6E-H). Taken together, these results suggest previously

unidentified sub-compartmental synaptic configuration organized across individual KCs.

Associative learning re-organizes sub-compartmental synaptic configuration

Sub-compartmental synaptic configuration may undergo experience-dependent changes, such as through associative learning (Turrel et al., 2022; Zhang et al., 2018). To examine such structural plasticity in local AZ configurations (Figure 6D, F-H), we visualized Brp::rGFP specifically in KCs using *R13F02-GAL4* (Figure 7A). This induction did not alter MB morphology and short-term memory compared to wild-type (Figure 7 – figure supplement 1-2). We presented flies with odor (4-methylcyclohexanol) with a concomitant (paired) or shifted (unpaired) electric shock (Fig. 7B). Since both paired and unpaired groups were exposed to the same odor and electric shock, the difference between these groups purely represents the effect of association. As single odor conditioning induced memory decaying gradually over one day (Figure 7B), we measured the correlation between the AZ density and local intensity of individual Brp::rGFP clusters (Figure 6F-H) at different time points after conditioning (from 3 min to 1 day). Strikingly, we found that associative learning induced a transient modification to the KC synaptic configurations in a compartment-specific manner (Figure 7C). This learning-induced structural plasticity was specific at 90 min after conditioning and disappeared within one day (Figure 7D), consistent with previous studies reporting short-lived AZ remodeling in the MBs by conditioning (Turrel et al., 2022; Zhang et al., 2018). These results showcase the advantage of split-GFP system in reporting the dynamics of local synapse subsets with high throughputs and contents. Altogether, we suggest that associative learning induces transient memory traces of local structural plasticity in AZ configuration.

Discussion

By leveraging the CRISPR/Cas9 genome editing of the *brp* locus and split-GFP technique, we systematically profiled presynaptic structures of MB neurons in a cell-type-specific manner. This approach enabled high-throughput analysis of multiple brain samples and detailed characterizations, revealing previously inaccessible structural features of AZs in CNS neurons. Cell-type specific labeling of other endogenous synaptic proteins, such as

vesicle-associated proteins, may provide complementary insights to AZ profiling and allow

molecular reconstruction of synapses.

Single-cell level analysis of AZ profiles revealed cell-type-specific stereotypy in

spatial regulation (Figure 4-6). We found that AZ density in the DPM neurons is

consistently high in the α'/β' lobes across individuals (Figure 5). This AZ stereotypy of

DPM explains the functional significance of the branches in the α'/β' lobes. Overexpression

of a Down syndrome cell adhesion molecule (DScam) variant in the DPM neurons was

shown to disrupt their innervation except in the α'/β' lobe, while memory formation

remained unaffected (Keene et al., 2006). Similarly, our analysis identified the intracellular

stereotypy in AZ distribution of the PPL1-α3 DAN within the α3 compartment (Figure 4).

This sub-compartmental organization of the single DAN and layered projections of α/β

KCs may underlie the differential dopaminergic modulation of KC subtypes in associative

learning (Perisse et al., 2013). While such stereotypy is less pronounced in APL, we found

strong correlations between Brp::rGFP intensities of neighboring AZs in contrast to DPM

(Figure 6D). This suggests that AZs in APL neurons are organized more locally, allowing

them to modulate microcircuits composed of KCs, DANs and MBONs (Amin et al., 2020).

Collectively, the spatial scale of AZ organizations depends on cell types, reflecting the

function of individual MB neurons.

We observed that Brp::rGFP intensity correlates strongly with the AZ size in KC

subtypes, in contrast to DPM and APL neurons (Figure 3). Previous studies using

localization microscopy showed that the molecular density of synaptic proteins at AZs can

be dynamically adjusted in motor neurons, potentially modulating the synaptic vesicle

release (Ghelani et al., 2023; Mrestani et al., 2021; Sachidanandan et al., 2023). The

variable Brp concentrations (i.e. the ratio of the Brp::rGFP intensity to the AZ size) in DPM

and APL neurons may represent AZs with distinct Brp densities and therefore different

functional modes (Figure 3) (Mrestani et al., 2021). Notably, a recent study successfully

predicted the neurotransmitter type of neurons using the ultrastructure of dense projections

as one of the parameters (Eckstein et al., 2024), suggesting neurotransmitter-specific AZ

structures. Interestingly, both the DPM and APL neurons are reported to release multiple

neurotransmitters (Davie et al., 2018; Haynes et al., 2015; Lee et al., 2011; Liu and Davis,
2009; Wu et al., 2013; Zeng et al., 2023). Distinct Brp concentrations at AZs might thus
support particular release modes or neurotransmitter types.

In KCs, the spatial analysis of local AZ configuration revealed strong correlations
between Brp::rGFP intensities at neighboring AZs as well as AZ "hot spots", regions with
high Brp localization and AZ densities (Figure 6). These results suggest previously
unidentified, highly localized sub-compartmental AZ structures, which are likely organized
across individual KCs. Presynaptic terminals of a single KC are found to be highly
heterogeneous in terms of their plasticity upon dopamine modulation and associative
learning (Bilz et al., 2020; Davidson et al., 2023). A recent study found the sub-
compartmentally heterogeneous activities of dopamine terminals explain memory
specificity (Yamagata et al., 2021). Such locally distinct dopamine input may be a source of
the AZ hot spot formation in KC terminals. This model is consistent with the localization of
two opposing dopamine receptors Dop1R1 and Dop2R to AZs (Hiramatsu et al., 2024).
Furthermore, we showed that the local configuration undergoes structural plasticity upon
associative learning that involves presynaptic dopaminergic modulation (Aso et al., 2012,
2010; Aso and Rubin, 2016; Burke et al., 2012; Liu et al., 2012; Yamagata et al.,
2015) (Figure 7). Since cAMP signaling plays a crucial role in the AZ structural plasticity
especially during associative memory formation (Baltruschat et al., 2021; Sachidanandan et
al., 2023; Wu et al., 2025), it may underlie learning-induced plasticity in sub-
compartmental AZ configuration (Turrel et al., 2022; Zhang et al., 2018).

Materials and Methods

Animals

Flies were maintained on standard cornmeal food at 25 °C under a 12:12 h light-dark cycle for all experiments. All flies used for experiments are 3-7 days old adult males. Flies were transferred to fresh food vials after hatching and flipped every two days before experiments. The GAL4-UAS system was used to express transgenes, and balancers were removed for animals used in all experiments. Fly strains and resources used in this study are as follow: *MB008B-GAL4* (BDSC 68291) (Aso et al., 2014), *MB009B-GAL4* (BDSC 68292) (Aso et al., 2014); *MB370B-GAL4* (BDSC 68319) (Aso et al., 2014), *MB504B-GAL4* (BDSC 68329) (Aso et al., 2014) , *VT64246-GAL4* (VDRC 204311) (Tirian and Dickson, 2017), *GH146-GAL4* (BDSC 30026) (Stocker et al., 1997), *R58E02-GAL4* (BDSC 41347) (Pfeiffer et al., 2008), *R13F02-GAL4* (BDSC 48571) (Pfeiffer et al., 2008), *R86E01-GAL4* (BDSC 45914) (Jenett et al., 2012)*, Amon-GAL4* (BDSC 30554) (Rhea et al., 2010) *, Rh6-GAL4* (BDSC 7464, by Dr. Claude Desplan), *UAS-CD4::tdTomato* (BDSC 35841, by Dr. Yhu Nung Jan & Chun Han), *UAS-Rab3-RNAi* (BDSC 34655) (Perkins et al., 2015), *brp::GFP$_{11}$* (This study and (Wu et al., 2025)), *UAS-GFP$_{1-10}$* (Kondo et al., 2020).

Generation of *brp::GFP$_{11}$*

CRISPR/Cas9-mediated homologous recombination was used to insert GFP$_{11}$ right before the stop codon of *brp*. The donor vector contained a left homology arm of 747 bp, the GFP11 sequence (Kondo et al., 2020), a floxed 3xP3-RFP marker and a right homology arm of 741 bp. A guide RNA sequence (TCGCAAAGCCACAGATACAC) targeting the *brp* locus was cloned into pBFv-U6.2 (Kondo and Ueda, 2013) to generate a gRNA expression vector. The donor and gRNA vectors were co-injected into fertilized eggs carrying the nos-Cas9 transgene (Kondo and Ueda, 2013). Successful transformants were identified by eye-specific RFP fluorescence. Flies were then crossed to a CyO-Cre balancer to remove the RFP marker. Primers used to clone homology arms are as followed (lowercase letters indicate homologous recombination regions with the doner vector): left

arm forward: gcttgatatcgaattcAAGGACATCGAGGAAAAGGAGAAGAAG; left arm

reverse: agttgggggcgtaggGAAAAAGCTCTTCAAGAAGCCAGCTGGTCC; right arm

forward: tagtataggaacttcTCGCAAAGCCACAGATACACACATCTTGG; right arm

reverse: cgggctgcaggaattcATCGTCGATAATTGTAAGTGCATGCTG

qRT-PCR experiment

Procedures are as previously described (Saito et al., 2025). For each sample, RNA extract

from ~20 heads was purified using TRIzol LS Reagent (Invitrogen) and Direct-zol RNA

Microprep kit (ZYMO RESEARCH), following the provided protocol. The concentration

of extracted RNA samples was then measured using the NanoDrop Spectrophotometer

(Thermo Fisher Scientific) and adjusted to a similar level using diluted water. Reverse

transcription was performed using ReverTra Ace qPCR RT Master Mix (TOYOBO, FSQ-

301). qRT-PCR was performed using iTaq Universal SYBR Green Supermix (BIO-RAD)

and CFX96 Touch Real-Time PCR Detection System (BIO-RAD). No template controls

were also prepared and measured alongside. Three primer pairs for *brp* were designed and

used in the experiment but only one pair that had Ct<30 for all samples was used. Primers

for *brp*, forward: AGCTCAAGGACCACATGGACATC; reverse:

GCGCCATATCCACCTGGTTGTC (Signma-Aldrich). Primers for *Ubiquitin-5E*, forward:

TCTTCACTTGGTCCTGCGTC; reverse: ATGGCTCGACCTCCAAAGTG (eurofins

Genomics). Primers for *αTubulin84B*, forward: GATCGTGTCCTCGATTACCGC;

reverse: GGGAAGTGAATACGTGGGTAGG (eurofins Genomics).

Light exposure experiment

Procedures are as previously described (Sugie et al., 2015). In brief, animals hatched within

one day were transferred to constant darkness (DD, 0 LUX) or constant light (LL, 2000-

4000 LUX by LED panels) environment at 25 °C and kept for 2-3 days. Animals were then

dissected and Brp::rGFP cluster number in individual R8 photoreceptor terminals in the

medulla was quantified.

Associative conditioning

For experiments in Figure 7C-D, adult flies at 3-7 days after eclosion were kept in a 25 °C incubator before and after conditioning. Approximately 50 flies were put in a training tube that is set inside the climate box for 2 minutes before the conditioning. Temperature in the climate box was set at 24 °C and the relative humidity was set to 70%. For the paired group, 2% 4-methycyclohexanol (4-MCH, Sigma-Aldrich, diluted in paraffin oil, Sigma-Aldrich) was presented in a cup with the diameter of 5 mm, together with 12 pulses of electric shocks at 90 V for 1 minute inside the training tube. The unpaired group received the same 1-minute electric shock first, followed by 1-minute rest, and then the same 1-minute 4-MCH presentation. Flies were dissected at 3 min, 20 min, 90 min, 270 min and 1 day after conditioning. Experiments were performed multiple times on different days for a single group.

For measuring the memory decay curve of single odor conditioning, 2% 4-MCH and pure paraffin oil were used for the training. One odor was presented with electric shocks for 1 minute, followed by 1 minute rest, then another odor was presented without electric shocks. For each reciprocal, preference was tested between 2% 4-MCH and paraffin oil. For the differential odor conditioning and short-term memory performance test, 10% 4-MCH and 10% 3-OCT were used for the conditioning, following the same training protocol. Short-term memory performance was tested immediately after the conditioning. Performance index was calculated as $\#(CS-) - \#(CS+)/\#(CS-) + \#(CS+)$.

Sample preparation

Animals used in experiments involving comparisons between control and experimental groups were dissected on the same day. Data were collected from multiple batches of experiments performed on different days. Flies were anesthetized by ice and placed on ice before dissection. The dissection was performed in cold PBS to collect the brain without the ventral nerve cord. The brains were subsequently fixed by 2% paraformaldehyde for 1 h and then washed 3×20 min by 0.1% PBT (0.1% Triton X-100 in PBS) in a PCR tube, typically 5-6 brains in one tube. Samples were mounted on microscope slides using

mounting medium SeeDB2S (Ke et al., 2016). For nc82 immunostaining, fixed brains were

washed 3×20 min with 0.1% PBT and incubated in 3% normal goat serum (NGS; Sigma-

Aldrich; G9023)-0.1% PBT blocking solution for 1 hour at room temperature. The nc82

antibody (DSHB) solution was diluted 1:20 in the blocking solution. Samples were

incubated in antibody solutions at 4°C for 48 hours for both primary and secondary

antibodies.

Confocal imaging

The image acquisition was performed using an Olympus FV1200 confocal microscope

platform equipped with both PMT and GaAsP high sensitivity detectors and a 60×/1.42 NA

oil immersion objective (PLAPON60XO, Olympus). The Kalman filter was turned on to 2×

averaging while scanning. For imaging Brp::rGFP clusters in MB-projecting neurons,

scanning speed was set at 2~4 µs/pixel with a voxel size of lateral 0.079 × 0.079 × axial

0.370 µm. For experimental point spread function (PSF) imaging, SeeDB2S immersed

beads were scanned in a setting that is 60×/1.42 NA oil immersion objective; 473 nm laser

power: 2.0%; 559 nm laser power: 1.5%; voxel size: 0.079 µm × 0.079 µm × 0.370 µm;

scanning speed: 4 µs/pixel, to produce multiple bead images.

PSF acquisition and image deconvolution

Sub-resolution fluorescent beads (Tetraspeck Microspheres 0.1 µm, Thermo Fisher

Scientific, T7279) were imaged to generate the PSF. The bead solution was diluted with

distilled water and sonicated several times to eliminate aggregation. PSF images acquired

were processed using Amira software (Thermo Fisher Scientific) with the Extract Point

Spread Function module. The PSFs extracted from each image were averaged to create a

single PSF for later image deconvolution (Resizing voxel size: 0.079 µm × 0.079 µm ×

0.370 µm; 32 pixels × 32 pixels × 21 slices of image size). Image deconvolution was

performed by using the Richardson-Lucy iterative non-blind algorithm in a Fiji plugin

DeconvolutionLab2 (Sage et al., 2017), or by CLIJ2 GPU-based Richardson-Lucy

deconvolution (Haase et al., 2020). Images were deconvolved 20 times iteratively to

improve image quality.

Brp clusters segmentation and data analysis

For Brp::rGFP cluster detection, the image stack size was adjusted to double the XY plate

pixel number using Fiji to increase the detection precision. The images were subsequently

processed using the "3D maxima finder" and the "3D spot segmentation" function of the

"3D suite" package in Fiji. 3D maxima were identified for each cluster and used as the

starting point for pixel clustering (Ollion et al., 2013). An intensity threshold (threshold)

was first set to reduce noise. Pixels having values below the threshold would become NaN

(not a number). The "noise tolerance" was then set in the 3D maxima finder to adjust its

sensitivity. To be noticed, threshold is only applied to images processed by the 3D maxima

finder to generates the "peak" image stack. 3D spot segmentation function uses both the

"peak" image and the deconvolved Brp::rGFP image to generate "3D ROIs". 3D ROIs were

then applied on the raw images to quantify the volume (total voxel number in ROI) and

Brp::rGFP intensity (total pixel value in ROI) of individual clusters using the "3D ROI

manager". For processing most images, both threshold and noise tolerance were set to

200~400, where the segmentation was optimal based on the F-score benchmarking. "Local

thresholding method" in 3D spot segmentation was set to "Gaussian fit", and "SD value"

was set to 2.

The above procedures may include a tiling and stitching step for images and data

because of the limited computer power. Typically, an image was tiled into 25 sub-stacks

with overlapped margins. After being processed separately, the data acquired from all tiled

sub-stacks was stitched back with a deletion of overlapping data points to generate the

complete detection result. Nearest neighbors were identified for each particle and spatial

distances in between (NND) were calculated using Python. For each sample, mean NND

was calculated separately. For a particular Brp::rGFP cluster, the number of surrounding

clusters within r ($r = 3 \times$ mean NND) spherically including itself is referred as the "AZ

density". The "local intensity" of a particular Brp::rGFP cluster is the average of Brp::rGFP

intensity of all clusters surrounding within r ($r = 3 \times$ mean NND) including itself. The plot

function of Python was used to plot the color-coded 3D reconstructions. A sample code for

analyzing NND, AZ density and local intensity is provided with a demonstration data sheet

(Supplementary file 1 and analysis code).

Detection optimization

The 3D detection parameters were optimized by matching the 3D maxima detection results

to the ground truths. Ground truths were defined by annotating AZs manually, according to

the size, position and signal pattern of Brp clusters in the images. The F-score was used to

assess detection precision. F-score is the harmonic mean of a system's precision and recall

value, calculated by the following formula: $2 \times$ [(Precision $\times$ Recall) / (Precision + Recall)].

The precision is the number of true positives divided by the total number of identified

positives, including incorrect identification. The recall is the number of true positives

divided by the total number of true data points defined as the ground truth. F-scores shown

in Figure S5 were automatically calculated using a customized Fiji Marco that aligned 3D

maxima with ground truth. Different combinations of noise threshold and noise tolerance

were tested for the test images cropped from raw images.

Statistical analysis

Statistical analysis was performed using GraphPad Prism version 8, 9 and 10 or Python.

For statistical comparisons, Kruskal-Wallis test with original False Discovery Rate method

of Benjamini and Hochberg correction was applied unless otherwise specified. The desired

false discovery rate was set to 0.05. Pearson's correlation coefficient (R) was calculated

using Python.

References

471
472

Akbergenova Y, Cunningham KL, Zhang YV, Weiss S, Littleton JT. 2018.
Characterization of developmental and molecular factors underlying release
heterogeneity at Drosophila synapses. *Elife* **7**:e38268. doi:10.7554/elife.38268

Amin H, Apostolopoulou AA, Suárez-Grimalt R, Vrontou E, Lin AC. 2020. Localized
inhibition in the Drosophila mushroom body. *Elife* **9**:e56954. doi:10.7554/elife.56954

Aso Y, Hattori D, Yu Y, Johnston RM, Iyer NA, Ngo T-T, Dionne H, Abbott L, Axel R,
Tanimoto H, Rubin GM. 2014. The neuronal architecture of the mushroom body
provides a logic for associative learning. *Elife* **3**:e04577. doi:10.7554/elife.04577

Aso Y, Herb A, Ogueta M, Siwanowicz I, Templier T, Friedrich AB, Ito K, Scholz H,
Tanimoto H. 2012. Three Dopamine Pathways Induce Aversive Odor Memories with
Different Stability. *PLoS Genet* **8**:e1002768. doi:10.1371/journal.pgen.1002768

Aso Y, Rubin GM. 2016. Dopaminergic neurons write and update memories with cell-type-
specific rules. *Elife* **5**:e16135. doi:10.7554/elife.16135

Aso Y, Siwanowicz I, Bräcker L, Ito K, Kitamoto T, Tanimoto H. 2010. Specific
Dopaminergic Neurons for the Formation of Labile Aversive Memory. *Curr Biol*
20:1445–1451. doi:10.1016/j.cub.2010.06.048

Baltruschat L, Prisco L, Ranft P, Lauritzen JS, Fiala A, Bock DD, Tavosanis G. 2021.
Circuit reorganization in the Drosophila mushroom body calyx accompanies memory
consolidation. *Cell Reports* **34**:108871. doi:10.1016/j.celrep.2021.108871

Bilz F, Geurten BRH, Hancock CE, Widmann A, Fiala A. 2020. Visualization of a
Distributed Synaptic Memory Code in the Drosophila Brain. *Neuron* **106**:963-976.e4.
doi:10.1016/j.neuron.2020.03.010

Burke CJ, Huetteroth W, Owald D, Perisse E, Krashes MJ, Das G, Gohl D, Silies M, Certel
S, Waddell S. 2012. Layered reward signalling through octopamine and dopamine in
Drosophila. *Nature* **492**:433–437. doi:10.1038/nature11614

Chen Y, Akin O, Nern A, Tsui CYK, Pecot MY, Zipursky SL. 2014. Cell-type-Specific
Labeling of Synapses In Vivo through Synaptic Tagging with Recombination. *Neuron*
81:280–293. doi:10.1016/j.neuron.2013.12.021

Davidson AM, Kaushik S, Hige T. 2023. Dopamine-Dependent Plasticity Is
Heterogeneously Expressed by Presynaptic Calcium Activity across Individual Boutons
of the Drosophila Mushroom Body. *eNeuro* **10**:ENEURO.0275-23.2023.
doi:10.1523/eneuro.0275-23.2023

Davie K, Janssens J, Koldere D, Waegeneer MD, Pech U, Kreft Ł, Aibar S, Makhzami S,
Christiaens V, González-Blas CB, Poovathingal S, Hulselmans G, Spanier KI, Moerman
T, Vanspauwen B, Geurs S, Voet T, Lammertyn J, Thienpont B, Liu S, Konstantinides
N, Fiers M, Verstreken P, Aerts S. 2018. A Single-Cell Transcriptome Atlas of the
Aging Drosophila Brain. *Cell* **174**:982-998.e20. doi:10.1016/j.cell.2018.05.057

Davis RL. 2023. Learning and memory using Drosophila melanogaster : a focus on
advances made in the fifth decade of research. *GENETICS*.
doi:10.1093/genetics/iyad085

Dorkenwald S, Matsliah A, Sterling AR, Schlegel P, Yu Szi-chieh, McKellar CE, Lin A,
Costa M, Eichler K, Yin Y, Silversmith W, Schneider-Mizell C, Jordan CS, Brittain D,
Halageri A, Kuehner K, Ogedengbe O, Morey R, Gager J, Kruk K, Perlman E, Yang R,
Deutsch D, Bland D, Sorek M, Lu R, Macrina T, Lee K, Bae JA, Mu S, Nehoran B,
Mitchell E, Popovych S, Wu J, Jia Z, Castro MA, Kemnitz N, Ih D, Bates AS, Eckstein
N, Funke J, Collman F, Bock DD, Jefferis GSXE, Seung HS, Murthy M, Dorkenwald S,
Matsliah A, Sterling AR, Schlegel P, Yu Szi-chieh, McKellar CE, Lin A, Costa M,
Eichler K, Yin Y, Silversmith W, Schneider-Mizell C, Jordan CS, Brittain D, Halageri
A, Kuehner K, Ogedengbe O, Morey R, Gager J, Kruk K, Perlman E, Deutsch D, Bland
D, Sorek M, Lu R, Macrina T, Lee K, Bae JA, Mu S, Nehoran B, Mitchell E, Popovych
S, Wu J, Jia Z, Castro MA, Kemnitz N, Ih D, Bates AS, Eckstein N, Funke J, Collman
F, Bock DD, Jefferis GSXE, Seung HS, Murthy M, Lenizo Z, Burke AT, Willie KP,
Serafetinidis N, Hadjerol N, Willie R, Silverman B, Ocho JA, Bañez J, Candilada RA,
Kristiansen A, Panes N, Yadav A, Tancontian R, Serona S, Dolorosa JI, Vinson KJ,
Garner D, Salem R, Dagohoy A, Skelton J, Lopez M, Capdevila LS, Badalamente G,
Stocks T, Pandey A, Akiatan DJ, Hebditch J, David C, Sapkal D, Monungolh SM, Sane
V, Pielago ML, Albero M, Laude J, Santos M dos, Vohra Z, Wang K, Gogo AM, Kind
E, Mandahay AJ, Martinez C, Asis JD, Nair C, Patel D, Manaytay M, Tamimi IFM, Lim
CA, Ampo PL, Pantujan MD, Javier A, Bautista D, Rana R, Seguido J, Parmar B,
Saguimpa JC, Moore M, Pleijzier MW, Larson M, Hsu J, Joshi I, Kakadiya D, Braun A,
Pilapil C, Gkantia M, Parmar K, Vanderbeck Q, Salgarella I, Dunne C, Munnelly E,
Kang CH, Lörsch L, Lee J, Kmecova L, Sancer G, Baker C, Joroff J, Calle S, Patel Y,
Sato O, Fang S, Salocot J, Salman F, Molina-Obando S, Brooks P, Bui M,
Lichtenberger M, Tamboboy E, Molloy K, Santana-Cruz AE, Hernandez A, Yu
Seongbong, Diwan A, Patel M, Aiken TR, Morejohn S, Koskela S, Yang T, Lehmann
D, Chojetzki J, Sisodiya S, Koolman S, Shiu PK, Cho S, Bast A, Reicher B, Blanquart
M, Houghton L, Choi H, Ioannidou M, Collie M, Eckhardt J, Gorko B, Guo L, Zheng Z,
Poh A, Lin M, Taisz I, Murfin W, Díez ÁS, Reinhard N, Gibb P, Patel N, Kumar S, Yun
M, Wang M, Jones D, Encarnacion-Rivera L, Oswald A, Jadia A, Erginkaya M,
Drummond N, Walter L, Tastekin I, Zhong X, Mabuchi Y, Santiago FJF, Verma U,
Byrne N, Kunze E, Crahan T, Margossian R, Kim H, Georgiev I, Szorenyi F, Adachi A,
Bargeron B, Stürner T, Demarest D, Gür B, Becker AN, Turnbull R, Morren A,
Sandoval A, Moreno-Sanchez A, Pacheco DA, Samara E, Croke H, Thomson A,
Laughland C, Dutta SB, Antón PGA de, Huang B, Pujols P, Haber I, González-Segarra

A, Choe DT, Lukyanova V, Mancini N, Liu Z, Okubo T, Flynn MA, Vitelli G, Laturney M, Li F, Cao S, Manyari-Diaz C, Yim H, Le AD, Maier K, Yu Seungyun, Nam Y, Bąba D, Abusaif A, Francis A, Gayk J, Huntress SS, Barajas R, Kim M, Cui X, Sterne GR, Li A, Park K, Dempsey G, Mathew A, Kim Jinseong, Kim T, Wu G, Dhawan S, Brotas M, Zhang C, Bailey S, Toro AD, Yang R, Gerhard S, Champion A, Anderson DJ, Behnia R, Bidaye SS, Borst A, Chiappe E, Colodner KJ, Dacks A, Dickson B, Garcia D, Hampel S, Hartenstein V, Hassan B, Helfrich-Forster C, Huetteroth W, Kim Jinseop, Kim SS, Kim Y-J, Kwon JY, Lee W-C, Linneweber GA, Maimon G, Mann R, Noselli S, Pankratz M, Prieto-Godino L, Read J, Reiser M, Reyn K von, Ribeiro C, Scott K, Seeds AM, Selcho M, Silies M, Simpson J, Waddell S, Wernet MF, Wilson RI, Wolf FW, Yao Z, Yapici N, Zandawala M. 2024. Neuronal wiring diagram of an adult brain. *Nature* **634**:124–138. doi:10.1038/s41586-024-07558-y

Eckstein N, Bates AS, Champion A, Du M, Yin Y, Schlegel P, Lu AK-Y, Rymer T, Finley-May S, Paterson T, Parekh R, Dorkenwald S, Matsliah A, Yu S-C, McKellar C, Sterling A, Eichler K, Costa M, Seung S, Murthy M, Hartenstein V, Jefferis GSXE, Funke J. 2024. Neurotransmitter classification from electron microscopy images at synaptic sites in Drosophila melanogaster. *Cell* **187**:2574-2594.e23. doi:10.1016/j.cell.2024.03.016

Ehmann N, Linde S van de, Alon A, Ljaschenko D, Keung XZ, Holm T, Rings A, DiAntonio A, Hallermann S, Ashery U, Heckmann M, Sauer M, Kittel RJ. 2014. Quantitative super-resolution imaging of Bruchpilot distinguishes active zone states. *Nat Commun* **5**:4650. doi:10.1038/ncomms5650

Fernández MP, Berni J, Ceriani MF. 2008. Circadian Remodeling of Neuronal Circuits Involved in Rhythmic Behavior. *PLoS Biol* **6**:e69. doi:10.1371/journal.pbio.0060069

Fouquet W, Owald D, Wichmann C, Mertel S, Depner H, Dyba M, Hallermann S, Kittel RJ, Eimer S, Sigrist SJ. 2009. Maturation of active zone assembly by Drosophila Bruchpilot. *J Cell Biol* **186**:129–145. doi:10.1083/jcb.200812150

Gao R, Asano SM, Upadhyayula S, Pisarev I, Milkie DE, Liu T-L, Singh V, Graves A, Huynh GH, Zhao Y, Bogovic J, Colonell J, Ott CM, Zugates C, Tappan S, Rodriguez A, Mosaliganti KR, Sheu S-H, Pasolli HA, Pang S, Xu CS, Megason SG, Hess H, Lippincott-Schwartz J, Hantman A, Rubin GM, Kirchhausen T, Saalfeld S, Aso Y, Boyden ES, Betzig E. 2019. Cortical column and whole-brain imaging with molecular contrast and nanoscale resolution. *Science* **363**. doi:10.1126/science.aau8302

Ghelani T, Escher M, Thomas U, Esch K, Lützkendorf J, Depner H, Maglione M, Parutto P, Gratz S, Matkovic-Rachid T, Ryglewski S, Walter AM, Holcman D, Giles KO, Heine M, Sigrist SJ. 2023. Interactive nanocluster compaction of the ELKS scaffold and Cacophony Ca2+ channels drives sustained active zone potentiation. *Sci Adv* **9**:eade7804. doi:10.1126/sciadv.ade7804

Gilestro GF, Tononi G, Cirelli C. 2009. Widespread Changes in Synaptic Markers as a
Function of Sleep and Wakefulness in Drosophila. *Science* **324**:109–112.
doi:10.1126/science.1166673

Graf ER, Daniels RW, Burgess RW, Schwarz TL, DiAntonio A. 2009. Rab3 Dynamically
Controls Protein Composition at Active Zones. *Neuron* **64**:663–677.
doi:10.1016/j.neuron.2009.11.002

Gratz SJ, Goel P, Bruckner JJ, Hernandez RX, Khateeb K, Macleod GT, Dickman D,
O'Connor-Giles KM. 2019. Endogenous tagging reveals differential regulation of Ca2+
channels at single AZs during presynaptic homeostatic potentiation and depression. *J*
Neurosci **39**:3068–18. doi:10.1523/jneurosci.3068-18.2019

Haase R, Royer LA, Steinbach P, Schmidt D, Dibrov A, Schmidt U, Weigert M, Maghelli
N, Tomancak P, Jug F, Myers EW. 2020. CLIJ: GPU-accelerated image processing for
everyone. *Nat Methods* **17**:5–6. doi:10.1038/s41592-019-0650-1

Hallermann S, Kittel RJ, Wichmann C, Weyhersmüller A, Fouquet W, Mertel S, Owald D,
Eimer S, Depner H, Schwärzel M, Sigrist SJ, Heckmann M. 2010. Naked Dense Bodies
Provoke Depression. *J Neurosci* **30**:14340–14345. doi:10.1523/jneurosci.2495-10.2010

Haynes PR, Christmann BL, Griffith LC. 2015. A single pair of neurons links sleep to
memory consolidation in Drosophila melanogaster. *eLife* **4**:e03868.
doi:10.7554/elife.03868

Hiramatsu S, Saito K, Kondo S, Katow H, Yamagata N, Wu C-F, Tanimoto H. 2024. Cell-
type-specific fluorescent tagging of endogenous target proteins reveals synaptic
enrichment and dynamic regulations of dopamine receptors. doi:10.7554/elife.98358.1

Huang S, Piao C, Beuschel CB, Götz T, Sigrist SJ. 2020. Presynaptic Active Zone
Plasticity Encodes Sleep Need in Drosophila. *Curr Biol* **30**:1077-1091.e5.
doi:10.1016/j.cub.2020.01.019

Jenett A, Rubin GM, Ngo T-TB, Shepherd D, Murphy C, Dionne H, Pfeiffer BD, Cavallaro
A, Hall D, Jeter J, Iyer N, Fetter D, Hausenfluck JH, Peng H, Trautman ET, Svirskas
RR, Myers EW, Iwinski ZR, Aso Y, DePasquale GM, Enos A, Hulamm P, Lam SCB, Li
H-H, Laverty TR, Long F, Qu L, Murphy SD, Rokicki K, Safford T, Shaw K, Simpson
JH, Sowell A, Tae S, Yu Y, Zugates CT. 2012. A GAL4-Driver Line Resource for
Drosophila Neurobiology. *Cell Rep* **2**:991–1001. doi:10.1016/j.celrep.2012.09.011

Kamiyama D, Sekine S, Barsi-Rhyne B, Hu J, Chen B, Gilbert LA, Ishikawa H, Leonetti
MD, Marshall WF, Weissman JS, Huang B. 2016. Versatile protein tagging in cells with
split fluorescent protein. *Nat Commun* **7**:11046. doi:10.1038/ncomms11046

Kawasaki F, Zou B, Xu X, Ordway RW. 2004. Active Zone Localization of Presynaptic
Calcium Channels Encoded by the cacophony Locus of Drosophila. *J Neurosci* **24**:282–
285. doi:10.1523/jneurosci.3553-03.2004

Ke M-T, Nakai Y, Fujimoto S, Takayama R, Yoshida S, Kitajima TS, Sato M, Imai T.
2016. Super-Resolution Mapping of Neuronal Circuitry With an Index-Optimized
Clearing Agent. *Cell Reports* **14**:2718–2732. doi:10.1016/j.celrep.2016.02.057

Keene AC, Krashes MJ, Leung B, Bernard JA, Waddell S. 2006. Drosophila Dorsal Paired
Medial Neurons Provide a General Mechanism for Memory Consolidation. *Curr Biol*
16:1524–1530. doi:10.1016/j.cub.2006.06.022

Keene AC, Stratmann M, Keller A, Perrat PN, Vosshall LB, Waddell S. 2004. Diverse
Odor-Conditioned Memories Require Uniquely Timed Dorsal Paired Medial Neuron
Output. *Neuron* **44**:521–533. doi:10.1016/j.neuron.2004.10.006

Kittel RJ, Wichmann C, Rasse TM, Fouquet W, Schmidt M, Schmid A, Wagh DA, Pawlu
C, Kellner RR, Willig KI, Hell SW, Buchner E, Heckmann M, Sigrist SJ. 2006.
Bruchpilot Promotes Active Zone Assembly, Ca^{2+} Channel Clustering, and Vesicle
Release. *Science* **312**:1051–1054. doi:10.1126/science.1126308

Kondo S, Takahashi T, Yamagata N, Imanishi Y, Katow H, Hiramatsu S, Lynn K, Abe A,
Kumaraswamy A, Tanimoto H. 2020. Neurochemical Organization of the Drosophila
Brain Visualized by Endogenously Tagged Neurotransmitter Receptors. *Cell Reports*
30:284-297.e5. doi:10.1016/j.celrep.2019.12.018

Kondo S, Ueda R. 2013. Highly Improved Gene Targeting by Germline-Specific Cas9
Expression in Drosophila. *Genetics* **195**:715–721. doi:10.1534/genetics.113.156737

Kremer MC, Christiansen F, Leiss F, Paehler M, Knapek S, Andlauer TFM, Förstner F,
Kloppenburg P, Sigrist SJ, Tavosanis G. 2010. Structural Long-Term Changes at
Mushroom Body Input Synapses. *Curr Biol* **20**:1938–1944.
doi:10.1016/j.cub.2010.09.060

Lee P-T, Lin H-W, Chang Y-H, Fu T-F, Dubnau J, Hirsh J, Lee T, Chiang A-S. 2011.
Serotonin–mushroom body circuit modulating the formation of anesthesia-resistant
memory in Drosophila. *Proc Natl Acad Sci* **108**:13794–13799.
doi:10.1073/pnas.1019483108

Lin AC, Bygrave AM, Calignon A de, Lee T, Miesenböck G. 2014. Sparse, decorrelated
odor coding in the mushroom body enhances learned odor discrimination. *Nat Neurosci*
17:559–568. doi:10.1038/nn.3660

Liu C, Plaçais P-Y, Yamagata N, Pfeiffer BD, Aso Y, Friedrich AB, Siwanowicz I, Rubin GM, Preat T, Tanimoto H. 2012. A subset of dopamine neurons signals reward for odour memory in Drosophila. *Nature* **488**:512–516. doi:10.1038/nature11304

Liu X, Davis RL. 2009. The GABAergic anterior paired lateral neuron suppresses and is suppressed by olfactory learning. *Nat Neurosci* **12**:53–59. doi:10.1038/nn.2235

Matkovic T, Siebert M, Knoche E, Depner H, Mertel S, Owald D, Schmidt M, Thomas U, Sickmann A, Kamin D, Hell SW, Bürger J, Hollmann C, Mielke T, Wichmann C, Sigrist SJ. 2013. The Bruchpilot cytomatrix determines the size of the readily releasable pool of synaptic vesicles. *J Cell Biol* **202**:667–683. doi:10.1083/jcb.201301072

Mosca TJ, Luo L. 2014. Synaptic organization of the Drosophila antennal lobe and its regulation by the Teneurins. *eLife* **3**:e03726. doi:10.7554/elife.03726

Mrestani A, Pauli M, Kollmannsberger P, Repp F, Kittel RJ, Eilers J, Doose S, Sauer M, Sirén A-L, Heckmann M, Paul MM. 2021. Active zone compaction correlates with presynaptic homeostatic potentiation. *Cell Rep* **37**:109770. doi:10.1016/j.celrep.2021.109770

Newman ZL, Bakshinskaya D, Schultz R, Kenny SJ, Moon S, Aghi K, Stanley C, Marnani N, Li R, Bleier J, Xu K, Isacoff EY. 2022. Determinants of synapse diversity revealed by super-resolution quantal transmission and active zone imaging. *Nat Commun* **13**:229. doi:10.1038/s41467-021-27815-2

Ollion J, Cochennec J, Loll F, Escudé C, Boudier T. 2013. TANGO: a generic tool for high-throughput 3D image analysis for studying nuclear organization. *Bioinformatics* **29**:1840–1841. doi:10.1093/bioinformatics/btt276

Paul MM, Pauli M, Ehmann N, Hallermann S, Sauer M, Kittel RJ, Heckmann M. 2015. Bruchpilot and Synaptotagmin collaborate to drive rapid glutamate release and active zone differentiation. *Front Cell Neurosci* **9**:29. doi:10.3389/fncel.2015.00029

Perisse E, Yin Y, Lin AC, Lin S, Huetteroth W, Waddell S. 2013. Different Kenyon Cell Populations Drive Learned Approach and Avoidance in Drosophila. *Neuron* **79**:945–956. doi:10.1016/j.neuron.2013.07.045

Perkins LA, Holderbaum L, Tao R, Hu Y, Sopko R, McCall K, Yang-Zhou D, Flockhart I, Binari R, Shim H-S, Miller A, Housden A, Foos M, Randkelv S, Kelley C, Namgyal P, Villalta C, Liu L-P, Jiang X, Huan-Huan Q, Wang X, Fujiyama A, Toyoda A, Ayers K, Blum A, Czech B, Neumuller R, Yan D, Cavallaro A, Hibbard K, Hall D, Cooley L, Hannon GJ, Lehmann R, Parks A, Mohr SE, Ueda R, Kondo S, Ni J-Q, Perrimon N.

2015. The Transgenic RNAi Project at Harvard Medical School: Resources and
Validation. *Genetics* **201**:843–852. doi:10.1534/genetics.115.180208

Pfeiffer BD, Jenett A, Hammonds AS, Ngo T-TB, Misra S, Murphy C, Scully A, Carlson
JW, Wan KH, Laverty TR, Mungall C, Svirskas R, Kadonaga JT, Doe CQ, Eisen MB,
Celniker SE, Rubin GM. 2008. Tools for neuroanatomy and neurogenetics in
Drosophila. *Proc Natl Acad Sci* **105**:9715–9720. doi:10.1073/pnas.0803697105

Pitman JL, Huetteroth W, Burke CJ, Krashes MJ, Lai S-L, Lee T, Waddell S. 2011. A Pair
of Inhibitory Neurons Are Required to Sustain Labile Memory in the Drosophila
Mushroom Body. *Curr Biol* **21**:855–861. doi:10.1016/j.cub.2011.03.069

Rhea JM, Wegener C, Bender M. 2010. The Proprotein Convertase Encoded by
amontillado (amon) Is Required in Drosophila Corpora Cardiaca Endocrine Cells
Producing the Glucose Regulatory Hormone AKH. *PLoS Genet* **6**:e1000967.
doi:10.1371/journal.pgen.1000967

Sachidanandan D, Aravamudhan A, Mrestani A, Nerlich J, Lamberty M, Hasenauer N,
Ehmann N, Pauls D, Seubert T, Maiellaro I, Selcho M, Heckmann M, Hallermann S,
Kittel RJ. 2023. Rab3 mediates cyclic AMP-dependent presynaptic plasticity and
olfactory learning. *bioRxiv* 2023.12.21.572589. doi:10.1101/2023.12.21.572589

Sachse S, Rueckert E, Keller A, Okada R, Tanaka NK, Ito K, Vosshall LB. 2007. Activity-
Dependent Plasticity in an Olfactory Circuit. *Neuron* **56**:838–850.
doi:10.1016/j.neuron.2007.10.035

Sage D, Donati L, Soulez F, Fortun D, Schmit G, Seitz A, Guiet R, Vonesch C, Unser M.
2017. DeconvolutionLab2: An open-source software for deconvolution microscopy.
Methods **115**:28–41. doi:10.1016/j.ymeth.2016.12.015

Saito K, Kanno M, Tanimoto H, Ichinose T. 2025. Ecdysteroid-DopEcR signaling in
neuronal and midgut cells mediates toxin avoidance and detoxification in Drosophila.
Curr Biol **35**:3418-3428.e4. doi:10.1016/j.cub.2025.06.023

Scheffer LK, Xu CS, Januszewski M, Lu Z, Takemura Shin-ya, Hayworth KJ, Huang GB,
Shinomiya K, Maitlin-Shepard J, Berg S, Clements J, Hubbard PM, Katz WT, Umayam
L, Zhao T, Ackerman D, Blakely T, Bogovic J, Dolafi T, Kainmueller D, Kawase T,
Khairy KA, Leavitt L, Li PH, Lindsey L, Neubarth N, Olbris DJ, Otsuna H, Trautman
ET, Ito M, Bates AS, Goldammer J, Wolff T, Svirskas R, Schlegel P, Neace E, Knecht
CJ, Alvarado CX, Bailey DA, Ballinger S, Borycz JA, Canino BS, Cheatham N, Cook
M, Dreher M, Duclos O, Eubanks B, Fairbanks K, Finley S, Forknall N, Francis A,
Hopkins GP, Joyce EM, Kim S, Kirk NA, Kovalyak J, Lauchie SA, Lohff A,
Maldonado C, Manley EA, McLin S, Mooney C, Ndama M, Ogundeyi O, Okeoma N,

Ordish C, Padilla N, Patrick CM, Paterson T, Phillips EE, Phillips EM, Rampally N, Ribeiro C, Robertson MK, Rymer JT, Ryan SM, Sammons M, Scott AK, Scott AL, Shinomiya A, Smith C, Smith K, Smith NL, Sobeski MA, Suleiman A, Swift J, Takemura Satoko, Talebi I, Tarnogorska D, Tenshaw E, Tokhi T, Walsh JJ, Yang T, Horne JA, Li F, Parekh R, Rivlin PK, Jayaraman V, Costa M, Jefferis GS, Ito K, Saalfeld S, George R, Meinertzhagen IA, Rubin GM, Hess HF, Jain V, Plaza SM. 2020. A connectome and analysis of the adult Drosophila central brain. *Elife* **9**:e57443. doi:10.7554/elife.57443

Schlegel P, Yin Y, Bates AS, Dorkenwald S, Eichler K, Brooks P, Han DS, Gkantia M, Santos Marcia dos, Munnelly EJ, Badalamente G, Capdevila LS, Sane VA, Fragniere AMC, Kiassat L, Pleijzier MW, Stürner T, Tamimi IFM, Dunne CR, Salgarella I, Javier A, Fang S, Perlman E, Kazimiers T, Jagannathan SR, Matsliah A, Sterling AR, Yu Szi-chieh, McKellar CE, Schlegel P, Yin Y, Dorkenwald S, Eichler K, Brooks P, Gkantia M, Badalamente G, Capdevila LS, Tamimi IFM, Salgarella I, Javier A, Fang S, Perlman E, Matsliah A, Yu Szi-chieh, McKellar CE, Kruk K, Bland D, Lenizo Z, Burke AT, Willie KP, Bates AS, Serafetinidis N, Hadjerol N, Willie R, Silverman B, Ocho JA, Bañez J, Candilada RA, Gager J, Kristiansen A, Panes N, Yadav A, Tancontian R, Serona S, Dolorosa JI, Vinson KJ, Garner D, Salem R, Dagohoy A, Skelton J, Lopez M, Stocks T, Pandey A, Akiatan DJ, Hebditch J, David C, Sapkal D, Monungolh SM, Sane V, Pielago ML, Albero M, Laude J, Santos Márcia dos, Deutsch D, Vohra Z, Wang K, Gogo AM, Kind E, Mandahay AJ, Martinez C, Asis JD, Nair C, Patel D, Manaytay M, Lim CA, Ampo PL, Pantujan MD, Bautista D, Rana R, Seguido J, Parmar B, Saguimpa JC, Moore M, Pleijzier MW, Larson M, Hsu J, Joshi I, Kakadiya D, Braun A, Pilapil C, Parmar K, Vanderbeck Q, Dunne C, Munnelly E, Kang CH, Lörsch L, Lee J, Kmecova L, Sancer G, Baker C, Joroff J, Calle S, Patel Y, Sato O, Salocot J, Salman F, Molina-Obando S, Bui M, Lichtenberger M, Tamboboy E, Molloy K, Santana-Cruz AE, Hernandez A, Yu Seongbong, Sorek M, Diwan A, Patel M, Aiken TR, Morejohn S, Koskela S, Yang T, Lehmann D, Chojetzki J, Sisodiya S, Koolman S, Shiu PK, Cho S, Bast A, Reicher B, Blanquart M, Houghton L, Choi H, Ioannidou M, Collie M, Eckhardt J, Gorko B, Guo L, Zheng Z, Poh A, Lin M, Taisz I, Murfin W, Díez ÁS, Reinhard N, Gibb P, Patel N, Kumar S, Yun M, Wang M, Jones D, Encarnacion-Rivera L, Oswald A, Jadia A, Erginkaya M, Drummond N, Walter L, Tastekin I, Zhong X, Mabuchi Y, Santiago FJF, Verma U, Byrne N, Kunze E, Crahan T, Margossian R, Kim H, Georgiev I, Szorenyi F, Adachi A, Bargeron B, Stürner T, Demarest D, Gür B, Becker AN, Turnbull R, Morren A, Sandoval A, Moreno-Sanchez A, Pacheco DA, Samara E, Croke H, Thomson A, Laughland C, Dutta SB, Antón PGA de, Huang B, Pujols P, Haber I, González-Segarra A, Lin A, Choe DT, Lukyanova V, Mancini N, Liu Z, Okubo T, Flynn MA, Vitelli G, Laturney M, Li F, Cao S, Manyari-Diaz C, Yim H, Le AD, Maier K, Yu Seungyun, Nam Y, Bąba D, Abusaif A, Francis A, Gayk J, Huntress SS, Barajas R, Kim M, Cui X, Sterling AR, Sterne GR, Li A, Park K, Dempsey G, Mathew A, Kim Jinseong, Kim T, Wu G, Dhawan S, Brotas M, Zhang C, Bailey S, Toro AD, Lee K, Macrina T, Schneider-Mizell C, Popovych S, Ogedengbe O, Yang R, Halageri A, Silversmith W, Gerhard S, Champion A, Eckstein N, Ih D, Kemnitz N, Castro M, Jia Z,

Wu J, Mitchell E, Nehoran B, Mu S, Bae JA, Lu R, Morey R, Kuehner K, Brittain D, Jordan CS, Anderson DJ, Behnia R, Bidaye SS, Borst A, Chiappe E, Collman F, Colodner KJ, Dacks A, Dickson B, Funke J, Garcia D, Hampel S, Hartenstein V, Hassan B, Helfrich-Forster C, Huetteroth W, Kim Jinseop, Kim SS, Kim Y-J, Kwon JY, Lee W-C, Linneweber GA, Maimon G, Mann R, Noselli S, Pankratz M, Prieto-Godino L, Read J, Reiser M, Reyn K von, Ribeiro C, Scott K, Seeds AM, Selcho M, Silies M, Simpson J, Waddell S, Wernet MF, Wilson RI, Wolf FW, Yao Z, Yapici N, Zandawala M, Costa M, Seung HS, Murthy M, Bock DD, Jefferis GSXE, Costa M, Seung HS, Murthy M, Hartenstein V, Bock DD, Jefferis GSXE. 2024. Whole-brain annotation and multi-connectome cell typing of Drosophila. *Nature* **634**:139–152. doi:10.1038/s41586-024-07686-5

Schlegel P, Yin Y, Bates AS, Dorkenwald S, Eichler K, Brooks P, Han DS, Gkantia M, Santos M dos, Munnelly EJ, Badalamente G, Capdevila LS, Sane VA, Pleijzier MW, Tamimi IFM, Dunne CR, Salgarella I, Javier A, Fang S, Perlman E, Kazimiers T, Jagannathan SR, Matsliah A, Sterling AR, Yu S, McKellar CE, Consortium F, Costa M, Seung HS, Murthy M, Hartenstein V, Bock DD, Jefferis GSXE. 2023. A consensus cell type atlas from multiple connectomes reveals principles of circuit stereotypy and variation. *bioRxiv* 2023.06.27.546055. doi:10.1101/2023.06.27.546055

Stocker RF, Heimbeck G, Gendre N, Belle JS de. 1997. Neuroblast ablation in Drosophila P[GAL4] lines reveals origins of olfactory interneurons. *J Neurobiol* **32**:443–456. doi:10.1002/(sici)1097-4695(199705)32:5<443::aid-neu1>3.0.co;2-5

Sugie A, Hakeda-Suzuki S, Suzuki E, Silies M, Shimozono M, Möhl C, Suzuki T, Tavosanis G. 2015. Molecular Remodeling of the Presynaptic Active Zone of Drosophila Photoreceptors via Activity-Dependent Feedback. *Neuron* **86**:711–725. doi:10.1016/j.neuron.2015.03.046

Takemura Shin-ya, Hayworth KJ, Huang GB, Januszewski M, Lu Z, Marin EC, Preibisch S, Xu CS, Bogovic J, Champion AS, Cheong HS, Costa M, Eichler K, Katz W, Knecht C, Li F, Morris BJ, Ordish C, Rivlin PK, Schlegel P, Shinomiya K, Stürner T, Zhao T, Badalamente G, Bailey D, Brooks P, Canino BS, Clements J, Cook M, Duclos O, Dunne CR, Fairbanks K, Fang S, Finley-May S, Francis A, George R, Gkantia M, Harrington K, Hopkins GP, Hsu J, Hubbard PM, Javier A, Kainmueller D, Korff W, Kovalyak J, Krzemiński D, Lauchie SA, Lohff A, Maldonado C, Manley EA, Mooney C, Neace E, Nichols M, Ogundeyi O, Okeoma N, Paterson T, Phillips E, Phillips EM, Ribeiro C, Ryan SM, Rymer JT, Scott AK, Scott AL, Shepherd D, Shinomiya A, Smith C, Smith N, Suleiman A, Takemura Satoko, Talebi I, Tamimi IF, Trautman ET, Umayam L, Walsh JJ, Yang T, Rubin GM, Scheffer LK, Funke J, Saalfeld S, Hess HF, Plaza SM, Card GM, Jefferis GS, Berg S. 2024. A Connectome of the Male Drosophila Ventral Nerve Cord. doi:10.7554/elife.97769.1

Tirian L, Dickson BJ. 2017. The VT GAL4, LexA, and split-GAL4 driver line collections for targeted expression in the Drosophila nervous system. *bioRxiv* 198648. doi:10.1101/198648

Turrel O, Ramesh N, Escher MJF, Pooryasin A, Sigrist SJ. 2022. Transient active zone remodeling in the Drosophila mushroom body supports memory. *Curr Biol*. doi:10.1016/j.cub.2022.10.017

Vogt K, Schnaitmann C, Dylla KV, Knapek S, Aso Y, Rubin GM, Tanimoto H. 2014. Shared mushroom body circuits underlie visual and olfactory memories in Drosophila. *Elife* **3**:e02395. doi:10.7554/elife.02395

Waddell S, Armstrong JD, Kitamoto T, Kaiser K, Quinn WG. 2000. The amnesiac Gene Product Is Expressed in Two Neurons in the Drosophila Brain that Are Critical for Memory. *Cell* **103**:805–813. doi:10.1016/s0092-8674(00)00183-5

Wagh DA, Rasse TM, Asan E, Hofbauer A, Schwenkert I, Dürrbeck H, Buchner S, Dabauvalle M-C, Schmidt M, Qin G, Wichmann C, Kittel R, Sigrist SJ, Buchner E. 2006. Bruchpilot, a Protein with Homology to ELKS/CAST, Is Required for Structural Integrity and Function of Synaptic Active Zones in Drosophila. *Neuron* **49**:833–844. doi:10.1016/j.neuron.2006.02.008

Wu C-L, Shih M-FM, Lee P-T, Chiang A-S. 2013. An Octopamine-Mushroom Body Circuit Modulates the Formation of Anesthesia-Resistant Memory in Drosophila. *Curr Biol* **23**:2346–2354. doi:10.1016/j.cub.2013.09.056

Wu H, Eno S, Jinnai K, Abe A, Saito K, Maekawa Y, Williams DW, Yamagata N, Kondo S, Tanimoto H. 2025. Octopamine signaling regulates the intracellular pattern of the presynaptic active zone scaffold within Drosophila mushroom body neurons. *PLOS Biol* **23**:e3003449. doi:10.1371/journal.pbio.3003449

Yamagata N, Ezaki T, Takahashi T, Wu H, Tanimoto H. 2021. Presynaptic inhibition of dopamine neurons controls optimistic bias. *Elife* **10**:e64907. doi:10.7554/elife.64907

Yamagata N, Ichinose T, Aso Y, Plaçais P-Y, Friedrich AB, Sima RJ, Preat T, Rubin GM, Tanimoto H. 2015. Distinct dopamine neurons mediate reward signals for short- and long-term memories. *Proc National Acad Sci* **112**:578–583. doi:10.1073/pnas.1421930112

Yu D, Keene AC, Srivatsan A, Waddell S, Davis RL. 2005. Drosophila DPM Neurons Form a Delayed and Branch-Specific Memory Trace after Olfactory Classical Conditioning. *Cell* **123**:945–957. doi:10.1016/j.cell.2005.09.037

Zeng J, Li X, Zhang R, Lv M, Wang Y, Tan K, Xia X, Wan J, Jing M, Zhang X, Li Yu,
Yang Y, Wang L, Chu J, Li Yan, Li Yulong. 2023. Local 5-HT signaling bi-
directionally regulates the coincidence time window for associative learning. *Neuron*
111:1118-1135.e5. doi:10.1016/j.neuron.2022.12.034

Zhang X, Li Q, Wang L, Liu Z-J, Zhong Y. 2018. Active Protection: Learning-Activated
Raf/MAPK Activity Protects Labile Memory from Rac1-Independent Forgetting.
Neuron **98**:142-155.e4. doi:10.1016/j.neuron.2018.02.025

Zheng Z, Lauritzen JS, Perlman E, Robinson CG, Nichols M, Milkie D, Torrens O, Price J,
Fisher CB, Sharifi N, Calle-Schuler SA, Kmecova L, Ali IJ, Karsh B, Trautman ET,
Bogovic JA, Hanslovsky P, Jefferis GSXE, Kazhdan M, Khairy K, Saalfeld S, Fetter
RD, Bock DD. 2018. A Complete Electron Microscopy Volume of the Brain of Adult
Drosophila melanogaster. *Cell* **174**:730-743.e22. doi:10.1016/j.cell.2018.06.019

Acknowledgments

We thank all lab members of Tanimoto Lab at Tohoku University for valuable discussions.

We thank Dr. Yuki Tsukada at Keio University for crucial advice on Fiji-based image

analysis. We thank Vienna Drosophila Resource Center and Bloomington Drosophila Stock

Center for transgenic flies.

Funding:

Japan Science and Technology Agency JPMJSP2114 (HW)

Tohoku University Research Program 'Frontier Research in Duo' (HT)

Ministry of Education, Culture, Sports, Science and Technology 20H00519 (HT)

Ministry of Education, Culture, Sports, Science and Technology 22H05481 (HT)

Ministry of Education, Culture, Sports, Science and Technology 22KK0106 (HT)

Author contributions:

Conceptualization: HW, HT

Methodology: HW, YM, NY, SE, SK

Investigation: HW, YM, SE

Visualization: HW, SE

Resources: SK

Supervision: HW, NY, HT

Writing—original draft: HY, HT

Writing—review & editing: HW, SE, HT

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

Further information and requests on resources and reagents should be directed to and will

be fulfilled by the corresponding author, Dr. Hiromu Tanimoto (hiromut@m.tohoku.ac.jp).

All data and codes needed to evaluate the conclusions in the paper are present in the paper.

Main figures

Figure 1 Spatial characterization of Brp::rGFP clusters at single-AZ level.

(**A**) Schematic of the split-GFP tagging strategy. The GFP$_{11}$ fragment is inserted by

CRISPR/Cas9 just prior to the stop codon of *brp*, while GFP$_{1-10}$ is expressed in specific cell

types via GAL4-UAS system.

(**B**) Brp::rGFP signals in DPM. The entire MB lobe structure is shown. The black box in

the left panel indicates the zoom-in area (right panel). GFP_{1-10} and CD4::tdTomato

(magenta) were co-expressed using *VT64246-GAL4*. Scale bar, 20 μm (left); 5 μm (right).

(**C**) Schematic showing brain regions and cell types analyzed in Figure 1C-F and H.

(**D**) Brp::rGFP (green) and anti-Brp immunostaining signals (blue) in the mushroom body

calyx. GFP_{1-10} was expressed using pan-neuronal driver *R57C10-GAL4*. Scale bar, 1 μm.

(**E**) Brp::rGFP (green) and Cac::tdTomato (orange) in PAM-γ5 dopamine neuron terminals.

GFP_{1-10} and Cac::tdTomato were co-expressed using *R15A04-GAL4.* Scale bar, 1 μm.

(**F**) Brp::rGFP (green) in R8 photoreceptors upon constant dark or light conditions. GFP_{1-10}

and CD4::TdTomato (magenta) were co-expressed using *Rh6-GAL4*. Brp::rGFP cluster

number of each R8 photoreceptor terminal in the medulla were quantified. n = 30 (3 brain

samples, 10 terminals from each brain). Welch's t test. Scale bar, 5 μm.

(**G**) Image processing pipeline. Raw images were processed with image deconvolution to

improve the signal quality. 3D maxima (green pixels) are detected for each Brp::rGFP

cluster. 3D ROIs are indicated by yellow circles. Nearest neighbor distance (NND), *r* is

defined as 3 × mean NND. Scale bar, 1 μm.

(**H**) Number of Brp::rGFP clusters detected in DPM and APL. The dashed lines indicate the

number of electron dense projections identified in the hemibrain connectome for DPM

(17530) and APL (10789) on the MB lobes and peduncle (Scheffer et al., 2020).

Schematics illustrate the innervation patterns of DPM and APL neurons. For both DPM and

APL, n = 5. Lower panels show the representative 3D reconstructions of Brp::rGFP clusters

in DPM and APL. Scale bars, 20 μm. Data were presented as mean ± s.e.m.

Figure 2 Compartmentalized AZ structures of KCs.

(**A**) Histograms of the signal intensity of individual Brp::rGFP clusters from three KC

subtypes. Values are normalized by the mean intensity in each data set (each sample). Each

line represents the histogram of one independent sample. γ KCs (*MB009B-GAL4*), n = 6;

α/β KCs (*MB370B-GAL4*), n = 5; α'/β' KCs (*MB008B-GAL4*), n = 5.

(**B**) 3D reconstructions of Brp::rGFP clusters in three KC subtypes, colored by the

Brp::rGFP intensity. The approximate locations of compartments are as indicated.

Schematics illustrate the innervation patterns of different KC subtypes. Min. and Max. were

set to represent the lowest and highest 5% of Brp::rGFP intensity value in the dataset

respectively. Scale bars, 20 μm.

(**C**) Signal intensity of Brp:rGFP clusters in each compartment. Brp::rGFP clusters were

quantified compartmentally. Medians in different compartments are showed as the ratio

918   against the average of five compartments in the corresponding KC subtype. Each line

919   represents one independent sample.

Figure 3 The variability of Brp::rGFP concentrations depends on cell type.

(**A**) Brp::rGFP clusters with distinct intensities but similar size in DPM. The left panel

shows a cropped image of Brp::rGFP in DPM. White boxes indicate areas zoomed-in in

panels 1 and 2. Transparent white lines in panels 1 and 2 show the lines on which the

intensity profiles were plotted. Intensity profiles were plotted for AZ1 (magenta) and AZ2

(blue) respectively.

(**B-D**) Scatter plots showing the correlation between the Brp::rGFP intensity and cluster

volume in DPM, APL and α/β KCs. Data of three representative samples were shown.

Pearson correlation coefficient R were calculated for each sample.

(**E**) Correlations between the intensity and volume of Brp::rGFP clusters in different cell

types. DPM (n = 6), APL (n = 5), γ KCs (n = 6, *MB009B-GAL4*), α'/β' KCs (n = 5,

MB370B-GAL4), α/β KCs (n = 5, *MB008B-GAL4*). DPM vs. α'/β' KCs: P = 0.0253; DPM

vs. α/β KCs: P = 0.0009; APL vs. α/β KCs: P = 0.0035; γ KCs vs. α/β KCs: P = 0.0418.

Values marked with different lowercase letters represent significant difference (P < 0.05);

Data were presented as box plots showing center (median), whiskers (Min. to Max.).

Figure 4 Stereotyped AZ distribution of PPL1-α3 DAN.

(**A**) Brp::rGFP in PPL1 DANs. UAS-GFP$_{1-10}$ was expressed using *MB504B-GAL4*. Dashed line marks the rough boundary between α3 and α2α'2 compartments. Schematic illustrates the innervation patterns of PPL1 DANs within the MB. Scale bar, 20 μm.

(**B**) Brp::rGFP intensity profiles of α3 and α2α'2 compartments. Left panels show the max-projection images of α3 and α2α'2 optical coronal sections. Transparent gray stripes indicate areas where intensity profiles are plotted in the right panels. Scale bars, 10 μm.

(**C**) 3D reconstruction colored by the AZ density in PPL-1 DANs. Color scale: Min. = 0, Max. = 40. Dashed line indicates the rough boundary between α3 and α2α'2 compartments. Scale bar, 20 μm.

(**D**) Stereotyped AZ distribution pattern in PPL1-α3 across individuals. 3D reconstructions show the AZ density across different brain samples. Scale bar, 20 μm.

Figure 5 Cell-type-specific stereotypy of AZ spatial distributions.

(**A**) 3D reconstructions of Brp::rGFP clusters in DPMs, colored by AZ density. Color scale:

Min. = 0, Max. = 35. Black arrows indicate consistently high AZ density regions across

brain samples. Scale bars, 20 μm.

(**B**) 3D reconstructions of Brp::rGFP clusters in APLs, colored by AZ density. Color scale:

Min. = 0, Max. = 30. Black dashed square indicates the area zoomed in. APL

reconstructions are arranged from left to right according to the overall AZ density in α3.

Figure 6 Local intensity analysis revealed sub-compartmental AZ structures.

(**A**) Brp::rGFP correlation analysis between nearest neighbors (NN). The intensity of a

cluster is plotted on the x-axis and the intensity of its nearest neighbor is plotted on the y-

axis. High correlation indicates nearest neighbors have similar Brp::rGFP intensity.

(**B**) Scatter plots showing the Brp::rGFP intensity correlation between nearest neighbors in

a representative sample of γ KCs. Values were normalized to the mean. Pearson correlation

coefficient R is shown.

(**C**) Correlation of Brp::rGFP intensities between nearest neighbor AZs in different cell

types. Brp::rGFP intensities were log transformed. DPM vs. γ KCs: P = 0.0066; DPM vs.

α'/β' KCs: P = 0.0027; DPM vs. α/β KCs: P = 0.0066; Scale bar: 1 μm.

(**D**) 3D reconstructions of Brp::rGFP clusters in a γ KC sample, colored by local intensity

and AZ density. Black arrows indicate areas with both high local intensities and high AZ

densities. Min. and Max. were set to represent the lowest and highest 5% in local intensity

value respectively. Color scale of AZ density: Min. = 0 and Max = 35.

(**E**) 3D reconstructions of Brp::rGFP clusters in a DPM neuron, colored by local intensity

and AZ density. Black triangle arrows indicate areas with high AZ densities but low local

intensities. Min. and Max. are set to represent the lowest and highest 5% in local intensity

value respectively. Color scale of AZ density: Min. = 0, Max. = 35.

(**F**) Correlation analysis between local intensity and AZ density. The local intensity of a

cluster is plotted on the x-axis and its AZ density is plotted on the y-axis.

(**G**) Scatter plots showing the correlation between Brp::rGFP local intensity and AZ density

in a representative sample of γ KCs. Values were normalized to the mean. Pearson

correlation coefficient R is shown.

(**H**) Correlations between AZ density and local intensity in different cell types. DPM (n =

6), APL (n = 5), γ KCs (n = 6, *MB009B-GAL4*), α'/β' KCs (n = 5, *MB370B-GAL4*), α/β KCs

(n = 5, *MB008B-GAL4*). DPM vs. γ KCs: P = 0.0171; DPM vs. α'/β' KCs: P = 0.0171; APL

vs. γ KCs: P = 0.0014; APL vs. α'/β' KCs: P = 0.0014; APL vs. α/β KCs: P = 0.0171.

Figure 7 Associative conditioning re-organizes sub-compartmental active zone

clusters.

(**A**) Brp::rGFP (green) and CD4::tdTomato cytoplasm membrane marker (magenta) in KCs,

visualized by using *R13F02-GAL4*.

(**B**) Single odor conditioning induces long-lasting memory. Left panel, the experimental

design of the aversive single odor conditioning. The paired group receives a concurrent

presentation of 2% 4-MCH and 90 V electric shock. The unpaired group first receives the

electric shock and then 4-MCH 1 min later. Right panel, preference of *Canton S* between

2% 4-MCH and paraffin oil at 3, 10, 30, 90, 270 minutes and 1 day after single odor

conditioning. n = 8 for all groups. Error bars show S.E.M.

(**C**) Correlation coefficient (AZ density vs. local intensity of individual Brp::rGFP clusters)

of each compartment at 90 min after conditioning. Pair (n = 11) vs. Unpair (n = 12), γ2 (P =

0.0261), β2 (P = 0.0060), α1 (P = 0.0026), α'3 (P = 0.0026).

(**D**) Heatmap showing the difference of correlation coefficient (AZ density vs. local

intensity of Brp::rGFP clusters) between the pair and unpair group at different time points

after conditioning. The color indicates the difference and the asterisks in the compartments

indicate the significant difference. For 3 min, Pair (n = 12), Unpair (n = 11); For 20 min,

Pair (n = 12), Unpair (n = 11); For 90 min, Pair (n = 11) vs. Unpair (n = 12), P = 0.0247;

For 270 min, Pair (n = 11), Unpair (n = 11), ; For 1 D, Pair (n = 12), Unpair (n = 9); For all

results involved statistical comparison, only significant results are shown. Scale bars, 20

μm. Mann-Whitney test with original False Discovery Rate method of Benjamini and

Hochberg correction. *P<0.05. Data were presented as box plots showing centre (median),

whiskers (Min. to Max.).

(**E**) Schematics showing learning induced local AZ remodeling. Single odor aversive

training transiently dismisses high Brp level – high AZ density hot spots in specific

compartments of KC terminals.

1017

1018 **Figure supplements**

1019

Figure 1 – figure supplement 1 Split-GFP tagging does not affect Brp neuronal

expression specificity and expression pattern in the brain.

(**A**) Fold change of *brp* mRNA level against reference genes, measured by qRT-PCR.

mRNA of *Ubiquitin-5E* and *αTubulin84B* were used as the reference. n = 3 for all groups.

Error bars show S.E.M. ns = non-significant. ***p ≤ 0.001. Mann-Whitney test.

(**B**) Anti-Brp (nc82) immunostaining signal in the MB of flies with or without GFP_{11}

insertion. Both have expressions of GFP_{1-10} by *R57C10-GAL4*. n = 11 for both groups.

Error bars show S.E.M.

(**C**) Anti-Brp immunostaining of brains of flies with or without pan-neuronal Brp::rGFP

tagging using *R57C10-GAL4*. Different planes of the image stack were shown. Scale bars,

100 μm.

1031

1032

1033 **Figure 1 – figure supplement 2 Neuron-specific assembly of Brp::rGFP.**

1034 Brp::rGFP and CD4::tdTomato were visualized in astrocyte-like glia using *R86E01-GAL4*,

1035 and in neuropeptide releasing neurons using *Amon-GAL4*, respectively. Images were

1036 acquired using comparable settings. Scale bars, 50 μm.

1037

1038

Figure 1 – figure supplement 3 Rab3 knock-down in KCs increases Brp::rGFP

intensity.

Rab3 was knocked-down in KCs using RNAi by *R13F02-GAL4*. The median of Brp::rGFP

intensity of individual clusters was calculated for each γ compartment. n = 6 for both

groups. Scale bars, 50 μm in the overview, 2 μm in the insets.

1044

1045

Figure 1 – figure supplement 4 Effect of iterative Richardson-Lucy image deconvolution.

1046

1047

(**A**) Brp::rGFP clusters of a DPM neuron in raw image, and in images deconvolved for

5~40 iterations. Max projections of the XY and XZ planes were shown. White transparent

lines indicate pixels where the intensity profiles were plotted in B. Numbers denote

different Brp::rGFP clusters. Scale bars, 1 μm.

(**B**) Upper panels show intensity profiles of Brp::rGFP clusters undergone different

iterations of deconvolution. Lower panels show the full width at half maximum (FWHM)

value of intensity profiles.

(**C**) 3D ROIs were generated using deconvolved images and then were used to analyze the

volume and intensity of individual Brp::rGFP clusters in raw images. Scale bars, 1 μm.

1057

1058

Figure 1 – figure supplement 5 Heatmaps of F-scores showing the performance of the pipeline in detecting individual Brp::rGFP clusters in different cell types.

1061 An intensity threshold (threshold) is applied to reduce background noise when detecting 3D

1062 maxima. Noise tolerance adjusts the sensitivity of 3D maxima detection. Color represents

1063 F-score value.

1064

R13F02>Brp::GFP$_{11}$, UAS-GFP$_{1-10}$

R13F02>UAS-GFP$_{1-10}$

1065

1066 **Figure 7 – figure supplement 1 Gross anatomy of KC terminals is not affected by**

1067 **Brp::rGFP tagging.**

1068 Morphology of KC terminals in flies with or without GFP$_{11}$ insertion. Both groups express

1069 GFP$_{1-10}$ and CD4::tdTomato, driven by *R13F02-GAL4*. Scale bars, 20 μm.

1070

1071

Figure 7 – figure supplement 2 Brp::rGFP tagging in KCs does not affect short-term memory performance.

10% 4-MCH or 10% 3-OCT was paired with 90 V electric shocks. Odor preference was tested immediately after conditioning. GFP reconstitution was induced in KCs using *R13F02-GAL4*. *R13F02>Brp::rGFP*, n = 10; *Canton S*, n = 8. ns = non-significant. Welch's t test.

1078

Supplementary files

Supplementary file 1

Python code used for nearest neighbor, local intensity and AZ density analysis.

Supplementary file 2

Example data of a DPM neuron generated by the 3D spot segmentation, before being processed by the analysis code.