TITLE

Octopamine signals coordinate the spatial pattern of presynaptic machineries within individual

Drosophila mushroom body neurons

AUTHORS

Hongyang Wu[1,6], Sayaka Eno[1,3,6], Kyoko Jinnai[2], Yoh Maekawa[3], Kokoro Saito[1], Ayako Abe[1], Darren

W Williams[4], Nobuhiro Yamagata[1,5], Shu Kondo[2], Hiromu Tanimoto[1,3,7*]

AFFILIATIONS

[1] Graduate School of Life Sciences, Tohoku University, Sendai, Japan.

[2] Department of Biological Science and Technology, Tokyo University of Science, Tokyo, Japan.

[3] Department of Biology, Faculty of Science, Tohoku University, Sendai, Japan.

[4] Centre of Developmental Neurobiology, Institute of Psychiatry, Psychology & Neuroscience, King's

College London, London, United Kingdom.

[5] Department of Life Science, Graduate School of Engineering Science, Akita University, Akita, Japan.

[6] These authors contributed equally.

[7] Lead contact.

*Correspondence: hiromut@m.tohoku.ac.jp (H.T.)

SUMMARY

Neurons need to adjust synaptic output according to the targets. However, the target-specific synaptic structures within individual neurons in the central nervous system remains unresolved. Applying the CRISPR/Cas9-mediated split-GFP tagging, we visualized the endogenous active zone scaffold protein, Bruchpilot (Brp), in specific cells. This technology enabled the spatial characterization of presynaptic machineries only within the Kenyon cells (KCs) of the *Drosophila* mushroom bodies. We found the patterned accumulation of Brp among the compartments of axon terminals, where a KC synapses onto different postsynaptic neurons. Mechanistically, the localized octopaminergic modulation along γ KC terminals regulate this compartmental Brp heterogeneity via Octβ2R and cAMP signaling. We further found that acute food deprivation reorganizes this spatial pattern in an octopamine-dependent manner. Such coordinated regulation of local synaptic machineries thus explains how the mushroom bodies integrate changing physiological states. This subcellular information processing represents an elegant solution to expand computational capacity of the circuit.

KEYWORDS:

Mushroom bodies; Kenyon cells; Bruchpilot; active zones; intracellular synaptic heterogeneity; target-specific synaptic structure; state-dependent synaptic plasticity; octopamine; cAMP signaling

INTRODUCTION

The molecular composition of chemical synapses determines the function of neurons and therefore exhibits a remarkable diversity among cell types [1,2]. As demonstrated by studies on the motor neuron, presynaptic molecular assemblies can be highly heterogeneous even within a single cell [3,4]. Critically, the difference of presynaptic structures may result in varying synaptic functions at the single active zone

(AZ) level, such as release probability [5–7]. Given that a single neuron in the central nervous system

(CNS) typically synapses onto multiple target cells, the spatial adaptation of output machineries could

differentiate the activities of post-synaptic cells. While such intracellular synaptic heterogeneity

influences complex computation of the circuit in the CNS, the target-specific AZ regulation is scarcely

studied.

To this end, neurons projecting to the *Drosophila* mushroom bodies (MBs) can serve as an

excellent model system. Kenyon cells (KCs), the major MB intrinsic neurons, synapse onto five sets of

post-synaptic partners in spatially segregated compartments [8,9]. Postsynaptic MB output neurons

(MBONs) have distinct and compartmentalized dendrites within the MB lobes [9], and their activities

collectively determine the MB output [10–14]. The compartmental distinctions in presynaptic assemblies

within individual KCs are likely to be critical in controlling MB-guided behaviors. However, it is

challenging to characterize AZs only in KCs without 3D electron microscopy, because the synapse

density in MB lobes is among the highest in the fly brain [15,16].

We here aim to characterize the compartmental distinctions of presynaptic structures of KCs by

visualizing endogenous Brp in a cell-type specific manner. *Drosophila* ELKS/CAST/ERC family

member Brp is a major scaffold protein forming the T-shape electron-dense projections decorating AZs

[17–20]. Brp clusters determine molecular assemblies at AZs such as the accumulation of calcium channels

and synaptic vesicles [17,21]. Therefore, Brp structures serve as a proxy for estimating the synapse function.

To label endogenous pre-synaptic proteins in designated cell types [22,23], we took advantage of the

CRISPR/Cas9-mediated split-GFP tagging strategy to target Brp in the dense network of MBs [24,25].

Profiling the character of individual Brp clusters only within KCs revealed the compartmental Brp

accumulation pattern along the MB lobes. We further report the state-dependent remodeling of this

intracellularly organized AZ structure and the regulatory mechanisms.

RESULTS

Cell-type specific visualization of endogenous active zone scaffold proteins Brp

Chemical tagging using Brp::SNAP [26] suggests endogenous Brp localizes heterogeneously within the MB (Fig. S1), which is constituted by pre-synapses of various cell types (Fig. S2). To visualize endogenous Brp only in designated cell types, we inserted the GFP_{11} fragment (the eleventh β-strand of the super-folder GFP) just prior to the stop codon of *brp* using CRISPR/Cas9. The self-assembly and reconstitution of GFP is induced by expressing the GFP_{1-10} fragments using the GAL4/UAS system (Fig. 1A).

We validated this system with GAL4 driver lines that label 7 different MB-projecting neuronal populations: 3 KC subtypes, 2 dopamine neuron clusters (PPL1 and PAM) and 2 single interneurons (DPM and APL neurons) [9,27]. Confocal microscopy detected reconstituted GFP (rGFP) signals only at restricted areas, in contrast to the dense signals in Brp::SNAP or Anti-Brp immunostaining (Fig. 1B, Fig. S1). The Brp::rGFP signal appeared as individual puncta and matched the areas of axon terminals of the cell types (Fig. 1B). Especially, Brp can be labeled at single-cell resolution for the APL and DPM neuron [8,28,29]. Strikingly, the Brp::rGFP signal intensity appeared to be heterogenous along the axon terminals of KCs (Fig. 1B). Since individual KCs arborize onto the entire lobe structure, these results suggest distinct accumulations of Brp intracellularly.

To test whether Brp::rGFP labels dynamic synaptic structures, we benchmarked its performance with different live-imaging contexts. The axon arborizations of pleural muscle motor neurons (PM-Mns) can be imaged on the abdominal body wall during pupal development [30]. Live-imaging Brp::rGFP in the PM-Mn revealed the formation of individual clusters at 48 hours after puparium formation and the increasing density in the same branch following two days (Fig. 1C). We further visualized Brp::rGFP

dynamics in the ex-vivo preparation of the larval brain, and found multiple fusion and fission events of

Brp clusters in the MB calyx (Fig. 1D). These results are consistent with the AZ assembly through

liquid-liquid phase separation [31] and validate the method to visualize the dynamics of endogenous Brp

structures.

Intracellular active zone heterogeneity among MB compartments

Brp::rGFP accumulation was heterogeneous among KC compartments, especially in γ KCs (Fig. 1B, 2A,

Fig. S3). To characterize this presynaptic Brp heterogeneity, an image-processing pipeline was

developed to better quantify the signal intensity of individual Brp::rGFP puncta. Optimizing the

parameter combination of 3D image deconvolution [32,33] and 3D spot segmentation enabled us to resolve

Brp::rGFP clusters that represent single AZs using conventional confocal microscopy [29]. By sampling a

volume of each compartment, we quantified the median of integrated Brp::rGFP intensity of clusters at

the AZs. Therefore, the quantification is independent of the AZ density or the sample volume. The

analysis revealed a striking compartmental heterogeneity of AZs (Fig. 2B). The median Brp::rGFP

intensity of clusters in the γ5 compartment was nearly three times as intense as that in γ1. These results

suggest that individual KCs tune their presynaptic structures according to compartments (Fig. 2C).

In motor neurons, Brp is well characterized to serve as a scaffold protein accumulating synapse

proteins such as voltage-gated Ca^{2+} channels [17,19,34]. Consistently, we found that the endogenous α

subunit of voltage-gated Ca^{2+} channel, Cacophony (Cac) co-localizes with Brp in the γ lobe (Fig. S4,

S5). Given the compartmental Brp heterogeneity (Fig. 2B-C), this correlation suggests that the Ca^{2+}

concentration is differentially set along the γ KCs. To measure the basal Ca^{2+} concentration nearby AZs,

we expressed Brp^{short}::mCherry::GCaMP6s, a ratiometric Ca^{2+} sensor fused to mCherry and truncated

Brp [35], in KCs. We live-imaged the MB and found that basal GCaMP signals were compartmentally

distinct and gradually increased towards the γ5 compartment (Fig. 2D). This compartmental pattern of

GCaMP signals nicely mirrored the Brp heterogeneity. Taken together the colocalization of Brp and Cac,

these results suggest that the Brp accumulation per AZ is set compartmentally, differentiating the basal

Ca^{2+} influx via Ca^{2+} channels at AZs [36].

The state-dependent regulation of Brp compartmental heterogeneity

As different MB compartments are functionally coordinated and integrate internal states such as

nutritional states, sleep need and aging [10,37–41], we examined whether KCs adapt the synaptic structures

upon physiological changes. We calculated the variance of the compartments' log-transformed medians

to measure the heterogeneity level. Therefore, the absolute Brp::rGFP intensity does not affect the

variance (Fig. 3A). Strikingly, the compartmental heterogeneity in γ KCs significantly decreased upon

food deprivation for 48 hours and recovered after refeeding (Fig. 3B, Fig. S8A). In contrast, we did not

find a significant effect by aging for 30 d (Fig. S6). These results suggest that acute stress can drive local

adjustment of AZ structures within KCs, reflecting reorganized compartmental activities.

Octopamine input underlies the Brp compartmental heterogeneity

Each compartment receives dopamine projections from structurally and functionally distinct subsets of

dopaminergic neurons (Fig. S7A) [9]. We therefore examined whether dopamine inputs determine the

compartmental Brp accumulation by knocking-down dopamine receptors (DopR1, DopR2 and D2R) in

KCs using RNA interference (RNAi) [42,43]. None of these disruptions showed a significant effect on the

compartmental heterogeneity (Fig. S7B, Fig. S8B).

The octopaminergic neurons (OANs), particularly OA-VPM3 and OA-VPM4 neurons, are

known to project to the MB and concentrate their innervations in the γ1 compartment [9,44]. We

characterized presynaptic components of OA-VPM3/4 visualized by Brp::rGFP and nSyb::CLIP

(neuronal synaptobrevin, nSyb; synaptic vesicle marker) and found they were densely localized to γ1

compared to the other compartments, complementary to the Brp compartmental heterogeneity of KCs

(Fig. 4A-4D). These results suggest localized synaptic output of OANs along the γ lobe, potentially

underlying the Brp compartmental heterogeneity.

To examine the contribution of octopamine to the Brp accumulation within KCs, we measured

the Brp compartmental heterogeneity in the mutants of tyramine β-hydroxylase (*Tβh*), the enzyme that

catalyzes the synthesis of octopamine from tyramine [45]. We generated two new *Tβh* null alleles:

insertion mutant *Tβh^{SK1-8}* and deletion mutant *Tβh^{SK2-4}* using CRISPR/Cas9-mediated mutagenesis. We

visualized Brp::rGFP in KCs of these mutants and found the significantly decreased compartmental

heterogeneity of the γ KCs (Fig. 4E, Fig. S8C). This suggests that octopamine directly controls the AZ

structure in KCs.

To identify the signaling pathway that regulates the Brp compartmental heterogeneity, we

knocked-down octopamine/tyramine receptors (including OAMB, Octα2R, Octβ1R, Octβ2R, Octβ3R,

OctTyrR, TyrR, TyrRII) in KCs using RNAi [42,46]. Among all the receptors tested, only Octβ2R

knockdown significantly decreased the compartmental heterogeneity (Fig. 5A, Fig. S8D-E). We could

not test the effect of Octβ1R and Octβ3R because their deficiency in KCs caused abnormal development

of the MB. Octβ2R is coupled to the Gs alpha subunit to stimulate cyclic adenosine monophosphate

(cAMP) production [47]. To test whether cAMP plays a role in Brp accumulation, we knocked-down the

adenylate cyclase Rutabaga (Rut) in KCs using RNAi [42,48–52]. The downregulation of Rut resulted in a

significant decrease of the compartmental heterogeneity (Fig. 5B, Fig. S8F), suggesting that octopamine

functions through Octβ2R-cAMP pathway.

Nutritional states adjust the Brp compartmental heterogeneity through octopamine signaling

The spontaneous firing of octopamine neurons was shown to be decreased upon starvation [53]. Therefore,

we hypothesized state-dependent AZ remodeling through octopamine (Fig. 6A). To test this hypothesis,

we starved the *Tβh* mutants for 48 hours and measured the Brp::rGFP heterogeneity in γ KCs. Indeed,

there was no significant state-dependent effect on the Brp compartmental heterogeneity in the *Tβh*

mutants in contrast to the control experiment (Fig. 6B, Fig. S8G-H). Taken together, we propose that

internal states adjust the synaptic output of MB compartments through octopamine signaling that

supervise the synaptic structures within KCs.

DISCUSSION

By labeling endogenous AZ scaffold protein Brp, we resolved individual AZs at the single-cell level.

The substantial intracellular heterogeneity of Brp accumulation within γ KCs is found to be regulated by

internal states of the animal. Other than the compartmental pattern of Brp clusters within KCs, our

experimental pipeline enables further profiling of other AZ characteristics such as the local AZ density

[29].

How does octopaminergic signaling modify Brp accumulation? This study identified the requirement of

Octβ2R and Rut in regulating the AZ structure in KCs (Fig. 5). Octβ2R was shown to stimulate cAMP

synthesis [47], suggesting that cAMP could play a determining role. Consistent with this idea, *rut* mutant

displays abnormal AZ structures [54]. We propose that the localized innervations of octopamine neurons

onto KCs (Fig. 4) results in different cAMP concentrations along the γ lobe. In both KCs and motor

neurons, cAMP concentration was shown to be locally regulated [55–57]. The affinity or stability of Brp to

AZs among different compartments is perhaps distinctly set by the cAMP signaling compartmentation

[58]. Interestingly, both DopR1 and Octβ2R stimulate cAMP biosynthesis, while the knock-down of DopR1 had no significant effect on the Brp heterogeneity (Fig. S7). This may be due to different downstream targets of the receptors including PKA [59].

AZ structures, especially Brp clusters, frequently serve as a proxy to explain synaptic functions [5–7,17,19,34,60]. Consistent with previous studies, our data suggest that Brp regulates the localization of Ca^{2+} channels in KCs [17,19,61]. The Brp heterogeneity may result in compartmental basal Ca^{2+} concentrations at AZs (Fig. 2D) and thus modulate spontaneous synaptic vesicle release via the stochastic activity of Ca^{2+} channels [36]. Indeed, the enrichment of the Ca^{2+} channel is associated to higher spontaneous release frequency in *Drosophila* NMJ [7,62]. Although the relationship between Brp accumulation and evoked release in KCs seems more complicated since it involves acute and compartmental dopaminergic modulations [13,63–65]. Unlike evoked activities, spontaneous release of synaptic vesicle persists in the absence of stimulation thereby affecting the states of the animal [10].

The Brp compartmental heterogeneity within KCs might directly modulate the MB output through controlling the activity of different MBONs. Each type of MBONs serves as an independent output unit, together regulating a variety of behaviors [11–14,40,66–68]. Starvation changes the activity balance between MBON-γ1 and MBON-γ5 thus enhances appetitive memory expression [66,68,69], and MBON- γ5 and MBON- γ2 oppositely control sleep amount [70]. Consistently, we found that the compartmental AZ heterogeneity within KCs, measured by Brp, decreased upon starvation (Fig. 3B). Considering the associations between octopamine and feeding behaviors [53,71–77], KCs generates behavior adaptations in response to changing internal states and environments by local AZ tuning [40,78,79].

200 **STAR METHOD**

201 **KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-Brp	DSHB	Cat#nc82; RRID: AB_2314866
AlexaFluor-568 goat anti-mouse	Invitrogen	Cat#A11004; RRID: AB_2534072
Chemicals		
SNAP-Surface Alexa Fluor 647	New England Biolabs	Cat#S9136S
CLIP-Surface Alexa Fluor 647	New England Biolabs	Cat#S9234S
OMNIPAQUE 350	GE Healthcare	Cat#22000AMX02447
Histodenz	Sigma	Cat#D2158
Tetraspeck Microspheres 0.1 mm	Thermo Fisher Scientific	Cat#T7279
NaCl	Nacalai Tesque	Cat#31320-05
KCl	Nacalai Tesque	Cat#28514-75
TES	Nacalai Tesque	Cat#329810-55
Trehalose dihydrate	Tokyo Kasei Kogyo	Cat#T0331
D-glucose	Nacalai Tesque	Cat#16806-25
$NaHCO_3$	Nacalai Tesque	Cat#31213-15
NaH_2PO_4	Merck KGaA	Cat#106346
$CaCl_2$	Sigma-Aldrich	Cat#C5080
$MgCl_2$	Sigma-Aldrich	Cat#M2670
Drosophila strains		
Brp::SNAP [26]	Bloomington Drosophila stock center (BDSC)	BDSC 58397
MB008B-GAL4 [9]	BDSC	BDSC 68291
MB009B-GAL4 [9]	BDSC	BDSC 68292
MB370B-GAL4 [9]	BDSC	BDSC 68319
VT64246-GAL4 [80]	VDRC	VDRC 204311
GH146-GAL4 [81]	BDSC	BDSC 30026
TH-GAL4 [82]	BDSC	BDSC 8848
R58E02-GAL4 [83]	BDSC	BDSC 41347
R13F02-GAL4 [83]	BDSC	BDSC 48571
OK371-GAL4 [84]	BDSC	BDSC 26160
UAS-Brp::mCherry::GCaMP6s [35]	BDSC	BDSC 77131
Cac::sfGFP	Gratz *et al.* [6]	N/A

Tdc2-GAL4 [85]	BDSC	BDSC 9313
MB022B-GAL4 [9]	BDSC	BDSC 68298
UAS-nSyn::CLIP [26]	BDSC	BDSC 58385
UAS-DopR1-RNAi [52]	BDSC	BDSC 55239
UAS-Dop2R-RNAi [52]	BDSC	BDSC 50621
UAS-DopR2-RNAi [52]	BDSC	BDSC 51423
UAS-OAMB-RNAi [46]	VDRC	VDRC 2861
UAS-Octα2R-RNAi [52]	BDSC	BDSC 50678
UAS-Octβ1R-RNAi [52]	BDSC	BDSC 58179
UAS-Octβ2R-RNAi [52]	BDSC	BDSC 34673
UAS-Octβ3R-RNAi [52]	BDSC	BDSC 62283
UAS-TyrR-RNAi [52]	BDSC	BDSC 25857
UAS-TyrR2-RNAi [52]	BDSC	BDSC 64964
UAS-OctTyrR-RNAi [52]	BDSC	BDSC 28332
UAS-Rut-RNAi [52]	BDSC	BDSC 80468
UAS-CD4::tdTomato	BDSC	BDSC 35841
UAS-CD8::GFP	BDSC	BDSC 32194
$T\beta h^{SK1-8}$	This study	N/A
$T\beta h^{SK2-4}$	This study	N/A
$brp::GFP_{11}$	This study	N/A
UAS-GFP$_{1-10}$	Kondo et al. [25]	N/A
Software		
Python	N/A	N/A
Graphpad PRISM	Graphpad software	https://www.graphpad.com/
Fiji/Image J	NIH	https://imagej.net/software/fiji/downloads
Amira Software	Thermo Fisher Scientific	thermofisher.com/amira-avizo
DeconvolutionLab2	Sage et al. [86]	https://bigwww.epfl.ch/deconvolution/deconvolutionlab2/
CLIJ2	Haase et al. [87]	https://clij.github.io/
3D ImageJ Suite	Ollion et al. [88]	https://imagej.net/plugins/3d-imagej-suite/

RESCOURCE AVAILIBILITY

Lead contact

205 Further information and requests on resources and reagents should be directed to and will be fulfilled by

206 the lead contact, Dr. Hiromu Tanimoto (hiromut@m.tohoku.ac.jp).

Material availability

209 All reagents generated in this study are available upon request from the lead contact.

Data and code availability:

212 This study did not generate any unique code. All data needed to evaluate the conclusions in the paper are

213 available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Animal husbandry

Flies were maintained on standard cornmeal food at 25 °C under a 12:12 hours light-dark cycle for all

experiments. All flies used for experiments are 3-7 days old adult males. After hatching, adult flies were

transferred to fresh food vials and flip every 2-3 days before experiments. In experiments where flies

were fasted, the same population of flies was separated before the experiment to ensure the

homogeneity. All flies were food-deprived for 48 hours for the experimental group, while provide

normal food supply for the control for the same time. The GAL4-UAS system was used to express

transgenes. For each transgenic control in RNAi experiments, crossing respective transgenic line with

Canton S strain, w1118, or y1w1118 to replace the balancer chromosomes. Flies for wild-type

characterization were also crossed to replace balancer chromosomes. Strains used in this study are as

indicated in the KEY RESOURCES TABLE.

METHOD DETAIL

Transgenic lines and mutants

Two independent null alleles of the *Tβh* gene were obtained by inducing frameshift indels by the

transgenic CRISPR/Cas9 system as previously described [89]. The following 20-bp gRNA sequences were

used:

gRNA#1: GATACGTACACCAGTCCGGA

gRNA#2: GGTGAGACGGGACTACCAGC

   Tβh^{SK1-8} was derived from mutagenesis by gRNA#1 and carries the following frameshift

insertion: CGTACACCAGgatggacactGGATGGACAG (the inserted sequence is indicated by lower-

case characters). *Tβh^{SK2-4-4}* was derived from mutagenesis by gRNA#2 and carries the following

frameshift deletion: TCCGGATGGA----------------------------CTGTGAGGTC (the gap is indicated as

hyphens). The *brp::GFP_{11}* line was generated by targeted insertion of a GFP_{11} cassette into the

endogenous *brp* locus just prior to the stop codon as previously described [25].

Sample preparation

Animals used in experiments involving comparisons between control and experimental groups were

dissected on the same day. Data were collected from multiple batches of experiments performed on

different days. All steps were performed at room temperature otherwise specified. Flies were

anesthetized on ice and placed on ice before dissection. The dissection was performed in ice PBS

solution to collect the brain without the ventral nerve cord. Brains are subsequently fixed by 2%

paraformaldehyde for 1 hour then washed 3×20 min with 0.1% PBT (0.1% Triton X-100 in PBS) in a

PCR tube, typically containing 5-6 brains.

For SNAP or CLIP chemical tagging, brains were incubated in SNAP-Surface 647 (1:1000; NEB; S9136S) or CLIP-Surface 647 (1:1000; NEB; S9234S) in 0.3% PBT for 15 min and washed with 3×20 min with 0.1% PBT. Samples are mounted on microscope slides using the appropriate mounting medium (SeeDB2S, SeeDB2G [90] or 86% glycerol according to the imaging condition). Brp::SNAP, Brp::rGFP, Cac::sfGFP, CD4::tdTomato and CD8::GFP samples were imaged without immunohistochemistry.

Immunohistochemistry procedures (for anti-Brp staining in Fig. 1B) were carried out as described [91]. In brief, fixed brains were washed 3×20 min with 0.1% PBT and incubated in 3% normal goat serum (NGS; Sigma-Aldrich; G9023) for 1 hour at room temperature. Antibodies used were as indicated in resource table (Table S1). Brains were incubated in antibody solutions at 4°C for 48 hours for both primary and secondary antibodies. Sub-resolution fluorescent beads (Tetraspeck Microspheres 0.1 mm, Thermo Fisher Scientific, T7279) were imaged for generating experimental point spread function (PSF). The beads solution was diluted with distilled water and sonicated multiple times to eliminate aggregation.

For *in vivo* imaging of PM-Mns in developing pupae, samples were prepared as previously described [30]. In brief, white prepupal stage pupae (~ 0 hour APF) were collected and incubated at 25 °C until the experiment. Before imaging, the puparium case was carefully removed with forceps, and the pupa was transferred to an imaging chamber sealed with a cover slip to create an imaging window on the abdomen. For ex vivo imaging of 3rd instar pupal CNS, the CNS was dissected directly from the animal in PBS at room temperature. The CNS sample was then mounted with a spacer between the microscopic glass and cover slip with PBS, The entire process, from dissection to imaging, was performed within 5 minutes.

Image acquisition

For fixed samples and *ex vivo* imaging, images are acquired using the Olympus FV1200 confocal microscope platform equipped with GaAsP high sensitivity detectors and a 60×/1.42 NA oil immersion objective (PLAPON60XO, Olympus) and a 30×/1.05 NA silicone immersion objective (UPLSAPO30XS, Olympus). For *in vivo* imaging on pupae, images were acquired using the Zeiss LSM 800 series confocal microscopy equipped with a 40×/1.3 NA objective (Plan-Apochromat 40×/1.3 Oil DIC (UV) VIS-IR M27).

For Brp::SNAP chemical tagging, 30×/1.05 NA objective is used with a voxel size of 0.53 µm × 0.53 µm × 0.84 µm (lateral × lateral × axial) to image the whole brain except for the optical lobes. For Brp::rGFP and UAS-CD4::tdTomato samples in all GAL4 types except Tdc2-GAL4, 60×/1.42 NA oil immersion objective was used and 473 nm laser power: 0.1%; 559 nm laser power: 0.1%; scanning speed: 2 µs/pixel with a voxel size of lateral 0.079 × 0.079 × axial 0.370 µm. For Brp::SNAP/Cac::sfGFP images, 60×/1.42 NA objective is used with a voxel size of 0.079 × 0.079 × 0.370 µm. For Tdc2-GAL4 driven UAS4-CD8::GFP and UAS-nSyb::CLIP samples, 30×/1.05 NA objective is used with a voxel size of 0.414 × 0.414 × 0.84 µm. For MB022B-GAL driven UAS-CD8::GFP and UAS-nSyb-CLIP samples, 30×/1.05 NA objective is used with a voxel size of 0.276 × 0.276 × 0.87 µm. For Tdc2-GAL4 driven Brp::rGFP and UAS-CD4::tdTomato, 60×/1.42 NA objective is used with a scanning voxel size of 0.132 × 0.132 × 0.34 µm. For Brp::rGFP *in vivo* imaging of PM-Mns, a 40×/1.3 NA objective is used with a voxel size of 0.0725 × 0.0725 × 0.460 µm. For Brp::rGFP ex vivo imaging, 60×/1.42 NA objective is used with a voxel size of 0.079 × 0.079 × 0.370 µm and 473 nm laser power: 2.0%. For experimental PSF imaging, SeeDB2S immerged beads were scanned in a setting that is 60×/1.42 NA oil immersion objective; 473 nm laser power: 2.0%; 559 nm laser power:

1.5%; voxel size: 0.079 μm × 0.079 μm × 0.370 μm; scanning speed: 4 μs/pixel, to produce multiple

bead images.

***In vivo* calcium imaging**

In vivo calcium imaging was performed following a previously described protocol [91]. Briefly, flies were

anesthetized on ice for 3 minutes and placed in a custom-made holding dish on a Peltier plate (CP-085,

Sinics) maintained at 5°C. The head capsule was fixed to the dish using UV curing optical adhesive

(NOA68, Thorlabs). To minimize brain movement, the proboscis was glued to the capsule A small

window was opened on the top of the head capsule, and the exposed area was filled with *Drosophila*

saline solution (final concentration: 103 mM NaCl, 3 mM KCl, 5 mM TES, 8 mM Trehalose dihydrate,

10 mM D-glucose, 26 mM $NaHCO_3$, 1 mM NaH_2PO_4, 1.5 mM $CaCl_2$, 4 mM $MgCl_2$, adjust to pH ~7.2).

Air sacs and fat bodies covering the brain surface were carefully removed. Live imaging was conducted

using a laser scanning confocal microscope equipped with GaAsP detectors (A1R, Nikon) and a 30×/1.1

NA water immersion objective (Apo LWD 25×, Nikon). GCaMP6s and mCherry were sequentially

excited at 488 and 561 nm, respectively. Emission light was collected using dichroic mirrors and

emission filters (BP500–550 and BP570-620) onto GaAsP detectors. The MB γ lobe was scanned at a

pixel size of 0.5μm/pixel (512 × 128 pixels) at a frame rate of 1 Hz using line scans with 16× averages

in the resonant scanning mode. The pinhole was set to 2.5 AU (561 nm). For recording, images were

sequentially acquired over 40 s and saved for later image processing.

Data processing and analysis

PSF images were processed in the Amira software (Thermo Fisher Scientific). Using the Extract Point

Spread Function module, PSFs extracted from each image were averaged into a single PSF, which was

later used for image deconvolution (resized voxel size: 0.079 µm × 0.079 µm × 0.370 µm; image size:

32 pixels × 32 pixels × 21 slices). Image deconvolution was performed using the Richardson-Lucy

iterative non-blind algorithm in the Fiji plugin DeconvolutionLab2 [86], or with CLIJ2 GPU-based

Richardson-Lucy deconvolution [87].

AZ detection was performed as described [29] in a compartmental manner. In brief, a sub-image

stacks were cropped from every MB compartment in the original image. Compartments were manually

identified referencing the CD4::tdTomato membrane signal. An intensity threshold was applied to

reduce background noise. Sub-stacks were processed by the 3D maxima finder and the 3D spot

segmentation function in the 3D suite plugin [88] in Fiji. 3D maxima were detected for each Brp::rGFP

cluster and used as start points for pixel clustering using the 3D spot segmentation function. This

process created 3D ROIs enclosing individual Brp clusters. ROIs were then used to extract Brp::rGFP

signal intensities independently for each cluster. Detection precision was optimized by comparing

detection results with manually defined ground truths as previously described [29].

For calcium imaging, ROIs were manually drawn for each compartment in time-series image

stacks. GCaMP6s signal was normalized by mCherry in each time-frame using the image calculator in

Fiji. The mean intensity was calculated in each compartment by averaging 7 frames without any odor

stimulation as the basal level.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analysis

Statistical analyses were performed using GraphPad Prism version 8, 9 and 10. Summarized data are

represented as box plots showing center (median), whiskers (Min. to Max.). For comparison between

more than two groups, Kruskal-Wallis test with original False Discovery Rate (FDR) method of

Benjamini and Hochberg correction was applied. Desired false discovery rate was set to 0.05. For

comparison between two groups, unpaired nonparametric two-sided Mann-Whitney U-test was used.

Pearson's correlation coefficient (R) was calculated in Python.

ACKNOWLEDGMENTS

We thank all lab members of Tanimoto Lab at Tohoku University for valuable discussions. We thank

Dr. Kate O'Connor-Giles (Brown University), Vienna Drosophila Resource Center (VDRC) and

Bloomington Drosophila Stock Center (BDRC) for transgenic flies, Ayano Wu (Tohoku University) for

graphical design.

FUNDING:

Ministry of Education, Culture, Sports, Science and Technology 22H05481 (HT)

Ministry of Education, Culture, Sports, Science and Technology 22KK0106 (HT)

Ministry of Education, Culture, Sports, Science and Technology 20H00519 (HT)

Japan Science and Technology Agency JPMJSP2114 (HW)

Tohoku University Research Program 'Frontier Research in Duo' (HT)

AUTHOR CONTRIBUTIONS:

Conceptualization, HW, NY and HT

Investigation, HW, SE, KS, AA, KJ

Methodology, HW, NY, SE, SK and YM

Writing – Original Draft, HW, SE and HT

Writing – Review & Editing, all authors

Funding Acquisition, HW and HT

Resources, KJ and SK

Supervision, DW, NY, SK and HT

COMPETING INTERESTS:

Authors declare that they have no competing interests.

REFERENCES

1. O'Rourke, N.A., Weiler, N.C., Micheva, K.D., and Smith, S.J. (2012). Deep molecular diversity of
mammalian synapses: why it matters and how to measure it. Nat Rev Neurosci *13*, 365–379.
https://doi.org/10.1038/nrn3170.

2. Cizeron, M., Qiu, Z., Koniaris, B., Gokhale, R., Komiyama, N.H., Fransén, E., and Grant, S.G.N.
(2020). A brainwide atlas of synapses across the mouse life span. Science *369*, 270–275.
https://doi.org/10.1126/science.aba3163.

3. Ehmann, N., Linde, S. van de, Alon, A., Ljaschenko, D., Keung, X.Z., Holm, T., Rings, A.,
DiAntonio, A., Hallermann, S., Ashery, U., et al. (2014). Quantitative super-resolution imaging of
Bruchpilot distinguishes active zone states. Nat Commun *5*, 4650. https://doi.org/10.1038/ncomms5650.

4. Paul, M.M., Pauli, M., Ehmann, N., Hallermann, S., Sauer, M., Kittel, R.J., and Heckmann, M. (2015).
Bruchpilot and Synaptotagmin collaborate to drive rapid glutamate release and active zone
differentiation. Front Cell Neurosci *9*, 29. https://doi.org/10.3389/fncel.2015.00029.

5. Akbergenova, Y., Cunningham, K.L., Zhang, Y.V., Weiss, S., and Littleton, J.T. (2018).
Characterization of developmental and molecular factors underlying release heterogeneity at Drosophila
synapses. Elife *7*, e38268. https://doi.org/10.7554/elife.38268.

6. Gratz, S.J., Goel, P., Bruckner, J.J., Hernandez, R.X., Khateeb, K., Macleod, G.T., Dickman, D., and
O'Connor-Giles, K.M. (2019). Endogenous tagging reveals differential regulation of Ca2+ channels at
single AZs during presynaptic homeostatic potentiation and depression. J Neurosci *39*, 3068–18.
https://doi.org/10.1523/jneurosci.3068-18.2019.

7. Newman, Z.L., Bakshinskaya, D., Schultz, R., Kenny, S.J., Moon, S., Aghi, K., Stanley, C., Marnani,
N., Li, R., Bleier, J., et al. (2022). Determinants of synapse diversity revealed by super-resolution
quantal transmission and active zone imaging. Nat Commun *13*, 229. https://doi.org/10.1038/s41467-
021-27815-2.

8. Tanaka, N.K., Tanimoto, H., and Ito, K. (2008). Neuronal assemblies of the Drosophila mushroom
body. J. Comp. Neurol. *508*, 711–755. https://doi.org/10.1002/cne.21692.

9. Aso, Y., Hattori, D., Yu, Y., Johnston, R.M., Iyer, N.A., Ngo, T.-T., Dionne, H., Abbott, L., Axel, R.,
Tanimoto, H., et al. (2014). The neuronal architecture of the mushroom body provides a logic for
associative learning. Elife *3*, e04577. https://doi.org/10.7554/elife.04577.

10. Suárez-Grimalt, R., Kadow, I.C.G., and Scheunemann, L. (2024). An integrative sensor of body
states: how the mushroom body modulates behavior depending on physiological context. Learn. Mem.
31, a053918. https://doi.org/10.1101/lm.053918.124.

11. Aso, Y., Sitaraman, D., Ichinose, T., Kaun, K.R., Vogt, K., Belliart-Guérin, G., Plaçais, P.-Y., Robie,
A.A., Yamagata, N., Schnaitmann, C., et al. (2014). Mushroom body output neurons encode valence and
guide memory-based action selection in Drosophila. eLife *3*, e04580. https://doi.org/10.7554/elife.04580.

12. Ichinose, T., Kanno, M., Wu, H., Yamagata, N., Sun, H., Abe, A., and Tanimoto, H. (2021).
Mushroom body output differentiates memory processes and distinct memory-guided behaviors. Curr
Biol *31*, 1294-1302.e4. https://doi.org/10.1016/j.cub.2020.12.032.

13. Cohn, R., Morantte, I., and Ruta, V. (2015). Coordinated and Compartmentalized Neuromodulation
Shapes Sensory Processing in Drosophila. Cell *163*, 1742–1755.
https://doi.org/10.1016/j.cell.2015.11.019.

14. Owald, D., Felsenberg, J., Talbot, C.B., Das, G., Perisse, E., Huetteroth, W., and Waddell, S. (2015).
Activity of Defined Mushroom Body Output Neurons Underlies Learned Olfactory Behavior in
Drosophila. Neuron *86*, 417–427. https://doi.org/10.1016/j.neuron.2015.03.025.

15. Scheffer, L.K., Xu, C.S., Januszewski, M., Lu, Z., Takemura, S., Hayworth, K.J., Huang, G.B.,
Shinomiya, K., Maitlin-Shepard, J., Berg, S., et al. (2020). A connectome and analysis of the adult
Drosophila central brain. Elife *9*, e57443. https://doi.org/10.7554/elife.57443.

16. Zheng, Z., Lauritzen, J.S., Perlman, E., Robinson, C.G., Nichols, M., Milkie, D., Torrens, O., Price,
J., Fisher, C.B., Sharifi, N., et al. (2018). A Complete Electron Microscopy Volume of the Brain of
Adult Drosophila melanogaster. Cell *174*, 730-743.e22. https://doi.org/10.1016/j.cell.2018.06.019.

17. Kittel, R.J., Wichmann, C., Rasse, T.M., Fouquet, W., Schmidt, M., Schmid, A., Wagh, D.A., Pawlu,
C., Kellner, R.R., Willig, K.I., et al. (2006). Bruchpilot Promotes Active Zone Assembly, Ca^{2+} Channel
Clustering, and Vesicle Release. Science *312*, 1051–1054. https://doi.org/10.1126/science.1126308.

18. Wagh, D.A., Rasse, T.M., Asan, E., Hofbauer, A., Schwenkert, I., Dürrbeck, H., Buchner, S.,
Dabauvalle, M.-C., Schmidt, M., Qin, G., et al. (2006). Bruchpilot, a Protein with Homology to
ELKS/CAST, Is Required for Structural Integrity and Function of Synaptic Active Zones in Drosophila.
Neuron *49*, 833–844. https://doi.org/10.1016/j.neuron.2006.02.008.

19. Fouquet, W., Owald, D., Wichmann, C., Mertel, S., Depner, H., Dyba, M., Hallermann, S., Kittel,
R.J., Eimer, S., and Sigrist, S.J. (2009). Maturation of active zone assembly by Drosophila Bruchpilot. J
Cell Biol *186*, 129–145. https://doi.org/10.1083/jcb.200812150.

20. Matkovic, T., Siebert, M., Knoche, E., Depner, H., Mertel, S., Owald, D., Schmidt, M., Thomas, U.,
Sickmann, A., Kamin, D., et al. (2013). The Bruchpilot cytomatrix determines the size of the readily
releasable pool of synaptic vesicles. J Cell Biol *202*, 667–683. https://doi.org/10.1083/jcb.201301072.

21. Hallermann, S., Kittel, R.J., Wichmann, C., Weyhersmüller, A., Fouquet, W., Mertel, S., Owald, D.,
Eimer, S., Depner, H., Schwärzel, M., et al. (2010). Naked Dense Bodies Provoke Depression. J
Neurosci *30*, 14340–14345. https://doi.org/10.1523/jneurosci.2495-10.2010.

22. Chen, Y., Akin, O., Nern, A., Tsui, C.Y.K., Pecot, M.Y., and Zipursky, S.L. (2014). Cell-type-
Specific Labeling of Synapses In Vivo through Synaptic Tagging with Recombination. Neuron *81*, 280–
293. https://doi.org/10.1016/j.neuron.2013.12.021.

23. Gärtig, P.-A., Ostrovsky, A., Manhart, L., Giachello, C.N.G., Kovacevic, T., Lustig, H., Chwalla, B.,
Cachero, S., Baines, R.A., Landgraf, M., et al. (2019). Motor circuit function is stabilized during
postembryonic growth by anterograde trans-synaptic Jelly Belly - Anaplastic Lymphoma Kinase
signaling. bioRxiv, 841106. https://doi.org/10.1101/841106.

24. Kamiyama, D., Sekine, S., Barsi-Rhyne, B., Hu, J., Chen, B., Gilbert, L.A., Ishikawa, H., Leonetti,
M.D., Marshall, W.F., Weissman, J.S., et al. (2016). Versatile protein tagging in cells with split
fluorescent protein. Nat. Commun. *7*, 11046. https://doi.org/10.1038/ncomms11046.

25. Kondo, S., Takahashi, T., Yamagata, N., Imanishi, Y., Katow, H., Hiramatsu, S., Lynn, K., Abe, A.,
Kumaraswamy, A., and Tanimoto, H. (2020). Neurochemical Organization of the Drosophila Brain
Visualized by Endogenously Tagged Neurotransmitter Receptors. Cell Reports *30*, 284-297.e5.
https://doi.org/10.1016/j.celrep.2019.12.018.

26. Kohl, J., Ng, J., Cachero, S., Ciabatti, E., Dolan, M.-J., Sutcliffe, B., Tozer, A., Ruehle, S., Krueger,
D., Frechter, S., et al. (2014). Ultrafast tissue staining with chemical tags. Proc. Natl. Acad. Sci. *111*,
E3805–E3814. https://doi.org/10.1073/pnas.1411087111.

27. Jenett, A., Rubin, G.M., Ngo, T.-T.B., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B.D.,
Cavallaro, A., Hall, D., Jeter, J., et al. (2012). A GAL4-Driver Line Resource for Drosophila
Neurobiology. Cell Rep. *2*, 991–1001. https://doi.org/10.1016/j.celrep.2012.09.011.

28. Waddell, S., Armstrong, J.D., Kitamoto, T., Kaiser, K., and Quinn, W.G. (2000). The amnesiac
Gene Product Is Expressed in Two Neurons in the Drosophila Brain that Are Critical for Memory. Cell
103, 805–813. https://doi.org/10.1016/s0092-8674(00)00183-5.

29. Wu, H., Maekawa, Y., Eno, S., Kondo, S., Yamagata, N., and Tanimoto, H. (2024). High-throughput
synapse profiling reveals cell-type-specific spatial configurations in the fly brain. bioRxiv,
2024.12.02.626511. https://doi.org/10.1101/2024.12.02.626511.

30. Constance, W.D., Mukherjee, A., Fisher, Y.E., Pop, S., Blanc, E., Toyama, Y., and Williams, D.W.
(2018). Neurexin and Neuroligin-based adhesion complexes drive axonal arborisation growth
independent of synaptic activity. Elife *7*, e31659. https://doi.org/10.7554/elife.31659.

31. McDonald, N.A., Fetter, R.D., and Shen, K. (2020). Assembly of synaptic active zones requires
phase separation of scaffold molecules. Nature *588*, 454–458. https://doi.org/10.1038/s41586-020-2942-
0.

32. Lucy, L.B. (1974). An iterative technique for the rectification of observed distributions.
Astronomical J *79*, 745. https://doi.org/10.1086/111605.

33. Richardson, W.H. (1972). Bayesian-Based Iterative Method of Image Restoration*. J Opt Soc Am
62, 55. https://doi.org/10.1364/josa.62.000055.

34. Ghelani, T., Escher, M., Thomas, U., Esch, K., Lützkendorf, J., Depner, H., Maglione, M., Parutto,
P., Gratz, S., Matkovic-Rachid, T., et al. (2023). Interactive nanocluster compaction of the ELKS
scaffold and Cacophony Ca2+ channels drives sustained active zone potentiation. Sci. Adv. *9*, eade7804.
https://doi.org/10.1126/sciadv.ade7804.

35. Kiragasi, B., Wondolowski, J., Li, Y., and Dickman, D.K. (2017). A Presynaptic Glutamate
Receptor Subunit Confers Robustness to Neurotransmission and Homeostatic Potentiation. Cell Reports
19, 2694–2706. https://doi.org/10.1016/j.celrep.2017.06.003.

36. Ermolyuk, Y.S., Alder, F.G., Surges, R., Pavlov, I.Y., Timofeeva, Y., Kullmann, D.M., and
Volynski, K.E. (2013). Differential triggering of spontaneous glutamate release by P/Q-, N- and R-type
Ca2+ channels. Nat. Neurosci. *16*, 1754–1763. https://doi.org/10.1038/nn.3563.

37. Huang, S., Piao, C., Beuschel, C.B., Zhao, Z., and Sigrist, S.J. (2022). A brain-wide form of
presynaptic active zone plasticity orchestrates resilience to brain aging in Drosophila. PLOS Biol. *20*,
e3001730. https://doi.org/10.1371/journal.pbio.3001730.

38. Huang, S., Piao, C., Beuschel, C.B., Götz, T., and Sigrist, S.J. (2020). Presynaptic Active Zone
Plasticity Encodes Sleep Need in Drosophila. Curr Biol *30*, 1077-1091.e5.
https://doi.org/10.1016/j.cub.2020.01.019.

39. Weiss, J.T., Blundell, M.Z., Singh, P., and Donlea, J.M. (2024). Sleep deprivation drives brain-wide
changes in cholinergic presynapse abundance in Drosophila melanogaster. Proc. Natl. Acad. Sci. *121*,
e2312664121. https://doi.org/10.1073/pnas.2312664121.

40. Weiss, J.T., and Donlea, J.M. (2021). Sleep deprivation results in diverse patterns of synaptic scaling
across the Drosophila mushroom bodies. Curr. Biol. *31*, 3248-3261.e3.
https://doi.org/10.1016/j.cub.2021.05.018.

41. Gupta, V.K., Pech, U., Bhukel, A., Fulterer, A., Ender, A., Mauermann, S.F., Andlauer, T.F.M.,
Antwi-Adjei, E., Beuschel, C., Thriene, K., et al. (2016). Spermidine Suppresses Age-Associated

Memory Impairment by Preventing Adverse Increase of Presynaptic Active Zone Size and Release. Plos Biol *14*, e1002563. https://doi.org/10.1371/journal.pbio.1002563.

42. Zirin, J., Hu, Y., Liu, L., Yang-Zhou, D., Colbeth, R., Yan, D., Ewen-Campen, B., Tao, R., Vogt, E., VanNest, S., et al. (2020). Large-Scale Transgenic Drosophila Resource Collections for Loss- and Gain-of-Function Studies. Genetics *214*, 755–767. https://doi.org/10.1534/genetics.119.302964.

43. Sun, H., Nishioka, T., Hiramatsu, S., Kondo, S., Amano, M., Kaibuchi, K., Ichinose, T., and Tanimoto, H. (2020). Dopamine receptor Dop1R2 stabilizes appetitive olfactory memory through the Raf/MAPK pathway in Drosophila. J Neurosci *40*, 1572–19. https://doi.org/10.1523/jneurosci.1572-19.2020.

44. Busch, S., Selcho, M., Ito, K., and Tanimoto, H. (2009). A map of octopaminergic neurons in the Drosophila brain. J Comp Neurol *513*, 643–667. https://doi.org/10.1002/cne.21966.

45. Monastirioti, M., Jr., C.E.L., and White, K. (1996). Characterization of Drosophila Tyramine β-HydroxylaseGene and Isolation of Mutant Flies Lacking Octopamine. J Neurosci *16*, 3900–3911. https://doi.org/10.1523/jneurosci.16-12-03900.1996.

46. Dietzl, G., Chen, D., Schnorrer, F., Su, K.-C., Barinova, Y., Fellner, M., Gasser, B., Kinsey, K., Oppel, S., Scheiblauer, S., et al. (2007). A genome-wide transgenic RNAi library for conditional gene inactivation in Drosophila. Nature *448*, 151–156. https://doi.org/10.1038/nature05954.

47. Maqueira, B., Chatwin, H., and Evans, P.D. (2005). Identification and characterization of a novel family of Drosophilaβ☐adrenergic☐like octopamine G☐protein coupled receptors. J Neurochem *94*, 547–560. https://doi.org/10.1111/j.1471-4159.2005.03251.x.

48. Levin, L.R., Han, P.-L., Hwang, P.M., Feinstein, P.G., Davis, R.L., and Reed, R.R. (1992). The Drosophila learning and memory gene rutabaga encodes a Ca2+calmodulin-responsive adenylyl cyclase. Cell *68*, 479–489. https://doi.org/10.1016/0092-8674(92)90185-f.

49. Livingstone, M.S., Sziber, P.P., and Quinn, W.G. (1984). Loss of calcium/calmodulin responsiveness in adenylate cyclase of rutabaga, a Drosophila learning mutant. Cell *37*, 205–215. https://doi.org/10.1016/0092-8674(84)90316-7.

50. Han, P.-L., Levin, L.R., Reed, R.R., and Davis, R.L. (1992). Preferential expression of the drosophila rutabaga gene in mushroom bodies, neural centers for learning in insects. Neuron *9*, 619–627. https://doi.org/10.1016/0896-6273(92)90026-a.

51. Thum, A.S., Jenett, A., Ito, K., Heisenberg, M., and Tanimoto, H. (2007). Multiple Memory Traces for Olfactory Reward Learning in Drosophila. J Neurosci *27*, 11132–11138. https://doi.org/10.1523/jneurosci.2712-07.2007.

52. Perkins, L.A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., Yang-Zhou, D., Flockhart, I., Binari, R., Shim, H.-S., et al. (2015). The Transgenic RNAi Project at Harvard Medical School: Resources and Validation. Genetics *201*, 843–852. https://doi.org/10.1534/genetics.115.180208.

53. LeDue, E.E., Mann, K., Koch, E., Chu, B., Dakin, R., and Gordon, M.D. (2016). Starvation-Induced
Depotentiation of Bitter Taste in Drosophila. Curr Biol *26*, 2854–2861.
https://doi.org/10.1016/j.cub.2016.08.028.

54. Renger, J.J., Ueda, A., Atwood, H.L., Govind, C.K., and Wu, C.-F. (2000). Role of cAMP Cascade
in Synaptic Stability and Plasticity: Ultrastructural and Physiological Analyses of Individual Synaptic
Boutons in Drosophila Memory Mutants. J Neurosci *20*, 3980–3992.
https://doi.org/10.1523/jneurosci.20-11-03980.2000.

55. Boto, T., Louis, T., Jindachomthong, K., Jalink, K., and Tomchik, S.M. (2014). Dopaminergic
Modulation of cAMP Drives Nonlinear Plasticity across the Drosophila Mushroom Body Lobes. Curr
Biol *24*, 822–831. https://doi.org/10.1016/j.cub.2014.03.021.

56. Wang, L., Wu, C., Peng, W., Zhou, Z., Zeng, J., Li, X., Yang, Y., Yu, S., Zou, Y., Huang, M., et al.
(2022). A high-performance genetically encoded fluorescent indicator for in vivo cAMP imaging. Nat.
Commun. *13*, 5363. https://doi.org/10.1038/s41467-022-32994-7.

57. Maiellaro, I., Lohse, M.J., Kittel, R.J., and Calebiro, D. (2016). cAMP Signals in Drosophila Motor
Neurons Are Confined to Single Synaptic Boutons. Cell Reports *17*, 1238–1246.
https://doi.org/10.1016/j.celrep.2016.09.090.

58. Zaccolo, M., Zerio, A., and Lobo, M.J. (2021). Subcellular Organization of the cAMP Signaling
Pathway. Pharmacol. Rev. *73*, 278–309. https://doi.org/10.1124/pharmrev.120.000086.

59. Gervasi, N., Tchénio, P., and Preat, T. (2010). PKA Dynamics in a Drosophila Learning Center:
Coincidence Detection by Rutabaga Adenylyl Cyclase and Spatial Regulation by Dunce
Phosphodiesterase. Neuron *65*, 516–529. https://doi.org/10.1016/j.neuron.2010.01.014.

60. Ehmann, N., Owald, D., and Kittel, R.J. (2017). Drosophila active zones: From molecules to
behaviour. Neurosci Res *127*, 14–24. https://doi.org/10.1016/j.neures.2017.11.015.

61. Cunningham, K.L., Sauvola, C.W., Tavana, S., and Littleton, J.T. (2022). Regulation of presynaptic
Ca2+ channel abundance at active zones through a balance of delivery and turnover. eLife *11*, e78648.
https://doi.org/10.7554/elife.78648.

62. Williams, C.L., and Smith, S.M. (2018). Calcium dependence of spontaneous neurotransmitter
release. J. Neurosci. Res. *96*, 335–347. https://doi.org/10.1002/jnr.24116.

63. Handler, A., Graham, T.G.W., Cohn, R., Morantte, I., Siliciano, A.F., Zeng, J., Li, Y., and Ruta, V.
(2019). Distinct Dopamine Receptor Pathways Underlie the Temporal Sensitivity of Associative
Learning. Cell *178*, 60-75.e19. https://doi.org/10.1016/j.cell.2019.05.040.

64. Noyes, N.C., and Davis, R.L. (2023). Innate and learned odor-guided behaviors utilize distinct
molecular signaling pathways in a shared dopaminergic circuit. Cell Reports *42*, 112026.
https://doi.org/10.1016/j.celrep.2023.112026.

65. Abe, T., Yamazaki, D., Hiroi, M., Ueoka, Y., Maeyama, Y., and Tabata, T. (2023). Revisiting the
role of cAMP in Drosophila aversive olfactory memory formation. bioRxiv, 2023.06.26.545795.
https://doi.org/10.1101/2023.06.26.545795.

66. Tsao, C.-H., Chen, C.-C., Lin, C.-H., Yang, H.-Y., and Lin, S. (2018). Drosophila mushroom bodies
integrate hunger and satiety signals to control innate food-seeking behavior. Elife 7, e35264.
https://doi.org/10.7554/elife.35264.

67. Hancock, C.E., Rostami, V., Rachad, E.Y., Deimel, S.H., Nawrot, M.P., and Fiala, A. (2022).
Visualization of learning-induced synaptic plasticity in output neurons of the Drosophila mushroom
body γ-lobe. Sci. Rep. 12, 10421. https://doi.org/10.1038/s41598-022-14413-5.

68. Perisse, E., Owald, D., Barnstedt, O., Talbot, C.B., Huetteroth, W., and Waddell, S. (2016). Aversive
Learning and Appetitive Motivation Toggle Feed-Forward Inhibition in the Drosophila Mushroom Body.
Neuron 90, 1086–1099. https://doi.org/10.1016/j.neuron.2016.04.034.

69. Pavlowsky, A., Schor, J., Plaçais, P.-Y., and Preat, T. (2018). A GABAergic Feedback Shapes
Dopaminergic Input on the Drosophila Mushroom Body to Promote Appetitive Long-Term Memory.
Curr Biol 28, 1783-1793.e4. https://doi.org/10.1016/j.cub.2018.04.040.

70. Sitaraman, D., Aso, Y., Jin, X., Chen, N., Felix, M., Rubin, G.M., and Nitabach, M.N. (2015).
Propagation of Homeostatic Sleep Signals by Segregated Synaptic Microcircuits of the Drosophila
Mushroom Body. Curr. Biol. 25, 2915–2927. https://doi.org/10.1016/j.cub.2015.09.017.

71. Berger, M., Auweiler, K., Tegtmeier, M., Dorn, K., Khadrawe, T.E., and Scholz, H. (2023).
Octopamine integrates the status of internal energy supply into the formation of food-related memories.
https://doi.org/10.7554/elife.88247.1.

72. Claßen, G., and Scholz, H. (2018). Octopamine Shifts the Behavioral Response From Indecision to
Approach or Aversion in Drosophila melanogaster. Front Behav Neurosci 12, 131.
https://doi.org/10.3389/fnbeh.2018.00131.

73. Scheiner, R., Steinbach, A., Claßen, G., Strudthoff, N., and Scholz, H. (2014). Octopamine indirectly
affects proboscis extension response habituation in Drosophila melanogaster by controlling sucrose
responsiveness. J Insect Physiol 69, 107–117. https://doi.org/10.1016/j.jinsphys.2014.03.011.

74. Schneider, A., Ruppert, M., Hendrich, O., Giang, T., Ogueta, M., Hampel, S., Vollbach, M.,
Büschges, A., and Scholz, H. (2012). Neuronal Basis of Innate Olfactory Attraction to Ethanol in
Drosophila. Plos One 7, e52007. https://doi.org/10.1371/journal.pone.0052007.

75. Yang, Z., Yu, Y., Zhang, V., Tian, Y., Qi, W., and Wang, L. (2015). Octopamine mediates
starvation-induced hyperactivity in adult Drosophila. Proc National Acad Sci 112, 5219–5224.
https://doi.org/10.1073/pnas.1417838112.

76. Youn, H., Kirkhart, C., Chia, J., and Scott, K. (2018). A subset of octopaminergic neurons that promotes feeding initiation in Drosophila melanogaster. Plos One *13*, e0198362. https://doi.org/10.1371/journal.pone.0198362.

77. Sayin, S., Backer, J.-F.D., Siju, K.P., Wosniack, M.E., Lewis, L.P., Frisch, L.-M., Gansen, B., Schlegel, P., Edmondson-Stait, A., Sharifi, N., et al. (2019). A Neural Circuit Arbitrates between Persistence and Withdrawal in Hungry Drosophila. Neuron *104*, 544-558.e6. https://doi.org/10.1016/j.neuron.2019.07.028.

78. Turrel, O., Ramesh, N., Escher, M.J.F., Pooryasin, A., and Sigrist, S.J. (2022). Transient active zone remodeling in the Drosophila mushroom body supports memory. Curr Biol. https://doi.org/10.1016/j.cub.2022.10.017.

79. Zhang, X., Li, Q., Wang, L., Liu, Z.-J., and Zhong, Y. (2018). Active Protection: Learning-Activated Raf/MAPK Activity Protects Labile Memory from Rac1-Independent Forgetting. Neuron *98*, 142-155.e4. https://doi.org/10.1016/j.neuron.2018.02.025.

80. Tirian, L., and Dickson, B.J. (2017). The VT GAL4, LexA, and split-GAL4 driver line collections for targeted expression in the Drosophila nervous system. bioRxiv, 198648. https://doi.org/10.1101/198648.

81. Stocker, R.F., Heimbeck, G., Gendre, N., and Belle, J.S. de (1997). Neuroblast ablation in Drosophila P[GAL4] lines reveals origins of olfactory interneurons. J. Neurobiol. *32*, 443–456. https://doi.org/10.1002/(sici)1097-4695(199705)32:5<443::aid-neu1>3.0.co;2-5.

82. Friggi☐Grelin, F., Coulom, H., Meller, M., Gomez, D., Hirsh, J., and Birman, S. (2003). Targeted gene expression in Drosophila dopaminergic cells using regulatory sequences from tyrosine hydroxylase. J. Neurobiol. *54*, 618–627. https://doi.org/10.1002/neu.10185.

83. Pfeiffer, B.D., Jenett, A., Hammonds, A.S., Ngo, T.-T.B., Misra, S., Murphy, C., Scully, A., Carlson, J.W., Wan, K.H., Laverty, T.R., et al. (2008). Tools for neuroanatomy and neurogenetics in Drosophila. Proc. Natl. Acad. Sci. *105*, 9715–9720. https://doi.org/10.1073/pnas.0803697105.

84. Mahr, A., and Aberle, H. (2006). The expression pattern of the Drosophila vesicular glutamate transporter: A marker protein for motoneurons and glutamatergic centers in the brain. Gene Expr. Patterns *6*, 299–309. https://doi.org/10.1016/j.modgep.2005.07.006.

85. Cole, S.H., Carney, G.E., McClung, C.A., Willard, S.S., Taylor, B.J., and Hirsh, J. (2005). Two Functional but Noncomplementing Drosophila Tyrosine Decarboxylase Genes DISTINCT ROLES FOR NEURAL TYRAMINE AND OCTOPAMINE IN FEMALE FERTILITY*. J. Biol. Chem. *280*, 14948–14955. https://doi.org/10.1074/jbc.m414197200.

86. Sage, D., Donati, L., Soulez, F., Fortun, D., Schmit, G., Seitz, A., Guiet, R., Vonesch, C., and Unser, M. (2017). DeconvolutionLab2: An open-source software for deconvolution microscopy. Methods *115*, 28–41. https://doi.org/10.1016/j.ymeth.2016.12.015.

87. Haase, R., Royer, L.A., Steinbach, P., Schmidt, D., Dibrov, A., Schmidt, U., Weigert, M., Maghelli,
N., Tomancak, P., Jug, F., et al. (2020). CLIJ: GPU-accelerated image processing for everyone. Nat.
Methods *17*, 5–6. https://doi.org/10.1038/s41592-019-0650-1.

88. Ollion, J., Cochennec, J., Loll, F., Escudé, C., and Boudier, T. (2013). TANGO: a generic tool for
high-throughput 3D image analysis for studying nuclear organization. Bioinformatics *29*, 1840–1841.
https://doi.org/10.1093/bioinformatics/btt276.

89. Kondo, S., and Ueda, R. (2013). Highly Improved Gene Targeting by Germline-Specific Cas9
Expression in Drosophila. Genetics *195*, 715–721. https://doi.org/10.1534/genetics.113.156737.

90. Ke, M.-T., Nakai, Y., Fujimoto, S., Takayama, R., Yoshida, S., Kitajima, T.S., Sato, M., and Imai, T.
(2016). Super-Resolution Mapping of Neuronal Circuitry With an Index-Optimized Clearing Agent.
Cell Reports *14*, 2718–2732. https://doi.org/10.1016/j.celrep.2016.02.057.

91. Yamagata, N., Ezaki, T., Takahashi, T., Wu, H., and Tanimoto, H. (2021). Presynaptic inhibition of
dopamine neurons controls optimistic bias. Elife *10*, e64907. https://doi.org/10.7554/elife.64907.

MAIN FIGURES

Figure 1. Cell-type specific dissection of the endogenous active zone scaffold

(**A**) Schematic of split-GFP tagging system. GFP_{11} was inserted prior to the stop codon of *brp*, while

GFP_{1-10} was expressed in specific cell types using the GAL4-UAS system. GFP_{11} and GFP_{1-10}

reconstitute to emit fluorescence. (**B**) Visualization of Brp::rGFP in a cell-type specific manner.

Immunostaining using antibody nc82 which labels all Brp in the brain, as a comparison to Brp::rGFP.

Diagrams in each panel show the innervation pattern of neurons in the MB. GAL4 lines used: γ KCs

(MB009B), α/β KCs (MB008B), α'/β' KCs (MB370B), PPL-1 (TH-GAL4), PAM (R58E02), DPM

(VT64246), APL (GH146). Scale bar, 20 μm. (**C**) Long-term live-imaging of Brp::rGFP (green) in a

growing motor neuron (magenta, CD4::tdTomato) from 48 to 96 hours APF. OK371-GAL4 was used to

express GFP_{1-10} and CD4::tdTomato. White boxes indicate zoomed-in areas shown in the lower panels.

Scale bars: upper panels, 10 μm; lower panels, 5 μm. (**D**) Ex vivo live-imaging of Brp::rGFP in 3^{rd} instar

larval KCs. GFP_{1-10} was expressed using R13F02-GAL4. The left panel shows the MB calyx region.

White boxes indicate zoomed-in areas shown on the right panels. Box 1 and 2 demonstrate Brp::rGFP

fission and fusion event. Whites arrows indicate Brp::rGFP clusters undergoing fission or fusion. Scale

bars: left panel, 10 μm; right panels, 200 nm.

Figure 2. Intracellular active zone heterogeneity among MB compartments

(**A**) Brp::rGFP visualized in KCs. The horizontal lobe of the MB was imaged. Brp::rGFP was visualized

using R13F02-GAL4.(**B**) Median intensity of Brp::rGFP clusters in each γ lobe compartment. R13F02-

GAL4 was used. Kruskal-Wallis test. n = 11. γ1 vs. γ2: P = 0.0012; γ1 vs. γ3: P = 0.0004; γ1 vs. γ4: P =

0.0004; γ1 vs. γ5: P < 0.0001. (**C**) Diagram showing compartments in the MB γ lobe and the

intracellular Brp localization in a single γ KC. (**D**) Basal Ca^{2+} concentration near the active zones varies

by compartments. The left panel shows the schematic of the Brp::mCherry::GCaMP6s ratio-matric

sensor. GCaMP6s sensor is fused to mCherry and Brp[short]. The sensor was expressed by R13F02-GAL4

and GCaMP6s signal is normalized by mCherry for analysis. Kruskal-Wallis test. n = 8. γ1 vs. γ3: P =

0.0494; γ1 vs. γ4: P = 0.0494; γ1 vs. γ5: P = 0.0155; γ2 vs. γ5: P = 0.0494; Scale bar, 10 μm. Data are

represented as box plots showing center (median), whiskers (Min. to Max.). Significant differences (P <

0.05) are indicated by distinct letters or *.

Figure 3 Acute starvation decreases the Brp::rGFP heterogeneity level among γ compartments

(**A**) Standard deviation (SD) as an indicative of Brp::rGFP heterogeneity level among compartments. The medians of Brp::rGFP intensity of clusters in each compartment were log-transformed to calculate the relative ratio of intensity among compartments. SD is calculated from five log-transformed medians for each brain sample. (**B**) Starvation for 48 hours reduced the Brp heterogeneity level in KCs. Pseudo color in the images represent the value of $\log_2$(pixel intensity/mean pixel intensity in the γ lobe). Pseudo color range: -1.2 to 1.2. The heterogeneity level (SD) was quantified for individual brain samples. Ordinary one-way ANOVA. Fed (n = 19) vs. Starved (n = 17): P = 0.0400; Starved (n = 17) vs. Refeed (n = 19): P = 0.0400; Fed (n = 19) vs. Refeed (n = 19): P = 0.7838; Scale bar, 10 μm. Data are represented as box plots showing center (median), whiskers (Min. to Max.). Significant differences (P < 0.05) are indicated by distinct letters or *.

Figure 4. Octopamine controls the AZ structure in KCs

(**A**) OANs in the adult brain. OANs were labeled by Tdc2-GAL4 driven CD4::tdTomato in the left

panel. OA-VPM3/4 neurons were labeled with MB022B-GAL4 [9] driven CD8::GFP (right panel).

Dashed lines indicate the MB. Scale bars, 20 µm. (**B**) AZ number and synaptic vesicle localization of

OANs along the γ lobe. GFP_{1-10} and CD4::tdTomato were expressed using Tdc2-GAL4. nSyb::CLIP

labeling synaptic vesicles and CD8::GFP were expressed using MB022B-GAL4. Scale bars, 2 µm.

Images were cropped from the same MB and are identical in size. (**C**) AZ density of OANs across γ

compartment. Brp::rGFP was visualized using Tdc2-GAL4, and cluster density was quantified. n = 6

brains. Kruskal-Wallis test. γ1 vs. γ2: P = 0.0066; γ1 vs. γ3: P =0.0319; γ1 vs. γ5: P =0.0004. (**D**)

Schematic showing the innervation pattern of OANs along the γ lobe. (**E**) Reduced Brp compartmental

heterogeneity level in *Tβh* mutants. Representative images of Brp::rGFP in control and *Tβh*$^{SK2-4}$

background are shown. Pseudo color in the images represent the value of log2(pixel intensity/mean pixel

705   intensity in the γ lobe). Pseudo color range: -1.0 to 1.0. Scale bars, 10 µm. Kruskal-Wallis test. Ctrl (n =

706   12) vs. $T\beta h^{SK1\text{-}8}$ (n = 11) and $T\beta h^{SK2\text{-}4}$ (n = 12). Ctrl vs. $T\beta h^{SK1\text{-}8}$: P = 0.0442; Ctrl vs. $T\beta h^{SK2\text{-}4}$: P =

707   0.0228. *P < 0.05. Data are represented as box plots showing center (median), whiskers (Min. to Max.).

708   Significant differences (P < 0.05) are indicated by distinct letters or *. Kruskal-Wallis test.

Figure 5. Octβ2R and cAMP underly the Brp compartmental heterogeneity

(**A**) Knockdown of Octβ2R reduced the Brp heterogeneity level in γ KCs. Receptors are knocked-down

in γ KCs specifically using RNAi with R13F02-GAL4. Kruskal-Wallis test. Ctrl (n = 13) vs. OAMB (n

= 11), Octα2R (n = 10) and Octβ2R (n = 12): P= 0.0228 for Ctrl vs. Octβ2R. Ctrl (n = 20) vs. TyrR (n =

22), TyrR2 (n = 22) and OctTyrR (n = 15). *P < 0.05; ns = not significant. (**B**) Knockdown of Rut

significantly decreased the Brp heterogeneity level in γ KCs. Representative images showing Brp::rGFP

in control and in Rut knockdown group. Pseudo color in the images represent the value of log2(pixel

intensity/mean pixel intensity in the γ lobe). Pseudo color range: -1.2 to 1.2. Mann-Whitney U-test. Ctrl

(n = 10) vs. Rut-RNAi (n = 10): P = 0.0011. Data are represented as box plots showing center (median),

whiskers (Min. to Max.). *P < 0.05 and ns = not significant. Kruskal-Wallis test (A) and Mann-Whitney

U-test (B).

Figure 6. Nutritional states adjust the Brp compartmental heterogeneity through octopamine

(**A**) Graphical summary of the working model. Localized innervation of OANs onto γ KCs regulates the

Brp accumulation and forms the Brp compartmental heterogeneity via Octβ2R and cAMP pathway.

Nutritional states may modulate OAN activity, thereby adjusting the Brp heterogeneity. (**B**) Starvation

for 48 hours did not affect the Brp::rGFP heterogeneity level in $T\beta h^{SK2-4}$ mutant. GFP_{1-10} was expressed

in KCs using R13F02-GAL4 in control, or $T\beta h^{SK2-4}$ background flies. Pseudo color in the images

represent the value of log2(pixel intensity/mean pixel intensity in the γ lobe). Pseudo color range: -1.2 to

1.2. For the control group, Fed (n = 16) vs. Starved (n = 17): P = 0.0187; For the $T\beta h^{SK2}$ group, Fed (n =

16) vs. starved (n = 17): P > 0.05. Data are represented as box plots showing center (median), whiskers

(Min. to Max.). *P < 0.05 and ns = not significant by Mann-Whitney U-test.

 SUPPLEMENTARY INFROAMTION

Figure S1. Heterogeneous Brp enrichment in the MB

(**A**) SNAP chemical tagging labeled endogenous Brp in the brain. Scale bar, 100 μm. The dash line area

in indicates the zoomed-in area showed in (B). (**B**) Intensity difference of Brp::SNAP between γ1 and γ5

compartment on the same imaging plate. Scale bar, 50 μm. (**C**) Schematic of the MB. The MB comprise

three lobes based on the projection patterns of three KC subtypes: γ KCs, α'/β' KCs and α/β KCs. (**D**)

Brp::SNAP intensity difference across compartments. Schematic drawings above indicate compartments

of each lobe. Data are represented as box plots showing center (median), whiskers (Min. to Max.).

Significant differences (P < 0.05) are indicated by distinct letters. Kruskal-Wallis test.

Figure S2. Pre-synapses in the MB are predominantly from KCs

Pie charts showing the composition of pre-synapses in each of the three MB lobes. Data is from the

hemibrain online data but not all the cell types are listed. The number of pre-synapse and the percentage

are indicated for each cell type, including KCs, MBONs PPL-1 neurons, PAM neurons, dorsal paired

medial (DPM) neuron and anterior paired lateral (APL) neuron.

Figure S3. Brp compartmental heterogeneity in different KC subtypes

The 3D reconstruction of Brp::rGFP clusters, colored by Brp::rGFP intensity. Brp::rGFP is visualized in

γ KCs using MB009B-GAL4, in α/β KCs using MB008B-GAL4 and in α'/β' KCs using MB0370B-

GAL4. Scale bars, 20 μm.

Figure S4. Brp is associated with Ca^{2+} channels in the fly brain

(**A**) Co-labeling of Cac::sfGFP (green) and Brp::SNAP (magenta) in an adult brain. The write box

indicates the zoomed-in areas shown in (**B**). Scale bar, 50 μm. (**B**) Co-localization of Brp::SNAP and

Cac::sfGFP signals. Signal intensity profiles of both Brp::SNAP (green) and Cac::sfGFP (magenta) in

the image were plotted below. Scale bar, 10 μm. (**C**) Correlation between Brp::SNAP (green) and

Cac::sfGFP (magenta) signal intensities. The image is a selected area from γ5. Yellow circles show the

3D ROIs generated by segmenting Brp::SNAP signals. A loose setting was applied to include

surrounding pixels. The same ROI set was used to quantify the signal density (total grey value divided

by the ROI volume) for both Brp::SNAP and Cac::sfGFP. (**D**) Scatter plot showing the correlation

between Brp::SNAP and Cac::sfGFP signal intensities in an image sample. A 180˚ rotated Cac::sfGFP

image was used as a control (see also Fig. S4). Pearson's correlation coefficient (R) is shown. (**E**)

Pearson correlation coefficient (R) from three individual γ lobes showing the correlation between

Brp::SNAP and Cac::sfGFP signal intensities. Data are represented as box plots showing center

(median), whiskers (Min. to Max.). Significant differences ($P < 0.05$) are indicated by distinct letters.

Kruskal-Wallis test.

Figure S5. Correlation between Brp::SNAP and Cac::sfGFP signals intensities

(**A**) Correlation analysis of Brp::SNAP and Cac::sfGFP signal intensities. ROIs are generated using 3D

spot segmentation method without a watershed process. Relatively loose setting was applied on

Brp::SNAP images to generate wide ROIs. The same ROI set was used to calculated signal intensities in

both Brp::SNAP and Cac::sfGFP channels. The 180˚ rotated version of Cac::sfGFP image was used as

the control. A total of 4000 ROIs were analyzed, and Pearson correlation coefficient (R) values were

calculated for each sample. Scale bars, 5 μm. (**B**) Scatter plots showing the correlation between

Brp::SNAP and Cac::sfGFP signal intensities (left) and control (right) in different brain samples. R

value is indicated for each sample.

Figure S6. Brp compartmental heterogeneity in aging flies

Median intensity of Brp::rGFP clusters and the Brp::rGFP heterogeneity level were compared between 3

days old and 30 days old flies. Mann-Whitney U-test, 3 d (n = 11) vs. 30 d (n = 10): p > 0.05. Data are

represented as box plots showing center (median), whiskers (Min. to Max.). ns = not significant by

Mann-Whitney U-test.

Figure S7. Dopamine receptors knockdown does not affect the Brp compartmental heterogeneity

(**A**) Schematic showing the innervation patterns of dopamine neurons. The γ lobe is innervated by

dopamine neurons from the PPL1 and PAM clusters. (**B**) Knockdown of dopamine receptors does not

significantly alter the Brp heterogeneity level. Three types of DA receptors, DopR1, DopR2 and D2R

were knocked down using RNAi in KCs specifically with R13F02-GAL4. Representative images of

Brp::rGFP from each group are shown. Pseudo color in the images represent the value of log2(pixel

intensity/mean pixel intensity in the γ lobe). Pseudo color range: -1.3 to 1.3. Scale bar: 20 μm. Ctrl (n =

24) vs. DopR1 (n = 7), DopR2 (n = 7) and D2R (n = 10): P > 0.05. Data are represented as box plots

showing center (median), whiskers (Min. to Max.). ns = not significant by Kruskal-Wallis test.

Figure S8. Median intensities of Brp::rGFP clusters in each experiment

(**A**) Starvation and refeeding experiment. (**B**) Dopamine receptor knockdown experiment. (**C**) $T\beta h$

mutant's experiment. (**D**) Octopamine receptor knockdown experiment. (**E**) Tyramine receptor

knockdown experiment. (**F**) Rut knockdown experiment. (**G**) Starvation experiment. (**H**) Starvation

experiment of $T\beta h^{SK2-4}$ mutants. Data are represented as box plots showing center (median), whiskers

(Min. to Max.).