1 *Centella asiatica* Tissue Culture-Derived Extracellular Vesicles: A
2 Multifaceted Approach to Skincare Applications
3

4 Tsong-Min Chang[1], Chung-Chin Wu[2], Huey-Chun Huang[3], Shr-Shiuan Wang[1], Ching-Hua Chuang[2],
5 Pei-Lun Kao[2], Wei-Hsuan Tang[2], Luke Tzu-Chi Liu[2], Wei-Yin Qiu[4], Ivona Percec[5], Charles Chen[2, 6],
6 Tsun-Yung Kuo[2*]

[1]Department of Applied Cosmetology, HungKuang University, Taichung City, Taiwan.

[2]Schweitzer Biotech Company, Taipei City, Taiwan

[3]Department of Medical Laboratory Science and Biotechnology, China Medical University, Taichung City, Taiwan

[4]Department of Clinical Application, Center for iPS Cell Research and Application (CiRA), Kyoto University, Kyoto, Japan

[5]Division of Plastic Surgery, Department of Surgery, University of Pennsylvania, Philadelphia, PA, USA

[6]College of Science and Technology, Temple University, Philadelphia, PA, USA

[*]Corresponding author: kingk66.tw@gmail.com

Abstract

Background: *Centella asiatica* has long been used as a medicinal herb in traditional Asian medicine. Its wound healing, skin improvement, and neuroprotective properties have been widely studied. Extracellular vesicles (EVs) are secreted by cells and contain bioactive components with therapeutic properties.

Objectives: This study aims to characterize EVs isolated from *C. asiatica* tissue culture and investigate their therapeutic properties using *in-vitro* assays and UVB-induced damage mouse model.

Methods: EVs were isolated from *C. asiatica* tissue culture and characterized by nanoparticle tracking analysis (NTA) and scanning electron microscopy (SEM). Cytotoxicity, anti-oxidation, anti-melanin, and anti-inflammation of the EVs were evaluated by MTT assay, tyrosinase assay and RT-qPCR in chemical or *in vitro* assays. A UVB-induced photodamage mouse model was established to assess the anti-inflammation effect of EVs *in vivo*. Gels with or without EVs were applied to the damage site and skin appearance was observed daily and skin histopathology was analyzed on day 7 by H&E and immunohistochemical staining.

Results: *C. asiatica* EVs were found to contain high levels of polyphenol and mitigated hydrogen peroxide-induced intracellular ROS. The EVs were further able to reduce intracellular melanin content and tyrosinase activity, and exhibited anti-inflammatory effects by downregulating expression of pro-inflammatory genes COX2 as well as nitric oxide production. In mice with UVB-induced skin damage, daily application of *C. asiatica* EV gel reduced skin epidermis thickness and inflammation compared to UVB-only or blank gel at seven days after UV irradiation.

Conclusions: The beneficial effects of *C. asiatica* EVs on skin quality warrant further studies as promising agents in skin care applications.

Keywords: *Centella asiatica*, skin care, extracellular vesicles, antioxidants, cosmetics,

anti-inflammatory, skin whitening, UV damage, photoaging

Introduction

Centella asiatica, also commonly known as Gotu Kola, is a traditional medicinal herb widely used

across Asia [1]. Its broad spectrum of therapeutic properties is attributed to a diverse phytochemical

profile, including triterpenoids, flavonoids, and polyphenols, which contribute to the plant's

anti-inflammatory, antioxidant, and wound-healing properties, as well as its neuroprotective and

cognitive-enhancing effects in animal studies [2-4]. Several key triterpenes, including madecassoside,

madecassic acid, asiaticoside, and asiatic acid, are well-studied compounds that play primary roles in

the above therapeutic effects [4, 5]. Due to its wide-ranging therapeutic benefits, *C. asiatica* has

garnered interest in modern pharmacology for potential applications as a natural treatment in chronic

inflammation, neurodegenerative diseases, and skin improvement.

Extracellular vesicles (EVs) are membrane-bound particles of various sizes and compositions

released by cells into the extracellular environment, serving as carriers of bioactive molecules such as

proteins, lipids, and nucleic acids [6]. EVs are classified based on size, origin, and markers: exosomes

(30–150 nm), which originate from the multivesicular bodies in the endosomal pathway; microvesicles

(100–1000 nm), which bud directly from the plasma membrane; apoptotic bodies (1000-5000 nm),

which are formed from membrane blebbing in apoptotic cells [6, 7]. These vesicles play essential roles

in intercellular communication, modulating physiological and pathological processes by delivering

molecular signals to recipient cells [6]. Due to their stability and biocompatibility, EVs have been

explored for applications ranging from drug delivery and diagnostics to regenerative medicine, as well

as skin therapies via non-invasive topical applications [7, 8].

Plant-derived EVs share similarities with mammalian EVs. For example, exosome-like particles in

plants also form from multivesicular bodies akin to processes in mammalian cells [9]. Recently,

plant-derived EVs have attracted considerable attention as an alternative to mammalian EVs due to

their comparable therapeutic potential and additional advantages in safety, scalability,

cost-effectiveness, and ethical sourcing [9, 10]. Plant-derived RNA, particularly microRNAs (miRNAs),

can be found in EVs and play a significant role by influencing gene expression in recipient cells,

particularly via cross-kingdom communication, where they can modulate the expression of target

genes in other organisms, including humans [11]. Recent studies have shown that exosome-like EVs

from plants, including *C. asiatica*, promote skin improvement and wound healing more effectively

than conventional plant extracts [12]. For instance, *C. asiatica* exosome-like EVs can upregulate the

COLA1 collagen gene and downregulate the MMP1 matrix metalloproteinase gene, and inhibit

melanogenesis genes [13]. Furthermore, *C. asiatica* exosome-like EVs can be absorbed into human live

cancer cell culture within 24 hours and exhibit anticancer effects by promoting apoptosis of cancerous

cells [14].

Many medicinal plants, including *C. asiatica*, are subject to overharvesting, which threatens

biodiversity and ecosystem health. By using tissue culture techniques, EVs can be obtained without

putting strain on ecological resources, supporting conservation efforts and ethical sourcing while
meeting production demand [15]. In light of the therapeutic potential of *C. asiatica* and the unique
properties of EVs as skin-penetrating biomolecular vehicles, this study aims to investigate the *in-vitro*
properties of EVs isolated from *C. asiatica* tissue culture as well as the effects of *C. asiatica* EV-based
gel on UV-induced skin damage in a mouse model.
82

Materials and Methods

Isolation and characterization of *Centella asiatica* extracellular vesicles

C. asiatica EVs were extracted and isolated by a multi-step filtration and centrifugation. Briefly,
tissue culture *C. asiatica* was homogenized, coarsely filtered to remove large debris, and centrifuged.
Pellets were collected and filtered through a 0.45 μm filter, extracted and finally filtered through a
0.45/0.22 μm membrane filter.

The size and morphology of the *C. asiatica* EVs were analyzed with nanoparticle tracking analysis
(NTA) and transmission electron microscopy (TEM). The NTA was performed with ZetaView® system
(Particle Metrix, Germany). A sample for TEM was prepared by dropping 10 μL of the specimen onto a
copper grid, followed by rest for 2 minutes and blotted dry with filter paper. Ten μL of uranyl acetate
was then applied to the grid and excess liquid was again dried with filter paper, and the process was
again repeated. Afterward, the grid was air-dried and subjected to the TEM procedure.

Cell culture

The B16F10 cell line (mouse melanoma cells, ATCC CRL6475, BCRC60031) was obtained from the
Bioresource Collection and Research Center (BCRC), Taiwan. The cells were maintained in Dulbecco's
Modified Eagle Medium (DMEM) (Hyclone, Logan, UT) supplemented with 10% fetal bovine serum
(FBS) and 1% antibiotics. Human keratinocyte line HaCaT (AddexBio T0020001) was grown in
high-glucose DMEM supplemented with 10% FBS, 4 mM/L-glutamine, and 1% antibiotic-antimycotic
solution. Human foreskin fibroblast line HS68 (BCRC 60038) was grown in high-glucose DMEM
supplemented with 10% FBS, 4 mM/L-glutamine, and 1% antibiotic-antimycotic solution. Human
keratinocytes line Detroit 551 (BCRC 60118) was grown in Minimum Essential Medium (MEM)
supplemented with 10% FBS, 0.1 mM non-essential amino acids, and 1.0 mM sodium pyruvate. All
cells were maintained at 37°C in a humidified atmosphere containing 5% CO_2.

Cell viability assay

The 3-(4,5-dimethylthiazol-2-yl)-2,5 diphenyltetrazolium bromide (MTT) method was performed
for B16F10 cells. Briefly, the cells were exposed to various concentrations of *C. asiatica* EVs for 24
hours, and the MTT solution was then added to the wells. The insoluble derivative of MTT produced
by intracellular dehydrogenase was solubilized with an ethanol-DMSO (1:1) mixture. The absorbance
of the wells at 570 nm was read using a microplate reader. For HaCaT and Detroit 551 cells, 1/10
volume of alamarBlue® (resazurin) reagent was added directly after 24 hours of EV treatment and
incubated for 4 hours at 37°C. The absorbance was then read at 570 nm with 600 nm as a reference.
In both assays, cell viability was calculated as the ratio of absorbance of the treated cells to control.

Thiazoline-6-sulfonic acid (ABTS) free radical scavenging assay

The ABTS free radical assays were conducted as previously outlined [16]. A 7 mM stock solution of ABTS was reacted with 2.45 mM potassium persulfate, and the mixture was left to stand in the dark for at least 6 hours before use. The absorbance at 734 nm was measured immediately after mixing various concentrations of *C. asiatica* EVs (4×10^6, 1×10^7, and 2×10^7 particles/mL) with 1 mL of ABTS solution to detect changes within 10 minutes. Vitamin C (0.9 mg/mL) and BHA (0.9 mg/mL) were used as positive controls.

Intracellular reactive oxygen species (ROS) assay

The intracellular ROS assay was performed using an established method [17]. B16F10 cells were cultured in 24-well plates and treated with various concentrations of *C. asiatica* EVs (1×10^9, 2.5×10^9, and 5×10^9 particles/mL), Trolox$^{®}$ (0.05 mg/mL, positive control), or none (negative control) for 24 hours. The cells were then incubated with 24 mM H_2O_2 for 30 minutes to induce oxidative stress. After incubation, 2', 7'-dichloro-fluorescein diacetate (DCFH-DA) was added to the cells and cultured for 30 minutes. After treatment, the cells were washed with PBS and trypsinized with trypsin/EDTA. The fluorescence intensities of DCF were measured at an excitation wavelength of 504 nm and an emission wavelength of 524 nm using a Fluoroskan Ascent fluorescent reader (Thermo Scientific, Vantaa, Finland).

Total polyphenol content assay

Different concentrations of *C. asiatica* EVs (2×10^7, 4×10^7, 5×10^7 particles/mL) and gallic acid standard were mixed with Folin-Ciocalteu reagent and incubated for 5 minutes. Sodium carbonate solution was added and incubated for 30 minutes. The absorbance at 765 nm was measured using a spectrophotometer and the gallic acid equivalent was calculated with the gallic acid standard curve [18].

Metal chelating assay

The metal chelating activity of the extracellular vesicles was performed with potassium ferricyanide ($K_3[Fe(CN)_6]$) assay [19]. Different concentrations of *C. asiatica* EVs (2×10^7, 5×10^7, 1×10^8 particles/mL) were mixed with 0.2 mM PBS and 1% (W/V) $K_3[Fe(CN)_6]$, with 2 mg/mL EDTA as the positive control and deionized water as the blank control. The mixtures were heated at 50°C for 20 minutes. After cooling to room temperature, 10% (W/V) TCA was added, and the mixtures were centrifuged at 12,000 rpm and 4°C for 5 minutes. The supernatants were transferred to a 96-well plate, followed by the addition of deionized water and 1% (W/V) $FeCl_3$. The plates were incubated in the dark for 30 minutes. Absorbance was measured at OD 700 nm to assess metal chelating activity.

Intracellular melanin content assay

The intracellular melanin content was measured as described previously [20]. B16F10 cells were treated with α-Melanocyte-stimulating hormone (α-MSH, 100 nM) for 24 h to induce melanin production. Following this, the cells were treated with different concentrations of *C. asiatica* EVs (1×10^9, 2.5×10^9, and 5×10^9 particles/mL), or the positive control, arbutin (0.54 mg/mL), for an additional 24 hours. Post-treatment, cell pellets containing a known number of cells were solubilized

in 1 N NaOH at 60°C for 60 minutes. The melanin content was then assayed at 405 nm.

<u>Intracellular tyrosinase activity assay</u>

The intracellular tyrosinase activity was determined as described in previous research [21].

B16F10 cells were treated with α-Melanocyte-stimulating hormone (α-MSH, 100 nM) for 24 hours to

induce melanin production. Following this, the cells were treated with different concentrations of *C.*

asiatica EVs (1×10^9, 2.5×10^9, and 5×10^9 particles/mL), or the positive control, arbutin (0.54 mg/mL),

for an additional 24 hours. After treatments, the cell extracts (100 μL) were mixed with freshly

prepared L-DOPA solution (0.1% in PBS) and incubated at 37°C. The absorbance at 490 nm was then

measured to assess the tyrosinase activity.

<u>Type I procollagen (PIP) ELISA assay</u>

Hs68 cells were seeded at 1×10^5 cells per well in 12 well plates and were cultured in 10% FBS

DMEM medium at 37°C with 5% CO_2 for 24 hours. After three washes with PBS, several different

concentrations (1×10^7, 1×10^8, 1×10^9) of *C. asiatica* EVs and 5 ng/mL TGF-β1 (AbCam Cat. #ab50036, as

positive control) were added and cultured for 24 hours. The level of type I procollagen in cell culture

supernatants was determined using a commercial PIP EIA Kit (TAKARA Cat. # MK101) according to the

manufacturer's instruction (2 step procedure).

<u>mRNA expression of aquaporin-3 (AQP3) and filaggrin (FLG) genes</u>

HaCat cells were dispensed into a 96-well plate and cultured for 24 hours under cell culture

conditions. Various concentrations of *C. asiatica* EVs and negative control (purified water only) were

added to the cells and cultured for 24 hours. Real-time PCR was performed to confirm the expression

of aquaporin-3 (AQP3) and filaggrin (FLG) at the gene level, and the test sequence is as follows. For

RNA isolation and cDNA synthesis, SuperPrep TM cell lysis & RT Kit for qPCR (TOYOBO, Cat. SCQ-101)

was used. To compare and analyze gene expression, the cDNA synthesized above was used as a

template, and Real-time PCR analysis was performed using Thunderbird TM SYBR qPCR Mix (TOYOBO,

Cat. QPS-201). The gene expression levels of the sample were quantified by normalization with

GAPDH using the $2^{-\triangle\triangle Ct}$ method.

<u>mRNA expression of pro-inflammatory genes after UVB induction</u>

HaCat cells (1×10^4/well for 96-well) or Hs68 cells (5×10^4/well for 24-well) were cultured for 24

hours and exposed to UVB at 90mJ dose. Various concentrations of *C. asiatica* EVs and negative

control (purified water only) were added to the cells and cultured for 4 hours. Real-time PCR was

performed to confirm the expression of cyclooxygenase-2 (COX) gene as above and the gene

expression levels were quantified by normalization with GAPDH using the $2^{-\triangle\triangle Ct}$ method.

<u>Measurement of nitric oxide production</u>

RAW 264.7 macrophage cells were seeded in 10% FBS DMEM medium in 24-well plates at a

density of 2×10^5 cells/well and were cultivated at 37°C with 5% CO_2 for 24 hours. Removing the

original medium, the cells were treated with 1×10^7, 1×10^8, 5×10^8, 1×10^9, 2×10^9, or 4×10^9 particles/mL

of *C. asiatica* EVs or gallic acid (10 and 20 μg/mL, Sigma-Aldrich Cat. #G7384) in serum-free DMEM

medium and incubated for 4 hours. After the pre-incubation, the cells were stimulated with 100

ng/mL LPS and incubated for 20 hours. The culture supernatants were harvested and centrifuged at
7,000 × g for 3 minutes. The samples were mixed with an equal volume (100 μL) of Griess reagent
(Sigma-Aldrich Cat. #G4410) and the absorbance was measured at 540 nm after reacting for 10
minutes.

<u>Animal study</u>

Twelve male ICR mice were obtained from BioLASCO Taiwan Co. Ltd. (Taipei City, Taiwan). The
mice were 7 weeks old and weighed from 31 to 33 g at the start of the study. The mice were randomly
assigned to four groups with three mice in each of the group as in Table 1:

Table 1. Group assignment for the animal study.

Groups	Description	# of mice
UVB	UVB exposure only	3
Control	UVB exposure + gel without active ingredients	3
Treatment 1	UVB exposure + gel with purified *C. asiatica* EVs	3
Treatment 2	UVB exposure + gel with purified *C. asiatica* EVs + TECA (Titrated extract of *C. asiatica*)	3

To remove hair for test gel administration, mice were anesthetized with Isoflurane (3% − 4%),
and back hair was shaved and further removed using hair removal cream. On Day 0, mice were
exposed to 120 mJ/cm² UVB for 5 minutes to induce skin damage. From Day 0, after UVB exposure,
the test gel was applied twice daily at a volume of 180–200 μL on the mice's back according to the
respective group assignments. Photographs of the hair removal area were taken before UVB exposure
on Day 0 and daily from Day 1 to Day 7 before applying the gel to observe skin repair. Body weight of
the animals were measured on Days 0, 1, 4, and 7. Animals were sacrificed on Day 7 for
histopathological analysis.

<u>Animal tissue histopathological analysis</u>

On the day of sacrifice, skin tissue samples were taken from back and fixed in 10% neutral
formalin for at least 24 hours, embedded, and sectioned. The long axis of the skin tissue section was
defined as the observation direction. After gradient alcohol dehydration, xylene permeation, and
paraffin embedding, the skin tissue was cut into several continuous thin sections, each 4–6 μm thick.
The first set of slides was stained with H&E, and five random fields were photographed at 200×
magnification. Epidermal thickness was measured six times per field. The second set of sections was
rehydrated, and antigen retrieval was performed using a pH 6 recovery solution.
Immunohistochemistry (IHC) staining was conducted using rabbit anti-mouse myeloperoxidase
antibody (Ab65871, Abcam Ltd, USA) at a concentration of 1 μg/mL, followed by goat anti-rabbit
secondary antibody (GTX83399, GeneTEX, Taiwan) to stain antigen-positive cells in the dermis.
Hematoxylin stain was then used as a counterstain. Six dermal areas were observed at 400×

magnification. Based on the presence of positive multinucleated cells, infiltrating cell counts were
categorized into five grades according to ISO 10993-6 2016: Table E1 [26]: 0 points (negative); 1 (1–5
positive cells per field); 2 (6–10 positive cells per field); 3 (abundant infiltration of positive cells); 4
(field densely packed with positive cells).

<u>Statistical analysis</u>

Results are presented as the mean of the values, with error bars representing the 95%
confidence intervals. The analysis package in GraphPad Prism 6.01 (Boston, MA) was used for
statistical analysis. Statistical results were calculated by one-way ANOVA with Dunnett's multiple
comparisons test.

For statistical analysis of tissue histopathology, Fisher's exact test was applied using the control
group as the reference. A threshold of grade 3 or above was set for classification. The odds ratio (O.R.)
was calculated using a one-tailed test. If the O.R. was less than 1, it indicated an effective treatment
outcome; if greater than 1, the treatment was considered ineffective.

<u>Ethical statements</u>

All animal works were supervised by a licensed veterinarian and performed in a GLP-compliant
animal testing facility. All animal studies were reviewed and approved by the Institutional Animal Care
and Use Committee (IACUC) of a qualified institute in Taiwan. All procedures involving study animals
were conducted in a manner that avoided or minimized discomfort, distress or pain to the animals.
Clinical symptoms were observed once per day on behavior, respiration, and appearance.

Results

<u>Isolation and characterization of *C. asiatica* EVs</u>

C. asiatica EVs were isolated from tissue culture *C. asiatica* by a centrifugation and filtration. The
size distribution of EVs harvested was analyzed by NTA and showed that the particle has a mean size
of 150 nm (Figure 1A). The morphological features shown by the TEM images can be observed as
vesicles of approximately 100-150 nm in size with a lipid bilayer membrane structure (Figure 1B).

<u>Cytotoxicity assessment</u>

To investigate the effect of *C. asiatica* EVs in cells, an MTT assay was used to evaluate the effect
of cell viability. B16F10 melanoma cells were treated with various concentrations of *C. asiatica* EVs
and MTT assay was performed. The results showed no significant differences compared to the control
group in this test (Figure 2). Thus, high concentrations of up to 5×10^9 particles/mL of *C. asiatica* EVs do
not exhibit cytotoxicity in B16F10 cells. In addition, no discernable differences in cell viability were
detected in HaCaT and Detroit 551 cell lines, confirming that the EVs do not exhibit cytotoxicity in
mouse or human cells (Figure S1).

<u>Antioxidant activity assessment</u>

A variety of naturally-occurring polyphenol compounds from *C. asiatica* extracts have0 been
studied, which contribute to its antioxidant properties [5]. Here we investigated the total polyphenol
content of *C. asiatica* EVs using an established method with the Folin-Ciocalteu reagent. The results
showed that the gallic acid equivalent of *C. asiatica* EVs increased with the particle number, indicating

that the antioxidant activity of the vesicles may be dose-dependent (Figure S2).

We also assessed the antioxidant activity of *C. asiatica* EVs by using two cell-free assays: ABTS

and $K_3[Fe(CN)_6]$ assays. At the higher concentrations, *C. asiatica* EVs exhibited comparable $ABTS^{+\cdot}$

scavenging capacity to positive controls vitamin C and BHA (Figure S3). For the $K_3[Fe(CN)_6]$ metal

chelating assay, the results showed that reducing capacities of EVs were also dose-dependent (Figure

S4).

To further confirm the antioxidant capacity of *C. asiatica* EVs in B16F10 cells, intracellular

reactive oxygen species (ROS) levels were measured by ROS scavenging rate using H_2O_2 at 20 mM to

induce oxidative stress. After treatment, the intracellular ROS clearance was 53.86 ± 0.94%, 59.32 ±

1.68% and 66.27 ± 1.10% for $1x10^9$, $2.5x10^9$ and $5x10^9$ (particles/mL) of *C. asiatica* EVs, respectively

(Figure 3). The results showed that all concentrations of *C. asiatica* EVs significantly increased ROS

scavenging activity compared to the control group (P<0.001), demonstrating effects similar to positive

control Trolox.

<u>Melanin inhibitory activity assessment</u>

In addition to its antioxidant properties, *C. asiatica* has potential for skin whitening application

through the inhibition of melanin synthesis [22]. First, we investigated its inhibition of melanin

production in B16F10 melanoma cells treated with α-melanocyte-stimulating hormone (α-MSH). After

C. asiatica EV treatment, the melanin content in the B16F10 cells was 93.55 ± 1.14%, 90.73 ± 1.17%

and 83.69 ± 4.31% of control at $1×10^9$, $2.5×10^9$ and $5×10^9$ particles/mL of *C. asiatica* EVs, respectively

(Figure 4A). For the positive standard arbutin (0.54 mg/mL), the remaining intracellular melanin

content was 71.95 ± 2.44% of the control. Tyrosinase catalyzes the first two steps of melanin synthesis

by hydroxylation of L-tyrosine to L-DOPA [23]. Therefore, to investigate the change in melanin

synthesis via tyrosinase activity, we have carried out intracellular tyrosinase activity in B16F10 cells

treated with α-MSH. After treatment, the remaining intracellular tyrosinase activity was 92.5 ±

6.77%, 80.2.7 ± 5.13% and 56 ± 3.83% for the $1x10^9$, $2.5x10^9$ and $5x10^9$ particles/mL of either *C.*

asiatica EVs, respectively (Figure 4B). In comparison, the remaining intracellular tyrosinase activity

was 66 ± 10.53% of control after the cells were treated with arbutin, a known skin-whitening

compound. The results indicated that $5x10^8$ particles/mL of *C. asiatica* EVs exhibited stronger

inhibitory effect on α-MSH induced tyrosinase activity in B16F10 cells than arbutin, albeit with higher

intracellular melanin content than arbutin.

<u>Effect of *C. asiatica* on skin integrity-related genes</u>

One of the major components in skin tissue is type I collagen, which is derived from its precursor

type I procollagen (PIP) secreted by skin fibroblasts, and lowered expression of PIP is associated with

skin UV damage and senescence leading to wrinkle formation [24]. *C. asiatica* EVs increased

production of PIP in Hs68 cells and induced a similar expression level of PIP to that of positive control

TGF-β1 at the highest concentration tested ($1x10^9$ particles/mL) (Figure S5). Aquaporin and filaggrin

are important in the barrier function of skin by regulating water transport and water retention in skin

[25]. Real-time PCR assay quantification of mRNA expression of aquaporin-3 (AQP3) and filaggrin (FLG)

genes in HaCaT keratinocyte cells treated with *C. asiatica* EVs showed that *C. asiatica* EVs could
upregulate the expression levels of AQP3 and FLG genes compared to the control group (Figure S6).

<u>Anti-inflammatory effects of *C. asiatica* EVs</u>

UV exposure of keratinocyte can induce COX2 expression that increases pro-inflammatory
prostalglandin synthesis, and *C. asiatica* triterpene madecassoside was found to inhibit UV-induced
inflammation [22]. Here we showed that addition of *C. asiatica* EVs can lower expression levels of
COX2 in skin fibroblast Hs68 and keratinocyte HaCaT cells exposed to UV (Figure S7). Nitric oxide (NO)
is an inflammatory response messenger secreted mainly by endothelial cells and macrophages, as well
as keratinocytes, and exposure of skin to UV irradiation results in elevated levels of NO production
[27]. We subjected *C. asiatica* EV-treated RAW 264.7 macrophage culture to lipopolysaccharide (LPS),
which is a TLR4 ligand and induces pro-inflammatory response [28]. The NO production was measured
by Griess reagent, which is converted to a pink color after reacting with nitrites. As shown in Figure 5,
the amount of nitrite decreases in a dose-dependent manner in cells treated with increasing
concentrations of EVs. Taken together, *C. asiatica* EVs were able to reduce UV-induced inflammatory
response in a variety of cell types tested.

<u>Effect of *C. asiatica* EV-based gel on UVB-induced skin damage in Mice</u>

Next, we investigated whether a *C. asiatica* EV-based gel formulation (with or without addition
of *C. asiatica* extract TECA) could alleviate skin damage induced by UVB exposure. First, we evaluated
the safety of the application of *C. asiatica* EV gel and we found no adverse effects in all groups that
were applied with gel, regardless of the presence of *C. asiatica* EV or TECA (Table S1). There were also
no significant weight differences between groups (p > 0.05), despite a slight decrease in body weight
during the study in all groups (Table S2 and Figure S8).

In tissue histopathological analysis, the epidermis measurement results are summarized in Table
2 and Figure 6. The UVB-only group had slightly less epidermis thickness (126.66 ± 47.73 µm)
compared to the control group (154.94 ± 58.14 µm), but the difference was not statistically significant.
Treatment groups 1 and 2 had average epidermal thicknesses of 87.66 ± 35.32 µm and 86.09 ± 25.74
µm, respectively. Compared to the control group (154.94 ± 58.14 µm), the epidermal thicknesses were
significantly (p < 0.01 for treatment 1 and p < 0.05 for treatment 2) reduced.

In addition, immunohistochemical staining of inflammation marker myeloperoxidase (MPO) was
used to observe positive multinucleated cells in the dermis of UV-damaged areas (Table 2 and Figure
7). In both UVB-only and control groups, 75% of the observed fields exhibited extensive infiltration of
positive multinucleated cells. The addition of *C. asiatica* EVs with or without TECA significantly
reduced cellular infiltration (8.3% in both treatment groups). The odds ratios were also significantly
lower in both treatment groups (0.03) compared to UVB and control groups (1.00) as calculated by
Fisher's exact test.

Table 2 Tissue section measurement and observation. Results are taken from measurements of two skin sections
(5 random fields for H&E stain, 6 random fields for IHC stain)

	UVB	Control	Treatment 1	Treatment 2
	H&E stain-epidermis thickness (mean ± SD in µm) (N = 10 random fields)			
Results	126.66 ±47.73[NS]	154.94 ±58.14	87.66 ±35.32[*]	86.09 ±25.74[**]
	IHC stain-Number of positives/Total number observed (N = 12 random fields)			
Myeloperoxidase-positive (score ≥3)	9/12[NS]	9/12	1/12[**]	1/12[**]
Odds ratio (95% CI)	1.00	1.00	0.03	0.03

[NS]: Statistically not significant compared to control group
[*]: Statistically significant compared to control group ($p < 0.01$).
[**]: Statistically significant compared to control group ($p < 0.05$).

Gross appearance examination of mouse skin UVB exposure site on Day 6 of the study is shown

in representative photographs in Figure 8. In UVB-only and blank gel control groups, inflammation and

redness are still apparent at the UV wound site, with only a few newly repaired skin and abundant

scarring. In contrast, for both treatment groups, the wound has been repaired and inflammation is

absent at the site with minimal scarring.

Discussion

In this study, we have isolated and characterized EVs secreted from *C. asiatica* tissue culture. The

EVs have a size of approximately 150 nm and a lipid bilayer morphology similar to that previously

reported for exosomes derived from *C. asiatica* and other plants [12, 13]. In a series of *in-vitro*

experiments in B16F10 cell culture, we have demonstrated that the EVs are safe for cells and that EVs

exhibited notable antioxidant, anti-melanogenic, and anti-inflammatory effects, suggesting potential

applications in skin care and therapeutic treatments to combat oxidative stress and

hyperpigmentation. Another study also isolated exosome-like EVs from *C. asiatica* but with a smaller

average particle size of 63.8 nm [14]. While the EVs from the above study exhibited cytotoxicity in

HepG2 human cancer cells, we did not observe any effect on cell viability in mouse melanoma B16F10

cells or HaCaT and Detroit 551 human cell lines tested (Figures 2 and S1) [14]. The therapeutic

properties of *C. asiatica* have mostly been attributed to the triterpenes and derivatives, including

madecassoside, madecassic acid, asiaticoside, and asiatic acid, and these are among the most

well-studied compounds in the plant [3-5]. In our study, we observed potent antioxidant effects from

C. asiatica, although further study is required to determine specifically how contents within EVs

(chemicals, proteins, and nucleic acids) can contribute to observed effects. In addition to

phytochemicals and proteins, small RNAs and miRNAs in EVs are important modulators of target effect

proteins, and RNA binding proteins, such as Argonaute 1, RNA helicases, and Annexins, may be

involved in packing RNA into EVs [10, 28] A recent preprint study has provided a comprehensive

analysis of the transcriptome human skin keratinocytes after treatment of *C. asiatica* exosomes,

revealing 11 novel miRNAs packed within the exosomes [13]. The above study also found that skin

integrity and barrier-related genes were upregulated after treatment by *C. asiatica* EVs, which we

have also observed in this study with increasing gene expression of PIP1, AQP3, and FLG (Figures S5

and S6) [13]. The study also found that *C. asiatica* exosomes induced higher gene expression than *C.*
asiatica extracts, possibly due to better penetration of skin cells by exosomes and the presence of
miRNAs [13]. Thus *C. asiatica* EVs could provide their effect via multitude of effectors such as
phytochemicals and miRNAs.

In *C. asiatica,* madecassoside by itself has the ability to inhibit UV-induced melanin synthesis in a
keratinocyte and melanocyte co-culture system as well as artificially tanned human skin [22]. Other
plant-derived EVs also have anti-melanogenic effect. For example, *Dendropanax morbifera* EVs can
inhibit the expression of melanogenesis genes such as MITF, TYR, TRP-1, and TRP-2 [9, 28]. In our
study, the EVs appeared to modulate melanogenesis pathways in cell culture, likely through the
inhibition of tyrosinase activity, which is a key enzyme in melanin synthesis [23]. The mechanism of
anti-melanogenesis by *C. asiatica* includes inhibition of inflammation, as observed in previous studies
[24]. and in our study, where we have observed that *C. asiatica* can reduce pro-inflammatory gene
expression, such as COX2 and decrease the production of NO (Figures 5 and S7). As melanin
overproduction contributes to skin conditions like freckles and nevus formation, and age spot
development, plant-derived EVs with anti-melanogenic properties represent an economical and
natural alternative to conventional treatments, which often come with side effects or limited efficacy
[29].

In the animal study, mice were exposed to UVB at 120 mJ/cm^2 for 5 minutes to induce
photoaging and skin damage. This was followed by daily application of EV gel for seven consecutive
days. Photoaging leads to degeneration of elastic fibers and collagen loss, resulting in epidermal
thickening of the stratum corneum [30]. In both treatment groups consist of *C. asiatica* EV gels with or
without TECA, we observed reduced epidermal thickening in skin sections and reduced cell infiltration
due to inflammation characterized using IHC staining, suggesting a potential skin repair effect.
However, in the control group with blank gel application, thicker, albeit not statistically significant,
epidermis layer was observed than UVB-only group, possibly due to mechanical irritation from the gel
application process. These preliminary results suggest that EVs may help to alleviate UV-induced
inflammation but are unable to reverse UV-induced epidermal damage.

The tissue culture micropropagation technology for *C. asiatica* has been established as a
sustainable source of *C. asiatica* [31]. This ensures its conservation in nature while providing a
constant yield of raw material, akin to the use of bioreactors in biopharmaceutical drug and vaccine
production. Using plant tissue culture as a source for EV production presents several advantages over
traditional methods of harvesting from wild plants. First, tissue culture allows for consistent, scalable
production of EVs with genetic uniformity, bypassing the variability associated with environmental
factors that affect wild plants, such as seasonal changes, soil quality, and climate/geographical
conditions [15, 29]. This standardization is particularly important for achieving reproducible results in
therapeutic and cosmetic applications, where consistent quality and efficacy are crucial in ensuring
the safety and efficacy of the product. Additionally, tissue culture can be optimized to enhance the
yield of specific bioactive compounds through controlled environmental conditions and the use of

elicitors [15].

In conclusion, this *in vitro* and animal model study demonstrated the antioxidant,
anti-melanogenic and anti-inflammatory potential of *C. asiatica*-derived EVs from tissue culture. The
advantages of tissue culture over wild plants in terms of sustainability, consistency, and the ability to
enhance bioactive compound production further support the use of this approach in developing
plant-based EV therapies. Future studies could focus on characterizing bioactive molecules and
miRNAs within these EVs, as well as exploring their safety and efficacy in *in-vivo* models. Ultimately,
this work paves the way for the sustainable production and utilization of plant-derived EVs in skin care
cosmetics and wound healing applications.

Acknowledgements

We thank the team of Schweitzer Biotech Company for reading and providing feedback for
preparation of the manuscript.

Competing Interests

C.-C. W., C.-H. C., P.-L. K., W.-H. T., L. T.-C. L., C. C., and T.-Y. K. are employees of Schweitzer
Biotech Company. W.-Y. Q. and I. P. are consultants employed by Schweitzer Biotech Company. T.-M. C.
and S.-S. W. have no competing interests to declare. This study is funded by Schweitzer Biotech
Company.

References

1.   Biswas D, Mandal S, Chatterjee Saha S, Tudu CK, Nandy S, Batiha GE, Shekhawat MS, Pandey
DK, Dey A (2021) Ethnobotany, phytochemistry, pharmacology, and toxicity of Centella asiatica
(L.) Urban: A comprehensive review. Phytother Res 35:6624–6654.
https://doi.org/10.1002/ptr.7248

2.   Sabaragamuwa R, Perera CO, Fedrizzi B (2018) Centella asiatica (Gotu kola) as a
neuroprotectant and its potential role in healthy ageing. Trends Food Sci Technol 79:88-97.
https://doi.org/10.1016/j.tifs.2018.07.024

3.   Bandopadhyay S, Mandal S, Ghorai M, Jha NK, Kumar M, Radha, Ghosh A, Proćków J, Perez de
la Lastra JM, Dey A (2023) Therapeutic properties and pharmacological activities of asiaticoside
and madecassoside: A review. J Cell Mol Med 27(5):593-608.
https://doi.org/10.1111/jcmm.17635

4.   Sun B, Wu L, Wu Y, Zhang C, Qin L, Hayashi M, Kudo M, Gao M, Liu T (2020) Therapeutic
potential of Centella asiatica and its triterpenes: a review. Front Pharmacol 11:568032.
https://doi.org/10.3389/fphar.2020.568032

5.   Diniz LR, Calado LL, Duarte AB, de Sousa DP (2023) Centella asiatica and its metabolite asiatic
acid: wound healing effects and therapeutic potential. Metabolites 13(2):276.
https://doi.org/10.3390/metabo13020276

6. Jeppesen DK, Zhang Q, Franklin JL, Coffey RJ (2023) Extracellular vesicles and nanoparticles:
emerging complexities. Trends Cell Biol 33(8):667-81.
https://doi.org/10.1016/j.tcb.2023.01.002

7. Moghassemi S, Dadashzadeh A, Sousa MJ, Vlieghe H, Yang J, León-Félix CM, Amorim CA (2024)
Extracellular vesicles in nanomedicine and regenerative medicine: A review over the last
decade. Bioact Mater 36:126-56. https://doi.org/10.1016/j.bioactmat.2024.02.021

8. Yang GH, Lee YB, Kang D, Choi E, Nam Y, Lee KH, You HJ, Kang HJ, An SH, Jeon H (2021)
Overcome the barriers of the skin: exosome therapy. Biomater Res 25(1):22.
https://doi.org/10.1186/s40824-021-00224-8

9. Li A, Li D, Gu Y, Liu R, Tang X, Zhao Y, Qi F, Wei J, Liu J (2023) Plant-derived nanovesicles: Further
exploration of biomedical function and application potential. Acta Pharm Sin B 13(8):3300-20.
https://doi.org/10.1016/j.apsb.2022.12.022

10. Zhu Y, Zhao J, Ding H, Qiu M, Xue L, Ge D, Wen G, Ren H, Li P, Wang J (2024) Applications of
plant-derived extracellular vesicles in medicine. MedComm. 2024 5(10):e741.
https://doi.org/10.1002/mco2.741

11. Urzì O, Gasparro R, Ganji NR, Alessandro R, Raimondo S (2022) Plant-RNA in extracellular
vesicles: the secret of cross-kingdom communication. Membranes 12(4):352.
https://doi.org/10.3390/membranes12040352

12. Cho JH, Hong YD, Kim D, Park SJ, Kim JS, Kim HM, Yoon EJ, Cho JS (2022) Confirmation of
plant-derived exosomes as bioactive substances for skin application through comparative
analysis of keratinocyte transcriptome. Appl Bio Chem 65(1):8.
https://doi.org/10.1186/s13765-022-00676-z

13. Kim MH, Yoon EJ, Kim JS, Park SJ, Lee H (2023) Comparative analysis of the transcriptome and
efficacy of bioactive Centella asiatica exosomes on skin cells. ResearchSquare
https://doi.org/10.21203/rs.3.rs-2787704/v1

14. Huang J, Cao X, Wu W, Han L, Wang F (2024) Investigating the proliferative inhibition of HepG2
cells by exosome-like nanovesicles derived from Centella asiatica extract through
metabolomics. Biomed Pharmacother 176:116855.
https://doi.org/10.1016/j.biopha.2024.116855

15. Kumari S, Kumar N, Trivedi MP (2022) Application of Plant Tissue Culture for Improvement of
Centella asiatica. In: Kumar N (ed) Biotechnology and Crop Improvement, 1st ed. CRC Press,
Boca Raton, FL, pp. 93-106. https://doi.org/10.1201/9781003239

16. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C (1999) Antioxidant activity
applying an improved ABTS radical cation decolorization assay. Free Radic Biol Med
26(9-10):1231-7. https://doi.org/10.1016/S0891-5849(98)00315-3

17. Murrant CL, Reid MB (2001) Detection of reactive oxygen and reactive nitrogen species in
skeletal muscle. Microsc Res Tech 55(4):236-48. https://doi.org/10.1002/jemt.1173

18. Singleton VL, Orthofer R, Lamuela-Raventós RM (1999) Analysis of total phenols and other

oxidation substrates and antioxidants by means of folin-ciocalteu reagent. In: Packer L (ed) Methods in enzymology. Academic Press, Cambridge MA, Vol. 299, pp. 152-178. https://doi.org/10.1016/S0076-6879(99)99017-1

19. Adjimani JP, Asare P. (2015) Antioxidant and free radical scavenging activity of iron chelators. Tox Rep 2:721-8. https://doi.org/10.1016/j.toxrep.2015.04.005

20. Tsuboi T, Kondoh H, Hiratsuka J, Mishima Y (1998) Enhanced melanogenesis induced by tyrosinase gene-transfer increases boron-uptake and killing effect of boron neutron capture therapy for amelanotic melanoma. Pigment Cell Res 11(5):275-82. https://doi.org/10.1111/j.1600-0749.1998.tb00736.x

21. Yang JY, Koo JH, Song YG, Kwon KB, Lee JH, Sohn HS, Park BH, Jhee EC, Park JW (2006) Stimulation of melanogenesis by scoparone in B16 melanoma cells 1. Acta Pharmacol Sin 27(11):1467-73. https://doi.org/10.1111/j.1745-7254.2006.00435.x

22. Jung E, Lee JA, Shin S, Roh KB, Kim JH, Park D (2013) Madecassoside inhibits melanin synthesis by blocking ultraviolet-induced inflammation. Molecules 18(12):15724-36. https://doi.org/10.3390/molecules181215724

23. Snyman M, Walsdorf RE, Wix SN, Gill JG (2024) The metabolism of melanin synthesis—From melanocytes to melanoma. Pigment Cell Melanoma Res 37(4):438-452. https://doi.org/10.1111/pcmr.13165

24. Talwar HS, Griffiths CE, Fisher GJ, Hamilton TA, Voorhees JJ (1995) Reduced type I and type III procollagens in photodamaged adult human skin. J Invest Dermatol 105(2):285-90. doi: 10.1111/1523-1747.ep12318471

25. Park CH, Min SY, Yu HW, Kim K, Kim S, Lee HJ, Kim JH, Park YJ (2020) Effects of apigenin on RBL-2H3, RAW264. 7, and HaCaT cells: anti-allergic, anti-inflammatory, and skin-protective activities. Int J Mol Sci 21(13):4620. https://doi.org/10.3390/ijms21134620

26. International Organization for Standardization. (2016) ISO 10993-6:2016(En). Biological Evaluation of Medical Devices — Part 6: Tests for Local Effects after Implantation. Geneva: ISO

27. Cals-Grierson MM, Ormerod AD (2004) Nitric oxide function in the skin. Nitric oxide 10(4):179-93. https://doi.org/10.1016/j.niox.2004.04.005

28. Mantovani A, Garlanda C (2023) Humoral innate immunity and acute-phase proteins. N Engl J Med 388(5):439-52. https://doi.org/10.1056/NEJMra220634

29. Lee R, Ko HJ, Kim K, Sohn Y, Min SY, Kim JA, Na D, Yeon JH (2020) Anti-melanogenic effects of extracellular vesicles derived from plant leaves and stems in mouse melanoma cells and human healthy skin. J Extracell Vesicles 9(1):1703480. https://doi.org/10.1080/20013078.2019.1703480

30. Liu H, Dong J, Du R, Gao Y, Zhao P. (2024) Collagen study advances for photoaging skin. Photodermatol Photoimmunol Photomed 40:e12931. https://doi:10.1111/phpp.12931

31. Mohapatra P, Ray A, Sandeep IS, Nayak S, Mohanty S (2021) Tissue-Culture-Mediated Biotechnological Intervention in Centella asiatica: A Potential Antidiabetic Plant. In: Gantait S,

521   Verma SK, Sharangi AB (eds) Biotechnology of Anti-diabetic Medicinal Plants. Spring, Singapore,

522   pp. 89-116. https://doi.org/10.1007/978-981-16-3529-8_4

523   32. Park JH, Choi JY, Son DJ, Park EK, Song MJ, Hellström M, Hong JT. (2017) Anti-inflammatory

524   effect of titrated extract of Centella asiatica in phthalic anhydride-induced allergic dermatitis

525   animal model. Int J Mol Sci 18(4):738. https://doi.org/10.3390/ijms18040738

Figures

A

B

Fig. 1 Size and morphology characterization of *C. asiatica*-derived EVs. A. Nanoparticle tracking analysis (NTA) of size and concentration distribution of EVs by using the ZetaView® system. B. Representative TEM images of purified *C. asiatica* EVs with red arrows showing vesicles with lipid bilayer membrane structure. Scale bar: 100 nm

Cell viability MTT assay

Fig. 2 Cell viability assay. B16F10 cells were treated with different concentrations of *C. asiatica* EVs. After 24h treatment, cell viability was measured by the MTT assay method. Data are presented as mean ± SD (n=3). Statistical results were calculated by one-way ANOVA with Dunnett's multiple comparisons test compared to the control group.

Intracellular ROS scavenging assay

Fig. 3 Intracellular ROS level. B16F10 cells were pretreated with various concentrations of *C. asiatica* EVs, Trolox (0.05 mg/mL, positive control), or left untreated as a control for 24 hours. The cells were then incubated with 24 mM H_2O_2 and DCFH-DA. The fluorescence intensities of DCF were measured to calculate the ROS levels. Data are presented as mean ± SD (n=3). Statistical analysis was calculated by one-way ANOVA with Dunnett's multiple comparisons test, compared to the control group. ****P<0.0001

Fig. 4 Effect of *C. asiatica* EVs on melanin production. Inhibitory effects of different concentrations of *C. asiatica* EVs on **A.** melanin content and **B.** intracellular tyrosinase activity in B16F10 cells. Data are presented as mean ± SD (n=3). Statistical results were calculated by one-way ANOVA with Dunnett's multiple comparisons test compared to the control group. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001

Fig. 5 Nitric oxide (NO) production in RAW 264.7 macrophages. Cells were treated with different concentrations of *C. asiatica* EVs or gallic acid after induction by LPS. The resulting NO produced was detected by the Griess reagent.

Fig. 6 Representative examples of histopathological analysis of skin tissue sections on Day 7 of the study. Arrows indicate thickness of the epidermis. H&E stain, 200x magnification (scale bar = 100 μm)

Fig. 7 Representative examples of IHC staining of myeloperoxidase (in brown, indicated by circles) in skin tissue sections on Day 7 of the study. Mouse back skin sections were stained with rabbit anti-mouse myeloperoxidase antibody and then detected by HRP goat anti-rabbit antibody. 400x magnification (scale bar = 20 μm)

Fig. 8 Representative examples of mouse skin appearance at the site of UVB exposure on Day 6 of the study.