Centella asiatica Tissue Culture-Derived Extracellular Vesicles: A Multifaceted Approach to Skincare Applications

Tsong-Min Chang[1], Chung-Chin Wu[2], Huey-Chun Huang[3], Shr-Shiuan Wang[1], Ching-Hua Chuang[2], Pei-Lun Kao[2], Wei-Hsuan Tang[2], Luke Tzu-Chi Liu[2], Wei-Yin Qiu[4], Ivona Percec[5], Charles Chen[2, 6], Tsun-Yung Kuo[2*]

[1]Department of Applied Cosmetology, HungKuang University, Taichung City, Taiwan.

[2]Schweitzer Biotech Company, Taipei City, Taiwan

[3]Department of Medical Laboratory Science and Biotechnology, China Medical University, Taichung City, Taiwan

[4]Department of Clinical Application, Center for iPS Cell Research and Application (CiRA), Kyoto University, Kyoto, Japan

[5]Division of Plastic Surgery, Department of Surgery, University of Pennsylvania, Philadelphia, PA, USA

[6]College of Science and Technology, Temple University, Philadelphia, PA, USA

[*]Corresponding author: kingk66.tw@gmail.com

Abstract

Background: *Centella asiatica* has long been used as a medicinal herb in traditional Asian medicine. Its wound healing, skin improvement, and neuroprotective properties have been widely studied. Extracellular vesicles (EVs) are secreted by cells and contain bioactive components with therapeutic properties.

Objectives: This study aims to characterize EVs isolated from *C. asiatica* tissue culture and investigate their therapeutic properties using *in-vitro* assays.

Methods: EVs were isolated from *C. asiatica* tissue culture and their properties were characterized by cell-free chemical assays which investigated free radical activity, metal chelation, and total polyphenol content. Mouse melanoma, human fibroblast and keratinocyte cell cultures were used to characterize cell viability, intracellular reactive oxygen species (ROS), intracellular melanin and tyrosinase assay, skin integrity and pro-inflammatory gene expression level by quantitative PCR.

Results: *C. asiatica* EVs were found to contain high levels of polyphenol, which contributed to their radical scavenging and metal chelating properties and mitigated hydrogen peroxide-induced intracellular ROS. The EVs were further able to reduce intracellular melanin content and tyrosinase activity, and upregulate skin barrier genes procollagen type I, aquaporin 3, and filaggrin. The inhibitory effect of *C. asiatica* EVs on inflammation was evident in downregulation of expression of pro-inflammatory genes COX2 as well as nitric oxide production.

Conclusions: The beneficial effects of *C. asiatica* EVs warrant further studies as promising formulations in skin care applications.

Keywords: *Centella asiatica*, traditional medicine, extracellular vesicles, antioxidants, cosmetics, anti-inflammatory, skin whitening

Introduction

Centella asiatica, also commonly known as Gotu Kola, is a traditional medicinal herb widely used across Asia [1]. Its broad spectrum of therapeutic properties is attributed to a diverse phytochemical profile, including triterpenoids, flavonoids, and polyphenols, which contribute to the plant's anti-inflammatory, antioxidant, and wound-healing properties, as well as its neuroprotective and cognitive-enhancing effects in animal studies [2-4]. Several key triterpenes, including madecassoside, madecassic acid, asiaticoside, and asiatic acid, are well-studied compounds that play primary roles in the above therapeutic effects [4, 5]. Due to its wide-ranging therapeutic benefits, C. asiatica has garnered interest in modern pharmacology for potential applications as a natural treatment in chronic inflammation, neurodegenerative diseases, and skin improvement.

Extracellular vesicles (EVs) are membrane-bound particles of various sizes and compositions released by cells into the extracellular environment, serving as carriers of bioactive molecules such as proteins, lipids, and nucleic acids [6]. EVs are classified based on size, origin, and markers: exosomes (30–150 nm), which originate from the multivesicular bodies in the endosomal pathway; microvesicles (100–1000 nm), which bud directly from the plasma membrane; apoptotic bodies (1000-5000 nm), which are formed from membrane blebbing in apoptotic cells [6, 7]. These vesicles play essential roles in intercellular communication, modulating physiological and pathological processes by delivering molecular signals to recipient cells [6]. Due to their stability and biocompatibility, EVs have been explored for applications ranging from drug delivery and diagnostics to regenerative medicine, as well as skin therapies via non-invasive topical applications [7, 8].

Plant-derived EVs share similarities with mammalian EVs. For example, exosome-like particles in plants also form from multivesicular bodies akin to processes in mammalian cells [9]. Recently, plant-derived EVs have attracted considerable attention as an alternative to mammalian EVs due to their comparable therapeutic potential and additional advantages in safety, scalability, cost-effectiveness, and ethical sourcing [9, 10]. Plant-derived RNA, particularly microRNAs (miRNAs), can be found in EVs and play a significant role by influencing gene expression in recipient cells, particularly via cross-kingdom communication, where they can modulate the expression of target genes in other organisms, including humans [11]. Recent studies have shown that exosome-like EVs from plants, including C. asiatica, promote skin improvement and wound healing more effectively than conventional plant extracts [12]. For instance, C. asiatica exosome-like EVs can upregulate the COLA1 collagen gene and downregulate the MMP1 matrix metalloproteinase gene, and inhibit melanogenesis genes [13]. Furthermore, C. asiatica exosome-like EVs can be absorbed into human live cancer cell culture within 24 hours and exhibit anticancer effects by promoting apoptosis of cancerous cells [14].

Many medicinal plants, including C. asiatica, are subject to overharvesting, which threatens biodiversity and ecosystem health. By using tissue culture techniques, EVs can be obtained without putting strain on ecological resources, supporting conservation efforts and ethical sourcing while meeting production demand [15]. In light of the therapeutic potential of C. asiatica and the unique

77 properties of EVs as skin-penetrating biomolecular vehicles, this study aims to investigate the *in-vitro*
78 properties of EVs isolated from *C. asiatica* tissue culture.
79
Materials and Methods

<u>Isolation and characterization of *Centella asiatica* extracellular vesicles</u>

C. asiatica EVs were extracted and isolated by a multi-step filtration and centrifugation process.
Briefly, tissue culture *C. asiatica* was homogenized, coarsely filtered to remove large debris, and
centrifuged. Pellets were collected and filtered through a 0.45 μm filter, extracted and finally filtered
through a 0.45/0.22 μm membrane filter.

The size and morphology of the *C. asiatica* EVs were analyzed with nanoparticle tracking analysis
(NTA) and transmission electron microscopy (TEM). The NTA was performed with ZetaView® system
(Particle Metrix, Germany). A sample for TEM was prepared by dropping 10 μL of the specimen onto a
copper grid, followed by rest for 2 minutes and blotted dry with filter paper. Ten μL of uranyl acetate
was then applied to the grid and excess liquid was again dried with filter paper, and the process was
again repeated. Afterward, the grid was air-dried and subjected to the TEM procedure.

<u>Cell culture</u>

The B16F10 cell line (mouse melanoma cells, ATCC CRL6475, BCRC60031) was obtained from the
Bioresource Collection and Research Center (BCRC), Taiwan. The cells were maintained in Dulbecco's
Modified Eagle Medium (DMEM) (Hyclone, Logan, UT) supplemented with 10% fetal bovine serum
(FBS) and 1% antibiotics. Human keratinocyte line HaCaT (AddexBio T0020001) was grown in
high-glucose DMEM supplemented with 10% FBS, 4 mM/L-glutamine, and 1% antibiotic-antimycotic
solution. Human foreskin fibroblast line HS68 (BCRC 60038) was grown in high-glucose DMEM
supplemented with 10% FBS, 4 mM/L-glutamine, and 1% antibiotic-antimycotic solution. Human
keratinocytes line Detroit 551 (BCRC 60118) was grown in Minimum Essential Medium (MEM)
supplemented with 10% FBS, 0.1 mM non-essential amino acids, and 1.0 mM sodium pyruvate. All
cells were maintained at 37°C in a humidified atmosphere containing 5% CO_2.

<u>Cell viability assay</u>

The 3-(4,5-dimethylthiazol-2-yl)-2,5 diphenyltetrazolium bromide (MTT) method was performed
for B16F10 cells. Briefly, the cells were exposed to various concentrations of *C. asiatica* EVs for 24
hours, and the MTT solution was then added to the wells. The insoluble derivative of MTT produced
by intracellular dehydrogenase was solubilized with an ethanol-DMSO (1:1) mixture. The absorbance
of the wells at 570 nm was read using a microplate reader. For HaCaT and Detroit 551 cells, 1/10
volume of alamarBlue® (resazurin) reagent was added directly after 24 hours of EV treatment and
incubated for 4 hours at 37°C. The absorbance was then read at 570 nm with 600 nm as a reference.
In both assays, cell viability was calculated as the ratio of absorbance of the treated cells to control.

<u>Thiazoline-6-sulfonic acid (ABTS) free radical scavenging assay</u>

The ABTS free radical assays were conducted as previously outlined [16]. A 7 mM stock solution
of ABTS was reacted with 2.45 mM potassium persulfate, and the mixture was left to stand in the dark

for at least 6 hours before use. The absorbance at 734 nm was measured immediately after mixing

various concentrations of *C. asiatica* EVs (4×10^6, 1×10^7, and 2×10^7 particles/mL) with 1 mL of ABTS

solution to detect changes within 10 minutes. Vitamin C (0.9 mg/mL) and BHA (0.9 mg/mL) were used

as positive controls.

Intracellular reactive oxygen species (ROS) assay

The intracellular ROS assay was performed using an established method [17]. B16F10 cells were

cultured in 24-well plates and treated with various concentrations of *C. asiatica* EVs (1×10^9, 2.5×10^9,

and 5×10^9 particles/mL), Trolox® (0.05 mg/mL, positive control), or none (negative control) for 24

hours. The cells were then incubated with 24 mM H_2O_2 for 30 minutes to induce oxidative stress. After

incubation, 2', 7'-dichloro-fluorescein diacetate (DCFH-DA) was added to the cells and cultured for 30

minutes. After treatment, the cells were washed with PBS and trypsinized with trypsin/EDTA. The

fluorescence intensities of DCF were measured at an excitation wavelength of 504 nm and an emission

wavelength of 524 nm using a Fluoroskan Ascent fluorescent reader (Thermo Scientific, Vantaa,

Finland).

Total polyphenol content assay

130 Different concentrations of *C. asiatica* EVs (2×10^7, 4×10^7, 5×10^7 particles/mL) and gallic acid

131 standard were mixed with Folin-Ciocalteu reagent and incubated for 5 minutes. Sodium carbonate

132 solution was added and incubated for 30 minutes. The absorbance at 765 nm was measured using a

133 spectrophotometer and the gallic acid equivalent was calculated with the gallic acid standard curve

134 [18].

Metal chelating assay

The metal chelating activity of the extracellular vesicles was performed with potassium

ferricyanide ($K_3[Fe(CN)_6]$) assay [19]. Different concentrations of *C. asiatica* EVs (2×10^7, 5×10^7, 1×10^8

particles/mL) were mixed with 0.2 mM PBS and 1% (W/V) $K_3[Fe(CN)_6]$, with 2 mg/mL EDTA as the

positive control and deionized water as the blank control. The mixtures were heated at 50°C for 20

minutes. After cooling to room temperature, 10% (W/V) TCA was added, and the mixtures were

centrifuged at 12,000 rpm and 4°C for 5 minutes. The supernatants were transferred to a 96-well plate,

followed by the addition of deionized water and 1% (W/V) $FeCl_3$. The plates were incubated in the

dark for 30 minutes. Absorbance was measured at OD 700 nm to assess metal chelating activity.

Intracellular melanin content assay

The intracellular melanin content was measured as described previously [20]. B16F10 cells were

treated with α-Melanocyte-stimulating hormone (α-MSH, 100 nM) for 24 h to induce melanin

production. Following this, the cells were treated with different concentrations of *C. asiatica* EVs

(1×10^9, 2.5×10^9, and 5×10^9 particles/mL), or the positive control, arbutin (0.54 mg/mL), for an

additional 24 hours. Post-treatment, cell pellets containing a known number of cells were solubilized

in 1 N NaOH at 60°C for 60 minutes. The melanin content was then assayed at 405 nm.

Intracellular tyrosinase activity assay

The intracellular tyrosinase activity was determined as described in previous research [21].

B16F10 cells were treated with α-Melanocyte-stimulating hormone (α-MSH, 100 nM) for 24 hours to

induce melanin production. Following this, the cells were treated with different concentrations of *C.*

asiatica EVs (1×10^9, 2.5×10^9, and 5×10^9 particles/mL), or the positive control, arbutin (0.54 mg/mL),

for an additional 24 hours. After treatments, the cell extracts (100 μL) were mixed with freshly

prepared L-DOPA solution (0.1% in PBS) and incubated at 37°C. The absorbance at 490 nm was then

measured to assess the tyrosinase activity.

<u>Type I procollagen (PIP) ELISA assay</u>

Hs68 cells were seeded at 1×10^5 cells per well in 12 well plates and were cultured in 10% FBS

DMEM medium at 37°C with 5% CO_2 for 24 hours. After three washes with PBS, several different

concentrations (1×10^7, 1×10^8, 1×10^9) of *C. asiatica* EVs and 5 ng/mL TGF-β1 (AbCam Cat. #ab50036, as

positive control) were added and cultured for 24 hours. The level of type I procollagen in cell culture

supernatants was determined using a commercial PIP EIA Kit (TAKARA Cat. # MK101) according to the

manufacturer's instruction (2 step procedure).

<u>mRNA expression of aquaporin-3 (AQP3) and filaggrin (FLG) genes</u>

HaCat cells were dispensed into a 96-well plate and cultured for 24 hours under cell culture

conditions. Various concentrations of *C. asiatica* EVs and negative control (purified water only) were

added to the cells and cultured for 24 hours. Real-time PCR was performed to confirm the expression

of aquaporin-3 (AQP3) and filaggrin (FLG) at the gene level, and the test sequence is as follows. For

RNA isolation and cDNA synthesis, SuperPrep TM cell lysis & RT Kit for qPCR (TOYOBO, Cat. SCQ-101)

was used. To compare and analyze gene expression, the cDNA synthesized above was used as a

template, and Real-time PCR analysis was performed using Thunderbird TM SYBR qPCR Mix (TOYOBO,

Cat. QPS-201). The gene expression levels of the sample were quantified by normalization with

GAPDH using the $2^{-\triangle\triangle Ct}$ method.

<u>mRNA expression of pro-inflammatory genes after UVB induction</u>

HaCat cells (1×10^4/well for 96-well) or Hs68 cells (5×10^4/well for 24-well) were cultured for 24

hours and exposed to UVB at 90mJ dose. Various concentrations of *C. asiatica* EVs and negative

control (purified water only) were added to the cells and cultured for 4 hours. Real-time PCR was

performed to confirm the expression of cyclooxygenase-2 (COX) gene as above and the gene

expression levels were quantified by normalization with GAPDH using the $2^{-\triangle\triangle Ct}$ method.

<u>Measurement of nitric oxide production</u>

RAW 264.7 macrophage cells were seeded in 10% FBS DMEM medium in 24-well plates at a

density of 2×10^5 cells/well and were cultivated at 37°C with 5% CO_2 for 24 hours. Removing the

original medium, the cells were treated with 1×10^7, 1×10^8, 5×10^8, 1×10^9, 2×10^9, or 4×10^9 particles/mL

of *C. asiatica* EVs or gallic acid (10 and 20 μg/mL, Sigma-Aldrich Cat. #G7384) in serum-free DMEM

medium and incubated for 4 hours. After the pre-incubation, the cells were stimulated with 100

ng/mL LPS and incubated for 20 hours. The culture supernatants were harvested and centrifuged at

7,000 × g for 3 minutes. The samples were mixed with an equal volume (100 μL) of Griess reagent

(Sigma-Aldrich Cat. #G4410) and the absorbance was measured at 540 nm after reacting for 10

minutes.

Results

Isolation and characterization of *C. asiatica* EVs

C. asiatica EVs were isolated from tissue culture *C. asiatica* by a centrifugation and filtration

process. The size distribution of EVs harvested was analyzed with ZetaView® NTA system and showed

that the particle has a mean size of 150 nm (Figure 1A). The morphological features shown by the TEM

images can be observed as vesicles of approximately 100 nm in size with a lipid bilayer membrane

structure (Figure 1B).

Cytotoxicity assessment

To investigate the effect of *C. asiatica* EVs in cells, an MTT assay was used to evaluate the effect

of cell viability. B16F10 melanoma cells were treated with various concentrations of *C. asiatica* EVs

and MTT assay was performed. The results showed no significant differences compared to the control

group in this test (Figure 2). Thus, high concentrations of up to 5×10^9 particles/mL of *C. asiatica* EVs do

not exhibit cytotoxicity in B16F10 cells. In addition, we investigated the cytotoxicity in two human cell

lines using the alamarBlue® reagent. No discernable differences in cell viability were detected in either

human cell line, confirming that the EVs do not exhibit cytotoxicity in mouse or human cells (Figure

S1).

Antioxidant activity assessment

Here we first assessed the antioxidant activity of *C. asiatica* EVs by using two cell-free assays:

ABTS (Thiazoline-6-sulfonic acid) and potassium ferricyanide ($K_3[Fe(CN)_6]$) assays. At the higher

concentrations, *C. asiatica* EVs exhibited comparable $ABTS^{+\cdot}$ scavenging capacity to positive controls

vitamin C and BHA (Figure S2). The results indicated that the *C. asiatica* EVs can scavenge a significant

amount of $ABTS^{+\cdot}$ radical. For the $K_3[Fe(CN)_6]$ metal chelating assay, the results of the reducing power

test showed that the reducing capacities of EVs at concentrations of 2×10^6, 5×10^6, and 1×10^7

particles/mL were 16.18 ± 0.4%, 45.98 ± 1.64%, and 84.75 ± 0.11%, respectively (Figure S3). The

results indicated that the reducing power of *C. asiatica* EVs increases with the particle concentration,

exhibiting a dose-dependent effect.

A variety of naturally-occurring polyphenol compounds from *C. asiatica* extracts has been

studied, which contribute to its antioxidant properties [5]. Here we investigated the total polyphenol

content of *C. asiatica* EVs using an established method with the Folin-Ciocalteu reagent. The results

showed that the gallic acid equivalent of *C. asiatica* EVs increased with the particle number, indicating

that the antioxidant activity of the vesicles may be dose-dependent (Figure 3).

To further confirm the antioxidant capacity of *C. asiatica* EVs in a cellular environment,

intracellular reactive oxygen species (ROS) levels were measured by ROS scavenging rate using H_2O_2 at

20 mM to induce oxidative stress. After treatment, the intracellular ROS clearance was 53.86 ±

0.94%, 59.32 ± 1.68% and 66.27 ± 1.10% for 1×10^9, 2.5×10^9 and 5×10^9 (particles/mL) of *C. asiatica*

EVs, respectively (Figure 4). The results showed that all concentrations of *C. asiatica* EVs significantly

increased ROS scavenging activity compared to the control group (P<0.001), demonstrating effects
similar to positive control Trolox.

<u>Melanin inhibitory activity assessment</u>

In addition to its antioxidant properties, *C. asiatica* has potential skin whitening application
through inhibition of melanin synthesis [22]. First, we investigated its inhibition of melanin production
in B16F10 melanoma cells treated with α-melanocyte-stimulating hormone (α-MSH). After *C. asiatica*
EV treatment, the melanin content in the B16F10 cells was 93.55 ± 1.14%, 90.73 ± 1.17% and 83.69 ±
4.31% of control at 1×10^9, 2.5×10^9 and 5×10^9 particles/mL of *C. asiatica* EVs, respectively (Figure 5).
For the positive standard arbutin (0.54 mg/mL), the remaining intracellular melanin content was 71.95
± 2.44% of the control. The results indicated that *C. asiatica* EVs could decrease the melanin content in
B16F10 melanoma cells. Tyrosinase catalyzes the first two steps of melanin synthesis by hydroxylation
of L-tyrosine to L-DOPA [23]. Therefore, to investigate the change in melanin synthesis via tyrosinase
activity, we have carried out intracellular tyrosinase activity in B16F10 cells treated with α-MSH. After
treatment, the remaining intracellular tyrosinase activity was 92.5 ± 6.77%, 80.2.7 ± 5.13% and 56
± 3.83% for the 1×10^9, 2.5×10^9 and 5×10^9 particles/mL of either *C. asiatica* EVs, respectively (Figure 6).
In comparison, the remaining intracellular tyrosinase activity was 66 ± 10.53% of control after the cells
were treated with arbutin, a known skin-whitening compound. The results indicated that 5×10^8
particles/mL of *C. asiatica* EVs exhibited stronger inhibitory effect on α-MSH induced tyrosinase
activity in B16F10 cells than arbutin, albeit with higher intracellular melanin content than arbutin
(Figures 5 and 6).

<u>Effect of *C. asiatica* on skin integrity-related genes</u>

One of the major component in skin tissue is type I collagen, which is derived from its precursor
type I procollagen (PIP) secreted by skin fibroblasts, and lowered expression of PIP is associated with
skin UV damage and senescence leading to wrinkle formation [24]. In this experiment Hs68 fibroblast
cells were used to investigate the effect of induction of PIP secretion by *C. asiatica* EVs. Results
showed increased production of PIP in Hs68 cells in a dose-dependent manner, and induced a similar
expression level of PIP to that of positive control TGF-β1 at the highest concentration tested (1×10^9
particles/mL) (Figure S4). Aquaporin and filaggrin are important in barrier function of skin by
regulating water transport and water retention in skin [25]. We used a real-time PCR assay to quantify
the expression of mRNA expression of aquaporin-3 (AQP3) and filaggrin (FLG) genes in HaCat
keratinocyte cells treated with *C. asiatica* EVs. The results show that the addition of *C. asiatica* EVs to
HaCaT cell cultures was found to upregulate the expression levels of AQP3 and FLG genes compared to
the control group (Figure S5).

<u>Anti-inflammatory effects of of *C. asiatica* EVs</u>

One mechanism of the anti-melanogenic effect by *C. asiatica* is the inhibition of UV-induced
inflammation by triterpene madecassoside [22]. Here we used the expression levels of
pro-inflammatory gene COX2 to gauge the modulation of inflammation by *C. asiatica* EVs. Skin
fibroblast Hs68 and keratinocyte HaCaT cells were subjected to UV-induced inflammation and *C.*

asiatica EVs were then added, and expression levels of COX2 gene were measured. In HaCaT cells, EVs
reduced COX2 expression in a dose-dependent manner; however, in Hs68 even though the overall COX
expression reduced when the cells were treated with EV, no dose-dependent effect was observed
(Figure S6). Furthermore, as nitric oxide (NO) is an inflammatory response messengers secreted by
endothelial cells and macrophage, we subjected *C. asiatica* EV-treated RAW 264.7 macrophage culture
to lipopolysacchiaride (LPS), which is a TLR4 ligand and induces pro-inflammatory response [26]. The
NO production was measured by Griess reagent, which is converted to a pink color after reacting with
nitrites. As shown in Figure 7, the amount of nitrite decreases in a dose-dependent manner in cells
treated with increasing concentrations of EVs. Taken together, *C. asiatica* EVs were able to reduce
inflammatory response in a variety of cell types tested.

Discussion

In this study, we have isolated and characterized EVs secreted from *C. asiatica* tissue culture. The
EVs have a size of approximately 160 nm and a lipid bilayer morphology similar to that previously
reported for exosomes derived from *C. asiatica* and other plants [12, 13]. In a series of *in-vitro*
experiments in B16F10 cell culture, we have demonstrated that the EVs are safe for cells and that EVs
exhibited notable antioxidant, anti-melanogenic, and anti-inflammatory effects, suggesting potential
applications in skin care and therapeutic treatments to combat oxidative stress and
hyperpigmentation. Another study also isolated exosome-like EVs from *C. asiatica* but with a smaller
average particle size of 63.8 nm [14]. While the EVs from the above study exhibited cytotoxicity in
HepG2 human cancer cells, we did not observe any effect on cell viability in mouse melanoma B16F10
cells or HaCaT and Detroit 551 human cell lines tested (Figures 2 and S1) [14]. The therapeutic
properties of *C. asiatica* have mostly been attributed to the triterpenes and derivatives, including
madecassoside, madecassic acid, asiaticoside, and asiatic acid, and these are among the most
well-studied compounds in the plant [3-5]. In our study, we observed potent antioxidant effects from
C. asiatica, although further study is required to determine specifically how contents within EVs
(chemicals, proteins, and nucleic acids) can contribute to observed effects. In addition to
phytochemicals and proteins, small RNAs and miRNAs in EVs are important modulators of target effect
proteins, and RNA binding proteins, such as Argonaute 1, RNA helicases, and Annexins, may be
involved in packing RNA into EVs [10, 28] A recent preprint study has provided a comprehensive
analysis of the transcriptome human skin keratinocytes after treatment of *C. asiatica* exosomes,
revealing 11 novel miRNAs packed within the exosomes [13]. The above study also found that skin
integrity and barrier-related genes were upregulated after treatment by *C. asiatica* EVs, which we
have also observed in this study with increasing gene expression of PIP1, AQP3, and FLG (Figures S4
and S5) [13]. The study also found that *C. asiatica* exosomes induced higher gene expression than *C.*
asiatica extracts, possibly due to better penetration of skin cells by exosomes and the presence of
miRNAs [13]. Thus *C. asiatica* EVs could provide its effect via multitude of effectors such as
phytochemicals and miRNAs.

In *C. asiatica*, madecassoside by itself has the ability to inhibit UV-induced melanin synthesis in a keratinocyte and melanocyte co-culture system as well as artificially tanned human skin [22]. Other plant-derived EVs also have anti-melanogenic effect. For example, *Dendropanax morbifera* EVs can inhibit the expression of melanogenesis genes such as MITF, TYR, TRP-1, and TRP-2 [9, 27]. In our study, the EVs appeared to modulate melanogenesis pathways in cell culture, likely through the inhibition of tyrosinase activity, which is a key enzyme in melanin synthesis [23]. The mechanism of anti-melanogenesis by *C. asiatica* include inhibition of inflammation, as observed in previous studies [24]. and in our study, where we have observed that *C. asiatica* can reduce pro-inflammatory gene expression, such as COX2 and decrease the production of NO (Figures 7 and S6). As melanin overproduction contributes to skin conditions like freckles and nevus formation, and age spot development, plant-derived EVs with anti-melanogenic properties represent an economical and natural alternative to conventional treatments, which often come with side effects or limited efficacy [27].

The tissue culture micropropagation technology for *C. asiatica* has been established as a sustainable source of *C. asiatica* [28]. This ensures its conservation in nature while providing a constant yield of raw material, akin to the use of bioreactors in biopharmaceutical drug and vaccine production. Using plant tissue culture as a source for EV production presents several advantages over traditional methods of harvesting from wild plants. First, tissue culture allows for consistent, scalable production of EVs with genetic uniformity, bypassing the variability associated with environmental factors that affect wild plants, such as seasonal changes, soil quality, and climate/geographical conditions [15, 28]. This standardization is particularly important for achieving reproducible results in therapeutic and cosmetic applications, where consistent quality and efficacy are crucial in ensuring the safety and efficacy of the product. Additionally, tissue culture can be optimized to enhance the yield of specific bioactive compounds through controlled environmental conditions and the use of elicitors [15].

In conclusion, this study investigated the antioxidant, anti-melanogenic and anti-inflammatory potential of *C. asiatica*-derived EVs from tissue culture. The advantages of tissue culture over wild plants in terms of sustainability, consistency, and the ability to enhance bioactive compound production further support the use of this approach in developing plant-based EV therapies. Future studies could focus on characterizing bioactive molecules and miRNAs within these EVs, as well as exploring their safety and efficacy in *in-vivo* models. Ultimately, this work paves the way for the sustainable production and utilization of plant-derived EVs in skin care cosmetics and wound healing applications.

Acknowledgements

We thank Dr. Meei-Yun Lin of Schweitzer Biotech Company for reading and providing feedback for preparation of the manuscript.

Competing Interests

C.-C. W., C.-H. C., P.-L. K., W.-H. T., L. T.-C. L., C. C., and T.-Y. K. are employees of Schweitzer Biotech Company. W.-Y. Q. and I. P. are consultants employed by Schweitzer Biotech Company. T.-M. C. and S.-S. W. have no competing interests to declare. This study is funded by Schweitzer Biotech Company.

References

1. Biswas D, Mandal S, Chatterjee Saha S, Tudu CK, Nandy S, Batiha GE, Shekhawat MS, Pandey DK, Dey A (2021) Ethnobotany, phytochemistry, pharmacology, and toxicity of Centella asiatica (L.) Urban: A comprehensive review. Phytother Res 35:6624–6654. https://doi.org/10.1002/ptr.7248

2. Sabaragamuwa R, Perera CO, Fedrizzi B (2018) Centella asiatica (Gotu kola) as a neuroprotectant and its potential role in healthy ageing. Trends Food Sci Technol 79:88-97. https://doi.org/10.1016/j.tifs.2018.07.024

3. Bandopadhyay S, Mandal S, Ghorai M, Jha NK, Kumar M, Radha, Ghosh A, Proćków J, Perez de la Lastra JM, Dey A (2023) Therapeutic properties and pharmacological activities of asiaticoside and madecassoside: A review. J Cell Mol Med 27(5):593-608. https://doi.org/10.1111/jcmm.17635

4. Sun B, Wu L, Wu Y, Zhang C, Qin L, Hayashi M, Kudo M, Gao M, Liu T (2020) Therapeutic potential of Centella asiatica and its triterpenes: a review. Front Pharmacol 11:568032. https://doi.org/10.3389/fphar.2020.568032

5. Diniz LR, Calado LL, Duarte AB, de Sousa DP (2023) Centella asiatica and its metabolite asiatic acid: wound healing effects and therapeutic potential. Metabolites 13(2):276. https://doi.org/10.3390/metabo13020276

6. Jeppesen DK, Zhang Q, Franklin JL, Coffey RJ (2023) Extracellular vesicles and nanoparticles: emerging complexities. Trends Cell Biol 33(8):667-81. https://doi.org/10.1016/j.tcb.2023.01.002

7. Moghassemi S, Dadashzadeh A, Sousa MJ, Vlieghe H, Yang J, León-Félix CM, Amorim CA (2024) Extracellular vesicles in nanomedicine and regenerative medicine: A review over the last decade. Bioact Mater 36:126-56. https://doi.org/10.1016/j.bioactmat.2024.02.021

8. Yang GH, Lee YB, Kang D, Choi E, Nam Y, Lee KH, You HJ, Kang HJ, An SH, Jeon H (2021) Overcome the barriers of the skin: exosome therapy. Biomater Res 25(1):22. https://doi.org/10.1186/s40824-021-00224-8

9. Li A, Li D, Gu Y, Liu R, Tang X, Zhao Y, Qi F, Wei J, Liu J (2023) Plant-derived nanovesicles: Further exploration of biomedical function and application potential. Acta Pharm Sin B 13(8):3300-20. https://doi.org/10.1016/j.apsb.2022.12.022

10. Zhu Y, Zhao J, Ding H, Qiu M, Xue L, Ge D, Wen G, Ren H, Li P, Wang J (2024) Applications of plant-derived extracellular vesicles in medicine. MedComm. 2024 5(10):e741.

https://doi.org/10.1002/mco2.741

11. Urzì O, Gasparro R, Ganji NR, Alessandro R, Raimondo S (2022) Plant-RNA in extracellular
vesicles: the secret of cross-kingdom communication. Membranes 12(4):352.
https://doi.org/10.3390/membranes12040352

12. Cho JH, Hong YD, Kim D, Park SJ, Kim JS, Kim HM, Yoon EJ, Cho JS (2022) Confirmation of
plant-derived exosomes as bioactive substances for skin application through comparative
analysis of keratinocyte transcriptome. Appl Bio Chem 65(1):8.
https://doi.org/10.1186/s13765-022-00676-z

13. Kim MH, Yoon EJ, Kim JS, Park SJ, Lee H (2023) Comparative analysis of the transcriptome
and efficacy of bioactive Centella asiatica exosomes on skin cells. ResearchSquare
https://doi.org/10.21203/rs.3.rs-2787704/v1

14. Huang J, Cao X, Wu W, Han L, Wang F (2024) Investigating the proliferative inhibition of
HepG2 cells by exosome-like nanovesicles derived from Centella asiatica extract through
metabolomics. Biomed Pharmacother 176:116855.
https://doi.org/10.1016/j.biopha.2024.116855

15. Kumari S, Kumar N, Trivedi MP (2022) Application of Plant Tissue Culture for Improvement
of Centella asiatica. In: Kumar N (ed) Biotechnology and Crop Improvement, 1st ed. CRC
Press, Boca Raton, FL, pp. 93-106. https://doi.org/10.1201/9781003239

16. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C (1999) Antioxidant
activity applying an improved ABTS radical cation decolorization assay. Free Radic Biol Med
26(9-10):1231-7. https://doi.org/10.1016/S0891-5849(98)00315-3

17. Murrant CL, Reid MB (2001) Detection of reactive oxygen and reactive nitrogen species in
skeletal muscle. Microsc Res Tech 55(4):236-48. https://doi.org/10.1002/jemt.1173

18. Singleton VL, Orthofer R, Lamuela-Raventós RM (1999) Analysis of total phenols and other
oxidation substrates and antioxidants by means of folin-ciocalteu reagent. In: Packer L (ed)
Methods in enzymology. Academic Press, Cambridge MA, Vol. 299, pp. 152-178.
https://doi.org/10.1016/S0076-6879(99)99017-1

19. Adjimani JP, Asare P. (2015) Antioxidant and free radical scavenging activity of iron chelators.
Tox Rep 2:721-8. https://doi.org/10.1016/j.toxrep.2015.04.005

20. Tsuboi T, Kondoh H, Hiratsuka J, Mishima Y (1998) Enhanced melanogenesis induced by
tyrosinase gene-transfer increases boron-uptake and killing effect of boron neutron capture
therapy for amelanotic melanoma. Pigment Cell Res 11(5):275-82.
https://doi.org/10.1111/j.1600-0749.1998.tb00736.x

21. Yang JY, Koo JH, Song YG, Kwon KB, Lee JH, Sohn HS, Park BH, Jhee EC, Park JW (2006)
Stimulation of melanogenesis by scoparone in B16 melanoma cells 1. Acta Pharmacol Sin
27(11):1467-73. https://doi.org/10.1111/j.1745-7254.2006.00435.x

22. Jung E, Lee JA, Shin S, Roh KB, Kim JH, Park D (2013) Madecassoside inhibits melanin
synthesis by blocking ultraviolet-induced inflammation. Molecules 18(12):15724-36.

419   https://doi.org/10.3390/molecules181215724

420 23. Snyman M, Walsdorf RE, Wix SN, Gill JG (2024) The metabolism of melanin synthesis—From
421    melanocytes to melanoma. Pigment Cell Melanoma Res 37(4):438-452.
422    https://doi.org/10.1111/pcmr.13165

423 24. Talwar HS, Griffiths CE, Fisher GJ, Hamilton TA, Voorhees JJ (1995) Reduced type I and type
424    III procollagens in photodamaged adult human skin. J Invest Dermatol 105(2):285-90.
425    doi: 10.1111/1523-1747.ep12318471

426 25. Park CH, Min SY, Yu HW, Kim K, Kim S, Lee HJ, Kim JH, Park YJ (2020) Effects of apigenin on
427    RBL-2H3, RAW264. 7, and HaCaT cells: anti-allergic, anti-inflammatory, and skin-protective
428    activities. Int J Mol Sci 21(13):4620. https://doi.org/10.3390/ijms21134620

429 26. Mantovani A, Garlanda C (2023) Humoral innate immunity and acute-phase proteins. N Engl
430    J Med 388(5):439-52. https://doi.org/10.1056/NEJMra220634

431 27. Lee R, Ko HJ, Kim K, Sohn Y, Min SY, Kim JA, Na D, Yeon JH (2020) Anti-melanogenic effects of
432    extracellular vesicles derived from plant leaves and stems in mouse melanoma cells and
433    human healthy skin. J Extracell Vesicles 9(1):1703480.
434    https://doi.org/10.1080/20013078.2019.1703480

435 28. Mohapatra P, Ray A, Sandeep IS, Nayak S, Mohanty S (2021) Tissue-Culture-Mediated
436    Biotechnological Intervention in Centella asiatica: A Potential Antidiabetic Plant. In: Gantait
437    S, Verma SK, Sharangi AB (eds) Biotechnology of Anti-diabetic Medicinal Plants. Spring,
438    Singapore, pp. 89-116. https://doi.org/10.1007/978-981-16-3529-8_4

Figures

A

B

Fig. 1 Size and morphology characterization of *C. asiatica*-derived EVs. A. Nanoparticle tracking analysis (NTA) of size and concentration distribution of EVs by using the ZetaView® system. B. Representative TEM images of purified *C. asiatica* EVs with red arrows showing vesicles with lipid bilayer membrane structure. Scale bar: 100 nm

Fig. 2 Cell viability assay. B16F10 cells were treated with different concentrations of *C. asiatica* EVs. After 24h treatment, cell viability was measured by the MTT assay method. Data are presented as mean ± SD (n=3). Statistical results were calculated by one-way ANOVA with Dunnett's multiple comparisons test compared to the control group.

Total polyphenol content

Fig. 3 Total polyphenol content. The total polyphenol content of *C. asiatica* EVs using the Folin-Ciocalteu method. Data are presented as mean ± SD (n=3). Statistical results were calculated by one-way ANOVA with Dunnett's multiple comparisons test compared to the control group). ****P<0.0001

Intracellular ROS scavenging assay

Fig. 4 Intracellular ROS level. B16F10 cells were pretreated with various concentrations of *C. asiatica* EVs, Trolox (0.05 mg/mL, positive control), or left untreated as a control for 24 hours. The cells were then incubated with 24 mM H_2O_2 and DCFH-DA. The fluorescence intensities of DCF were measured to calculate the ROS levels. Data are presented as mean ± SD (n=3). Statistical analysis was calculated by one-way ANOVA with Dunnett's multiple comparisons test, compared to the control group. ****P<0.0001

Fig. 5 Intracellular melanin content. Inhibitory effects of different concentrations of *C. asiatica* EVs on melanin content in B16F10 cells. Data are presented as mean ± SD (n=3). Statistical results were calculated by one-way ANOVA with Dunnett's multiple comparisons test compared to the control group). *P<0.05, **P<0.01, ****P<0.0001

Fig. 6 Intracellular tyrosinase activity. Inhibitory effects of different concentrations of *C. asiatica* EVs on intracellular tyrosinase activity in B16F10 cells. Data are presented as mean ± SD (n=3). Statistical results were calculated by one-way ANOVA with Dunnett's multiple comparisons test (compared to the control group). *P<0.05, ***P<0.001, ****P<0.0001

Fig. 7 Nitric oxide (NO) production in RAW 264.7 macrophages. Cells were treated with different concentrations of *C. asiatica* EVs or gallic acid after induction by LPS. The resulting NO produced was detected by the Griess reagent.