The oncogene SLC35F2 is a high-specificity transporter for the micronutrients queuine and queuosine

Lyubomyr Burtnyak[a], Yifeng Yuan[b], Xiaobei Pan[c], Lankani Gunaratne[d,e], Gabriel Silveira d'Almeida[d], Maria Martinelli[b&], Colbie Reed[c$], Jessie Fernandez Garcia[b], Bhargesh Indravadan Patel[f], Isaac Marquez[g], Ann E. Ehrenhofer-Murray[f], Manal A. Swairjo[g], Juan D. Alfonzo[d], Brian D. Green[c], Vincent P. Kelly[a*] and Valérie de Crécy-Lagard[b,g*]

[a]School of Biochemistry & Immunology, Trinity Biomedical Sciences Institute, Trinity College Dublin, Dublin 2, Ireland

[b]Department of Microbiology and Cell Science, University of Florida, Gainesville, Florida 32611, USA

[c]School of Biological Sciences, Institute for Global Food Security, Queen's University Belfast, Belfast, UK

[d]Department of Molecular Biology, Cell Biology and Biochemistry and the Brown RNA Center, Brown University, Providence, Rhode Island 02903, USA

[e]Ohio State Biochemistry Program, The Ohio State University, Columbus, OH, USA

[f]Institut für Biologie, Lebenswissenschaftliche Fakultät, Humboldt-Universität zu Berlin, 10115 Berlin, Germany

[g]Department of Chemistry and Biochemistry, San Diego State University, San Diego, California 92182, USA

[h]Genetic Institute, University of Florida, Gainesville, Florida 32611, USA

[&]Current address: eSTEAMed Learning Inc., Maitland, FL 32751, USA

[$]Current address: Data Science Institute, Medical College of Wisconsin, Milwaukee, Wisconsin 53226, USA

Corresponding authors: Valérie de Crécy-Lagard and Vincent P. Kelly

Email: vcrecy@ufl.edu; kellyvp@tcd.ie

Author Contributions:

Conceptualization, VdC-L, VPK; methodology, VdL, VPK, JA, MAS, AEM; formal analysis VdC-L, VPK; investigation, LB, YY, XP, BDG, LG,GSdA, MM, JFG, CR. BIP, IM; writing original draft, YY, LB, AEM, VdC-L and VPK.; funding acquisition, VdC-L, VPK, MAS, JA, BDG, AEM.

Competing Interest Statement: The authors declare no competing interests.

Classification:

Major Classification: Biological Sciences

Minor Classification: Biochemistry

Keywords:

Queuine, Queuosine, transporter, *SPCC320.08*, Qtp1

This PDF file includes:

    Main Text
    Figures 1 to 4

Abstract

The nucleobase queuine (q) and its nucleoside queuosine (Q) are micronutrients derived from bacteria that are acquired from the gut microbiome and/or diet in humans. Following cellular uptake, Q is incorporated at the wobble base (position 34) of tRNAs with a GUN anticodon, which is important for efficient translation. Early studies suggested that cytosolic uptake of queuine is mediated by a selective transporter that is regulated by mitogenic signals, but the identity of this transporter has remained elusive. Here, through a cross-species bioinformatic search and genetic validation, we have identified the solute carrier family member SLC35F2 as a unique transporter for both queuine and queuosine in *Schizosaccharomyces pombe* and *Trypanosoma brucei*. Furthermore, gene disruption in HeLa cells revealed that SLC35F2 is the sole transporter for queuosine in HeLa cells (K_m 174 nM) and a high-affinity transporter for the queuine nucleobase (K_m 67 nM), with the presence of another low-affinity transporter (K_m 259 nM) in these cells. Competition uptake studies show that SLC35F2 is not a general transporter for other canonical ribonucleobases or ribonucleosides, but selectively imports q and Q. The identification of SLC35F2, an oncogene, as the transporter of both q and Q advances our understanding of how intracellular levels of queuine and queuosine are regulated and how their deficiency contributes to a variety of pathophysiological conditions, including neurological disorders and cancer.

Significance Statement

The discovery of SLC35F2 as the eukaryotic transporter of queuine and queuosine is key to understanding how these micronutrients are salvaged from the human gut and distributed to different body tissues. Queuosine modification of tRNAs enhances the accuracy and efficiency of codon-anticodon pairing and regulates a range of biological and pathophysiological states, including oxidative stress responses, cancer, learning, memory, and gut homeostasis.

Introduction

Human health is deeply intertwined with the acquisition of sufficient daily nutrients. This encompasses not only adequate caloric intake, but also a diverse array of essential vitamins, minerals, and other micronutrients necessary for optimal cellular function, disease prevention, and overall well-being (1). Queuine (q) and its nucleoside derivative queuosine (Q) are one such group of micronutrients that are synthesized exclusively by bacteria and salvaged by eukaryotic organisms including plants, fungi, yeast, and metazoan species, with q undergoing subsequent incorporation into tRNAs decoding His, Tyr, Asp, and Asn codons and serving as a substrate for further glycosylation, turnover, and salvage (2) (Fig. 1). Currently, no transporters for intracellular uptake of Q nor q have been identified in any eukaryote and, in metazoans, the mechanism of uptake from the gut and redistribution throughout the body is poorly understood.

In humans, the circulating levels of q are estimated to be in the range of 1-10 nM (3) recent studies indicating a mean serum concentration of 8 nM in females and 6.8 nM in males (4). Early studies in fibroblasts and mammalian cell lines showed that a specific cellular uptake mechanism exists to transport q from the extracellular space to the cytosol. Using a tritiated q derivative (rQT_3), it was shown that uptake in human foreskin fibroblasts can occur *via* two different transporters, a low K_m transporter (30 nM) that reached saturation after 2 minutes (13.5 pmol/10^6 cells/h) and a slower, high K_m mechanism (350 nM) that displayed a linear uptake after 3-4 hrs (2.3 pmol/10^6 cells/h). Moreover, competitive uptake experiments with a variety of purine and purine-derivatives did not affect q transport, suggesting that the transporter is specific to q (5).

Earlier studies indicated that mitogenic signalling pathways could modulate q transport
(6). In fibroblasts, low doses and short-term exposure to the Protein Kinase C (PKC) activator
phorbol-12, 13-didecanoate (PDD) led to increased rQT_3 uptake, whereas extended exposure
durations inhibited uptake. In addition, the protein phosphatase inhibitors calyculin A and okadaic
acid were found to increase rQT_3 uptake but avoidance of excess exposure time and
concentrations were necessary [6]. Later, other PKC activators, including phosphatidylserine,
calcium ionophore and diolein were also found to increase rQT_3 uptake (7). Furthermore, growth
factors such as epidermal growth factor, fibroblast growth factor and platelet-derived growth
factor were also shown to increase q uptake, both alone and in combination with PKC activators
(7). Notably, although a major portion of q transport appears to be regulated by PKC and
mitogenic signalling, a substantial basal rate of q uptake exists, which operates independently of
exogenous signalling. This implies the presence of two or more transporters for q uptake, one of
which may be PKC- and mitogen-regulated.

Here we show that SLC35F2 functions as the exclusive transporter for q and Q in
Schizosaccharomyces pombe and the human parasite *Trypanosoma brucei.* In the human HeLa
cell line, it serves as the primary high-affinity transporter for q and as the unique transporter for Q.
Competitive uptake studies reveal that SLC35F2 does not recognize canonical nucleobases nor
nucleosides, since they were unable to disrupt q/Q import into HeLa cells. Defining the role of the
SLC35F2 transporter, an identified oncogene, as the q/Q transporter has significance for
understanding the uptake, distribution and function of this micronutrient in eukaryotes.

Results

Prediction of a candidate Q/q transporter family by comparative genomics

The Q modification of tRNA has been independently lost in several eukaryotic clades including
the model organism *Saccharomyces cerevisiae*, while being present in most sequenced fungal
genomes including *S. pombe* (8). The presence/absence of Q can be inferred by the
presence/absence of the signature gene *Qtrt1*, which encodes tRNA guanine transglycosylase
that exchanges G for Q at position 34 in tRNA. Previous phyletic profile searches to identify the
missing q/Q transporter gene(s) by searching for protein families that co-distribute with *Qtrt1* were
performed across three eukaryotic kingdoms (fungi, viridiplantae and metazoa) (8). However,
these searches only returned candidate genes for the already characterized *Qtrt1* and *Qng1*
genes, the latter being responsible for the release of q from the Q precursor queuosine 5′-
monophosphate (Q-5′MP). As an alternate strategy, we analyzed protein families containing
transmembrane domains separately in four different taxonomic clades (Sar, Viridiplantae,
Metazoa and Fungi), each including around 200 organisms encoding QTRT1 (QTRT[+]) and at
least 10 organisms that are QTRT1 deficient (QTRT[-]) (Fig. S1 Step I). The proteins in each of
these taxonomic groups containing at least four transmembrane (TM) domains covering 25% or
more of their length were clustered based on sequence similarity as described in the methods
section (Fig. S1 Step II). The choice of four TM domains was because 96.4% of all known
electrochemical potential-driven transporters have at least four TM domains (9). Each cluster
within a given group was given a probability score that would be 1 if all the proteins of the cluster
were encoded by QTRT[+] organisms and absent in QTRT[-] organisms as described in the methods
section and in Fig. S1 (Step III).

The top candidate in the fungi clade, which also scored highly in the other three analyzed
taxonomic groups, was the IPR052221 annotated in InterPro database as the "SLC35F solute
transporter family" (Table S1**),** but corresponds to the "solute carrier family 35 member F1/2" or
the "SLC35F1/F2" KEGG orthology group (K15287)(10). This family of transporters contains 10
transmembrane helices (Fig. S1A**)** and is a subgroup of the SLC35 superfamily (11). Most
members of this family transport nucleoside di-phosphate (NDP) sugars, but a few transport

vitamins such as thiamin (12). To date, the substrate specificity of the SLC35F1/F2 subgroup is
not known (13).

SLC35F1/F2 proteins were present in 184 out of 189 QTRT[+] fungi and absent in all of
189 QTRT[-] Fungi. The gene encoding a SLC35F protein in *S. pombe*, *SPCC320.08*, is co-
expressed with ribosome biogenesis genes (https://coxpresdb.jp/locus/?gene_id=2539011).
Another published co-expression analysis along the *S. pombe* cell cycle (14) found that
SPCC320.08 was the only transporter gene in a cluster of co-expressed genes highly enriched
(21 out of 24, Table S2) in rRNA processing and translation related proteins.

We constructed a sequence similarity network (SSN) of the IPR052221 family proteins to
dissect their functions (Fig. 2A). The Chordata F1 and F2 paralogous subgroups could not be
separated with an alignment score threshold (AST) of 110 (Fig. 2A), but the divergence becomes
evident with an AST of 125 (Fig. 2B), indicating similar functions. Members of this family separate
mostly following taxonomic lineages except for the Viridiplantae SLC35F proteins, which are split
in two groups: one that contains members from QTRT[-] organisms, such as *Arabidopsis thaliana*
(**Table S3**), and one that lacks *A. thaliana* homologs but contains members only from QTRT[+]
organisms, such as *Zea mays.* Indeed*,* the *Z. mays* genome encodes the full Q salvage pathway
(Uniprot accession: QTRT1, A0A3L6E559; QTRT2, A0A8J8Y6E7 and Qng1, A0A3L6FZJ1) while
A. thaliana is known to lack Q in tRNA, as this wobble base modification has been lost in
cruciferous plants (8). The identification of distinct Viridiplantae paralogous groups, one with no *A.*
thaliana member, suggested that the SLC35F family is not isofunctional and could also explain
why this transporter family was not identified in previous phylogenetic distribution searches that
were filtering for transporter families present in *Zea mays* and absent in *A. thaliana* (8). Analyses
of the co-expression data available for the two human SLC35F genes link SLC35F2 to ribosome
biogenesis and RNA processing, but associate SLC35F1 to neural and synaptic function (Fig.
S2). In addition, overexpression of SLC35F1 in HeLa cells recently showed that the transporter
resides exclusively on recycling endosomes (15). This would suggest that SLC35F1 is likely not
the plasma membrane transporter for q/Q. The combination of *in silico-derived* evidence
(presence/absence correlated with QTRT1, co-expression with RNA processing proteins and
membrane protein topology) led us to test the hypothesis that SLC35F2 is one of the missing
human Q/q transporters.

The human *SLC35F2* gene encodes the Q/q transporter.

To investigate whether the SLC35F2 protein is required for q or Q transport in humans, a
knockout approach was adopted. The SLC35F2 gene is located on the reverse strand of
chromosome 11, comprising eight exons that encode a 374-amino acid protein. CRISPR-Cas9
was used to insert a promoter trap into exon two of the gene by homologous recombination —
providing puromycin resistance under the control of the SLC35F2 promoter (Fig. S3A). From the
clones surviving puromycin treatment, four were selected and genotyped. All four clones were
homozygous knockouts for SLC35F2, with either no PCR amplicon observed for the wild-type
allele or a band of the incorrect size (Fig. S3B). Strikingly, 3-AcrylamidoPhenylBoronic acid (APB)
Northern blot analysis, which can separate Q modified from unmodified tRNAs (16), revealed that
in the absence of SLC35F2, all four HeLa cell lines were devoid of detectable Q modification in
tRNA[His] after incubation with Q (25 nM), contrasting with wildtype cells, where approximately fifty
percent of the tRNA was Q-modified (Fig. 3A, upper panel). In the case of q nucleobase (25 nM),
in wild-type cells, all the tRNA was found to be fully Q-modified with the addition of the
micronutrient at a concentration of 25 nM, but a dramatic reduction in modification was observed
in the SLC35F2 knockout clones (Fig. 3A, lower panel). To validate and extend this finding, Clone
1 was selected for further analysis by treating cells with increasing concentrations (25 to 250 nM)
of either the q-base or Q-nucleoside (Fig. 3B). Notably, at all concentrations tested Q failed to
modify the tRNA in the SLC35F2 knockout cells (upper panel), whereas in the case of q, almost a

ten-fold higher concentration was required, relative to wild-type cells, to completely modify the
tRNA (lower panel). These data show that SLC35F2 is a unique transporter for Q-nucleoside in
human cells and the primary high-affinity transporter for the uptake of q base.

The SLC35F2 homologs of *T. brucei* and *S. pombe* are the sole q/Q transporters in these
organisms.

Having shown that SLC35F2 was required for both q and Q transport in human cells, we
investigated if this was also the case in eukaryotes that encode only one SLC35F protein (Fig.
3C-D). We focused on two single-cell eukaryotes where Q synthesis and function have been
extensively studied: the fission yeast *S. pombe* (8, 17–20) and the parasite *T. brucei* (21–23).

The fission yeast *S. pombe* encodes an SLC35F1/F2 homolog SPCC320.08, (here
renamed Qtp1 [**Q**ueuosine and queuine **t**rans**p**orter 1]), which has no apparent homolog in
Saccharomyces cerevisiae (PomBase,(24)). To test whether Qtp1 transports q and/or Q in *S.*
pombe, a *qtp1Δ* strain was constructed, and Q levels analyzed via APB Northerns in tRNA^{Asp}
extracted from cells grown in medium supplemented with either q or Q (**Fig. 3C**). In the presence
of 10 nM q, the wildtype cells showed partial Q modification of tRNA^{Asp}, and 100 nM q resulted in
full modification, as determined by APB Northern blotting. In contrast, in the *qtp1Δ* strain, tRNA^{Asp}
showed no Q modification of the tRNA even at the higher q concentration. Similarly, the *qtp1Δ*
strain showed no Q modification of tRNA^{Asp} when provided with the Q nucleoside, whereas the
wildtype strain showed full modification.

To study the function of the SLC35F2 homolog in *T. brucei* (ORF Tb927.4.320), here
renamed Qtp1 as above for *S. pombe*, both alleles of the corresponding gene were deleted using
the CRISPR/Cas9 system (Fig. S4). This gene was not essential in *T. brucei* and growth was not
affected (Fig. S5) in agreement with the fact that Q in tRNA is not essential for growth in this
organism (21–23). We then extracted total RNA from wild type and knockout cells lines and
analyzed Q-content in tRNA^{Tyr} via APB Northern blotting (Fig. 3D**)**. For this analysis, Q-content
from cells grown in complete media (non-dialyzed serum) were compared to cells grown in media
containing dialyzed serum. No Q-tRNA was detected in cells grown with dialyzed serum,
regardless of the presence or absence of Qtp1. This is expected given that serum is the only
source of q/Q for *T. brucei* when grown in culture. Similar experiments were then performed by
supplementing the dialyzed serum with q or Q (at 25 nM concentration, respectively). In this case,
only wild-type cells showed detectable Q-tRNA, while Q-tRNA was undetectable in the *qtp1*
knockout (Fig. 3D) even when the cultures were supplemented with 250 nM q (Fig. S6) or when
the knockout cells were grown in complete media (**Fig. 3D**).

Taken together, these experiments show that in two organisms encoding only one
member of the SLC35F2/Qtp1 family, this protein is the sole protein responsible for q and Q
import.

230
Kinetic analysis of SLC35F2 q/Q transporter activity.
Recent studies to determine the circulating levels of q in humans indicate a range of 1-10 nM,
with higher levels reported in females (4, 25) but a surprising absence of detectable Q [4]. It
therefore was of interest to determine the substrate affinity and specificity of SLC35F2, to provide
an indication of its physiological activity. Of note, the SLC35F2 solute carrier has previously been
identified as a unique, high-specificity transporter for the small molecule anti-cancer agent YM155
(sepantronium bromide), with retroviral disruption in KBM7 cells conferring drug resistance to this
molecule (26).
To negate the interference of multiple downstream steps, the uptake kinetics of SLC35F2
for Q was determined in HeLa cells that were knocked out for QNG1 and therefore are incapable

of converting Q to q (Fig. 4A, upper panel). Intracellular Q and Q-5′MP were quantified by LC-
MS/MS and their additive concentration was used to measure uptake kinetics, providing a K_m of
174 nM and a Vmax of 0.87 pmol/min (lower panel). To examine q uptake, HeLa cells deficient in
Qtrt1 were exploited (Fig. 4B, upper panel), providing a K_m of 67 nM for q uptake through both the
SLC35F2 transporter and the unidentified secondary mechanism (lower panel). To obtain further
data on the uptake kinetics of the unidentified transporter, we took advantage of the fact that
YM155 is a high-specificity substrate for SLC35F2 uptake and thus capable of competitively
inhibiting the transporter (Fig. 4C, upper panel) (26). Saturating concentrations of YM155 were
added to the Qtrt1 deficient HeLa cells to inhibit q uptake, revealing that the second transporter
for q has a higher K_m of 260 nM (lower panel), which agrees with previous literature wherein both
a low-K_m and high-K_m transporter for the q nucleobase had been demonstrated (5). Together, the
data indicate that the SLC35F2 transporter is uniquely tasked for Q uptake in humans with a K_m
that is comparatively high relative to that of q and which could indicate differential roles for the
transporter in the uptake of each form of the micronutrient.
255
The SLC35F2 transporter is highly specific for q/Q micronutrient uptake.
Earlier studies showed that transporters for q are specific, and that uptake cannot be
outcompeted by a variety of nucleobases or nucleotides (5). Using the level of Q-modified tRNA
as an indirect indicator of the ability of cells to import Q or q, competitive uptake studies were
performed using a range of nucleosides and nucleobases. In all cases in wild-type cells, none of
the canonical nucleosides were capable of competing against Q-nucleoside uptake (Fig. 4D).
Similarly, none of the tested nucleobases competed against q uptake (Fig. 4E) at saturating
concentrations (25 μM versus 25 nM) except for the anti-cancer drug YM155, which effectively
outcompeted Q/q import. This further validates SLC35F2 as the high-specificity transporter for
both q and Q, both of which possess a cyclopentene-diol moiety not shared with canonical
nucleobase or nucleoside molecules.
267
Structural insights into SLC35F2/Qtp1 proteins.
Human SLC35F2, *T. brucei* Qtp1 (ORF Tb927.4.320), and *S. pombe* Qtp1 (*SPCC320.08)* are all
members of the SLC35F subfamily of the large solute carrier family 35 (SLC35). SLC35 proteins
are highly conserved TM proteins typically composed of 10 α-helices (10-TM) arranged in two
bundles of five helices each (12). These bundles are related by a pseudo-twofold symmetry axis
running parallel to the membrane. Architecturally, they are part of the Drug/ Metabolite Exporter
(DME) structural superfamily (27) and operate using an alternating-access mechanism to
facilitate solute exchange across the membrane. This process involves two gates, formed by
specific helical pairs, that alternately open and close on the opposite sides of the DME fold,
enabling the central solute-binding site to access solutes from either side of the membrane. To
date, the SLC35 family comprises 32 members classified into seven subfamilies (SLC35A-G)
based on sequence similarity and phylogenetic analyses, with members within a subfamily
differentiated by the solutes they transport. For the majority of SLC35 proteins (21 of the 32
members), the substrates are unknown (orphan transporters) [27], while most of the functionally
characterized members (mostly in subfamilies A-D) are Nucleotide Sugar Transporters (NSTs)
located in the Golgi and endoplasmic reticulum membranes, where they transport specific
nucleotide sugars into the lumens of these organelles to serve as substrates for glycosylation
reactions (28–31). NSTs use an antiport mechanism and typically utilize the corresponding
luminal nucleoside monophosphates as their antiport molecules (32). Additionally, members of
the SLC35B subfamily include an ATP/ADP exchanger (33) and 3′-phosphoadenosine 5′-
phosphosulfate (PAPS) transporters (34). In the SLC35F subfamily, which comprises six
members, none have been reported to function as an NST, and only one has been functionally
characterized and found to transport thiamine across the plasma membrane (SLC35F3, (35)).
We built a structure-based multisequence alignment of human, *S. pombe* and *T. brucei*
SLC35F2/Qtp1 proteins based on the superposition of tertiary structures predicted by the AI-
based tool AlphaFold (36) (**Fig. S7**). Overall, the three proteins exhibit 14% sequence identity and
31% similarity. Their predicted structures share the common 10-TM transporter core and possess

different, putatively extra-membranous, N- and C-terminal regions that could not be modeled with confidence. The 10-TM core is most similar to the crystal structure of the CMP-sialic acid transporter SLC35A1 (12, 37) (e.g., mouse CMP-sialic acid transporter, PDB ID 6OH3, r.m.s.d. 3.3 Å over 247 Cα atoms, 13% sequence identity (38)). (e.g., mouse CMP-sialic acid transporter, PDB ID 6OH3, r.m.s.d. 3.3 Å over 247 Cα atoms, 13% sequence identity (38)) In terms of their topology and orientation in the membrane, SLC35 proteins are classified as type III integral membrane proteins, i.e. their N- and C-termini face the cytoplasm (12). Notably, the predicted structures of human, *S. pombe* and *T. brucei* SLC35F2/Qtp1 proteins exhibit a biased distribution of positively charged amino acids (**Fig. S7B**), consistent with the "positive-inside rule" for integral membrane proteins, whereby basic amino acids cluster at the cytoplasmic interface (39). Analysis of the topology of the core membrane-spanning domain using the online tool Deep TMHMM (40) suggests that the N- and C-termini are located on the cytoplasmic side (**Fig. S7A**). These observations are consistent with the orientation of a type III integral membrane protein (**Fig. S7A**).

Discussion

SLC35F2 belongs to the SLC35 solute superfamily and is a part of the F subfamily, which mostly consists of orphan transporters whose substrates and physiological function(s) are yet to be determined (30). Herein, we demonstrate by homozygous disruption in human HeLa cells that SLC35F2 functions as a key transporter for queuine and queuosine. We show that it is the sole high-affinity plasma membrane transporter for the Q nucleoside and the primary high-affinity transporter for the nucleobase q in human cells. Furthermore, it is the exclusive transporter of both q and Q in *T. brucei* and *S. pombe* and thus has been renamed Qtp1 in those two species.

Previous literature has shown that q uptake can occur by two different transporters in human cells, a low K_m (30 nM) component demonstrating rapid saturation and a high K_m component (350 nM) showing a slower linear uptake, but their identity was not determined (5). Here we demonstrate that SLC35F2 accounts for high affinity K_m q transporter activity (67 nM), and that there exists an independent lower affinity transporter for q (259 nM) that is evident from competitive inhibition of SLC35F2 with saturating concentrations of the anti-cancer compound YM155. Notably, the K_m for q is higher than the maximum levels determined in human serum under normal conditions of ~25 nM (4, 25) and could therefore indicate that the levels of q in serum are capable of reaching higher levels, such as after a meal replete with q/Q. In the case of Q, its levels have not been detectable in human plasma (4). Therefore, the observed K_m of 174 nM could suggest SLC35F2's role in Q uptake is limited to environments that are rich in this micronutrient, such as the human gut, where Q is liberated from tRNA turnover either from the microbiome or from food. In this regard, an analysis of tissue-specific expression data ((41) and Table S4) finds that there is a heavy propensity for SLC35F2 to be expressed in the human alimentary canal, suggesting that SLC35F2 is responsible for the active transport of Q/q during the digestive process. In our previous work, we observed that the liver has distinctly higher levels of the Q salvage enzyme QNG1 (18). It could therefore be envisaged that Q is transported from the gut by SLC35F2 and delivered to the liver for cleavage by QNG1, which then serves to liberate the q base for subsequent release into the serum and wider distribution throughout the body. Whether the SLC35F2 transporter also functions in a reversible manner to release q to the extracellular space is an outstanding question.

Similar to other eukaryotes, the blood borne parasite *T. brucei* is auxotrophic for q/Q, and it is interesting that it exploits a homologous transporter for uptake (termed Qtp1 in this work). In this regard, tRNA from blood-stream trypanosomes shows a greater level of Q-modification relative to the procyclic form that resides in the midgut of the tsetse fly (22), a difference that could potentially be explained by differential expression of the Qtp1 transporter.

The SLC35F2 gene was first identified during the construction of a transcription map of the ATM gene locus located on chromosome region 11q23, which is associated with the genetic

disorder ataxia telangiectasia (42). Subsequently, an analysis of SLC35F2 mRNA expression by
RT-PCR revealed that it is highest in human adult salivary glands (41). Previous studies on
SLC35F2 have linked its expression to the progression of several forms of cancer (43). For
example, high levels of SLC35F2 expression have been identified in non-small cell lung
carcinoma tissue (under the name LSCC-3), suggesting its use as a prognostic biomarker (44).
Moreover, elevated levels of SLC35F2 have been associated with papillary thyroid carcinoma
(PTC) and inhibiting SLC35F2 expression was shown to suppress the malignant phenotype of
PTC (45). Also, SLC35F2 shows oncogenic behavior, promoting cell proliferation and metastasis
in H1299 lung cancer cells (46). Furthermore, SLC35F2 is highly expressed in human pluripotent
stem cells, which underlies the high cytotoxicity of YM155 in these cells (47). Given our
identification of SLC35F2 as an import molecule for q and Q, one interpretation of its oncogenic
activity is that its overexpression increases cellular levels of q/Q and thus promotes higher Q34
modification of tRNAs. This in turn may give malignant cells a selective advantage over normal
cells. Elucidating the role of SLC35F2 in q/Q import will deepen our understanding of its
involvement in cancer progression and open new avenues for the development of targeted anti-
cancer therapies.

Materials and Methods

364   The *in silico* methods are given below. The growth conditions, mutant constructions, and
365   analytical methods for Q/q detection and transport assays for HeLa, *T. brucei* and *S. pombe* cells
366   are given in the supplemental methods section.

Bioinformatic Analyses

The workflow for searching for candidates of Q precursor transmembrane transporters comprised
three steps. First, QTRT$^-$ and QTRT$^+$ organisms were selected for comparative genomic analysis
(Fig. S1 Step I). To identify the QTRT$^-$ Eukaryotes, 245,896,766 protein sequences and metadata
were retrieved from the Uniprot database (www.uniprot.org/help/downloads). Homologs of the
human QTRT1 (Uniprot ID Q9BXR0) were searched from the proteomes of 10,768 taxa that
contained at least 5000 protein entries using DIAMOND v2.1.8 (48) with the setting "--very-
sensitive --matrix BLOSUM45 --evalue 0.001 -k0". Four taxa of the Eukaryota superkingdom
(including Sar, Viridiplantae, Fungi and Metazoa) that comprised at least ten QTRT$^-$ organisms
(Table S51) were subjected to further analyses. To ensure that all *Qtrt1* genes were called, *Qtrt1*
coding sequences were retrieved using tBLASTn (49) searching the NCBI nt and wgs databases
(50) limited to genomes from these four taxa. Organisms without hits at an E-value of 0.001 and
20% minimum identity were retained as QTRT$^-$. Organisms encoding QTRT1 proteins between
300-500 amino acid in length were identified as QTRT$^+$. These organisms were then sorted in
descending order based on the number of proteins in any given taxa with at least four
transmembrane domains, and this information was used to select approx. 200 organisms in each
rank with criteria in Table S6. The QTRT$^-$ and QTRT$^+$ organisms are listed in Tables S3 and S7-9.
Potential transporter families from both QTRT$^+$ and QTRT- organisms were then clustered based
on sequence similarity (using MMseqs2(51) (ref) with --cov-mode 0; Fig. S1 Step II). Clustering
criteria for minimum coverage and sequence identity were tested in increments from 5 to 50%. A
30% threshold for both criteria was chosen for separation between homologous families as the
number of proteins in each cluster sharply decreased at this cut-off (Table S10). Finally, the
potential transporter subfamilies clusters were given scores as follows: the presence of a family
member in a given QTRT$^+$ organism or its absence in a given QTRT$^-$ organism were scored by
multiplying by 1; while the presence of a family member in a QTRT$^-$ organism or its absence in a
QTRT$^+$ organism were penalized by multiplying by -1 (Fig. S1 Step III). The score of each family
was normalized by the expected maximum score (present in all QTRT$^+$ organisms and absent in

QTRT$^-$ organisms) so it fell within the range of -1 to 1. The scores of the top 200 candidates for
the four ranks are listed in Table S1. The scripts used in the workflow is available at
https://github.com/vdclab/published_scripts.

Sequence Similarity Networks (SSNs)

SSNs were generated using the Enzyme Function Initiative (EFI) analytic suite (52) and
visualized using Cytoscape (3.10.1) (53). 5782 sequences for the SLC35F1/F2 solute transporter
family in Eukaryota were retrieved with the InterPro accession (54) of "IPR052221" through the
"Family" option of EFI-EST (EFI Enzyme Similarity Tool)(52). The initial SSN was generated with
an alignment score threshold (AST) set such that each connection (edge) represented a
sequence identity above 40%. The obtained SSN was first colored by the presence and absence
of QTRT in each organism. More stringent SSNs were then created by gradually increasing the
AST in increments (usually by 10). This process was iterated until clusters containing nodes from
QTRT$^-$ organisms were separated from those comprising nodes from QTRT$^+$ organisms. Clusters
with less than 10 sequences were hidden for better visualization. The UniProt IDs of IPR05221
family sequences in the SSN are listed in Table S11.

Phylogenetic Analysis of SLC35F1/F2 family Proteins

34 IPR05221 family sequences across different clades/kingdoms and three outgroup sequences,
SLC35B1-4/HUT1 were aligned using MUSCLE v5.2 (55) and trimmed using BMGE v1.12 (56)
with matrix BLOSUM30. The maximum likelihood tree was generated using FastTree v2.1.11 (57)
with -lg -cat 20 -gamma model and bootstrap (1000 replicates) and visualized by iTOL v6.9.1 (58)
The QTRT$^-$ organisms are colored in red. The branches were colored by clades or kingdoms, and
those with bootstrap support values higher than 0.7 were indicated by blue dots. The sequences
used in the tree are listed in Table S12.

Sequence and Structure Alignment

422 Structural models of SLC35F2 sequences from human, *S. pombe* strain 972 / ATCC 24843, and
423 *T. brucei* strain 927/4 (Uniprot IDs Q8IXU6, O59785 and Q57UU3, respectively) were generated
424 using AlphaFold v4 (36) and aligned in PROMALS3D (59). The resulting structure-based multi-
425 sequence alignment was visualized using ESPript server v3.0 (60).

Funding

National Institutes of Health [GM132254 to V.d.-L., M.A.S., J.A.; GM146075 to M.A.S], Science
Foundation Ireland [HRB-USIRL-2019–2 to V.P.K.]; Medical Research Council [MC PC 18038 to
B.D.G.]; Health and Social Care in Northern Ireland (HSCNI) [STL/5460/18 to BDG]; Deutsche
Forschungsgemeinschaft (EH237/ 19-1) to A.E.-M., Deutscher Akademischer Austauschdienst
DAAD to B.I.P.

Figures

Figure 1. Transport of q and Q into eukaryotes and subsequent intracellular utilization. The
q/Q micronutrients are imported into the cell via (an) unidentified transporter(s). Q is
phosphorylated by an unknown kinase to Q-5′-monophosphate (Q-5′MP) and subsequently
hydrolyzed by QNG1 to release the q nucleobase. Queuine serves as a substrate of the
eukaryotic tRNA-guanine transglycosylase (TGT) heterodimer (QTRT1 and QTRT2) for
incorporation into tRNA of the GUN anticodon, generating tRNAs modified with Q (Q_{34}-tRNA) at
position 34. The Q in tyrosyl- and aspartyl-tRNA can be modified further by galactosylation and
mannosylation by the enzymes QTMAN and QTGAL, respectively (61). Q-5′MP is the biologically
relevant substrate of QNG1, a queuosine-5′-monophosphate hydrolase (18).

Figure 2. The SLC35F transporter family is functionally diverse. (A) The SSN of SLC35F family proteins in Eukaryota. Each node in the network represents one SLC35F protein. An edge (represented as a line) is drawn between two nodes with a BLAST E-value cutoff of better than 10^{-110} (alignment score threshold of 110). The shape of nodes is based on the presence (circles) and absence (diamonds) of QTRT1 in the organism. The nodes of selected model organisms are highlighted in different colors. The borders of SLC35F1 and F2 annotated in Metazoa are highlighted in orange and pink, respectively. For better visualization, clusters with less than ten sequences are hidden. (B) The sub-SSN of the cluster of phylum Chordata. An edge (represented as a line) is drawn between two nodes with a BLAST E-value cutoff of better than 10-125 (alignment score threshold of 125). (C) The maximum likelihood tree of 34 SLC35F proteins. SLC35B1-4/HUT1 proteins were used as the outgroup. The branches are colored by taxonomic ranks and branches with a 0.7 or more bootstrap value are indicated by blue dots. The QTRT$^+$ and QTRT$^-$ organisms are shown in black and red, respectively.

Figure 3. SLC35F2 is the transmembrane transporter responsible for Q/q uptake in humans, *S. pombe* and *T. brucei*. (A) tRNA[His] is Q-modified in wild-type (WT) HeLa cells, but not in four SLC35F2 knockout candidate clones (1-4), as determined by APB Northern blot. Cells were incubated with Q (25 nM) or q (25 nM) for 6 hrs; **(B)** SLC35F2 knock-out clone 1 was incubated with increasing concentrations of q and Q as indicated for 6 hrs, and the Q-modification status of tRNA[His] was analyzed; **(C)** WT and a *qtp1Δ S. pombe* strain were cultured in the absence (-) or presence of q/Q at the indicated concentrations, tRNA extracted and analyzed by APB Northern using a tRNA[Asp] probe; **(D)** APB Northern analysis of Q-modification status of tRNA[Tyr] in WT and *Δqtp1 T. brucei* strains that were cultured in media supplemented with either complete or dialyzed Fetal Bovine Serum (FBS) supplemented when noted with 25 nM Q or q. tRNA extracted from a strain deleted for the gene encoding TGT2, one paralog of TGT in *T. brucei* (21).

Figure 4. Analysis of q/Q uptake kinetics and specificity of SLC35F2 in human cells. (A)
Uptake of Q was examined by adding increasing concentrations of Q (25, 50, 100, 250 nM) over
four timepoints (15, 30, 45 and 60 mins) to QNG1-knockout HeLa cells, followed by
methanol:water extraction from cell pellets and LC-MS/MS analysis to quantify the intracellular Q
and Q-5′MP concentrations. (**B-C**) Similar analysis was performed for q in *Qtrt1*-knockout HeLa
cells that were untreated (B) or administered the known SLC35F2 inhibitor YM155 (C) at
saturating concentrations, before methanol:water extraction and LC-MS/MS analysis. **(D-E)**
Import of q and Q by SLC35F2 was inhibited by YM155, but not other nucleobases or
nucleosides. To wild-type or SLC35F2 knockout HeLa cells, a fixed concentration of Q (25 nM) or
q (25 nM) was added, as indicated, in the absence (-) or presence of saturating concentrations of
nucleobases or nucleosides (25 μM) or YM155 (25 μM). Total RNA was extracted and the Q-
modification status of tRNA[His] analysed by APB Northern blotting. Nucleobases - guanine (G),
adenine (A), cytosine (C), thymine (T), uracil (U), hypoxanthine (H). Nucleosides - guanosine
(Guo), adenosine (Ado), cytidine (Cyd), thymidine (Thy), uridine (Urd), inosine (Ino).

References

495   1.   B. N. Ames, Prolonging healthy aging: Longevity vitamins and proteins. *Proc Natl Acad*
496   *Sci U S A* **115**, 10836–10844 (2018).

497   2.   C. Fergus, D. Barnes, M. A. Alqasem, V. P. Kelly, The queuine micronutrient: Charting a
498   course from microbe to man. *Nutrients* **7**, 2897–2929 (2015).

499   3.   J. R. Katze, Queuosine Metabolism: Possible relation to B-cell activation by C8
500   derivatives of guanosine. *Proc Soc Exp Biol Med* **179**, 492–496 (1985).

501   4.   X. Pan, *et al.*, Development, validation and application of an LC–MS/MS method
502   quantifying free forms of the micronutrients queuine and queuosine in human plasma
503   using a surrogate matrix approach. *Anal Bioanal Chem* **416**, 5711–5719 (2024).

504   5.   M. S. Elliott, R. W. Trewyn, J. R. Katze, Inhibition of Queuine uptake in cultured human
505   fibroblasts by phorbol-12,13-didecanoate1. *Cancer Res* **45**, 1079–1085 (1985).

506   6.   R. C. Morris, B. J. Brooks, K. L. Hart, M. S. Elliott, Modulation of queuine uptake and
507   incorporation into tRNA by protein kinase C and protein phosphatase. *Biochim Biophys*
508   *Acta Mol Cell Res* **1311**, 124-132(1996).

509   7.   M. S. Elliott, D. L. Crane, Protein kinase C modulation of queuine uptake in cultured
510   human fibroblasts. *Biochem Biophys Res Commun* **171**, 393–400 (1990).

511   8.   R. Zallot, *et al.*, Plant, animal, and fungal micronutrient queuosine is salvaged by
512   members of the DUF2419 protein family. *ACS Chem Biol* **9**, 1812–1825 (2014).

513   9.   M. H. Saier, *et al.*, The Transporter Classification Database (TCDB): 2021 update.
514   *Nucleic Acids Res* **49**, D461–D467 (2021).

515   10.   M. Kanehisa, Y. Sato, M. Kawashima, M. Furumichi, M. Tanabe, KEGG as a reference
516   resource for gene and protein annotation. *Nucleic Acids Res* **44**, D457–D462 (2016).

517   11.   A. Orellana, C. Moraga, M. Araya, A. Moreno, Overview of nucleotide sugar transporter
518   gene family functions across multiple species. *J Mol Biol* **428**, 3150-3165(2016).

519   12.   S. Kamiyama, H. Sone, Solute Carrier Family 35 (SLC35)—An overview and recent
520   progress. *Biologics* **4**, 242–279 (2024).

521   13.   R. Fredriksson, K. J. V. Nordström, O. Stephansson, M. G. A. Hägglund, H. B. Schiöth,
522   The solute carrier (SLC) complement of the human genome: Phylogenetic classification
523   reveals four major families. *FEBS Lett* **582**, 3811–3816 (2008).

524   14.   P. R. Bushel, *et al.*, Dissecting the fission yeast regulatory network reveals phase-
525   specific control elements of its cell cycle. *BMC Syst Biol* **3**, 93 (2009).

526   15.   F. Van den Bossche, *et al.*, Residence of the Nucleotide Sugar Transporter Family
527   Members SLC35F1 and SLC35F6 in the Endosomal/Lysosomal Pathway. *Int J Mol Sci*
528   **25**, 6718 (2024).

529   16.   G. L. Igloi, H. Kossel, Affinity electrophoresis for monitoring terminal phosphorylation
530   and the presence of queuosine in RNA. Application of polyacrylamide containing a
531   covalently bound boronic acid. *Nucleic Acids Res* **13**, 6881–6898 (1985).

17.   M. Müller, *et al.*, Queuine links translational control in eukaryotes to a micronutrient from
bacteria. *Nucleic Acids Res* **47**, 3711–3727 (2019).

18.   S. H. Hung, *et al.*, Structural basis of Qng1-mediated salvage of the micronutrient
queuine from queuosine-5′-monophosphate as the biological substrate. *Nucleic Acids*
Res **51**, 935–951 (2023).

19.   B. I. Patel, M. Heiss, A. Samel-Pommerencke, T. Carell, A. E. Ehrenhofer-Murray,
Queuosine salvage in fission yeast by Qng1-mediated hydrolysis to queuine. *Biochem*
Biophys Res Commun **624**, 146–150 (2022).

20.   A. E. Ehrenhofer-Murray, Cross-Talk between Dnmt2-dependent tRNA methylation and
queuosine modification. *Biomolecules* **7**, 14 (2017).

21.   S. Kulkarni, *et al.*, Preferential import of queuosine-modified tRNAs into *Trypanosoma*
brucei mitochondrion is critical for organellar protein synthesis. *Nucleic Acids Res* **49**,
8247–8260 (2021).

22.   S. Dixit, *et al.*, Dynamic queuosine changes in tRNA couple nutrient levels to codon
choice in *Trypanosoma brucei*. *Nucleic Acids Res* **49**, 12986–12999 (2021).

23.   A. C. Kessler, *et al.*, Retrograde nuclear transport from the cytoplasm is required for
tRNATyr maturation in *T. brucei*. *RNA Biol* **15**, 528–536 (2018).

24.   K. M. Rutherford, M. Lera-Ramírez, V. Wood, PomBase: a Global Core Biodata
Resource—growth, collaboration, and sustainability. *Genetics* **227**, iyae007 (2024).

25.   P. Richard, *et al.*, Queuine, a bacterial-derived hypermodified nucleobase, shows
protection in in vitro models of neurodegeneration. *PLoS One* **16**, e0253216 (2021).

26.   G. E. Winter, *et al.*, The solute carrier SLC35F2 enables YM155-mediated DNA damage
toxicity. *Nat Chem Biol* **10**, 768–773 (2014).

27.   P. M. Berninsone, C. B. Hirschberg, Nucleotide sugar transporters of the Golgi
apparatus. *Curr Opin Struct Biol* **10**, 542–547 (2000).

28.   K. T. Schjoldager, Y. Narimatsu, H. J. Joshi, H. Clausen, Global view of human protein
glycosylation pathways and functions. *Nat Rev Mol Cell Biol* **21**, 729-749(2020).

29.   Z. Song, Roles of the nucleotide sugar transporters (SLC35 family) in health and
disease. *Mol Aspects Med* **34**, 590-600(2013).

30.   B. Hadley, *et al.*, Nucleotide Sugar Transporter SLC35 family structure and function.
Comput Struct Biotechnol J **17**, 1123-1134(2019).

31.   S. Mikkola, Nucleotide Sugars in Chemistry and Biology. *Molecules* **25**, 5755 (2020).

32.   D. L. Jack, N. M. Yang, M. H. Saier, The drug/metabolite transporter superfamily. *Eur J*
Biochem **268**, 3620–3639 (2001).

33.   M.-C. Klein, *et al.*, AXER is an ATP/ADP exchanger in the membrane of the
endoplasmic reticulum. *Nat Commun* **9**, 3489 (2018).

34.   C. B. Hirschberg, P. W. Robbins, C. Abeijon, Transporters of nucleotide sugars, ATP,
and nucleotide sulfate in the endoplasmic reticulum and Golgi apparatus. *Annu Rev*
Biochem **67**, 49–69 (1998).

35. K. Zhang, *et al.*, Genetic implication of a novel thiamine transporter in human
hypertension. *J Am Coll Cardiol* **63**, 1542–1555 (2014).

36. J. Jumper, *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature*
596, 583–589 (2021).

37. E. Nji, A. Gulati, A. A. Qureshi, M. Coincon, D. Drew, Structural basis for the delivery of
activated sialic acid into Golgi for sialylation. *Nat Struct Mol Biol* **26**, 415–423 (2019).

38. S. Ahuja, M. R. Whorton, Structural basis for mammalian nucleotide sugar transport.
Elife **8,** e45221 (2019).

39. C. Baeza-Delgado, M. A. Marti-Renom, I. Mingarro, Structure-based statistical analysis
of transmembrane helices. *Eur Biophys J* **42**, 199–207 (2013).

40. J. Hallgren, *et al.*, DeepTMHMM predicts alpha and beta transmembrane proteins using
deep neural networks. *bioRxiv* 2022.04.08.487609 (2022).

41. M. Nishimura, S. Suzuki, T. Satoh, S. Naito, Tissue-Specific mRNA Expression Profiles
of Human Solute Carrier 35 Transporters. *Drug Metab Pharmacokinet* **24**, 91–99 (2009).

42. T. Stankovic, *et al.*, Construction of a transcription map around the gene for ataxia
telangiectasia: identification of at least four novel genes. *Genomics* **40**, 267–276 (1997).

43. R. Kotolloshi, M. Hölzer, M. Gajda, M. O. Grimm, D. Steinbach, SLC35F2, a transporter
sporadically mutated in the untranslated region, promotes growth, migration, and
invasion of bladder cancer cells. *Cells* **10**, 1–12 (2021).

44. L. Bu, G. Jiang, F. Yang, J. Liu, J. Wang, Highly expressed SLC35F2 in non-small cell
lung cancer is associated with pathological staging. *Mol Med Rep* **4**, 1289–1293 (2011).

45. J. He, *et al.*, Solute carrier family 35 member F2 is indispensable for papillary thyroid
carcinoma progression through activation of transforming growth factor-β type I
receptor/apoptosis signal-regulating kinase 1/mitogen-activated protein kinase signaling
axis. *Cancer Sci* **109**, 642–655 (2018).

46. X. Li, *et al.*, Influence on the behavior of lung cancer H1299 cells by silencing SLC35F2
expression. *Cancer Cell Int* **13**, 1–6 (2013).

47. K.-T. Kim, *et al.*, Safe scarless cassette-free selection of genome-edited human
pluripotent stem cells using temporary drug resistance. *Biomaterials* **262**, 120295
(2020).

48. B. Buchfink, K. Reuter, H. G. Drost, Sensitive protein alignments at tree-of-life scale
using DIAMOND. *Nat Methods* **18**, 366–368 (2021).

49. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment
search tool. *J Mol Biol* **215**, 403–410 (1990).

50. E. W. Sayers, *et al.*, Database resources of the national center for biotechnology
information. *Nucleic Acids Res* **50**, D20–D26 (2022).

51. M. Steinegger, J. Söding, MMseqs2 enables sensitive protein sequence searching for
the analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028(2017).

52.   R. Zallot, N. Oberg, J. A. Gerlt, The EFI web resource for genomic enzymology tools:
leveraging protein, genome, and metagenome databases to discover novel enzymes
and metabolic pathways. *Biochemistry* **58**, 4169–4182 (2019).

53.   P. Shannon, *et al.*, Cytoscape: A software environment for integrated models of
biomolecular interaction networks. *Genome Res* **13**, 2498–2504 (2003).

54.   T. Paysan-Lafosse, *et al.*, InterPro in 2022. *Nucleic Acids Res* **51**, D418–D427 (2023).

55.   R. C. Edgar, Muscle5: High-accuracy alignment ensembles enable unbiased
assessments of sequence homology and phylogeny. *Nature Commun* **13**, 1–9 (2022).

56.   A. Criscuolo, S. Gribaldo, BMGE (Block Mapping and Gathering with Entropy): A new
software for selection of phylogenetic informative regions from multiple sequence
alignments. *BMC Evol Biol* **10**, 1–21 (2010).

57.   M. N. Price, P. S. Dehal, A. P. Arkin, FastTree 2 – Approximately maximum-likelihood
trees for large alignments. *PLoS One* **5**, e9490 (2010).

58.   I. Letunic, P. Bork, Interactive Tree of Life (iTOL) v6: Recent updates to the
phylogenetic tree display and annotation tool. *Nucleic Acids Res* **52**, W78–W82 (2024).

59.   J. Pei, M. Tang, N. V. Grishin, PROMALS3D web server for accurate multiple protein
sequence and structure alignments. *Nucleic Acids Res* **36**, W30–W34 (2008).

60.   X. Robert, P. Gouet, Deciphering key features in protein structures with the new
ENDscript server. *Nucleic Acids Res* **42**, W320–W324 (2014).

61.   X. Zhao, *et al.*, Glycosylated queuosines in tRNAs optimize translational rate and post-
embryonic growth. *Cell* **186**, 5517-5535.e24 (2023).