The mammalian SKI complex is a broad-spectrum antiviral drug target that upregulates cellular cholesterol to inhibit viral replication

Short Title: Drug targeting the SKI complex increases cellular cholesterol

Stuart Weston[a], Lauren Baracco[a], Louis Taylor[a], Alison Scott[b,c], Gaurav Kumar[d], Paul Shapiro[d], Alexander D. MacKerell Jr. [d], and Matthew B. Frieman[a*]

[a] Center for Pathogen Research, Department of Microbiology and Immunology, University of Maryland School of Medicine, Baltimore, MD, 21201

[b] Department of Microbial Pathogenesis, University of Maryland School of Dentistry, Baltimore, MD, 21201

[c] Department of Microbiology and Immunology, University of Maryland School of Dentistry, Baltimore, MD, 21201

[d] Department of Pharmaceutical Sciences, University of Maryland School of Pharmacy, Baltimore, MD, 20201

* Corresponding author: MFrieman@som.umaryland.edu

Abstract

There is a need for the development of broad-spectrum antiviral compounds that can act as first line therapeutic countermeasures to emerging viral infections. Host-directed approaches present a promising avenue of development and carry the benefit of mitigating risks of viral escape mutants. We have previously found the SKI (super killer) complex to be a broad-spectrum, host-target with our lead compound ("UMB18") showing activity against influenza, coronaviruses, and filoviruses. The SKI complex is a cytosolic RNA helicase and we previously found that targeting it with UMB18 inhibited viral RNA production but did not further define the mechanism. Here, transcriptomic analysis of UMB18 treated A549 cells revealed an upregulation of genes in the mevalonate pathway which drives cholesterol synthesis. Further investigation validated the genetic upregulation and confirmed an increase in total cellular cholesterol. This upregulation was dependent on the SKI complex, the sterol regulatory element binding proteins (SREBPs) and their regulator SCAP, the major regulators for cholesterol and fatty acid synthesis. Depletion of the SREBPs or SCAP with siRNA, or extraction of cholesterol with methyl β-cyclodextrin attenuated UMB18 antiviral activity, emphasizing the role of increased cholesterol synthesis in this mechanism of action. Our findings further define the antiviral mechanism of a developmental host-directed therapeutic approach with broad applicability against emerging viral pathogens.

Author Summary

The COVID-19 pandemic has underscored the urgent need for effective countermeasures to novel and emerging viral pathogens. Our research presented here builds upon our previously published data on an experimental novel antiviral compound termed UMB18. We have found this compound capable of inhibiting replication of influenza A virus, coronaviruses and the filoviruses Marburg and Ebola virus, but did not fully define a mechanism of action. In this work, we demonstrate that UMB18 exerts antiviral activity by modulating cellular cholesterol levels. By targeting the SKI complex, UMB18 triggers an increase in endogenous cellular cholesterol which disrupts the fine balance viruses rely on for efficient infection. We demonstrate that this mechanism inhibits replication of SARS-CoV-2, revealing a previously undescribed host-directed strategy for antiviral intervention. These findings highlight UMB18's potential as a broad-spectrum antiviral agent and pave the way for further research into its mechanism and therapeutic applications, offering a promising avenue for development of antiviral countermeasures to current, novel and emerging pathogens.

Introduction

The COVID-19 pandemic has highlighted the lack of broadly acting therapeutics to treat coronaviruses as well as other emerging pathogens. When severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) emerged and spread globally there were few countermeasures available for treating coronavirus disease-2019 (COVID-19) resulting in a reliance on non-pharmacological interventions of physical distancing and isolation to control the outbreak. While those measures helped to some degree, the continued pandemic spread of the virus highlighted the urgent need for novel therapeutics against emerging viral disease. Fortunately, vaccines were developed at a record pace and contributed significantly to the management of COVID-19. However, the ongoing threat of emerging viral pathogens necessitates the development of complementary strategies. Broad-spectrum antiviral chemical compounds could serve as rapid, first-line countermeasures against emerging pathogens, filling a crucial gap before specific measures can be developed. Addressing this need, our research is focused on a host-directed approach of targeting the SKI (super killer) complex[1].

While using host-directed therapeutics may come at a cost of having certain side effects, there is a significantly lower risk of viral escape mutants developing owing to the varying mutation rate of host versus virus. The risk of virus evolution to escape countermeasures is perfectly demonstrated by the dwindling number of effective monoclonal antibodies available to treat COVID-19 patients through mutation of the spike protein in SARS-CoV-2 variants[2]. Direct acting nucleoside analogues have been implicated in increasing the risk of viral mutagenesis posing concerns for vaccine and

monoclonal antibody treatments(3). Targeting host factors that do not directly affect the

viral genome could alleviates these concerns.

We focus on the SKI complex, a eukaryotic protein complex we previously identified

through a yeast-based screening system to be a proviral factor(1). Comprising SKIC2,

SKIC3, and SKIC8 (formerly SKIV2L, TTC37, and WDR61, respectively) the SKI

complex acts as a cofactor to the cytosolic RNA exosome to facilitate 3' to 5' RNA

degradation(4–6). SKIC2 is a member of the DExD/H helicase family, while SKIC3 and

SKIC8 have structural and regulatory functions(7). The complex was first defined in

yeast where it was implicated in resistance to the "super killer" toxin of a yeast double

stranded RNA virus, hence the naming convention of SKI(8,9). Our previous study

demonstrated that siRNA depletion of the SKI complex inhibited replication of influenza

A virus (IAV) and Middle East respiratory syndrome coronavirus (MERS-CoV). We

proceeded to investigate chemical compounds that were *in silico* predicted to bind to

SKIC8 and potentially disrupt the proviral activity of the SKI complex. This work

generated multiple chemical scaffold structures that inhibited both IAV and MERS-CoV,

in agreement with our siRNA studies, but additionally inhibited SARS-CoV, SARS-CoV-

2, and the filoviruses Ebola and Marburg. We demonstrated that our lead compound,

termed UMB18, did not significantly inhibit IAV entry to cells, but potently inhibited RNA

and subsequent protein production in single-cycle infections and therefore imposed

antiviral activity. Our previous work defined UMB18 as a host-directed antiviral, targeting

post-entry steps prior to RNA production, with broad-spectrum activity(1).

This study aimed to elucidate the mechanism of the antiviral activity of UMB18. Using

RNA sequencing of cells treated with UMB18, with and without SARS-CoV-2 infection,

we identified a significant upregulation of genes in the mevalonate pathway, which is

crucial for cholesterol synthesis, regardless of viral infection status. Cholesterol is an

essential lipid found throughout cellular membranes which at high levels causes an

increased rigidity, it is also a major part of hormone and oxysterol production.

Cholesterol can be obtained from external sources through low-density lipoprotein (LDL)

receptors or endogenously synthesized through the mevalonate pathway(10–12). This

cholesterol synthesis pathway has many other useful metabolic products for the cell and

is involved with protein prenylation, such that there must be tight regulation between

uptake from external sources and endogenous synthesis of cholesterol. Activation of

this pathway by UMB18 suggests a previously unknown antiviral mechanism of action

for our developmental compound.

Sterol regulatory element binding proteins (SREBPs) are master regulators of

cholesterol and fatty acid synthesis through upregulating genes involved in both

synthesis pathways(13). SREBPs comprise three isoforms, SREBP1a, SREBP1c, and

SREBP2, expressed from two genes *Srebf1* and *Srebf2*(14). Each isoform has an

equivalent tripartite structure with an N-terminal basic-helix-loop-helix leucine zipper

family transcriptional domain, a short hairpin transmembrane domain, and a C-terminal

domain involved in protein-protein interactions(13,14). SREBP1a and SREBP1c are

produced from alternate transcriptional start sites which alters their N-terminal domains

and their sterol regulator element (SRE) binding specificity(15). SREBP2 is roughly 50%

identical to SREBP1 at the amino acid level. It is generally considered that SREBP1c is

mostly involved in expression of lipogenic genes, SREPB1a can regulate both lipid and

cholesterol synthesis genes, while SREBP2 is mostly involved in cholesterol

synthesis(16,17).

Srebf genes produce inactive precursor proteins that localize to the endoplasmic

reticulum (ER) where they associate to SREBP cleavage-activating protein (SCAP).

Cholesterol binding to SCAP causes a conformation that results in retention of the

SCAP-SREBP complex at the ER. However, low cholesterol levels cause a

conformational change to SCAP, exposing a MELADL motif that directs COPII-coated

vesicle trafficking of the SCAP-SREBP complex to the Golgi apparatus(18,19). At the

Golgi, SREBP proteins are cleaved by signal peptidase 1 and 2 to free the N-terminal

transcription factor domain for nuclear translocation(13,20). Cleavage-activated SREBP

binds SREs and initiates transcription of genes in both fatty acid and cholesterol

synthesis pathways.

In the present study, transcriptomic analysis revealed that our SKI-targeting compound

UMB18 caused an upregulation of genes involved in cholesterol synthesis, a finding we

confirmed with qRT-PCR. In agreement with the RNA-based increase in cholesterol

synthesis genes we also saw an increase in total cellular cholesterol using mass

spectrometry. UMB18 was found to bind SKIC8 through *in vitro* expression and binding

assays and the transcriptional changes were dependent on the presence of SKI

complex since these were reduced by siRNA-mediated depletion of SKI genes, where

SKIC8 depletion had the greatest effect. Further, depletion of *Srebf1* or *Srebf2* reduced

UMB18-mediated increases in mevalonate pathway genes, indicating that UMB18 acts

through the canonical pathway for endogenous cholesterol synthesis. Notably, UMB18

antiviral activity against SARS-CoV-2 was significantly reduced with either depletion of

the *Scap* or *Srebf* genes or extraction of cholesterol with methyl-β cyclodextrin. These

results underscore the importance of increased total cellular cholesterol in the

mechanism of action for UMB18 as an antiviral compound.

Overall, the work presented here builds upon our previous study demonstrating the

promise of UMB18 as a broad-spectrum, host-directed antiviral compound and

elucidates its mechanism of action through upregulation of cellular cholesterol synthesis

by targeting the SKI complex.

Results

Transcriptomic profiles show UMB18 upregulates genes involved in cholesterol

synthesis.

Having previously described the broad-spectrum antiviral activity of the SKI-targeting

compound UMB18(1), we aimed to better identify the mechanism of action for the

compound. To this end we utilized RNA transcriptomic analysis of treated and/or SARS-

CoV-2 infected cells *in vitro*. A549 cells (human lung alveolar adenocarcinoma cells)

overexpressing human angiotensin converting enzyme 2 (A549-ACE2) to allow for

SARS-CoV-2 infection were used for the transcriptomic analysis and four conditions

were analyzed: 10μM UMB18 treatment, 0.1% DMSO treatment (carrier for UMB18),

10µM UMB18 with SARS-CoV-2 infection or 0.1% DMSO with SARS-CoV-2 infection.

We previously found 10µM to be inhibitory to viral replication with no cytotoxicity(1). All

infections were at multiplicity of infection (MOI) 0.1 and across two experiments with

either 24-hour (h) and 48 h or just a 24 h time point were analyzed. In uninfected cells,

UMB18 treatment resulted in increased expression of mevalonate pathway genes at 24

h and 48 h compared to uninfected, DMSO-treated cells (Fig. 1A). Pathway analysis

revealed an enrichment in pathways associated with cholesterol biosynthesis, which

were predicted to have increased activity based on the patterns of differential

expression of genes within the pathway (Fig. 1B). Mevalonate pathway genes are

upregulated with UMB18 treatment at both 24 h and 48 h relative to DMSO-treated

controls, regardless of infection with SARS-CoV-2 (Fig. 1C). SARS-CoV-2 infection had

minimal impact on the mevalonate pathway, probably because we used bulk RNA

sequencing and 10µM UMB18 caused a roughly 90% inhibition of infection meaning

only low levels of SARS-CoV-2 replication are occurring in the treated cells (for example

see Fig. 5F). This analysis demonstrated upregulation of genes involved in the

mevalonate pathway after treatment with UMB18, potentially linking the SKI complex

with cholesterol synthesis.

Increase in cholesterol levels in UMB18 treated cells.

The upregulation of the mevalonate pathway through the transcriptomic analysis

suggested that cholesterol synthesis may be affected by UMB18. We validated the

results of the transcriptomic analysis using qRT-PCR with primers targeting a subset of

the genes in the mevalonate pathway. Identical to the previous experiment, A549-ACE2

cells were treated with 10µM UMB18 or 0.1% DMSO as the vehicle control for 24 or

48h. Additionally, A549 cells not expressing ACE2 were subject to the same treatments

to further validate the results and ensure that ACE2 overexpression is not impacting the

transcriptional changes. Seven genes for enzymes involved in various stages of the

mevalonate pathway were analyzed to get a broad view of transcriptional changes

(pathway schematic in Fig. 2A, yellow boxes are selected genes for analysis). Relative

to DMSO treatment, UMB18 caused an increase in expression of genes in the

mevalonate pathway, ranging from approximately 2- to 6-fold. In the A549-ACE2 cells

(Fig. 2B), the induction of the selected genes peaked at 24h and then declined by 48h,

yet remain above the levels observed in DMSO-treated cells, while in the A549 cells

(Fig. 2C) there was an increase or at least stability of the change observed across the

full 48 h. Even though there was a minor difference in the temporal dynamics, the

magnitude of change was largely equivalent between the two cell lines, and we suspect

these differences are artifacts of using cell culture systems with the base A549 cells

having two different backgrounds (see Materials and Methods), rather than an impact of

ACE2 expression. Overall, these qRT-PCR data suggest that UMB18 caused an

increase in genes that regulate cholesterol synthesis through the mevalonate pathway

and that the expression of ACE2, which allows for infection by SARS-CoV-2 (which will

be used later in this study), does not significantly alter the magnitude of change.

These experiments were performed in serum supplemented media and therefore have

exogenous cholesterol available. Cells can endocytose cholesterol through the LDL-

receptor and activate a negative feedback loop to inhibit the mevalonate pathway. It

therefore appears UMB18 may activate this metabolic pathway even in the presence of

exogenous cholesterol. To further analyze the system, we treated A549-ACE2 cells with

UMB18 or DMSO in serum-containing or serum-free media, harvested at 24 hours, and

performed qRT-PCR for a subset of the mevalonate pathway genes. In serum-

containing media, UMB18 treatment resulted in a similar increased gene expression.

However, no significant differences were noted in serum-free media (Fig. S1A). In the

absence of exogenous cholesterol, cells likely activate the mevalonate pathway to

promote endogenous synthesis. Analysis of DMSO treated cells demonstrated a 6-7-

fold increase in expression of three mevalonate pathway genes in serum-free compared

to serum-containing media (Fig. S1B). We hypothesize that the lack of UMB18 effect in

serum-free media (Fig. S1A) is due to the pathway already being activated in response

to cholesterol-deprivation resulting in UMB18 having minimal additional impact on this

pathway. Supporting this hypothesis, UMB18 treatment in serum-free media showed

only a 2-fold increase in gene expression compared to serum-containing media, in

contrast to the 6-7-fold increase with DMSO (Fig. S1B). Consequently, since UMB18

and serum depravation both increase mevalonate pathway gene expression, their

combination results in minimal additive effects on the mevalonate pathway gene

expression. These results, which have implications for viral replication (Fig. S1C) to be

discussed later, demonstrate that UMB18 treatment induces expression of mevalonate

pathway genes in a manner comparable to the absence of serum.

With there being an increase in the expression of genes involved in the mevalonate

pathway, we next investigated whether total cellular cholesterol was increased in the

cells. To this end, we utilized silver-assisted laser desorption ionization mass

spectrometry imaging (Ag-LDI MSI), a highly sensitive method of detecting total cellular

cholesterol, of A549-ACE2 cells treated with UMB18 or DMSO. Cell culture monolayers

treated with UMB18 showed a higher intensity of cholesterol signal compared to DMSO-

treated cells. Cholesterol was detected and visualized in the monolayers as the ion *m/z*

493.260 +/- 2.2 mDa ([cholesterol+Ag$_{107}$]$^+$) ion (Fig. 2D) and the silver isotopologue ion

[cholesterol+Ag$_{109}$]$^+$ data were in agreement. Treatment with UMB18 resulted in a

significant increase in relative cholesterol abundance, showing a nearly 2-fold increase

in pixel intensities. Further, a receiver-operator characteristic analysis showed an area-

under-the-curve value of 0.804 for cholesterol in UMB18 treated cells, making

cholesterol abundance a strong discriminator over DMSO-treated cells. These data

demonstrate that UMB18 treatment of A549-ACE2 cells results in significantly and

discriminatorily higher amounts of cholesterol compared to control.

UMB18 directly binds SKIC8 and the SKI complex is required for increasing

mevalonate pathway genes.

Since UMB18 treatment caused an increase in cellular cholesterol through

transcriptional regulation of genes involved in the mevalonate pathway, we wanted to

ensure that this response was through specificity of UMB18 targeting the SKI complex.

We previously developed the compound using an *in silico* modeling approach, with the

compound predicted to bind to SKIC8 (formerly WDR61), and showed that the genes of

the SKI complex were required in cells for the antiviral activity, thus suggesting

specificity(1). However, whether the increase in cholesterol is a consequence of

targeting the SKI complex needs to be established.

In our previous work, UMB18 was modeled to target SKIC8 at the binding interface with

SKIC3 (formerly TTC37). We wanted to establish that the predicted binding is seen *in*

vitro and to that end, expressed and purified SKIC8 from bacterial cells and analyzed

UMB18 binding by surface plasmon resonance (SPR). The fast association and

dissociation rate constants observed at various compound concentrations indicated

binding to the SKIC8 target with a relatively low binding affinity, $K_D = 81$ µM (Fig. 3A).

We next tested whether the SKI complex was important for the observed increase in

mevalonate pathway genes. A549-ACE2 cells were transfected with siRNA sequences

targeting each of the three SKI genes, SKIC2 (formerly SKIV2L), SKIC3, and SKIC8, for

individual knockdown and then treated with UMB18 or DMSO before analysis of

mevalonate pathway genes by qRT-PCR. Cells were siRNA transfected for 72 h and

treated with UMB18 or DMSO for 24 h before collection for qRT-PCR with three select

target genes (MVD, FPDS and MSMO1), which target different parts of the pathway as

displayed in Fig. 2A. The SKI siRNA knockdowns were stable throughout the 96 h of

experiment time and were gene-specific (Fig. 3B). Moreover, UMB18 treatment of cells

did not impact the knockdown nor expression of the SKI genes as compared to DMSO

(Fig. 3B), with no significant different seen between UMB18 and DMSO treated siRNA

transfected cells. In each of the paired UMB18 or DMSO knockdowns, a relative level of

expression of 33 and 35% was achieved for SKIC2 (UMB18 and DMSO respectively),

20 and 20% for SKIC3 and 21% and 20% for SKIC8, all relative to scrambled siRNA

with DMSO.

Transfection of scrambled siRNA did not alter the ability of UMB18 to increase the level

of transcript for the mevalonate pathway genes. There were statistically significant

differences in the ΔCt values between UMB18 and DMSO treated conditions for all

three genes (Fig. 3C-E). For analyzing the effect of SKI siRNA transfection on changes

in mevalonate genes, the ΔΔCT was calculated and plotted as relative fold change to

scrambled siRNA (Scr) transfected cells treated with DMSO treatment (Scr/DMSO) as

the control for all conditions (Fig. 3C). There was still an increase in the mevalonate

genes in the SKI siRNA cells compare to DMSO. However, the only statistically

significant differences were seen for MVD and MSMO1 in the context of SKIC3 siRNA.

No statistically significant differences were seen in any of the SKIC2 or SKIC8 pairwise

comparisons by two-way ANOVA analysis. This lack of statiscally significant difference

suggests that the loss of the SKI complex removes the change caused by UMB18

treatment, even though there is still an increase in expression. To better analyze these

changes, the fold change relative to Scr/DMSO of Fig. 3C can be set as fold change of

UMB18 in SKI siRNA compared to scrambled siRNA to assess the impact that

knockdown of SKI genes has. In this analysis, if the relative fold change is 1, there

would be no change from the knockdown of SKI gene expression, were it to be greater

than 1, the loss of the SKI genes would be enhancing the effect, and were it be below 1,

this suggests that loss of the SKI genes is attenuating the UMB18-mediated increase in

mevalonate pathway genes. Knockdown of the enzymatic SKIC2 had little impact on the

increase of the mevalonate genes, yet there is a reduction in the UMB18-induced

increase when SKIC3 and SKIC8 are reduced. SKIC8 had the most pronounced effect,

causing roughly a 40% decline in the increase in mevalonate gene expression. Since

UMB18 targets SKIC8 at the interface of SKIC3 and SKCI8 we interpret these data as

evidence that the SKI complex is indeed involved in UMB18's induction of the

mevalonate pathway.

SCAP and SREBP are required for UMB18-induced increases in mevalonate

pathway genes.

Our data suggest that UMB18 binds to the SKI complex and consequently increases

expression of genes in the mevalonate pathway leading to an increase in total cellular

cholesterol. We set out to determine what cellular pathways the SKI complex might be

interacting with to mediate this effect. First, we looked at the mammalian target of

rapamycin (mTOR) complex because of recently published findings that demonstrated

that a loss of SKIC2 activity was connected to an increase in mTORC1 signaling(21).

The mTOR complex is a master regulator of numerous aspects of cellular metabolism

and of interest to our research it can regulate activation of the sterol regulatory element

binding proteins (SREBPs) which promote expression of the mevalonate genes. We

analyzed the ribosomal protein S6 kinase 1 (S6K1) which is phosphorylated of when the

mTOR pathway is activated. Using treatment with epidermal growth factor (EGF), a

known activator of the mTOR pathway, we were able to detect phospho-S6K1 (pS6K1),

demonstrating that our cells show a canonical marker for mTOR signaling (Fig. S2B).

We proceeded to treat cells with UMB18 for various time points, either longer times that

match our other treatments (Fig. S2B) or shorter times (Fig. S2C). However, none of

these experiments showed any significant changes to phospho-S6K1, suggesting that

UMB18 wasn't causing a detectable activation of mTOR, at least based on

phosphorylation of S6K1.

While UMB18 did not showing any clear activation of mTOR, we next investigated the

downstream target that was more directly relevant to our data of increased mevalonate

pathway gene expression by looking at the SREBPs. There are three SREBP proteins

in cells produced from the genes *Srebf1* and *Srebf2* (SREBP-1a, SREBP-1c, and

SREBP2). These proteins are usually maintained as inactive precursors in the ER, but

upon appropriate signaling of a requirement for increased lipogenesis are trafficked to

the Golgi to be cleaved, releasing the active form of the protein that can migrate to the

nucleus and activate genes associated with sterol and fatty acid synthesis. Transport of

the SREBPs from the ER to the Golgi is regulated by the actions of SREBP cleavage

activating protein (SCAP). We validated siRNA sequences to target each of these three

genes, three for *Srebf1* and two for each of *Srebf2* and *Scap* by transfection into A549-

ACE2 cells for 72 h prior to treatment with DMSO for 24 h prior to collection for qRT-

PCR for each of the siRNA targets and the same three mevalonate pathway genes as

used in Fig. 3 (Fig. 4A-C).

As shown in Fig. 4A-C, the siRNAs for each gene caused varying levels of knockdown

for each of the target genes in DMSO conditions, but all were effective (white bars in

each panel). For *SCAP*, siRNA sequence 1 and 2 caused a relative level of expression

of 24% and 6%, respectively, compared to scrambled siRNA with DMSO. *Srebf1* had a

relative expression of 33, 47 and 48% for the three siRNAs, while *Srebf2* had a relative

level of expression at 11% and 29% for siRNA sequence 1 and 2, respectively. The loss

of each of these genes has some impact on the expression of mevalonate genes at

baseline levels since there are minor alterations in MVD, MSMO1, and FDPS

expression levels in DMSO conditions, but no statistically significant changes (Fig. 4A-

C). As described previously, to analyze the role of each gene in the UMB18-mediated

increase in mevalonate pathway genes, the ΔΔCT was calculated and plotted as

relative fold change compared to Scr/DMSO (Fig. 4D-F). UMB18 treatment of siScr cells

resulted in approximately a 2- and 12-fold increase in each of the three mevalonate

pathway genes, similar to data from previous figures, but it does not appear to alter

expression of either of the SREBPs nor SCAP (Fig. 4D-F).

However, when cells were transfected with siRNAs targeting *SCAP*, the fold changes in

expression of *MVD*, *FDPS*, and *MSMO1* from UMB18 treatment were all minimally

changed to Scr/DMSO (Fig. 4D). There are significant differences in the fold change

between the UMB18 treated siScr transfected cells and the si*SCAP* transfected cells,

with siRNA-1 giving significant difference for all three genes and siRNA-2 giving

significant differnces for *FDPS* (Fig. 4D). This suggested that the loss of SCAP

attenuated the impact of UMB18 treatment. When the fold change between UMB18

treated siScr and si*SCAP* was analyzed there was a greater than 50% drop in the

impact from UMB18 treatment (Fig. 4G). SREBP2 is generally considered the main

regulator for cholesterol synthesis genes, but SREBP1 can also regulate this pathway.

Our results suggest that loss of SREBP2 does not have as large of an effect on UMB18-

mediated increases in mevalonate pathway genes, with the fold change in Fig. 4H being

close to 1 when comparing UMB18 in siScr and si*SREBF2*. However, knockdown of

SREBF1 caused attenuation of the UMB18-mediated increases similarly to *SCAP*,

again, with varying diegrees of significance seen (Fig. 4F). Overall, these data point

towards SCAP and SREBP1 being the main regulators of the increase in mevalonate

pathway genes following UMB18 treatment of A549-ACE2 cells, with SREBP2 having a

comparatively more minor role.

Increased cellular cholesterol is required for UMB18-mediated antiviral activity.

We have found that UMB18 caused an increase in mevalonate pathway gene

expression and that the canonical signaling pathway mediated by SCAP and SREBPs is

involved. This transcriptional response increased total cellular cholesterol, leading to the

question of whether the increase in cholesterol is important for antiviral activity. To

investigate this, we again used siRNA to target SCAP and the SREBP genes and

treated cells with UMB18 while also infecting with SARS-CoV-2. A549-ACE2 cells were

plated and transfected for 72 h as in other experiments and then were infected with

SARS-CoV-2 at MOI 0.1 and treated with 10μM UMB18 or 0.1% DMSO for 24 h.

Supernatant was collected, and viral titer determined by plaque assay on VeroE6 cells

expressing transmembrane protease serine 2 (TMPRSS2). With scrambled siRNA there

is approximately a 10-fold drop in viral titer (Fig. 5A-C), similar to the effects we

previously published(1). However, when *Srebf1* was knocked down by siRNA, UMB18

lowered viral titer by 1.23-2-fold, compared to DMSO (Fig. 5C), suggesting SREBP1 is

involved in UMB18-mediated antiviral activity. For *Srebf2* siRNA, one sequence

demonstrated a 2-fold inhibition with UMB18, while the other sequence had less effect

at a 7.5-fold decrease and the SCAP siRNAs were at 4.3- and 2.9-fold decrease in viral

titer (Fig. 5C). In all cases, viral titer inhibition due to UMB18 is attenuated by loss of the

SREBPs and SCAP, with varying degrees of significance. Since the loss of these genes

causes a decrease in mevalonate pathway gene expression induced by UMB18 (Fig. 4),

these data collectively suggest that increased cellular cholesterol plays a role in

UMB18-mediated antiviral activity.

UMB18 appears to be exerting antiviral activity through increasing expression of genes

involved in the mevalonate pathway. In previously described experiments, we found that

culturing cells in serum-free media produced a similar increase in gene expression as

UMB18. We therefore investigated whether viral replication was attenuated in serum-

free media, regardless of UMB18 treatment. A549-ACE2 cells were infected with SARS-

CoV-2 at MOI 0.1 in serum-free or serum-containing media with 10μM UMB18 or 0.1%

DMSO for 24 h prior to collection of supernatant and titration by plaque assay. UMB18

is equally effective at inhibiting viral replication regardless of the presence of serum

(Sup Fig. 1C). In serum-free media with DMSO there is a significant drop in viral

replication compared to serum-containing media suggesting that the increased

expression of mevalonate pathway genes may impact replication (Fig. S1C). The

inhibition is not as pronounced as UMB18 treatment and we speculate that this may be

a temporal effect with UMB18 causing a more rapid increase in expression of the

genes.

We also investigated the effects of methyl-β cyclodextrin (MβCD), a chemical that

extracts cholesterol from cells, to see if this would disrupt UMB18 antiviral activity.

Initially, A549-ACE2 cells were treated with a range of concentrations between 39 and

5000 µg/ml to investigate toxicity. The higher concentrations resulted in cell death based

on automated counting of nuclei in a 96-well plate format (Fig. 5D). At 625 µg/ml and

lower, there were no signs of toxicity. A549-ACE2 cells were then infected with SARS-

CoV-2 at MOI 3 and treated with MβCD at non-cytotoxic concentrations. At 24 hours

post-infection cells were fixed and immunofluorescent labeled for the viral N protein and

high content imaged to determine the percentage infection. Fig. 5E demonstrates that

MβCD has some level of antiviral activity with 625, 312.5, and 156.25 µg/ml causing

roughly 30-40% fewer infected cells compared to DMSO-treated cells. However, at 39

µg/ml, there was no antiviral activity from MβCD alone. We, therefore, combined 39

µg/ml MβCD with 10µM UMB18 to investigate whether extracting cholesterol from cells

treated with UMB18 would reduce the antiviral activity. Fig. 5F demonstrates that

UMB18 is a potent inhibitor infection reducing percentage infection of cells by about

90% compared to DMSO-treated control cells. When combined with MβCD, the

percentage inhibition is approximately 40%, demonstrating that MβCD significantly

reduced the antiviral activity of UMB18. In sum, these data suggest that reducing the

increase in mevalonate pathway genes with siRNA to *Srebf* genes or *SCAP*, or by

treating cells with a cholesterol extracting chemical both significantly reduce the antiviral

activity of UMB18, pointing towards cholesterol playing a role in the antiviral mechanism

of our chemical compound.

Discussion

Building upon our identification of the SKI complex as a potential host-directed antiviral

drug target(1), here we have further investigated the mechanism of action of our lead

compound UMB18. The SKI complex is a cytosolic RNA helicase comprising SKIC2,

SKIC3, and SKIC8 (SKIC2 being the enzymatic core) that is a cofactor for the cytosolic

RNA exosome that drives 3' to 5' RNA degradation(4–6). Our previous study

demonstrated the SKI complex is a proviral factor as depletion of its component parts

reduced replication of influenza A virus (IAV) and Middle East respiratory syndrome

coronavirus (MERS-CoV). Using an *in silico* modeling approach we developed chemical

compounds that had broad spectrum antiviral activity. These chemicals were modeled to

target the SKI complex at the interface between the structural units SKIC3 and SKIC8,

with the binding pocket on SKIC8. Various chemical scaffolds were found to be antiviral,

and that activity was against pathogenic human coronaviruses (SARS-CoV, SARS-CoV-

2, and MERS-CoV), IAV, and the filoviruses Ebola and Marburg viruses. In that study,

we demonstrated UMB18 minimally impacts IAV entry yet potently inhbited RNA and

subsequent protein production to cause antiviral activity. However, beyond this

explanation for inhibiting viral replication, we did not further characterize the mechanism

of antiviral activity. Here, we identify that the antiviral activity of UMB18 is mediated by

an increase in cellular cholesterol.

We initially used RNA transcriptomics that were validated by qRT-PCR to demonstrate

that UMB18 treatment, regardless of SARS-CoV-2 infection status, caused significant

upregulation of genes involved in cholesterol synthesis and the mevalonate pathway

(Fig. 1 and 2). In agreement with the increased gene expression, we found total cellular

cholesterol to be increased (Fig. 2). Interestingly, these results were in the context of

serum supplementation and therefore cholesterol containing media which can inhibit

activation of the mevalonate pathway through a negative feedback loop. We found that

cells cultured in serum-free media activated the mevalonate pathway and addition of

UMB18 had minimal additive effect, suggesting the pathway was not activated further

(Fig. S1A and S1B). We propose that the lack of increase results from the pathway

already being activate and UMB18 cannot further increase signaling, however, we

cannot rule out alternative hypotheses that UMB18 may be modified by components of

serum or may not be effectively taken into cells in the absence of serum, or may have

impacts on roles adjact to the mevalonate pathway such as protein prenylation. We do

suspect either of these may to some extent play a role since virus replication is

attenuated in serum-free media but UMB18 still provides a greater level of inhibition

(Fig. S1C). As this compound is further developed, biochemical experiments to look at

the impact of serum will be performed.

Using surface plasmon resonance we demonstrated that UMB18 is indeed binding to *in*

vitro expressed SKIC8 and if the SKI complex is depleted by siRNA the impact of

UMB18 to increase genes in the mevalonate pathway is attenuated (Fig. 3), suggesting

specificity to the SKI complex. Although we cannot rule out the possibility of UMB18

interacting with other targets, we take the *in vitro* binding data combined with the

diminished effect of UMB18 when SKI genes were depleted to support direct and

specific action on the complex. We note that the loss of effect from UMB18 when each

individual SKI gene was knocked down was small, reaching roughly a 40% decrease

from depletion of SKIC8 as the most effective. We attempted to perform double or triple

knockdown of the genes to improve these results but could not produce reliable data

owing to cytopathic effect in these experiments. However, overall, these results suggest

that UMB18 is specifically targeting the SKI complex and increasing mevalonate gene

expression that causes an elevation of total cellular cholesterol.

To investigate how targeting the SKI complex with UMB18 causes an increase in

cholesterol synthesis we focused on the SREBPs and their regulator SCAP, which are

master regulators of sterol and fatty acid synthesis(13). Using siRNA, we knocked down

each of *Srebf1*, *Srebf2,* and *SCAP* individually and then treated cells with UMB18.

Depletion of any of these three genes resulted in an attenuation of UMB18-induced

upregulation of mevalonate pathway genes, indicating that UMB18's effect is through

this canonical signaling pathway (Fig. 4). Crucially, the antiviral activity of UMB18

against SARS-CoV-2 was markedly decrease when the SREBP or SCAP genes were

depleted from cells, arguing that the increased expression of mevalonate pathway

genes is essential to the mechanism of antiviral activity (Fig 5). Consistently, extraction

of cellular cholesterol using MβCD also attenuated the antiviral activity of UMB18,

reinforcing the importance of increased cellular cholesterol in mediating antiviral activity

(Fig. 5).

Our findings collectively reveal that UMB18 binds the SKI complex and activates

SREBP-mediated gene expression, leading to an increase in cellular cholesterol which

is crucial for antiviral efficacy of the compound. We have shown that UMB18 and similar

SKI-targeting compounds exhibit broad-spectrum activity against enveloped viruses,

including coronaviruses, orthomyxoviruses, and filoviruses(1), all of which rely on

manipulation of cellular membranes for efficient replication. We hypothesize that

UMB18-mediated increase in total cellular cholesterol, which causes an increase in

membrane rigidity, impedes the ability of these viruses to manipulate membranes and

thus disrupts efficient replication. Previous studies have suggested SARS-CoV-2

infection may downregulate expression of mevalonate pathway genes associated with

cholesterol synthesis(22,23). It would therefore be reasonable to hypothesize that

increasing expression of these genes would have antiviral activity, in alignment with our

data. However, contrasting studies have suggested that SARS-CoV-2 and other

coronaviruses cause an increase in lipid metabolism, mediated by the SREBPs, which

may make our results seem counter-intuitive(24–26). We propose that there may be a

delicate equilibrium in cholesterol levels, or a "Goldilocks" range of cellular cholesterol

necessary to support viral replication; deviations from this balance, whether through

insufficient or excessive cholesterol disrupts membrane dynamics and adversely affects

viral replication. Indeed, our data somewhat support this notion that a delicate balance

of cholesterol is important and both an increase and decrease in cellular cholesterol can

impact viral replication from our data showing that MβCD, which extracts cellular

cholesterol, has antiviral activity (Fig. 5E).

Our results are looking at cellular level changes in cholesterol synthesis, rather than the

context of a high cholesterol organism. It has been suggested that lowered cellular

cholesterol can be antiviral(27,28). But, again, based on our data we suggest there may

be a delicate balance and the disruption of cellular cholesterol either to increase or

decrease can be detrimental to SARS-CoV-2 replication. It is worth noting that many

studies have implicated high cholesterol with increased COVID-19 severity. We do not

think these necessarily impact the potential efficacy of our development antiviral though

since the increased cholesterol seen in humans usually comes from, or with various

other underlying health conditions which can all impact viral disease pathogenesis. In

future work we intend to further study the role these cellular changes have at an

organismal level using animal models of UMB18 treatment.

One hypothesis for the mechanism through which UMB18 is causing the SREBP-

mediated transcriptional changes draws from work by Yang *et al.*(21), which linked

SKIC2 deletion in mice to an increase in mTORC1 signaling. Despite efforts to assess

whether UMB18 activated mTORC1 as assess by phosphorylation levels of S6K1, we

observed no significant changes (Fig. S2). The activation of mTORC1 drives many

cellular changes, one of which is the activation of the SREBP proteins, and thus a link

between the SKI complex and mTORC1 activation could make sense in the context of

our data. It is possible that UMB18 could activate mTORC1 but not at a level to observe

changes to the phosphorylation status of S6K1.

Alternatively, UMB18 may alter the activity of the SKI complex as an RNA helicase and

impact genes that regulate SREBP proteins independently of mTORC1. For instance,

the SKI complex has been shown to have a role in resolution of stalled ribosomes(29)

which could be hypothesized to be disrupted by treatment with UMB18. Stalled

ribosomes can activate ZAKα, which can in turn activate JNK signaling(30) and JNK

signaling has been linked to activation of SREBPs(31). Future research will explore the

interaction between UMB18, the SKI complex and these various cellular signaling

pathways.

Notably, our findings demonstrate that *Srebf1* knockdown appeared to more potently

attenuate the antiviral activity of UMB18 compared to *Srebf2*. It is generally considered

that SREBP2 is primarily involved in expression of genes in the cholesterol synthesis

pathway, while SREBP1 predominantly influences fatty acid synthesis genes(16,17).

However, there are two isoforms of SREBP1 generated from alternative start sites,

SREBP1a and SREBP1c(15). Studies have suggested that SREBP1a can play roles in

promoting expression of genes involved in cholesterol synthesis, along with lipogenic

genes(16). Thus, our data may point towards a nuanced interaction between the SKI

complex and SREBP activation, warranting further research into how SREBP isoforms

contribute to cholesterol regulation and the antiviral activity of UMB18.

This study advances our understanding of our developmental host-directed, broad-

spectrum antiviral therapeutic targeting the SKI complex. Building on our previous work

which established the SKI complex as a host-directed target to inhibit diverse highly

pathogenic human viruses, we reveal a crucial role for increased cellular cholesterol in

mediating this antiviral effect. Our findings prompt further investigation into the

mechanisms by which the SKI complex, an RNA helicase associated with 3' to 5' RNA

degradation, influences metabolic pathways crucial for lipid and cholesterol synthesis, a

finding that will be the focus of future work.

Material and Methods

Mammalian cell culture

A549 cells and A549 cells overexpressing human ACE2 (A549-ACE2, kindly provided

by Dr. Brad Rosenberg (MSSM, NYC)) were cultured in DMEM (Quality Biologicals),

supplemented with 10% (v/v) heat inactivated fetal bovine serum (FBS; Sigma) and 1%

(v/v) penicillin/streptomycin (pen/strep, 10,000 U/ml / 10 mg/ml; Gemini Bio-Products).

VeroE6 cells overexpressing TMPRSS2 (VeroT, kindly provided by Shutoku Matsuyama

(National Institute of Infectious Diseases, Japan - Shirato2018_Virology) were cultured

in DMEM, supplemented with 10% FBS, 1% pen/strep and 1% (v/v) L-glutamine (2 mM

final concentration, Gibco). Cells were maintained at 37°C and 5% CO_2.

Chemical compounds

SKI targeting compounds were synthesized by, and purchased from, Dalriada Drug

Discovery and were previously described(1). Methyl-beta-cyclodextrin (MβCD) was

purchased from Sigma.

Viruses and infections

Details on stock production have been described previously(1). SARS-CoV-2 strain

Washington 1 (WA1) was initially provided by the CDC (BEI #NR-52281). All

coronavirus work was performed in a Biosafety Level 3 laboratory and approved by our

Institutional Biosafety Committee. In all cases of infection for experiments, cells were

plated one day prior to infection to be at a desired confluence for infection. All viral titer

were determined using plaque assays on VeroT cells as described(32). For

experimental infections, virus was diluted to indicated MOI in media used for culture of

A549 cells with addition of chemical compounds or controls as appropriate. For

experiments using MβCD, 50 mM Hepes (Gibco) was additionally added to the media.

RNA extraction and qRT-PCR

RNA extractions and cDNA reactions were performed as described previously(1) using

Direct-zol RNA miniprep kit (Zymo Reserach) and RevertAid RT Kit (Thermo Scientific).

The primers used are listed in the table below. A QuantStudio 5 (Applied Biosystems)

was used for the qRT-PCR reactions. Loading was normalized using GAPDH and fold

change was determined by calculating ΔΔCT after normalization.

RT-PCR primers

All primers for qRT-PCR were purchased from Integrated DNA Technologies (IDT) using

their pre-designed qPCR assay primers:

Gene target	IDT Assay ID
SKIC2 (A.K.A SKIV2L)	Hs.PT.58.14722008
SKIC3 (A.K.A TTC37)	Hs.PT.58.26762302
SKIC8 (A.K.A WDR61)	Hs.PT.58.157001
HMGCS1	Hs.PT.58.4084870

HMGCR	Hs.PT.58.41105491
MVK	Hs.PT.58.40379926
MVD	Hs.PT.58.39035173
IDI1	Hs.PT.58.1930362
FDPS	Hs.PT.58.27570524
MSMO1	Hs.PT.58.3747602
SCAP	Hs.PT.58.45442299
SREBF1	Hs.PT.58.3359761
SREBF2	Hs.PT.58.45335433
GAPDH	Hs.PT.39a.22214836

Transcriptomic sequencing and analysis

Library preparation and sequencing were carried out by the University of Maryland

Institute of Genomic Sciences (IGS; Baltimore, MD, USA) on an Illumina NovaSeq

6000; 100bp paired-end sequencing on an S4 flow cell (Illumina, San Diego, CA, USA).

Reads were preprocessed using cutadapt v3.4(33), then aligned to the *Mus musculus*

genome (assembly GRCm38) with STAR v2.7.8(34). Gene expression analysis was

carried out using DESeq2 v4.1.0(35) in R (Rstudio, Boston, MA, USA). Pathway

analysis was carried out using Ingenuity Pathway Analysis (QIAGEN, Hilden, Germany)

using an alpha value of $p_{adj} < 0.01$ to call a gene significantly differentially expressed.

Raw sequence data is available in the NCBI Sequence Read Archive under the

accession number PRJNA1055076.

Mass spectrometry imaging

A modified cell culture apparatus was used for compatibility with mass spectrometry

imaging (MSI). Briefly, Superfrost plus gold microscope slides were autoclaved with

flexible silicon 12-well transwell adapters (Flexiperm, Starstedt), dried completely, and

the transwell adapters mounted. Wells were coated with 0.1 mg/mL poly-l-lysine (0.1 M

borate buffer pH 8.5) overnight at 37°C, aspirated, and dried. Cells were grown in the

transwells overnight and subsequently treated with 10μM UB18 or 0.1% DMSO for 24h.

At the end of the incubation, supernatants were aspirated, the transwell mount was

removed, and the slides were submerged in ice-cold 50 mM ammonium formate (pH

6.7) for 1 minute with gentle rocking. Excess buffer was removed by tilting and using

absorbent paper. Slides were dried, flat, in a desiccator for 20 minutes. Silver-assisted

laser desorption ionization (Ag-LDI) was used to ionize cholesterol on a quadropole time

of flight (qTOF) mass spectrometer in imaging mode (Yang2020_JMassSpectrom).

Silver nitrate (8.5 mg/mL in 0.1% trifluoroacetic acid/methanol) was sprayed on the

slides using an HTX M5 Matrix Sprayer with the following settings: 45°C nozzle, 40 mm

height, 24 passes, 0.075 mL/min flow, 500 mm/min speed, 4 mm track spacing, 20 psi,

2 L/min N2, in "CC" pattern, with 0 s dry time between passes. Cholesterol imaging data

were captured from an area of uniform monolayer (>80% confluence) integrity guided by

an optical pre-scan using a Bruker Daltonics timsTOF Flex (Billerica, MA) in positive ion

mode with a 50 μM scan area (11 μM x 11 μM beamscan), over range *m/z* 100-1000

with enhanced quadratic fit calibration to Agilent Tune-Mix, 500 shots per pixel at 10 kHz

with 5-point setting in M5 small laser focus. Total scan area per sample was ~3 mm^2

resulting in ~1200 pixels per replicate, with quadruplicate samples per condition. Data

analyzed in SCiLS Lab (Bruker Daltonics), normalized (total ion current), projected with

hotspotting on, and visualized as the ion *m/z* 493.260 +/-2.2 mDa ([cholesterol+Ag$_{107}$]$^+$)

ion; the [cholesterol+Ag$_{109}$]$^+$ ion data was in agreement. Student's T-test and a receiver-

operator characteristic were performed between vehicle-treated and UMB18-treated

cells on quadruplicate samples.

Expression and purification of SKIC8

Transformed *E. coli* BL21 (DE3) cells containing the full-length human *SKIC8* gene with

a HisX6 tag in the pET-28a(+) vector were grown in Luria-Bertani (LB)-Kanamycin broth

at 37ºC until the optical density at 600 nm equaled 0.8. SKIC8 protein expression was

induced with 1 mM Isopropyl β-D-1-thiogalactopyranoside for 20 h at 15°C. Cells were

lysed using Bugbuster (Novagen) supplemented with protease and phosphatase

inhibitors (Thermo Scientific). Lysates were clarified by centrifugation at 14,000 rpm for

30 min. and the supernatant was applied to a Talon Metal Affinity Resin (Clontech)

column equilibrated in 50 mM sodium phosphate (pH 7.8) containing 10 mM imidazole

and 300 mM NaCl. The column was washed with the same sodium phosphate buffer

containing 20 mM imidazole. SKIC8 was eluted in 50 mM sodium phosphate (pH 7.8)

containing a 50-250 mM imidazole gradient and 300 mM NaCl. The fractions containing

SKIC8, as determined following SDS-PAGE, were pooled, and concentrated with an

Amicon ultrafiltration cell (Millipore, Bedford, MA). The concentrated protein was

dialyzed against 20 mM Tris-HCl (pH 7.8) containing 150 mM NaCl.

Surface Plasmon Resonance (SPR)

SKIC8 and test compound interactions were evaluated using a Biacore T-200

instrument (GE Healthcare). A carboxymethylated dextran chip (CM5, GE Healthcare)

was used to immobilize SKIC8. A net positive charge is required for direct amine

coupling; therefore, the proteins were diluted to a final concentration of 30 µg/ml in 10

mM sodium acetate buffer pH 4.0. The protein was injected at 2 µl/min until

approximately a 4500 resonance/response unit level of immobilized SKIC8 was

reached, and then the flow cell was deactivated with 1 M ethanolamine HCl (pH 8.5).

Degassed 20 mM Hepes (pH 7.4) containing 5 mM MgCl2, 100 mM NaCl, and 0.005%

Nonidet P40 was utilized as continuous running buffer for all experiments. Test

compound binding interactions with SKIC8 were investigated by injecting test compound

(1.0 –50 µM) at a flow rate of 20 µl/min. The resulting sensorgrams were analyzed using

BIAevaluation 3.1 software (GE Healthcare) to determine the dissociation constant (K_D)

as previously described(36).

RNAi knockdown

Cells were seeded to 24-well plates at a density of 4.4e4 cells/well one day prior to

transfection. Transfections were performed using Oligofectamine (Thermo Scientific) to

the manufacture's methods. Briefly, per well of transfection, 4.4µl Opti-MEM (Gibco) was

mixed with 2.2µl Oligofectamine and incubated at room temperature (RT°C) for 5 min,

while 35.5µl Opti-MEM was mixed with 0.8µl 50mM siRNA in a separate tube. The two

tubes were combined and incubated at RT°C for 20 minutes prior to addition of 177µl

Opti-MEM. Cells were washed 1x with PBS and 1x with OptiMEM then 200µl of

transfection mixture was added per well. All siRNAs were purchased from Sigma using

their Rosetta prediction system for predesigned sequences. MISSION siRNA Universal

Negative Control #1 (Sigma) was used as the scrambled control sequence. Sequences

are displayed in table:

<u>RNAi sequences</u>

Gene name	siRNA ID
SKIC2 (A.K.A SKIV2L)	SASI_Hs01_00196881
SKIC3 (A.K.A TTC37)	SASI_Hs01_00010354
SKIC8 (A.K.A WDR61)	SASI_Hs01_00155941
SCAP si-1	SASI_Hs01_00032110
SCAP si-2	SASI_Hs01_00032111
SREBF1 si-1	SASI_Hs01_00051828
SREBF1 si-2	SASI_Hs01_00051829
SREBF2 si-1	SASI_Hs01_00075424
SREBF2 si-2	SASI_Hs01_00075425
SREBF2 si-3	SASI_Hs01_00075426

Western blotting

Whole cell lysates were made using Triton-X100 (Tx100) lysis buffer lysis buffer (1%

Tx100 (v/v), 150mM NaCl , 50mM Tris (all Sigma), pH8). In all cases 1x cOmplete Mini,

EDTA-Free protease inhibitor cocktail (Roche) was added to the lysis buffer and for

experiments involving S6K and phospho-S6K (pS6K), Phosphatase Inhibitor Cocktail 2

(Sigma) was additionally used. Western blotting procedures have been previously

described(37). Primary antibodies used as follows: p70 S6K (Cell Signaling

Technologies #9202S, 1:1000 dilution), phospho-p70 S6K (T421/S424) (Cell Signaling

Technologies #9204S, 1:1000 dilution) and mouse anti-tubulin (clone DMA1A, Sigma,

1:1000). Secondary antibodies were used as follows: goat anti-rabbit HRP and goat

anti-mouse HRP (both 0.8 mg/ml, Thermo Scientific, 1:10,000 diluted) which were

detected using ECL Prime (Amersham).

Data analysis

All graphs produced and statistical analyese performed using GraphPad Prism 10.3

software.

730 Acknowledgements

731 Acknowledgments: We thank We would like to thank all of the Frieman, Jackson and

732 Coughlan lab members for their help in this study.

733 Funding: MBF is supported by an endowment from The Alicia and Yaya Foundation.

734 The funders had no role in study design, data collection and analysis, decision to

735 publish, or preparation of the manuscript.

737 References

1. Weston S, Baracco L, Keller C, Matthews K, McGrath ME, Logue J, et al. The SKI

complex is a broad-spectrum, host-directed antiviral drug target for coronaviruses,

influenza, and filoviruses. Proc Natl Acad Sci U S A. 2020 Dec 1;117(48):30687–98.

2. Cox M, Peacock TP, Harvey WT, Hughes J, Wright DW, Willett BJ, et al. SARS-CoV-

2 variant evasion of monoclonal antibodies based on in vitro studies. Nat Rev

Microbiol. 2023 Feb;21(2):112–24.

3. Sanderson T, Hisner R, Donovan-Banfield I, Hartman H, Løchen A, Peacock TP, et

al. A molnupiravir-associated mutational signature in global SARS-CoV-2 genomes.

Nature. 2023 Nov;623(7987):594–600.

4. Houseley J, Tollervey D. The Many Pathways of RNA Degradation. Cell. 2009 Feb

20;136(4):763–76.

5. Schneider C, Tollervey D. Looking into the barrel of the RNA exosome. Nat Struct

Mol Biol. 2014 Jan;21(1):17–8.

6. Januszyk K, Lima CD. The eukaryotic RNA exosome. Curr Opin Struct Biol. 2014

Feb;24:132–40.

7. Johnson SJ, Jackson RN. Ski2-like RNA helicase structures: common themes and

complex assemblies. RNA Biol. 2013 Jan;10(1):33–43.

8. Toh-E A, Guerry P, Wickner RB. Chromosomal superkiller mutants of

Saccharomyces cerevisiae. J Bacteriol. 1978 Dec;136(3):1002–7.

9. Widner WR, Wickner RB. Evidence that the SKI antiviral system of Saccharomyces

cerevisiae acts by blocking expression of viral mRNA. Mol Cell Biol. 1993

Jul;13(7):4331–41.

10. Goldstein JL, Brown MS. Regulation of the mevalonate pathway. Nature. 1990

Feb;343(6257):425–30.

11. Luo J, Yang H, Song BL. Mechanisms and regulation of cholesterol homeostasis.

Nat Rev Mol Cell Biol. 2020 Apr;21(4):225–45.

12. Cerqueira NMFSA, Oliveira EF, Gesto DS, Santos-Martins D, Moreira C, Moorthy

HN, et al. Cholesterol Biosynthesis: A Mechanistic Overview. Biochemistry. 2016 Oct

4;55(39):5483–506.

13. Brown MS, Goldstein JL. The SREBP Pathway: Regulation of Cholesterol

Metabolism by Proteolysis of a Membrane-Bound Transcription Factor. Cell. 1997

May 2;89(3):331–40.

14. Hua X, Nohturfft A, Goldstein JL, Brown MS. Sterol resistance in CHO cells traced to

point mutation in SREBP cleavage-activating protein. Cell. 1996 Nov 1;87(3):415–

26.

15. Shimomura I, Shimano H, Horton JD, Goldstein JL, Brown MS. Differential

expression of exons 1a and 1c in mRNAs for sterol regulatory element binding

protein-1 in human and mouse organs and cultured cells. Journal of Clinical

Investigation. 1997 Mar 3;99(5):838.

16. Horton JD, Goldstein JL, Brown MS. SREBPs: activators of the complete program of

cholesterol and fatty acid synthesis in the liver. J Clin Invest. 2002 May

1;109(9):1125–31.

17. Xu X, So JS, Park JG, Lee AH. Transcriptional control of hepatic lipid metabolism by

SREBP and ChREBP. Semin Liver Dis. 2013 Nov;33(4):301–11.

18. Nohturfft A, DeBose-Boyd RA, Scheek S, Goldstein JL, Brown MS. Sterols regulate

cycling of SREBP cleavage-activating protein (SCAP) between endoplasmic

reticulum and Golgi. Proceedings of the National Academy of Sciences. 1999 Sep

28;96(20):11235–40.

19. Sun LP, Li L, Goldstein JL, Brown MS. Insig required for sterol-mediated inhibition of

Scap/SREBP binding to COPII proteins in vitro. J Biol Chem. 2005 Jul

15;280(28):26483–90.

20. Nagoshi E, Imamoto N, Sato R, Yoneda Y. Nuclear import of sterol regulatory

element-binding protein-2, a basic helix-loop-helix-leucine zipper (bHLH-Zip)-

containing transcription factor, occurs through the direct interaction of importin beta

with HLH-Zip. Mol Biol Cell. 1999 Jul;10(7):2221–33.

21. Yang K, Han J, Asada M, Gill JG, Park JY, Sathe MN, et al. Cytoplasmic RNA quality

control failure engages mTORC1-mediated autoinflammatory disease. Journal of

Clinical Investigation. 2022 Jan 18;132(2):e146176.

22. Gomez Marti JL, Wells A, Brufsky AM. Dysregulation of the mevalonate pathway

during SARS-CoV-2 infection: An in silico study. Journal of Medical Virology.

2021;93(4):2396–405.

23. Wang X, Zhao Y, Yan F, Wang T, Sun W, Feng N, et al. Viral and Host

Transcriptomes in SARS-CoV-2-Infected Human Lung Cells. J Virol. 2021 Aug

25;95(18):e0060021.

24. Dias SSG, Soares VC, Ferreira AC, Sacramento CQ, Fintelman-Rodrigues N,

Temerozo JR, et al. Lipid droplets fuel SARS-CoV-2 replication and production of

inflammatory mediators. PLOS Pathogens. 2020 Dec 16;16(12):e1009127.

25. Gao P, Ji M, Liu X, Chen X, Liu H, Li S, et al. Apolipoprotein E mediates cell

resistance to influenza virus infection. Science Advances. 2022 Sep

21;8(38):eabm6668.

26. Yuan S, Chu H, Chan JFW, Ye ZW, Wen L, Yan B, et al. SREBP-dependent

lipidomic reprogramming as a broad-spectrum antiviral target. Nat Commun. 2019

Jan 10;10(1):120.

27. Wing PAC, Schmidt NM, Peters R, Erdmann M, Brown R, Wang H, et al. An ACAT

inhibitor suppresses SARS-CoV-2 replication and boosts antiviral T cell activity.

PLOS Pathogens. 2023 May 3;19(5):e1011323.

28. Wang H, Yuan Z, Pavel MA, Jablonski SM, Jablonski J, Hobson R, et al. The role of

high cholesterol in SARS-CoV-2 infectivity. Journal of Biological Chemistry [Internet].

2023 Jun 1 [cited 2024 Aug 28];299(6). Available from:

https://www.jbc.org/article/S0021-9258(23)01791-X/abstract

29. Zinoviev A, Ayupov RK, Abaeva IS, Hellen CUT, Pestova TV. Extraction of mRNA

from Stalled Ribosomes by the Ski Complex. Molecular Cell. 2020 Mar

19;77(6):1340-1349.e6.

30. Snieckute G, Genzor AV, Vind AC, Ryder L, Stoneley M, Chamois S, et al. Ribosome

stalling is a signal for metabolic regulation by the ribotoxic stress response. Cell

Metabolism. 2022 Dec;34(12):2036-2046.e8.

31. Kotzka J, Knebel B, Haas J, Kremer L, Jacob S, Hartwig S, et al. Preventing

Phosphorylation of Sterol Regulatory Element-Binding Protein 1a by MAP-Kinases

Protects Mice from Fatty Liver and Visceral Obesity. PLOS ONE. 2012 Feb

27;7(2):e32609.

32. Weston S, Coleman CM, Haupt R, Logue J, Matthews K, Li Y, et al. Broad Anti-

coronavirus Activity of Food and Drug Administration-Approved Drugs against

SARS-CoV-2 In Vitro and SARS-CoV In Vivo. J Virol. 2020 Oct 14;94(21):e01218-

20.

33. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnet.journal. 2011 May 2;17(1):10–2.

34. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:

ultrafast universal RNA-seq aligner. Bioinformatics. 2013 Jan 1;29(1):15–21.

35. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biology. 2014 Dec 5;15(12):550.

36. Burkhard KA, Chen F, Shapiro P. Quantitative analysis of ERK2 interactions with

substrate proteins: roles for kinase docking domains and activity in determining

binding affinity. J Biol Chem. 2011 Jan 28;286(4):2477–85.

37. Weston S, Matthews KL, Lent R, Vlk A, Haupt R, Kingsbury T, et al. A Yeast

Suppressor Screen Used To Identify Mammalian SIRT1 as a Proviral Factor for

Middle East Respiratory Syndrome Coronavirus Replication. J Virol. 2019 Aug

15;93(16):e00197-19.

Author Contributions:

-    Conceptualization: SW, MBF

-    Data curation: LT,

-    Formal analysis: SW, LT, AS, PS,

-    Funding acquisition: MBF

-    Investigation: SW, LB, AS, GK,

-    Methodology: SW, LT, AS, PS,

-    Resources: ADM

-    Supervision: MBF

-    Visualization: SW, LT, AS, PS,

-    Writing – original draft: SW, AS, PS, MBF

-    Writing – review & editing: SW, MFB

Competing Interests: The authors declare the following competing interesting: M.B.F is

the lead inventor along with S.W. and A.D.M as contributing inventors for patents

held for development of SKI targeting compounds (of which UMB18 is one) with

antiviral activity. The patent filings were through University of Maryland, Baltimore

and listed as PCT/US20/36483 and PCT/US2021/1061863. There has been a

corporate partnership with Aikido Pharmaceuticals for developmental research on

the compound along with others that target the SKI complex. This developmental

work has not influenced the academic research presented in this paper.

Patents: Filed patents: Patents on UMB18 and related compounds have been filed

through University of Maryland, Baltimore and listed as PCT/US20/36483 and

PCT/US2021/1061863.

Data and Material Availability: All data needed to evaluate the conclusions in this paper

are present in the paper and supplementary materials. Data are also deposited at

https://osf.io/63h8t/.

Figure Legend

Figure 1 – Transcriptomic analysis of UMB18 treated cells.

A) Volcano plots showing differential expression of genes at 24h (top) and 48h (bottom)

in uninfected A549 cells treated with UMB18 compared to untreated, uninfected A549

cells. Cholesterol biosynthesis genes are colored blue and labeled. A horizontal dashed

line is drawn at p_{adj} = 0.01, vertical dashed lines are drawn at +/- 2-fold change. B) Plot

showing significantly enriched pathways in uninfected A549 cells treated with UMB18

compared to untreated, uninfected A549 cells at 24h. Orange bars show pathways with

predicted increased pathway output (z > 2); white bars show pathways with no predicted

884 activity pattern ($z = 0$); blue bars indicate a predicted decrease in pathway output ($z < -$

885 2). C) Venn Diagram comparing upregulation of cholesterol biosynthesis genes in

886 UMB18-treated A549 cells against timepoint- and infection-matched, untreated controls.

887 The number and color intensity in each area corresponds to the number of cholesterol

888 biosynthesis genes that are significantly upregulated ($p_{adj} < 0.01$).

Figure 2 – UMB18 causes increased expression of mevalonate pathway genes

and cellular cholesterol.

A) Schematic representation of the mevalonate pathway leading to cholesterol

synthesis. In blue are metabolic products of the pathway and in black are the enzymes

that regulate the various steps with those that are highlighted being analyzed by qRT-

PCR through the paper. B and C) A549-ACE2 cells (B) or A549 cells (C) were treated

with 10µM UMB18 or 0.1% DMSO vehicle control and collected at 24 h or 48 h in TRIzol

for qRT-PCR analysis of expression of various genes throughout the mevalonate

pathway. The CT values were set relative to GAPDH as a housekeeping gene control

and then relative fold change calculated between UMB18 treated cells and DMSO

treated cells. D) Relative intensity cloud projection with box plot of *m/z* 493.260 +/- 2.2

mDa ([cholesterol+Ag$_{107}$]$^+$) ion per image pixel for UMB18- and DMSO-treated cells with

area-under-the-curve (AOC) from a receiver-operator characteristic analysis. Student's

T-test base on quadruplicate samples.

Figure 3 – UMB18 directly binds the SKI complex and that is required for increased cellular cholesterol.

A) Binding of UMB18 to SKIC8. The binding affinity of immobilized SKIC8 to UMB18 at various concentrations was quantified by SPR as described in the methods. B) A549-ACE2 cells were transfected with siRNAs targeting each of the three SKI complex genes (SKIC2, SKIC3, and SKIC8) for 72h prior to treatment with 10µM UMB18 or 0.1% DMSO for 24 h. Cells were collected in TRIzol and qRT-PCR performed for each of the three SKI genes in all of the various conditions after normalization with GAPDH. Scrambled siRNA (siScr) cells treated with DMSO (siScr/DMSO) were used as the control to determine relative fold change. C-E) Using the same RNA samples and qRT-PCR data in B, the Ct values were normalized to GAPDH and the ΔCt values for three mevalonate pathway genes (MVD, FDPS and MSMO1, respectively) from the siScr/DMSO samples treated with either UMB18 or DMSO are plotted. Student's t-test was performed and the p-values are labled for the comparison. F) From the same RNA samples, the relative fold change for the same three mevalonate pathway genes compared to siScr/DMSO in all other conditions was determined. Two-way ANOVA testing was performed for all comparisons, but only the comparisons between UMB18 or DMSO treatment for each siRNA treatment are shown. Unless otherwise indicated, there were no significant differences (ns), based on p value < 0.05. G) To further analyze the impact of SKI siRNA transfection on UMB18 treatment the fold change for siScr/UMB18 was set relative to the fold change in the SKI siRNA transfected cells, treated with UMB18. In this comparison a fold change of 1 would mean no difference between SKI siRNA and scrambled siRNA with UMB18 and a fold change below 1

931   means a lower change from UMB18 treatment in SKI siRNA. All RNA data are from two

932   experiments performed in duplicate wells.

Figure 4 – UMB18 requires SCAP and SREBP genes to increase mevalonate

pathway gene expression.

A549-ACE2 cells were transfected with unique siRNA sequences targeting *SCAP* (A, D,

G)*, Srebf2* (B, E and H) or *Srebf 1* (C, F and I). The cells were then treated with 10µM

UMB18 or 0.1% DMSO for 24 h and collected for qRT-PCR targeting the specified

genes with normalization to GAPDH as a housekeeping gene. Fold change in gene

expression was calculated compared to scrambled siRNA (siScr) transfected cells

treated with DMSO (siScr/DMSO) for each of the conditions. A-C) Fold change relative

to siScr/DMSO for each of the siRNA transfection conditions with DMSO treatment.

Two-way ANOVA analysis was performed for comparison of all conditions and

significant p values (<0.05) have been plotted. For all other comparisons within each

gene target there were no significant differences between the siRNA sequences. D-F)

Fold change relative to siScr/DMSO for UMB18 treatment siRNA transfected cells. Two-

way ANOVA analysis were performed for all conditions and significant p-values (<0.05)

within each gene target are plotted. Where no p-value has been plotted there were no

significant differences within each gene comparison. G-F) As described in Figure 3G,

the fold change of UMB18 treated siRNA samples was set relative to the fold change of

UMB18 treated siScr and plotted, such that a fold change of 1 would mean no

difference between *SCAP/Srebf2/Srebf1* siRNA and scrambled siRNA with UMB18 and

a fold change below 1 means a lower change from UMB18 treatment in the targeted

siRNA samples. Data are from three independent experiments performed in triplicate in

all cases.

Figure 5 – Increased cellular cholesterol is required for UMB18-mediated antiviral activity.

A549-ACE2 cells were transfected with siRNA sequences targeting *Srebf1* (A), *SCAP* or *Srebf2* (sequences as used in Figure 4, *SCAP* and *Srebf2* transfections were experimentally performed together while *Srebf1* were performed separately). Following transfection, cells were infected with SARS-CoV-2 at MOI 0.1 and treated with 10µM UMB18 or 0.1% DMSO for 24 h. Supernatant was collected from the cells and viral titer determined by plaque assay on VeroE6 cells expressing TMPRSS2. The titer is plotted as plaque forming units/ml from each of the different transfection and treatment conditions. Using scrambled siRNA (Scr) as the control. Two-way ANOVA analysis was performed and the p-values are plotted. Data are from three independent experiemnts

performed in triplicate. C) Fold change between titer of DMSO treated and UMB18

treated in each transfection condition summarized in a table format. D) A549-ACE2 cells

were treated with the indicated concentrations of methyl-beta-cyclodextrin (MβCD) for

24 h prior to fixation with neutral buffered formalin. Cells were stained with Hoechest to

label nuclei and cell count per well (96 well format) was quantified with a Celigo high

content imager. Reported is the percentage total cell count compared to untreated cells

as a measure of cytotoxicity from MβCD treatment. E) A549-ACE2 cells were infected

with SARS-CoV-2 at MOI 3 and treated with the indicated concentrations of MβCD for

24 h. Cells were similarly fixed with neutral buffered formalin then immunofluorescence

labeled for SARS-CoV-2 N protein and stained with Hoechest. High content imaging

was used to quantify percentage infection and the plotted data are relative parentage

infection compared to untreated control cells. F) As in E, cells were infected with SARS-

CoV-2 at MOI 3 and treated with or without MβCD with either 10µM UMB18 or 0.1%

DMSO. Cells were similarly fixed and labeled for high content imaging of percentage

infection. Plotted are the relative infection percentage to 0.1% DMSO treated cells for

each of the different conditions. An unpaired t-test was performed for for each condition

and the p-valus are reported. Data are from three independent experiments.

Supplemental Figure Captions

Supplemental Figure 1 – Serum impacts UMB18 induced transcriptional changes

and virus replication.

A and B) A549-ACE2 cells were treated with 10 µM UMB18 or 0.1% DMSO for 24 hours

in cell culture media supplemented with serum or without supplementation. Cells were

harvested in TRIzol and qRT-PCR performed for the indicated genes. A) Fold change in

gene transcript in UMB18 treated cells relative to DMSO treated. B) The same set of

experimental data with fold change being in serum free media relative to serum

containing media. C) A549-ACE2 cells were infected with SARS-CoV-2 at MOI 0.5 in

serum containing or serum free media with 10 µM UMB18 or 0.1% DMSO, or neither

and left in just media. 24-hour post infection supernatant was collected and used to titer

virus by plaque assay on VeroTMPRSS2 cells. Significance calculated by two-way

ANOVA with a Tukey correction. Only statically significant (p < 0.05) comparisons are

shown on the graph.

1002

Supplemental Figure 2 – UMB18 does not appear to activate mTOR signaling as

assessed by phospho-S6K1.

A) A549-ACE2 cells were plated to plastic and the following day were treated with

100ng recombinant EGF, or PBS carrier alone. At time points between 5 and 30 min,

cells were collected in Triton-X100 lysis buffer and prepared for western blotting (see

Materials and Methods). Samples were separated on gels and blotted for phospho-

S6K1 or S6K1 and tubulin was used as a loading control. All probes were detected with

HRP. Density of bands was determined and quantified by setting relative to tubulin

control and then reported as a relative band density of EGF treated compared to PBS

treated at each time point. B and C) A549-ACE2 cell were plated to plastic and the

following day treated with 10µM UMB18 or 0.1% DMSO for the indicated time points or

left untreated (for 4 hours in B). At each timepoint, cells were collected, and western

blots run similarly to A. All gels were imaged with HRP, except tubulin in B which was

visualized by AlexaFluor 488. Density of UMB18 compared to DMSO at each time point

is reported as in A.

1018

1019