1

Trehalose supports the growth of *Aedes aegypti* cells and modifies gene expression and dengue virus replication

Andrew D. Marten[1], Douglas P. Haslitt[1], Chad A. Martin[1], Daniel H. Swanson[2], Karishma Kalera[2,3], Ulysses G. Johnson[2,3], Benjamin M. Swarts[2,3], and Michael J. Conway[1]

[1]Foundational Sciences, Central Michigan University College of Medicine, Mount Pleasant, MI 48859, USA.

[2]Department of Chemistry and Biochemistry, Central Michigan University, Mount Pleasant, MI 48859, USA.

[3]Biochemistry, Cell, and Molecular Biology Graduate Programs, Central Michigan University, Mount Pleasant, MI 48859, USA.

16
17

Address correspondence to: Foundational Sciences, Central Michigan University

College of Medicine, Mount Pleasant, MI, 48859

E-mail: michael.conway@cmich.edu. Phone: (989) 774-3930, Fax: (989) 774-3462

22
23
24

Short title: Trehalose metabolism in *Aedes aegypti*

26
27
28
29

Abstract

Trehalose is a non-reducing disaccharide that is the major sugar found in insect hemolymph fluid. Trehalose provides energy, and promotes growth, metamorphosis, stress recovery, chitin synthesis, and insect flight. Trehalase is the only enzyme responsible for the hydrolysis of trehalose, which makes it an attractive molecular target. Here we show that *Aedes aegypti* (Aag2) cells express trehalase and that they can grow on trehalose-containing cell culture media. Trehalase activity was confirmed by treating Aag2 cells with trehalase inhibitors, which inhibited conversion of trehalose to glucose and reduced cell proliferation. Cell entry of a fluorescent trehalose probe was dependent on trehalose concentration, suggesting that trehalose moves across the cell membrane via passive transport. Culturing Aag2 cells with trehalose-containing cell culture media led to significant changes in gene expression, intracellular lipids, and dengue virus replication and specific infectivity, and increased their susceptibility to trehalase inhibitors. These data describe an *in vitro* model that can be used to rapidly screen novel trehalase inhibitors and probes and underscores the importance of trehalose metabolism in *Ae. aegypti* physiology and transmission of a mosquito-borne virus.

Introduction

Aedes aegypti is the primary vector of several important global human pathogens including dengue virus (DENV), Zika virus (ZIKV), and chikungunya virus (CHIKV). There are no targeted antivirals for these viruses and the currently licensed DENV vaccine is restricted to a subset of the human population due to the potential of eliciting antibody-dependent enhancement (ADE) in those who have not previously been infected (1-6). Further, resistance to pesticides including carbamates, organochlorines, organophosphates, and pyrethroids has been detected in the Americas, Africa, and Asia. New strategies to control disease vectors are desperately needed (7-13).

Trehalose is a non-reducing disaccharide and is the most prevalent sugar found in insect hemolymph fluid. Trehalose provides energy but also promotes growth, metamorphosis, stress recovery, chitin synthesis, and insect flight (14-28). Previous research has shown that trehalose is synthesized in the insect fat body from glucose 6-phosphate and UDP-glucose by the conserved enzymes trehalose 6-phosphate synthase (TPS) and trehalose 6-phosphate phosphatase (TPP) (16, 22, 29, 30). The hydrolysis of trehalose is under the enzymatic control of trehalase, which catalyzes the conversion of trehalose into two glucose molecules (22, 30). There are two forms of insect trehalase. Trehalase 1 (Tre-1) is soluble and has been purified from hemolymph, midgut goblet cells, and eggs. Trehalase 2 (Tre-2) is a membrane bound form and has been identified in flight muscle, follicle cells, ovary cells, spermatophore, midgut, and brain tissue (22). Although it is known that the facilitated trehalose transporter (TRET1) transfers newly synthesized

trehalose from the fat body across the cellular membrane and into the hemolymph, it is

unclear if trehalose transporters exist in peripheral tissues and import this sugar (22).

Trehalose is not synthesized by human cells, but it is present in varying quantities

in certain foods such as mushrooms, yeast, honey, beans, seaweeds, and shellfish, and

synthesized by certain bacterial species that comprise the microbiome (31). Humans

express a membrane-bound trehalase enzyme in the small intestine's brush border cells,

which catalyzes dietary trehalose to glucose (31). Mammalian cells do not normally engage

with high concentrations of trehalose, and previous research indicates that treatment of

mammalian cells with trehalose induces autophagy through an MTOR-independent

mechanism. This has been associated with both neuroprotective and antiviral activities (32-

39).

Our previous research revealed that treatment of *Ae. aegypti* with the trehalase

inhibitor validamycin A (ValA) promoted hypoglycemia and trehalose accumulation, and

corresponded with delayed larval and pupal development, ultimately preventing flight in

adult mosquitoes (23). Unfortunately, a novel trehalose analogue called 5-ThioTre was

unable to recapitulate the activity of ValA (23). We sought to develop an *in vitro* model

that would facilitate screening of novel trehalose analogues and to determine the impact of

trehalose metabolism on a mosquito-borne virus. Here, we show that the *Ae. aegypti* cell

line (Aag2) expresses a secreted trehalase isoform, and that this enzyme facilitates growth

when cell culture media glucose is exchanged for trehalose. Specifically, the trehalase

inhibitor validamycin A (ValA) and novel trehalase inhibitors prevented conversion of cell

culture media trehalose to glucose, and treatment with ValA significantly reduced cell

proliferation. Cell entry of a fluorescent trehalose probe was dependent on trehalose

concentration, suggesting that trehalose moves across the cell membrane via passive

transport. Importantly, RNA-Seq analysis of Aag2 cells grown in trehalose-containing cell

culture media revealed significant changes in gene expression in multiple cellular

pathways, including those involved in integral/intrinsic membrane components. Trehalose

also increased the concentration of intracellular lipids and viral RNA (vRNA), shedding of

infectious virus, and the specific infectivity of DENV virions, while increasing the

susceptibility of Aag2 cells and DENV to trehalase inhibitors. This manuscript describes

the use of an *in vitro* model that can be used to interrogate trehalose metabolism in insect

cells and reveals how trehalose modulates mosquito cell biology and virus infection.

Methods

Cells and reagents. The *Ae. aegypti* cell line, Aag2 (ATCC, VA), was used for *in vitro*

studies. Aag2 cells were maintained at 28 °C with 5% CO_2 in complete cell culture media

containing DMEM no glucose media supplemented with 4,500 mg/L glucose, 10% heat-

inactivated fetal bovine serum (FBS) (Gemini, CA), 1% penicillin-streptomycin (Thermo

Fisher Scientific, MA) and 1% tryptose phosphate broth (Sigma-Aldrich, MO). Aag2 cells

were also grown using the above cell culture media with equimolar concentrations of

trehalose (2,250 mg/L) (Sigma-Aldrich, MO) instead of glucose. It is important to note that

the trehalose-containing cell culture media retained low concentrations of glucose from

FBS and tryptose phosphate broth. The glucose concentration in our trehalose-containing

cell culture media was approximately 125 mg/L but rapidly increased in glucose

concentration after inoculation onto Aag2 cells. The virus strain used was dengue virus

type 2 New Guinea C (DENV2 NGC). DENV2 stocks were made by infecting Aag2 cells

at a MOI of 1.0 and cell-free virus stocks were harvested 8 days post-infection (dpi) and

stored at −80 °C until use.

Trehalose analogues. Validamycin A (ValA) (Cayman Chemical, MI) was the only

commercially available trehalase inhibitor available at the time of this study. The synthesis

of trehalose analogues was carried out using a chemoenzymatic method as previously

reported for 5-ThioTre (40) and 6-TreNH$_2$ (41). The water solubility of 5-ThioTre and 6-

TreNH$_2$ is predicted to be very close to trehalose and stocks were generated in water at 100

mM.

Synthesis of fluorescent trehalose derivative NBD-Trehalose. To a 25 mL round bottom

flask flushed under argon was added anhydrous *N,N*-dimethylformamide (10 mL), 6-

TreNH$_2$ (10.9 mg, 0.032 mmol), 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole (7.6 mg, 0.038

mmol), and *N,N*-diisopropylethylamine (8 μL, 0.043 mmol). After stirring at room

temperature under argon overnight, thin-layer chromatography indicated that the reaction

was complete. The crude product was concentrated by rotary evaporation and purified by

reverse phase chromatography on a Biotage Isolera One automated flash chromatography

system (10 g C18 column; 0% CH$_3$CN in H$_2$O → 80% CH$_3$CN in H$_2$O) to give NBD-

Trehalose (12.9 mg, 80%) as an orange solid. [1]H NMR (500 MHz, D$_2$O): δ 8.38 (d, *J* = 9.0

Hz, 1 H), 6.38 (d, *J* = 9.0 Hz, 1 H), 5.18 (d, *J* = 4.0 Hz, 1 H), 5.06 (d, *J* = 3.5 Hz, 1 H),

4.11-4.07 (m, 1 H), 3.94 (bs, 1 H), 3.87-3.76 (m, 5 H), 3.71 (dd, $J = 5.5, 12.0$ Hz, 1 H),

3.67 (dd, $J = 4.0, 10.0$ Hz, 1 H), 3.47 (t, $J = 10.0$ Hz, 1 H), 3.46 (dd, $J = 3.5, 10.0$ Hz, 1 H),

3.38 (t, $J = 10.0$ Hz, 1 H). ^{13}C NMR (125.7 MHz, D$_2$O): δ 146.46, 144.32, 143.82, 120.62,

138.70, 100.64, 93.47, 93.33, 72.58, 72.51, 71.17, 71.34, 70.93, 70.89, 70.51, 69.61, 60.49,

48.59. HRMS (ESI-TOF) m/z: [M + Na]$^+$ Calcd for C$_{18}$H$_{24}$N$_4$O$_{13}$Na 527.1238; Found

527.1249. Stocks of NBD-Trehalose were generated in dimethylsulfoxide (DMSO) at 100

mM.

Cell proliferation and glucose assays. Cell proliferation assays were performed by

seeding 100 Aag2 cells into a 96 well plate in triplicate. The total number of viable cells

per well were quantified using trypan blue exclusion each day starting on day 0 and ending

on day 7. This assay was repeated using cells that were fed either glucose or trehalose-

containing cell culture media in the presence or absence of 1 mM validamycin A (ValA).

A Glucose Colorimetric Assay Kit (Cayman Chemical, MI) was used to assay glucose

concentration in Aag2 cell-free supernatant from cells grown in either glucose or trehalose-

containing cell culture media each day starting on day 0 and ending on day 3. Glucose

concentration was also assayed in Aag2 cell-free supernatant from cells grown in trehalose-

containing cell culture media and in the presence of increasing concentrations of ValA, and

trehalose analogues 5-ThioTre and 6-TreNH2, which are described above. Glucose

concentration was also assayed in Aag2 cell-free supernatant from cells that were infected

with 1 MOI DENV2 every other day starting 1-day post-infection (dpi) and ending 7 dpi.

NBD-Tre probe assays. Aag2 cells were grown to confluence in trehalose-containing cell culture media using the Nunc Lab-Tek Chamber Slide System (Thermo Fisher Scientific, MA). Cell culture media was removed and replaced with cell culture media spiked with 0.1 mM NBD-Tre probe with and without 1 mM ValA. Negative controls included wells treated with equimolar concentrations of the DMSO vehicle control. Treated cells were incubated for 4 hours, cell culture media was then removed, and cells were stained with the nuclear stain Hoescht (Thermo Fisher Scientific, MA). Cells were imaged immediately using a Leica DM5000 B, Leica DFC550 digital camera, and Leica CTR5000 control unit. Mean fluorescence intensity (MFI) per cell was quantified using ImageJ and included 10 separate images per condition and the average MFI of 5 cells per image. This experiment was repeated by treating Aag2 cells with 0.1 mM NBD-Tre probe in the presence of increasing concentrations of glucose or trehalose and on uninfected and DENV2-infected cells.

Nile Red assay. Aag2 cells were grown to confluence in either glucose or trehalose-containing cell culture media using the Nunc Lab-Tek Chamber Slide System. A 30 mM Nile Red (Cayman Chemical, MI) stock in DMSO was diluted 1:3000 in either glucose or trehalose-containing cell culture medium and incubated for 30 minutes at room temperature. Cells were fixed and counterstained with the nuclear stain Hoescht. Cells were imaged immediately using a Leica DM5000 B, Leica DFC550 digital camera, and Leica CTR5000 control unit. Corrected total cell fluorescence (CTCF) was quantified using integrated density, area, and mean background fluorescence data obtained from 3 separate images per condition and 20 cells per image.

RNA-Seq. RNA Sequencing (RNASeq) was conducted using a service offered by Novogene, USA (California UC Davis) on Aag2 cells that were either maintained in glucose or trehalose-containing cell culture media in triplicate. RNA was extracted using the Qiagen RNeasy Plus RNA Extraction kit with gDNA eliminator columns. RNA integrity was assessed using the RNA Nano 6000 Assay Kit of the Bioanalyzer 2100 system (Agilent Technologies, CA, USA). Total RNA was used as input material for the RNA sample preparations. Briefly, mRNA was purified from total RNA using poly-T oligo-attached magnetic beads. Fragmentation was carried out using divalent cations under elevated temperature in First Strand Synthesis Reaction Buffer(5X). First strand cDNA was synthesized using random hexamer primer and M-MuLV Reverse Transcriptase(RNase H-). Second strand cDNA synthesis was subsequently performed using DNA Polymerase I and RNase H. Remaining overhangs were converted into blunt ends via exonuclease/polymerase activities. After adenylation of 3' ends of DNA fragments, Adaptor with hairpin loop structure were ligated to prepare for hybridization. In order to select cDNA fragments of preferentially 370~420 bp in length, the library fragments were purified with AMPure XP system (Beckman Coulter, Beverly, USA). Then PCR was performed with Phusion High-Fidelity DNA polymerase, Universal PCR primers and Index (X) Primer. At last, PCR products were purified (AMPure XP system) and library quality was assessed on the Agilent Bioanalyzer 2100 system. The clustering of the index-coded samples was performed on a cBot Cluster Generation System using TruSeq PE Cluster Kit v3-cBot-HS (Illumia) according to the manufacturer's instructions. After cluster generation, the library preparations were sequenced on an Illumina Novaseq

platform and 150 bp paired-end reads were generated. Raw data (raw reads) of fastq format

were firstly processed through in-house perl scripts. In this step, clean data (clean reads)

were obtained by removing reads containing adapter, reads containing ploy-N and low-

quality reads from raw data. At the same time, Q20, Q30 and GC content the clean data

were calculated. All the downstream analyses were based on the clean data with high

quality. Reference genome and gene model annotation files were downloaded from

genome website directly. Index of the reference genome was built using Hisat2 v2.0.5 and

paired-end clean reads were aligned to the reference genome using Hisat2 v2.0.5. We

selected Hisat2 as the mapping tool for that Hisat2 can generate a database of splice

junctions based on the gene model annotation file and thus a better mapping result than

other non-splice mapping tools. featureCounts v1.5.0-p3 was used to count the reads

numbers mapped to each gene. FPKM of each gene was calculated based on the length of

the gene and reads count mapped to this gene. FPKM, expected number of Fragments Per

Kilobase of transcript sequence per Millions base pairs sequenced, considers the effect of

sequencing depth and gene length for the reads count at the same time, and is currently the

most commonly used method for estimating gene expression levels. Differential expression

analysis of the two conditions/groups was performed using the DESeq2 R package (1.20.0).

DESeq2 provide statistical routines for determining differential expression in digital gene

expression data using a model based on the negative binomial distribution. The resulting

P-values were adjusted using the Benjamini and Hochberg's approach for controlling the

false discovery rate. Genes with an adjusted P-value $< = 0.05$ found by DESeq2 were

assigned as differentially expressed. Functional annotations of differentially expressed

genes were performed using the g:Cost Functional Profiling tool at g:Profiler and NCBI's

DAVID Functional Annotation Bioinformatics tool. FASTQ files for each sample are

available in the NCBI SRA database (BioProject ID: PRJNA936836 and PRJNA882813).

Focus-forming unit assays. Aag2 cells were seeded into 96 well plates and grown to ~70%

confluence in glucose-containing cell culture media. Wells were then inoculated with 20

focus-forming units (FFUs) of DENV2 in either glucose or trehalose-containing cell

culture media for 1 hour at 27°C. Unbound virus was removed and fresh glucose-

containing cell culture media was added. At 3 dpi, confluent Aag2 monolayers were fixed

in 4% paraformaldehyde in phosphate buffered saline (PBS) for 10 minutes, cell

membranes were permeabilized with 0.1% Triton X-100, and DENV2 antigen was stained

with 1:200 anti-DENV2 envelope antibody (3H5-1, EMD Millipore) in PBS with 1%

bovine serum albumin (BSA). The total number of DENV-positive foci were revealed

using a 1:200 horseradish peroxidase-conjugated secondary antibody in PBS plus 1% BSA,

and an AEC Peroxidase Substrate kit (Vector Laboratories). This experiment was repeated

using cells that were maintained in either glucose or trehalose-containing cell culture media

throughout the experiment, and digital images were collected to quantify the total focus-

forming units/well and total number of DENV2-positive cells/well. FFU assays were also

performed using cell free supernatants isolated from DENV2-infected Aag2 cells that were

maintained in either glucose or trehalose-containing cell culture media, examining cell free

supernatants at 0, 1, and 3 dpi. Cell free supernatants were stored at -80°C until use and

then used to inoculate subconfluent Aag2 cell monolayers that were maintained in glucose-

containing cell culture media. FFU assays were also used to quantify the impact of ValA

treatment on DENV2 infection. For this experiment, Aag2 cells were inoculated with 40

FFU/well DENV2 and co-treated with 0, 0.1, 0.2, and 0.5 mg/mL ValA. Digital images

were captured using the Evos XL Core Imaging System.

RT-qPCR-based infectivity assays.

Aag2 cells were seeded into 96 well plates and grown to ~70% confluence in either glucose

or trehalose-containing cell culture media. Cells were then mock infected or inoculated

with 1×10^6 genomic equivalents of DENV2 in either glucose or trehalose-containing cell

culture media in triplicate. For the attachment assay, incubation of DENV2 with cells took

place at 4°C for 1 hr. and then unbound virus was removed, cells were washed with PBS,

and total RNA was extracted and quantified by RT-qPCR. DENV2 viral RNA (vRNA) was

normalized by total nanograms of cellular RNA. For the cell entry and replication assay,

incubation of DENV2 with cells took place at 25°C for 1 hr. and then unbound virus was

removed and replaced with either glucose or trehalose-containing cell culture media. Total

RNA was extracted 3 dpi, quantified by RT-qPCR, and normalized using total nanograms

of cellular RNA. DENV2 stocks were also generated using Aag2 cells maintained in either

glucose or trehalose-containing cell culture media, frozen at -80°C, and then RNA was

extracted and quantified using RT-qPCR. FFUs were also measured in these samples,

which allowed for determination of the specific infectivity of virions produced from

glucose and trehalose-fed cells. Specific infectivity was represented as FFU/vRNA.

Amplification of the viral target was performed using a singleplex format in 48-well plates

(Illumina) with a total reaction volume of 10 μl. Reverse transcription and quantitative PCR

were performed in the same closed tube with 50 ng of total RNA per reaction using the

Quantitect RT-qPCR Kit (Qiagen). All primers were used at final concentrations of 4 μM.

For amplification of DENV2 vRNA (nt 87–207), Forward: 5′ CAG ATC TCT GAT GAA

TAA CCA ACG 3′ and Reverse: 5′ CAT TCC AAG TGA GAA TCT CTT TGT CA 3′

DENV2-specific primers were used. The DENV2-specific probe, 5'/56-FAM/CTG CCG

ATC/ZEN/TTG CAC ATT ACC ACC GC/3IABkFQ/3′, was used for added specificity.

All RT-qPCR reactions were performed using an Eco Illumina and each time point was

measured in triplicate. Cycling conditions were 50 °C for 30 min (reverse transcription)

and 95 °C for 15 min, followed by 45 cycles of 94 °C for 15 s, 55 °C for 30 s and 72 °C

for 30 s. Relative quantities of target cDNA were determined using the Pfaffl method.

Student's t tests were performed between groups to determine statistical significance.

Results

Aag2 cells express a secreted trehalase enzyme that can convert trehalose to glucose.

RNA-Seq datasets that characterize gene expression of Aag2 cells reveal high

expression of trehalose-6-phosphate synthase (TPS) and trehalase (TREH) homologues

(42). Expression of a trehalose 6-phosphate phosphatase (TPP) homologue is low but not

absent (Figure 1A). The amino acid sequence of TREH was analyzed using SignalP 6.0

and indicated a signal peptide cleavage site between amino acid positions 30 and 31

(probability 0.97812) (Figure 1B). PSORT Wolf analysis of the TREH amino acid

sequence predicted that it is located in the endoplasmic reticulum, extracellular space,

peroxisome, and plasma. Together, these data indicated that Aag2 TREH is a secreted form

of the enzyme and exists in the extracellular environment.

Most cell culture media used to support insect cell growth contains glucose. We

reasoned that if Aag2 expressed TREH, and if this enzyme was secreted, it would convert

trehalose to glucose and similarly support cell growth. Aag2 cells were grown in both

glucose and trehalose-containing cell culture media and passaged at least two times before

experiments. Aag2 cells proliferated equally in glucose and trehalose-containing cell

culture media (Figure 2A-B). Importantly, over a 3-day time course Aag2 metabolized

glucose when maintained on glucose-containing cell culture media, and generated glucose

when maintained on trehalose-containing cell culture media (Figure 2C-D).

***In vitro* trehalase inhibitor assays show reduction in cell proliferation and reveal**

differences in inhibitor efficacy.

Previous *in vivo* research showed that treatment with trehalase inhibitor ValA at

0.2, 0.4, and 1.0 mM led to dose-dependent changes in *Ae. aegypti* development (23). Cell

proliferation assays were performed to determine if trehalase inhibitors interfered with

Aag2 cell metabolism when maintained in trehalose-containing cell culture media. There

was no reduction in cell proliferation when Aag2 cells maintained in glucose-containing

cell culture media were treated with 1 mM ValA (Figure 2E). However, when Aag2 cells

maintained in trehalose-containing cell culture media were treated with 1 mM ValA, a

significant reduction in cell proliferation was evident 3 days post-treatment, which

persisted throughout the time course (Figure 2E). To directly test the impact of trehalose

analogues on trehalase activity, a dose-dependence assay was performed that quantified

conversion of trehalose to glucose. ValA maintained ~75% activity as a trehalase inhibitor

at the lowest concentration tested (i.e., 0.008 mM) (Figure 2F). Although 5-ThioTre and 6-

TreNH$_2$ maintained ~75% activity at the highest concentration tested (i.e., 1.0 mM), they

had no activity at the two lowest concentrations (Figure 2F).

RNA-Seq analysis identifies differentially expressed genes involved in trehalose

metabolism and innate immunity.

Previous research confirmed that key enzymes involved in trehalose metabolism

are expressed in Aag2 cells (42). This was somewhat unexpected given that Aag2 cells do

not derive from fat body tissue. Very little is known about the response of non-fat body

cells and tissues to trehalose. RNA-Seq was performed to characterize the differentially

expressed genes in Aag2 cells maintained in glucose and trehalose-containing cell culture

media.

Significant differences in gene expression were observed. There were 157

upregulated and 118 downregulated genes in Aag2 cells that were maintained in trehalose-

containing cell culture media (Figure 3A). Although TREH expression did not change in

the RNA-Seq dataset (Supplementary Table 1) or in a follow-up RT-qPCR validation study

(Figure 3B), trehalose transporter 1 Tret1 (AAEL026825) was upregulated 2.7-fold in the

presence of trehalose (Supplementary Table 2). Functional annotations of differentially

expressed genes also showed significant changes in genes involved in composition of the

cell membrane, lipid biosynthesis, sugar conjugation, extracellular matrix, calcium

signaling, signal transduction, and mitochondrial function Figure 3C-D). Significant

changes in the expression of some innate immunity genes were also noted (Table 1).

Passive uptake of a fluorescently labeled trehalose probe.

Considering that Aag2 cells increase expression of Tret1 in response to trehalose while poorly expressing TPP, it is reasonable to expect that these cells can acquire trehalose from the extracellular environment (Supplementary Tables 1-2). To enable testing of whether Aag2 cells can accumulate extracellular trehalose, we synthesized a novel fluorescent trehalose derivative, NBD-Tre, which consists of trehalose conjugated to a fluorescent 2-(7-<u>n</u>itro-2,1,3-<u>b</u>enzoxa<u>d</u>iazol-4-yl) group (Figure 4A). Aag2 cells that were maintained in trehalose-containing cell culture media were treated with 1.0 mM NBD-Tre probe or a vehicle control for 4 hours, excess probe was removed, cells were washed in cell culture media, and then digital images were taken using a fluorescent microscope. 1.0 mM ValA was used to determine if trehalase inhibition would prevent probe degradation and improve cell labeling. Quantifying the mean fluorescent intensity/cell showed a statistically significant increase in signal when cells were treated with the NBD-Tre probe compared to DMSO vehicle control. The addition of trehalase inhibitor ValA did not influence cell labeling (Figure 4B) suggesting that trehalase does not degrade the fluorescent molecule.

To test if trehalose entered into Aag2 cells using a passive/facilitated process, 0.1 mM NBD-Tre probe was suspended in cell culture media with a range of glucose and trehalose concentrations: 0.0, 0.1, 1.0, 10.0, and 25.0 mM. A lower concentration of NBD-Tre probe was used in this experiment because the labeling efficiency of 0.1 mM NBD-Tre in sugar-free cell culture media was equivalent to 1.0 mM NBD-Tre in high glucose or trehalose cell culture media. The mean fluorescent intensity per cell decreased ~10-20% at

the highest two concentrations of glucose (Figures 4C,E). In contrast, the mean fluorescent

intensity per cell increased at the lowest concentration of trehalose and decreased 40-75%

at the three highest concentrations of trehalose (Figures 4D,F). These results support that

the trehalose probe entered Aag2 cells through a process of facilitated diffusion, which is

consistent with the function of known trehalose transporters such as Tret1 (43).

Maintaining Aag2 cells in trehalose-containing cell culture media increases DENV2

shedding and specific infectivity.

Maintaining Aag2 cells in trehalose-containing cell culture media led to significant

changes in gene expression, and trehalose used a facilitated/passive process to accumulate

inside of these cells. Transcriptomic changes, which include differential gene expression

of key innate immunity genes (Table 1), might influence virus infection. Similarly, an

influx of trehalose into intracellular compartments may influence different aspects of the

virus life cycle. Dengue virus type 2 (DENV2) was used as a model vector-borne disease

to test the hypothesis that trehalose can modify virus infection in insect cells.

Aag2 cells maintained in glucose-containing cell culture media were inoculated

with 20 focus-forming units (FFUs) of DENV2 and the infection was allowed to spread

throughout the monolayer for 5 days. Cells were also mock-infected as a negative control.

Infected and mock infected cells were then treated with either no glucose cell culture media

as a vehicle control or with 1 mM NBD-Tre probe for 4 hours. Aag2 cells were then fixed

and stained using an anti-DENV2 3H5.1 Mab and counterstained with Hoescht.

Immunofluorescence microscopy revealed diffuse cytoplasmic localization of the NBD-

Tre probe and discrete localization of DENV2 envelope (E) protein outside of the nucleus

(Figure 5). There was no clear co-localization between these molecules.

Although the immunofluorescence analysis did not reveal evidence of specific co-

localization between trehalose and DENV2 E, the diffuse localization of trehalose Aag2

may provide opportunities for interaction with other DENV2 proteins or to influence virus

replication. We tested this hypothesis using Aag2 cells maintained in glucose-containing

cell culture media, which were inoculated with 20 focus-forming units (FFUs) of DENV2

in either glucose or trehalose-containing cell culture media for 1 hour. Unbound virus and

cell culture media was removed and replaced with glucose-containing cell culture media.

Three days later cell monolayers were fixed and stained with anti-DENV2 MAb and the

total number of FFUs per well were quantified. Pre-treatment with either glucose or

trehalose-containing cell culture media had no impact on DENV2 infection (Figure 6A).

In contrast, when the same experiment was performed using Aag2 cells that were

maintained in either glucose or trehalose-containing cell culture media for the duration of

the focus forming unit assay, there was a 2-fold increase in FFUs per well in the trehalose-

maintained cells (Figures 6B,D). The total number of DENV2 positive cells were

quantified in each well, which also revealed a 2-fold increase in the trehalose-maintained

cells (Figures 6C,D).

To determine the stage of the virus life cycle that was affected by trehalose, an

attachment assay was performed using Aag2 cells that were maintained in either glucose

or trehalose-containing cell culture media. Cells were inoculated with 20 FFUs DENV2 in

glucose-containing cell culture media at 4 °C for 1 hour, unbound virus was removed, cells

were washed with cell culture media, and total RNA was extracted from cell monolayers.

RT-qPCR analysis showed no difference in the attachment of DENV2 to Aag2 cells

maintained in either glucose or trehalose-containing cell culture media (Figure 6E). A cell

entry assay was then performed, which allowed DENV2 to attach and enter Aag2 cells at

37 °C for 4 hours. RT-qPCR analysis showed a statistically significant increase in cell entry

of DENV2 inside of Aag2 cells that were maintained in trehalose-containing cell culture

media (Figure 6F). Supernatants were then collected from DENV2-infected Aag2 cells that

were maintained in either glucose or trehalose-containing cell culture media and stored as

viral stocks. Viral stocks were tested to determine the total FFUs/well in a 5 μL inoculum

onto Aag2 cells that were maintained in glucose-containing cell culture media. Consistent

with previous experiments, significantly more DENV2 was shed from Aag2 cells that were

maintained in trehalose-containing cell culture media by day 3 post-infection (dpi) (Figure

6G). To test the quality of virions produced by Aag2 cells maintained in either glucose or

trehalose-containing cell culture media, virus stocks were prepared from each condition at

5 dpi, and then FFU and RT-qPCR assays were performed. The total FFUs per mL were

normalized by vRNA (FFU/vRNA) to determine specific infectivity of virus stocks. The

specific infectivity of DENV2 produced from Aag2 cells maintained in trehalose-

containing cell culture media was 2-fold greater (Figure 6H).

Trehalose promotes accumulation of intracellular lipids.

Previous research suggests that trehalose induces autophagy in mammalian cells,

although autophagy's role in virus infection is varied (32-36, 39, 44). The consensus is that

DENV benefits from autophagy in mammalian cells by enhancing cell survival and

promoting virus maturation (45-49). The role of autophagy in arbovirus infection in insect

cells is less clear, and conventional autophagy inducers and inhibitors may respond

differently in insect cells (44, 50). Trehalose can also be converted to glucose and used as

a substrate for lipogenesis (42).

RNA-Seq analysis revealed significant changes in gene expression related to

integral/intrinsic membrane components, and the main function of autophagy is to

accumulate and utilize lipids to enhance cell survival. We used Nile Red staining to

determine if trehalose modified the lipid environment of Aag2 cells. We identified a slight,

yet significant increase in intracellular lipid accumulation in Aag2 cells that were

maintained in trehalose-containing cell culture media (Figure 7A-B).

DENV2 infection does not impact trehalose metabolism.

Previous research has shown that DENV manipulates the metabolism of its host

cell to promote replication. Different parameters of trehalose metabolism were tested

during DENV2 infection to determine if this pathway is manipulated in Aag2 cells. Aag2

cells maintained in trehalose-containing cell culture media were either mock infected or

infected with DENV2 and conversion of trehalose to glucose was assessed on 1, 3, 5, and

7 dpi. DENV2 infection did not influence trehalase activity (Figure 8A). The impact of

DENV2 infection on trehalose uptake was also tested by infecting Aag2 cells that were

maintained in glucose-containing cell culture media with 0, 1, 10, 100, and 500 FFUs and

then labeling cells with 0.1 mM NBD-Tre 3 dpi. There was no difference in mean

fluorescence intensity per cell, therefore DENV2 infection did not influence trehalose

uptake (Figure 8B).

Trehalase inhibition reduces cell viability and DENV infection in trehalose-dependent

Aag2 cells.

We previously showed that trehalase inhibition reduced cell proliferation when

cells were maintained in trehalose-containing cell culture media (Figure 2E). We

hypothesized that this activity would reduce DENV productivity. As expected, ValA

treatment did not reduce cell viability or DENV productivity in Aag2 cells that were

maintained in glucose-containing cell culture media, and surprisingly increased DENV

productivity by 50% (Figure 9A-C). In contrast, ValA treatment significantly reduced cell

viability and DENV productivity in Aag2 cells that were maintained in trehalose-

containing cell culture media (Figure 9D-F). It is important to note that a FFU assay was

not performed directly on Aag2 cells that were maintained on trehalose-containing cell

culture media because the addition of ValA prevents establishment of a monolayer, which

is required for the assay. Instead, cell-free supernatants were harvested from ValA-treated

cells and used for downstream FFU assays performed on Aag2 cells that were maintained

on glucose-containing cell culture media.

Discussion

Control of *Aedes* spp. disease vectors is an important area of ongoing research due

to concerns regarding vaccine limitations, pesticide resistance, and increased geographic

spread (6). Trehalase inhibitors have been previously characterized as a novel class of

putative insecticides, potentially effective for controlling *Aedes* spp. disease vectors due to

the vital importance of trehalose as a blood sugar during mosquito development (14, 15,

17, 18, 20, 24, 25, 30, 40). Our previous research has revealed that the commercially

available trehalase inhibitor ValA negatively impacts mosquito development *in vivo* when

applied directly to the larval environment by showing dose-dependent activity delaying

egg hatch, pupation, eclosure, and flight of adult *Ae. Aegypti* mosquitoes (23).

Here, we describe the use of an *in vitro* model to screen for trehalase inhibitors in

Ae. aegypti and determine the impact of trehalose metabolism on a mosquito-borne virus.

ValA and novel trehalase inhibitors (5-ThioTre, 6-TreNH$_2$) prevented conversion of cell

culture media trehalose to glucose. ValA had the strongest activity, which significantly

reduced cell proliferation. Differences in the activity of trehalose analogues seen *in vitro*

and *in vivo* suggest that there are opportunities to optimize trehalase inhibitors for target

species (23). Our *in vitro* assay is a platform that could be used for high throughput studies

to develop targeted trehalose analogues.

RNA-Seq analysis showed that Tret1 expression was induced in cells that were

maintained in trehalose-containing cell culture media. The Aag2 cell line was originally

generated from whole homogenized embryos. Cells within the culture exhibit differing

morphologies, and it has been suggested that the varying morphologies of mosquito cells

in culture may be indicative of the presence of a diversity of embryonic and differentiated

cell types (51). Aag2 cells were not derived from fat body cells or tissue, which is where

Tret1 is concentrated *in vivo* (43). The role of Tret1 in peripheral cells and tissues is

unknown. Considering that the machinery for trehalose synthesis was lacking in Aag2

cells, we hypothesized that these cells may use Tret1 for trehalose uptake rather than

excretion. Previous research has shown that Tret1 is a facilitated/passive transporter (43).

We confirmed that Aag2 cells could accumulate a novel fluorescent trehalose probe, NBD-

Tre, and that uptake was prevented if unlabeled trehalose was present. This supports

facilitated/passive uptake of trehalose in Aag2 cells.

Entry of trehalose into Aag2 cells may have led to differential gene expression as

noted in the RNA-Seq dataset, although the presence of trehalose also induced expression

of integrin proteins, which mediate signaling from the cell surface. Significant differences

in gene expression were present in genes involved in composition of the cell membrane,

lipid biosynthesis, sugar conjugation, extracellular matrix, calcium signaling, signal

transduction, and mitochondrial function. These global changes in gene expression may

explain why the physiological role of trehalose extends beyond a simple source of fuel.

Trehalase inhibition can impact a wide array of physiological functions in insects including

chitin synthesis, stress protection, larval and pupal development (14, 17-20, 24-26, 28).

Considering that trehalose enters Aag2 cells and leads to significant changes in

gene expression, we hypothesized that the environment of a trehalose-fed insect cell was

different enough to influence virus infection. This was supported by previous data

suggesting that trehalose can induce autophagy and that DENV2 benefits from autophagy

by prolonging cell survival and enhancing virus maturation (45-49). We found that Aag2

cells that were maintained in trehalose-containing cell culture media promoted DENV2

cell entry, which led to a higher number of infected cells, and increased virus shedding. In

mammalian cells, trehalose is an inducer of autophagy and can block autophagic flux from

autophagosomes to autolysosomes (39). DENV2 also induces the autophagy pathway, and

infection is reduced when autophagy is inhibited (45-49). It is possible that trehalose

increases DENV2 infection in Aag2 cells by promoting autophagy and enhancing virus

cell entry through modification of cell membranes. Previous research suggests that the

relationship between autophagy and DENV infection in Aag2 cells is complicated and that

conventional autophagy inhibitors may not work as expected in these cells (44). Instead,

we used Nile Red staining as a proxy to quantify changes to the intracellular lipid

environment. Nile Red staining revealed an increase in intracellular lipids in Aag2 cells

maintained on trehalose-containing cell culture media. It is important to note that trehalose

is the main blood sugar in insects, and it may not have the same influence on autophagy as

seen in mammalian cells.

Trehalose can also be converted to glucose, which is a substrate for lipogenesis. It

is possible that trehalose stimulates lipogenesis and the production of triglycerides and

associated fatty acids that promote virus replication and synthesis of more infectious viral

particles. RNA-Seq analysis showed that trehalose-fed cells had a 55% downregulation of

phosphoenolpyruvate carboxykinase (i.e., AAEL000006) and a 20% upregulation of fatty

acid synthase (i.e., AAEL001194) (Supplementary Table 1). Further, one of the most

upregulated genes (714%) was elongation of very long chain fatty acids protein 4

(AAEL004947). Together, these changes in gene expression may have been responsible

for the increase in intracellular lipids and perhaps the increase in virus replication and

shedding.

The specific infectivity of DENV2 also increased when virus stocks were produced

from Aag2 cells that were maintained in trehalose-containing cell culture media. The

mechanism of action that led to increased specific infectivity is unknown, although

trehalose is widely known for its ability to prevent heat and cold shock and it is used as a

cryoprotectant (52-58). Trehalose may provide some membrane or protein stabilizing

activity that either allows virus production and shedding to occur more efficiently or it

simply allows more virus to survive a freeze-thaw event. Autophagy has also been linked

to DENV maturation, which suggests that trehalose may facilitate production of more

mature virions (49). It is important to note that the trehalose concentration used in our cell

culture media is 20-50-fold less than what is typically used for cryoprotection (52-54, 59-

61).

Trehalase inhibition interferes with multiple physiological pathways in insects, and

we found that ValA reduced cell proliferation in Aag2 cells that were maintained in

trehalose-containing cell culture media. It was reasonable to hypothesize that trehalase

inhibition would also reduce DENV productivity. We showed that 1.0 mM ValA causes a

90% reduction in cell viability 5 days post-treatment in Aag2 cells that were maintained in

trehalose-containing cell culture media. The reduction in cell viability correlated with an

approximately 75% decrease in DENV vRNA shedding. *In vivo* studies show deficits in

Ae. aegypti egg hatch rate, larval development, pupation, and eclosure (23). Considering

that vertical transmission is less important for the maintenance and transmission of DENV

in a natural environment, trehalase inhihibition in larval habitats would likely not impact

DENV in a meaningful way. It is unclear how trehalase inhibition impacts adult *Ae. aegypti*

at this time, although these data suggest that targeting adults could simultaneously reduce

mosquito and virus fitness.

This manuscript describes an *in vitro* model that can be used to rapidly screen novel

trehalase inhibitors and probes and underscores the importance of trehalose metabolism in

Ae. aegypti physiology and the transmission of a mosquito-borne virus. These data

facilitate the future development of targeted trehalase analogues that will reduce

populations of disease vectors and related viruses with minimal off-target effects.

Acknowledgements

This work was supported through start-up funds from Central Michigan University College

of Medicine. BMS was supported by a grant from the National Institutes of Health

(R15AI117670).

Figure legends

Figure 1. Trehalose metabolism genes expressed in Aag2 cells. (A) Trehalose pathway

showing *Ae. aegypti* enzymes involved in synthesis: TPS; AAEL006446 and TPP;

AAEL010684 and catabolism: TREH; AAEL009658. (B) SignalP 6.0 prediction of a

signal peptide in TREH; AAEL009658.

Figure 2. Trehalose supports the growth of Aag2 cells. (A) Representative image of
Aag2 cells at 20X magnification maintained in either glucose (GLU) or trehalose (TRE)-
containing cell culture media. (B) Cell proliferation assay of Aag2 cells maintained in
either glucose (circles) or trehalose (squares)-containing cell culture media was performed
for 7 days (D0-D7). (C) Glucose assay of Aag2 cell supernatant at days 0, 1, 2, and 3 post-
feeding (D0-D3) with glucose-containing cell culture media. (D) Glucose assay of Aag2
cell supernatant at days 0, 1, 2, and 3 (D0-D3) post-feeding with trehalose-containing cell
culture media. (E) Cell proliferation assay of Aag2 cells grown in either glucose-containing
cell culture media with (square) or without (circle) 1.0 mM ValA or trehalose-containing
cell culture media with (upside down triangle) or without (triangle) 1.0 mM ValA. (F)
Glucose assay of Aag2 cell supernatants from cells maintained in trehalose-containing cell
culture media and treated with 0.008-1 mM ValA, 5-ThioTre, and 6-TreNH2. Assays were
performed in triplicate. Unpaired Student's t tests were performed between groups to assess
statistical significance. Standard deviations are shown.

Figure 3. RNA-Seq analysis and differentially expressed genes. (A) Volcano plot of
upregulated (157) and downregulated (118) differentially expressed genes present in Aag2
cells that were maintained in trehalose-containing cell culture media. (B) RT-qPCR
analysis of TREH expression in Aag2 cells that were maintained in either glucose (G) or
trehalose (T)-containing cell culture media on 0-3 dpt (D0-D3). (C) Functional annotation
categories present in upregulated genes. (D) Functional annotation categories in
downregulated genes.

Figure 4. NBD-Tre probe uptake in Aag2 cells. (A) Chemical structure of fluorescently labeled trehalose probe NBD-Tre. (B) Aag2 cells were treated with either vehicle control (DMSO), 1.0 mM NBD-Tre probe, or with 1.0 mM NBD-Tre probe and 1.0 mM ValA in glucose free cell culture media for 4 hours and then mean fluorescence intensity/cell was determined. (C-D) Aag2 cells were treated with 1.0 mM NBD-Tre probe in the presence of (C) 0, 0.1, 1, 10, and 25 mM glucose and (D) 0, 0.1, 1, 10, and 25 mM trehalose for 4 hours and then mean fluorescence intensity/cell was determined. (E-F) Representative images of Aag2 cells that were treated with 1.0 mM NBD-Tre probe in the presence of (E) 0, 0.1, 1, 10, and 25 mM glucose and (F) 0, 0.1, 1, 10, and 25 mM trehalose with a Hoescht nuclear counterstain. Assays were performed in triplicate. Unpaired Student's t tests were performed between groups to assess statistical significance. Standard deviations are shown.

Figure 5. Co-localization of NBD-Tre and DENV2 in Aag2 cells. Aag2 cells were uninfected (Mock) or infected with DENV2 for 5 days and then treated with no glucose cell culture media alone (Vehicle) or 1 mM NBD-Tre probe for 4 hours. Aag2 cells were then fixed and stained with anti-DENV2 Mab and digital images were obtained at 20X magnification. Hoescht nuclear stain was added to each condition.

Figure 6. Trehalose promotes DENV2 cell entry and virus shedding. Aag2 cells were Mock infected or infected with DENV in either glucose (GLU) or trehalose (TRE)-containing cell culture media. (A) Trehalose-containing cell culture media was only present during inoculation and was then exchanged for glucose-containing cell culture media and

focus-forming units were quantified 3 dpi. (B) Trehalose-containing cell culture media was

present during inoculation and for the three days prior to a focus-forming unit assay. (C)

The total number of DENV2 positive cells per well were quantified from (B). (D)

Representative images from (B) showing distinct foci are shown. (E) An attachment assay

was performed by incubating DENV2 with Aag2 cells at 4°C for 1 hour. Unbound virus

was removed, cells were washed with PBS, and total RNA was extracted and quantified by

RT-qPCR. DENV2 viral RNA (vRNA) was normalized by total nanograms of cellular

RNA (F) A cell entry/replication assay was performed by incubating DENV2 with Aag2

cells at 25°C for 1 hour. Unbound virus was removed and replaced with either glucose or

trehalose-containing cell culture media. Total RNA was extracted 3 dpi, quantified by RT-

qPCR, an DENV2 vRNA was normalized using total nanograms of cellular RNA. (G) A

virus shedding assay was performed by collecting cell free supernatants at 0, 1, and 3 dpi

from Aag2 cells that were inoculated with DENV2 in either glucose or trehalose-containing

cell culture media. Cell free supernatants were used to inoculate monolayers of Aag2 cells

grown in glucose-containing cell culture media and focus forming unit assays were used to

quantify the infectious units present in each sample. (H) DENV2 stocks were generated

from Aag2 cells that were grown in either glucose or trehalose-containing cell culture

media. Focus forming unit assays were performed to quantify the infectivity of each stock.

Total RNA was also extracted from virus stocks and DENV2 vRNA was quantified using

RT-qPCR. Specific infectivity was derived by normalizing the total number of focus

forming units in a sample by its vRNA content (i.e., FFU/vRNA). Assays were performed

in triplicate. Unpaired Student's t tests were performed between groups to assess statistical

significance. Standard deviations are shown.

Figure 7. Trehalose promotes intracellular lipid accumulation in Aag2 cells. (A) Intracellular lipids were labeled in Aag2 cells maintained in either glucose (GLU) or trehalose (TRE) cell culture media using either vehicle control (DMSO) or Nile Red. Cells were fixed and digital images were obtained at 20X magnification. Hoescht nuclear stain was added to each condition. (B) The corrected total cell fluorescence (CTCF) was quantified using the integrated density of 60 individual cells per condition. Assays were performed in triplicate. Unpaired Student's t tests were performed between groups to assess statistical significance. Standard deviations are shown.

Figure 8. DENV2 infection does not influence trehalase activity or trehalose uptake. (A) A glucose assay was performed on cell-free supernatants taken from Aag2 cells maintained in trehalose-containing cell culture media that were either uninfected (Mock) or infected with DENV2 for 1, 3, 5, and 7 dpi. (B) A trehalose probe uptake assay was performed on Aag2 cells that were either uninfected (Mock) or infected with 1, 10, 100, and 500 FFUs of DENV2 5 dpi. 1.0 mM NBD-Tre probe was added to cells in glucose free cell culture media for 4 hours. Unbound probe was removed, and then digital images were obtained at 20X magnification. Mean fluorescence intensity was determined for 50 individual cells per condition. Assays were performed in triplicate. Unpaired Student's t tests were performed between groups to assess statistical significance. Standard deviations are shown.

Figure 9. Validamycin A reduces cell viability and DENV2 productivity when Aag2 cells are maintained in trehalose-containing cell culture media. (A) Subconfluent Aag2 cells maintained in glucose-containing cell culture media were treated with 0, 0.2, 0.4, and 1.0 mM ValA for 5 days and then trypan blue exclusion assays were performed to quantify percent cell viability. (B) Subconfluent Aag2 cells maintained in glucose-containing cell culture media were inoculated with 20 FFUs of DENV for 1 hour, followed by treatment with 0, 0.2, 0.4, and 1.0 mM ValA for 5 days. vRNA was extracted from equivalent volumes of cell free supernatants and quantified by RT-qPCR. (C) Subconfluent Aag2 cells maintained in glucose-containing cell culture media were inoculated with 20 FFUs of DENV for 1 hour, followed by treatment with 0, 0.2, 0.4, and 1.0 mM ValA for 3 days. FFU assays were performed to quantify the total FFUs present in each well. (D) Subconfluent Aag2 cells maintained in trehalose-containing cell culture media were treated with 0, 0.2, 0.4, and 1.0 mM ValA for 5 days and then trypan blue exclusion assays were performed to quantify percent cell viability. (E) Subconfluent Aag2 cells maintained in trehalose-containing cell culture media were inoculated with 20 FFUs of DENV for 1 hour, followed by treatment with 0, 0.2, 0.4, and 1.0 mM ValA for 5 days. vRNA was extracted from equivalent volumes of cell free supernatants and quantified by RT-qPCR. (F) Subconfluent Aag2 cells maintained in trehalose-containing cell culture media were inoculated with 20 FFUs of DENV for 1 hour, followed by treatment with 0, 0.2, 0.4, and 1.0 mM ValA for 3 days. Cell free supernatants were harvested and used for a downstream FFU assay performed on Aag2 cells maintained on glucose-containing cell culture media, which quantified the total FFUs present in each well. Assays were performed in triplicate.

Unpaired Student's t tests were performed between groups to assess statistical significance.

Standard deviations are shown.

Supplementary Table 1. Complete readcount and FPKM list of all genes in GLU vs

TRE Aag2 cells.

Supplementary Table 2. Complete readcount and FPKM list of upregulated genes in

GLU vs TRE Aag2 cells.

Supplementary Table 3. Complete readcount and FPKM list of all downregulated in

GLU vs TRE Aag2 cells.

References

1.   Paul LM, Carlin ER, Jenkins MM, Tan AL, Barcellona CM, Nicholson CO, et al.
Dengue virus antibodies enhance Zika virus infection. Clin Transl Immunology.
2016;5(12):e117.
2.   Masel J, McCracken MK, Gleeson T, Morrison B, Rutherford G, Imrie A, et al.
Does prior dengue virus exposure worsen clinical outcomes of Zika virus infection? A
systematic review, pooled analysis and lessons learned. PLoS Negl Trop Dis.
2019;13(1):e0007060.
3.   Guzman MG, Alvarez M, Halstead SB. Secondary infection as a risk factor for
dengue hemorrhagic fever/dengue shock syndrome: an historical perspective and role of
antibody-dependent enhancement of infection. Arch Virol. 2013;158(7):1445-59.
4.   Obi JO, Gutiérrez-Barbosa H, Chua JV, Deredge DJ. Current Trends and
Limitations in Dengue Antiviral Research. Trop Med Infect Dis. 2021;6(4).
5.   Dighe SN, Ekwudu O, Dua K, Chellappan DK, Katavic PL, Collet TA. Recent
update on anti-dengue drug discovery. Eur J Med Chem. 2019;176:431-55.
6.   Conway MJ, Haslitt DP, Swarts BM. Targeting Aedes aegypti Metabolism with
Next-Generation Insecticides. Viruses. 2023;15(2).
7.   Rose RI. Pesticides and public health: integrated methods of mosquito
management. Emerg Infect Dis. 2001;7(1):17-23.

8.   Moyes CL, Vontas J, Martins AJ, Ng LC, Koou SY, Dusfour I, et al. Contemporary
status of insecticide resistance in the major Aedes vectors of arboviruses infecting humans.
PLoS Negl Trop Dis. 2017;11(7):e0005625.
9.   Zulfa R, Lo WC, Cheng PC, Martini M, Chuang TW. Updating the Insecticide
Resistance Status of Aedes aegypti and Aedes albopictus in Asia: A Systematic Review
and Meta-Analysis. Trop Med Infect Dis. 2022;7(10).
10.   Roberts DR, Andre RG. Insecticide resistance issues in vector-borne disease
control. Am J Trop Med Hyg. 1994;50(6 Suppl):21-34.
11.   Kasai S, Itokawa K, Uemura N, Takaoka A, Furutani S, Maekawa Y, et al.
Discovery of super-insecticide-resistant dengue mosquitoes in Asia: Threats of
concomitant knockdown resistance mutations. Sci Adv. 2022;8(51):eabq7345.
12.   Du Y, Nomura Y, Zhorov BS, Dong K. Sodium Channel Mutations and Pyrethroid
Resistance in Aedes aegypti. Insects. 2016;7(4).
13.   Amelia-Yap ZH, Chen CD, Sofian-Azirun M, Low VL. Pyrethroid resistance in the
dengue vector Aedes aegypti in Southeast Asia: present situation and prospects for
management. Parasit Vectors. 2018;11(1):332.
14.   Zhang L, Qiu LY, Yang HL, Wang HJ, Zhou M, Wang SG, et al. Study on the
Effect of Wing Bud Chitin Metabolism and Its Developmental Network Genes in the
Brown Planthopper, Nilaparvata lugens, by Knockdown of TRE Gene. Front Physiol.
2017;8:750.
15.   Xia Y, Clarkson JM, Charnley AK. Trehalose-hydrolysing enzymes of
Metarhizium anisopliae and their role in pathogenesis of the tobacco hornworm, Manduca
sexta. J Invertebr Pathol. 2002;80(3):139-47.
16.   Wyatt GR, Kale GF. The chemistry of insect hemolymph. II. Trehalose and other
carbohydrates. J Gen Physiol. 1957;40(6):833-47.
17.   Wegener G, Macho C, Schloder P, Kamp G, Ando O. Long-term effects of the
trehalase inhibitor trehazolin on trehalase activity in locust flight muscle. J Exp Biol.
2010;213(Pt 22):3852-7.
18.   Wegener G, Tschiedel V, Schloder P, Ando O. The toxic and lethal effects of the
trehalase inhibitor trehazolin in locusts are caused by hypoglycaemia. J Exp Biol.
2003;206(Pt 7):1233-40.
19.   Thorat LJ, Gaikwad SM, Nath BB. Trehalose as an indicator of desiccation stress
in Drosophila melanogaster larvae: a potential marker of anhydrobiosis. Biochem Biophys
Res Commun. 2012;419(4):638-42.
20.   Tatun N, Wangsantitham O, Tungjitwitayakul J, Sakurai S. Trehalase activity in
fungus-growing termite, Odontotermes feae (Isoptera: Termitideae) and inhibitory effect
of validamycin. J Econ Entomol. 2014;107(3):1224-32.
21.   Tang B, Yang M, Shen Q, Xu Y, Wang H, Wang S. Suppressing the activity of
trehalase with validamycin disrupts the trehalose and chitin biosynthesis pathways in the
rice brown planthopper, Nilaparvata lugens. Pestic Biochem Physiol. 2017;137:81-90.
22.   Shukla E, Thorat LJ, Nath BB, Gaikwad SM. Insect trehalase: physiological
significance and potential applications. Glycobiology. 2015;25(4):357-67.
23.   Marten AD, Stothard AI, Kalera K, Swarts BM, Conway MJ. Validamycin A
Delays Development and Prevents Flight in Aedes aegypti (Diptera: Culicidae). J Med
Entomol. 2020;57(4):1096-103.

24.   Katagiri N, Ando O, Yamashita O. Reduction of glycogen in eggs of the silkworm,
Bombyx mori, by use of a trehalase inhibitor, trehazolin, and diapause induction in
glycogen-reduced eggs. J Insect Physiol. 1998;44(12):1205-12.
25.   Liebl M, Nelius V, Kamp G, Ando O, Wegener G. Fate and effects of the trehalase
inhibitor trehazolin in the migratory locust (Locusta migratoria). J Insect Physiol.
2010;56(6):567-74.
26.   Chen J, Zhang D, Yao Q, Zhang J, Dong X, Tian H, et al. Feeding-based RNA
interference of a trehalose phosphate synthase gene in the brown planthopper, Nilaparvata
lugens. Insect Mol Biol. 2010;19(6):777-86.
27.   Chen J, Tang B, Chen H, Yao Q, Huang X, Zhang D, et al. Different functions of
the insect soluble and membrane-bound trehalase genes in chitin biosynthesis revealed by
RNA interference. PLoS One. 2010;5(4):e10133.
28.   Asano N, Yamaguchi T, Kameda Y, Matsui K. Effect of validamycins on
glycohydrolases of Rhizoctonia solani. J Antibiot (Tokyo). 1987;40(4):526-32.
29.   Murphy TA, Wyatt GR. ENZYMATIC REGULATION OF TREHALOSE AND
GLYCOGEN SYNTHESIS IN THE FAT BODY OF AN INSECT. Nature.
1964;202:1112-3.
30.   Becker A, Schloder P, Steele JE, Wegener G. The regulation of trehalose
metabolism in insects. Experientia. 1996;52(5):433-9.
31.   Arguelles JC. Why can't vertebrates synthesize trehalose? J Mol Evol. 2014;79(3-
4):111-6.
32.   Yuan S, Zhang ZW, Li ZL. Trehalose May Decrease the Transmission of Zika
Virus to the Fetus by Activating Degradative Autophagy. Front Cell Infect Microbiol.
2017;7:402.
33.   Wu Q, Jiang D, Huang C, van Dyk LF, Li L, Chu HW. Trehalose-mediated
autophagy impairs the anti-viral function of human primary airway epithelial cells. PLoS
One. 2015;10(4):e0124524.
34.   Clark AE, Zhu Z, Krach F, Rich JN, Yeo GW, Spector DH. Zika virus is transmitted
in neural progenitor cells via cell-to-cell spread and infection is inhibited by the autophagy
inducer trehalose. J Virol. 2021;95(5).
35.   Clark AE, Sabalza M, Gordts P, Spector DH. Human Cytomegalovirus Replication
Is Inhibited by the Autophagy-Inducing Compounds Trehalose and SMER28 through
Distinctively Different Mechanisms. J Virol. 2018;92(6).
36.   Belzile JP, Sabalza M, Craig M, Clark AE, Morello CS, Spector DH. Trehalose, an
mTOR-Independent Inducer of Autophagy, Inhibits Human Cytomegalovirus Infection in
Multiple Cell Types. J Virol. 2016;90(3):1259-77.
37.   Rodríguez-Navarro JA, Rodríguez L, Casarejos MJ, Solano RM, Gómez A,
Perucho J, et al. Trehalose ameliorates dopaminergic and tau pathology in parkin
deleted/tau overexpressing mice through autophagy activation. Neurobiol Dis.
2010;39(3):423-38.
38.   Perucho J, Casarejos MJ, Gomez A, Solano RM, de Yébenes JG, Mena MA.
Trehalose protects from aggravation of amyloid pathology induced by isoflurane
anesthesia in APP(swe) mutant mice. Curr Alzheimer Res. 2012;9(3):334-43.
39.   Zhang X, Chen S, Song L, Tang Y, Shen Y, Jia L, et al. MTOR-independent,
autophagic enhancer trehalose prolongs motor neuron survival and ameliorates the

autophagic flux defect in a mouse model of amyotrophic lateral sclerosis. Autophagy.
2014;10(4):588-602.
40. Danielson ND, Collins J, Stothard AI, Dong QQ, Kalera K, Woodruff PJ, et al.
Degradation-resistant trehalose analogues block utilization of trehalose by hypervirulent
Clostridioides difficile. Chem Commun (Camb). 2019;55(34):5009-12.
41. Kalera K, Liu R, Lim J, Pathirage R, Swanson DH, Johnson UG, et al. Targeting
Mycobacterium tuberculosis Persistence through Inhibition of the Trehalose Catalytic
Shift. ACS Infect Dis. 2024;10(4):1391-404.
42. Marten AD, Tift CT, Tree MO, Bakke J, Conway MJ. Chronic depletion of
vertebrate lipids in A*edes aegypti* cells dysregulates lipid metabolism
and inhibits innate immunity without altering dengue
infectivity. In Press ed. PLoS Neglected Tropical Diseases2022.
43. Kikawada T, Saito A, Kanamori Y, Nakahara Y, Iwata K, Tanaka D, et al.
Trehalose transporter 1, a facilitated and high-capacity trehalose transporter, allows
exogenous trehalose uptake into cells. Proc Natl Acad Sci U S A. 2007;104(28):11585-90.
44. Brackney DE, Correa MA, Cozens DW. The impact of autophagy on arbovirus
infection of mosquito cells. PLoS Negl Trop Dis. 2020;14(5):e0007754.
45. Heaton NS, Perera R, Berger KL, Khadka S, Lacount DJ, Kuhn RJ, et al. Dengue
virus nonstructural protein 3 redistributes fatty acid synthase to sites of viral replication
and increases cellular fatty acid synthesis. Proc Natl Acad Sci U S A. 2010;107(40):17345-
50.
46. Lee YR, Lei HY, Liu MT, Wang JR, Chen SH, Jiang-Shieh YF, et al. Autophagic
machinery activated by dengue virus enhances virus replication. Virology.
2008;374(2):240-8.
47. Heaton NS, Randall G. Dengue virus-induced autophagy regulates lipid
metabolism. Cell Host Microbe. 2010;8(5):422-32.
48. McLean JE, Wudzinska A, Datan E, Quaglino D, Zakeri Z. Flavivirus NS4A-
induced autophagy protects cells against death and enhances virus replication. J Biol Chem.
2011;286(25):22147-59.
49. Mateo R, Nagamine CM, Spagnolo J, Méndez E, Rahe M, Gale M, Jr., et al.
Inhibition of cellular autophagy deranges dengue virion maturation. J Virol.
2013;87(3):1312-21.
50. Chen TY, Smartt CT. Activation of the autophagy pathway decreases dengue virus
infection in Aedes aegypti cells. Parasit Vectors. 2021;14(1):551.
51. Fredericks AC, Russell TA, Wallace LE, Davidson AD, Fernandez-Sesma A,
Maringer K. Aedes aegypti (Aag2)-derived clonal mosquito cell lines reveal the effects of
pre-existing persistent infection with the insect-specific bunyavirus Phasi Charoen-like
virus on arbovirus replication. PLoS Negl Trop Dis. 2019;13(11):e0007346.
52. Yoshida K, Ono F, Chouno T, Perocho BR, Ikegami Y, Shirakigawa N, et al.
Cryoprotective enhancing effect of very low concentration of trehalose on the functions of
primary rat hepatocytes. Regen Ther. 2020;15:173-9.
53. Erdag G, Eroglu A, Morgan J, Toner M. Cryopreservation of fetal skin is improved
by extracellular trehalose. Cryobiology. 2002;44(3):218-28.
54. Beattie GM, Crowe JH, Lopez AD, Cirulli V, Ricordi C, Hayek A. Trehalose: a
cryoprotectant that enhances recovery and preserves function of human pancreatic islets
after long-term storage. Diabetes. 1997;46(3):519-23.

55.   Elbein AD, Pan YT, Pastuszak I, Carroll D. New insights on trehalose: a
multifunctional molecule. Glycobiology. 2003;13(4):17R-27R.

56.   Liu K, Dong Y, Huang Y, Rasgon JL, Agre P. Impact of trehalose transporter
knockdown on Anopheles gambiae stress adaptation and susceptibility to Plasmodium
falciparum infection. Proc Natl Acad Sci U S A. 2013;110(43):17504-9.

57.   Breeuwer P, Lardeau A, Peterz M, Joosten HM. Desiccation and heat tolerance of
Enterobacter sakazakii. J Appl Microbiol. 2003;95(5):967-73.

58.   Versele M, Thevelein JM. Lre1 affects chitinase expression, trehalose
accumulation and heat resistance through inhibition of the Cbk1 protein kinase in
Saccharomyces cerevisiae. Mol Microbiol. 2001;41(6):1311-26.

59.   Hu Y, Liu X, Liu F, Xie J, Zhu Q, Tan S. Trehalose in Biomedical
Cryopreservation-Properties, Mechanisms, Delivery Methods, Applications, Benefits, and
Problems. ACS Biomater Sci Eng. 2023;9(3):1190-204.

60.   Solocinski J, Osgood Q, Wang M, Connolly A, Menze MA, Chakraborty N. Effect
of trehalose as an additive to dimethyl sulfoxide solutions on ice formation, cellular
viability, and metabolism. Cryobiology. 2017;75:134-43.

61.   Stokich B, Osgood Q, Grimm D, Moorthy S, Chakraborty N, Menze MA.
Cryopreservation of hepatocyte (HepG2) cell monolayers: impact of trehalose.
Cryobiology. 2014;69(2):281-90.
902

A.

TPS
AAEL006446

TPP
AAEL010684?

TREH
AAEL009658

Glucose 6-phosphate → Trehalose 6-phosphate → Trehalose → 2 Glucose

UDP-glucose   UDP

H_2O iP

H_2O

B.

SignalP 6.0 prediction: Sequence

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9