1

Trehalose supports the growth of *Aedes aegypti* cells and modifies gene expression and dengue virus replication

Andrew D. Marten[1], Douglas P. Haslitt[1], Chad A. Martin[1], Daniel H. Swanson[2], Karishma Kalera[2,3], Ulysses G. Johnson[2,3], Benjamin M. Swarts[2,3], and Michael J. Conway[1]

[1]*Foundational Sciences, Central Michigan University College of Medicine, Mount Pleasant, MI 48859, USA.*

[2]*Department of Chemistry and Biochemistry, Central Michigan University, Mount Pleasant, MI 48859, USA.*

[3]*Biochemistry, Cell, and Molecular Biology Graduate Programs, Central Michigan University, Mount Pleasant, MI 48859, USA.*

16
17

Address correspondence to: Foundational Sciences, Central Michigan University

College of Medicine, Mount Pleasant, MI, 48859

E-mail: michael.conway@cmich.edu. Phone: (989) 774-3930, Fax: (989) 774-3462

22
23
24
Short title: Trehalose metabolism in *Aedes aegypti*
26
27
28
29

Abstract

Trehalose is a non-reducing disaccharide that is the major sugar found in insect hemolymph fluid. Trehalose provides energy, and promotes growth, metamorphosis, stress recovery, chitin synthesis, and insect flight. Trehalase is the only enzyme responsible for the hydrolysis of trehalose, which makes it an attractive molecular target. Here we show that *Aedes aegypti* (Aag2) cells express trehalase and that they can grow on trehalose-containing cell culture media. Trehalase activity was confirmed by treating Aag2 cells with trehalase inhibitors, which inhibited conversion of trehalose to glucose and reduced cell proliferation. Cell entry of a fluorescent trehalose probe was dependent on trehalose concentration, suggesting that trehalose moves across the cell membrane via passive transport. Culturing Aag2 cells with trehalose-containing cell culture media led to significant changes in gene expression, intracellular lipids, and dengue virus replication and specific infectivity, and increased their susceptibility to trehalase inhibitors. These data describe an *in vitro* model that can be used to rapidly screen novel trehalase inhibitors and probes and underscores the importance of trehalose metabolism in *Ae. aegypti* physiology and transmission of a mosquito-borne virus.

Introduction

Aedes aegypti is the primary vector of several important global human pathogens including dengue virus (DENV), Zika virus (ZIKV), and chikungunya virus (CHIKV). There are no targeted antivirals for these viruses and the currently licensed DENV vaccine is restricted to a subset of the human population due to the potential of eliciting antibody-dependent enhancement (ADE) in those who have not previously been infected (1-6). Further, resistance to pesticides including carbamates, organochlorines, organophosphates, and pyrethroids has been detected in the Americas, Africa, and Asia. New strategies to control disease vectors are desperately needed (7-13).

Trehalose is a non-reducing disaccharide and is the most prevalent sugar found in insect hemolymph fluid. Trehalose provides energy but also promotes growth, metamorphosis, stress recovery, chitin synthesis, and insect flight (14-28). Previous research has shown that trehalose is synthesized in the insect fat body from glucose 6-phosphate and UDP-glucose by the conserved enzymes trehalose 6-phosphate synthase (TPS) and trehalose 6-phosphate phosphatase (TPP) (16, 22, 29, 30). The hydrolysis of trehalose is under the enzymatic control of trehalase, which catalyzes the conversion of trehalose into two glucose molecules (22, 30). There are two forms of insect trehalase. Trehalase 1 (Tre-1) is soluble and has been purified from hemolymph, midgut goblet cells, and eggs. Trehalase 2 (Tre-2) is a membrane bound form and has been identified in flight muscle, follicle cells, ovary cells, spermatophore, midgut, and brain tissue (22). Although it is known that the facilitated trehalose transporter (TRET1) transfers newly synthesized trehalose from the fat body across the cellular membrane and into the

hemolymph, it is unclear if trehalose transporters exist in peripheral tissues and import

this sugar (22).

Trehalose is not synthesized by human cells, but it is present in varying quantities

in certain foods such as mushrooms, yeast, honey, beans, seaweeds, and shellfish, and

synthesized by certain bacterial species that comprise the microbiome (31). Humans

express a membrane-bound trehalase enzyme in the small intestine's brush border cells,

which catalyzes dietary trehalose to glucose (31). Mammalian cells do not normally

engage with high concentrations of trehalose, and previous research indicates that

treatment of mammalian cells with trehalose induces autophagy through an MTOR-

independent mechanism. This has been associated with both neuroprotective and antiviral

activities (32-39).

Our previous research revealed that treatment of *Ae. aegypti* with the trehalase

inhibitor validamycin A (ValA) promoted hypoglycemia and trehalose accumulation, and

corresponded with delayed larval and pupal development, ultimately preventing flight in

adult mosquitoes (23). Unfortunately, a novel trehalose analogue called 5-ThioTre was

unable to recapitulate the activity of ValA (23). We sought to develop an *in vitro* model

that would facilitate screening of novel trehalose analogues and to determine the impact

of trehalose metabolism on a mosquito-borne virus. Here, we show that the *Ae. aegypti*

cell line (Aag2) expresses a secreted trehalase isoform, and that this enzyme facilitates

growth when cell culture media glucose is exchanged for trehalose. Specifically, the

trehalase inhibitor validamycin A (ValA) and novel trehalase inhibitors prevented

conversion of cell culture media trehalose to glucose, and treatment with ValA

significantly reduced cell proliferation. Cell entry of a fluorescent trehalose probe was

dependent on trehalose concentration, suggesting that trehalose moves across the cell

membrane via passive transport. Importantly, RNA-Seq analysis of Aag2 cells grown in

trehalose-containing cell culture media revealed significant changes in gene expression in

multiple cellular pathways, including those involved in integral/intrinsic membrane

components. Trehalose also increased the concentration of intracellular lipids and viral

RNA (vRNA), shedding of infectious virus, and the specific infectivity of DENV virions,

while increasing the susceptibility of Aag2 cells and DENV to trehalase inhibitors. This

manuscript describes the use of an *in vitro* model that can be used to interrogate trehalose

metabolism in insect cells and reveals how trehalose modulates mosquito cell biology and

virus infection.

Methods

Cells and reagents. The *Ae. aegypti* cell line, Aag2 (ATCC, VA), was used for *in vitro*

studies. Aag2 cells were maintained at 28 °C with 5% CO_2 in complete cell culture

media containing DMEM no glucose media supplemented with 4,500 mg/L glucose, 10%

heat-inactivated fetal bovine serum (FBS) (Gemini, CA), 1% penicillin-streptomycin

(Thermo Fisher Scientific, MA) and 1% tryptose phosphate broth (Sigma-Aldrich, MO).

Aag2 cells were also grown using the above cell culture media with equimolar

concentrations of trehalose (2,250 mg/L) (Sigma-Aldrich, MO) instead of glucose. It is

important to note that the trehalose-containing cell culture media retained low

concentrations of glucose from FBS and tryptose phosphate broth. The glucose

concentration in our trehalose-containing cell culture media was approximately 125 mg/L

but rapidly increased in glucose concentration after inoculation onto Aag2 cells. The

virus strain used was dengue virus type 2 New Guinea C (DENV2 NGC). DENV2 stocks

were made by infecting Aag2 cells at a MOI of 1.0 and cell-free virus stocks were

harvested 8 days post-infection (dpi) and stored at -80 °C until use.

Trehalose analogues. Validamycin A (ValA) (Cayman Chemical, MI) was the only

commercially available trehalase inhibitor available at the time of this study. The

synthesis of trehalose analogues was carried out using a chemoenzymatic method as

previously reported for 5-ThioTre (40) and 6-TreNH$_2$ (41). The water solubility of 5-

ThioTre and 6-TreNH$_2$ is predicted to be very close to trehalose and stocks were

generated in water at 100 mM.

Synthesis of fluorescent trehalose derivative NBD-Trehalose. To a 25 mL round

bottom flask flushed under argon was added anhydrous *N*,*N*-dimethylformamide (10

mL), 6-TreNH$_2$ (10.9 mg, 0.032 mmol), 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole (7.6

mg, 0.038 mmol), and *N*,*N*-diisopropylethylamine (8 μL, 0.043 mmol). After stirring at

room temperature under argon overnight, thin-layer chromatography indicated that the

reaction was complete. The crude product was concentrated by rotary evaporation and

purified by reverse phase chromatography on a Biotage Isolera One automated flash

chromatography system (10 g C18 column; 0% CH$_3$CN in H$_2$O $\rightarrow$ 80% CH$_3$CN in H$_2$O)

to give NBD-Trehalose (12.9 mg, 80%) as an orange solid. ^{1}H NMR (500 MHz, D$_2$O): δ

8.38 (d, J = 9.0 Hz, 1 H), 6.38 (d, J = 9.0 Hz, 1 H), 5.18 (d, J = 4.0 Hz, 1 H), 5.06 (d, J =

3.5 Hz, 1 H), 4.11-4.07 (m, 1 H), 3.94 (bs, 1 H), 3.87-3.76 (m, 5 H), 3.71 (dd, J = 5.5,

12.0 Hz, 1 H), 3.67 (dd, J = 4.0, 10.0 Hz, 1 H), 3.47 (t, J = 10.0 Hz, 1 H), 3.46 (dd, J =

3.5, 10.0 Hz, 1 H), 3.38 (t, J = 10.0 Hz, 1 H). ^{13}C NMR (125.7 MHz, D$_2$O): δ 146.46,

144.32, 143.82, 120.62, 138.70, 100.64, 93.47, 93.33, 72.58, 72.51, 71.17, 71.34, 70.93,

70.89, 70.51, 69.61, 60.49, 48.59. HRMS (ESI-TOF) m/z: [M + Na]$^+$ Calcd for

C$_{18}$H$_{24}$N$_4$O$_{13}$Na 527.1238; Found 527.1249. Stocks of NBD-Trehalose were generated in

dimethylsulfoxide (DMSO) at 100 mM.

Cell proliferation and glucose assays. Cell proliferation assays were performed by

seeding 100 Aag2 cells into a 96 well plate in triplicate. The total number of viable cells

per well were quantified using trypan blue exclusion each day starting on day 0 and

ending on day 7. This assay was repeated using cells that were fed either glucose or

trehalose-containing cell culture media in the presence or absence of 1 mM validamycin

A (ValA). A Glucose Colorimetric Assay Kit (Cayman Chemical, MI) was used to assay

glucose concentration in Aag2 cell-free supernatant from cells grown in either glucose or

trehalose-containing cell culture media each day starting on day 0 and ending on day 3.

Glucose concentration was also assayed in Aag2 cell-free supernatant from cells grown in

trehalose-containing cell culture media and in the presence of increasing concentrations

of ValA, and trehalose analogues 5-ThioTre and 6-TreNH2, which are described above.

Glucose concentration was also assayed in Aag2 cell-free supernatant from cells that

were infected with 1 MOI DENV2 every other day starting 1-day post-infection (dpi) and

ending 7 dpi.

NBD-Tre probe assays. Aag2 cells were grown to confluence in trehalose-containing cell culture media using the Nunc Lab-Tek Chamber Slide System (Thermo Fisher Scientific, MA). Cell culture media was removed and replaced with cell culture media spiked with 0.1 mM NBD-Tre probe with and without 1 mM ValA. Negative controls included wells treated with equimolar concentrations of the DMSO vehicle control. Treated cells were incubated for 4 hours, cell culture media was then removed, and cells were stained with the nuclear stain Hoescht (Thermo Fisher Scientific, MA). Cells were imaged immediately using a Leica DM5000 B, Leica DFC550 digital camera, and Leica CTR5000 control unit. Mean fluorescence intensity (MFI) per cell was quantified using ImageJ and included 10 separate images per condition and the average MFI of 5 cells per image. This experiment was repeated by treating Aag2 cells with 0.1 mM NBD-Tre probe in the presence of increasing concentrations of glucose or trehalose and on uninfected and DENV2-infected cells.

Nile Red assay. Aag2 cells were grown to confluence in either glucose or trehalose-containing cell culture media using the Nunc Lab-Tek Chamber Slide System. A 30 mM Nile Red (Cayman Chemical, MI) stock in DMSO was diluted 1:3000 in either glucose or trehalose-containing cell culture medium and incubated for 30 minutes at room temperature. Cells were fixed and counterstained with the nuclear stain Hoescht. Cells were imaged immediately using a Leica DM5000 B, Leica DFC550 digital camera, and Leica CTR5000 control unit. Corrected total cell fluorescence (CTCF) was quantified

using integrated density, area, and mean background fluorescence data obtained from 3

separate images per condition and 20 cells per image.

RNA-Seq. RNA Sequencing (RNASeq) was conducted using a service offered by

Novogene, USA (California UC Davis) on Aag2 cells that were either maintained in

glucose or trehalose-containing cell culture media in triplicate. RNA was extracted using

the Qiagen RNeasy Plus RNA Extraction kit with gDNA eliminator columns. RNA

integrity was assessed using the RNA Nano 6000 Assay Kit of the Bioanalyzer 2100

system (Agilent Technologies, CA, USA). Total RNA was used as input material for the

RNA sample preparations. Briefly, mRNA was purified from total RNA using poly-T

oligo-attached magnetic beads. Fragmentation was carried out using divalent cations

under elevated temperature in First Strand Synthesis Reaction Buffer(5X). First strand

cDNA was synthesized using random hexamer primer and M-MuLV Reverse

Transcriptase(RNase H-). Second strand cDNA synthesis was subsequently performed

using DNA Polymerase I and RNase H. Remaining overhangs were converted into blunt

ends via exonuclease/polymerase activities. After adenylation of 3' ends of DNA

fragments, Adaptor with hairpin loop structure were ligated to prepare for hybridization.

In order to select cDNA fragments of preferentially 370~420 bp in length, the library

fragments were purified with AMPure XP system (Beckman Coulter, Beverly, USA).

Then PCR was performed with Phusion High-Fidelity DNA polymerase, Universal PCR

primers and Index (X) Primer. At last, PCR products were purified (AMPure XP system)

and library quality was assessed on the Agilent Bioanalyzer 2100 system. The clustering

of the index-coded samples was performed on a cBot Cluster Generation System using

TruSeq PE Cluster Kit v3-cBot-HS (Illumia) according to the manufacturer's

instructions. After cluster generation, the library preparations were sequenced on an

Illumina Novaseq platform and 150 bp paired-end reads were generated. Raw data (raw

reads) of fastq format were firstly processed through in-house perl scripts. In this step,

clean data (clean reads) were obtained by removing reads containing adapter, reads

containing ploy-N and low-quality reads from raw data. At the same time, Q20, Q30 and

GC content the clean data were calculated. All the downstream analyses were based on

the clean data with high quality. Reference genome and gene model annotation files were

downloaded from genome website directly. Index of the reference genome was built

using Hisat2 v2.0.5 and paired-end clean reads were aligned to the reference genome

using Hisat2 v2.0.5. We selected Hisat2 as the mapping tool for that Hisat2 can generate

a database of splice junctions based on the gene model annotation file and thus a better

mapping result than other non-splice mapping tools. featureCounts v1.5.0-p3 was used to

count the reads numbers mapped to each gene. FPKM of each gene was calculated based

on the length of the gene and reads count mapped to this gene. FPKM, expected number

of Fragments Per Kilobase of transcript sequence per Millions base pairs sequenced,

considers the effect of sequencing depth and gene length for the reads count at the same

time, and is currently the most commonly used method for estimating gene expression

levels. Differential expression analysis of the two conditions/groups was performed using

the DESeq2 R package (1.20.0). DESeq2 provide statistical routines for determining

differential expression in digital gene expression data using a model based on the

negative binomial distribution. The resulting P-values were adjusted using the Benjamini

and Hochberg's approach for controlling the false discovery rate. Genes with an adjusted

P-value $<=0.05$ found by DESeq2 were assigned as differentially expressed. Functional

annotations of differentially expressed genes were performed using the g:Cost Functional

Profiling tool at g:Profiler and NCBI's DAVID Functional Annotation Bioinformatics

tool. FASTQ files for each sample are available in the NCBI SRA database (BioProject

ID: PRJNA936836 and PRJNA882813).

Focus-forming unit assays. Aag2 cells were seeded into 96 well plates and grown to

~70% confluence in glucose-containing cell culture media. Wells were then inoculated

with 20 focus-forming units (FFUs) of DENV2 in either glucose or trehalose-containing

cell culture media for 1 hour at 27°C. Unbound virus was removed and fresh glucose-

containing cell culture media was added. At 3 dpi, confluent Aag2 monolayers were fixed

in 4% paraformaldehyde in phosphate buffered saline (PBS) for 10 minutes, cell

membranes were permeabilized with 0.1% Triton X-100, and DENV2 antigen was

stained with 1:200 anti-DENV2 envelope antibody (3H5-1, EMD Millipore) in PBS with

1% bovine serum albumin (BSA). The total number of DENV-positive foci were

revealed using a 1:200 horseradish peroxidase-conjugated secondary antibody in PBS

plus 1% BSA, and an AEC Peroxidase Substrate kit (Vector Laboratories). This

experiment was repeated using cells that were maintained in either glucose or trehalose-

containing cell culture media throughout the experiment, and digital images were

collected to quantify the total focus-forming units/well and total number of DENV2-

positive cells/well. FFU assays were also performed using cell free supernatants isolated

from DENV2-infected Aag2 cells that were maintained in either glucose or trehalose-

containing cell culture media, examining cell free supernatants at 0, 1, and 3 dpi. Cell

free supernatants were stored at -80°C until use and then used to inoculate subconfluent

Aag2 cell monolayers that were maintained in glucose-containing cell culture media.

FFU assays were also used to quantify the impact of ValA treatment on DENV2

infection. For this experiment, Aag2 cells were inoculated with 40 FFU/well DENV2 and

co-treated with 0, 0.1, 0.2, and 0.5 mg/mL ValA. Digital images were captured using the

Evos XL Core Imaging System.

RT-qPCR-based infectivity assays.

Aag2 cells were seeded into 96 well plates and grown to ~70% confluence in either

glucose or trehalose-containing cell culture media. Cells were then mock infected or

inoculated with 1×10^6 genomic equivalents of DENV2 in either glucose or trehalose-

containing cell culture media in triplicate. For the attachment assay, incubation of

DENV2 with cells took place at 4°C for 1 hr. and then unbound virus was removed, cells

were washed with PBS, and total RNA was extracted and quantified by RT-qPCR.

DENV2 viral RNA (vRNA) was normalized by total nanograms of cellular RNA. For the

cell entry and replication assay, incubation of DENV2 with cells took place at 25°C for 1

hr. and then unbound virus was removed and replaced with either glucose or trehalose-

containing cell culture media. Total RNA was extracted 3 dpi, quantified by RT-qPCR,

and normalized using total nanograms of cellular RNA. DENV2 stocks were also

generated using Aag2 cells maintained in either glucose or trehalose-containing cell

culture media, frozen at -80°C, and then RNA was extracted and quantified using RT-

qPCR. FFUs were also measured in these samples, which allowed for determination of

the specific infectivity of virions produced from glucose and trehalose-fed cells. Specific

infectivity was represented as FFU/vRNA. Amplification of the viral target was

performed using a singleplex format in 48-well plates (Illumina) with a total reaction

volume of 10 μl. Reverse transcription and quantitative PCR were performed in the same

closed tube with 50 ng of total RNA per reaction using the Quantitect RT-qPCR Kit

(Qiagen). All primers were used at final concentrations of 4 μM. For amplification of

DENV2 vRNA (nt 87–207), Forward: 5′ CAG ATC TCT GAT GAA TAA CCA ACG 3′

and Reverse: 5′ CAT TCC AAG TGA GAA TCT CTT TGT CA 3′ DENV2-specific

primers were used. The DENV2-specific probe, 5'/56-FAM/CTG CCG ATC/ZEN/TTG

CAC ATT ACC ACC GC/3IABkFQ/3′, was used for added specificity. All RT-qPCR

reactions were performed using an Eco Illumina and each time point was measured in

triplicate. Cycling conditions were 50 °C for 30 min (reverse transcription) and 95 °C for

15 min, followed by 45 cycles of 94 °C for 15 s, 55 °C for 30 s and 72 °C for 30 s.

Relative quantities of target cDNA were determined using the Pfaffl method. Student's t

tests were performed between groups to determine statistical significance.

Results

Aag2 cells express a secreted trehalase enzyme that can convert trehalose to glucose.

RNA-Seq datasets that characterize gene expression of Aag2 cells reveal high

expression of trehalose-6-phosphate synthase (TPS) and trehalase (TREH) homologues

(42). Expression of a trehalose 6-phosphate phosphatase (TPP) homologue is low but not

absent (Figure 1A). The amino acid sequence of TREH was analyzed using SignalP 6.0

and indicated a signal peptide cleavage site between amino acid positions 30 and 31

(probability 0.97812) (Figure 1B). PSORT Wolf analysis of the TREH amino acid

sequence predicted that it is located in the endoplasmic reticulum, extracellular space,

peroxisome, and plasma. Together, these data indicated that Aag2 TREH is a secreted

form of the enzyme and exists in the extracellular environment.

Most cell culture media used to support insect cell growth contains glucose. We

reasoned that if Aag2 expressed TREH, and if this enzyme was secreted, it would convert

trehalose to glucose and similarly support cell growth. Aag2 cells were grown in both

glucose and trehalose-containing cell culture media and passaged at least two times

before experiments. Aag2 cells proliferated equally in glucose and trehalose-containing

cell culture media (Figure 2A-B). Importantly, over a 3-day time course Aag2

metabolized glucose when maintained on glucose-containing cell culture media, and

generated glucose when maintained on trehalose-containing cell culture media (Figure

2C-D).

***In vitro* trehalase inhibitor assays show reduction in cell proliferation and reveal**

differences in inhibitor efficacy.

Previous *in vivo* research showed that treatment with trehalase inhibitor ValA at

0.2, 0.4, and 1.0 mM led to dose-dependent changes in *Ae. aegypti* development (23).

Cell proliferation assays were performed to determine if trehalase inhibitors interfered

with Aag2 cell metabolism when maintained in trehalose-containing cell culture media.

There was no reduction in cell proliferation when Aag2 cells maintained in glucose-

containing cell culture media were treated with 1 mM ValA (Figure 2E). However, when

Aag2 cells maintained in trehalose-containing cell culture media were treated with 1 mM

ValA, a significant reduction in cell proliferation was evident 3 days post-treatment,

which persisted throughout the time course (Figure 2E). To directly test the impact of

trehalose analogues on trehalase activity, a dose-dependence assay was performed that

quantified conversion of trehalose to glucose. ValA maintained ~75% activity as a

trehalase inhibitor at the lowest concentration tested (i.e., 0.008 mM) (Figure 2F).

Although 5-ThioTre and 6-TreNH$_2$ maintained ~75% activity at the highest concentration

tested (i.e., 1.0 mM), they had no activity at the two lowest concentrations (Figure 2F).

RNA-Seq analysis identifies differentially expressed genes involved in trehalose

metabolism and innate immunity.

Previous research confirmed that key enzymes involved in trehalose metabolism

are expressed in Aag2 cells (42). This was somewhat unexpected given that Aag2 cells

do not derive from fat body tissue. Very little is known about the response of non-fat

body cells and tissues to trehalose. RNA-Seq was performed to characterize the

differentially expressed genes in Aag2 cells maintained in glucose and trehalose-

containing cell culture media.

Significant differences in gene expression were observed. There were 157

upregulated and 118 downregulated genes in Aag2 cells that were maintained in

trehalose-containing cell culture media (Figure 3A). Although TREH expression did not

change in the RNA-Seq dataset (Supplementary Table 1) or in a follow-up RT-qPCR

validation study (Figure 3B), trehalose transporter 1 Tret1 (AAEL026825) was

upregulated 2.7-fold in the presence of trehalose (Supplementary Table 2). Functional

annotations of differentially expressed genes also showed significant changes in genes

involved in composition of the cell membrane, lipid biosynthesis, sugar conjugation,

extracellular matrix, calcium signaling, signal transduction, and mitochondrial function

Figure 3C-D). Significant changes in the expression of some innate immunity genes were

also noted (Table 1).

Passive uptake of a fluorescently labeled trehalose probe.

Considering that Aag2 cells increase expression of Tret1 in response to trehalose

while poorly expressing TPP, it is reasonable to expect that these cells can acquire

trehalose from the extracellular environment (Supplementary Tables 1-2). To enable

testing of whether Aag2 cells can accumulate extracellular trehalose, we synthesized a

novel fluorescent trehalose derivative, NBD-Tre, which consists of trehalose conjugated

to a fluorescent 2-(7-nitro-2,1,3-benzoxadiazol-4-yl) group (Figure 4A). Aag2 cells that

were maintained in trehalose-containing cell culture media were treated with 1.0 mM

NBD-Tre probe or a vehicle control for 4 hours, excess probe was removed, cells were

washed in cell culture media, and then digital images were taken using a fluorescent

microscope. 1.0 mM ValA was used to determine if trehalase inhibition would prevent

probe degradation and improve cell labeling. Quantifying the mean fluorescent

intensity/cell showed a statistically significant increase in signal when cells were treated

with the NBD-Tre probe compared to DMSO vehicle control. The addition of trehalase

inhibitor ValA did not influence cell labeling (Figure 4B) suggesting that trehalase does

not degrade the fluorescent molecule.

To test if trehalose entered into Aag2 cells using a passive/facilitated process, 0.1

mM NBD-Tre probe was suspended in cell culture media with a range of glucose and

trehalose concentrations: 0.0, 0.1, 1.0, 10.0, and 25.0 mM. A lower concentration of

NBD-Tre probe was used in this experiment because the labeling efficiency of 0.1 mM

NBD-Tre in sugar-free cell culture media was equivalent to 1.0 mM NBD-Tre in high

glucose or trehalose cell culture media. The mean fluorescent intensity per cell decreased

~10-20% at the highest two concentrations of glucose (Figures 4C,E). In contrast, the

mean fluorescent intensity per cell increased at the lowest concentration of trehalose and

decreased 40-75% at the three highest concentrations of trehalose (Figures 4D,F). These

results support that the trehalose probe entered Aag2 cells through a process of facilitated

diffusion, which is consistent with the function of known trehalose transporters such as

Tret1 (43).

Maintaining Aag2 cells in trehalose-containing cell culture media increases DENV2

shedding and specific infectivity.

Maintaining Aag2 cells in trehalose-containing cell culture media led to

significant changes in gene expression, and trehalose used a facilitated/passive process to

accumulate inside of these cells. Transcriptomic changes, which include differential gene

expression of key innate immunity genes (Table 1), might influence virus infection.

Similarly, an influx of trehalose into intracellular compartments may influence different

aspects of the virus life cycle. Dengue virus type 2 (DENV2) was used as a model vector-

borne disease to test the hypothesis that trehalose can modify virus infection in insect

cells.

Aag2 cells maintained in glucose-containing cell culture media were inoculated

with 20 focus-forming units (FFUs) of DENV2 and the infection was allowed to spread

throughout the monolayer for 5 days. Cells were also mock-infected as a negative

control. Infected and mock infected cells were then treated with either no glucose cell

culture media as a vehicle control or with 1 mM NBD-Tre probe for 4 hours. Aag2 cells

were then fixed and stained using an anti-DENV2 3H5.1 Mab and counterstained with

Hoescht. Immunofluorescence microscopy revealed diffuse cytoplasmic localization of

the NBD-Tre probe and discrete localization of DENV2 envelope (E) protein outside of

the nucleus (Figure 5). There was no clear co-localization between these molecules.

Although the immunofluorescence analysis did not reveal evidence of specific co-

localization between trehalose and DENV2 E, the diffuse localization of trehalose Aag2

may provide opportunities for interaction with other DENV2 proteins or to influence

virus replication. We tested this hypothesis using Aag2 cells maintained in glucose-

containing cell culture media, which were inoculated with 20 focus-forming units (FFUs)

of DENV2 in either glucose or trehalose-containing cell culture media for 1 hour.

Unbound virus and cell culture media was removed and replaced with glucose-containing

cell culture media. Three days later cell monolayers were fixed and stained with anti-

DENV2 MAb and the total number of FFUs per well were quantified. Pre-treatment with

either glucose or trehalose-containing cell culture media had no impact on DENV2

infection (Figure 6A). In contrast, when the same experiment was performed using Aag2

cells that were maintained in either glucose or trehalose-containing cell culture media for

the duration of the focus forming unit assay, there was a 2-fold increase in FFUs per well

in the trehalose-maintained cells (Figures 6B,D). The total number of DENV2 positive

cells were quantified in each well, which also revealed a 2-fold increase in the trehalose-

maintained cells (Figures 6C,D).

To determine the stage of the virus life cycle that was affected by trehalose, an

attachment assay was performed using Aag2 cells that were maintained in either glucose

or trehalose-containing cell culture media. Cells were inoculated with 20 FFUs DENV2

in glucose-containing cell culture media at 4 °C for 1 hour, unbound virus was removed,

cells were washed with cell culture media, and total RNA was extracted from cell

monolayers. RT-qPCR analysis showed no difference in the attachment of DENV2 to

Aag2 cells maintained in either glucose or trehalose-containing cell culture media (Figure

6E). A cell entry assay was then performed, which allowed DENV2 to attach and enter

Aag2 cells at 37 °C for 4 hours. RT-qPCR analysis showed a statistically significant

increase in cell entry of DENV2 inside of Aag2 cells that were maintained in trehalose-

containing cell culture media (Figure 6F). Supernatants were then collected from

DENV2-infected Aag2 cells that were maintained in either glucose or trehalose-

containing cell culture media and stored as viral stocks. Viral stocks were tested to

determine the total FFUs/well in a 5 µL inoculum onto Aag2 cells that were maintained

in glucose-containing cell culture media. Consistent with previous experiments,

significantly more DENV2 was shed from Aag2 cells that were maintained in trehalose-

containing cell culture media by day 3 post-infection (dpi) (Figure 6G). To test the

quality of virions produced by Aag2 cells maintained in either glucose or trehalose-

containing cell culture media, virus stocks were prepared from each condition at 5 dpi,

and then FFU and RT-qPCR assays were performed. The total FFUs per mL were

normalized by vRNA (FFU/vRNA) to determine specific infectivity of virus stocks. The

specific infectivity of DENV2 produced from Aag2 cells maintained in trehalose-

containing cell culture media was 2-fold greater (Figure 6H).

Trehalose promotes accumulation of intracellular lipids.

Previous research suggests that trehalose induces autophagy in mammalian cells,

although autophagy's role in virus infection is varied (32-36, 39, 44). The consensus is

that DENV benefits from autophagy in mammalian cells by enhancing cell survival and

promoting virus maturation (45-49). The role of autophagy in arbovirus infection in

insect cells is less clear, and conventional autophagy inducers and inhibitors may respond

differently in insect cells (44, 50). Trehalose can also be converted to glucose and used as

a substrate for lipogenesis (42).

RNA-Seq analysis revealed significant changes in gene expression related to

integral/intrinsic membrane components, and the main function of autophagy is to

accumulate and utilize lipids to enhance cell survival. We used Nile Red staining to

determine if trehalose modified the lipid environment of Aag2 cells. We identified a

slight, yet significant increase in intracellular lipid accumulation in Aag2 cells that were

maintained in trehalose-containing cell culture media (Figure 7A-B).

DENV2 infection does not impact trehalose metabolism.

Previous research has shown that DENV manipulates the metabolism of its host

cell to promote replication. Different parameters of trehalose metabolism were tested

during DENV2 infection to determine if this pathway is manipulated in Aag2 cells. Aag2

cells maintained in trehalose-containing cell culture media were either mock infected or

infected with DENV2 and conversion of trehalose to glucose was assessed on 1, 3, 5, and

7 dpi. DENV2 infection did not influence trehalase activity (Figure 8A). The impact of

DENV2 infection on trehalose uptake was also tested by infecting Aag2 cells that were

maintained in glucose-containing cell culture media with 0, 1, 10, 100, and 500 FFUs and

then labeling cells with 0.1 mM NBD-Tre 3 dpi. There was no difference in mean

fluorescence intensity per cell, therefore DENV2 infection did not influence trehalose

uptake (Figure 8B).

Trehalase inhibition reduces cell viability and DENV infection in trehalose-

dependent Aag2 cells.

We previously showed that trehalase inhibition reduced cell proliferation when

cells were maintained in trehalose-containing cell culture media (Figure 2E). We

hypothesized that this activity would reduce DENV productivity. As expected, ValA

treatment did not reduce cell viability or DENV productivity in Aag2 cells that were

maintained in glucose-containing cell culture media, and surprisingly increased DENV

productivity by 50% (Figure 9A-C). In contrast, ValA treatment significantly reduced

cell viability and DENV productivity in Aag2 cells that were maintained in trehalose-

containing cell culture media (Figure 9D-F). It is important to note that a FFU assay was

not performed directly on Aag2 cells that were maintained on trehalose-containing cell

culture media because the addition of ValA prevents establishment of a monolayer,

which is required for the assay. Instead, cell-free supernatants were harvested from ValA-

treated cells and used for downstream FFU assays performed on Aag2 cells that were

maintained on glucose-containing cell culture media.

Discussion

Control of *Aedes* spp. disease vectors is an important area of ongoing research due

to concerns regarding vaccine limitations, pesticide resistance, and increased geographic

spread (6). Trehalase inhibitors have been previously characterized as a novel class of

putative insecticides, potentially effective for controlling *Aedes* spp. disease vectors due

to the vital importance of trehalose as a blood sugar during mosquito development (14,

15, 17, 18, 20, 24, 25, 30, 40). Our previous research has revealed that the commercially

available trehalase inhibitor ValA negatively impacts mosquito development *in vivo*

when applied directly to the larval environment by showing dose-dependent activity

delaying egg hatch, pupation, eclosure, and flight of adult *Ae. Aegypti* mosquitoes (23).

Here, we describe the use of an *in vitro* model to screen for trehalase inhibitors in

Ae. aegypti and determine the impact of trehalose metabolism on a mosquito-borne virus.

ValA and novel trehalase inhibitors (5-ThioTre, 6-TreNH$_2$) prevented conversion of cell

culture media trehalose to glucose. ValA had the strongest activity, which significantly

reduced cell proliferation. Differences in the activity of trehalose analogues seen *in vitro*

and *in vivo* suggest that there are opportunities to optimize trehalase inhibitors for target

species (23). Our *in vitro* assay is a platform that could be used for high throughput

studies to develop targeted trehalose analogues.

RNA-Seq analysis showed that Tret1 expression was induced in cells that were

maintained in trehalose-containing cell culture media. The Aag2 cell line was originally

generated from whole homogenized embryos. Cells within the culture exhibit differing

morphologies, and it has been suggested that the varying morphologies of mosquito cells

in culture may be indicative of the presence of a diversity of embryonic and differentiated

cell types (51). Aag2 cells were not derived from fat body cells or tissue, which is where

Tret1 is concentrated *in vivo* (43). The role of Tret1 in peripheral cells and tissues is

unknown. Considering that the machinery for trehalose synthesis was lacking in Aag2

cells, we hypothesized that these cells may use Tret1 for trehalose uptake rather than

excretion. Previous research has shown that Tret1 is a facilitated/passive transporter (43).

We confirmed that Aag2 cells could accumulate a novel fluorescent trehalose probe,

NBD-Tre, and that uptake was prevented if unlabeled trehalose was present. This

supports facilitated/passive uptake of trehalose in Aag2 cells.

Entry of trehalose into Aag2 cells may have led to differential gene expression as

noted in the RNA-Seq dataset, although the presence of trehalose also induced expression

of integrin proteins, which mediate signaling from the cell surface. Significant

differences in gene expression were present in genes involved in composition of the cell

membrane, lipid biosynthesis, sugar conjugation, extracellular matrix, calcium signaling,

signal transduction, and mitochondrial function. These global changes in gene expression

may explain why the physiological role of trehalose extends beyond a simple source of

fuel. Trehalase inhibition can impact a wide array of physiological functions in insects

including chitin synthesis, stress protection, larval and pupal development (14, 17-20, 24-

26, 28).

Considering that trehalose enters Aag2 cells and leads to significant changes in

gene expression, we hypothesized that the environment of a trehalose-fed insect cell was

different enough to influence virus infection. This was supported by previous data

suggesting that trehalose can induce autophagy and that DENV2 benefits from autophagy

by prolonging cell survival and enhancing virus maturation (45-49). We found that Aag2

cells that were maintained in trehalose-containing cell culture media promoted DENV2

cell entry, which led to a higher number of infected cells, and increased virus shedding.

In mammalian cells, trehalose is an inducer of autophagy and can block autophagic flux

from autophagosomes to autolysosomes (39). DENV2 also induces the autophagy

pathway, and infection is reduced when autophagy is inhibited (45-49). It is possible that

trehalose increases DENV2 infection in Aag2 cells by promoting autophagy and

enhancing virus cell entry through modification of cell membranes. Previous research

suggests that the relationship between autophagy and DENV infection in Aag2 cells is

complicated and that conventional autophagy inhibitors may not work as expected in

these cells (44). Instead, we used Nile Red staining as a proxy to quantify changes to the

intracellular lipid environment. Nile Red staining revealed an increase in intracellular

lipids in Aag2 cells maintained on trehalose-containing cell culture media. It is important

to note that trehalose is the main blood sugar in insects, and it may not have the same

influence on autophagy as seen in mammalian cells.

Trehalose can also be converted to glucose, which is a substrate for lipogenesis. It

is possible that trehalose stimulates lipogenesis and the production of triglycerides and

associated fatty acids that promote virus replication and synthesis of more infectious viral

particles. RNA-Seq analysis showed that trehalose-fed cells had a 55% downregulation of

phosphoenolpyruvate carboxykinase (i.e., AAEL000006) and a 20% upregulation of fatty

acid synthase (i.e., AAEL001194) (Supplementary Table 1). Further, one of the most

upregulated genes (714%) was elongation of very long chain fatty acids protein 4

(AAEL004947). Together, these changes in gene expression may have been responsible

for the increase in intracellular lipids and perhaps the increase in virus replication and

shedding.

The specific infectivity of DENV2 also increased when virus stocks were

produced from Aag2 cells that were maintained in trehalose-containing cell culture

media. The mechanism of action that led to increased specific infectivity is unknown,

although trehalose is widely known for its ability to prevent heat and cold shock and it is

used as a cryoprotectant (52-58). Trehalose may provide some membrane or protein

stabilizing activity that either allows virus production and shedding to occur more

efficiently or it simply allows more virus to survive a freeze-thaw event. Autophagy has

also been linked to DENV maturation, which suggests that trehalose may facilitate

production of more mature virions (49). It is important to note that the trehalose

concentration used in our cell culture media is 20-50-fold less than what is typically used

for cryoprotection (52-54, 59-61).

Trehalase inhibition interferes with multiple physiological pathways in insects,

and we found that ValA reduced cell proliferation in Aag2 cells that were maintained in

trehalose-containing cell culture media. It was reasonable to hypothesize that trehalase

inhibition would also reduce DENV productivity. We showed that 1.0 mM ValA causes a

90% reduction in cell viability 5 days post-treatment in Aag2 cells that were maintained

in trehalose-containing cell culture media. The reduction in cell viability correlated with

an approximately 75% decrease in DENV vRNA shedding. *In vivo* studies show deficits

in *Ae. aegypti* egg hatch rate, larval development, pupation, and eclosure (23).

Considering that vertical transmission is less important for the maintenance and

transmission of DENV in a natural environment, trehalase inhihibition in larval habitats

would likely not impact DENV in a meaningful way. It is unclear how trehalase

inhibition impacts adult *Ae. aegypti* at this time, although these data suggest that targeting

adults could simultaneously reduce mosquito and virus fitness.

This manuscript describes an *in vitro* model that can be used to rapidly screen

novel trehalase inhibitors and probes and underscores the importance of trehalose

metabolism in *Ae. aegypti* physiology and the transmission of a mosquito-borne virus.

These data facilitate the future development of targeted trehalase analogues that will

reduce populations of disease vectors and related viruses with minimal off-target effects.

Acknowledgements

This work was supported through start-up funds from Central Michigan University

College of Medicine. BMS was supported by a grant from the National Institutes of

Health (R15AI117670).

Figure legends

Figure 1. Trehalose metabolism genes expressed in Aag2 cells. (A) Trehalose pathway

showing *Ae. aegypti* enzymes involved in synthesis: TPS; AAEL006446 and TPP;

AAEL010684 and catabolism: TREH; AAEL009658. (B) SignalP 6.0 prediction of a

signal peptide in TREH; AAEL009658.

Figure 2. Trehalose supports the growth of Aag2 cells. (A) Representative image of

Aag2 cells at 20X magnification maintained in either glucose (GLU) or trehalose (TRE)-

containing cell culture media. (B) Cell proliferation assay of Aag2 cells maintained in

either glucose (circles) or trehalose (squares)-containing cell culture media was

performed for 7 days (D0-D7). (C) Glucose assay of Aag2 cell supernatant at days 0, 1, 2,

and 3 post-feeding (D0-D3) with glucose-containing cell culture media. (D) Glucose

assay of Aag2 cell supernatant at days 0, 1, 2, and 3 (D0-D3) post-feeding with trehalose-

containing cell culture media. (E) Cell proliferation assay of Aag2 cells grown in either

glucose-containing cell culture media with (square) or without (circle) 1.0 mM ValA or

trehalose-containing cell culture media with (upside down triangle) or without (triangle)

1.0 mM ValA. (F) Glucose assay of Aag2 cell supernatants from cells maintained in

trehalose-containing cell culture media and treated with 0.008-1 mM ValA, 5-ThioTre,

and 6-TreNH2. Assays were performed in triplicate. Unpaired Student's t tests were

performed between groups to assess statistical significance. Standard deviations are

shown.

Figure 3. RNA-Seq analysis and differentially expressed genes. (A) Volcano plot of

upregulated (157) and downregulated (118) differentially expressed genes present in

Aag2 cells that were maintained in trehalose-containing cell culture media. (B) RT-qPCR

analysis of TREH expression in Aag2 cells that were maintained in either glucose (G) or

trehalose (T)-containing cell culture media on 0-3 dpt (D0-D3). (C) Functional annotation

categories present in upregulated genes. (D) Functional annotation categories in

downregulated genes.

Figure 4. NBD-Tre probe uptake in Aag2 cells. (A) Chemical structure of fluorescently

labeled trehalose probe NBD-Tre. (B) Aag2 cells were treated with either vehicle control

(DMSO), 1.0 mM NBD-Tre probe, or with 1.0 mM NBD-Tre probe and 1.0 mM ValA in

glucose free cell culture media for 4 hours and then mean fluorescence intensity/cell was

determined. (C-D) Aag2 cells were treated with 1.0 mM NBD-Tre probe in the presence

of (C) 0, 0.1, 1, 10, and 25 mM glucose and (D) 0, 0.1, 1, 10, and 25 mM trehalose for 4

hours and then mean fluorescence intensity/cell was determined. (E-F) Representative

images of Aag2 cells that were treated with 1.0 mM NBD-Tre probe in the presence of

(E) 0, 0.1, 1, 10, and 25 mM glucose and (F) 0, 0.1, 1, 10, and 25 mM trehalose with a

Hoescht nuclear counterstain. Assays were performed in triplicate. Unpaired Student's t

tests were performed between groups to assess statistical significance. Standard

deviations are shown.

Figure 5. Co-localization of NBD-Tre and DENV2 in Aag2 cells. Aag2 cells were

uninfected (Mock) or infected with DENV2 for 5 days and then treated with no glucose

cell culture media alone (Vehicle) or 1 mM NBD-Tre probe for 4 hours. Aag2 cells were

then fixed and stained with anti-DENV2 Mab and digital images were obtained at 20X

magnification. Hoescht nuclear stain was added to each condition.

Figure 6. Trehalose promotes DENV2 cell entry and virus shedding. Aag2 cells were

Mock infected or infected with DENV in either glucose (GLU) or trehalose (TRE)-

containing cell culture media. (A) Trehalose-containing cell culture media was only

present during inoculation and was then exchanged for glucose-containing cell culture

media and focus-forming units were quantified 3 dpi. (B) Trehalose-containing cell

culture media was present during inoculation and for the three days prior to a focus-

forming unit assay. (C) The total number of DENV2 positive cells per well were

quantified from (B). (D) Representative images from (B) showing distinct foci are shown.

(E) An attachment assay was performed by incubating DENV2 with Aag2 cells at 4°C

for 1 hour. Unbound virus was removed, cells were washed with PBS, and total RNA was

extracted and quantified by RT-qPCR. DENV2 viral RNA (vRNA) was normalized by

total nanograms of cellular RNA (F) A cell entry/replication assay was performed by

incubating DENV2 with Aag2 cells at 25°C for 1 hour. Unbound virus was removed and

replaced with either glucose or trehalose-containing cell culture media. Total RNA was

extracted 3 dpi, quantified by RT-qPCR, an DENV2 vRNA was normalized using total

nanograms of cellular RNA. (G) A virus shedding assay was performed by collecting cell

free supernatants at 0, 1, and 3 dpi from Aag2 cells that were inoculated with DENV2 in

either glucose or trehalose-containing cell culture media. Cell free supernatants were used

to inoculate monolayers of Aag2 cells grown in glucose-containing cell culture media and

focus forming unit assays were used to quantify the infectious units present in each

sample. (H) DENV2 stocks were generated from Aag2 cells that were grown in either

glucose or trehalose-containing cell culture media. Focus forming unit assays were

performed to quantify the infectivity of each stock. Total RNA was also extracted from

virus stocks and DENV2 vRNA was quantified using RT-qPCR. Specific infectivity was

derived by normalizing the total number of focus forming units in a sample by its vRNA

content (i.e., FFU/vRNA). Assays were performed in triplicate. Unpaired Student's t tests

were performed between groups to assess statistical significance. Standard deviations are

shown.

Figure 7. Trehalose promotes intracellular lipid accumulation in Aag2 cells. (A)

Intracellular lipids were labeled in Aag2 cells maintained in either glucose (GLU) or

trehalose (TRE) cell culture media using either vehicle control (DMSO) or Nile Red.

Cells were fixed and digital images were obtained at 20X magnification. Hoescht nuclear

stain was added to each condition. (B) The corrected total cell fluorescence (CTCF) was

quantified using the integrated density of 60 individual cells per condition. Assays were

performed in triplicate. Unpaired Student's t tests were performed between groups to

assess statistical significance. Standard deviations are shown.

Figure 8. DENV2 infection does not influence trehalase activity or trehalose uptake.

(A) A glucose assay was performed on cell-free supernatants taken from Aag2 cells

maintained in trehalose-containing cell culture media that were either uninfected (Mock)

or infected with DENV2 for 1, 3, 5, and 7 dpi. (B) A trehalose probe uptake assay was

performed on Aag2 cells that were either uninfected (Mock) or infected with 1, 10, 100,

and 500 FFUs of DENV2 5 dpi. 1.0 mM NBD-Tre probe was added to cells in glucose

free cell culture media for 4 hours. Unbound probe was removed, and then digital images

were obtained at 20X magnification. Mean fluorescence intensity was determined for 50

individual cells per condition. Assays were performed in triplicate. Unpaired Student's t

tests were performed between groups to assess statistical significance. Standard

deviations are shown.

Figure 9. Validamycin A reduces cell viability and DENV2 productivity when Aag2

cells are maintained in trehalose-containing cell culture media. (A) Subconfluent

Aag2 cells maintained in glucose-containing cell culture media were treated with 0, 0.2,

0.4, and 1.0 mM ValA for 5 days and then trypan blue exclusion assays were performed

to quantify percent cell viability. (B) Subconfluent Aag2 cells maintained in glucose-

containing cell culture media were inoculated with 20 FFUs of DENV for 1 hour,

followed by treatment with 0, 0.2, 0.4, and 1.0 mM ValA for 5 days. vRNA was extracted

from equivalent volumes of cell free supernatants and quantified by RT-qPCR. (C)

Subconfluent Aag2 cells maintained in glucose-containing cell culture media were

inoculated with 20 FFUs of DENV for 1 hour, followed by treatment with 0, 0.2, 0.4, and

1.0 mM ValA for 3 days. FFU assays were performed to quantify the total FFUs present

in each well. (D) Subconfluent Aag2 cells maintained in trehalose-containing cell culture

media were treated with 0, 0.2, 0.4, and 1.0 mM ValA for 5 days and then trypan blue

exclusion assays were performed to quantify percent cell viability. (E) Subconfluent

Aag2 cells maintained in trehalose-containing cell culture media were inoculated with 20

FFUs of DENV for 1 hour, followed by treatment with 0, 0.2, 0.4, and 1.0 mM ValA for

5 days. vRNA was extracted from equivalent volumes of cell free supernatants and

quantified by RT-qPCR. (F) Subconfluent Aag2 cells maintained in trehalose-containing

cell culture media were inoculated with 20 FFUs of DENV for 1 hour, followed by

treatment with 0, 0.2, 0.4, and 1.0 mM ValA for 3 days. Cell free supernatants were

harvested and used for a downstream FFU assay performed on Aag2 cells maintained on

glucose-containing cell culture media, which quantified the total FFUs present in each

well. Assays were performed in triplicate. Unpaired Student's t tests were performed

between groups to assess statistical significance. Standard deviations are shown.

Supplementary Table 1. Complete readcount and FPKM list of all genes in GLU vs

TRE Aag2 cells.

Supplementary Table 2. Complete readcount and FPKM list of upregulated genes in

GLU vs TRE Aag2 cells.

Supplementary Table 3. Complete readcount and FPKM list of all downregulated in

GLU vs TRE Aag2 cells.

References

1.   Paul LM, Carlin ER, Jenkins MM, Tan AL, Barcellona CM, Nicholson CO, et al.
Dengue virus antibodies enhance Zika virus infection. Clin Transl Immunology.
2016;5(12):e117.
2.   Masel J, McCracken MK, Gleeson T, Morrison B, Rutherford G, Imrie A, et al.
Does prior dengue virus exposure worsen clinical outcomes of Zika virus infection? A
systematic review, pooled analysis and lessons learned. PLoS Negl Trop Dis.
2019;13(1):e0007060.
3.   Guzman MG, Alvarez M, Halstead SB. Secondary infection as a risk factor for
dengue hemorrhagic fever/dengue shock syndrome: an historical perspective and role of
antibody-dependent enhancement of infection. Arch Virol. 2013;158(7):1445-59.
4.   Obi JO, Gutiérrez-Barbosa H, Chua JV, Deredge DJ. Current Trends and
Limitations in Dengue Antiviral Research. Trop Med Infect Dis. 2021;6(4).
5.   Dighe SN, Ekwudu O, Dua K, Chellappan DK, Katavic PL, Collet TA. Recent
update on anti-dengue drug discovery. Eur J Med Chem. 2019;176:431-55.
6.   Conway MJ, Haslitt DP, Swarts BM. Targeting Aedes aegypti Metabolism with
Next-Generation Insecticides. Viruses. 2023;15(2).
7.   Rose RI. Pesticides and public health: integrated methods of mosquito
management. Emerg Infect Dis. 2001;7(1):17-23.
8.   Moyes CL, Vontas J, Martins AJ, Ng LC, Koou SY, Dusfour I, et al.
Contemporary status of insecticide resistance in the major Aedes vectors of arboviruses
infecting humans. PLoS Negl Trop Dis. 2017;11(7):e0005625.
9.   Zulfa R, Lo WC, Cheng PC, Martini M, Chuang TW. Updating the Insecticide
Resistance Status of Aedes aegypti and Aedes albopictus in Asia: A Systematic Review
and Meta-Analysis. Trop Med Infect Dis. 2022;7(10).
10.   Roberts DR, Andre RG. Insecticide resistance issues in vector-borne disease
control. Am J Trop Med Hyg. 1994;50(6 Suppl):21-34.
11.   Kasai S, Itokawa K, Uemura N, Takaoka A, Furutani S, Maekawa Y, et al.
Discovery of super-insecticide-resistant dengue mosquitoes in Asia: Threats of
concomitant knockdown resistance mutations. Sci Adv. 2022;8(51):eabq7345.
12.   Du Y, Nomura Y, Zhorov BS, Dong K. Sodium Channel Mutations and
Pyrethroid Resistance in Aedes aegypti. Insects. 2016;7(4).
13.   Amelia-Yap ZH, Chen CD, Sofian-Azirun M, Low VL. Pyrethroid resistance in
the dengue vector Aedes aegypti in Southeast Asia: present situation and prospects for
management. Parasit Vectors. 2018;11(1):332.
14.   Zhang L, Qiu LY, Yang HL, Wang HJ, Zhou M, Wang SG, et al. Study on the
Effect of Wing Bud Chitin Metabolism and Its Developmental Network Genes in the
Brown Planthopper, Nilaparvata lugens, by Knockdown of TRE Gene. Front Physiol.
2017;8:750.
15.   Xia Y, Clarkson JM, Charnley AK. Trehalose-hydrolysing enzymes of
Metarhizium anisopliae and their role in pathogenesis of the tobacco hornworm,
Manduca sexta. J Invertebr Pathol. 2002;80(3):139-47.

16.   Wyatt GR, Kale GF. The chemistry of insect hemolymph. II. Trehalose and other
carbohydrates. J Gen Physiol. 1957;40(6):833-47.
17.   Wegener G, Macho C, Schloder P, Kamp G, Ando O. Long-term effects of the
trehalase inhibitor trehazolin on trehalase activity in locust flight muscle. J Exp Biol.
2010;213(Pt 22):3852-7.
18.   Wegener G, Tschiedel V, Schloder P, Ando O. The toxic and lethal effects of the
trehalase inhibitor trehazolin in locusts are caused by hypoglycaemia. J Exp Biol.
2003;206(Pt 7):1233-40.
19.   Thorat LJ, Gaikwad SM, Nath BB. Trehalose as an indicator of desiccation stress
in Drosophila melanogaster larvae: a potential marker of anhydrobiosis. Biochem
Biophys Res Commun. 2012;419(4):638-42.
20.   Tatun N, Wangsantitham O, Tungjitwitayakul J, Sakurai S. Trehalase activity in
fungus-growing termite, Odontotermes feae (Isoptera: Termitideae) and inhibitory effect
of validamycin. J Econ Entomol. 2014;107(3):1224-32.
21.   Tang B, Yang M, Shen Q, Xu Y, Wang H, Wang S. Suppressing the activity of
trehalase with validamycin disrupts the trehalose and chitin biosynthesis pathways in the
rice brown planthopper, Nilaparvata lugens. Pestic Biochem Physiol. 2017;137:81-90.
22.   Shukla E, Thorat LJ, Nath BB, Gaikwad SM. Insect trehalase: physiological
significance and potential applications. Glycobiology. 2015;25(4):357-67.
23.   Marten AD, Stothard AI, Kalera K, Swarts BM, Conway MJ. Validamycin A
Delays Development and Prevents Flight in Aedes aegypti (Diptera: Culicidae). J Med
Entomol. 2020;57(4):1096-103.
24.   Katagiri N, Ando O, Yamashita O. Reduction of glycogen in eggs of the
silkworm, Bombyx mori, by use of a trehalase inhibitor, trehazolin, and diapause
induction in glycogen-reduced eggs. J Insect Physiol. 1998;44(12):1205-12.
25.   Liebl M, Nelius V, Kamp G, Ando O, Wegener G. Fate and effects of the
trehalase inhibitor trehazolin in the migratory locust (Locusta migratoria). J Insect
Physiol. 2010;56(6):567-74.
26.   Chen J, Zhang D, Yao Q, Zhang J, Dong X, Tian H, et al. Feeding-based RNA
interference of a trehalose phosphate synthase gene in the brown planthopper,
Nilaparvata lugens. Insect Mol Biol. 2010;19(6):777-86.
27.   Chen J, Tang B, Chen H, Yao Q, Huang X, Zhang D, et al. Different functions of
the insect soluble and membrane-bound trehalase genes in chitin biosynthesis revealed by
RNA interference. PLoS One. 2010;5(4):e10133.
28.   Asano N, Yamaguchi T, Kameda Y, Matsui K. Effect of validamycins on
glycohydrolases of Rhizoctonia solani. J Antibiot (Tokyo). 1987;40(4):526-32.
29.   Murphy TA, Wyatt GR. ENZYMATIC REGULATION OF TREHALOSE AND
GLYCOGEN SYNTHESIS IN THE FAT BODY OF AN INSECT. Nature.
1964;202:1112-3.
30.   Becker A, Schloder P, Steele JE, Wegener G. The regulation of trehalose
metabolism in insects. Experientia. 1996;52(5):433-9.
31.   Arguelles JC. Why can't vertebrates synthesize trehalose? J Mol Evol. 2014;79(3-
4):111-6.
32.   Yuan S, Zhang ZW, Li ZL. Trehalose May Decrease the Transmission of Zika
Virus to the Fetus by Activating Degradative Autophagy. Front Cell Infect Microbiol.
2017;7:402.

33. Wu Q, Jiang D, Huang C, van Dyk LF, Li L, Chu HW. Trehalose-mediated
autophagy impairs the anti-viral function of human primary airway epithelial cells. PLoS
One. 2015;10(4):e0124524.
34. Clark AE, Zhu Z, Krach F, Rich JN, Yeo GW, Spector DH. Zika virus is
transmitted in neural progenitor cells via cell-to-cell spread and infection is inhibited by
the autophagy inducer trehalose. J Virol. 2021;95(5).
35. Clark AE, Sabalza M, Gordts P, Spector DH. Human Cytomegalovirus
Replication Is Inhibited by the Autophagy-Inducing Compounds Trehalose and SMER28
through Distinctively Different Mechanisms. J Virol. 2018;92(6).
36. Belzile JP, Sabalza M, Craig M, Clark AE, Morello CS, Spector DH. Trehalose,
an mTOR-Independent Inducer of Autophagy, Inhibits Human Cytomegalovirus
Infection in Multiple Cell Types. J Virol. 2016;90(3):1259-77.
37. Rodríguez-Navarro JA, Rodríguez L, Casarejos MJ, Solano RM, Gómez A,
Perucho J, et al. Trehalose ameliorates dopaminergic and tau pathology in parkin
deleted/tau overexpressing mice through autophagy activation. Neurobiol Dis.
2010;39(3):423-38.
38. Perucho J, Casarejos MJ, Gomez A, Solano RM, de Yébenes JG, Mena MA.
Trehalose protects from aggravation of amyloid pathology induced by isoflurane
anesthesia in APP(swe) mutant mice. Curr Alzheimer Res. 2012;9(3):334-43.
39. Zhang X, Chen S, Song L, Tang Y, Shen Y, Jia L, et al. MTOR-independent,
autophagic enhancer trehalose prolongs motor neuron survival and ameliorates the
autophagic flux defect in a mouse model of amyotrophic lateral sclerosis. Autophagy.
2014;10(4):588-602.
40. Danielson ND, Collins J, Stothard AI, Dong QQ, Kalera K, Woodruff PJ, et al.
Degradation-resistant trehalose analogues block utilization of trehalose by hypervirulent
Clostridioides difficile. Chem Commun (Camb). 2019;55(34):5009-12.
41. Kalera K, Liu R, Lim J, Pathirage R, Swanson DH, Johnson UG, et al. Targeting
Mycobacterium tuberculosis Persistence through Inhibition of the Trehalose Catalytic
Shift. ACS Infect Dis. 2024;10(4):1391-404.
42. Marten AD, Tift CT, Tree MO, Bakke J, Conway MJ. Chronic depletion of
vertebrate lipids in A*edes aegypti* cells dysregulates lipid metabolism
and inhibits innate immunity without altering dengue
infectivity. In Press ed. PLoS Neglected Tropical Diseases2022.
43. Kikawada T, Saito A, Kanamori Y, Nakahara Y, Iwata K, Tanaka D, et al.
Trehalose transporter 1, a facilitated and high-capacity trehalose transporter, allows
exogenous trehalose uptake into cells. Proc Natl Acad Sci U S A. 2007;104(28):11585-
90.
44. Brackney DE, Correa MA, Cozens DW. The impact of autophagy on arbovirus
infection of mosquito cells. PLoS Negl Trop Dis. 2020;14(5):e0007754.
45. Heaton NS, Perera R, Berger KL, Khadka S, Lacount DJ, Kuhn RJ, et al. Dengue
virus nonstructural protein 3 redistributes fatty acid synthase to sites of viral replication
and increases cellular fatty acid synthesis. Proc Natl Acad Sci U S A.
2010;107(40):17345-50.
46. Lee YR, Lei HY, Liu MT, Wang JR, Chen SH, Jiang-Shieh YF, et al. Autophagic
machinery activated by dengue virus enhances virus replication. Virology.
2008;374(2):240-8.

47. Heaton NS, Randall G. Dengue virus-induced autophagy regulates lipid
metabolism. Cell Host Microbe. 2010;8(5):422-32.
48. McLean JE, Wudzinska A, Datan E, Quaglino D, Zakeri Z. Flavivirus NS4A-
induced autophagy protects cells against death and enhances virus replication. J Biol
Chem. 2011;286(25):22147-59.
49. Mateo R, Nagamine CM, Spagnolo J, Méndez E, Rahe M, Gale M, Jr., et al.
Inhibition of cellular autophagy deranges dengue virion maturation. J Virol.
2013;87(3):1312-21.
50. Chen TY, Smartt CT. Activation of the autophagy pathway decreases dengue
virus infection in Aedes aegypti cells. Parasit Vectors. 2021;14(1):551.
51. Fredericks AC, Russell TA, Wallace LE, Davidson AD, Fernandez-Sesma A,
Maringer K. Aedes aegypti (Aag2)-derived clonal mosquito cell lines reveal the effects of
pre-existing persistent infection with the insect-specific bunyavirus Phasi Charoen-like
virus on arbovirus replication. PLoS Negl Trop Dis. 2019;13(11):e0007346.
52. Yoshida K, Ono F, Chouno T, Perocho BR, Ikegami Y, Shirakigawa N, et al.
Cryoprotective enhancing effect of very low concentration of trehalose on the functions
of primary rat hepatocytes. Regen Ther. 2020;15:173-9.
53. Erdag G, Eroglu A, Morgan J, Toner M. Cryopreservation of fetal skin is
improved by extracellular trehalose. Cryobiology. 2002;44(3):218-28.
54. Beattie GM, Crowe JH, Lopez AD, Cirulli V, Ricordi C, Hayek A. Trehalose: a
cryoprotectant that enhances recovery and preserves function of human pancreatic islets
after long-term storage. Diabetes. 1997;46(3):519-23.
55. Elbein AD, Pan YT, Pastuszak I, Carroll D. New insights on trehalose: a
multifunctional molecule. Glycobiology. 2003;13(4):17R-27R.
56. Liu K, Dong Y, Huang Y, Rasgon JL, Agre P. Impact of trehalose transporter
knockdown on Anopheles gambiae stress adaptation and susceptibility to Plasmodium
falciparum infection. Proc Natl Acad Sci U S A. 2013;110(43):17504-9.
57. Breeuwer P, Lardeau A, Peterz M, Joosten HM. Desiccation and heat tolerance of
Enterobacter sakazakii. J Appl Microbiol. 2003;95(5):967-73.
58. Versele M, Thevelein JM. Lre1 affects chitinase expression, trehalose
accumulation and heat resistance through inhibition of the Cbk1 protein kinase in
Saccharomyces cerevisiae. Mol Microbiol. 2001;41(6):1311-26.
59. Hu Y, Liu X, Liu F, Xie J, Zhu Q, Tan S. Trehalose in Biomedical
Cryopreservation-Properties, Mechanisms, Delivery Methods, Applications, Benefits,
and Problems. ACS Biomater Sci Eng. 2023;9(3):1190-204.
60. Solocinski J, Osgood Q, Wang M, Connolly A, Menze MA, Chakraborty N.
Effect of trehalose as an additive to dimethyl sulfoxide solutions on ice formation,
cellular viability, and metabolism. Cryobiology. 2017;75:134-43.
61. Stokich B, Osgood Q, Grimm D, Moorthy S, Chakraborty N, Menze MA.
Cryopreservation of hepatocyte (HepG2) cell monolayers: impact of trehalose.
Cryobiology. 2014;69(2):281-90.
918

A.

B.

	Hoescht	NBD-Tre	DENV2	Merge
Vehicle + Mock				
Vehicle + DENV2				
NBD-Tre + Mock				
NBD-Tre + DENV2				

A.

	Hoescht	Nile Red	Merge
Vehicle + GLU			
Nile Red+ GLU			
Vehicle + TRE			
Nile Red + TRE			

B.

A.

B.