Expanding the HP1a-binding consensus and molecular grammar for heterochromatin

assembly

Serafin U. Colmenares[1,†], Shingo Tsukamoto[2,†], Collin Hickmann[1], Lucy D. Brennan[1],

Mohammad Khavani[2], Mohammad R. K. Mofrad[2,3,*], Gary H. Karpen[1,4,*]

[1]Department of Molecular and Cell Biology, University of California, Berkeley

[2]Molecular Cell Biomechanics Laboratory, Departments of Bioengineering and Mechanical

Engineering, University of California, Berkeley

[3]Molecular Biophysics and Integrative BioImaging Division, Lawrence Berkeley National Lab

[4]Department of BioEngineering and BioMedical Sciences, Lawrence Berkeley National Lab

[†]These authors contributed equally to this work.

* Correspondence: gkarpen@berkeley.edu

ABSTRACT

The recruitment of Heterochromatin Protein 1 (HP1) partners is essential for heterochromatin assembly and function, yet our knowledge regarding their organization in heterochromatin remains limited. Here we show that interactors engage the Drosophila HP1 (HP1a) dimer through a degenerate and expanded form of the previously identified PxVxL motif, which we now term HP1a Access Codes (HACs). These HACs reside in disordered regions, possess high conservation among Drosophila homologs, and contain alternating hydrophobic residues nested in a cluster of positively charged amino acids. These findings and molecular dynamics simulations identify key electrostatic interactions that modulate HP1a-binding strength and provide a dramatically improved HP1a-binding consensus motif that can reveal protein partners and the molecular grammar involved in heterochromatin assembly. We propose HP1a acts as a scaffold for other heterochromatin components containing HAC motifs, which in turn may regulate the function and higher order structure of the heterochromatin compartment.

INTRODUCTION

Heterochromatin forms an evolutionarily conserved nuclear compartment critical for genome

integrity. In humans and Drosophila, heterochromatin houses 25-30% of the genome,

respectively, that is primarily composed of repetitive DNA such as transposons and satellite

sequences (Hoskins *et al.*, 2007) enriched in gene-poor pericentric and telomeric regions of

chromosomes. Important for transposon suppression, chromosome segregation, and telomere

protection, heterochromatin displays physical, molecular, and functional attributes distinct

from gene-rich euchromatin (Janssen, Colmenares and Karpen, 2018). Constitutively dense

through interphase, the heterochromatin compartment is transcriptionally repressive, replicates

later in S phase in many organisms, and utilizes alternative DNA repair pathways (Chiolo *et*

al., 2011; Janssen, Colmenares and Karpen, 2018).

Various factors converge in heterochromatin to perform its many functions. Transcriptional

repression, as assayed by the insertion of reporter genes or chromosome rearrangements, have

been attributed to many heterochromatin factors (HP1a, Su(var)3-9, HP2) (Eissenberg *et al.*,

1992; Schotta *et al.*, 2002; Shaffer *et al.*, 2002). Interestingly, hundreds of genes reside in

Drosophila heterochromatin, some of which are essential for viability (Weiler and Wakimoto,

1995; Dimitri *et al.*, 2003), suggesting the presence of an unknown alternative transcriptional

program specific for the heterochromatin landscape. Meanwhile, transcription of repetitive

DNA has been shown in many organisms to be instrumental in heterochromatin

establishment/maintenance and suppression of transposable elements by providing a non-

coding scaffold for small RNAs that recruit heterochromatin complexes (Piwi, Ago2)

(Brennecke *et al.*, 2007; Czech *et al.*, 2008). Heterochromatin also employs factors that inhibit

replication fork progression and suppress polytenization (Rif1, SuUR) (Makunin *et al.*, 2002;

Seller and O'Farrell, 2018), or direct double-strand breaks to the periphery for homologous

recombination repair (Smc5/6) (Chiolo *et al.*, 2011). Finally, the heterochromatin subnuclear

compartment houses components of sister chromatid cohesion (Dmt) (Yamada *et al.*, 2017)

necessary for the coordinated distribution of sister chromatids during mitosis and undergoes

dissolution and re-assembly every cell cycle. How such various factors involved in disparate

functions are recruited and organized within the same 3-D heterochromatin compartment

remains largely unknown.

Central to heterochromatin organization and function is Heterochromatin Protein 1, or HP1,

that specifically binds and propagates the heterochromatic histone mark di- and tri-methylated

H3 K9 by recruiting methyltransferases. HP1 also binds multiple proteins, many of which have

been shown to require HP1 for enrichment in heterochromatin. HP1 is comprised of two

globular structured domains: a chromodomain (CD) that binds methylated H3 K9 (Bannister

et al., 2001; Lachner *et al.*, 2001) and a chromoshadow domain (CSD) important for

dimerization and protein partner binding (Brasher *et al.*, 2000). Other than short, disordered

regions at the flanks of the protein, HP1 also possesses a single unstructured hinge region

between the two globular domains that binds nucleic acid (Muchardt *et al.*, 2002). Previously,

we and others have shown that the Drosophila homolog HP1a co-purifies with dozens to

hundreds of proteins (Alekseyenko *et al.*, 2014; Swenson *et al.*, 2016), and that some partners

display a remarkable variation of enrichment patterns, or "subdomains," that indicate the

presence of a higher-order structure within heterochromatin (Swenson *et al.*, 2016). Whether

the large number of partners utilize individual mechanisms to target HP1a or a common motif

to access the heterochromatin domain is an important fundamental question for understanding

heterochromatin structure and organization.

A pentameric HP1-binding motif (PxVxL) was previously identified through a phage display

screen to bind the HP1 CSD dimer. Based on the peptide consensus sequence, this pentamer

contains alternating hydrophobic amino acids of proline, valine, and leucine that give rise to

its name (Smothers and Henikoff, 2000). Structural studies showed that PxVxL peptides bind

asymmetrically to a hydrophobic pocket formed by the HP1 CSD dimer, with the central valine

occupying the cleft formed by the two CSD C-termini (Thiru *et al.*, 2004). Other domains of

HP1, such as the CD and hinge regions, have also been implicated in binding other proteins

(Badugu *et al.*, 2005). However, most examples of HP1 partners identified across many species

from fission yeast to humans utilize the PxVxL motif, signifying its evolutionary importance

(Thiru *et al.*, 2004; Coustham *et al.*, 2006; Honda and Selker, 2008).

The recent discovery that heterochromatin domains display biophysical properties of

biocondensates *in vivo* (Strom *et al.*, 2017) has also prompted a re-evaluation of the PxVxL

motif as a potentially important contributor to emergent properties and functions. As

biocondensates form membraneless compartments through multivalency and the cumulative

weak interactions among components, we propose that HP1 acts as a scaffold that recruits

multiple heterochromatin components through the PxVxL motif. In addition, peptides

containing PxVxL motifs have been shown to enhance HP1 dimerization (Mendez *et al.*, 2011)

and promote or inhibit phase separation *in vitro* and *in silico* (Thiru *et al.*, 2004; Larson *et al.*,

2017; Her *et al.*, 2022). Thus, the PxVxL motif may represent not only a common HP1a

interaction interface but also a mechanism to modulate local or global HP1a dynamics and

condensate composition and biophysical properties.

Variants of the PxVxL motif, such as LxVxL, LxVxI, and CxVxL, have also been identified

(Bassett *et al.*, 2008; Mendez *et al.*, 2011; Mendez, Mandt and Elgin, 2013; Meyer-Nava *et al.*,

2020). Interestingly, a non-canonical motif (LxVxI) displayed a higher binding affinity to the

HP1a CSD dimer than canonical (PxVxL) and near-canonical (PxVxV) motifs (Mendez *et al.*,

2011). The canonical PxVxL motif is also not universally sufficient for binding, as evidenced

by mammalian STAM2a, which has a PxVxL motif but fails to bind the CSD (Lechner *et al.*,

2005). Furthermore, the HP1a C-terminal extension (CTE) and the flanking residues of the

PxVxL motif are implicated in HP1a-partner binding selectivity and interaction strength (Thiru

et al., 2004; Mendez *et al.*, 2011; Mendez, Mandt and Elgin, 2013; Liu *et al.*, 2017). Deletion

of the last 3 residues on the CTE enhanced HP1a dimerization but severely reduced the binding

of PxVxL peptides, whereas chemical shift perturbations were detected at HP1a residues that

contact positions beyond the PxVxL pentamer core (Mendez *et al.*, 2011). Altogether, these

findings raise questions about the combination of hydrophobic residues sufficient for PxVxL

function, and whether additional molecular features exist beyond the pentapeptide motif.

Here we utilize a combination of evolutionary comparisons, *in vivo* imaging, and full atomistic

molecular dynamics (MD) simulations to comprehensively define the molecular signature of

the PxVxL motif in Drosophila and determine the specific interactions that regulate HP1a

binding. Our analyses show that PxVxL motifs of HP1a-binding partners are highly conserved

across Drosophila species, embed in mainly disordered domains of HP1a partner proteins, and

are driven by charged residues that we now propose to be a critical component of the molecular

grammar for heterochromatin targeting. As a result, we have expanded the number of known

functional HP1a-binding sites and provide an improved signature with higher predictive power,

which we refer to as the HP1a Access Code, or "HAC". We have also determined that HACs

exhibit a range of affinity for HP1a, which are modulated by electrostatic interactions. We

propose that HACs embedded in disordered domains are an important part of the molecular

language used to assemble the heterochromatin condensate, and that differences in HP1a

affinity among protein partners are critical for the composition, ultrastructure, and function of

heterochromatin.

RESULTS

Imaging peptide localization in cells identifies new HP1a-binding sequences

Discovery of the foundational HP1a-binding pentamer sequence, PxVxL, showed that it

consists of alternating nonpolar residues with the central valine, designated as position 0,

binding the junction of two CSD domains in an HP1a dimer (**Figure 1A**). Correspondingly,

the proline (P) was designated as position -2 and leucine (L) as position +2, while positions -1

and +1 displayed higher variability, and were thus represented as 'x'. Together, the trio of

nonpolar residues occupy a hydrophobic pocket formed by the HP1a CSD dimer, as shown by

structural studies (Thiru *et al.*, 2004), and is thought to mediate the ability of HP1a to associate

with protein complexes.

HP1a binding sites have been mapped in many proteins, but of those only 6 in Drosophila have

been shown to contain functional PxVxL motifs, as demonstrated by disrupting HP1a

association through mutation of the central Val residue. These include the SNF2-like chromatin

remodeler XNP (Bassett *et al.*, 2008), histone H3 K36 demethylase KDM4A (Lin *et al.*, 2008),

zinc finger protein Odj (Kasinathan *et al.*, 2020), structural heterochromatin protein HP2

(Stephens *et al.*, 2005), sister chromatid cohesion regulator Dmt (Yamada *et al.*, 2017), and

transposon repressor Piwi (Brower-Toland *et al.*, 2007) (**Supplementary Figure 1A**). A

sequence alignment showed that while position 0 is consistently valine, the other positions vary

between 3 hydrophobic residues (proline, leucine, and cysteine at -2 and leucine, isoleucine,

valine at +2) (**Figure 1B**). Using the expanded consensus signature of HP1a-binding domains

at positions -2 and +2, we probed the Drosophila proteome, as well as proteins annotated to

localize to the nucleus or cytoplasm, for potential HP1a interactors (**Supplementary Figure**

1B). Not surprisingly, such a small motif identified over 50% of the proteome as HP1a

interactors, including more than 50% of proteins annotated as cytoplasmic, where HP1a is not

normally found. This led us to investigate whether the consensus sequence and our ability to

predict HP1a binding domains can be enhanced. We note, for example, that positively charged

residues frequently occupy the -1 and +1 positions along with flanking regions (**Figure 1B**),

suggesting that other positions on the HP1a-binding sequence contribute to their function and

could improve our understanding of the functional 'HP1a Access Code', hereafter referred to

as "HAC."

In order to discover additional HAC features, we needed to expand the number of sequences

that exhibit HP1a association and differentiate them from those that lack such function. We

assessed diverse HAC candidate sequences, surveying known HP1a-interacting proteins for a

more degenerate pentamer sequence consisting of any hydrophobic amino acid at positions -2

and +2, and a central valine or isoleucine. The inclusion of isoleucine is based on sequence

alignment of Dmt homologs, some of which were found to contain this residue instead of valine

(**Supplementary Figure 2B**). We also focused our search on HP1a interactors that have been

partially mapped for HP1a-interacting regions (Pindyurin *et al.*, 2008; Schwendemann *et al.*,

2008). Twenty putative HACs were identified in 11 proteins (**Figure 1C**), including HP2 and

Odj that were already known to have at least one functional HAC. To test for HAC

functionality, we employed an *in vivo*, imaging-based approach to screen 25-amino acid

peptides from the HAC candidates for enrichment within the heterochromatin subnuclear

compartment, as marked by fluorescently tagged HP1a. Similarly-sized peptides were

previously shown to bind HP1a *in vitro* (Mendez *et al.*, 2011), and the relatively short

sequences minimize the possibility of including additional domains that may confound results.

Thus, this assay indirectly tests HP1a binding through the ability of HAC-containing peptides

to localize to the HP1a-rich heterochromatin, but also exposes candidate sequences to

physiological conditions not possible with *in vitro* binding assays.

For our experiment, we tested peptides containing our known and candidate HACs, comprised

of the hydrophobic pentameric core containing a PxVxL-like consensus plus 10-amino acid

sequences on each flank, by fusing them to a fluorescent tag (mNeonGreen) and a myc nuclear

localization signal. Co-transfection into Drosophila S2R+ cells with mScarlet-I-tagged HP1a

showed co-enrichment of HP1a with known HACs, indicating that HAC peptides are sufficient

to bind HP1a *in vivo* (**Supplementary Figure 1C**). In contrast, expression of the fluorescent

tag and the NLS without any HAC peptide failed to concentrate at HP1a-rich nuclear regions.

We also mutated the central HAC residue of several peptides, resulting in loss of enrichment

in heterochromatin (**Supplementary Figure 1C**). Based on these results, we conclude that the

peptide-based imaging approach is sufficient to detect sequences capable of associating with

HP1a, as measured by fold-enrichment within the heterochromatin domain over the total nuclear signal.

Next, we tested putative HACs from several heterochromatin-interacting proteins and identified sequences from SuUR (Pindyurin *et al.*, 2008), Lhr (Brideau and Barbash, 2011), HP4 (Schwendemann *et al.*, 2008), Mu2 (Dronamraju and Mason, 2011), ADD1 (Alekseyenko *et al.*, 2014), Ciz1 (Alekseyenko *et al.*, 2014), HP2 (Stephens *et al.*, 2005), Odj (Kasinathan *et al.*, 2020), Mod (Perrin *et al.*, 1998), MED26 (Marr *et al.*, 2014), and Su(var)3-9 (Schotta *et al.*, 2002) that did or did not accumulate in heterochromatin (**Figure 1D**). Heterochromatin-enriched peptides measured significantly higher levels of fluorescence intensity at regions of high HP1a enrichment relative to expression of the fluorescent tag without a HAC sequence (**Figure 1E**). This expands the number of known HACs from 6 to 16, while eliminating 10 sequences as putative HACs, or at the very least, HACs with HP1a affinities too low to detect with this assay. Image analysis also revealed varying levels of HAC co-enrichment with heterochromatin, with XNP displaying the highest co-enrichment with HP1a, and Piwi exhibiting the lowest. Mutation of the central residue of 4 of these peptides severely impaired heterochromatin localization, confirming that these comprise functional HAC sequences (**Supplementary Figure 1D**). We conclude that the *in vivo* peptide imaging approach can successfully differentiate varying levels of HP1a affinity among HACs, and that motifs resembling a PxVxL are frequently found in many proteins but poorly predict heterochromatin localization.

Finally, we determined whether the results of our peptide imaging experiment can be recapitulated with full-length proteins. Central residue mutations of HAC sequences in SuUR,

Lhr and Mu2 abrogated enrichment to the heterochromatin domain and resulted in lower

protein levels (**Supplementary Figure 1E-G**). This indicates that loss of binding with HP1a

destabilizes these proteins, as has been shown for other complex-forming proteins (Hsu, Yen

and Yeang, 2022). In contrast, corresponding mutations of non-functional sequences displayed

no effect on heterochromatin co-localization or protein levels. Meanwhile, mutations in both

the HAC and the H3K9me3-binding chromodomain (Wang *et al.*, 2012) were necessary to

abolish heterochromatin co-localization of Su(var)3-9 (**Supplementary Figure 1H**),

highlighting how other domains also play roles in heterochromatin enrichment. Finally, HP4

required the simultaneous mutation of all 3 central HAC residues to fully abrogate the

localization of HP4 to heterochromatin (**Supplementary Figure 1I**), confirming their

redundant functions to target HP1a. We therefore conclude that new HACs identified by

peptide localization correspond to functional heterochromatin localization domains in full-

length proteins.

Figure 1 (A) Alphafold-generated model of a HP1a CSD dimer and a peptide containing the PxVxL motif. Sidechains of PxVxL positions -2, 0, and +2 are shown in green. (B) Motif analysis of original peptide sequences identified by phage display (Smothers et al, 2000, top) and the 6 sequences previously identified as HP1a-binding in *Drosophila melanogaster* (bottom). Core PxVxL residues -2, 0, and +2 are marked. (C) Schematic diagram of HP1a-binding proteins and their corresponding known (yellow triangles) or putative (black triangles)

HACs. Numbers indicate positions of putative HACs on full-length proteins from N- to C-
terminus, except for known HACs, which are automatically numbered 1. (D) Schematic of
expression constructs mScarlet-I-tagged HP1a and mNeonGreen-tagged HACs (top) and
representative images of constructs after transient transfection into cultured S2R+ cells
(bottom). Scale bars = 1 um. (E) Fold enrichment of known and candidate HACs in
heterochromatin are calculated by the ratio of mean intensity in heterochromatin to the mean
intensity of the non-nucleolar nucleus. IDs of candidate HACs that fail to enrich in
heterochromatin are highlighted in blue; known HACs are marked in red, and candidate HACs
that display enrichment in heterochromatin are in orange. Also shown is the heterochromatin
enrichment of the mNeonGreen tag, marked with a diamond, which equals about 1 and thus is
used as the baseline to test significant enrichment of putative HACs by Student's t-test. *** =
p-value<0.001; ** = p-value<0.01; *=p-value<0.05. Determination of which candidate HACs
are enriched in heterochromatin was also assessed manually as some candidates that did not
visually display heterochromatin enrichment produced higher and lower enrichment values
than the mNeonGreen control, likely due to some peptides accumulating at the nucleolus or
being excluded from heterochromatin, respectively. HAC peptides tested for central residue
mutation effects in this study are marked with dotted circles; HACs shown to be functional in
the full-length protein are marked with empty circles.

Supplementary Figure 1 (A) Sequence alignment of 6 known PxVxL domains (highlighted in yellow) from published research with positions -2, 0, and +2 marked. (B) Venn diagram of proteins containing degenerate PxVxL consensus and their association with GO-annotations of nucleus/chromosome or cytoplasm components. Number of proteins within each group is shown as well as those containing the motif and annotated to be nuclear/chromosomal or cytoplasmic. (C) Representative images of mNeonGreen-tagged wildtype and mutant XNP HAC peptide in relation to mScarlet-I- tagged HP1a in S2R+ cells (top). Mutant XNP HAC has its central residue mutated from Val to Glu (V0E). Quantitation of fold-enrichment in

HP1a-rich heterochromatin condensates by XNP HAC constructs and a control containing only
the fluorescent tag (bottom). *** = p-value<0.001; ** = p-value<0.01. (D) Validation of
putative Lhr and Su(var)3-9 HACs by mutation of central Ile to Glu (I0E). Representative
images of full-length SuUR (E), full-length Lhr (F), and full-length Mu2 (G) with and without
V0E mutations at putative HAC sites. A longer exposure for mutant Mu2 is included. (H)
Representative images of full-length Su(var)3-9 with or without HAC and chromodomain (CD)
mutations. (I) Representative images of full-length HP4 with or without individual or
combinatorial HAC mutations. Scale bars = 1 um.

Drosophila HACs are conserved yet embedded in disordered regions.

In order to further define the sequence features that mediate HP1a binding and

heterochromatin-targeting, we investigated whether functional HACs are more evolutionarily-

conserved compared to non-functional peptides. Sequence alignment of HP1a CSD and C-

terminal tails for 12 Drosophila species across 40 million years of evolution show remarkable

conservation (91-100% identity in the CSD, 100% in the C-terminal tail) (**Supplementary**

Figure 2A). Sequence alignments of heterochromatin proteins harboring previously known

HACs and their homologs also show a high degree of conservation in terms of containing a

degenerate HAC pentamer sequence at the same position in the protein (**Figure 2A left,**

Supplementary Figure 2B). Kdm4A, Dmt, HP2, and Odj HACs are preserved across all 12

Drosophila species, though the Piwi HAC appears to be more recently evolved. Similar results

were obtained with our newly identified HACs, with 6 of 10 conserved across Drosophilids

(**Figure 2A middle, Supplementary Figure 2B**). In contrast, the non-functional candidate

sequences, as a group, were generally less conserved compared to functional ones (mean

conservation = 70% vs 87%) (**Figure 2A right, Supplementary Figure 2B**). However, the

non-functional candidate sequences with high conservation could be part of other domains.

The higher degree of conservation among functional HAC sequences suggests that they may

also share an ordered structure. However, disorder prediction analysis of the entire protein

sequence using PONDR (Romero *et al.*, 2001) showed that known HACs are either embedded

within disordered domains, or in transition regions between ordered and disordered regions

(**Figure 2B left**). Similarly, newly identified HACs showed a similar association with

disordered regions (**Figure 2B middle**). In contrast, the group of non-functional sequences

included both disordered and highly ordered structures (**Figure 2B right**). This indicates that

functional HACs, despite their localization in disordered regions, which are often associated

with higher mutation rates than globular domains, may be subjected to higher purifying

selection for conservation. This also suggests that HACs inherently occur in the more solvent-

accessible unstructured regions of proteins to facilitate binding to the HP1a CSD dimer.

Overall, our results indicate that high evolutionary conservation and association with

disordered domains are two shared features of HACs.

Figure 2 (A) Conservation of HAC sequences tested in this study among Drosophila species
spanning up to 40 million years of evolution (tree indicated on the left). Positive identification
was assessed by presence of [hydrophobic]x[V/I]x[hydrophobic] pattern after alignment to *D.*
melanogaster sequences among previously known HACs (red circles), as well as positively

identified HACs (yellow circles) and sequences that did not display HP1a co-enrichment (blue
circles) from Figure 1E. (B) Heat map showing disorder prediction per residue of HAC
sequences in (A) as obtained from the PONDR-VLXT algorithm. Prediction scores indicate
presence of intrinsic disorder (purple) to globular structures (white). Core HAC residues at -
12, -2, 0, +2, and +12 positions are indicated.

A.

```
                              Chromoshadow                              Tail
D. sechellia     GLEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSSVANEKIPRMVIHFYEERLSWYSDNED
D. simulans      GLEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSSVANEKIPRMVIHFYEERLSWYSDNED
D. melanogaster  GLEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSSVANEKIPRMVIHFYEERLSWYSDNED
D. yakuba        GLEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSSVANQKIPQMVIHFYEERLSWYSDNED
D. erecta        GLEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSSVANQKIPQMVIHFYEERLSWYSDNED
D. ananassae     GLEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSTVANVKIPQMVIHFYEERLSWYSDNED
D. pseudoobscura GQEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSSIANLKIPQMVIGFYEERLSWYSDNED
D. persimilis    GQEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSSIANLKIPQMVIGFYEERLSWYSDNED
D. willistoni    GLEAEKILGASDNNGRLTFLIQFKGVDQAEMVLSTVANVKIPQMVIRFYEERLSWYSDNED
D. mojavensis    GLEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSTVANVKIPQMVIRFYEERLSWYSDNED
D. virilis       GLEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSTVANVKIPQMVIRFYEERLSWYSDNED
D. grimshawi     GLEAEKILGASDNNGRLTFLIQFKGVDQAEMVPSTVANLAIPQMVIRFYEERLSWYSDNED
                 * ************************* *::** **.*** *************
                 |                                                          |
                 146                                                        206
```

B.

```
HP2:
Drosophila_mojavensis    AAVRPLPPLVPIMPAKLVQKEKDLQLPPQPLPPKLRKLRVRLNRSVIPNWLKSLK-----...PQVKIPIVNSVPL-PVPLPVLEVKP
Drosophila_virilis       TITKAPAPS--IPSEKPMQKPKELQLPSQPLPQKTRKLRVRLNRNMIKNWVLGLQ-----...QLLKAPVVRSMPL-PVPLPVLEVKA
Drosophila_grimshawi     AATATETTKQQCVLPAKHPTTNKLHLHLEPQLRTPKLRKLHVRLNRNCFKNWLKAHQ-----...----KSVPLPVPLPSL---------
Drosophila_willistoni    RVEVDIPQQ--------------QVQQNEMQLE------------------...LVEEVRTIRRCRVRLNLPPQVPPTR
Drosophila_pseudoobscura AVSSKSQPI--------------TKPKLRKCLVKINRGVIKKWMEA--------...PVAPVDKSDSVSVSVLAQP------
Drosophila_persimilis    PVSSKSQPI--------------TKPKLRKCLVKINRGVVKKWMEA--------...PVAPVDKSDSVSVSVLAQP------
Drosophila_ananassae     IATQPAKPV--------------QKEKKLCVKINRRAFNKWMEQQQGPQAT..---PAPVPAPQPTASAPDPKAPINP
Drosophila_melanogaster  RSTGNTPVQ--------------NISPRKLCVKINRRPYNKWLRSTQ--ERN..-----AKATLNTALCPSTEKHLPDEP
Drosophila_simulans      HSTGNTVPL--------------IVNSRKLCVKINRRPYNKWLRSTQ--ERN.......AKTTLKTGICSLTEQHLPDDP
Drosophila_sechellia     HSTGNIVPD--------------IVNPRKLCVKINRRPYNKWLRSTQ--ERN.......AKTTLKTGICSLTEQHLPDDP
Drosophila_erecta        HSTGNTVLQ--------------KAKLRKLCVKINRMPFHKWLRSTQ--E-N...PVLPAKTPLKTAIYPLTENQLPDES
Drosophila_yakuba        RTTGNIVPQ--------------KAKLRKLCVKMNRRPFNKWLRSTQ--EKN...VSPPAKTALKTAIYPLTEKKLPDET

KDM4A:
Drosophila_pseudoobscura TSTSSL-RLKQAVVKLRKLPQSAL-----------VS
Drosophila_persimilis    TSTSSL-RLKQAVVKLRKLPQSAM-----------VS
Drosophila_willistoni    SSSSSA-CLKKLVVKVRKLPPRDPEPAAVNSEASAIT
Drosophila_ananassae     SSYSSS-RQLQPVVKIRKLPQKPI-----------VP
Drosophila_yakuba        SSYNSC-RQLQPVVKLRKLPPTAS-----------VP
Drosophila_erecta        SSYNSC-RQLQPVVKLRKLPPTAS-----------VP
Drosophila_melanogaster  SSYSSC-RQLQPVVKLRKLPTIAS-----------VP
Drosophila_sechellia     SSYSSC-RQLQPVVKLRKLPTIAS-----------VP
Drosophila_simulans      SSYSSC-RQLQPVVKLRKLPTIAS-----------VP
Drosophila_grimshawi     SV-----NCKQPVVKLRKLPVATP--LTTQTANPQIE
Drosophila_virilis       SSRSSA-NLKQAVVKLRKLPTVVP---TAESAPVEAA
Drosophila_mojavensis    SSRSSCSNLKQAVVKLRKLPSTVT---------DAQP

XNP:
Drosophila_willistoni    -------------PIRKEKEQASSRENNRTYPTLKRCAVRLQRVKLPLT--------SEP
Drosophila_grimshawi     -----------------------------------AVGTLAAATTETATTEESATAATI
Drosophila_mojavensis    --------------------------------RRCVVRLKRVKWLTAAPISRETASSA
Drosophila_virilis       -------------------------------LRRCVVRLKRLKLPTA-----SAAEAG
Drosophila_persimilis    RPRRDLTLSKEKHPTKPSATSVLAQQQQQQQHRLKTCLVRVKRIKASEPEPRPSIKEPEP
Drosophila_pseudoobscura RPRRDSTLSKEKHPPKPSATSVLAQQQQQQQHRLKTCLVRVKRIKASEPEPRPSIKEAEP
Drosophila_ananassae     ------------TLSV----------------KRCLVRLKRVSL--------------
Drosophila_melanogaster  ------------PPRP----------------KRCVVRLKRVSL-------------P
Drosophila_simulans      ------------PPRP----------------KRCVVRLKRVSL-------------A
Drosophila_sechellia     ------------PPRP----------------KRCVVRLKRVSL-------------P
Drosophila_yakuba        ------------PPRP----------------KRCVVRLKRVSL-------------T
Drosophila_erecta        ------------PLRP----------------KRCVVRLKRVSL-------------P

Odj:
Drosophila_willistoni    ----PLTSNYPRVLIKRLPSPVKKI
Drosophila_grimshawi     ----TQSHLSPRVRIMRCRHPSEAL
Drosophila_mojavensis    ----SPRHGSPRVRLLRCRDPNNGR
Drosophila_virilis       ----SSSHLSPRVRLVRCRDPSTPR
Drosophila_persimilis    ----NPRSDYPRVVKRYRMPVATT
Drosophila_pseudoobscura ----NPRSDYPRVVVKRYRMPVATT
Drosophila_ananassae     ----KHNMQSPRVRLKRFHLPAKSE
Drosophila_yakuba        EAQKLPNSQSPRVRVKRLRVQEVEK
Drosophila_erecta        EAQQLPNSQSPRVRVKRLRVEEIEK
Drosophila_melanogaster  EDQNFPNRQSPRVRVKRLRVPVVEK
Drosophila_simulans      ETQNLSNSQSPRVRVKRLRVPVVEK
Drosophila_sechellia     ETQNLSNSQSPRVRVKRLRVPVVEK

PIWI:
Drosophila_grimshawi     --DVHQRSGQEPQSKIFRNQSDRHTPGDR----------DRSDVSSSRR-ASTS-----HG
Drosophila_mojavensis    --DISPESVQQPRPKIFRGDGFKSDDRDRSEASSSRGTSRSHAPSTSHAASTSYAATTLHA
Drosophila_virilis       --DRSPRSVQEPRAKVFRDNGPHNR--------------DRSEASRS------Y-----RG
Drosophila_willistoni    S-DEGRKRAEEPSAKFYKCHSDGSDPSRIPPKASSDAQPSSSSSSSRGRSFVPNE-----PR
Drosophila_melanogaster  S-STSRGSGDGPRVKVFRG---------------------SSSG------D-----PR
Drosophila_simulans      S-STSRGSGDGPRVKVFRA---------------------SSSG------G-----PR
Drosophila_sechellia     S-STSRGSGDGPRVKVFRA---------------------SSSG------D-----PR
Drosophila_yakuba        P-SNSRGSDDGPRVKVFRG---------------------SSSD------G-----PR
Drosophila_erecta        P-SNSGGSCDGPREKIFRA---------------------SSSD------G-----SR
Drosophila_ananassae     P-STSHEGREEPRAKIFKGPSDPRRE-----------------GSRG------D-----PR
Drosophila_persimilis    PGSSTDEKRDEPRAKVFKGQSEA--------------------STSG-------------
Drosophila_pseudoobscura PGSSTDEKRDEPRAKVFKGQSEA--------------------STSG--------------

Dmt:
Drosophila_grimshawi        NSQSRYRIRKFVVRIKRIK----------QQETIV
Drosophila_virilis          INQVNC-VRKCYVRLKRLK----------LPTNNA
Drosophila_mojavensis       KRNNLVRVRNCIVRIKRLKLNARDPTTYDVNKTLT
Drosophila_willistoni       VTNQMKSIKKCVVKIKRLK---------NLPKSHT
Drosophila_pseudoobscura    ANLPNATISPFRIRIKKLP----------VPK---
Drosophila_persimilis       ANLPNATISPFRIRIKKLP----------VPK---
Drosophila_ananassae        TPIP---LRKCVVKIRRLR----------VQENTP
Drosophila_melanogaster_Dmt LNEPECRIRKCVVKIRRIK----------VKNGVV
Drosophila_sechellia        PNVPECRLRKCVVKIRRIK----------VKNSVA
Drosophila_simulans         PNVPECRIRKCVVKIRRIK----------VKNAVA
Drosophila_yakuba           LTEPECRIRKCVVKLKRCK----------VQEGGA
Drosophila_erecta           LTEPERRIRKCVVKIRRIK----------VQKGVG
```

HP4:

```
Drosophila_willistoni      -MSQRSVQNKMKKMVVRIPRHPYLNLHGDHLLQ-----ENRVRKCRVNLERLKMS---PKQ--LKPKHVKVCVCRLPEYFFTERSE
Drosophila_pseudoobscura   ----MTPP-KIKKVVIKIPRHPYFAVQVDQGNM----VEYQLRKCRVNLERLKPPTPFALQLKPKPKRCIVRVARLPD---VAR--
Drosophila_persimilis      ----MTPP-KIKKVVIKIPRHPYFAVQVDQGNM----VEYQLRKCRVNLERLKPPTPFVLQLKPKPKRCIVRVARLPD---VAR--
Drosophila_ananassae       ----MSPKAKKMKMVVKIPRHPYFTVQSDSMQEHMRAVEYQVRKCRVNLERIKSSNAFPLK--PKPKRCIVRVARLPD---VERSE
Drosophila_melanogaster    ----MSP--KTKKMIVKIPRHPYFNVQGERSVE--REVEYQVRKCRVNLERIKSSTPFPLK--PKPKRCIVRVARLPD---VERSE
Drosophila_simulans        ----MSP--KTKKMIVKIPRHPYFNVQGDRSVE--REVEYQVRKCRVNLERIKSSTPFPLK--PKPKRCIVRVARLPD---VERSE
Drosophila_sechellia       ----MSP--KTKKMIVKIPRHPYFNVQGDRSVE--REVEYQVRKCRVNLERIKSSTPFPLK--PKPKRCIVRVARLPD---VERSE
Drosophila_erecta          ----MSP--KTKKMVVKIPRHPYFNVQVERSVE--REVEYQVRKCRVNLERIKSSNPFPLK--PKPKICIVRVARLPD---VERSE
Drosophila_yakuba          ----MLP--KTKKMVVKIPRHPYFNVQGEGSVE--REVEYQVRKCRVNLERIKSSTPFPLK--PKPKRCIVRVARLPD---VERSE
Drosophila_mojavensis      MLTKMSPQNKLKKMIVKLPRHPLIMANSDFVYQ----ECMQIRKCRVNLERIKGQTPFFPLK--SKPKRCCVRVARLPD---VERSE
Drosophila_grimshawi       ----MSHQNKIKKMIIKIPRHPFIIDQSNVNYQ----ENISLRKCRVNLERINGQQSFPLK--AKPKRCCVRVARLPN---VERNE
Drosophila_virilis         ----MSPQNKIKKMVIKIPRHPYIIAQGEFDYQ----ERVQIRKCRVNLERMKGQQPFPLK--PKPKRCCVRVARLPN---VERSE
```

Su(var)3-9:

```
Drosophila_grimshawi       VRFKNELERNITIKLERLSESKIQK
Drosophila_mojavensis      VRFKNELERNITIKLERLSEMKIQK
Drosophila_virilis         VRFKNELERNITIKLERLSEKKIQK
Drosophila_willistoni      VRFKNELERNITIKLERLSEKKIQK
Drosophila_pseudoobscura   VRFKNELERNITIKLERLSEKKIKL
Drosophila_persimilis      VRFKNELERNITIKLERLSEKKIKL
Drosophila_ananassae       VRFKNELERNITIKLERLSEKKIQK
Drosophila_sechellia       VRFKNELERNITIKLERLSEKKIKK
Drosophila_simulans        VRFKNELERNITIKLERLSEKKIKK
Drosophila_melanogaster    VRFKNELERNITIKLERLSEKKIKN
Drosophila_yakuba          VRFKNELERNITIKLERLSEKKIEK
Drosophila_erecta          VRFKNELERNITIKLERLSEKKLKK
```

ADD1:

```
Drosophila_willistoni      KTTNKPNPALLQLEKQRLEELKNSDPKLKMKVVVKLRRAEEDFEVARKWMAQH-HKKHTL
Drosophila_mojavensis      KTTHKPNPALLQLERQRMDEMKRNDSKLKMKPIVRLVRAEEQFEVARKWVEKYGQKKSKV
Drosophila_grimshawi       KMTNKPNPALLHLERQRIEETKRNDAKLKRKFVVKLRRAEKDFDVARKWVEENNKLNNLN
Drosophila_virilis         KMTNKPNPALLHLERQRMEETKRNDAKLKMKVVVKLRRAEEDFEVARKWMDM--EHAKKL
Drosophila_pseudoobscura   KTTNKPNPALLQLERQRMEEIKHTDAKLKMKVRVVLRRAEEEFEVARKWAKNRDKQN---
Drosophila_persimilis      KTTNKPNPALLQLERQRMEEIKHTDAKLKMKVRVVLRRAEEEFEVARKWAKNRDKQN---
Drosophila_ananassae       KTTNKPNPALLQLERKRMEEMKKSDAKLKMKVVVKLRRAEDEFEVARLWAEEENKKRQQDS
Drosophila_melanogaster    KTTNKPNPALLQLERQRMEDMKKSDAKLKMKVMVKLKRAEDEFEVARKWVEDM-KQGQEL
Drosophila_sechellia       KTTNKPNPALLQLERQRMEDMKKSDAKLKMKVMVKLKRAEDEFEVARKWVEDM-KQKQEL
Drosophila_simulans        KTTNKPNPALLQLERQRMEDMKKSDAKLKMKVMVKLKRAEDEFEVARKWVEDM-KREQEL
Drosophila_yakuba          KTTNKPNPALLQLERQRMEDMKKFDAKLKMKVMVKLRRAEDEFEVARKWVEECRKKEQEL
Drosophila_erecta          KTTNKPNPALLQLERQRMEDMKKSDAKLKMKVMVKLKRAEDEFEVARKWVEECRKREQEL
```

Lhr:

```
Drosophila_ananassae       IKSAKYYIKPVRVRLKRLDLDDYIS
Drosophila_melanogaster    LKSAKYYIKRCRIRLKRVEIDDYLP
Drosophila_sechellia       LKSAKYYIKSCRIRVKRVEIDDYLP
Drosophila_simulans        LKSAKYYIKSCRIRLKRVEIDDYLP
Drosophila_yakuba          LKSAKYYIKSCRIRVKRVDFEDLLP
Drosophila_erecta          LKSAKYYIKSCRIRVKRVDFKDYVP
Drosophila_mojavensis      LDSVRLYVKKVRVRVKRLDLADYVR
Drosophila_pseudoobscura   IKSAKYYTKKLIVRVKRIDLDEYLP
Drosophila_persimilis      IKSAKYYTKKLIVRVKRIDLDEYLP
Drosophila_willistoni      IKSAKYYVKKVRVLMKRQNLDDFIP
Drosophila_virilis         LKSAKYYVKKVQVRLKRLDLADYMP
Drosophila_grimshawi       ------------------------
```

Ciz1:

```
Drosophila_melanogaster    KATKPGPASTKRNVLVSVKRTSAAAIKAATKS
Drosophila_simulans        KAAKPGPASTKRNVLVSVKRTSAAAIKAATKS
Drosophila_sechellia       KAAKPGPASTKRNVLVSVKRTSAAAIKAATKS
Drosophila_erecta          KAAKPGPASTKRNVLVSVKRTSAAAIKAATKS
Drosophila_yakuba          KAAKPGPASTKRNVLVSVKRTSAAAIKAATKS
Drosophila_persimilis      KVGKPGPASTKRNVLVNVKRTSAAAIKAATKS
Drosophila_pseudoobscura   KVGKPGPASTKRNVLVNVKRTSAAAIKAATKS
Drosophila_mojavensis      KANRPGPASKKRQVLVSLKRTSAAAIKAATNS
Drosophila_virilis         KANRPGPASTKRQVLVSLKRTSAATIKAATKS
Drosophila_ananassae       KAAKPGPASTKRNVVVSVKRTSNATIKAATKS
Drosophila_grimshawi       ATSRPGPASTKRQVLVNGKRTPAAIIKAATKT
Drosophila_willistoni      KGAKPGPASTKRNVVVNVKRT--VAIKTGNKS
```

```
SuUR:

Drosophila_grimshawi      PS-----------------SPSKIFA-RKLNVKINRLHLPKMQKNVLAAISSKIPTNQLTKPAPAPREQPTPLPVKSPVKAKPMPEPN
Drosophila_mojavensis     PNTST-------------ITRKVLNVRL---RRASIQID----------------------PTNSAKASELPTKAVKASPKKSPNS
Drosophila_virilis        RPTPKIKGKKLNVRLSRVCLPQK----DLNIIAVTNSHKPKEQAATNSHKPKEQASKPGQEVSKKPPEEITPLEIKLLRAKPNNKVKL
Drosophila_willistoni     METVK-------------HPSP----VL--------------------------------A----TRNVNIRLRRT--------
Drosophila_pseudoobscura  DSSRQ---------KTRS----TP---AAAAVSPT----------------------EK----TRKVNVRLRRM--------
Drosophila_persimilis     DSSRQ---------KTRS----TPAAAAAAVSPT----------------------EK----TRKVNVRLRRM--------
Drosophila_ananassae      EKVDN---------NPKP----IT---PPKKVATP----------------------KK----AMKLNIRLKR---------
Drosophila_melanogaster   ATDNM---------PPRK----RT---RAATVHLT----------------------PK----TRRLNVRILRV--------
Drosophila_sechellia      AMDNM---------PPKK----RT---RAATVHLT----------------------PK----TRRLNVRILRC--------
Drosophila_simulans       A---M---------PPRK----RT---RAATVHLT----------------------PK----TRRLNVRILRC--------
Drosophila_yakuba         ALT-I---------SPRK----RT---RAATVHFT----------------------PK----TRRLNVRILRV--------
Drosophila_erecta         GLT-M---------PPRK----ST---RAATVHFT----------------------PK----TRRLNVRLRV--------
                                                           :  .

Mu2:

Drosophila_pseudoobscura  --------KIGNTPEEGVVEKIASKRVGRPRK...NANTSAVAKRRVGRPRKSQTPTEDK
Drosophila_persimilis     --------KIGNTPEEGVVEKIASKRVGRPRK...NANTSAVAKRRVGRPRKSQTPTEDK
Drosophila_grimshawi      -----------------------------...NASSKRAARVR--------------
Drosophila_mojavensis     -----------------------------...KAGSARAATRKSDSAKPTLKNVEDT
Drosophila_virilis        K----------------------------...PAPSLRSFRKSTSKTDLTEETKEDI
Drosophila_willistoni     R---------------------------S...SEEKDVKGRKLKVNVKNITSQGNKQ
Drosophila_ananassae      TKKAKESSKAEQPKIQKRVTRLNSANESVESH...GSTKLGPRKGSALPGSNEEEPLDAK
Drosophila_melanogaster   V-------KTGGALIKRAVVRISRVSIEQLES...PSSSSAAAKEPNVSTATTR------
Drosophila_simulans       V-------KVTGALIKRAVVRISRVSIEQLEA...PSSSSAAAKGPNVSTATTRATRTTR
Drosophila_sechellia      V-------KAAGALIKRAVVRISRVSIEQLES...PSSSSAAAKGPNVSTAITRTTRTTR
Drosophila_yakuba         V-------KPADALIKRAVVRISRASIENPAS...PSSSTAAAKGPNVS-ATTRAPKTVR
Drosophila_erecta         V-------KPEGAIFKRAVVRISKISIENPES...PSSSSAAAKGPNVS-GTTRTARSAR

HP2:

Drosophila_mojavensis     STC---S------------ ...--------SNSSCRKLRILIK------- ...LEDPTYPSDQSVC---------HDVGDKA
Drosophila_virilis        SSSSSGSSRRKLRILLRKV ...ALEQISVDNPLDNRGDLVVNEDSVDQA ...PTGQV--------------------EVN
Drosophila_grimshawi      ------------------ ...-----------------------... .------------------------EKP
Drosophila_willistoni     ------------------ ...EGKISENSTIEKKRLR----------- ...KKGRQNDNESEI----LVQAESTQEIA
Drosophila_pseudoobscura  SEA---TELEGPPSTTPTL ...ESKQTSESNPSSEKSK---EPQPEKQP ...PSVPITPCTAPCRQLRVLMRRTPSSKSSV
Drosophila_persimilis     SEA---TELEGPPSTKPTL ...ESKQTSESNPSSEKSK---EPQPETQP ...PSVPITPCTAPCRQLRVLMRRTPSSKSSV
Drosophila_ananassae      --------------TPQK ...--------TPSCRKLRILIKRTPNTKY ...ESPDVPPTATEMEEPPQTSLDIKPPDDTE
Drosophila_melanogaster   TKS---SDQTKNSFITPNR ...VQKELRTPTASCRKLRVLIKRTPTSSL ...SLGEVNPDSDPVAAES--VAVLHES-DRD
Drosophila_simulans       TKS---SDQT-NSFITPIR ...EQKELRTPTASCRKLRVLIKRTPTSNL ...SSDEINPDSDPVAAESLTVAILHDS-DRD
Drosophila_sechellia      TKS---SDQT-NSFITPNR ...EQNELRTPTASCRKVRVLIKRTPTSNL ...SSDEINPDSDPVAGESLPVAILHDS-DRD
Drosophila_erecta         TKS---SDQTKNTFITPNR ...LEKELGTPTASCRKLRVLIKRTPTSNL ...SSGEINPESDP--------------DKD
Drosophila_yakuba         TKS---SDQTKNTIITPKK ...IQKELSSPTASCRKLRVLIKRTPTSDL ...SSGVLNPESDPVADL---LPVAVTESDKE
```

Supplementary Figure 2: (A) Sequence alignment of HP1a CSD (highlighted in blue) and
CTE (highlighted in yellow) for 12 Drosophila species. Divergent residues are marked in red.
(B) Alignments of HP1a-binding proteins containing HACs across 12 Drosophila species.
Pentameric core domains of HACs are highlighted in blue; HACs tested to accumulate in
heterochromatin are highlighted in red. Some HAC sequences were manually curated as
alignments of SuUR, Mu2, and HP2 homologs did not align with the Drosophila melanogaster
HAC sequences and may represent paralogous HACs.

The pentameric core requires flanking residues for heterochromatin localization

To identify the amino acids and chemical interactions responsible for binding of HACs with

HP1a dimers, we performed full atomistic molecular dynamics (MD) simulations on 25-amino

acid HAC sequences containing the hydrophobic core at the center. We confirmed that 15 of

the 16 HACs showed high prediction accuracy with pLDDT (predicted Local Distance

Difference Test) (**Supplementary Method Table 1**), and the expected positioning of their core

region at the center of the CSD dimer in AlphaFold models and throughout the 250 ns MD

simulations (**Figure 3A, Supplementary Figure 3A-O**). To investigate whether the core

region is sufficient to explain the variable fold-enrichment of HACs in heterochromatin

(**Figure 1E**), we compared the root mean square deviation (RMSD) of HAC sequences from -

2 to +2 in the last 50 ns of the MD simulations (**Figure 3B**). Though Piwi, with the lowest fold

enrichment score, showed very high fluctuation, the RMSD of the core regions of other

peptides showed poor correlation with the ranking of HP1a co-enrichment scores from

experimental observations (**Figures 1E, 3B,** $R^2 = 0.07$). To expand and elucidate key residues

of HACs, the root mean square fluctuation (RMSF) was calculated for Piwi and XNP, which

had the lowest and highest fold enrichment scores from the peptide imaging assay, respectively

(**Figure 3C**). XNP showed overall lower RMSF values than Piwi as well as a longer region of

low RMSF values ($\leq$RMSF values of the Piwi core region, dotted horizontal lines, **Figure 3C**),

consistent with the stronger enrichment of the XNP HAC with HP1a in cells (**Figure 1E**). As

expected, the core regions (-2 to +2) exhibited the lowest RMSF values, indicating the part of

the HAC with the most stable association with the CSD binding pocket (**Figure 3C**).

Interestingly, the proximal flanking sequences on both sides (-7 to -3 and +3 to +7) also showed

low RMSF values and no significant difference from the core regions of both XNP and Piwi

(**Figure 3C**). While Piwi showed no significant RMSF difference between the proximal

flanking sequences and distal flanking sequences (-12 to -8 and +8 to +12), XNP had a

significant difference between the proximal and distal flanking sequences (**Figure 3C**). These

RMSF results from the MD simulations suggest the potentially crucial role of the proximal

flanking regions in determining the strength of the HP1a-peptide interactions and accessibility

to heterochromatin.

We tested different lengths of flanking sequences required for the central core of HACs to

maintain heterochromatin enrichment *in vivo*. Lack of any flanking sequences abrogated the

ability of three different HACs to localize to heterochromatin at levels indistinguishable from

negative controls (**Figure 3D**, **Supplementary Figure 3P and Q**). In contrast, the retention of

5 amino acids flanking each side of the central pentamer maintained enrichment with HP1a,

indicating that flanking residues are crucial to HAC association with HP1a (**Figure 3D,**

Supplementary Figure 3P). In conclusion, both MD simulations and *in vivo* peptide

localization demonstrate that the traditional PxVxL motif is insufficient for heterochromatin

enrichment, and additional residues within 5 amino acids on both sides of the hydrophobic core

are required for HAC function.

Figure 3 (A) Visualization of the CSD dimer with the XNP HAC. Left panel: side view. Right Panel: Zoomed top view. Each CSD monomer is colored cyan and purple with a green and blue colored C-terminal tail, respectively. XNP peptide is colored yellow. Residues of the central hydrophobic core at the -2, 0, and +2 positions (in licorice drawing style) are colored orange. (B) Root mean square deviation (RMSD) ± SD [nm] of the core sequences from -2 to +2 of 15 HACs between 200 to 250 ns of the MD simulations. The peptides are ranked according to the experimental ranking of fold HP1a co-enrichment from Piwi, the lowest, to XNP, the strongest, based on results from Figure 1E. The x-axis values for the linear regression calculation are assigned from 1 (Piwi) to 15 (XNP). The R-squared value is shown on the graph. (C) Left: Root mean square fluctuation (RMSF) [nm] of 25-residue for XNP and Piwi HAC sequences (-12 to +12) between 200 to 250 ns of the MD simulations. The gray dash line at the -2 and +2 is drawn to indicate the core pentamer. The blue and red dashed lines indicate the span of sequences that consecutively display RMSF values less than the combined average of Piwi and XNP = 0.32 nm. Right: Averaged root mean square fluctuation (RMSF) ± SD [nm] of the XNP and Piwi distal flanking sequences (positions -12 to -8 and +8 to +12), proximal flanking sequences (positions -7 to -3 and +3 to +7), and core sequences (positions -2 to +2). Asterisks denote p-values < 0.05 (*) (D) Left: Representative localization images of XNP HAC constructs with different amounts of flanking residues around the hydrophobic core as illustrated in the right panel. As controls, fluorescent tags with different lengths of linker regions were also tested. Right: Quantitation of HP1a co-enrichment at heterochromatin domains for the tested HAC constructs. N/S = not significant; *** = p-value < 0.001.

396
Supplementary Figure 3: (A-O) The Alphafold Multimer-predicted computational model
(left) and visualized MD simulation image of the last frame at 250 ns (right) for the CSD dimer
and 15 peptides. The Alphafold Multimer-predicted structure is colored based on pLDDT
(predicted local distance difference test) scores. Red: pLDDT<50, Orange: 50<pLDDT<60,
Yellow: 60<pLDDT<70, Green: 70<pLDDT<80, Skyblue: 80<pLDDT<90, and Blue:
90<pLDDT. CSD homodimer and the peptide are colored cyan and yellow, respectively.
Residues at the -2, 0, and +2 positions (in licorice drawing style) are colored orange, with the
amino acid code indicated. (P-Q) Quantitation of HP1a co-enrichment at heterochromatin

domains for the tested KDM4A and HP2-1 HAC constructs, respectively. 25AA = full-length
native sequence containing core region surrounded by 10 residues on each flank; 15AA = core
region surrounded by 5 residues on each side; 5AA = core region only. *** = p-value < 0.001;
** = p-value < 0.01; * = p-value < 0.05.

Electrostatic forces surrounding the hydrophobic pocket drive HAC enrichment to

heterochromatin

To identify the driving force of the interactions between HACs and HP1a, the types of amino

acids in HACs and their molecular properties were analyzed *in silico*. Sequence alignment of

all 16 HACs displayed high frequencies of hydrophobic amino acids at positions -7, -5, -1, +5,

and +7 in addition to the core hydrophobic residues (-2, 0, and +2) (**Figure 4A**). Amino acids

with positively charged side chains (Lys and Arg) are enriched in the flanks and interior of the

central hydrophobic core (-4, -3, +1, +3, and +4) when compared to non-HAC peptides (**Figure**

4A, Supplementary Figure 4A). In contrast, negatively charged residues (Asp and Glu) are

relatively under-represented among HAC peptides. The sequence alignment analysis thus

suggests that hydrophobic and positively charged amino acids contribute significantly to HP1a

interactions at specific positions of HACs.

We first investigated the effects of the hydrophobic properties of HACs using MD simulations.

The hydrophobic pocket formed by the central pentamer of HACs and the HP1 CSD dimer was

previously identified to be the main interaction region. The flanking hydrophobic residues,

such as -5 and +5, were also thought to contribute to hydrophobic interactions (Thiru *et al.*,

2004). We compared the hydrophilic surface area of the HAC sequences from -7 to +7 obtained

from the MD analysis with the experimental ranking of HP1a co-enrichment (**Figure 4B, 1E**).

Surprisingly, despite some variations, the hydrophilic surface area values did not correlate with

the fold enrichment ranking (**Figure 4B, Supplementary Figure 4B,** R^2 = 0.14). The

hydrophilic surface area of only the HAC central pentamer from -2 to +2 also showed no

correlation with the fold enrichment scores (**Supplementary Figure 4D,** R^2 = 0). Comparing

the hydrophilic surface area frequencies of the top two and bottom two ranking peptides

showed that Piwi had the highest and lowest hydrophilic surface area in the HAC core pentamer

(-2 to +2) and the proximal flanking region (-7 to +7), respectively (**Supplementary Figure**

4C, E). This indicates that Piwi possesses the lowest hydrophobic interaction between the CSD

dimer and the HAC core pentamer, but the highest hydrophobicity between the CSD and the

proximal flanking sequence (**Supplementary Figure 4C, E**). However, Su(var)3-9 and Dmt,

which had the second lowest and second highest fold enrichment scores, respectively, did not

show a corresponding trend of hydrophilic surface area values in relation to Piwi and XNP

(**Supplementary Figure 4C, E**).

To investigate how the hydrophilicity and hydrophobicity of each residue on the HAC peptide

participate in the interaction, we visualized the hydrophilic and hydrophobic distribution

properties of the HP1a-interacting XNP regions (**Figure 4C**). As expected, there is a

hydrophobic pocket between the -2 to +2 amino acid sequences of the HAC and HP1 CSD

dimer (**Figure 4C**). On the contrary, the nearby flanking residues tend to have high

hydrophilicity rather than hydrophobicity (**Figure 4C**). The hydrophobic pocket of the peptide

and **α**-helices of the CSD dimer were surrounded by a high hydrophilic region composed of -

4, -3, +1, +3, and +4 positions, indicating the possible existence of other types of interactions

that could modulate interactional strength with HP1a. These results also suggest that the

hydrophobic interactions of the HACs alone are insufficient to describe the fold enrichment of

HAC-containing peptides with HP1a.

Given the lack of correlation between the hydrophobic properties and strength of HP1a co-

localization *in vivo*, we then investigated the electrostatic contributions of the core and flanking

amino acids using MD simulations. Electrostatic interactions, which include salt bridges, pi-

cation interactions, and hydrogen bonding, represent another key type of interaction in protein

structure and binding (Zhou and Pang, 2018). Salt bridge interactions, in particular, are known

to be one of the most robust forms of residue-residue interactions (Xie *et al.*, 2015). Salt bridge

interactions form when the distance between the side-chain carbonyl oxygen of Asp or Glu and

the side-chain nitrogen atoms of Arg, Lys, or His is within 0.4 nm, facilitated by Coulomb

attraction with water molecules (Kumar and Nussinov, 2002; Pylaeva, Brehm and Sebastiani,

2018). The high hydrophilic and positively charged amino acid-conserved regions were seen

at -4, -3, +1, +3, and +4 (**Figure 4A, C**). Therefore, we calculated Coulomb energy levels

between the HACs from +4 to +4 and the HP1a CSD dimer with C-terminal tails. Notably,

Coulomb energy levels generally decreased among peptides ranked from low to high HP1a co-

enrichment, indicating that attractive electrostatic force followed the trend of the *in vivo* fold

enrichment ranking (**Figure 4D, Supplementary Figure F**, R^2 = 0.46). Comparing the

electrostatic energy of the top two and bottom two ranking peptides similarly showed a

correlation between Coulomb energy levels and heterochromatin enrichment (**Supplementary**

Figure G). The Coulomb energy between the HACs from -7 to +7 and the HP1a CSD dimer

with C-terminal tails showed less correlation (**Supplementary Figure H, I**, R^2 = 0.28),

implying the -4 to +4 residues of HACs have more important contributions to the fold

enrichment scores. The visualization of the electrical charge distribution of the HP1a-

interacting XNP regions showed a high charge density at -4, -3, +1, +3, and +4 amino acids

outside the hydrophobic pocket of the CSD, but did not show a charge density outside -4 and

+4 regions, emphasizing the importance of electrically charged amino acids between -4 and +4

in the electrostatic interactions with HP1a (**Figure 4E**).

Overall, our observations indicate that electrostatic interactions between the HAC residues

from -4 to +4 and the HP1a CSD dimer with C-terminal tails are crucial determinants in the

binding of HAC-containing peptides with HP1a dimers. Of course, hydrophobic and

electrostatic residues beyond the ranges of positions analyzed here may also contribute to HP1a

binding, but these positions, such as distant flanking sequences (-12 to -8 and +8 to +12), are

more variable in residue content and experience more fluctuation (as shown by RMSF variation

in Figure 3).

A.

B.

C.

D.

E.

Figure 4 (A) Frequency of occupancy by hydrophobic (pink), positively charged (blue), and negatively charged (red) select groups of amino acids at each position among tested HACs (top) and non-HACs (bottom). Numbering of positions are in relation to the central residue at position 0. (B) Hydrophilic surface area $\pm$ SD [nm^2] of sequences from -7 to +7 of 15 HACs between 200 to 250 ns of the MD simulations. The peptides are ranked according to the experimental ranking of fold HP1a co-enrichment from Piwi, the lowest peptide, to XNP, the strongest peptide. The x-axis values for the linear regression calculation are assigned from 1 (Piwi) to 15 (XNP). The R-squared value is shown. (C) Visualizations of XNP HAC surface hydrophilicity and hydrophobicity in the side view (above) and the top view (bottom) were calculated using the Discovery Studio software. The hydrophilic to hydrophobic properties are colored from green to pink. XNP backbone is colored yellow. (D) Coulomb energy $\pm$ SD [kJ/mol] of 15 HAC sequences from -4 to +4 between 200 to 250 ns of the MD simulations. A ranking similar to that of (A) is used to order the peptides. The R-squared value is shown. (E) Visualization of the XNP HAC surface charge in the side view (above) and the top view (bottom). The negatively to positively charged properties are colored from red to blue. XNP is colored yellow.

Supplementary Figure 4: (A) Bar graphs showing differences in percent frequency of positively charged residues (left in blue), negatively charged residues (center in red) and hydrophobic residues (right in pink) between HAC peptides and non-HACs. (B) Hydrophilic surface area of sequences from -7 to +7 of 15 HACs between 200 to 250 ns of the MD simulations are plotted according to the fold HP1a co-enrichment values (Figure 1E). The R-squared value is shown. (C) Histogram showing the percentage frequency of hydrophilic surface area for sequences from -7 to +7 of HACs for Piwi, Su(var)3-9, Dmt, and XNP between 200 to 250 ns of the MD simulations. Piwi: Blue with a dash line. Su(var)3-9: Green with a dotted line. Dmt: Pink with a dash and dotted line. XNP: Orange with a solid line. Histogram was fitted by kernel density estimation (KDE). (D) Hydrophilic surface area ± SD [nm^2] of sequences from -2 to +2 of 15 HACs between 200 to 250 ns of the MD simulations. The peptides are plotted according to ranking in terms of fold HP1a co-enrichment. The R-squared value is shown. (E) Histogram showing the percentage frequency of hydrophilic surface area for pentameric core sequences from -2 to +2 of HACs for Piwi, Su(var)3-9, Dmt, and XNP between 200 to 250 ns of the MD simulations, using the same format as C. (F) Coulomb energy of 15 HAC sequences from -4 to +4 between 200 to 250 ns of the MD simulations are plotted according to HP1a co-enrichment values. The R-squared value is shown. (G) Histogram showing the percentage frequency of Coulomb energy for 9 residues from -4 to +4 of HACs for Piwi, Su(var)3-9, Dmt, and XNP between 200 to 250 ns of the MD simulations, using the same format as C. (H) Coulomb energy ± SD [kJ/mol] of 15 HAC sequences from -7 to +7 between 200 to 250 ns of the MD simulations. The peptides are plotted according to ranking in terms of fold HP1a co-enrichment. The R-squared value is shown. (I) Histogram showing the percentage frequency of Coulomb energy from positions -7 to +7 of Piwi, Su(var)3-9, Dmt, and XNP HACs between 200 to 250 ns of the MD simulations, using the same format as C.

Distinct roles of positively charged amino acids in HAC binding

To reveal the molecular roles of positively charged amino acids in the HAC peptide binding and structure, we analyzed the individual Coulomb energy from positions -7 to +7 for several HACs in association with the HP1a CSD dimer and C-termini in the MD simulations (**Figure 5A, C**). Three peptides were chosen for the electrostatic analysis: XNP as the strongest interacting peptide, Piwi as the weakest interacting peptide lacking a positively charged amino acid at the +3 position, and Ciz1 as the relatively weak interacting peptide lacking a positive amino acid at the +1 position. **Table 1** shows the quantitative values of Coulomb energy for -7 to +7 sequences of these three peptides. Overall, -4, -3, +3, +4 HAC amino acids had higher

attractive Coulomb energy than -7 to -5 and +5 to +7 regions, quantitatively confirming the

important contributions of -4 to +4 amino acids suggested in Figure 4.

The highest attractive energy to the CSD dimer was observed at the +4 position for all three

peptides (**Table 1, Figure 5A**), where a salt bridge interaction with Asp157^{II} (of CSD^{II}) was

detected (**Figure 5B, Supplementary Figure 5A**). Similarly, the -4 position of the XNP and

Ciz1 HACs formed a salt bridge interaction with Asp157^{I}, but was not present for Piwi due to

the lack of a positively charged amino acid at the -4 position (**Figure 5A**). All 16 peptides that

exhibited heterochromatin enrichment have a positively charged amino acid at the +4 position

(**Figure 4A**), suggesting the +4 positively charged amino acid comprises a basic feature of the

HAC motif.

In contrast, the residues with the highest attractive Coulomb energy with the HP1a C-terminus

varied among the three peptides (**Table 1**). The Coulomb energy at the +3 position displayed

high variance among the three peptides (-164±70.42, -47.39±57.50, and -3.01±5.15 [kJ/mol]

for XNP, Ciz1, and Piwi, respectively). Unlike the other two, Piwi lacks a positively charged

amino acid at the +3 position. Interestingly, there is also a discrepancy between XNP and Ciz1,

even though both XNP and Ciz1 have lysine at the +3 position. Molecular interaction analysis

showed that the Arg+3 of XNP formed a salt bridge or H-bond with HP1a Glu205^{I} or Asp203^{I}

(**Figure 5D, Supplementary Figure 5B**). However, Arg+3 of Ciz1 did not form such

interactions and instead had an unfavorable interaction with HP1a Arg197^{II} (**Figure 5D, F,**

Supplementary Figure 5C). This difference is attributed to interactions between +1 of the

HACs and Arg197^{II} of the CSD: Arg+1 of XNP formed a maximum of three H-bonds with

Arg197[II], including its side chain (**Figure 5E, Supplementary 5C**), while Ser+1 of Ciz1 had

only one H-bond with the carboxylic acid group of Arg197[II], leading to a repulsive interaction

between the Arg197[II] side chains and Lys+3, and a H-bond between Arg197[II] and Glu205[I]

(**Figure 5F, Supplementary 5C**). These results indicate that Arg+1 plays an important role in

both the H-bond interaction with Arg197[II] and the maintenance of the electrostatic force field

to develop the salt bridge and H-bond among +3, Asp203[I], and Glu203[I]. We therefore tested

the effects of a Ser+1Arg Ciz1 mutation *in vivo* to determine if increasing electrostatic

interactions at this position will enhance HP1a co-enrichment. The Ciz1 mutant modestly

improved the fold enrichment score compared to a wildtype Ciz1 HAC, indicating the

deterministic effect of +1 in controlling the strength of the HP1a and HAC peptide interaction

(**Figure 5G, Supplementary Figure 5D**).

As previously noted, Piwi lacks a positively charged amino acid at +3, but our analysis showed

that the Lys+1 side chain can form an H-bond with Ser199[I] or salt bridge interaction with

Asp203[II] (**Figure 5H, Supplementary Figure 5E**). A positively charged amino acid at position

+1 may therefore compensate for the lack of a positively charge at +3 by forming a salt bridge

or hydrogen bond to attain sufficient HP1a interaction strength. There are likely many other

compensatory mechanisms for different HACs, as positively charged residues occupy variable

positions within and surrounding the hydrophobic core (**Supplementary Figure 5F**).

To validate the computational theory of the electrostatic effects of positive amino acids on

HAC peptide and HP1 interactions, we performed single and combinatorial amino acid

mutation experiments with XNP, which had the highest Coulomb energy for the C-terminus

and CSD dimer (**Table 1, Figure 5A, C**). First, single mutation experiments showed that

Arg+1Ala and Arg+4Ala mutations significantly decreased the fold enrichment in

heterochromatin (**Figure 5I, Supplementary Figure 5G**). Although high electrostatic energy

was observed for Arg+3 in the MD simulation, the Arg+3Ala mutation did not show a

significant difference in HP1a co-enrichment. The compensatory mechanism of the Piwi Arg+1

side chain for the lack of a positively charged amino acid at +3 (**Figure 5H**) might explain the

lack of effect of Arg+3Ala in the XNP experiment. Nevertheless, the double mutations of

Lys+3 and Arg+4, as well as double mutations of Lys-4 and Arg-3, greatly diminished XNP

HAC enrichment in heterochromatin compared to the single mutants. Furthermore, a quadruple

mutation of -4, -3, +3, and +4 nearly abrogated all enrichment, suggesting the basic residues

act cooperatively to stabilize HAC enrichment to heterochromatin (**Figure 5I, Supplementary**

Figure 5H). These results show that positively charged amino acids immediately flanking the

pentameric core are essential for HAC function. Further, they suggest that the interaction

strength between the HAC and HP1a can be controlled by selecting appropriate mutations of

positively charged amino acids in the HAC sequence.

To obtain a comprehensive view of the molecular forces that govern HAC binding to HP1a,

we analyzed the XNP HAC for electrostatic and hydrophobic interactions in simulations

(**Figure 5J, Supplementary Figure 5I**). Limiting the scope of analysis to between HAC

residues -7 and +7, we identified salt bridges forming at positions -4, +3, and +4 with both the

CSD and C-terminal tail, as previously discussed. We also detected pi-cation interactions at

the basic positions -4, +1, and +4 with the hydrophobic pocket residues of the CSD dimer.

Meanwhile, hydrophobic interactions were predicted at positions -5, -2, -1, 0, +2, and +5,

consistent with previous studies. Hydrogen bonds were also formed by both nonpolar and basic

amino acids at 9 of the 15 residues analyzed. Of important note is the hydrogen bonding

detected at position -2 unique to cysteine, due to its polar, hydrophobic sidechain. This confers

an extra electrostatic interaction absent from other hydrophobic residues that occupy that

position in other HACs, and thus provides higher affinity for HP1a. We predict that other HACs

follow a general pattern of molecular interactions contained in this XNP map, though

alternative interactions are likely to occur due to the positional variations of basic and

hydrophobic residues of other HACs that impact HP1a affinity.

Table 1 Sequences and coulomb energy ± SD [kJ/mol] among the CSD dimer or CTE and -7
to +7 residues of XNP, Ciz1, and Piwi. The highest coulomb energy among the -7 to +7 is
highlighted in red.

Residue	XNP	Ciz1	Piwi	CSD_XNP	CSD_Ciz1	CSD_Piwi	C-terminus_XNP	C-terminus_Ciz1	C-terminus_Piwi
-7	P	S	G	-1.02 ± 4.99	0.17 ± 0.89	-7.34 ± 8.71	-0.04 ± 0.13	-2.79 ± 9.95	0.02 ± 2.09
-6	R	T	S	-9.27 ± 11.92	-1.94 ± 6.99	-0.89 ± 2.73	-46.09 ± 62.83	-20.22 ± 19.17	-5.35 ± 13.21
-5	P	K	G	-5.45 ± 2.38	-2.55 ± 4.81	0.09 ± 0.55	0.60 ± 1.37	-41.43 ± 39.44	-0.10 ± 1.04
-4	K	R	D	-105.93 ± 48.80	-90.84 ± 59.05	1.06 ± 5.62	-49.02 ± 59.89	-40.17 ± 16.80	0.00 ± 7.94
-3	R	N	G	-4.58 ± 5.44	-8.29 ± 2.93	1.32 ± 4.32	-50.86 ± 45.06	-59.87 ± 18.65	0.23 ± 0.22
-2	C	V	P	-59.14 ± 10.41	-14.00 ± 3.01	-14.79 ± 9.36	-0.84 ± 1.17	-0.58 ± 2.31	-0.01 ± 0.05
-1	V	L	R	-41.48 ± 9.21	-38.41 ± 7.45	-41.34 ± 12.06	-0.35 ± 0.75	-0.33 ± 0.62	-32.02 ± 43.60
0	V	V	V	-34.87 ± 7.38	-34.82 ± 9.73	-32.84 ± 9.36	-0.13 ± 0.27	-0.02 ± 0.03	0.04 ± 0.05
+1	R	S	K	-66.45 ± 25.07	-27.53 ± 9.84	-60.91 ± 32.91	2.45 ± 6.20	-0.19 ± 1.72	-32.08 ± 50.42
+2	L	V	V	-20.24 ± 5.45	-15.57 ± 5.31	-18.88 ± 7.60	-13.30 ± 4.62	-14.29 ± 4.19	-16.89 ± 4.62
+3	K	K	F	7.16 ± 5.26	4.82 ± 4.27	-21.33 ± 5.79	-161.04 ± 70.42	-47.46 ± 57.53	-3.01 ± 5.15
+4	R	R	R	-133.15 ± 21.40	-132.20 ± 21.07	-81.99 ± 52.34	-3.96 ± 9.15	-29.54 ± 23.95	-6.04 ± 11.63
+5	V	T	G	-16.68 ± 5.73	-18.45 ± 8.98	-15.14 ± 6.16	-0.05 ± 1.17	-12.96 ± 23.31	0.04 ± 0.38
+6	S	S	S	-0.93 ± 1.01	-0.69 ± 2.24	-1.21 ± 3.60	-6.33 ± 14.80	-18.59 ± 30.59	-0.11 ± 0.86
+7	L	A	S	1.49 ± 0.99	-0.37 ± 2.36	-1.34 ± 6.45	0.83 ± 2.43	-0.28 ± 1.13	-0.01 ± 0.23

Figure 5: (A) Coulomb energy ± SD [kJ/mol] between the CSD dimer and -7 to +7 residues of XNP, Ciz1, and Piwi. Orange circle: XNP, Cyan diamond: Ciz1, and Blue square: Piwi. (B) Visualization of the salt bridge interaction between Arg+4 and Asp157^{II}. Arg+4 and Asp157^{II} in the Corey-Pauling-Koltun (CPK) drawing style are colored yellow and purple, respectively.

(C) Coulomb energy ± SD [kJ/mol] between the HP1a C-terminal tails and -7 to +7 residues of XNP, Ciz1, and Piwi. Orange circle: XNP, Cyan diamond: Ciz1, and Blue square: Piwi. (D) Histograms showing the percentage frequency of Coulomb energy between HP1a Asp203[I]/Glu205[I] and +3 of XNP and Ciz1 between 200 to 250 ns of the MD simulations. The histogram was fitted by kernel density estimation (KDE). XNP: Orange with a solid line. Ciz1: Cyan with a dashed line. (E) Visualizations of XNP and (F) Ciz1 with blue, orange, and red circles with arrows indicating a hydrogen bond, salt bridge, and unfavorable interactions, respectively. (G) Quantification of HP1a co-enrichment of Ciz1 HAC and mutations either disabling the central valine or adding a positive charge at the +1 position. (H) Visualization of H-bond interaction between Piwi Lys+1 and Ser199[I] (left) and salt bridge interaction between Piwi Lys+1 and Asp203[II] (right). H-bond and salt bridge are highlighted in green and orange, respectively. (I) Quantification of HP1a co-enrichment of XNP HACs with single alanine substitutions at positions -4, -3, +1, +3, or +4 (top) or double substitutions at positions -4 and -3 or +3 and +4 or quadruple substitutions at all positions (bottom). Asterisks denote p-values < 0.05 (*), <0.01 (**), <0.001 (***). (J) 2D molecular interactional map of HP1a CSD dimer with C-terminal tails and XNP HAC. Each CSD monomer is colored cyan and purple, respectively. XNP is colored yellow. Orange, green, pink, and blue dashed lines represent a salt bridge interaction, hydrogen bond, hydrophobic interaction, and pi-cation interaction.

I.

Supplementary Figure 5: (A) Visualization of salt bridge interactions between XNP HAC Arg+4 and Glu157 of HP1a CSDII (left). Histograms of the percentage frequency of salt bridge formation between the guanidinium and a guanidino group of the XNP HAC Arg+4 sidechain and 2 O atoms of HP1a Glu157[II] (right). (B) Histograms showing the percentage frequency of the salt bridges between Asp203[I] and +3 (left) and between Glu205[I] and +3 (right) of XNP as orange with a solid line, and of Ciz1 as cyan with a dashed line, between 200 to 250 ns of the MD simulations. (C) Interaction models of HP1a Arg197 showing 3 hydrogen bonds forming with Arg+1 of the XNP HAC (top), and a single hydrogen bond forming between Ser+1 plus charge repulsion with Lys+3 of the Ciz1 HAC (bottom). (D) Representative images of Ciz1 HAC and mutations either disabling central valine or adding a positive charge at the +1 position. (E) Histogram showing the percentage frequency of the Coulomb energy between Asp203[II] or Ser199[I] and +1 of XNP as orange with a solid line and Piwi as blue with a dashed line, between 200 to 250 ns of the MD simulations. All histograms were fitted by kernel density

estimation (KDE). (F) Alignment of all 16 HACs with pentameric core highlighted in yellow and marked with -2, 0, and +2 positions, and positively charged residues marked in red. (G) Representative images of mNeonGreen-tagged XNP HACs with single alanine substitutions at positions -4, -3, +1, +3, or +4 co-transfected with mScarlet-I-tagged HP1a (H) Representative images of XNP HACs containing combinations of alanine mutations at either -4 and -3, +3 and +4, or all four positions. (I) Interactional profiles of XNP from -7 to +7 generated by Discovery Studio. Orange, green, and pink dashed lines represent an electrostatic interaction, hydrogen bond, and hydrophobic interaction. Grey, blue, and red atoms of amino acid structures indicate carbon, nitrogen, and oxygen, respectively.

Generation of an improved HAC molecular signature using homolog analysis

Our discovery of new HAC sequences, coupled with their high conservation between

Drosophila species, amount to 173 peptide sequences that could be used to improve our

understanding of the HAC molecular signature. Color-coded sequence alignment stacks of

HACs and their homologs between hydrophobic, positively charged, and negatively charged

residues reveal high conservation within each group of HAC homologs (**Figure 6A** and

Supplementary Figure 6A). This suggests that HAC organization for each protein possesses

specific features that contribute to its affinity for HP1a. For example, HACs containing a

proline at -2 position often lack positively charged residues at the -3 and -4 positions, which

may indicate a proline-imparted kinked structure that prohibits positively charged residues at

those positions or render them unnecessary. We also identified rare, new residues at key

positions such as leucine or asparagine at position +4, of *D. willistoni* HP2-1 and Mu2-3,

respectively, capable of supporting HP1a co-enrichment (**Supplementary Figure 6B**), despite

the other 171 sequences containing positively charged residues at that position.

To identify unifying features of HACs, the occupancy of the same groups of amino acids were

quantified for each position (**Figure 6B**). Aside from the expected 100% occupancy of

nonpolar residues at positions -2, 0, and +2, there is nearly 80% occupancy at position +5 and

about 60% at positions -7, -5, -1, and +7. This indicates that hydrophobic forces in the flanking

regions of some HAC pentamers also contribute to functional HP1a binding directly or

indirectly. Consistent with our analysis in previous figures, positively charged residues

predominate at positions -6, -4, -3, +1, +3, and +4, which also correspond to a dearth of

negatively charged residues. We hypothesize that the less frequent but conserved occupations

of negatively charged residues in the flanking regions among specific HAC homologs reflect

mechanisms to partially destabilize HP1a affinity and/or utilize alternative mechanisms of

binding HP1a at those positions. This domination of HAC organization by nonpolar and basic

residues is highlighted by the corresponding dramatic drop of amino acid heterogeneity

between positions -6 and +6 (**Supplementary Figure 6C**). Finally, HAC organization across

various Drosophila species is also marked by association with intrinsically disordered regions

(**Figure 6C**), indicating that properties of these regions contribute to HAC function or that

HAC sequences impart disorder to the surrounding protein structure.

To summarize our findings, conservation analysis of Drosophila homolog sequences revealed

evolutionary positional bias for nonpolar and basic residues that correspond to various

molecular interactions with the HP1a dimer (**Figure 5I**). We refine the original Drosophila

HAC signature to consist of nearly any hydrophobic residue at -2, an invariant valine or

isoleucine at the central position, and nearly any aliphatic amino acid at +2. Expanding this

molecular signature, we also identify a nearly invariant basic residue at +4 (**Figure 6D**). We

also observe basic residues to be favored at -4, -3, +1, and +3, while hydrophobic interactions

are favored at positions -7, -5, -1, +5, and +7. However, due to the relative heterogeneity of

residues found at these positions, they likely modulate HP1a interactions, as we showed

previously for positions +1 and +3. Instead, we propose an enrichment of 3-6 basic residues

between positions -4 to +4 as a key signature of HACs (**Figure 6D**). Additionally, no acidic

residues are found at -6, -3, -1, +1 and +5, and no aromatic residues are located at -4 and +6,

suggesting these are detrimental to HAC function.

Figure 6 (A) Compilation of alignments for each HAC homolog across Drosophila evolution
showing positioning of basic (red), hydrophobic (blue), and acidic (green) residues. Position
numbers are indicated on the top and gene names are provided on the right of each stack of
alignments. (B) Percent occupancy of basic (red), hydrophobic (dotted), and acidic (green)
residues is calculated across Drosophila evolution for each position in all HACs. (C)
Compilation of disorder prediction scores are shown for each position across evolution for each
HAC. (D) Diagram of the evolutionarily conserved HAC molecular signature shows amino

 acids compatible for specific positions (top row), incompatible amino acids (second row),
 sidechain group bias (third row).

 Supplementary Figure 6. (A) HAC sequence pileup using MEME Suite 5.5.7 showing high
 conservation of specific residues among each group of homologs. (B) Imaging panels showing
 enrichment of putative Mu2 HAC from *Drosophila willistoni* containing a unique Asn+4. Scale
 bar = 1 um. (C) Positional bias for basic (red dots) and hydrophobic (blue dots) residues across
 Drosophila evolution of HACs is contrasted with the heterogeneity of amino acids (purple) for

each position.

Previously, we showed that the PxVxL motif based on known HP1a-binding sites is found in

nearly half of all proteins, including those annotated to be in the nucleus or chromosomes

(Supplementary Figure 1B). In contrast, the updated HAC molecular signature reduces the

number of potential candidates over 10-fold, even without accounting for disorder content

(Figure 7A). GO analysis of the proteins that contain sequences matching the updated HAC

signature yielded heterochromatin and chromocenter as the top annotations (**Figure 7B**). In

contrast, these categories were not significantly represented among the protein hits for the

PxVxL motif search, indicating that our updated HAC signature vastly improves the prediction

of HP1a-binding motifs.

To test the ability of the updated HAC signature to identify *bona fide* HP1a binding sites, we

used a known HP1a interactor with no known binding sites, HP5, as a test case (**Figure 7C**).

We identified two sequences in HP5 that matched our updated HAC signature (marked in pink),

whereas 6 other sequences fit a more inclusive search comprised of a hydrophobic-x-valine-x-

aliphatic signature (marked in yellow). HP5 was then subjected to a systematic fragment

analysis to identify functional HP1a-binding regions in an unbiased manner, using imaging for

HP1a co-localization as the readout (**Figure 7C**). Our results showed that only fragments

containing the two binding sites predicted by our updated signature co-localized with

heterochromatin **(Figure 7D),** suggesting these sequences are *bona fide* HP1a-binding sites.

Imaging analysis of 25-mer peptides harboring these putative HACs displayed remarkably high

enrichment at HP1a domains **(Figure 7D)**. Whereas individual mutations of the central residues

of these two HACs (V159E or V294E) did not impact association of full-length HP5-PA with

heterochromatin **(Supplementary Figure 7)**, a double mutation (V159E and V294E)

dramatically reduced HP5 expression and abrogated all association with heterochromatin (see

longer exposure). Therefore, our results indicate that HP5 is completely dependent on the

predicted HAC motifs for associating with the HP1a condensate (**Supplementary Figure 7**).

In conclusion, we propose that this updated signature, when combined with disorder prediction

and conservation analysis, will be invaluable for identifying other HP1a interactors and

regulators of the heterochromatin biocondensate.

Figure 7: (A) Venn diagram showing the number of proteins harboring a PxVxL motif as defined by the indicated molecular signature versus those with the updated HAC signature. Compare with Supplementary Figure 1B. (B) GO analysis results showing the top annotations of "heterochromatin" and "chromocenter" among proteins containing the updated HAC signature, but not among proteins harboring the degenerate PxVxL motif; FE = fold enrichment above expected results, FDR = false discovery rate, N/S = not significant. (C) Diagram of HP5 isoform PB with locations of putative HP1a-binding sites based on either the PxVxL motif of hydrophobic-x-Val-x-hydrophobic (in pink) or the new HAC signature (in yellow) described in Figure 6B. Green rectangles indicate intact gene or fragments that show co-localization with HP1a in S2R+ cells after transient transfection; gray rectangles indicate lack of co-localization.

774 (D) Representative images showing enrichment of HAC candidate peptides at HP1a-enriched
775 heterochromatin (left). Quantification of HP1a co-enrichment of candidate HP5 HAC peptides
776 (right). Scale bars = 2 microns.

777
778 **Supplementary Figure 7.** Representative images showing enrichment of wildtype or mutant
779 full-length HP5-PA isoform at HP1a biocondensates. A longer exposure of the HP5 double
780 mutant was necessary to visualize the loss of HP1a co-enrichment and accumulation at
781 nucleoli. Scale bars = 2 microns.
782

DISCUSSION:

The HAC motif provides a widespread and evolutionarily conserved mechanism for HP1a

binding and targeting of the heterochromatin domain. The results of this investigation expand

our understanding of this HP1a-binding motif by exposing (1) the range of degeneracy of the

core hydrophobic sequences that insert into the CSD dimer cleft, (2) the requirement of basic

residues to stabilize HP1a association in Drosophila, and (3) the molecular interactions that

modulate HAC affinity with the HP1a CSD and C-terminal tail. This led to expansion and

modification of the original pentameric PxVxL motif into a larger consensus region **(Figure**

6D) consisting of 3-6 positively charged residues that interlace and surround a core of three

alternating hydrophobic residues containing a central valine or isoleucine. This 9-mer basic

consensus pattern is further flanked by a moderate frequency of both hydrophobic and

positively charged residues, as well as low frequency of negatively charged and aromatic

residues, over a 15-mer span that, altogether, generate the molecular interface for binding to

the HP1a CSD dimer and CTE.

The HAC signature was determined through positional analysis of 6 previously identified

PxVxL sequences in *Drosophila melanogaster*, 10 more sequences identified from HP1a-

binding proteins through this study, and 157 sequences acquired from alignments of HP1a

partner homologs in 11 other Drosophila species. We note that though most of the HAC

homologs were untested in the peptide imaging assay, the positional and sequence conservation

between most HAC homologs were readily distinguished in alignments, suggesting that

protein-specific molecular interactions between HAC and HP1a are preserved between species.

For example, the Su(var)3-9 HAC is 100% identical over its 15-mer length among 12 species

of Drosophila, and the HP2-1 HAC is distinctive for containing a highly conserved Asn at the

+3 position (**Supplementary Figure 6A**). We also validated the HP1a co-enrichment of 4 HAC

homologs that represent some of the more divergent sequences (**Supplementary Figure 6B**).

Finally, the HAC-interacting residues of the HP1a CSDs and C-termini are preserved among

all Drosophila species included in this study (**Supplementary Figure 8A**). For these reasons,

we conclude that HAC homologs represent functionally conserved HP1a-binding domains that

greatly inform the HAC consensus signature. We do note that the high similarities in amino

acid sequences within each group of HAC homologs cause overrepresentation of specific

residues in the consensus analysis. Therefore, a more refined signature could be achieved

through the inclusion of more homologs from other Drosophila species or discovery of more

HACs in other HP1a-binding proteins.

Another predominant feature of HACs is their location within or in proximity to intrinsically

disordered regions of proteins. Disorder prediction was conducted using the widely used

PONDR VL-XT algorithm that integrates three neural networks trained on long (40+ residues),

N-terminal, and C-terminal disordered regions of X-ray crystallographic data (Li *et al.*, 1999;

Romero *et al.*, 2001). Though we cannot rule out the possibility that some HACs are located

on exposed surfaces of globular secondary structures, we hypothesize that HAC association

with intrinsically disordered domains confers flexibility and accessibility (Holehouse and

Kragelund, 2024) that may be important for targeting the CSD dimer. The AlphaFold-Multimer

model for the non-functional HAC candidate ADD1-1, which is predicted to contain a highly

ordered structure, fails to position the central hydrophobic core of ADD1-1 at the center of the

HP1a CSD dimer (**Supplementary Figure 8B**), despite containing a canonical motif (PxVxL).

The binding region exhibited lower prediction confidence (pLDDT < 50) than functional

peptides that contain highly disordered regions (pLDDT scores > 90). These structural

prediction results suggest that the flexibility of the peptide, attributed to its disordered regions,

may be necessary for proper fitting within the HP1a CSD dimer. We also detect the

conservation of disorder correlation among putative HACs of other Drosophila species. HACs

may therefore represent short linear motifs (SLiMs) or molecular recognition features (MoRFs)

within disordered regions that mediate diverse functions (Hsu *et al.*, 2013). Unlike well-defined

globular structures, SLiMs and MoRFs possess weaker conservation due to the higher mutation

rates found in disordered regions. Nevertheless, we identified positional biases in the

surrounding residues of the central hydrophobic core of HACs that contribute to HAC binding

with HP1a (**Figure 6D**). Aside from the functional role of flanking basic residues in modulating

interactional strength with HP1a, we also observed a strong bias against acidic and aromatic

residues when basic ones are absent, suggesting that their presence at these positions interfere

with HAC function. Additionally, the location of HACs at highly mutable disordered domains

and at paralogous locations in some protein homologs indicates that HACs may undergo

relatively rapid evolutionary selection but become stably inherited once established. Our

findings highlight the functional importance of intrinsically disordered domains as sites for

protein targeting and the evolution of HACs.

Our findings also expand the molecular signature of the original PxVxL motif that forms the

hydrophobic core of functional HACs. Though a trio of alternating hydrophobic residues is the

classical feature of the PxVxL motif, there is currently confusion regarding the types of

hydrophobic residues that can occupy each position in the motif, and whether this pentamer is

sufficient for HP1a binding (Lechner *et al.*, 2005; Meyer-Nava *et al.*, 2020). Our search for the

HAC molecular signature uncovers the range of amino acids that occupy each position of the

hydrophobic core residues. Expectedly, we identified mostly valines at the central position of

HACs, extending the previous structural prediction that only valine can be accommodated by

the hydrophobic pocket of the Drosophila HP1a CSD dimer (Thiru *et al.*, 2004). However, we

also identified isoleucines at the central positions of Su(var)3-9 and Lhr HACs of Drosophila

melanogaster, as well as in HACs of other homologs. Mutation of the central valine residue to

glutamate or alanine has previously been shown to eliminate HP1a binding, and we

demonstrate that mutation of a central isoleucine of a HAC similarly abrogates heterochromatin

targeting (**Supplementary Figure 1C**). Meanwhile, we identify that the -2 position utilizes a

much wider variety of hydrophobic residues, including polar cysteines, bulky phenylalanines,

and conformationally rigid prolines. Conservation analysis of HACs across Drosophila

evolution suggests that only the smallest and largest sidechain volumes of hydrophobic

residues are excluded at the -2 position, namely glycine, tyrosine, and tryptophan. Additionally,

our MD simulation shows that the sulfur in the cysteine sidechain at this position provides an

additional hydrogen bond that may enhance HP1a binding. In contrast, we identify only small

hydrophobic residues such as leucine, valine, and isoleucine to occupy the +2 position, though

alignment analysis also identifies rare homologs containing glycine and methionine instead.

This suggests that the +2 position is more sterically restrictive to larger hydrophobic residues

compared to the -2 position. Altogether, this trio of residues comprise an asymmetric

hydrophobic plug that is essential to recognize the conformation of the CSD dimer pocket, but

which we also show is insufficient to target heterochromatin effectively *in vivo*. Our HAC

fragment analysis (**Figure 3**) not only showed that the hydrophobic pentamer is insufficient for

heterochromatin targeting, but also that the flanking 5 residues on each side of the pentamer

are crucial for stable HAC association with heterochromatin.

Finally, we determined that electrostatic interactions are required for effective HAC-HP1a

dimer binding, in addition to hydrophobic interactions. Despite clear exceptions, it is

remarkable that there is a clear correlation between predicted Coulombic energy and measured

fold-enrichment (**Figure 4D**), given the higher complexity of *in vivo* conditions and potential

contributions from other factors not included in the simulations. In fact, the exceptions are

informative about potential complexities in HP1-partner interactions. For example, Lhr-2 has

larger attractive Coulomb energy (**Figure 4D**) but also greater hydrophilic surface areas than

higher-ranked HP2-1 and HP4-3 peptides (**Figure 4B, Supplementary Figure 4D**). This

suggests Lhr-2 contains a less stable hydrophobic pocket that could weaken HP1a interactions,

despite its high electrostatic energy. Meanwhile, other exceptions, such as HP2-1, HP4-3, and

Kdm4A, show high fold enrichment values despite lower Coulomb energy compared to other

HACs (**Figure 4D**). HP2-1 possesses L-2, V0, and I+2, the top three amino acids on the

hydropathy index (Kyte and Doolittle, 1982), whereas HP4-3 exhibits high hydrophobicity at

L+5, and Kdm4A displays significant hydrophobicity at the flanking region, specifically at L-

4 and L+5 (**Supplementary Figure 8C**). These peptides likely engage in more hydrophobic-

driven interactions that compensate for their lower electrostatic energy. We should also note

that electrostatic and hydrophobic interactions can negatively affect each other and be

competitive (Dzubiella and Hansen, 2003, 2004; Wang, Friesner and Berne, 2010). For

example, the Dmt HAC had weaker attractive Coulomb energy at the +1 and +4 positions than

XNP and Piwi, likely due to the presence of isoleucine, which has the highest hydropathy index

(Kyte and Doolittle, 1982), at the adjacent +2 and +5 positions (**Supplementary Figure 8D**).

Electrostatic interactions from the +1 and +4 positively charged amino acids are key

contributors to the HAC and CSD interaction (**Figure 5**). These results suggest that HAC

peptides and CSD dimer interactions are mediated by both hydrophobic and electrostatic

interactions, and that maintaining a moderate balance between hydrophobicity and charge is

crucial to prevent competition between these types of interactions. Therefore, the contributions

of electrostatic and hydrophobic interactions vary between HACs, and such structural

flexibility could provide the variations in HP1a affinity that are important for each

heterochromatin protein function.

Conservation analysis and MD simulations both indicate that positions -4, -3, -1, +1, +3, and

+4 contain a strong bias for basic residues (lysine or arginine) that make electrostatic

interactions with the HP1a CSD and C-terminal tail. In particular, we showed that the positively

charged amino acids ±4 form a salt bridge interaction with Asp157 (**Figure 5A, B**). The

positively charged amino acid at +1 plays an important role in maintaining the electrostatic

force field by forming an H-bond with Arg197 of the CSD dimer and enabling the +3 positively

charged amino acid to form salt bridges with Asp203 and Glu205 of the CSD (**Figure 5C, D,**

and E). Additionally, we identified compensatory mechanisms when specific electrostatic

interactions are absent at a given locus (+1, +3) (**Figure 5D, F**). The high conservation of

positive charges in Drosophila HACs is made possible by the absolute conservation of

interacting residues in Drosophila HP1a homologs, namely Asp157 of the CSD and Asp203

and Glu205 of the C-terminal tail (**Supplementary Figure 8A**). Acidic residues at positions

157 and 203 are also observed in HP1a homologs of humans, frogs, and fish, suggesting that

HACs of these organisms also utilize basic residues for targeting heterochromatin

(**Supplementary Figure 8E**). We note that, though HP1 C-terminal tails between these

homologs are highly variable in length and sequence, they nevertheless are highly enriched for

acidic residues that may partner with different positions of HACs for electrostatic interactions.

We therefore propose that the inclusion of 3-6 basic residues within HAC positions -4 to +4 is

an integral part of the HAC molecular signature in Drosophila HP1a, which may vary in content

and position in non-Drosophila HACs.

Supplementary Figure 8: (A) Alignment of CSD and C-terminal tails of *D. melanogaster*
HP1a homologs with other Drosophila species. Salt bridge-forming residues are marked with
orange dots, pi-cation residues in blue, hydrogen bonds in green. Residues that interact with
the hydrophobic core of HACs are marked with pink dots, whereas those that interact with
flanking hydrophobic residues are marked with purple dots. Conservation is marked with
purple highlight. HP1a CSD and CTE regions are marked with blue and yellow lines,
respectively. (B) The Alphafold Multimer-predicted computational model (top) and VMD-
visualized image of the model for the CSD dimer with C-terminal tails and ADD1-1 (bottom).
The Alphafold Multimer-predicted structure is colored based on pLDDT (predicted local

distance difference test) scores. Red: pLDDT<50, Orange: 50<pLDDT<60, Yellow:
60<pLDDT<70, Green: 70<pLDDT<80, Skyblue: 80<pLDDT<90, and Blue: 90<pLDDT.
CSD homodimer and the ADD1-1 are colored cyan and yellow, respectively. Residues at the -
2, 0, +2, +3, +4, +5, +6, +7, and +8 positions (in licorice drawing style) are colored orange,
with the amino acid code indicated. (C) Visualization of surface hydrophilicity and
hydrophobicity of XNP, H2-1, HP4-3, and Kdm4A in the side view were calculated using the
Discovery Studio software. The hydrophilic to hydrophobic properties are colored from blue
to brown. Each peptide is colored yellow. (D) Coulomb energy ± SD [kJ/mol] among the CSD
dimer and -7 to +7 residues of XNP, Dmt, and Piwi. Orange circle: XNP, Green diamond: Dmt,
and Blue square: Piwi. Electrostatic sequences at +1 and +4 are highlighted in orange;
hydrophobic sequences at +2 and +5 are highlighted in pink. (E) Alignment of CSD and C-
terminal tails of *D. melanogaster* HP1a with homologs from other species, including Homo
sapiens, Xenopus, zebrafish, worm, and yeast, using a similar format as shown in (A).

Variations in the number and positions of basic residues in different HACs indicate that they

act to modulate HP1a interactional strength. Though it is possible that there are HACs with

fewer basic residues, we predict that these will exhibit a much lower affinity for HP1a and

cannot be detected by our live imaging assay. Identification of an even more refined molecular

signature may also be achieved with a wider conservation analysis beyond the 12 Drosophila

species used here and more MD analysis for other types of peptides. To closely examine the

molecular interactions of a HAC, we used the highest-enriching sequence of XNP to be a model

peptide, by combining visualizations of interactions for residues from -7 to +7 with statistical

observations of the MD simulations (**Figure 5I** and **Supplementary Figure 5I**). The last frame

was initially used to make the interaction map, which was then adjusted by comparing it with

the statistical results, such as Coulomb energy, to obtain a more accurate interpretation. Based

on the interaction map, conservation analysis, and Coulomb energy, we can infer that the HAC

positions between -4 and +4 form the strongest molecular interactions, extending the

previously recognized 5-mer signature of PxVxL. As shown in **Figure 5I**, each amino acid has

distinct roles in the interactions between the peptide and the CSD dimer+CTE, and these roles

can change depending on the neighboring sequences. We note, for example, that homologs of

each HAC tend to possess additional conserved molecular features within their group that are

beyond the HAC consensus sequence (**Supplementary Figure 6A**), pointing to protein-

specific HP1a interfaces. A more comprehensive mutational study of such protein-specific

residues could uncover additional mechanisms of peptide interaction. Alternatively, factors

other than hydrophobicity and electrostatic forces could also be contributing to HAC-HP1a

interactions, such as post-translational modifications of peptides or HP1a.

But why would an affinity hierarchy exist for protein interactions with HP1a? HAC structures

likely reflect the specific requirements of heterochromatin proteins to achieve different levels

of enrichment or mixing in heterochromatin. For example, rare proteins may contain high-

affinity HACs for efficient targeting, whereas more abundant proteins, or those containing

additional domains that interact with other heterochromatin factors, will not require such high

affinity. Alternatively, structurally important heterochromatin proteins likely contain HACs

that stably associate with HP1a, whereas enzymes or proteins that only interface transiently

with heterochromatin may require lower affinities. For example, HP5 with its two strong HACs

could form highly stable heterochromatin structures or even crosslink two HP1a dimers,

whereas the Su(var)3-9 methyltransferase likely requires only transient association with HP1a

as its chromodomain is expected to bind H3K9me2/3 in heterochromatin (Müller *et al.*, 2016).

HAC affinities could therefore facilitate how HP1a organizes heterochromatin architecture by

directing the activity of multiple complexes to execute diverse functions in this complex,

dynamic biocondensate.

A hallmark of biocondensates is the accumulation of multivalent interactions between their

constituents to drive partitioning from the surrounding environment (Banani *et al.*, 2017). In

heterochromatin, this is achieved by the relatively abundant HP1 protein that centrally

organizes the accumulation of other structural condensate proteins through its abilities to bind

the heterochromatin-specific histone mark H3K9me2/3, homodimerize, and recognize HACs

(Janssen, Colmenares and Karpen, 2018). This inherent demixing ability also effectively

diminishes, if not outright excludes, other nuclear proteins from the condensate (Banani *et al.*,

2017). Nevertheless, the underlying heterochromatic DNA polymer is not precluded from pan-

nuclear processes such as transcription, replication, and repair. Therefore, heterochromatin

must either (1) employ dedicated factors to carry out such functions, (2) recruit factors from

such processes to access the interior, (3) transiently or locally dismantle the condensate for said

processes to proceed, or 4) a combination of these mechanisms. HACs can provide the means

to carry out at least the first two potential mechanisms, though controlled destabilization of

heterochromatin may also utilize an activatable HP1a-interacting protein. The HAC-containing

proteins SuUR and Mu2 appear to represent such factors that regulate DNA replication and

repair in heterochromatin, respectively, as SuUR is known to suppress polytenization of

heterochromatic domains (Belyaeva *et al.*, 2006) and Mu2 is an early DNA damage marker

that recruits downstream repair proteins (Dronamraju and Mason, 2009).

1009

HACs contribute to the multivalency of heterochromatin by facilitating the mixing of HP1a

with multiple proteins. Conversely, HP1a CSD dimerization is enhanced through binding

HACs, as has been previously shown with HP2-1 and Piwi peptides (Mendez *et al.*, 2011).

Whether higher HP1a affinity correlates with enhanced dimerization is currently unknown, but

HACs with different binding strengths, in combination with properties of linked domains in

full length versions, could regulate heterochromatin condensate formation, dissolution, and/or

material properties. Interestingly, mammalian HP1 phase separation properties were shown to

be differentially regulated by 2 different PxVxL-containing peptides, further indicating that

HAC-binding alters the biophysical properties of HP1 directly (Larson *et al.*, 2017). Our results

also indicate that HACs show enrichment throughout the heterochromatin domain, even though

the full-length proteins encoding certain HACs, such as SuUR and XNP, normally display

smaller subdomains of enrichment within heterochromatin (Swenson *et al.*, 2016). Subdomain

formation within heterochromatin must therefore be mediated by the concerted action of HACs

and other types of interaction domains present in full length proteins. One possible mechanism

is that such proteins utilize HACs to gain access to the heterochromatin interior and then,

through strong self-interacting domains or protein partners, form subdomain condensates

within heterochromatin. Through a similar pairing of self-interacting domains and particularly

weak affinity HACs, other nuclear bodies, such as those that facilitate transcription, replication,

and repair, may also gain access to the heterochromatin periphery that, upon activation, seed

the formation of complexes to carry out functions (Rajshekar *et al.*, 2023).

1030

We foresee that our improvements in identifying HAC molecular signatures will help identify

such 'seed' components for various nuclear processes, as well as focus attention on intrinsically

disordered regions of various proteins to contain access codes to different biocondensates in

the nucleus (King *et al.*, 2024). Whether all condensates contain short linear motifs to facilitate

the targeting of their components is not known, but our experimental and computational

approaches could provide a framework to identify molecular grammar utilized by other types

of biocondensates. Based on our findings with HACs, we propose that affinity hierarchies by

short disordered motifs may comprise a general mechanism by which various biocondensates

coordinate structural and transient components, as well as access shared nuclear components

and even interface with other condensates.

1041

1042   **ONLINE METHODS**

1043

Plasmid Construction

Oligonucleotides of putative HAC sequences (IDT) were cloned into a pCOPIA vector

containing a N-terminal mNeonGreen fluorescent tag and a myc nuclear localization signal.

HACs used for testing variable lengths of flanking sequence contained GGGSG in place of

sequences -12 to -8 and lacked positions +8 to +12, or GGGSGGSGGG in place of -12 to -3

and lacked positions +3 to +12. Corresponding control constructs contained GGGSG or

GGGSGGSGGG after the mNeonGreen tag and myc NLS. Full-length genes for HP1a, XNP,

SuUR, Lhr, Su(var)3-9, HP4, Mu2, and HP5, along with HP5 gene fragments, were cloned into

pCOPIA vectors containing N-terminal mCherry, mScarlet-1, mCitrine, or mGFP fluorescent

tags as described previously (Colmenares *et al.*, 2017). Mutants for HACs and full-length genes

were generated using a one-step directed mutagenesis protocol (Liu and Naismith, 2008).

1055

Bioinformatics

25-amino acid sequences containing a central pattern of hydrophobic-x-Val/Ile-x-hydrophobic

consensus were identified as putative HACs using custom Perl scripts. Hydrophobic residues

included Ala, Cys, Phe, Ile, Leu, Met, Pro, Val, Trp, and Tyr, whereas x = any amino acid.

Homologs of *Drosophila melanogaster* protein sequences were obtained using standard NCBI

protein-protein BLAST by retrieval of sequences with the highest homology scores for each of

the other 11 species of Drosophila (taxid:7215), using the non-redundant protein database.

Conservation of putative HACs was determined by testing whether the same consensus pattern

is detected to align with the putative *Drosophila melanogaster* HAC sequence after multiple

sequence alignment of full-length homologous proteins with Clustal Omega (Madeira *et al.*, 2024). Initially, only sequences containing a hydrophobic-x-Val/Ile-x-hydrophobic pattern that spatially aligned with the putative Drosophila melanogaster HAC, as determined by Clustal Omega with default settings, were scored positively to be conserved (see Supplementary Figure 2) and are shown on Figure 2. However, re-analysis of HP1a-binding protein homologs for the final HAC consensus motif (see below) revealed additional putative HAC sequences in other parts of SuUR, Mu2, and HP2 homologs that do not align with the Drosophila melanogaster HAC and may represent paralogous HACs. These additional sequences were included in the analysis for Figure 6 and Supplementary Figure 6. Disorder prediction was analyzed using PONDR-VLXT (Romero *et al.*, 2001). Positional frequency of each amino acid was analyzed among groups of HACs using custom Perl scripts. The final HAC consensus used for motif search from positions -7 to +7 are as follows: [^WF] [^DEW] [^WY] [^STFWY] [^DEFW] [FPLICVMA] [^DEFWY] [VI] [^DEFWY] [LIVGM] [^WY] [KRLN] [^DEW] [^FWY], where brackets contain all possible amino acids at each position, or all the excluded amino acids if marked with "^."

Tissue culture manipulation and image acquisition

S2R+ embryonic tissue culture cells were maintained in Schneider's medium (Thermo) with 10% fetal bovine serum (Gemini) and 1% antibiotic-antimycotic (Gibco) at 25°C. Transient co-transfections with mNeonGreen-tagged constructs and mScarlet-I-tagged HP1a were conducted using the TransIT-2020 reagent (Mirus). Cells were mounted 72 hours later on chambered coverslip slides (Ibidi) and live imaging was performed using a Zeiss LSM880 Airyscan microscope with 63X oil immersion objective at room temperature. A mNeonGreen tag containing only a myc NLS was co-transfected with mScarlet-I-HP1a as a negative control.

1089 Transient co-transfections of fluorescently-tagged full-length HP1a-interacting proteins with

1090 HP1a were similarly conducted and images were collected using an Applied Precision

1091 Deltavision microscope or a Zeiss LSM880 Airyscan microscope with 63X oil immersion

1092 objective. All results are based from at least two biological replicates.

1093

Image analysis

Raw images from the Zeiss Airyscan were processed using ZEN black edition software,

whereas Deltavision images were processed using SoftWoRx. All representative manuscript

images were formatted using Fiji software. To calculate heterochromatin enrichment, 3-D

Airyscan images of HP1a were segmented into nucleus, chromocenter, and nucleolus volume

masks by manually setting the intensity thresholds in Arivis Vision4D software. A low

intensity threshold setting selected the nucleus, a high setting was used to segment the

chromocenter, while the negative space lacking HP1a signal adjacent to the chromocenter was

isolated by the subtraction of HP1a nuclear masks with holes from those without holes. The

nucleolar mask was dilated 3-4 pixels to avoid potentially measuring nucleolar intensity signal

of mNeonGreen, which sometimes accumulated in the nucleolus in certain constructs,

including the negative control. The mean chromocenter signal intensity of mNeonGreen was

therefore measured only in chromocenter regions that did not overlap the dilated nucleolar

volume, and its chromocenter enrichment was calculated by dividing the mean chromocenter

intensity with the mean nuclear intensity. 10-36 nuclei were measured for each construct tested

from 2 biological replicates. Student's t test, assuming two-sample tails and unequal variance,

was calculated.

1111

Alphafold Multimer-predicted computational model

AlphaFold-Multimer (Evans *et al.*, 2022) was employed to develop the computational models

of the HAC peptide and HP1a CSD dimer with C-terminal tails. The sequences for the

Drosophila HP1a CSD dimer with C-terminal tails and flanking residues, sourced from the

protein database (PDB ID: 3P7J) (Mendez *et al.*, 2011), served as inputs for AlphaFold-

Multimer   computational   model   predictions:

MKKHHHHHHEQDTIPVSGSTGFDRGLEAEKILGASDNNGRLTFLIQFKGVDQAEMV

PSSVANEKIPRMVIHFYEERLSWYSDNEDKK. The 16 heterochromatin-enriched peptides

identified by our experiments (**Figure 1E**) were selected for computational modeling to

investigate the key features of the HP1a-peptide binding. The 25-amino-acid sequences for all

16 peptides are shown in Supplementary Method Table 1. The "pdb100" was used for the

template mode. The model type was set to "alphafold2_mmultimer_v3". The highest-ranked

model for each peptide, based on pLDDT (predicted Local Distance Difference Test) scores,

was used for the following computational analysis. All predicted models, except for Odj-1,

showed high pLDDT (predicted Local Distance Difference Test, scaled 0-100) and pTM

(predicted Template Modelling, scaled 0-1) scores, representing confidence in the local and

overall structure of the complex, respectively (**Supplementary Method Table 1**). The 15 out

of 16 peptide models positioned their central core region at the center of the CSD dimer (**Figure**

3A, Supplementary Figure 3A-O). While the disordered edge regions of the complex showed

low prediction accuracy, the key binding sites between the peptide and CSD dimer had very

high confidence with over 90 pLDDT scores (**Supplementary Figure 3A-O**). Odj-1 was the

only model that mispositioned the peptide and had low prediction accuracy even in the binding

sites with 60-80 pLDDT scores. Due to its low prediction confidence, the Odj-1 model was

excluded from further computational analysis. The 15 peptide models, except for Odj-1, were

used for the following molecular dynamics simulation analysis.

Supplementary Method Table 1: The 25-amino-acid sequences of 16 heterochromatin-enriched peptides, with the central valine (V) designated as position 0. The table also includes AlphaFold-Multimer prediction scores: pLDDT (predicted Local Distance Difference Test, scaled from 0 to 100) and pTM (predicted Template Modelling, scaled from 0 to 1) were shown.

Peptide Name	-12	-11	-10	-9	-8	-7	-6	-5	-4	-3	-2	-1	0	1	2	3	4	5	6	7	8	9	10	11	12	pLDDT	pTM
XNP	R	A	E	S	P	P	R	P	K	R	C	V	V	R	L	K	R	V	S	L	P	K	T	K	P	76.7	0.706
Dmt	L	N	E	P	E	C	R	I	R	K	C	V	V	K	I	R	R	I	K	V	K	N	G	V	V	75.9	0.696
Odj-1	E	D	Q	N	F	P	N	R	Q	S	P	R	V	R	V	K	R	L	R	V	P	V	V	E	K	72	0.655
HP2-1	T	V	P	Q	N	I	S	P	R	K	L	C	V	K	I	N	R	R	P	Y	N	K	W	L	R	77.3	0.729
HP4-3	P	F	P	L	K	P	K	P	K	R	C	I	V	R	V	A	R	L	P	D	V	E	R	S	E	76.2	0.705
Lhr-2	L	K	S	A	K	Y	Y	I	K	R	C	R	I	R	L	K	R	V	E	I	D	D	Y	L	P	80.1	0.761
Kdm4A	S	S	Y	S	S	C	R	Q	L	Q	P	V	V	K	L	R	K	L	P	T	I	A	S	V	P	75.7	0.714
HP2-2	L	R	T	P	T	A	S	C	R	K	L	R	V	L	I	K	R	T	P	T	S	S	L	S	L	77.1	0.732
HP4-2	E	R	E	V	E	Y	Q	V	R	K	C	R	V	N	L	E	R	I	K	S	S	T	S	P	N	75.6	0.687
SuUR-2	T	V	H	L	T	P	K	T	R	R	L	N	V	R	I	L	R	V	S	L	D	T	L	S	T	77.6	0.735
Ciz1	K	P	G	P	A	S	T	K	R	N	V	L	V	S	V	K	R	T	S	A	A	A	I	K	A	75.4	0.719
HP4-1				M	S	P	K	T	K	K	M	I	V	K	I	P	R	H	P	Y	F	N	V	Q	G	78.6	0.727
Mu2-2	V	K	T	G	G	A	L	I	K	R	A	V	V	R	I	S	R	V	S	I	E	Q	L	E	S	77.8	0.732
ADD1-3	K	K	S	D	A	K	L	K	M	K	V	M	V	K	L	K	R	A	E	D	E	F	E	V	A	79.4	0.741
Su(var)3-9	V	R	F	K	N	E	L	E	R	N	I	T	I	K	L	E	R	L	S	E	K	K	I	K	N	78.2	0.74
Piwi	S	S	T	S	R	G	S	G	D	G	P	R	V	K	V	F	R	G	S	S	S	G	D	P	R	75.4	0.688

1142

1143

Supplementary Method Figure 1: (A) The Alphafold Multimer-predicted computational model and (B) visualized image of the model for the CSD dimer with C-terminal tails and Odj-1. The Alphafold Multimer-predicted structure is colored based on pLDDT (predicted local distance difference test) scores. Red: pLDDT<50, Orange: 50<pLDDT<60, Yellow: 60<pLDDT<70, Green: 70<pLDDT<80, Skyblue: 80<pLDDT<90, and Blue: 90<pLDDT. CSD homodimer and the Odj-1 are colored cyan and yellow, respectively. Residues at the -2, 0, +2, +5, and +7 positions (in licorice drawing style) are colored orange, with the amino acid code indicated. (C and D) Root mean square deviation (RMSD) of (C) the HP1a CSD dimer (amino acids 141-201) in Piwi and XNP peptide models and (D) the HAC peptides of Piwi and XNP from -7 to +7 amino acids throughout the 250 ns simulations.

1154

Molecular dynamics simulation

GROMACS 2020.4 software package (Pronk *et al.*, 2013; Abraham *et al.*, 2015; Páll *et al.*,

2015; Lindahl *et al.*, 2020) was used for the all-atom molecular dynamics (MD) simulation

with the CHARMM36 force field (Best et $al.$, 2012; Huang and Alexander D MacKerell, 2013).

The developed model of the HAC peptide and HP1a CSD dimer with C-terminal tails was

inserted into a cube box ($13*13*13$ nm^3) with periodic boundary conditions applied in all three

directions. The CHARMM-modified TIP3P water model was employed to solvate the system

(Boonstra, Onck and van der Giessen, 2016). The sodium and chloride counter ions were added

to the system to obtain the 150 mM concentration of NaCl, a condition where chromatin is

known to be aggregated (Huang and Cole, 1984; Shakya et $al.$, 2020). Energy minimization

was conducted for 50,000 steps with an energy tolerance of 1000 $kJ \cdot mol^{-1} nm^{-1}$ to eliminate no

steric clashes and ensure proper geometry of the system. The target temperature and

equilibration of the solvent and ions in the system were achieved by employing NVT

(isothermal-isochoric) and NPT (isothermal-isobaric) ensembles, each for 5 ns with a time step

of 1 fs, with V-rescale thermostat at a constant temperature of 298 K. The Parrinello-Rahman

barostat was used during NPT ensembles to maintain a pressure of 1 atm. The damping

parameters were set to 0.1 ps for the V-rescale thermostat and 2 ps for the Parrinello-Rahman

barostat. The heavy atom's positions of the protein were constrained during NVT and NPT

equilibration to prevent the additional variable of structural changes in the protein.

Subsequently, protein relaxation was employed for 250 ns (with 2 fs time steps) as the product

step without any position restraints in the equilibrated system. The MD simulations were

performed using NVIDIA V100 SMX2 GPU at the San Diego Supercomputer Center. The

particle mesh Ewald method was used for long-range electrostatic interactions. The 1.0 nm

cutoff was used for short-range electrostatic and van der Waals interactions.

1179

Analysis of Molecular Dynamics Simulations

The trajectory files saved every 10 ps were used for the analyses. The HP1a CSD dimer (amino

acids 141-201) in Piwi and XNP peptide models was stable throughout the 250 ns simulations

(**Supplementary Method Figure 1C**). The HAC peptides of Piwi and XNP from -7 to +7

amino acids reached a stable state after around 175 ns (**Supplementary Method Figure 1D**).

Data from 200 ns to 250 ns after reaching the stable state was used for computational analysis.

The simulation was visualized using VMD software (Humphrey, Dalke and Schulten, 1996).

All analyzed results were plotted using Python.

Root Mean Square Deviation (RMSD): RMSD of the HAC central pentamer from -2 to +2 for

the last 50 ns was calculated using "gmx rms" GROMACS tool built-in function.

Root Mean Square Fluctuation (RMSF): RMSF of the XNP and Piwi of 25-residue HAC

sequences (-12 to +12) for the last 50 ns was calculated using "gmx rmsf" GROMACS tool

built-in function. Averaged RMSF of the distal flanking 10 sequences (-12 to -8 and +8 to +12),

nearby flanking 10 sequences (-7 to -3 and +3 to +7), and central 5 sequences (-2 to +2) for

XNP and Piwi was calculated, respectively. Tukey test was employed to compare differences

between multiple groups (p-values < 0.05).

Hydrophobic and charge distribution: The hydrophilic and hydrophobic distribution and

electrical charge distribution of the HP1a-interacting XNP regions were visualized using

BIOVIA Discovery Studio (Dassault Systèmes, USA) (*Discovery Studio*, 2023).

Hydrophilic surface area: The hydrophilic surface area of the HAC central pentamer (-2 to +2)

and the pentamer with nearby flanking (-7 to +7) were computed for the last 50 ns of the MD

simulations using "gmx sasa" GROMACS tool built-in function for oxygen and nitrogen

atoms.

Coulomb energy: The short-range Coulomb energy between the HAC peptide and HP1a CSD

dimer with C-terminal tails with a 1.0 nm cutoff distance was calculated for the last 50 ns of

the MD simulations using "gmx energy" GROMACS tool built-in function. The Coulomb

energy for each individual residue was calculated in the same manner. The frequencies of the

coulombic energy with kernel density estimation were plotted using Python.

BIOVIA (*Discovery Studio*, 2023) visualization of hydrophilicity/hydrophobicity and surface

charge distributions: The hydrophobicity on the peptide is derived from the Kyte-Doolittle

scale. For the charge calculation, partial charges assigned to each surface atom are used. The

algorithm details can be found in BIOVIO Discovery Studio (*Discovery Studio*, 2023).

VMD (Humphrey, Dalke and Schulten, 1996) visualization of hydrogen bonds (H-bonds) and

salt bridge interactions: H-bonds were colored in green when the distance between the possible

acceptor (OH and NH groups) and donor (O and N atoms) pair was within 0.35 nm, and the

angle between the hydrogen, oxygen, and nitrogen was below 30°. Salt bridge interactions were

colored in orange when the distance between the side-chain carbonyl oxygen of Asp or Glu

and the side-chain nitrogen atoms of Arg, Lys, or His is within 0.4 nm.

2D molecular interaction map: The 2D molecular interactional map of XNP and HP1a CSD

dimer with C-terminal tails was created using Adobe Illustrator (Adobe Inc., USA) (*Vector*

Graphics Software – Adobe Illustrator, no date) based on the 2D diagrams for each residue

analyzed by BIOVIA Discovery Studio (Dassault Systèmes, USA) (*Discovery Studio*, 2023).

1222

Acknowledgements:

Many thanks to members of the Karpen lab and Dr. Grace Y.C. Lee for their valuable feedback.

We thank Dr. Ksenia V Krasileva and Dr. Daniil M. Prigozhint for their helpful discussion

regarding AlphaFold-Multimer modeling, the Japan Student Services Organization (JASSO)

Student Exchange Support Program (Graduate Scholarship for Degree Seeking Students), the

SAITAMA GO GLOBAL STUDENT scholarship, and the Hearts To Humanity Eternal

(H2H8) graduate research grant for supporting Shingo Tsukamoto's doctoral studies. This

research used resources of the Advanced Cyberinfrastructure Coordination Ecosystem:

Services & Support (ACCESS) super-computing facilities, supported by the National Science

Foundation (NSF) Grant No. BIO230220. The Karpen lab is supported by the National

Institutes of Health (R35GM139653).

1234

<u>References.</u>

Abraham, M.J. *et al.* (2015) 'GROMACS: High performance molecular simulations through
multi-level parallelism from laptops to supercomputers', *SoftwareX*, 1–2, pp. 19–25. Available
at: https://doi.org/10.1016/j.softx.2015.06.001.

Alekseyenko, A.A. *et al.* (2014) 'Heterochromatin-associated interactions of Drosophila HP1a
with dADD1, HIPP1, and repetitive RNAs', *Genes & Development*, 28(13), pp. 1445–1460.
Available at: https://doi.org/10.1101/gad.241950.114.

Badugu, R. *et al.* (2005) 'Mutations in the heterochromatin protein 1 (HP1) hinge domain
affect HP1 protein interactions and chromosomal distribution', *Chromosoma*. Available at:
https://doi.org/10.1007/s00412-004-0324-2.

Banani, S.F. *et al.* (2017) 'Biomolecular condensates: organizers of cellular biochemistry',
Nature Reviews. Molecular Cell Biology, 18(5), pp. 285–298. Available at:
https://doi.org/10.1038/nrm.2017.7.

Bannister, A.J. *et al.* (2001) 'Selective recognition of methylated lysine 9 on histone H3 by the
HP1 chromo domain.', *Nature*. Available at: https://doi.org/10.1038/35065138.

Bassett, A.R. *et al.* (2008) 'The chromatin remodelling factor dATRX is involved in
heterochromatin formation', *PLoS ONE*. Available at:
https://doi.org/10.1371/journal.pone.0002099.

Belyaeva, E.S. *et al.* (2006) 'DNA underreplication in intercalary heterochromatin regions in
polytene chromosomes of Drosophila melanogaster correlates with the formation of partial
chromosomal aberrations and ectopic pairing', *Chromosoma*. Available at:
https://doi.org/10.1007/s00412-006-0063-7.

Best, R.B. *et al.* (2012) 'Optimization of the additive CHARMM all-atom protein force field
targeting improved sampling of the backbone ϕ, ψ and side-chain $\chi(1)$ and $\chi(2)$ dihedral
angles', *Journal of Chemical Theory and Computation*, 8(9), pp. 3257–3273. Available at:
https://doi.org/10.1021/ct300400x.

Boonstra, S., Onck, P.R. and van der Giessen, E. (2016) 'CHARMM TIP3P Water Model
Suppresses Peptide Folding by Solvating the Unfolded State', *The Journal of Physical
Chemistry B*, 120(15), pp. 3692–3698. Available at:
https://doi.org/10.1021/acs.jpcb.6b01316.

Brasher, S.V. *et al.* (2000) 'The structure of mouse HP1 suggests a unique mode of single
peptide recognition by the shadow chromo domain dimer', *The EMBO journal*, 19(7), pp.
1587–1597. Available at: https://doi.org/10.1093/emboj/19.7.1587.

Brennecke, J. *et al.* (2007) 'Discrete small RNA-generating loci as master regulators of
transposon activity in Drosophila', *Cell*, 128(6), pp. 1089–1103. Available at:
https://doi.org/10.1016/j.cell.2007.01.043.

Brideau, N.J. and Barbash, D.A. (2011) 'Functional conservation of the Drosophila hybrid
incompatibility gene Lhr', *BMC Evolutionary Biology*, 11(1), p. 57. Available at:
https://doi.org/10.1186/1471-2148-11-57.

Brower-Toland, B. *et al.* (2007) 'Drosophila PIWI associates with chromatin and interacts
directly with HP1a', *Genes & Development*, 21(18), pp. 2300–2311. Available at:
https://doi.org/10.1101/gad.1564307.

Chiolo, I. *et al.* (2011) 'Double-Strand Breaks in Heterochromatin Move Outside of a Dynamic
HP1a Domain to Complete Recombinational Repair', *Cell*. Available at:
https://doi.org/10.1016/j.cell.2011.02.012.

Colmenares, S.U. *et al.* (2017) 'Drosophila Histone Demethylase KDM4A Has Enzymatic and
Non-enzymatic Roles in Controlling Heterochromatin Integrity', *Developmental Cell*, 42(2),
pp. 156-169.e5. Available at: https://doi.org/10.1016/j.devcel.2017.06.014.

Coustham, V. *et al.* (2006) 'The *C. elegans* HP1 homologue HPL-2 and the LIN-13 zinc finger
protein form a complex implicated in vulval development', *Developmental Biology*, 297(2), pp.
308–322. Available at: https://doi.org/10.1016/j.ydbio.2006.04.474.

Czech, B. *et al.* (2008) 'An endogenous small interfering RNA pathway in Drosophila', *Nature*,

453(7196), p. 798. Available at: https://doi.org/10.1038/nature07007.

Dimitri, P. *et al.* (2003) 'Vital genes in the heterochromatin of chromosomes 2 and 3 of
Drosophila melanogaster', *Genetica*.

Discovery Studio (2023) *Dassault Systèmes*. Available at:
https://www.3ds.com/products/biovia/discovery-studio (Accessed: 23 October 2024).

Dronamraju, R. and Mason, J.M. (2009) 'Recognition of double strand breaks by a mutator
protein (MU2) in Drosophila melanogaster', *PLoS Genet*. Available at:
https://doi.org/10.1371/journal.pgen.1000473.

Dronamraju, R. and Mason, J.M. (2011) 'MU2 and HP1a regulate the recognition of double
strand breaks in Drosophila melanogaster', *PLoS One*. Available at:
https://doi.org/10.1371/journal.pone.0025439.

Dzubiella, J. and Hansen, J.-P. (2003) 'Reduction of the hydrophobic attraction between
charged solutes in water', *The Journal of Chemical Physics*, 119(23), pp. 12049–12052.
Available at: https://doi.org/10.1063/1.1632902.

Dzubiella, J. and Hansen, J.-P. (2004) 'Competition of hydrophobic and Coulombic
interactions between nanosized solutes', *The Journal of Chemical Physics*, 121(11), pp. 5514–
5530. Available at: https://doi.org/10.1063/1.1783274.

Eissenberg, J.C. *et al.* (1992) 'The heterochromatin-associated protein HP-1 is an essential
protein in Drosophila with dosage-dependent effects on position-effect variegation', *Genetics*.

Evans, R. *et al.* (2022) 'Protein complex prediction with AlphaFold-Multimer'. bioRxiv, p.
2021.10.04.463034. Available at: https://doi.org/10.1101/2021.10.04.463034.

Her, C. *et al.* (2022) 'Molecular interactions underlying the phase separation of HP1α: role of
phosphorylation, ligand and nucleic acid binding', *Nucleic Acids Research*, 50(22), pp. 12702–
12722. Available at: https://doi.org/10.1093/nar/gkac1194.

Holehouse, A.S. and Kragelund, B.B. (2024) 'The molecular basis for cellular function of
intrinsically disordered protein regions', *Nature Reviews. Molecular Cell Biology*, 25(3), pp.
187–211. Available at: https://doi.org/10.1038/s41580-023-00673-0.

Honda, S. and Selker, E.U. (2008) 'Direct interaction between DNA methyltransferase DIM-
2 and HP1 is required for DNA methylation in Neurospora crassa', *Molecular and Cellular*
Biology, 28(19), pp. 6044–6055. Available at: https://doi.org/10.1128/MCB.00823-08.

Hoskins, R.A. *et al.* (2007) 'Sequence finishing and mapping of Drosophila melanogaster
heterochromatin', *Science (New York, NY)*. Available at:

https://doi.org/10.1126/science.1139816.

Hsu, K.-L., Yen, H.-C.S. and Yeang, C.-H. (2022) 'Cooperative stability renders protein
complex formation more robust and controllable', *Scientific Reports*, 12(1), p. 10490. Available
at: https://doi.org/10.1038/s41598-022-14362-z.

Hsu, W.-L. *et al.* (2013) 'Exploring the binding diversity of intrinsically disordered proteins
involved in one-to-many binding', *Protein Science: A Publication of the Protein Society*, 22(3),
pp. 258–273. Available at: https://doi.org/10.1002/pro.2207.

Huang, H.C. and Cole, R.D. (1984) 'The distribution of H1 histone is nonuniform in
chromatin and correlates with different degrees of condensation.', *Journal of Biological*
Chemistry, 259(22), pp. 14237–14242. Available at: https://doi.org/10.1016/S0021-
9258(18)89883-0.

Huang, J. and Alexander D MacKerell, J. (2013) 'CHARMM36 all-atom additive protein force
field: Validation based on comparison to NMR data', *Journal of computational chemistry*,
34(25), p. 2135. Available at: https://doi.org/10.1002/jcc.23354.

Humphrey, W., Dalke, A. and Schulten, K. (1996) 'VMD: visual molecular dynamics', *Journal*
of Molecular Graphics, 14(1), pp. 33–38, 27–28. Available at: https://doi.org/10.1016/0263-
7855(96)00018-5.

Janssen, A., Colmenares, S.U. and Karpen, G.H. (2018) 'Heterochromatin: Guardian of the
Genome', *Annual Review of Cell and Developmental Biology*, 34, pp. 265–288. Available at:
https://doi.org/10.1146/annurev-cellbio-100617-062653.

Kasinathan, B. *et al.* (2020) 'Innovation of heterochromatin functions drives rapid evolution of
essential ZAD-ZNF genes in Drosophila', *eLife*, 9, p. e63368. Available at:
https://doi.org/10.7554/eLife.63368.

King, M.R. *et al.* (2024) 'Macromolecular condensation organizes nucleolar sub-phases to set
up a pH gradient', *Cell*, 187(8), pp. 1889-1906.e24. Available at:
https://doi.org/10.1016/j.cell.2024.02.029.

Kumar, S. and Nussinov, R. (2002) 'Close-Range Electrostatic Interactions in Proteins',
ChemBioChem, 3(7), pp. 604–617. Available at: https://doi.org/10.1002/1439-
7633(20020703)3:7<604::AID-CBIC604>3.0.CO;2-X.

Kyte, J. and Doolittle, R.F. (1982) 'A simple method for displaying the hydropathic character
of a protein', *Journal of Molecular Biology*, 157(1), pp. 105–132. Available at:
https://doi.org/10.1016/0022-2836(82)90515-0.

Lachner, M. *et al.* (2001) 'Methylation of histone H3 lysine 9 creates a binding site for HP1

proteins.', *Nature*. Available at: https://doi.org/10.1038/35065132.

Larson, A.G. *et al.* (2017) 'Liquid droplet formation by HP1alpha suggests a role for phase
separation in heterochromatin', *Nature*. Available at: https://doi.org/10.1038/nature22822.

Lechner, M.S. *et al.* (2005) 'The mammalian heterochromatin protein 1 binds diverse nuclear
proteins through a common motif that targets the chromoshadow domain', *Biochemical and*
Biophysical Research Communications, 331(4), pp. 929–937. Available at:
https://doi.org/10.1016/j.bbrc.2005.04.016.

Li, X. *et al.* (1999) 'Predicting Protein Disorder for N-, C-, and Internal Regions', *Genome*
Informatics. Workshop on Genome Informatics, 10, pp. 30–40.

Lin, C.-H. *et al.* (2008) 'Heterochromatin protein 1a stimulates histone H3 lysine 36
demethylation by the Drosophila KDM4A demethylase', *Molecular Cell*. Available at:
https://doi.org/10.1016/j.molcel.2008.11.008.

Lindahl *et al.* (2020) 'GROMACS 2020 Manual'. Available at:
https://doi.org/10.5281/zenodo.3562512.

Liu, H. and Naismith, J.H. (2008) 'An efficient one-step site-directed deletion, insertion, single
and multiple-site plasmid mutagenesis protocol', *BMC Biotechnology*, 8(1), p. 91. Available
at: https://doi.org/10.1186/1472-6750-8-91.

Liu, Y. *et al.* (2017) 'Peptide recognition by heterochromatin protein 1 (HP1) chromoshadow
domains revisited: Plasticity in the pseudosymmetric histone binding site of human HP1',
Journal of Biological Chemistry, 292(14), pp. 5655–5664. Available at:
https://doi.org/10.1074/jbc.M116.768374.

Madeira, F. *et al.* (2024) 'The EMBL-EBI Job Dispatcher sequence analysis tools framework
in 2024', *Nucleic acids research*, 52(W1), pp. W521–W525. Available at:
https://doi.org/10.1093/nar/gkae241.

Makunin, I.V. *et al.* (2002) 'The Drosophila suppressor of underreplication protein binds to
late-replicating regions of polytene chromosomes', *Genetics*.

Marr, S.K. *et al.* (2014) 'The Metazoan-Specific Mediator Subunit 26 (Med26) Is Essential for
Viability and Is Found at both Active Genes and Pericentric Heterochromatin in Drosophila
melanogaster', *Molecular and Cellular Biology*, 34(14), pp. 2710–2720. Available at:
https://doi.org/10.1128/MCB.01365-13.

Mendez, D.L. *et al.* (2011) 'The HP1a disordered C terminus and chromo shadow domain
cooperate to select target peptide partners', *Chembiochem: A European Journal of Chemical*
Biology, 12(7), pp. 1084–1096. Available at: https://doi.org/10.1002/cbic.201000598.

Mendez, D.L., Mandt, R.E. and Elgin, S.C.R. (2013) 'Heterochromatin Protein 1a (HP1a)
partner specificity is determined by critical amino acids in the chromo shadow domain and C-
terminal extension', *The Journal of Biological Chemistry*, 288(31), pp. 22315–22323. Available
at: https://doi.org/10.1074/jbc.M113.468413.

Meyer-Nava, S. *et al.* (2020) 'Insights into HP1a-Chromatin Interactions', *Cells*, 9(8), p. 1866.
Available at: https://doi.org/10.3390/cells9081866.

Muchardt, C. *et al.* (2002) 'Coordinated methyl and RNA binding is required for
heterochromatin localization of mammalian HP1alpha.', *EMBO reports*. Available at:
https://doi.org/10.1093/embo-reports/kvf194.

Müller, M.M. *et al.* (2016) 'A two-state activation mechanism controls the histone
methyltransferase Suv39h1', *Nature Chemical Biology*, 12(3), pp. 188–193. Available at:
https://doi.org/10.1038/nchembio.2008.

Páll, S. *et al.* (2015) 'Tackling Exascale Software Challenges in Molecular Dynamics
Simulations with GROMACS', in S. Markidis and E. Laure (eds) *Solving Software Challenges*
for Exascale. Cham: Springer International Publishing (Lecture Notes in Computer Science),
pp. 3–27. Available at: https://doi.org/10.1007/978-3-319-15976-8_1.

Perrin, L. *et al.* (1998) 'Dynamics of the sub-nuclear distribution of Modulo and the regulation
of position-effect variegation by nucleolus in Drosophila.', *Journal of cell science*.

Pindyurin, A.V. *et al.* (2008) 'Interaction between the Drosophila heterochromatin proteins
SUUR and HP1', *J Cell Sci*. Available at: https://doi.org/10.1242/jcs.018655.

Pronk, S. *et al.* (2013) 'GROMACS 4.5: a high-throughput and highly parallel open source
molecular simulation toolkit', *Bioinformatics (Oxford, England)*, 29(7), pp. 845–854. Available
at: https://doi.org/10.1093/bioinformatics/btt055.

Pylaeva, S., Brehm, M. and Sebastiani, D. (2018) 'Salt Bridge in Aqueous Solution: Strong
Structural Motifs but Weak Enthalpic Effect', *Scientific Reports*, 8(1), p. 13626. Available at:
https://doi.org/10.1038/s41598-018-31935-z.

Rajshekar, S. *et al.* (2023) 'Affinity hierarchies and amphiphilic proteins underlie the co-
assembly of nucleolar and heterochromatin condensates', *bioRxiv: The Preprint Server for*
Biology, p. 2023.07.06.547894. Available at: https://doi.org/10.1101/2023.07.06.547894.

Romero, P. *et al.* (2001) 'Sequence complexity of disordered protein', *Proteins: Structure,*
Function, and Bioinformatics, 42(1), pp. 38–48. Available at: https://doi.org/10.1002/1097-
0134(20010101)42:1<38::AID-PROT50>3.0.CO;2-3.

Schotta, G. *et al.* (2002) 'Central role of Drosophila SU(VAR)3-9 in histone H3-K9

methylation and heterochromatic gene silencing', *The EMBO journal*. Available at:
https://doi.org/10.1093/emboj/21.5.1121.

Schwendemann, A. *et al.* (2008) 'Hip, an HP1-interacting protein, is a haplo- and triplo-
suppressor of position effect variegation', *Proceedings of the National Academy of Sciences of*
the United States of America, 105(1), pp. 204–209. Available at:
https://doi.org/10.1073/pnas.0705595105.

Seller, C.A. and O'Farrell, P.H. (2018) 'Rif1 prolongs the embryonic S phase at the Drosophila
mid-blastula transition', *PLoS biology*, 16(5), p. e2005687. Available at:
https://doi.org/10.1371/journal.pbio.2005687.

Shaffer, C.D. *et al.* (2002) 'Heterochromatin protein 2 (HP2), a partner of HP1 in Drosophila
heterochromatin', *Proceedings of the National Academy of Sciences of the United States of*
America. Available at: https://doi.org/10.1073/pnas.212458899.

Shakya, A. *et al.* (2020) 'Liquid-Liquid Phase Separation of Histone Proteins in Cells: Role in
Chromatin Organization', *Biophysical Journal*, 118(3), pp. 753–764. Available at:
https://doi.org/10.1016/j.bpj.2019.12.022.

Smothers, J.F. and Henikoff, S. (2000) 'The HP1 chromo shadow domain binds a consensus
peptide pentamer', *Current biology: CB*, 10(1), pp. 27–30. Available at:
https://doi.org/10.1016/s0960-9822(99)00260-2.

Stephens, G.E. *et al.* (2005) 'Interaction of heterochromatin protein 2 with HP1 defines a novel
HP1-binding domain', *Biochemistry*, 44(40), pp. 13394–13403. Available at:
https://doi.org/10.1021/bi051006+.

Strom, A.R. *et al.* (2017) 'Phase separation drives heterochromatin domain formation', *Nature*.
Available at: https://doi.org/10.1038/nature22989.

Swenson, J.M. *et al.* (2016) 'The composition and organization of Drosophila heterochromatin
are heterogeneous and dynamic', *eLife*, 5, p. e16096. Available at:
https://doi.org/10.7554/eLife.16096.

Thiru, A. *et al.* (2004) 'Structural basis of HP1/PXVXL motif peptide interactions and HP1
localisation to heterochromatin', *The EMBO journal*, 23(3), pp. 489–499. Available at:
https://doi.org/10.1038/sj.emboj.7600088.

Vector Graphics Software – Adobe Illustrator (no date). Available at:
https://www.adobe.com/products/illustrator.html (Accessed: 23 October 2024).

Wang, L., Friesner, R.A. and Berne, B.J. (2010) 'Competition of electrostatic and hydrophobic
interactions between small hydrophobes and model enclosures', *The journal of physical*

chemistry. B, 114(21), pp. 7294–7301. Available at: https://doi.org/10.1021/jp100772w.

Wang, T. *et al.* (2012) 'Crystal Structure of the Human SUV39H1 Chromodomain and Its
Recognition of Histone H3K9me2/3', *PLoS ONE*, 7(12), p. e52977. Available at:
https://doi.org/10.1371/journal.pone.0052977.

Weiler, K.S. and Wakimoto, B.T. (1995) 'Heterochromatin and gene expression in
Drosophila.', *Annual review of genetics*. Available at:
https://doi.org/10.1146/annurev.ge.29.120195.003045.

Xie, N.-Z. *et al.* (2015) 'Exploring Strong Interactions in Proteins with Quantum Chemistry
and Examples of Their Applications in Drug Design', *PLOS ONE*, 10(9), p. e0137113.
Available at: https://doi.org/10.1371/journal.pone.0137113.

Yamada, T. *et al.* (2017) 'Drosophila Dalmatian combines sororin and shugoshin roles in
establishment and protection of cohesion', *The EMBO journal*, 36(11), pp. 1513–1527.
Available at: https://doi.org/10.15252/embj.201695607.

Zhou, H.-X. and Pang, X. (2018) 'Electrostatic Interactions in Protein Structure, Folding,
Binding, and Condensation', *Chemical Reviews*, 118(4), pp. 1691–1741. Available at:
https://doi.org/10.1021/acs.chemrev.7b00305.

1467