Comparative genomics of *Rickettsiella* bacteria reveal variable metabolic traits involved in symbiotic interactions
with arthropods

Authors

Anna M. Floriano[1], Adil El-Filali[1], Julien Amoros[2], Marie Buysse[2], Hélène Jourdan-Pineau[3,4], Hein Sprong[5], Robert
Kohl[5], Ron P. Dirks[6], Peter Schaap[7,8], Jasper Koehorst[7,8], Bart Nijsse[7,8], Didier Bouchon[9], Vincent Daubin[1], Fabrice
Vavre[1], Olivier Duron[2]

1 UMR 5558, Laboratoire de Biométrie et Biologie Evolutive, Université Lyon 1, CNRS, VetAgroSup, Villeurbanne,
France.
2 MIVEGEC, University of Montpellier, Centre National de la Recherche Scientifique (CNRS), Institut de Recherche
pour le Développement (IRD), Montpellier, France.
3 CIRAD, UMR ASTRE, F-34398 Montpellier, France.
4 ASTRE, CIRAD, INRAE, Université de Montpellier, Montpellier, France.
5 Centre for Zoonoses and Environmental Microbiology, Netherlands National Institute for Public Health and the
Environment (RIVM), Bilthoven, Netherlands.
6 Future Genomics Technologies BV, Leiden, Netherlands.
7 UNLOCK Large Scale Infrastructure for Microbial Communities, Wageningen University & Research, Wageningen,
The Netherlands
8 Laboratory of Systems and Synthetic Biology, Wageningen University & Research, Wageningen, The Netherlands
9 EBI, University of Poitiers, UMR CNRS 7267, Poitiers, France.

* Corresponding authors: Anna M. Floriano (annamaria.floriano01@gmail.com), Olivier Duron (olivier.duron@cnrs.fr)

Abstract

Members of the *Rickettsiella* genus (Legionellales) are emerging as widespread bacteria associated with insects, arachnids, and crustaceans. While some *Rickettsiella* strains are highly virulent entomopathogens, others are maternally inherited endosymbionts that manipulate arthropod phenotypes, including the induction of defensive symbiosis and cytoplasmic incompatibility. However, the genomic diversity of *Rickettsiella* remains largely unexplored, and their genetic potential to induce complex phenotypes in arthropods is only partially understood. In this study, we sequenced five novel *Rickettsiella* genomes isolated from three tick species. Through comparative genomics, we observed that *Rickettsiella* members share similar metabolic capabilities, and collectively lack virulence genes from pathogenic Legionellales. Additional analysis of *Rickettsiella* genomes revealed significant variability in metabolic properties related to endosymbiosis. Specifically, their capacity to biosynthesize certain B vitamins and heme varies, suggesting a functional role of some *Rickettsiella* strains in the nutrition of their arthropod hosts. Some *Rickettsiella* genomes harbour homologs of *Wolbachia cif* genes, encoding for the effectors of cytoplasmic incompatibility, suggesting that *Rickettsiella* may use a similar molecular mechanism similar to manipulate the reproduction of their arthropod hosts. Phylogenomics further revealed that tick-borne *Rickettsiella* exhibit distinct evolutionary origins within the genus, indicating that *Rickettsiella* have undergone repeated horizontal transfers between ticks and other arthropods.

Keywords: symbiosis, *cif* genes, cytoplasmic incompatibility, ticks, entomopathogenicity

Introduction

Bacteria of the genus *Rickettsiella* (Legionellales: Coxiellaceae) are emerging as both widespread and genetically diverse in arthropods [1–7]. Historically, *Rickettsiella* have been identified as contagious entomopathogens infecting a broad range of insects, but various pathotypes have also been reported in other arthropods, including arachnids and crustaceans [1–3, 8–11]. In infected arthropods, *Rickettsiella*-induced diseases often replace body cavity tissue with a white iridescent liquid, leading to the death of hosts [9–11], which has spurred research into using entomopathogenic *Rickettsiella* as a biocidal agent [12],[13]. However, further surveys have revealed that non-pathogenic *Rickettsiella* are common in field populations of arthropods [14–18]. Most of these non-pathogenic *Rickettsiella* are predominantly maternally transmitted, and have a variety of repercussions on the arthropod's phenotypes, ranging from providing fitness advantages to manipulating reproduction [16, 19-21]. In aphids, *Rickettsiella viridis* can alter body colour, reducing attractiveness to predators and parasitoids [16, 21], or directly protect from entomopathogenic fungi, significantly lowering aphid mortality [16, 19]. In the poultry red mite *Dermanyssus gallinae*, *Rickettsiella* may be an essential nutritional endosymbiont providing B vitamins that are lacking in the blood diet of its host [22]. In the agricultural spider *Mermessus fradeorum*, a *Rickettsiella* endosymbiont induces cytoplasmic incompatibility (CI), that results in embryonic mortality in the offspring of matings between infected males and uninfected females [20]. This reproductive manipulation reduces the production of uninfected progeny and increases the proportion of infected females (*i.e.,* the transmitting sex), thereby promoting the spread of CI-inducing bacteria within the host population [23]. Overall, members of the genus *Rickettsiella* have evolved complex lifestyles that can significantly impact the ecology and evolution of their arthropod hosts [24].

Rickettsiella is remarkably common in the two main tick families, Ixodidae (hard ticks) and Argasidae (soft ticks), with

at least 15% of species infected and prevalence ranging from 5% to 100% at the population level [4, 5, 25–31].

Rickettsiella also exhibit significant genetic diversity both within and between tick species [4, 5, 25–31]. Indeed,

genetically distinct strains of *Rickettsiella* were detected in populations of the castor bean tick *Ixodes ricinus* and the

polar seabird tick *Ixodes uriae* [25, 26]. The *Rickettsiella*-induced phenotypes in ticks have not been clearly identified

yet, but a *Rickettsiella* endosymbiont was observed in a parthenogenetic laboratory colony of the wood tick *Ixodes*

woodi [30, 31]. Both males and females are normally present in this tick species, hence *Rickettsiella* might manipulate

the tick reproduction to induce asexuality [30]. A bacterium was identified in *I. ricinus*, and further in the blacklegged

tick *Ixodes scapularis* [32] as a new genus and species, *Diplorickettsia massiliensis* [33], but it was further

synonymised as *Rickettsiella massiliensis* [34] since the bacterium is nested within the current *Rickettsiella* genus [31].

Rickettsiella massiliensis has been proposed as a potential agent of opportunistic infections in humans [35], but no clear

clinical cases have been reported to date.

Rickettsiella is currently among the lesser-known tick-borne bacteria [24], whereas other Legionellales, particularly

those that infect humans, are better understood. The order *Legionellales* includes two families, *Legionellaceae* and

Coxiellaceae, with several clinically significant microorganisms, such as *Legionella pneumophila,* the causative agent

of Legionnaires' disease, and *Coxiella burnetii,* responsible for Q fever, both of which have been the focus of extensive

genomic studies [24, 36]. These pathogenic *Legionellales* have evolved specific mechanisms to infect mammalian

alveolar macrophages, such as a *Dot/Icm* type IV secretion system to translocate effectors and inhibit host cell apoptosis

[37, 38]. However, not all Legionellales are pathogens: Some are B vitamin provisioning endosymbionts essential for

the nutrition of arthropods with an obligate blood-feeding habit [39, 40], including *Coxiella*-like endosymbionts in ticks

[41–43] and *Legionella polyplacis* in rat lice [22, 24, 36, 44]. Blood is nutritionally unbalanced, and these obligate

blood feeders have evolved narrow associations with B vitamin provisioning symbionts, as they themselves cannot

synthesise the essential cofactors and vitamins that are lacking in their diet [39, 40]. Additionally, *Coxiella*-like

endosymbionts produce chorismate, a tryptophan precursor that regulates serotonin biosynthesis, actively promoting

tick feeding activity and blood intake, and modifying behaviour [45].

Compared to other Legionellales, few *Rickettsiella* genomes have been sequenced. They have circular 1.3-1.9 Mb

genomes, smaller than the genomes of other *Legionellales* such as the pathogens *L. pneumophila* (3.4 Mb) and *C.*

burnetii (2.0 Mb), and the members of the genera *Aquicella* (2.5 Mb) and *Berkiella* (3 - 3.6 Mb), associated with

amoebae living in aquatic environments [24, 34, 46]. While *Rickettsiella* genomes are known to encode 1,400-2,200

protein-coding genes [2, 22, 47, 48], no comparative studies have been conducted. So far, the genomes have been used

in phylogenomic studies [34, 42], and a few investigations have been conducted focusing on single strains. Indeed, the

Rickettsiella genome of the poultry red mite *D. gallinae* contains more than 300 pseudogenized protein-coding

sequences, but has retained several B vitamin biosynthesis pathways, suggesting the importance of these pathways in

evolution of a nutritional endosymbiosis [22]. Other examples include the genomes of *R. grylli,* infecting isopods, and

R. viridis, infecting aphids, which both contain a complete *Dot/Icm* type IV secretion system [47, 48]. However, many

phenotypically characterised *Rickettsiella* strains, as the one inducing CI in spiders [20], have never been sequenced.

Among tick-borne *Rickettsiella*, only *R. massiliensis* has had its genome sequenced, though it was published as a brief

genome announcement that included only a basic genome analysis [2].

In this study, we present five novel genomes of *Rickettsiella* naturally infecting ticks, retrace their evolutionary history,

characterise their genomic content, and infer their metabolic capacities. To this aim, we sequenced five microbial

metagenomes, including three from *I. ricinus*, one from *Ornithodoros erraticus* and one from *Ornithodoros*

phacochoerus. We then compared these genomes with published genomes of *Rickettsiella* and representative

Legionellales, both pathogenic (*L. pneumophila, C. burnetii*) and symbiotic (*Coxiella*-like endosymbionts of ticks), to

assess gene content and metabolic similarities. We further used these genomic analyses to predict the putative

phenotypes of these *Rickettsiella* on their arthropod hosts.

Methods

Sample collection, DNA extraction and sequencing

Specimens of three tick species, *I. ricinus, O. erraticus* and *O. phacochoerus*, were collected in this study. Different

methods for DNA extraction and sequencing were applied depending on tick species. For *I. ricinus*, questing ticks were

collected in September 2020 by dragging in "De Dorschkamp", a deciduous forest east of Wageningen, the Netherlands

(51°58'39.5"N, 5°41'59.3"E). Species and stage of collected ticks were morphologically identified using an

identification key [49]. A first metagenome was sequenced from a pool of twelve males, leading to the recovery of a

genome of *Rickettsiella* later designated as Iric. Another batch of questing *I. ricinus* was collected in September 2021

by dragging in "De Amsterdamse Waterleiding Duinen", a coastal sand dune area in the Netherlands (52°32'79.9"N,

4°54'38.6"E). From this batch, DNA from two females was separately extracted and sequenced, and two *Rickettsiella*

genomes, designated as Ird1 and Ird6, were recovered among genomes of other tick-borne bacteria [50]. *Ornithodoros*

phacochoerus specimens were sampled from warthog (*Phacochoerus africanus*) burrows in the Coutada 9 Game

Reserve in Mozambique (17°42'33.1"S 33°50'29.5"E) in 2021. *Ornithodoros erraticus* specimens were obtained from a

laboratory colony derived from field species collected in a pigsty in the Alentejo region of Portugal in 2016. For *O.*

phacochoerus (Opha) and *O. erraticus* (Oerr), genomic DNA extractions enriched in endosymbiont DNA were obtained

as previously described [41, 51]. Briefly, Malpighian tubules and ovaries were dissected from four ticks of *O.*

phacochoerus and three ticks of *O. erraticus*. The organs were pooled and homogenised in 100µl sterile double-distilled

water. The obtained homogenate was diluted and incubated for 1h at 20°C in 10ml sterile double-distilled water. To

remove host nuclei and other cell debris, the homogenate was filtered using a 5µm Minisartfilter (Sartorius). The

remaining cells in the homogenate were pelleted by centrifugation (15min at 20,000 x g at 4°C). Total genomic DNA

(gDNA) was extracted from the obtained pellet using the DNeasy Blood and Tissue Kit (Qiagen). The obtained gDNA

was quantified on Qubit using the dsDNA high-sensitivity kit (Invitrogen).

In all cases, a shotgun metagenomics approach was applied, obtaining reads belonging to the ticks, the *Rickettsiella*

strains, and any other microbial organisms infecting the same ticks. For the metagenomes Iric, Ird1, and Ird6, DNA

extracted from whole tick bodies was first sequenced using PromethION flow cells (FLO-PRO002) with the Ligation

Sequencing Kit (SQK-LSK110) from Oxford Nanopore Technologies, and simultaneously sequenced on the NovaSeq

6000 platform with the Nextera DNA Flex Library Prep Kit (Illumina). For the metagenomes Oerr and Opha,

endosymbionts DNA enriched samples were sequenced using HiSeq2000 technology using the TruSeq Nano DNA

library construction and HiSeq SBS v3 kits (Illumina).

Genome assembly

For the Oerr and Opha metagenomes, the reads were subjected to a modified version of the Blobology pipeline [52],
available on github. The modified version allows an initial filtering of the host reads using the genome of *I. scapularis*
(GCA_016920785.2) as reference, to ease the toll on computing resources, and includes the use of Python3, SPAdes
v.3.15.4, bowtie2 v.2.1.0, blast v.2.12.0. Finally, the obtained genome assemblies were refined with Bandage v.0.8.1
[53], by manual removal of contigs unrelated to the assembly graph with no BLAST hits against the *R. viridis* genome
as reference (GCF_003966755.1).
For the metagenomes Iric, Ird1 and Ird6, Minimap2 vs. 2.24 was used to map all nanopore reads against five known
Rickettsiella genome sequences (GCF_000168295.1_ASM16829v1, GCF_000257395.1_ASM25739v1,
GCF_001881495.1_ASM188149v1, NZ_AP018005.1 and NZ_CP079094.1). Reads that mapped against the
Rickettsiella reference genomes were collected using Samtools vs. 1.15.1 and then *de novo* assembled using Flye vs.
2.9.2_b1786 (error rate 0.03), which resulted in 43 contigs with sizes ranging from 0.5 kb to 1.81 Mb. Medaka vs. 1.7.2
was used to polish the contigs based on the filtered *Rickettsiella* nanopore reads. Blastn against the *Rickettsiella*
reference genomes confirmed that the 1.81 Mb contig corresponded to a near-complete genome assembly.
147
The completeness of retrieved Metagenome-Assembled Genomes (MAGs hereafter) was estimated using miComplete
v.1.1.1 [54] (--hmms Bact105). The completeness estimation was performed on a dataset including 12 representative
species of *Legionellales* available on NCBI and using the larger genome of the aquatic *Legionellales* "*Candidatus*
Berkiella aquae" (GCF_001431295.2) as reference (Supplementary Table 1). This dataset of organisms was used in
several analyses in our study and we will therefore refer to it as "Dataset 1". The creation of this dataset was completed
on February 1st, 2024. Three *R. isopodorum* and one *R. grylli* genome sequences were excluded from the dataset as they
do not show significant differences in terms of sequence similarity (ANI higher than 98, calculated with the ANI
calculator implemented in EZBioCloud [55]) and gene content (no strain specific ortholog clusters) when compared to
their reference genomes. A summary of the organisms included in Dataset 1 and their hosts is presented in Table 1,
while more information about the dataset, including the average nucleotide identity among genomes, can be found in
Supplementary Table 1.
159
160

Organism	Host	Host generic
Aquicella lusitana	Aquatic environment	Aquatic
"Candidatus Berkiella aquae"	*Acanthamoeba polyphaga*	Amoeba
"Candidatus Berkiella cookevillensis"	*Acanthamoeba polyphaga*	Amoeba
Coxiella burnetii RSA 493	*Dermacentor andersoni*	Tick
Coxiella endosymbiont of Amblyomma sculptum strain CeAS-UFV	*Amblyomma sculptum*	Tick
Coxiella endosymbiont of Ornithodoros maritimus CLEOmar	*Ornithodoros maritimus*	Tick
Legionella pneumophila	*Homo sapiens*	Human
Rickettsiella endosymbiont of Dermanyssus gallinae	*Dermanyssus gallinae*	Mite
Rickettsiella grylli	Armadillium sp.	Isopod
Rickettsiella isolated from Ixodes ricinus, Ird1 strain	*Ixodes ricinus*	Tick
Rickettsiella isolated from Ixodes ricinus, Ir_d6 strain	*Ixodes ricinus*	Tick
Rickettsiella isolated from Ixodes ricinus, Iric strain	*Ixodes ricinus*	Tick

Rickettsiella isolated from Ornithodoros erraticus, Oerr strain	*Ornithodoros erraticus*	Tick
Rickettsiella isolated from Ornithodoros phacochoerus, Opha strain	*Ornithodoros phacochoerus*	Tick
"Candidatus Rickettsiella isopodorum"	*Trachelipus rathkii*	Isopod
Rickettsiella massiliensis 20B	*Ixodes ricinus*	Tick
"Candidatus Rickettsiella viridis"	*Acyrthosiphon pisum*	Hemipteran

Table 1: Summary of Dataset 1.

Genome annotation

Genomes were annotated with Prokka v.1.14.6 [56] using the "compliant" mode and default parameters. The resulting

annotations in GenBank format, together with the other 12 annotated *Legionellales* of Dataset 1 downloaded from

NCBI, were subjected to pseudogene analysis with PseudoFinder v.1.1.0 [57] with default parameters. The counts of

intact and pseudo- genes have been graphically represented with an in-house R script available on github. KEGG

orthology assignments of intact genes and pseudogenes of each genome were predicted using BlastKoala v.3.0 [58].

The outputs were then merged using in-house scripts available on github and github. Finally, we screened for mobile

elements with MobileElementFinder v.1.1.2 [59] using the database MGEdb v.1.1.1 and default parameters.

Phylogeny of *Rickettsiella* organisms

The phylogenetic relationships within Dataset 1 were inferred through a phylogenomic approach on single copy

panorthologous nucleotide sequences using *Buchnera aphidicola* (GCF_003099975.1) as an outgroup. The pipeline is

available on github and includes the use of the tools OrthoFinder v.2.5. [60], SeaView v.5.0.5 [61], ModelTest-NG

v0.2.0 [62], RAxML (raxmlHPC-PTHREADS v.8.2.12, [63], 100 bootstraps), and in-house scripts in Python3, bash,

and R. All of the softwares have been run in default settings, except for RAxML, where the model was set to GTR+G4.

Additionally, in order to catch a glimpse on *Rickettsiella* evolution on a larger scale, a second dataset (Dataset 2) was

created by adding 16S rRNA sequences (n = 446) of members of the *Rickettsiella*, *Aquicella*, and *Berkiella* genera.

These sequences were downloaded from SILVA RefNR SSU r138.1 to the 16S rRNA sequences of the organisms of

Dataset 1, the 16S rRNA sequence from the CI-inducing *Rickettsiella* strain from spiders (MK627517.1), and the 16S

rRNA sequence of *Buchnera aphidicola* str. Sg (NR_145527.1) predicted with barrnap v.0.9 [64]. The sequences were

aligned with MUSCLE v3.8.1551 [65] and the phylogeny was inferred with RAxML (100 bootstraps, rapid bootstrap

random number seed: 1234, random number seed for the parsimony inferences: 123). The obtained 16S rRNA-based

tree was then pruned using iTOL v.6.9 [66] to ease tree visualisation.

Comparative genomics

In order to obtain a description of common and strain-specific genomic features and metabolic capabilities of

Rickettsiella, we analysed the clusters of orthologous intact genes detected for Dataset 1, regardless of phylogenetic

relationships. The OrthoFinder output table from which the single copy panorthologs used in the phylogenomics-based

approach for the phylogeny inference (Orthogroups.GeneCount.tsv, Supplementary Table 2) was analysed for

intersections using the R package UpSetR [67], and the results were graphically represented using the same tool. All the

commands are available on github.

Particular attention was given to clusters uniquely shared by (i) all *Rickettsiella* species and strains, (ii) *Rickettsiella*

strains isolated from hematophagous arthropods, for traces of adaptation to host blood feeding (iii) *Rickettsiella* strains

isolated from phytophagous/detritivorous hosts, for traces of adaptation to other diets (iv) *Rickettsiella* isolated from the

same host, for adaptation to hosts/environments and (v) *Rickettsiella* and *Coxiella* isolated from ticks, for traces of

adaptation to tick hosts. We also investigated gene orthology in respect to the phylogenetic clusters that appeared well-

defined in the phylogenetic trees. Orthology was performed using OrthoFinder on all translated proteins encoded in

genomes, as well as on translated sequences of intact genes and pseudogenes as previously predicted with

PseudoFinder, in order to obtain more information about the evolution of the strains. The results were graphically

represented as Venn diagrams using the R package VennDiagram [68], and the KEGG and Prokka annotations of the

clusters were obtained (commands available on GitHub).

We also analysed specific pathways that could be involved in nutritional symbioses with blood-feeding arthropods (B

vitamins, chorismate and heme metabolism [45, 51, 69, 70]), using subsets of Supplementary Table 3 to obtain

graphical representation of the gene presence/absence/pseudogenization along the phylogenomics-obtained tree with an

in-house R script available on github. The analysis showed differences in some of the pathways, we thus aligned the

amino acid sequences of the said genes, with MUSCLE and inferred their respective phylogeny with RAxML (100

bootstraps, rapid bootstrap random number seed: 1234, random number seed for the parsimony inferences: 123).

We further checked for the presence of virulence factors known in *Legionellales* [37, 38]. Protein sequences of

virulence factors annotated in *Legionella longbeachae* NSW150 (NC_013861), *L. pneumophila* str. Corby

(NC_009494) and *L. pneumophila* subsp. *pneumophila* str. Philadelphia 1 (NC_002942) were downloaded from VFDB

[71] and used as queries in a blast search against the translated sequences of intact genes and pseudogenes of Dataset 1.

The phylogeny of the identified virulence factors was inferred using the same pipeline applied for the phylogeny of the

pathways potentially involved in the nutritional symbiosis (see above).

Finally, we also looked for the presence of CI genes. The CI phenotype is encoded by a pair of syntenic genes, *cifA* and

cifB, whose direct genotype-phenotype association has been demonstrated in *Wolbachia* [72, 73]. *cifA* and *cifB* possess

homologs that have been categorised into five phylogenetic clusters [74–77], although type V has recently been split

into type V-to-X [77]. To determine the presence of *cif* gene homologs on the *Rickettsiella* genomes, we used

Orthofinder to search for specific orthologs of the *cif* gene groups identified in a previous study [74]. These searches

were conducted on a database of CDS from Dataset 1, as well as from published *Wolbachia,* known to induce CI, and

Rickettsia genomes (Supplementary Table 4). Protein domains were predicted for the *cifA* and *cifB* homologs detected

using the HHpred webserver (https://toolkit.tuebingen.mpg.de/#/tools/hhpred/, [78]) with default parameters [79], and

using the following databases: SCOPe70 v.2.08, Pfam v.36.0, SMART v.6.0, and COG/KOG v.1.0. The sequences of

each detected domain were retrieved and manually aligned using Clustal Omega [80], independently for each *cifA* and

cifB homologs, to complete the domain detection of some false negatives generated by HHpred. Visualisation of protein

domains within the *cif* genes was performed using RStudio v4.3.1 with the ggplot2 v.3.4.4 [81], cowplot v1.1.1 [82],

and gridExtra v.2.3 [83] packages. A maximum likelihood phylogeny of *cifA* and *cifB* pair genes has been inferred on

the basis of three conserved domains (RNA-binding-like, AAA-ATPases-like and PD-(D/E)XK nucleases)

independently aligned and concatenated.

Results

Microbial metagenomes

We sequenced and assembled three microbial metagenomes from *I. ricinus* (Ird1, Ird6 and Iric), one from O. *phacochoerus* (Opha) and one from *O. erraticus* (Oerr). We obtained an average of 71.86 Gb of paired-end reads (from 2.2 gb to 192 bp) of 150 nt in length for each sample (see Supplementary Table 1 for details). For each metagenome, we extracted the reads belonging to the target organisms and obtained the complete sequences of five novel *Rickettsiella* MAGs (see Supplementary Table 1 for details). The *Rickettsiella* MAGs are all of similar size (~1.7 Mb), with the exception of *Rickettsiella* Oerr MAG (~1.3 Mb), and they are all estimated to be complete. However, the estimation attributes to *Rickettsiella* strain Oerr MAG lower completeness scores (0.9048; weighted completeness 0.8109) among the MAGs sequenced in this study. Lower scores are expected for endosymbiotic, host-restricted bacteria, whose genomes undergo erosion [84–87], and observed in *Rickettsiella* relatives, as the *Coxiella*-like-endosymbionts of ticks. The completeness score of *Rickettsiella* strain Oerr MAG thus reflects more the genomic erosion in *Rickettsiella* than a misassembled genome. The GC content varies between 36% and 39%, values also consistent with those from endosymbiotic bacteria. Metagenomes also contain reads of other bacterial endosymbionts as *Midichloria* and *Francisella*-like endosymbiont for *I. ricinus* and *O. phacochoerus*, respectively (data not shown). These endosymbionts were not further considered in this study. No other endosymbiont than *Rickettsiella* was detected for *O. erraticus*.
The annotation of the five *Rickettsiella* MAGs revealed a somewhat similar number of functional genes, ranging from 1,268 to 1,473 (Supplementary Table 1). Including all *Rickettsiella* genomes shows that the number of functional genes is between 1,189 for *R. isopodorum* and 1,707 for the *Rickettsiella* endosymbiont of *D. gallinae* (Supplementary Table 1, Supplementary Figure 1). *Rickettsiella* genomes display similar levels of pseudogenization to other *Legionellales*, though some variation is observed. For instance, higher numbers of pseudogenes were predicted for the *Rickettsiella* MAGs Iric and Oerr (340 and 237 respectively), *R. massiliensis* 20B (508) and a few other *Legionellales*, including *Coxiella*-like endosymbiont of *Ornithodoros maritimus* strain CLEOmar (778). In contrast, lower numbers of pseudogenes were predicted in other *Legionellales*, such as the *Coxiella*-like endosymbiont of *Amblyomma sculptum* strain CeAS-UFV (31) (Supplementary Table 1, Supplementary Figure 1). Concerning mobile elements, only one was detected in the analysed dataset, and only in *R. grylli* (Supplementary Table 1). Both recently host-restricted bacteria and long term symbionts generally possess a low number of mobile elements in their genomes, although recent symbionts are usually characterised by a greater number. However, long term symbionts are additionally characterised by fewer pseudogenes and smaller genome sizes [85] than the *Rickettsiella* strains here investigated. These features taken together suggest a relatively recent infection of the hosts by *Rickettsiella*.
Most Average Nucleotide Identities (ANIs) between the five *Rickettsiella* MAGs range in similar values, varying from 69.39 to 72.61%, suggesting that they may be distinct species (Supplementary Table 1). A notable exception is the 99.77% ANI between Iric and Ird6, showing that these two *Rickettsiella* MAGs are very similar. The five *Rickettsiella* MAGs have ANIs ranging from 69.36-to-98.46% with other *Rickettsiella* genomes. Indeed, ANI is 98.46% between Iric/Ird6 and *R. massiliensis*, and 98.26% between Ird1 and *R. isopodorum* (Supplementary Table 1).

Phylogeny of *Rickettsiella*

Phylogenomic analysis based on 92 single-copy panorthologs showed that *Rickettsiella, Coxiella, Legionella, Aquicella,* and *Berkiella* genera cluster together in a robust monophyletic clade forming the Legionellales order (Figure 1). The inner tree topology is also congruent with the known genera of Legionellales [24]. Indeed, genomes of *Rickettsiella, Coxiella,* and *Berkiella* cluster together according to their assigned genus and form robust monophyletic subclades, with *Aquicella* and *Coxiella* forming sister genera of *Rickettsiella*. Two *Rickettsiella* MAGs obtained in this study from *I. ricinus,* Iric and Ird6, cluster with *R. massiliensis* sharing a high ANI from 98.5 to 99.8% (node 15, Figure 1B). The three other *Rickettsiella* MAGs obtained in this study cluster with diverse *Rickettsiella* species: Opha with the defensive symbiont of aphid, *R. viridis,* and the putative nutritional *Rickettsiella* symbiont of poultry red mite (node 13), Ird1 with *R. isopodorum* (node 12) and Ierr with a group formed by *R. grylli, R. isopodorum* and Irid6 (node 10, Figure 1B). The *Rickettsiella* found in different tick species are thus distantly related and do not form a tick-specific clade (Figure 1).

The 16S rRNA-based phylogeny shows a well-supported monophyletic *Rickettsiella* clade, confirming the robustness of the genus (Supplementary Figure 2). No *Rickettsiella* subclade specific to ticks are evident along the tree: the *Rickettsiella* found in ticks are rather scattered among *Rickettsiella* of woodlice, spiders and insects and associated with diverse phenotypic effect as nutritional symbiosis, defensive symbiosis and entomopathogenicity (Supplementary Figure 2). Particularly striking is the separation of *Rickettsiella* strain Ird1, isolated from an *I. ricinus* individual, and the other strains isolated from *I. ricinus* individuals (Iric, Ird6, and *R. massiliensis* 20B [2]). The Ird1 *Rickettsiella* strain appears to be phylogenetically closer to a *Rickettsiella* strain isolated from *Proasellus aquaticus,* an aquatic crustacean with a detritivorous diet. However, the inner topology of the genus *Rickettsiella* based on 16S rRNA gene sequences remains too poorly resolved in many other cases (as shown by low support values) for inferring the exact relatedness between all *Rickettsiella* from different arthropod species.

Genome comparisons

***Rickettsiella* genome stability**

No mobile elements were detected by MobileElementFinder in the *Rickettsiella* genomes apart from one insertion sequence in *R. grylli*.

Orthology analysis

The analysis of the intact orthologous genes detected for Dataset 1 (Figure 2, Supplementary Figure 3, Supplementary Table 2) indicates huge variation among Legionellales, with just 135 clusters shared among all *Legionellales,* most of which involved in basic bacterial processes. Only 25 clusters of intact genes are shared exclusively by members of the genus *Rickettsiella,* most of them associated with no function (Supplementary Table 2). No clusters of intact genes specific to *Rickettsiella* and *Coxiella* strains isolated from hematophagous arthropod hosts were detected. We further performed a phylogeny-driven analysis of Dataset 1, focusing on the phylogenetic clusters that could be identified on the phylogenomics-based tree shown in Figure 1. The core *Rickettsiella* genome appears to contain 606 clusters of orthologs, that drop to 362 when pseudogenes are removed from the analysis; core pseudogenes in the *Rickettsiella* genus (29 clusters of pseudogenized sequences) appear to be mainly associated with transcription and regulation functions (Supplementary Table 5).

In Figure 1, node 10 supports the cluster encompassing two previously described and potentially pathogenic *Rickettsiella* species isolated from isopods (*R. grylli* [3] and *R. isopodorum* [88]), and two of the newly *Rickettsiella* strains identified in ticks. Notably, these two new strains were isolated from very distant ticks, both in terms of host species and in geographic location: *Rickettsiella* Ird1 from the hard tick *I. ricinus* collected in the Netherlands and *Rickettsiella* Oerr from the soft tick *O. erraticus* collected in Mozambique (Supplementary Table 1). The core genome of these organisms contains 823 clusters of orthologs, which drop to 704 when pseudogenes are not considered in the analysis (Figure 2). Furthermore, when only intact genes are retained in the orthology inference, *R. isopodorum* does not present any specific genes, the number of specific clusters of orthologs slightly decreases in the *Rickettsiella* strain Oerr (16 to 13), whereas the number of clusters of orthologs specific to *Rickettsiella* strain Ird1 increases, and remains unvaried for *R. grylli*. Of the 704 clusters of intact genes shared by the members of this node, 198 are uncharacterised proteins, and 27 appear to be involved in secretion of effectors and pumps. Furthermore, five were annotated by Prokka as multidrug resistance proteins. However, the putative pathogen *R. grylli* possesses 22 additional transport proteins compared to the rest of this dataset. In addition, *Rickettsiella* Oerr uniquely possesses 35 uncharacterised proteins (Supplementary Table 5). Interestingly, *Rickettsiella* Oerr and Ird1 exclusively share 15 clusters of intact orthologs, and 9 clusters of intact orthologs. The intact shared orthologs include 13 uncharacterised protein sequences, a transposase, a methylase, and the T4SS components VirB4 and VirB9; whereas the shared clusters of pseudogenes include one uncharacterised protein and the Cell division coordinator CpoB (Supplementary Table 5).

Node 13 supports the cluster encompassing the host colour-altering and defensive symbiont *R. viridis* (aphid host, [14]), the *Rickettsiella* endosymbiont of *D. gallinae* (poultry red mite host, [89]), and the newly sequenced *Rickettsiella* str. Oerr isolated from the soft tick *O. phacochoerus*. The core genome of this node contains 935 clusters of orthologous intact genes (1061 clusters when pseudogenes are maintained in the analysis; Figure 1B). All the *Rickettsiella* strains encompassed in this node appear to maintain a discrete, albeit low, number of unique clusters. Among others, the core intact genes include T4SS components, ATPase subunits, biotin genes, 79 hypothetical proteins, four ankyrin repeat domain-containing proteins, and 14 DUF domain-containing proteins. Notably, *Rickettsiella* strain Opha appears to share more clusters of orthologous genes with the endosymbiont of *D. gallinae* (65, encoding for hypothetical proteins, several ATPases, DUF containing-proteins, and transcription regulators) than with *R.viridis* (37, most of them being potentially involved in adhesion). However, the *Rickettsiella* endosymbiont of *D. gallinae* and *R. viridis* share a large number of clusters as well (90, including 69 with unknown function). Interestingly, the core pseudogenes include many response regulators and translocators, and the *Rickettsiella* endosymbiont of *D. gallinae* and the *Rickettsiella* strain Opha share no additional pseudogenes, whereas the *Rickettsiella* endosymbiont of *D. gallinae* and *R. viridis* share 9 more clusters of pseudogenes. Finally, the aerobic respiration control sensor protein ArcB is pseudogenized in the strain Opha only, and the endosymbiont of *D. gallinae* contains 10 more unique pseudogenes, that include regulators and transposases (Supplementary Table 5).

Finally, Node 15 encompasses three *Rickettsiella* strains isolated from different individuals of the hard tick *I. ricinus*: *R. massiliensis*, *Rickettsiella* Iric and Ird6. The core of 929 clusters of intact genes (1384 clusters when pseudogenes are included in the analysis) contains mainly hypothetical proteins and house-keeping genes, but also genes involved in energy production. Interestingly, *Rickettsiella* strain Iric exhibits distinctive clusters of orthologous sequences exclusively when pseudogenes are analysed, revealing just a single cluster associated with amino sugar and nucleotide sugar metabolism (Supplementary Table 5). However, the other strains possess only five unique clusters of orthologous intact sequences as well. A relatively high number of intact sequences is shared by any two organisms in these subset of

genomes: 74 clusters between *R. massiliensis* and *Rickettsiella* str. Ird6, 82 between *R. massiliensis* and *Rickettsiella* str.

Iric, and 314 between *Rickettsiella* str. Iric and *Rickettsiella* str. Ird6. Most sequences shared by *R. massiliensis* and the

strain Iric are annotated as hypothetical proteins, whereas most sequences shared by *R. massiliensis* and the strain Ird6

have an annotated function, most of them being housekeeping genes and genes encoding for chaperons. The

Rickettsiella Iric and Ird6 share mostly uncharacterised proteins, but also several dehydrogenases, ATPase synthase

subunits A and B, and a queuosine precursor transporter (Supplementary Table 5).

Metabolic capabilities

The annotation of metabolic pathways (Supplementary Table 3) shows overall similar metabolic capabilities among the

members of the *Rickettsiella* genus. Major differences can be identified, however, in specific pathways: the respiration

and ATP production appears to be most degraded in *Rickettsiella* strain Iric and R. *massiliensis* 20B, *i.e.*, the two

Rickettsiella strains with the higher number of predicted pseudogenes that form a monophyletic cluster together with the

Rickettsiella Ird6. Furthermore, B vitamin and heme pathways show significant differences, appearing most degraded in

R. *massiliensis* 20B, but still relatively conserved in *Rickettsiella* strain Iric (*e.g.*, biotin, folate). Strikingly, the heme

biosynthesis pathway is absent in *Rickettsiella* strain Oerr, which has also lost most of the thiamine biosynthesis

pathway, yet retains a unique segment of this pathway not found in other *Rickettsiella* strains sequenced in this study.

However, although it is not possible to fully reconstruct the evolution of these sequences, an online blast search and

inferred phylogenies for each gene shows that they are of Gammaproteobacteria origin (Supplementary Table 3, five

best hits shown, and Supplementary Figure 4), often found in *Legionella* spp. and in two additional *Rickettsiella*

organisms. This suggests either multiple horizontal transfers from other Gammaproteobacteria or Legionellales to these

Rickettsiella strains, or a loss of the genes in the other *Rickettsiella* strains here analysed.

Finally, while *Coxiella*-like endosymbionts can produce a tryptophan precursor, chorismate, which regulates serotonin

biosynthesis in ticks [44], *Rickettsiella* genomes do not harbour chorismate biosynthesis genes, suggesting that

Rickettsiella infection do not impact tick behaviour through this mechanism.

Identification of virulence factors

Further genomic comparisons showed that abundance of virulence genes varies between Legionellales members (Table

2, Supplementary Table 6). While large repertories of virulence genes associated with adherence, secretion systems,

effectors and exotoxin RtxA are common in *L. pneumophila* genomes, these genes are rare or absent in most other

Legionellales as *Coxiella*-like endosymbionts associated with ticks. Indeed, some virulence genes are absent in

Rickettsiella genomes including the exotoxin rtxA, which is involved in adherence, cytotoxicity, and pore formation in

vertebrate cells by *Legionella* species [90]. *Aquicella lusitana* do not possess the exotoxin as well, suggesting that the

loss of this gene might be ancestral to *Rickettsiella* and *Aquicella*. Of the other factors associated with virulence in

Legionella, *Rickettsiella* genomes conserve the ones involved in adherence, in regulation, and in nutritional metabolism

(particularly, heme metabolism cofactors [91]). Furthermore, the *Rickettsiella* strains maintain the effector delivery

systems, but do not retain the secreted effectors associated with virulence identified in *Legionella*, suggesting that this

machinery could be exploited by *Rickettsiella* to perform different functions. The same occurs in *Aquicella, Berkiella*

and *C. burnetii*, whereas the two *Coxiella*-like endosymbionts present neither the delivery system nor the genes

encoding for its virulence secreted effectors. Furthermore, all the investigated *Legionellales*, with the exception of

Legionella and several *Rickettsiella* strains, do not possess the Type IV A secretion system (TIVASS). The strains that maintain most of this system are *R. viridis* and the tick strains Ird1, Oerr and Opha, whereas the *Rickettsiella* endosymbiont of *D. gallinae, R. grylli* and *R. isopodorum* maintain only a few genes involved in this machinery. No evidence of horizontal gene transfer could be obtained. Finally, the enhancer-binding protein (Enh) loci are not found in *Rickettsiella*, whereas they can be found in *Aquicella, Berkiella, C. burnetii*, and *Legionella*.

Category	*Aquicella*	*Berkiella*	*Coxiella burnetii*	*Coxiella endosymbionts*	*Rickettsiella strains*	*Legionella*
Adherence	■	■	■		■	■
Dot/Icm	■	■	■	■		■
Dot/Icm secretedEffectors			■			■
lsp_TIISS	■	■	■		■	■
LvhTIVASS			■		■	■
Exotoxin RtxA		■	■			■
Nutritional Metabolic Function	■	■	■		■	■
PostTranslational Modification	■	■	■		■	■
Regulation	■	■	■		■	■
Enh_loci	■	■	■			■

Table 2: Simplified scheme of presence (light blue) and absence (white) of virulence factors categories in the organisms of Dataset 1. The organisms were grouped by genus, separating only *Coxiella* endosymbionts from the proved pathogenic relative *C. burnetii*. The virulence factor category was considered present if at least three of the representatives of the *Rickettsiella* strains and one of the *Berkiella* and *Coxiella* strains contained at least half of the virulence factors of that category. The search was performed using reference *Legionella* sequences from VFDB as queries in a local blast search (default settings, results were manually curated). Details on the annotation can be found in Supplementary table 6.

Rickettsiella and cytoplasmic incompatibility

We detected homologs of the CI genes *cifA* and *cifB* in the genomes of (i) *Rickettsiella* strain Opha, and (ii) *R. isopodorum* where a CifB-like protein has been previously identified (Figure 3, [92]). Both *Rickettsiella* strains harbour the RNA-binding-like domain in *cifA*, and the AAA-ATPase-like and PD-(D/E)XK nuclease domains in *cifB*, both conserved in all *cif* genes [74, 93]. Additionally, the *cifB* gene in *Rickettsiella* strain Opha has a weakly aligned latrotoxin-like fragment, while *R. isopodorum* has a large TcdA/TcdB pore-forming toxin domain, also found in other *cifB* homologs of *Wolbachia* and *Rickettsia*. Furthermore, we also identified a unique *cifB* homolog (without *cifA*) in a *Rickettsiella* sp. genome reconstructed from eDNA isolated from hot springs and volcanic lakes (GCA_037439325.1) and in the distant relative "*Ca.* Aquirickettsiella gammari" (GCA_002290645.2).

Both the detected *cifA* and *cifB* homologs of *Rickettsiella* cluster within the Type V *cif* paraphyletic group, which is the most diverse in terms of size and domain composition [74, 93]. Both the *cif* genes from *Rickettsiella* strain Opha and *R. isopodorum* group together in a highly supported clade comprising *cif* genes from *Rickettsia* and *Wolbachia*,

407 originating from at least two supergroups (*w*Meg and *w*Con from supergroup A; *w*Oegib and *w*Bor from supergroup B),
408 supporting the occurrence of gene transfers between these phylogenetically distinct endosymbionts. However, the long
409 branch lengths and low intra-group bootstrap values create uncertainty regarding the exact position of the *Rickettsiella*
410 *cif* genes, challenging the reconstruction of their evolutionary history and origin.

Discussion

We show here that members of the genus *Rickettsiella* share a set of similar metabolic capabilities, as pathways
involved in energy production, including glycolysis and a normal TCA cycle. Most notably, the *Rickettsiella* species
and strains lack most key virulence genes responsible for persistence and replication of pathogenic *Legionella* and
Coxiella in vertebrate cells. Indeed, while the *Rickettsiella* genomes harbour the virulence genes encoding the *Dot*/*Icm*
secretion system, the genes encoding the Legionellales *Dot*/*Icm* secreted effectors and the exotoxin RtxA have been
pseudogenized or are completely absent. However, examination of *Rickettsiella* genomes further reveals a greater
variability in metabolic properties associated with endosymbiosis: They have variable capacity to biosynthesize *de novo*
certain B vitamins (biotin, riboflavin, pyridoxal phosphate, pantothenate) and heme, and some harbour CI effectors,
suggesting that they are engaged in complex interactions with their respective arthropod hosts. Phylogenomics further
confirmed that members of the genus *Rickettsiella* form a robust monophyletic clade of Legionellales associated with
arthropods but that tick-borne *Rickettsiella* do not cluster within a tick-specific subclade. Tick-borne *Rickettsiella* rather
exhibit distinct evolutionary origins in the genus, revealing that *Rickettsiella* have undergone repeated horizontal
transfers between ticks and other arthropods, including aphids and woodlice.

The virulence factors presence/absence pattern across Legionellales reveals key past events in the evolutionary history
of *Rickettsiella*. Indeed, the absence of major Legionellales virulence factors in all *Rickettsiella* strains, including the
putative human pathogen *R. massiliensis* 20B, suggests that these microorganisms are not strongly pathogenic to
mammals as *L. pneumophila* and *C. burnetii*. Concerning the exotoxin RtxA, the closest relative of *Rickettsiella*,
Aquicella lusitana, does not possess the exotoxin as well, suggesting that the loss of this gene might be ancestral to
Rickettsiella and *Aquicella* and extend deep in the evolutionary past of Coxiellaceae. Several *Legionella* virulence
factors are conserved in *Rickettsiella* and to some extent in *Aquicella*, though they may have alternative functions.
Notably, traits such as adherence, stress response regulation, nutritional metabolism (particularly involving heme
cofactors), and the type IV secretion systems are common to both pathogenic and mutualistic bacteria [87,90-97] the
Dot/*Icm* effector delivery system has been exclusively associated only with pathogenesis thus far [42]. Indeed, genes
encoding the *Dot*/*Icm* system are either absent or pseudogenized in *Coxiella*-like endosymbionts associated with
nutritional symbiosis in ticks [42,98]. The conservation of the system *Dot*/*Icm* in *Rickettsiella* suggests either that this
machinery is being exploited by *Rickettsiella* to perform functions other than virulence by secreting different molecular
effectors, or that *Rickettsiella* maintain a degree of virulence but using different virulence effectors than *Legionella*.
Alternatively, since the *Rickettsiella Dot*/*Icm* system lacks a few genes compared to *L. pneumophila* and *C. burnetii*,
this machinery may undergone a gradual degradation and they mirrored a global genome decay, as commonly observed
in endosymbionts of arthropods [84, 85, 87, 99, 100]. Whatever the explanation, virulence against mammals is clearly
being gradually lost in *Rickettsiella,* leaving opportunities for the evolution of other interactions with their arthropod
hosts.

The genomes of *Rickettsiella* consistently harbour genes involved in narrow interactions with arthropods, although

these genes vary between *Rickettsiella* strains and species. Indeed, the biosynthesis genes of several B vitamins (biotin,

riboflavin, and folate) have been found all intact in most *Rickettsiella* genomes, suggesting that they could provide these

nutrients to their arthropod hosts. In this context, *Rickettsiella* may be essential mutualistic partners for hematophagous

arthropods, such as ticks and certain mites, which need these essential B vitamins for their growth and to achieve their

life cycle [39]. Most tick species harbour *Coxiella*-like endosymbionts that ensure obligate B vitamins provisioning,

while certain tick species lacking *Coxiella*-LE instead host *Francisella*-like endosymbionts [39, 101] or *Midichloria*

[102, 103] as we observed in this study in *O. phacocherus* and *I. ricinus*, respectively. However, co-infections (i.e., the

simultaneous infection of *Rickettsiella* with one or more different microorganisms) have been previously documented in

ticks [4, 5, 101], hence, the identification of *Rickettsiella* tick species previously associated with other symbionts is not

surprising. However, in *O. erraticus*, we observed no other endosymbionts than *Rickettsiella* suggesting that this

bacterium is a obligate nutritional partner of this tick species. In addition, certain *Rickettsiella* genomes possess

biosynthesis genes for heme, a nutrient which is essential for regulating protein synthesis and cell differentiation in

animals. Unlike other eukaryotes, ticks lack most of the genes encoding proteins necessary for heme production and

degradation [104–106]. Some *Rickettsiella* endosymbionts could thus potentially influence tick metabolism through the

production of this essential cofactor.

Further examinations of *Rickettsiella* genomes also suggest that at least two *Rickettsiella* strains may be reproductive

parasites of the soft tick *O. phacochoerus* and the woodlouse *Trachelipus rathkei*. Although no phenotypic data are

currently available for these arthropod species, their *Rickettsiella* possess CI-inducing genes with key functional

domains, including a RNA-binding-like domain in *cifA*, a the AAA-ATPase-like and PD-(D/E)XK nuclease domains in

cifB, as observed in all *cif* genes of CI-inducing *Wolbachia* [74, 93]. The observed pattern suggests acquisitions of *cif*

genes in *Rickettsiella* through lateral gene transfers from phylogenetically distant endosymbionts also inhabiting

arthropods, *Rickettsia* and *Wolbachia*. Both are common in arthropods [107, 108], especially *Rickettsia* in ticks [4, 109]

in which co-infections with *Rickettsiella* have been reported [4]. These co-infections create physical proximity between

these endosymbionts and may have facilitated lateral DNA transfers between them [110, 111], which could explain the

intertwined evolutionary history of *cifA* and *cifB* genes among endosymbionts of arthropods.

The phylogenetic diversity pattern strongly suggests that horizontal transfers between arthropod species are key drivers

of *Rickettsiella* spread, dictating global incidence of infections across arthropod communities. Repeated horizontal

transfers results to a widespread and global distribution in diverse arthropod hosts, but also to multiple acquisitions of

unrelated *Rickettsiella* strains in the same host species as in *I. ricinus* (this study) and the polar seabird tick *Ixodes uriae*

[25]. The capacity of *Rickettsiella* to move between host species can thus explain why no clusters specific to tick-borne

Rickettsiella were identified. However, exceptions exist: one uncharacterised protein is shared by all *Rickettsiella*

isolated from *I. ricinus*, the fatty acid oxidation complex subunit alpha is shared by all strains isolated from *I. ricinus*

with the exception of *R. massiliensis* 20 B, five clusters of hypothetical proteins specific to the Oerr strain, and nine

clusters of uncharacterised proteins specific to the Opha strain. While most of these proteins have not been associated

with any function yet, the fatty acid oxidation complex is quite interesting. Indeed, bacteria may exploit fatty acids from

the host and utilise them for their own sustainment [112]. On the other hand, the oxidation of fatty acids might allow

Rickettsiella to produce energy-rich compounds that could be used by *Rickettsiella* itself but also its tick host, aiding the

host's metabolism, as previously suggested for the tick microbiome [113, 114] and *Coxiella*-like endosymbionts [41,
115].

Conclusions

Rickettsiella is a diverse group of endosymbiotic bacteria that exhibit a range of adaptations, from pathogenic to
mutualistic, enabling them to thrive within arthropod cells and manipulate their hosts' biology. Over the past decade,
significant progress has been made in understanding their impact on arthropods and introducing these bacteria into the
field of genomics. However, *Rickettsiella* remains an understudied aspect of tick biology, and a comprehensive
approach is needed to fully understand its interactions with various arthropod hosts. Key questions persist, such as:
How are the global abundance, diversity, and distribution of *Rickettsiella* maintained? Can *Rickettsiella* be effectively
used in pest control? How do they manipulate host reproduction? What roles do they play in the evolution of their
arthropod hosts? Importantly, future research is needed to more thoroughly investigate how *Rickettsiella* affects the
fitness of arthropods, including ticks, and to what extent they can cause diseases in humans.

List of abbreviations

Ca.	*Candidatus*
CI	Cytoplasmic incompatibility
R.	*Rickettsiella*
TCA	Tricarboxylic acid (cycle)
TIVASS	Type IV A secretion system
min	minutes
nt	Nucleotides
ANI	Average Nucleotide Identity

Declarations

The authors have complied with all ethical standards required for conducting this research. Consents and approvals are
not applicable to this research. The genomes are available on ENA under project number PRJEB70514 (Ird1 and Ird6),
PRJEB82506 (Oerr and Opha), and PRJNA816462 (Iric). All the bioinformatics pipelines are available on GitHub
(https://github.com/annamariafloriano/RickettsiellaComparative). This work was funded by French Agence Nationale
de la Recherche (ANR, France, ref. ANR-21-CE02-0002, Laboratoire d'Excellence CEBA, ref. ANR-10-LABX-25-01
and LabEx CeMEB, ref. ANR-10-LABX-04-01), and by the United States Department of Agriculture (USDA, USA,
grant no. 2019-67015-28981; https://www.asf-nifnaf.org/).

Acknowledgements

We thank Carlos João Quembo, Florian Taraveau and Maxime Duhayon for their contribution in the sample collection.

References

1. Leclerque A, Kleespies RG, Ritter C, Schuster C, Feiertag S. Genetic and electron-microscopic characterization of "Rickettsiella agriotidis", a new Rickettsiella pathotype associated with wireworm, Agriotes sp. (Coleoptera: Elateridae). Curr Microbiol. 2011;63:158–63.

2. Mathew MJ, Subramanian G, Nguyen T-T, Robert C, Mediannikov O, Fournier P-E, et al. Genome sequence of Diplorickettsia massiliensis, an emerging Ixodes ricinus-associated human pathogen. J Bacteriol. 2012;194:3287.

3. Leclerque A. Whole genome-based assessment of the taxonomic position of the arthropod pathogenic bacterium Rickettsiella grylli. FEMS Microbiol Lett. 2008;283:117–27.

4. Duron O, Binetruy F, Noël V, Cremaschi J, McCoy KD, Arnathau C, et al. Evolutionary changes in symbiont community structure in ticks. Mol Ecol. 2017;26:2905–21.

5. Duron O, Noël V, McCoy KD, Bonazzi M, Sidi-Boumedine K, Morel O, et al. The Recent Evolution of a Maternally-Inherited Endosymbiont of Ticks Led to the Emergence of the Q Fever Pathogen, Coxiella burnetii. PLoS Pathog. 2015;11:e1004892.

6. Yaripour S, Bandekar M, Vihavainen J, More K, Eronen A, Durão MF, et al. Effects of Nanoplastics on the Swimming Behaviour and Gut Microbiome of the Freshwater Amphipod Gammarus Pulex. 2024. https://doi.org/10.2139/ssrn.4876517.

7. Akter R, Punom NJ, Enan Eshik MM, Hossain MS, Momtaz S, Sultana S, et al. Gut Microbiota Exhibits Discrete Patterns in the Presence of Wssv and Vpahpnd in Black Tiger Shrimp Penaeus Monodon. 2024. https://doi.org/10.2139/ssrn.4846014.

8. Cordaux R, Paces-Fessy M, Raimond M, Michel-Salzat A, Zimmer M, Bouchon D. Molecular characterization and evolution of arthropod-pathogenic Rickettsiella bacteria. Appl Environ Microbiol. 2007;73:5045–7.

9. Zchori-Fein K. Rickettsiella, Intracellular Pathogens of Arthropods. In: Taylor & Francis. 1st Edition. CRC Press; 2011.

10. Crotti E, Rizzi A, Chouaia B, Ricci I, Favia G, Alma A, et al. Acetic acid bacteria, newly emerging symbionts of insects. Appl Environ Microbiol. 2010;76:6963–70.

11. Bouchon D, Zimmer M, Dittmer J. The Terrestrial Isopod Microbiome: An All-in-One Toolbox for Animal-Microbe Interactions of Ecological Relevance. Front Microbiol. 2016;7:1472.

12. Zchori-Fein E, Bourtzis K. Manipulative Tenants: Bacteria Associated with Arthropods. CRC Press; 2011.

13. Website. https://www.researchgate.net/publication/285958041_Emerging_entomopathogenic_bacteria_for_insect_pest_management.

14. Tsuchida T, Koga R, Fujiwara A, Fukatsu T. Phenotypic effect of "Candidatus Rickettsiella viridis," a facultative symbiont of the pea aphid (Acyrthosiphon pisum), and its interaction with a coexisting symbiont. Appl Environ Microbiol. 2014;80:525–33.

15. Nikoh N, Tsuchida T, Maeda T, Yamaguchi K, Shigenobu S, Koga R, et al. Genomic Insight into Symbiosis-Induced Insect Color Change by a Facultative Bacterial Endosymbiont, " Rickettsiella viridis." MBio. 2018;9.

16. Tsuchida T, Koga R, Horikawa M, Tsunoda T, Maoka T, Matsumoto S, et al. Symbiotic bacterium modifies aphid
body color. Science. 2010;330:1102–4.

17. Iasur-Kruh L, Weintraub PG, Mozes-Daube N, Robinson WE, Perlman SJ, Zchori-Fein E. Novel Rickettsiella
bacterium in the leafhopper Orosius albicinctus (Hemiptera: Cicadellidae). Appl Environ Microbiol. 2013;79:4246–52.

18. Gauthier J-P, Outreman Y, Mieuzet L, Simon J-C. Bacterial Communities Associated with Host-Adapted
Populations of Pea Aphids Revealed by Deep Sequencing of 16S Ribosomal DNA. PLoS One. 2015;10:e0120664.

19. Łukasik P, van Asch M, Guo H, Ferrari J, Godfray HCJ. Unrelated facultative endosymbionts protect aphids against
a fungal pathogen. Ecol Lett. 2013;16:214–8.

20. Rosenwald LC, Sitvarin MI, White JA. Endosymbiotic causes cytoplasmic incompatibility in a spider host. Proc
Biol Sci. 2020;287:20201107.

21. Polin S, Le Gallic J-F, Simon J-C, Tsuchida T, Outreman Y. Conditional Reduction of Predation Risk Associated
with a Facultative Symbiont in an Insect. PLoS One. 2015;10:e0143728.

22. Price DRG, Bartley K, Blake DP, Karp-Tatham E, Nunn F, Burgess STG, et al. A Endosymbiont Is a Potential
Source of Essential B-Vitamins for the Poultry Red Mite,. Front Microbiol. 2021;12:695346.

23. Engelstädter J, Hurst GDD. The ecology and evolution of microbes that manipulate host reproduction. Annu Rev
Ecol Evol Syst. 2009;40:127–49.

24. Duron O, Doublet P, Vavre F, Bouchon D. The Importance of Revisiting Legionellales Diversity. Trends Parasitol.
2018;34:1027–37.

25. Duron O, Cremaschi J, McCoy KD. The High Diversity and Global Distribution of the Intracellular Bacterium
Rickettsiella in the Polar Seabird Tick Ixodes uriae. Microb Ecol. 2016;71:761–70.

26. Garcia-Vozmediano A, Tomassone L, Fonville M, Bertolotti L, Heylen D, Fabri ND, et al. The Genetic Diversity of
Rickettsiella Symbionts in Ixodes ricinus Throughout Europe. Microb Ecol. 2021;84:613–26.

27. Vilcins I-ME, Old JM, Deane E. Molecular detection of Rickettsia, Coxiella and Rickettsiella DNA in three native
Australian tick species. Exp Appl Acarol. 2009;49:229–42.

28. Anstead CA, Chilton NB. Discovery of novel Rickettsiella spp. in ixodid ticks from Western Canada. Appl Environ
Microbiol. 2014;80:1403–10.

29. Carpi G, Cagnacci F, Wittekindt NE, Zhao F, Qi J, Tomsho LP, et al. Metagenomic profile of the bacterial
communities associated with Ixodes ricinus ticks. PLoS One. 2011;6:e25604.

30. Kurtti TJ, Palmer AT, Oliver JH Jr. Rickettsiella-like bacteria in Ixodes woodi (Acari: Ixodidae). J Med Entomol.
2002;39:534–40.

31. Leclerque A, Kleespies RG. A Rickettsiella bacterium from the hard tick, Ixodes woodi: molecular taxonomy
combining multilocus sequence typing (MLST) with significance testing. PLoS One. 2012;7:e38062.

32. Merenstein C, Ward J, Allen D. Diplorickettsia Bacteria in an Ixodes scapularis Tick, Vermont, USA - Volume 26,
Number 5—May 2020 - Emerging Infectious Diseases journal - CDC. https://doi.org/10.3201/eid2605.191135.

33. Mediannikov O, Sekeyová Z, Birg M-L, Raoult D. A novel obligate intracellular gamma-proteobacterium
associated with ixodid ticks, Diplorickettsia massiliensis, Gen. Nov., Sp. Nov. PLoS One. 2010;5:e11478.

34. Saini N, Gupta RS. A robust phylogenetic framework for members of the order Legionellales and its main genera
(Legionella, Aquicella, Coxiella and Rickettsiella) based on phylogenomic analyses and identification of molecular
markers demarcating different clades. Antonie Van Leeuwenhoek. 2021;114:957–82.

35. Subramanian G, Mediannikov O, Angelakis E, Socolovschi C, Kaplanski G, Martzolff L, et al. Diplorickettsia

massiliensis as a human pathogen. Eur J Clin Microbiol Infect Dis. 2011;31:365–9.

36. Duron O, Sidi-Boumedine K, Rousset E, Moutailler S, Jourdain E. The Importance of Ticks in Q Fever

Transmission: What Has (and Has Not) Been Demonstrated? Trends Parasitol. 2015;31:536–52.

37. van Schaik EJ, Chen C, Mertens K, Weber MM, Samuel JE. Molecular pathogenesis of the obligate intracellular

bacterium Coxiella burnetii. Nat Rev Microbiol. 2013;11:561–73.

38. Isberg RR, O'Connor TJ, Heidtman M. The Legionella pneumophila replication vacuole: making a cosy niche

inside host cells. Nat Rev Microbiol. 2008;7:13–24.

39. Duron O, Gottlieb Y. Convergence of Nutritional Symbioses in Obligate Blood Feeders. Trends Parasitol.

2020;36:816–25.

40. Tamarit D. Lifestyle transitions in endosymbiosis. Peer Community in Microbiology. 2023;1:100003.

41. Gottlieb Y, Lalzar I, Klasson L. Distinctive Genome Reduction Rates Revealed by Genomic Analyses of Two

Coxiella-Like Endosymbionts in Ticks. Genome Biol Evol. 2015;7:1779–96.

42. Santos-Garcia D, Morel O, Henri H, El Filali A, Buysse M, Noël V, et al. Genomic changes during the evolution of

the *Coxiella* genus along the parasitism-mutualism continuum. Peer Community Journal. 2023;3.

43. Guizzo MG, Parizi LF, Nunes RD, Schama R, Albano RM, Tirloni L, et al. A Coxiella mutualist symbiont is

essential to the development of Rhipicephalus microplus. Sci Rep. 2017;7:17554.

44. Ríhová J, Nováková E, Husník F, Hypša V. Legionella Becoming a Mutualist: Adaptive Processes Shaping the

Genome of Symbiont in the Louse Polyplax serrata. Genome Biol Evol. 2017;9:2946–57.

45. Zhong Z, Zhong T, Peng Y, Zhou X, Wang Z, Tang H, et al. Symbiont-regulated serotonin biosynthesis modulates

tick feeding activity. Cell Host Microbe. 2021;29:1545–57.e4.

46. Graells T, Ishak H, Larsson M, Guy L. The all-intracellular order Legionellales is unexpectedly diverse, globally

distributed and lowly abundant. FEMS Microbiol Ecol. 2018;94.

47. Nikoh N, Tsuchida T, Maeda T, Yamaguchi K, Shigenobu S, Koga R, et al. Genomic Insight into Symbiosis-

Induced Insect Color Change by a Facultative Bacterial Endosymbiont, "Candidatus Rickettsiella viridis." MBio. 2018.

https://doi.org/10.1128/mbio.00890-18.

48. Wang Y, Chandler C. Candidate pathogenicity islands in the genome of" Rickettsiella isopodorum", an intracellular

bacterium infecting terrestrial isopod crustaceans. PeerJ. 2016;4:e2806.

49. Hillyard PD. Ticks of North-West Europe: Keys and Notes for Identification of the Species. 1996.

50. Lesiczka PM, Azagi T, Krawczyk AI, Scott WT, Dirks RP, Šimo L, et al. Deep sequencing of 16 Ixodes ricinus

ticks unveils insights into their interactions with endosymbionts. bioRxiv. 2024;:2024.04.22.590557.

51. Duron O, Morel O, Noël V, Buysse M, Binetruy F, Lancelot R, et al. Tick-Bacteria Mutualism Depends on B

Vitamin Synthesis Pathways. Curr Biol. 2018;28:1896–902.e5.

52. Kumar S, Jones M, Koutsovoulos G, Clarke M, Blaxter M. Blobology: exploring raw genome data for

contaminants, symbionts and parasites using taxon-annotated GC-coverage plots. Front Genet. 2013;4:237.

53. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of de novo genome assemblies.

Bioinformatics. 2015;31:3350–2.

54. Hugoson E, Lam WT, Guy L. miComplete: weighted quality evaluation of assembled microbial genomes.

Bioinformatics. 2020;36:936–7.

55. Yoon S-H, Ha S-M, Lim J, Kwon S, Chun J. A large-scale evaluation of algorithms to calculate average nucleotide

identity. Antonie Van Leeuwenhoek. 2017;110:1281–6.

56. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30:2068–9.

57. Syberg-Olsen MJ, Garber AI, Keeling PJ, McCutcheon JP, Husnik F. Pseudofinder: Detection of Pseudogenes in

Prokaryotic Genomes. Mol Biol Evol. 2022;39.

58. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG Tools for Functional Characterization

of Genome and Metagenome Sequences. J Mol Biol. 2016;428:726–31.

59. Johansson MHK, Bortolaia V, Tansirichaiya S, Aarestrup FM, Roberts AP, Petersen TN. Detection of mobile

genetic elements associated with antibiotic resistance in Salmonella enterica using a newly developed web tool:

MobileElementFinder. J Antimicrob Chemother. 2020;76:101–9.

60. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics. Genome Biol.

2019;20:238.

61. Gouy M, Tannier E, Comte N, Parsons DP. Seaview Version 5: A Multiplatform Software for Multiple Sequence

Alignment, Molecular Phylogenetic Analyses, and Tree Reconciliation. Methods Mol Biol. 2021;2231:241–60.

62. Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T. ModelTest-NG: A New and Scalable Tool for

the Selection of DNA and Protein Evolutionary Models. Mol Biol Evol. 2020;37:291–4.

63. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies.

Bioinformatics. 2014;30:1312–3.

64. GitHub - tseemann/barrnap: :microscope: Bacterial ribosomal RNA predictor. GitHub.

https://github.com/tseemann/barrnap. Accessed 28 May 2024.

65. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res.

2004;32:1792–7.

66. Letunic I, Bork P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display and annotation

tool. Nucleic Acids Res. 2024. https://doi.org/10.1093/nar/gkae268.

67. Conway JR, Lex A, Gehlenborg N. UpSetR: an R package for the visualization of intersecting sets and their

properties. Bioinformatics. 2017;33:2938–40.

68. Chen H, Boutros PC. VennDiagram: a package for the generation of highly-customizable Venn and Euler diagrams

in R. BMC Bioinformatics. 2011;12:35.

69. Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. Wolbachia as a bacteriocyte-associated nutritional

mutualist. Proc Natl Acad Sci U S A. 2010;107:769–74.

70. Buysse M, Floriano AM, Gottlieb Y, Nardi T, Comandatore F, Olivieri E, et al. A dual endosymbiosis supports

nutritional adaptation to hematophagy in the invasive tick. Elife. 2021;10.

71. Liu B, Zheng D, Zhou S, Chen L, Yang J. VFDB 2022: a general classification scheme for bacterial virulence

factors. Nucleic Acids Res. 2022;50:D912–7.

72. LePage DP, Metcalf JA, Bordenstein SR, On J, Perlmutter JI, Shropshire JD, et al. Prophage WO genes recapitulate

and enhance Wolbachia-induced cytoplasmic incompatibility. Nature. 2017;543:243–7.

73. Beckmann JF, Bonneau M, Chen H, Hochstrasser M, Poinsot D, Merçot H, et al. The Toxin-Antidote Model of

Cytoplasmic Incompatibility: Genetics and Evolutionary Implications. Trends Genet. 2019;35:175–85.

74. Martinez J, Klasson L, Welch JJ, Jiggins FM. Life and Death of Selfish Genes: Comparative Genomics Reveals the

Dynamic Evolution of Cytoplasmic Incompatibility. Mol Biol Evol. 2021;38:2–15.

75. Martinez J, Ant TH, Murdochy SM, Tong L, da Silva Filipe A, Sinkins SP. Genome sequencing and comparative

analysis of Wolbachia strain wAlbA reveals Wolbachia-associated plasmids are common. PLoS Genet.

2022;18:e1010406.

76. Vancaester E, Blaxter M. Phylogenomic analysis of Wolbachia genomes from the Darwin Tree of Life biodiversity

genomics project. PLoS Biol. 2023;21:e3001972.

77. Tan Y, Aravind L, Zhang D. Genomic Underpinnings of Cytoplasmic Incompatibility: CIF Gene-Neighborhood

Diversification Through Extensive Lateral Transfers and Recombination in Wolbachia. Genome Biol Evol. 2024;16.

78. Söding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology detection and structure

prediction. Nucleic Acids Res. 2005;33 Web Server issue:W244–8.

79. Evolutionary Genetics of Cytoplasmic Incompatibility Genes cifA and cifB in Prophage WO of Wolbachia.

Genome Biol Evol. 2019;11:1320.

80. Sievers F, Higgins DG. Clustal Omega for making accurate alignments of many protein sequences. Protein Sci.

2018;27:135–45.

81. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer Science & Business Media; 2009.

82. Wilke CO. Streamlined Plot Theme and Plot Annotations for "ggplot2" [R package cowplot version 1.1.3]. 2024.

83. Auguie B. Miscellaneous Functions for "Grid" Graphics [R package gridExtra version 2.3]. 2017.

84. Moran NA. Microbial minimalism: genome reduction in bacterial pathogens. Cell. 2002;108:583–6.

85. McCutcheon JP, Moran NA. Extreme genome reduction in symbiotic bacteria. Nat Rev Microbiol. 2011;10:13–26.

86. Boscaro V, Kolisko M, Felletti M, Vannini C, Lynn DH, Keeling PJ. Parallel genome reduction in symbionts

descended from closely related free-living bacteria. Nat Ecol Evol. 2017;1:1160–7.

87. Kashkouli M, Castelli M, Floriano AM, Bandi C, Epis S, Fathipour Y, et al. Characterization of a novel Pantoea

symbiont allows inference of a pattern of convergent genome reduction in bacteria associated with Pentatomidae.

Environ Microbiol. 2021;23:36–50.

88. Kleespies RG, Federici BA, Leclerque A. Ultrastructural characterization and multilocus sequence analysis (MLSA)

of "Candidatus Rickettsiella isopodorum", a new lineage of intracellular bacteria infecting woodlice (Crustacea:

Isopoda). Syst Appl Microbiol. 2014;37:351–9.

89. De Luna CJ, Moro CV, Guy JH, Zenner L, Sparagano OAE. Endosymbiotic bacteria living inside the poultry red

mite (Dermanyssus gallinae). Exp Appl Acarol. 2009;48:105–13.

90. Cirillo SL, Bermudez LE, El-Etr SH, Duhamel GE, Cirillo JD. Legionella pneumophila entry gene rtxA is involved

in virulence. Infect Immun. 2001;69:508–17.

91. Shevket SH, Gonzalez D, Cartwright JL, Kleanthous C, Ferguson SJ, Redfield C, et al. The CcmC-CcmE

interaction during cytochrome maturation by System I is driven by protein-protein and not protein-heme contacts. J Biol

Chem. 2018;293:16778–90.

92. Gillespie JJ, Driscoll TP, Verhoeve VI, Rahman MS, Macaluso KR, Azad AF. A Tangled Web: Origins of

Reproductive Parasitism. Genome Biol Evol. 2018;10:2292–309.

93. Beliavskaia A, Tan K-K, Sinha A, Husin NA, Lim FS, Loong SK, et al. Metagenomics of culture isolates and insect

tissue illuminate the evolution of , and symbionts in spp. fleas. Microb Genom. 2023;9.

94. Lomovatskaya LA, Romanenko AS. Secretion Systems of Bacterial Phytopathogens and Mutualists (Review). Appl

Biochem Microbiol. 2020;56:115–29.

95. Costa TRD, Harb L, Khara P, Zeng L, Hu B, Christie PJ. Type IV secretion systems: Advances in structure,

function, and activation. Mol Microbiol. 2021;115:436–52.

96. Cao Y, Miller SS, Dornbusch MR, Castle SS, Lenz P, Ferguson J, et al. Widespread occurrence of Sinorhizobium

meliloti strains with a type IV secretion system. Symbiosis. 2018;75:81–91.

97. Floriano AM, Batisti Biffignandi G, Castelli M, Olivieri E, Clementi E, Comandatore F, et al. The evolution of

intramitochondriality in Midichloria bacteria. Environ Microbiol. 2023;25:2102–17.

98. Buysse M, Duron O. Evidence that microbes identified as tick-borne pathogens are nutritional endosymbionts. Cell.

2021;184:2259–60.

99. Klasson L, Andersson SGE. Evolution of minimal-gene-sets in host-dependent bacteria. Trends Microbiol.

2004;12:37–43.

100. Bennett GM, Moran NA. Heritable symbiosis: The advantages and perils of an evolutionary rabbit hole. Proc Natl

Acad Sci U S A. 2015;112:10169–76.

101. Binetruy F, Buysse M, Lejarre Q, Barosi R, Villa M, Rahola N, et al. Microbial community structure reveals

instability of nutritional symbiosis during the evolutionary radiation of Amblyomma ticks. Mol Ecol. 2020;29:1016–29.

102. Duron O. Nutritional symbiosis in ticks: singularities of the genus Ixodes. Trends Parasitol. 2024;0.

103. Duron O. Nutritional symbiosis in ticks: singularities of the genus Ixodes: (Trends in Parasitology, Published

online June 27, 2024). Trends Parasitol. 2024;40:767.

104. Jia N, Wang J, Shi W, Du L, Sun Y, Zhan W, et al. Large-Scale Comparative Analyses of Tick Genomes Elucidate

Their Genetic Diversity and Vector Capacities. Cell. 2020;182:1328–40.e13.

105. Gulia-Nuss M, Nuss AB, Meyer JM, Sonenshine DE, Roe RM, Waterhouse RM, et al. Genomic insights into the

Ixodes scapularis tick vector of Lyme disease. Nat Commun. 2016;7:10507.

106. Hatalová T, Erhart J, Kopáček P, Perner J. On the haem auxotrophy of the soft tick Ornithodoros moubata. Ticks

Tick Borne Dis. 2023;14:102170.

107. Weinert LA, Araujo-Jnr EV, Ahmed MZ, Welch JJ. The incidence of bacterial endosymbionts in terrestrial

arthropods. Proceedings of the Royal Society B: Biological Sciences. 2015. https://doi.org/10.1098/rspb.2015.0249.

108. Duron O, Bouchon D, Boutin S, Bellamy L, Zhou L, Engelstädter J, et al. The diversity of reproductive parasites

among arthropods: Wolbachia do not walk alone. BMC Biol. 2008;6:1–12.

109. Binetruy F, Buysse M, Barosi R, Duron O. Novel Rickettsia genotypes in ticks in French Guiana, South America.

Sci Rep. 2020;10:1–11.

110. Bordenstein SR, Wernegreen JJ. Bacteriophage Flux in Endosymbionts (Wolbachia): Infection Frequency, Lateral

Transfer, and Recombination Rates. Mol Biol Evol. 2004;21:1981–91.

111. Duron O. Lateral transfers of insertion sequences between Wolbachia, Cardinium and Rickettsia bacterial

endosymbionts. Heredity . 2013;111:330–7.

112. Sonenshine DE, Michael Roe R. Biology of Ticks Volume 2. Oxford University Press; 2013.

113. Narasimhan S, Fikrig E. Tick microbiome: the force within. Trends Parasitol. 2015;31:315–23.

114. Institute of Medicine, Board on Population Health and Public Health Practice, Committee on Lyme Disease and

Other Tick-Borne Diseases: The State of the Science. Critical Needs and Gaps in Understanding Prevention,

Amelioration, and Resolution of Lyme and Other Tick-Borne Diseases: The Short-Term and Long-Term Outcomes:

Workshop Report. National Academies Press; 2011.

115. Bonnet SI, Binetruy F, Hernández-Jarguín AM, Duron O. The Tick Microbiome: Why Non-pathogenic

Microorganisms Matter in Tick Biology and Pathogen Transmission. Front Cell Infect Microbiol. 2017;7:236.

Keywords

Symbiosis; comparative genomics; evolutionary biology; phylogeny; ticks; bacterial communities

Figure captions

Figure 1: Phylogeny of Legionellales (Dataset 1) focusing on the Rickettsiella genus inferred with RAxML on 92 single copy panorthologous intact genes (100 bootstraps, evolutionary model: GTR+G4, rapid bootstrap random number seed: 1234, the random number seed for the parsimony inferences: 123), using B. aphidicola as outgroup to root the tree. Only bootstrap supports lower than 98 are shown. The Rickettsiella organisms whose genomes were sequenced in this study are indicated in bold. On the right, the presence (green), absence (grey) and pseudogenization (yellow) of genes of specific metabolic pathways annotated with KEGG is represented as a heatmap. B. aphidicola sequences were not annotated. On the bottom, Venn diagrams of clusters of orthologs of all encoded genes, intact genes, and pseudogenes is provided for the specified nodes of the tree (Node 10, Node 13, Node 15).

Figure 2: Intersection plot displaying shared and specific clusters of orthologous proteins between Rickettsiella and other Legionellales. Intersection plot is based on Dataset 1 and was produced with the R package UpSetR. In the left side, the horizontal bar plot represents the number of orthogroups present in each organism of the dataset. In the top, the vertical bar plot represents the number of orthogroups that belong to specific intersections of organisms in the dataset. The height of each bar indicates how many orthogroups are in that particular intersection, i.e., shared by the combination of organisms indicated by the intersection matrix below. In the intersection matrix, on the bottom, each row represents an organism and each column represents a possible intersection of these organisms. Filled circles indicate that the organisms in that row are part of the intersection shown in the corresponding column. The lines connecting the filled circles indicate which organisms are included in each specific intersection. This plot is limited to six orthogroups in the intersections, for the full plot of all intersections until one shared orthogroup please see Supplementary Figure 2.

Figure 3: Phylogeny and domain architecture composition of cifA and cifB homologs in Rickettsiella and other endosymbionts. The maximum-likelihood tree was reconstructed from an alignment concatenated with RNA-binding-like, AAA-ATPase-like, and PD-(D/E)XK nuclease domain sequences (191 amino acid positions, JTT+G substitution model). The tree is midpoint rooted. The type I-V cif representation, as well as the sizes and domain compositions of the cifA and cifB genes, were drawn relative to those described in Martinez et al. (2021, 2022). The scale bar represents the inferred number of substitutions per site. Bootstrap values were estimated from 1000 replicates. Accession numbers of the Rickettsiella, Wolbachia, and Rickettsia genomes used in this analysis are listed in Supplementary Table S4.

Table 1: Simplified scheme of presence (✓) and absence (X) of virulence factors categories in the organisms of Dataset 1. The organisms were grouped by genus, separating only Coxiella endosymbionts from the proved pathogenic relative C. burnetii. The virulence factor category was considered present if at least three of the representatives of the Rickettsiella strains and one of the Berkiella and Coxiella strains contained at least half of the virulence factors of that category. The search was performed using reference Legionella sequences from VFDB as queries in a local blast search (default settings, results were manually curated). Details on the annotation can be found in Supplementary table 6.

Supplementary Figure 1: barplot of intact genes and pseudogenes detected by PseudoFinder in each organism of Dataset 1. The height of each bar represents the number of intact genes (blue) and pseudogenes (magenta).

Supplementary Figure 2: 16S rRNA based phylogeny of Rickettsiella and other Legionellales. The dataset includes the sequences of Dataset 1 (indicated in bold), all the available Diplorickettsiales 16S rRNA sequences on SILVA RefNR SSU r138.1 (446 at 10th April 2024), and the 16S rRNA of the outgroup B. aphidicola. The phylogenetic inference was

performed using RAxML (100 bootstraps, rapid bootstrap random number seed: 1234, random number seed for the
parsimony inferences: 123, default setting for model of nucleotide substitution). The tree was pruned with iTOL
collapsing nodes distantly related to the target strains from Dataset 1 (bootstrap supports of each node, including the
collapsed ones, were maintained). The Rickettsiella strains sequenced in this study are indicated in bold red. The arrow
indicates the phylogenetic cluster supported by Node 15 in the phylogenomics-based tree (Figure 1) and the arrowhead
indicates the phylogenetic cluster supported by Node 10 in the phylogenomics-based tree (Figure 1).
799
Supplementary Figure 3: Intersection plot of Dataset 1 produced with the R package UpSetR, where all intersections are
presented. In the left side, the horizontal bar plot represents the number of orthogroups present in each organism of the
dataset. In the top, the vertical bar plot represents the number of orthogroups that belong to specific intersections of
organisms in the dataset. The height of each bar indicates how many orthogroups are in that particular intersection, i.e.,
shared by the combination of organisms indicated by the intersection matrix below. In the intersection matrix, on the
bottom, each row represents an organism and each column represents a possible intersection of these organisms. Filled
circles indicate that the organisms in that row are part of the intersection shown in the corresponding column. The lines
connecting the filled circles indicate which organisms are included in each specific intersection.
808
809 Supplementary Figure 4: Phylogenies of the thiamine genes present in Oerr and not in the other Rickettsiella strains of
810 Dataset 1, in order: thiD, thiDE, thiO, thiG, thiS. Although it is not possible to clearly establish the evolutionary origin
811 of these sequences, it is clear that they are of Gammaproteobacteria/Pseudomonadota origin, often found in Legionella
812 and in two other Rickettsiella organisms (See also Supplementary Table 3).
813
814 Supplementary Table 1: Metadata, genome statistics and number of intact and pseudo- genes of the organisms included
815 in Dataset 1. The ANI values for Dataset 1 and for the redundant genomes that were excluded from the analyses are also
816 presented, as well as the mobile elements for each genome in Dataset 1.
817
Supplementary Table 2: Orthogroups.GeneCount.tsv contains the orthologous clusters matrix produced with
OrthoFinder on the translated intact genes of Dataset 1 and on the translated gene sequences of the outgroup B.
aphidicola. This matrix was used to select the clusters for the phylogenomics-based tree and for the intersection plots. In
the following sheets, the intersections of clusters of specific organisms of the datasets are presented, together with their
KEGG and Prokka annotations for one of the organisms in the intersection (sequence ID reported). SharedByAll:
clusters of intact genes shared by all organisms in the dataset, i.e. Dataset 1 and B. aphidicola; AllNoBuchnera: shared
by all Legionellales of Dataset 1 and not present in B. aphidicola; SharedByAllRickettsiella: shared by all Rickettsiella
strains of Dataset 1; SharedAllIricinus: shared by all Rickettsiella strains isolated from I. ricinus included in Dataset 1;
SharedAllIricinus_NoRmassiliensis: as SharedAllIricinus, but not present in the putative pathogen R. massiliensis 20B;
Shared_MostPSstrains_IricRmassiliensis: shared exclusively by the Rickettsiella strains with the highest number of
pseudogenes, i.e. R. massiliensis 20B and Rickettsiella strain Iric; SpSpecific_RDG: clusters exclusively present in the
genome of the Rickettsiella endosymbiont of D. gallinae; SpSpecific_Rgrylli: clusters exclusively present in the
genome of R. grylli; SpSpecific_Ird1: exclusively present in the genome of Rickettsiella strain Ird1; SpSpecific_Oerr:
exclusively present in the genome of Rickettsiella strain Oerr; SpSpecific_Opha: exclusively present in the genome of
Rickettsiella strain Opha; SpSpecific_Rmassiliensis: exclusively present in the genome of R. massiliensis 20B;
SpSpecific_Rviridis: exclusively present in the genome of R. viridis.
834
Supplementary Table 3: KEGG annotation of the organisms of Dataset 1 obtained through BlastKoala (default settings).
The column Knumbers indicates the KEGG ID. The values indicate presence (2), pseudogenization (1), and absence (0)
in each organism. The sheets Summary NutrMutPaths, NutrMutPaths and EnergyPaths are focused on specific
metabolic pathways potentially involved in nutritional mutualisms (B vitamins and heme metabolism, energy
production); in these tables, "B"s mean null values in correspondence of the pathway names and were inserted as
information for the script to obtain the heatmap shown in Figure 1. Thiamine_Oerr contains the amino acid sequences
of the thiamine pathway components identified in Rickettsiella strain Oerr and not in the other Rickettsiella strains
included in Dataset 1, together with the five best-hits of blast against the non-redundant NCBI database, ordered by
identity percentage (ID of the hit sequences, identity percentage, query cover, E value and available taxonomic
annotation of the hit organism reported). The organisms and their sequence accession number used to infer the

phylogenies of the Oerr thiamine genes (Supplementary Figure 4) are reported in the sheet Thiamine Phylogenies
datasets.
847
848 Supplementary Table 4: Accession number of the Rickettsiella, Wolbachia, and Rickettsia genomes from public
849 databases used to screen the cifA and cifB genes pair homologs.
850
851 Supplementary Table 5: Intersections of clusters of all encoded sequences, all intact sequences, and all pseudogenes of
852 specific phylogenetic clades represented in Figure 1. A reference sequence ID is provided for each cluster, together with
853 the KEGG (obtained with BlastKoala, default settings) and Prokka annotation.
854
855 Supplementary Table 6: Annotation of putative virulence factors in the organisms of Dataset 1, annotated using amino
856 acid Legionella queries downloaded from VFDB as a reference in a local blast search (default parameters) against
857 translated all genes of the organisms of Dataset 1. Sequence IDs are reported for each organism, cells are highlighted in
858 yellow when the sequence was identified as a pseudogene by PseudoFinder. The hyphen ("-") indicates complete
859 absence of the sequence.
860
861

Buchnera aphidicola
Uncultured Legionellales sp. HQ225336.1.1510
Uncultured Legionellales sp. FM242264.1.1505
Legionella pneumophila NZ_CP015941.1

Berkiella cookevillensis EF492067
Berkiella aquae AY741401
Uncultured Legionellales bacterium KR110095.1.1504
Coxiella endosymbiont of *Amblyomma sculptum* strain CeasUFV CP033868.1
Coxiella burnetii RSA 493 NC002971.1
Coxiella endosymbiont of *Ornithodoros maritimus* CLEOmar GCF907164965.1

Uncultured Legionellales bacterium KJ752786.1.1458

Uncultured Legionellales bacterium MF085232.1.1543

Uncultured Legionellales bacterium KC541146.1.1542
Uncultured Legionellales bacterium AB630758.1.1506
Rickettsiella endosymbiont of *Dermanyssus gallinae* GCF019285595.1
Rickettsiella viridis GCF003966755.1

Uncultured *Rickettsiella* sp. KC764412.1.1530
Rickettsiella massiliensis GQ857549.1.1488
Rickettsiella massiliensis strain 20B GCF000257395.1
Rickettsiella isolated from *Ixodes ricinus* strain Ird6
Rickettsiella isolated from *Ixodes ricinus* strain Iric
Uncultured *Rickettsiella* bacterium EF451626.1.1272
Rickettsiella sp. clone RL1 MK627517.1

Uncultured *Rickettsiella* bacterium GU906684.1.1529
Rickettsiella bacterium from *Cecidotrioza sozanica* AF286124.1.1470
Uncultured *Rickettsiella* sp. KT767597.1.1207
Rickettsiella isolated from *Ornithodoros phacochoerus* strain Opha
Uncultured *Rickettsiella* bacterium HM329706.1.1329
Rickettsiella grylli GCF000168295.1
Rickettsiella grylli AAQJ02000001.271236.272713
Rickettsiella grylli AAQJ02000001.641042.642519

Rickettsiella grylli U97547.1.1478
Rickettsiella sp. GSU AF383621.1.1538
Rickettsiella isolated from *Ornithodoros erraticus* strain Oerr

Rickettsiella isopodorum GCF001881495.1
Uncultured Legionellales bacterium EU800584.1.1495
Rickettsiella isolated from *Ixodes ricinus* strain Ird1
Rickettsiella isolated from *Proasellus assaforensis* HAEP01038506.44.1584

Uncultured *Rickettsiella* sp. EU800447.1.1527
Rickettsiella symbiont of *Asellus aquaticus* AY447040.1.1494
Uncultured *Rickettsiella* bacterium JN983644.1.1524