RNA-programmable cell type monitoring and manipulation in the human cortex with CellREADR

Elizabeth A. Matthews,[1,2] Jeffery B. Russ,[1,3] Yongjun Qian,[2,5] Shengli Zhao,[2] Peyton Thompson,[1,2] Muhib Methani,[1,2] Matthew L. Vestal,[1,6] Z. Josh Huang,[2,4]* and Derek G. Southwell[1,2,4],**

[1]Department of Neurosurgery, Duke University School of Medicine, Durham, NC USA
[2]Department of Neurobiology, Duke University School of Medicine, Durham, NC USA
[3]Department of Pediatrics, Division of Neurology, Duke University School of Medicine, Durham, NC, USA
[4]Department of Biomedical Engineering, Duke University, Durham, NC USA
[5]Present address: College of Future Technology, Peking-Tsinghua Cener for Life Sciences, IDG/McGovern Institute for Brain Research,
Beijing Advanced Center of RNA, Peking University, Peking China
[6]Present address: Department of Neurosurgery, Dartmouth Hitchcock Medical Center, Lebanon, NH USA
*Correspondence: josh.huang@duke.edu
**Correspondence: derek.southwell@duke.edu
10

SUMMARY

Reliable and systematic access to diverse cell types is necessary for understanding the organization,
function, and pathophysiology of human neural circuits. Methods for targeting human neural populations
are scarce and currently center on identifying transcriptional enhancers and engineering viral capsids.
Here we demonstrate the utility of CellREADR, a programmable RNA sensor-effector technology that
couples cellular RNA sensing to effector protein translation, for accessing, monitoring, and manipulating
specific neuron types in human cortex, *ex vivo*. We designed CellREADRs to target two subpopulations,
CALB2 GABAergic interneurons and *FOXP2* glutamatergic projection neurons, then validated targeting
specificity using histological, electrophysiological, and transcriptomic methods. CellREADR expression of
channelrhodopsin and GCamp enabled the manipulation and monitoring of these populations in live
cortical microcircuits. By demonstrating specific, reliable, and programmable experimental access to
human neuronal subpopulations, our results highlight CellREADR's potential for studying neural circuits
and treating brain disorders.

KEYWORDS

25 Human neuroscience, organotypic slice culture, cortex, cellular access, RNA sensor, interneuron,
26 excitatory neuron, PatchSeq, calcium imaging, optogenetics

INTRODUCTION

The human brain has evolved divergent and species-specific properties that underlie our unique cognitive
and behavioral capabilities. While approaches such as MRI, EEG, and post-mortem neuroanatomical
studies have identified many brain areas, functional networks, and fiber tracts involved in human
sensorimotor, cognitive, and emotional functions and their related disorders, brain function fundamentally
emerges from neural circuit computations performed by diverse, synaptically interacting cell populations.
To better understand the neural underpinnings of human behaviors and neurological conditions, it is thus
essential to investigate the fundamental elements of the brain - the diverse human neural cell types - and
to dissect their functional features and dynamic interactions at the spatiotemporal resolution of neural
circuit operation[1,2]. This requires 1) identifying and categorizing the diverse cell populations that comprise
the brain, 2) understanding the input-output connectivities of these cell types, and 3) characterizing their
physiological functions in circuit activity. Tools that allow for specific and systematic access to diverse
human cell types are essential to these efforts.

In recent years, single-cell RNA sequencing (scRNA-Seq) studies have yielded transcriptomic cell type
taxonomies of the rodent, non-human primate (NHP), and human brains[3–12]. In particular, scRNA-seq
data have demonstrated conserved as well as divergent features of human cell types, brain areal
specializations in cell composition[4,5], and evidence for genetic adaptations that may confer human-

specific neural circuit architectures[3,7,9,13]. Importantly, these transcriptional data also serve as a catalog of
genetic access points (i.e., "marker genes") that may enable the systematic targeting of human cell types
for integrated multimodal analyses. What has been lacking, however, are reliable, facile and scalable
tools that leverage gene expression to provide experimental access to marker-defined cell populations for
human neuroscience[2,14].

Traditionally, cell type experimental access has relied on germline engineering approaches in model
organisms (e.g., transgenic mouse driver lines)[15–18], which are inapplicable to human research[15–17]. More
recently, enhancer-based viral vectors have been advanced as an alternative, cross-species approach to
targeting neural cell types[19–22]. However, because enhancers work through complex DNA-protein
interactions across complex chromatin landscapes, they often require large-scale efforts to identify and
validate[21,23,24]. Furthermore, as enhancer tools are typically validated in mouse models, their
generalizability and applicability to human cell types remain to be established. Adeno-associated viral
capsid engineering is another potential strategy for cell type access[25], but capsid tropism at the resolution
of specific cell types has yet to be demonstrated for neurons, particularly in human brain tissues[26–30].

Neurosurgical tissue specimens represent a promising experimental platform for tool development and
human neuroscience research. Resected cortical tissues can be maintained *ex vivo* for weeks in
organotypic culture preparations, which maintain local tissue cytoarchitecture, cell transcriptomic profiles,
morphological features, and neurophysiological activity[31–33]. *Ex vivo* human tissues have been used for a
variety of live, semi-chronic research applications, such as the characterization of human disease
pathogenesis[34], testing of emerging therapeutic approaches[35,36], and validation of novel experimental
tools[21,22,37]. They have also been increasingly utilized for morphological and electrophysiological studies
of human neurons[38–42] with, in some cases, subsequent analyses of cell transcriptomic profiles[6,33,43–45]. By
enabling researchers to overlay cellular anatomical and physiological data onto transcriptomic datasets,
this set of approaches has enabled an initial multi-modal profiling of some human brain cell types.
However, previous studies have restricted neuronal sampling based on visible cellular features, such as
soma shape and laminar position[6,46,47], or through viral targeting with broadly active enhancers[19–21].
Altogether a lack of specific and effective methods for accessing molecularly defined cell types has
limited the basic and pre-clinical potential of human *ex vivo* tissue preparations.

We recently developed a technology orthogonal to enhancers, CellREADR, which uses a RNA and
translation-based method to achieve cell type-specific targeting across diverse organisms[48]. CellREADR -
Cell access through RNA sensing by Endogenous ADAR (adenosine deaminase acting on RNA) - is a
programmable RNA sensor-effector technology that couples the detection of a cell-defining RNA to the
translation of a user-selected effector protein; this coupling is achieved through an ADAR-mediated A-to-I
editing mechanism ubiquitous to metazoan cells. Two independent teams have described essentially
similar methods, and termed them RADARS[49] and RADAR[50]. In a prior study we characterized
CellREADR sensor and effector properties in human cell lines, demonstrated CellREADR-mediated
monitoring and manipulation of neurons in behaving mice, and provided initial proof-of-principle for its
application in *ex vivo* human brain tissues[48]. These initial results highlighted CellREADR's potential as a
specific, scalable, and versatile genetic tool for studying human brain cell types and neural circuits.

In this study we demonstrate CellREADR's utility for accessing, monitoring, and manipulating marker-
defined cell types in *ex vivo* human brain. We designed CellREADR constructs to provide experimental
access to *CALB2* (calretinin) GABAergic interneurons and *FOXP2* (forkhead box protein P2)
glutamatergic projection neurons in the temporal neocortex. Following histological, electrophysiological
and transcriptomic characterizations of CellREADR-labeled neurons, we used PatchSeq techniques to
cross-validate targeting specificity and perform molecular alignment of the experimentally accessed cells
onto published transcriptomic reference atlases. We further performed optical stimulation and calcium
imaging in *CALB2* and *FOXP2* populations to probe human cortical microcircuits. Together, our
experiments highlight CellREADR as a powerful approach for cell type-specific access, monitoring, and
manipulation in human brain circuits, *ex vivo*.

RESULTS

Deploying CellREADR to target RNA-defined cell types in ex vivo human cortex
We used an organotypic slice preparation to study CellREADR applications in human cortical tissues
(Figure 1A). Live tissue specimens were donated by epilepsy surgical patients and collected from non-

diseased areas adjacent to known seizure foci (proximal, non-diseased tissues are minimally removed
during surgery to access epileptogenic areas). Donor demographics and specimen details are presented
in Table 1.

We designed both singular and binary CellREADR vectors for use in human tissues (Figure 1B). The
singular vector system drives expression of the sensor and, upon sense-editing by endogenous ADAR,
expression of the effector RNA; this vector was used to directly couple RNA sensing to the translation of
an mNeonGreen (mNeon) effector. The binary system was designed to utilize two vectors, one, a READR
vector that expressed the transcription activator, tTA2, upon sense-editing, and the other, a reporter
vector containing a TRE-driven effector. The binary system enables amplification of effector payload
expression and also allows pairing of READR and reporter vectors for differential expression of effectors
in separate target populations. The singular vector was used in some histological studies (Figure 1) and a
subset of the patch clamp studies (Figures 2 and 3; *CALB2* only), while binary vectors were used for all
other experiments, including histological studies (Figure 1; effector, mNeon), patch clamp
characterizations of cellular properties (Figures 2 and 3; mNeon), optogenetic interrogations (Figure 4;
ChIEF), and calcium imaging (Figure 5; GCaMP7f).

We first produced CellREADR sensors for various interneuron and projection neuron marker genes
(*VGAT* (n=2 sensors), *LAMP5* (n=1), *CALB2* (n=3)*, SST* (n=2), *PVALB* (n=3)*, UNC5B* (n=1)*, CUX1* (n=1)*,
FOXP2* (n=2)*;* a total of 15 sensors) and used them in binary vector designs to express mNeon in cells of
the human cortex. In screening these designs on human tissues, we observed READR activity with 13
sensors. Labeled cells exhibited diverse morphological features that were consistent with glutamatergic
and GABAergic neurons, depending on the sensor (example labeling, Figure 1C). We elected to pursue
more detailed investigations using two screened sensors, one for *FOXP2* and the other for *CALB2*, given
their robust expression in human slices and prior data suggesting these RNAs may be expressed in cell
populations that confer or exhibit human specializations.

FOXP2 is implicated in human language disorders[51,52], aberrant song learning in songbirds[53], and
impairments in fine motor learning in mice[54]. In rodents, *FOXP2* expression is restricted to cortical
excitatory neurons in deep layers[55], however, *FOXP2* is expressed more widely across cortical lamina in
humans[48,56]. Consistent with this, after applying a binary *FOXP2* CellREADR system to neocortical slices,
we observed mNeon epifluorescence in pyramidal cells of both deep and superficial cortical layers
(Figures 1C-E). To assess the specificity of *FOXP2* CellREADR targeting, we performed
immunohistochemistry to quantify co-labeling of READR mNeon with cellular FOXP2 7 days after virus
application (Figure 1F). Across tissues from 5 subjects, the mean specificity of the *FOXP2* READR
labeling (percentage of mNeon+ cells co-labeled with FOXP2) was 75.4 ± 2.2% (Figure 1F-G).

In the human cortex, *CALB2* expression is largely restricted to supragranular GABAergic interneurons
that originate from the caudal ganglionic eminence (CGE), primarily *VIP* subclass interneurons[57–59]. There
is evidence for an increased abundance of CGE interneurons (versus medial ganglionic eminence
interneurons) in humans and non-human primates, as compared to mice[3,7,9], with some studies
suggesting a specific expansion of *CALB2* populations[57,60,61]. Using a binary *CALB2* READR system in
human neocortical slices, we observed a pattern of mNeon labeling concordant with prior descriptions of
CALB2 expression. Compared to *FOXP2* READR labeling, *CALB2* READR expression was generally
observed in superficial cortical layers (Figures 1C-E). *CALB2* READR-labeled cells also exhibited
morphologies distinct from *FOXP2* READR-labeled cells, with small somata and vertically oriented,
bipolar or multipolar dendritic processes (Figure 1C). The specificity of the binary *CALB2* READR
transduction, as tested by immunohistochemistry in tissues from 7 subjects, was 72.2 ± 5.5%; specificity
of the singular *CALB2* READR was 73.1 ± 3.0% (3 subjects; Figures 1F-G). Quantification of
immunostaining signals in FOXP2 and CALB2 cells indicated that CellREADR sense-editing did not
disturb CALB2 expression in targeted cells, whereas FOXP2 expression was found to be slightly lower in
cells targeted by the *FOXP2* READR (Supplemental Figure 1).

To benchmark targeting by the *CALB2* CellREADRs, we compared READR labeling to that achieved by
an enhancer virus that has been recently used for human pan-interneuron targeting, DLX2.0[20] (Figure
1H). Consistent with prior evidence that DLX2.0 transduces broadly across GABAergic cell subclasses[33],
43.1 ± 6.0% of DLX2.0-YFP+ cells co-expressed CALB2 (Figure 1J),. Compared to *CALB2* CellREADR

labeling, which was noted primarily in supragranular layers, DLX2.0 virus expression was also more
evenly distributed across the depth of the cortical gray matter (Figure 1I). Although the binary and
singular *CALB2* CellREADR vectors theoretically targeted a more restricted population of interneurons,
we found that they labeled greater numbers of cells than the DLX2.0 virus, suggesting their greater
targeting efficiency (Figure 1K; Binary *CALB2*-READR, 34.5 ± 8.0 cells/mm^2, Singular *CALB2*-READR,
20.1 ± 3.5 cells/mm^2, DLX2.0-YFP, 14.8 ± 2.8 cells/mm^2; 1-way ANOVA, p = 0.04).

PatchSeq validation of CALB2 CellREADR targeting
To further validate CellREADR targeting and gain a multimodal characterization of some of the cells
accessed by the *CALB2* READR, we performed READR-targeted PatchSeq on a limited set of cells. Four
to 12 days after the application of singular or binary *CALB2* READR vectors, we targeted mNeon+ cells
for patch clamp electrophysiology, biocytin filling, and nuclear extraction. After performing RNA-seq on
the nuclear aspirate, we applied an open sequencing analysis pipeline[62] to map queried cells onto a
reference transcriptomics-based taxonomy of human middle temporal gyrus (MTG) neurons[59] (Figure 2A,
Supplemental Figure 2A).

As expected, our RNA-seq analysis detected *CALB2* transcripts in nearly all mNeon+ patch-targeted cells
(18/19, 94.7%; Figure 2B). Notably, this exceeded the specificity of *CALB2* READR targeting measured
by immunohistochemistry (Figure 1G), suggesting that incomplete sensitivity of CALB2 immunostaining
may have underestimated the specificity of the *CALB2* READR. These RNA-seq results also indicate the
CellREADR sense-edit mechanism did not grossly disrupt expression of the *CALB2* target gene. Based
on existing transcriptomics data[59] (Figure 2A), which demonstrate that *CALB2* is expressed almost
exclusively by *VIP* subclass interneurons, we hypothesized that *CALB2* READR-targeted cells would also
express the interneuron class marker, *GAD1*, as well as *VIP*, but not markers of other cortical interneuron
subclasses, such as *LAMP5*, *SST*, and *PVALB*. In accordance, transcripts of *GAD1* were detected in the
majority of sampled cells (14/19, 73.7%), while *VIP* was detected in a somewhat smaller fraction (9/19;
47.4%); by contrast, transcripts of *LAMP5, SST, PVALB* were rarely observed in *CALB2* READR-targeted
cells (Figure 2B). Looking beyond single gene expression, essentially all cells (18/19; 94.7%) mapped
transcriptomically to types within the *VIP* subclass of MTG interneurons (Figures 2A and 2B). Of cells that
mapped to the *VIP* subclass, one-third (6/18; 33.3%) were categorized as *PAX6 SYT6* type, while the
remaining two-thirds (12/18; 66.7%) mapped to other *VIP* types. In total, 10/21 previously annotated
transcriptomic types of *VIP* interneurons[63] were represented amongst the 19 cells sampled by *CALB2*
READR targeting. These results together indicate that CellREADR specifically targeted *CALB2*
populations and, thereby, effectively accessed *VIP* interneurons (Supplemental Figure 2B). Expected
transcriptomic alignment of *CALB2* READR-targeted cells onto this existing database (which was
produced without CellREADR) suggest CellREADR did not significantly alter target cell transcriptomes,
consistent with immunohistochemistry data provided in Supplemental Figure 1 and results we have
presented in a previous study[48].

Electrophysiology data from 17/19 cells studied with RNAseq passed quality checks, and the
morphologies of 7 cells were recovered for tracing. *CALB2* READR-targeted neurons largely showed a
steep gain relationship between injected current and firing frequency, with many cells exhibiting firing
frequencies exceeding 40Hz (Figure 2C). Qualitatively, cells that clustered within the same transcriptomic
type nevertheless exhibited differences in their electrophysiological features. For example, within
READR-targeted cells identified as *PAX6 SYT6* type, we observed two distinct patterns of input-output
responses (Figure 2C), as well as similarly variable responses to hyperpolarization (Figure 2D). These
basic phenotypic observations were consistent with recent human interneuron PatchSeq data[33], and
generally align with prior observations that cell transcriptomic types do not necessarily express type-
specific physiological features. As expected, however, we did observe some relationships between ion
channel subunit expression and membrane physiology[38]. The expression of BK channel subunits, for
example, correlated with both the depth of the fast afterhyperpolarization and the action potential
duration, as would be expected given the function of this channel[64,65] (Supplemental Figure 3).
Morphologically, reconstructed neurons were located in Layers 2-5 and exhibited diverse dendritic and
axonal distributions. Six of 7 cells projected dendrites vertically, while the other recovered cell's dendrites
were oriented horizontally (Figure 2E). These morphologies are likewise generally consistent with *VIP*
subclass morphologies[33].

CALB2- and FOXP2-targeted neurons exhibit distinct physiological profiles and spontaneous activities

After gaining evidence for CellREADR specificity through immunostaining and sequencing analyses, we proceeded to a more detailed physiological characterization of neurons targeted by *CALB2* and *FOXP2* READRs. Applying a panel of electrophysiological readouts used to construct prior human cellular atlases[33,43,66], we recorded electrophysiological properties from 60 *CALB2* READR-targeted cells (n=8 subjects) and 43 *FOXP2*-targeted cells (n=7 subjects). We first measured intrinsic membrane properties and noted group differences in membrane capacitances (*CALB2*: 31.72 ± 3.35 pF; *FOXP2*: 88.46 ± 9.05 pF, Mann-Whitney p<0.0001) and input resistances (*CALB2*: 0.280 ± 0.041 MΩ; *FOXP2*: 0.082 ± 0.018 MΩ, Mann-Whitney p<0.0001; Figure 3A). Hyperpolarizing sag ratios (*CALB2*: 0.18 ± 0.02; *FOXP2*: 0.15 ± 0.02) and resting membrane potentials (*CALB2*: -55.07 ± 1.04 mV; *FOXP2*: -57.30 ± 1.77 mV) did not differ between targeted cells (Figure 3A). We also characterized active membrane properties by quantifying cellular input-output responses (rheobase, gain, and maximal firing frequency; Figure 3B). *CALB2*-targeted cells showed more variability in the shapes of their input-output functions, with many neurons exhibiting low rheobase, high gain, and high maximal firing frequency (41/59 (65%) with maximum firing frequency >40Hz), and a smaller subset demonstrating high rheobase, low gain, and low maximal firing frequency. *FOXP2*-targeted cells were of more uniform firing behaviors, with relatively shallow gain slopes and maximal firing frequencies typically under 40Hz (11/46 (24%) with maximum firing frequency >40Hz). Statistically, rheobases (*CALB2*: 37.9 ± 5.9 pA; *FOXP2*: 110.8 ± 14.0 pA, paired Mann-Whitney p<0.0001), gains (*CALB2*: 0.55 ± 0.04 Hz/pA; *FOXP2*: 0.18 ± 0.03 Hz/pA, paired t-test, p<0.0001), and maximal firing frequencies (*CALB2*: 53.94 ± 2.57Hz, *FOXP2*: 36.15 ± 4.03Hz, Welch's t-test p=0.001) differed between *CALB2*- and *FOXP2*-targeted cells (Figure 3B).

We performed brief current injections (2 ms) to measure evoked single action potential parameters, including threshold, peak, half-width, and fast afterhyperpolarization from targeted cells (Figure 3C). All single action potentials from the *CALB2* and *FOXP2* cells were aligned on the rising slope and averaged. *CALB2*-targeted cells had narrower action potentials (half-width, *CALB2*: 0.83 ± 0.03 ms; *FOXP2*: 1.30 ± 0.13 ms, Welch's t-test p=0.002; duration (not shown), *CALB2*: 1.99 ± 0.08 ms; *FOXP2*: 2.78 ± 0.44 ms, Mann-Whitney p=0.003). The *CALB2* population had a larger fast afterhyperpolarization (fAHP; *CALB2*: -21.9 ± 0.75 mV; *FOXP2*: -18.29 ± 1.13 mV, Welch's t-test p=0.009); a subset of the *FOXP2*-targeted neurons were notable for their pronounced afterdepolarizations. Neither the action potential thresholds (*CALB2*: -42.77 ± 0.98 mV; *FOXP2*: -45.36 ± 1.53 mV) nor the action potential heights (*CALB2*: 43.95 ± 1.60 mV; *FOXP2*: 41.98 ± 1.92 mV) statistically differed between the targeted populations. In an alternate stimulation paradigm (500 ms current injections), we also observed these differences in action potential waveforms between *CALB2*- and *FOXP2*-targeted neurons (Supplemental Figures 4A and 4B). The observed differences in intrinsic properties, firing phenotypes, and evoked action potential dynamics also indicate the *CALB2* and *FOXP2* READRS targeted distinct populations of interneurons and projection neurons, respectively[67,68]. Additionally, and consistent with these findings and the results in Figure 1, reconstructions of the patched cells also demonstrated corresponding morphological differences between the targeted populations (Supplemental Figure 5).

Consistent with previous results[33], we did not find evidence that intrinsic properties measured at later time points (8-12 days after slice preparation) differed from data collected at earlier points (Supplemental Table 1). Likewise, the *CALB2* READR-targeted cells exhibited physiological properties that resembled those of *VIP* subclass human interneurons collected with an alternate targeting approach[33] (either without the use of viruses, or with a DLX2.0 virus; Supplemental Table 2). These data indicate that over the period of slice culture CellREADR also did not grossly disturb cellular physiology.

We also used patch clamp to measure activity from *CALB2*- and *FOXP2*-targeted cells at rest, in order to assess synaptic inputs and cellular activity in the slice network. In both target populations, we observed excitatory and inhibitory postsynaptic potentials (EPSPs and IPSPs; Figure 3D), as well as action potential firing (Figure 3E). *CALB2*-targeted neurons often fired from resting membrane potential, whereas a smaller fraction of *FOXP2*-targeted cells exhibited this behavior (*CALB2*: 45/66 cells (69.7%), *FOXP2*: 14/50 cells (28.0%); Fisher's exact test p<0.001; Figure 3F). Spontaneous activity, measured as the frequency of spontaneous action potential firing, differed between *CALB2-* and *FOXP2*-targeted populations (*CALB2*: 2.65 ± 0.53 Hz; *FOXP2*: 1.60 ± 0.69 Hz, Mann-Whitney p<0.0001, Figure 3G). The properties of spontaneous action potentials did not differ between groups (Supplemental Figures 4C and

4D). Overall, these observed differences in spontaneous activity were consistent with our intrinsic
physiological measures (Figure 3), which indicated the *CALB2*-targeted population is more excitable, with
lower rheobase and higher input resistance.

In a subset of spontaneous action potential recordings, we monitored firing before and after application of
glutamatergic blockers (CNQX/AP-5; Figure 3H). Glutamatergic blockade resulted in increased (1/6 cells),
abolished (3/6), or partially reduced firing (2/6; Figure 3I). As some cells exhibited spontaneous firing
while under excitatory blockade, the baseline activity measured at rest reflected a mixture of synaptic
recruitment by excitatory inputs as well as spontaneous action potential generation.

Cell type specific optical manipulation of human neurons with CellREADR
We next explored CellREADR functionality for cellular manipulation and interrogation of human
microcircuits. We designed a binary vector system consisting of either *CALB2*-tTA2 or *FOXP2*-tTA2
paired with a TRE-driven channelrhodopsin variant, ChIEF[69], and then delivered viruses to neocortical
slices. We first used patch clamp to assess optically evoked action potential firing in *CALB2*-ChIEF and
FOXP2-ChIEF cells. In all *CALB2*-targeted cells (n=6), we observed reliable, light-entrained action
potentials (Figure 4A); each of these cells was able to follow light pulses with action potentials at
frequencies 25-50Hz (data not shown). By contrast, optical stimulation of *FOXP2*-ChIEF cells evoked
membrane depolarizations of 2-9mV, but it did not elicit action potentials (n=10 cells; Figure 4A). These
differences in light responsiveness were somewhat expected based on our intrinsic physiological
characterizations of these populations (Figure 3), which indicated that *FOXP2*-targeted cells are less
excitable than *CALB2*-targeted cells. We thus focused on additional optogenetic experiments with *CALB2*
READRs.

It has been reported that calretinin interneurons project onto local inhibitory and excitatory populations in
rodent cortex[70] and hippocampus[71]. In non-human primate visual cortex CALB2-immunostained cells
have been found to target glutamatergic neurons in deeper layers[72]. Here, to functionally interrogate the
synaptic targets of *CALB2* neurons in human neocortex, we recorded from pyramidal neurons and
interneurons while optically stimulating *CALB2*-ChIEF cells. Putative postsynaptic cells were labeled with
complementary DLX2.0-YFP (Figure 4B) or hSyn-mCherry AAV vectors to label YFP+ interneurons and
mCherry+ pyramidal neurons, respectively, for patch clamp recordings (Figure 4B). At short latencies
following *CALB2*-ChIEF stimulation, we observed optical inhibitory PSCs (oIPSCs) in slice interneurons
and pyramidal neurons, which were absent at the reversal potential for GABA and blocked by bicuculline
(Figure 4C). We also detected optically evoked EPSCs in pyramidal neurons but not interneurons; these
events were absent at the reversal potential for glutamate, and they were not blocked by bicuculline.
These oEPSCs were only seen in hSyn-mCherry labeled pyramidal neurons (Figure 4C). Representative
recordings of 10 trials from a single postsynaptic DLX2.0-YFP cell are shown in Figure 4D. The first
stimulus in a 25 Hz train had a failure rate of 10%, while the fourth stimulus had a failure rate of 40%;
these results are aligned with the failure rates for *CALB2* cell activation shown in Figure 4A. The average
amplitude oIPSCs in both post-synaptic populations was 3.30 ± 0.4pA (Figure 4E); the latencies, rise
times, and durations of oIPSCs (normalized to event peak amplitude) indicated that light stimulation
produced monosynaptic GABA release onto the patched cells (Figure 4D).

We observed optical PSCs in both YFP+ interneurons and mCherry+ pyramidal neurons (6/16 DLX2.0-
YFP+ cells (37.5%) and 3/14 hSyn-mCherry+ cells (21.4%)). When performing voltage clamp measures,
we found that *CALB2*-ChIEF activation evoked inhibitory, but not excitatory, responses in interneurons
(6/16 DLX2.0-YFP+ cells (33.3%) exhibited inhibitory responses and 0/16 cells exhibited excitatory
responses upon *CALB2*-ChIEF activation; Figure 4F). By contrast, *CALB2*-ChIEF activation evoked
inhibitory responses in some pyramidal neurons, and excitatory responses in others (3/14 hSyn-
mCherry+ pyramidal cells (21.4%) exhibited inhibitory responses and 2/14 cells (14.3%) exhibited
excitatory responses *CALB2*-ChIEF activation; Figure 4F). It is unclear whether this observation of
excitatory responses in pyramidal neurons reflected glutamate release from a subset of *CALB*+ cells or
off-target expression of the CellREADR ChIEF. Notably, our PatchSeq experiments indicated that at least
a subset of *CALB2*-targeted cells, particularly those that align with the PAX6 SYT6 transcriptomic
subtype, appear to express both vesicular glutamate transporter (*SLC17A7*) and glutamate
decarboxylase transcripts (*GAD1* and *GAD2*); Figure 2B)); additionally, a small subset of human MTG
excitatory neurons express low levels of *CALB2* (Supplemental Figure 2A)[59]. These results may indicate

that neurotransmitter signaling from human *CALB2* neurons may differ from that produced by rodent
Calb2 neurons. We did not observe both excitatory and inhibitory responses to *CALB2*-ChIEF activation
in the same pyramidal cell.

Cell type-specific monitoring of population activity with CellREADR

Last, to investigate CellREADR functionality for imaging cell type population activity, we used binary
vectors to express the genetically encoded calcium indicator, GCamp7f, in *CALB2* and *FOXP2* target
populations. To establish how calcium signals reported membrane depolarizations and action potentials,
we first measured fluorescence signals from individual GCaMP7f+ cells while performing current
injections with patch clamp (Figure 5A). In both target populations we observed a linear relationship (R^2 of
the linear fit *CALB2*: 0.84; *FOXP2*: 0.93) between the GCaMP $\Delta F/F$ measure and the frequency of
evoked action potentials. During current clamp recordings, we also observed both prolonged
spontaneous subthreshold membrane depolarizations and spontaneous action potentials in the same cell;
these events were reflected by distinct fluorescence signals with different rise times (Tau_{3AP} = 390 ms,
$Tau_{Depolarization}$ = 1760 ms, Tau_{1AP} = 202 ms, Figure 5B).

As we had observed frequent spontaneous action potential firing in *CALB2*- and *FOXP2*-targeted neurons
under patch clamp (Figure 3), we expected GCaMP imaging would detect population activity in the slice
preparation. Indeed, when performing wide field imaging of neocortical slices without stimulating agents,
we observed baseline spontaneous activity in both populations (representative images of GCaMP
labeling, Figures 5C). Representative time series of $\Delta F/F$ signals for *CALB2*- and *FOXP2*-targeted
populations are shown in Figure 5D. Qualitatively, spontaneous population activity patterns varied
between slices, even across slices that were prepared from the same subject's tissue. Action potentials
(reflected by fast-rising calcium transients of varying amplitudes (as in Figures 5A-B)) were observed in
multiple cells from the READR-targeted populations (Figure 5D). We also observed prolonged calcium
signals (as in Figure 5B) that occurred simultaneously in many cells (Figure 5D). These latter events may
reflect the influence of local neuromodulatory signals or enhanced within-population connectivity across
the imaged population. Coordinated activity was more prominent in *FOXP2*-targeted populations than in
CALB2-targeted populations, as reflected by a cross-wise correlation analysis of calcium signal patterns
(Pearson's rho value for population correlation, 0.639 for the *FOXP2-targeted* cells, 0.402 for the *CALB*-
targeted cells; translated into Fisher's Z values for statistical comparison, p=0.03; Figure 5E).

Differential gene expression analyses of an existing RNAseq database[59] indicate the expression of
nicotinic acetylcholine receptor subunits differs significantly between *CALB2* and *FOXP2* neurons. In
particular, cholinergic receptor nicotinic alpha 2 subunit (*CHRNA2)* is expressed in a higher percentage of
CALB2+ neurons (52.5%) than *FOXP2*+ neurons (1.1%), and at an approximately 1000-fold higher
expression level (p-adj = 0), though *CHRNA4*, *CHRNA6*, and *CHRNA7* also showed enrichment in
CALB2+ over *FOXP2*+ neurons (p-adj = 8.75×10^{-7}, 3.22×10^{-94}, 1.27×10^{-191}, respectively). Of nicotinic
acetylcholine receptor subunits, only *CHRNA1* showed enrichment in *FOXP2* neurons (2.8% of FOXP2+
neurons, with approximately 100-fold enrichment over *CALB2* neurons, p-adj = 6×10^{-3}). We thus
hypothesized that calcium signal responses to nicotinic acetylcholine receptor signaling would differ
between these populations. We compared transients from cells in the absence and presence of bath
nicotine, and observed a trend towards increased engagement of *CALB2*-targeted cells in population
activity (calcium signaling events were detected in 19/100 *CALB2*-targeted cells (19%) under baseline
conditions, and in 30/100 cells (30%) following nicotine application; calcium signals were detected in
32/65 *FOXP2*-targeted cells at baseline (49.2%) and in 33/65 cells (50.8%) following nicotine application
(n=3 slices for each); *CALB2,* p=0.07 and *FOXP2,* p=0.999 by Fisher exact test, Figure 5G top). In active
CALB2-targeted cells, nicotine application was found to increase calcium transients relative to baseline
conditions (*CALB2*-targeted cells: 2.4 ± 1.1 $\Delta F/F$ under baseline conditions, 6.0 ± 2.6 $\Delta F/F$ following
nicotine application, 2.52 ± 0.9 fold increase; *FOXP2*-targeted cells: 1.6 ± 0.4 $\Delta F/F$ under baseline
conditions, 1.0 ± 0.4 $\Delta F/F$ following nicotine application, 0.94 ± 0.6 fold change; 2-way ANOVA
interaction, p=0.045, Figure 5G bottom).

DISCUSSION

359   To advance our understanding of the human brain and devise more effective neurotherapies, it is
360   essential to investigate the organization and function of human neural circuits at cell type-resolution. *Ex*
361   *vivo* human tissues hold significant potential as a platform for studying human neural circuits, but the lack

of facile and scalable tools for targeting cell types limits the discovery and translational impacts of this model system. Here we demonstrate that the CellREADR RNA sensor-effector technology enables specific, effective, and programmable recording and manipulation of marker-defined neuronal types in *ex vivo* human brain tissues. Our findings highlight CellREADR's promise as a genetic tool for human neuroscience and therapeutic applications.

Specificity is of foremost importance for cell type-targeting technologies. While we found that *CALB2* and *FOXP2* sensors exceeded 75% specificity, based on immunohistochemical validations (Figure 1), these measures may be underestimates, as immunohistochemistry is technically challenging in human slice cultures due to factors such as lipofuscin autofluorescence and reagent availability (e.g., many commercial antibodies are not raised against human antigens). Indeed, our single-cell PatchSeq, which assessed comprehensive transcriptional profiles of READR-labeled cells, demonstrated 95% specificity (18/19 cells tested) for the *CALB2* sensor (Figure 2). Our ongoing testing of multiple sensors for *PVALB*, *SST*, and *UNC5B* has yielded promising READRs for other cell types (Figure 1C), but quantitative validation awaits further testing through RNA *in situ* hybridization, antibody staining, or RNA-seq analysis. Future studies will implement large-scale screens of sensor libraries in human cell culture systems and *ex vivo* tissues. Such efforts will facilitate sensor identification and design algorithms and ultimately establish comprehensive RNA sensor libraries for human neuron types.

Payload expression level is also critical to the functionality of a cell-targeting tool. While CellREADR can potentially be delivered as *in vitro* transcribed mRNAs using lipid nanoparticles[73,74], here we focused on DNA and AAV-based delivery, which may be better-suited for long-term payload expression in brain cells. In our singular vector design, direct coupling of payload translation to the sensor yielded mNeonGreen fluorescence in live tissues 6-9 days after virus application. The binary vector design substantially enhanced payload expression through tTA-TRE transcriptional amplification, achieving live mNeonGreen labeling within 3 days of virus application. Also, the binary vector system expressed fluorescent proteins, calcium indicators and opsins at levels that enabled targeted patch clamp (Figures 2 and 3), optogenetic manipulation (Figure 4), and calcium imaging (Figure 5) of human cell types. As CellREADR is orthogonal to and complementary with transcriptional enhancer-based tools, we were able to use it together with an interneuron class-specific enhancer, DLX-2.0, to investigate human interneuron synaptic connectivity (Figure 4). Our optogenetic data extend from previous studies[19,37,39] to provide the first proof-of-principle for cell type-specific optical interrogation of synaptic connectivity in human tissues. Likewise, our CellREADR calcium imaging experiments (Figure 5) also stand as the first demonstration of cell type-specific calcium imaging in human microcircuits[39].

Capsid (i.e., serotype) selection is prerequisite to effective viral delivery and expression of the genetic payload in the target population, and should be validated from the outset of CellREADR applications[25]. The AAV-PHP.eB (as in Lee et al.[33]) and rAAV2-retro (here) appear to have broad tropism for human neuron classes, and thus can be paired with CellREADR for targeting neuronal types. In addition, the promoter used to drive READR expression must be active in the targeted cells. Here we used a human synapsin promoter[75], which appears to be broadly active in mammalian neurons. Prior to pairing with CellREADR, candidate promoters should likewise be validated in populations that include the target cell type.

To date, transcriptional enhancers are the best-described tools for targeting NHP and human brain cell populations. For example, the DLX2.0 enhancer[21] provides cross-species (NHP, human, as well as mouse) access to a broad set of cortical GABAergic interneurons[33], and the mouse *Scn1a* enhancer targets the parvalbumin+ subclass in several mammalian species, including humans[19]. Additional interneuron subclass-specific enhancers have recently been identified through screening and validation in mice, and some of these may be of utility for targeting human interneurons[76]. With advances in single cell epigenome analysis and large-scale enhancer discovery, this approach will continue to yield useful cell type targeting tools, which could also be combined with CellREADR sensors in the same vector for increased specificity. On the other hand, as enhancer screening and validation are typically carried out in mice, the degree to which they are conserved and thus effective in human cell "homologs" remains to be established, though some encouraging results have emerged[19,21,22,76]. Also, recent findings indicate the human brain contains transcriptomic cell types that are unique from those in other mammals[9,46,47]. Studying these human-unique cell types and their related circuits may require dedicated tools; in this context, CellREADR RNA sensors may be better suited to directly target these cells' specific RNA

markers. Alternatively, sensors can be selected for conserved RNA targets, thereby enabling utilization across model organisms. The *CALB2* and *FOXP2* sensors used here were designed specifically for human sequences, but future sensor designs may prioritize such cross-species functionality.

In summary, our results demonstrate that the CellREADR RNA sensor-effector technology can achieve specific, effective, and versatile monitoring and manipulation of human neuron types in *ex vivo* brain tissues. Significant efforts, including large-scale sensor screens in cell cultures and organotypic brain preparations, will be necessary to establish comprehensive sensor tools for human brain cell types. Future efforts will also be directed to refining the specificity of cell type access through the intersectional targeting of 2 or more markers and devising multiplex strategies for differential recording and manipulation of multiple cell types in the same preparation. Other applications may pair CellREADR with a range of tool sets for circuit breaking and cellular control, including neurotransmitter reporters, voltage indicators, retrograde transsynaptic tracers and sonogenetic effectors. Synergistic with and complementary to capsid engineering and enhancer-based approaches, CellREADR will elevate the human *ex vivo* tissue system as a foundational platform and provide a translational bridge towards cell type and circuit specific therapeutics in the human brain.

RESOURCE AVAILABILITY

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Derek Southwell (derek.southwell@duke.edu)

Materials availability
All unique materials generated for this study are available from the lead contact.

Data and code availability
Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

ACKNOWLEDGMENTS

This study was supported in part by NIH grants 1DP2MH140149 (D.G.S.), 1DP1MH129954-01 (Z.J.H.), and 1K08NS133292 (J.B.R). J.B.R. was supported by the NINDS K12 Child Neurologist Career Development Program. D.G.S was supported by the NINDS K12 Neurosurgery Research Career Development Program, The Klingenstein-Simons Fellowship Award in Neuroscience, The Whitehall Foundation, and the Ruth K. Broad Foundation for Biomedical Research. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We thank the Duke Light Microscopy Core Facility and the Duke Sequencing and Genomics Core for their support in data acquisition and analysis. We are grateful to the patients who donated surgical tissue specimens for this research.

AUTHOR CONTRIBUTIONS

Concept and methodology: E.A.M., D.G.S., Z.J.H., J.B.R., Y.Q.
Data acquisition and analysis: E.A.M., J.B.R., S.Z., P.T., M.M.
Resources: D.G.S., M.L.V. (surgical tissue specimens); Y.Q., S.Z., Z.J.H. (CellREADR viruses)
Writing: D.G.S., E.A.M., Z.J.H., J.B.R.
Funding: D.G.S., Z.J.H.

DECLARATION OF INTERESTS

Z.J.H. has filed patents for CellREADR (US patent #: PCT/US22/79004 & PCT/US22/7900). Z.J.H is a co-founder of Doppler Bio. All other authors have no conflicts of interest to declare.

SUPPLEMENTAL INFORMATION

Document S1. Figures S1–S5 and Table S1 & S2

FIGURE TITLES AND LEGENDS

Figure 1. Deploying CellREADR to target RNA-defined cell types in ex vivo human cortex

(A) Human organotypic slice model. Human neocortical tissue specimens (here, middle and inferior temporal gyri; left) were sectioned into 350 μm slices and cultured on semi-permeable membranes for up to 16 days *in vitro* (DIV; right). CellREADR AAV vectors were directly applied to slices on DIV 0. Orientation: A, anterior; D, dorsal; P, posterior; V, ventral.

(B) Schematic of CellREADR design. In the single virus design (left), binding of the sensor RNA sequence to the target leads to ADAR-mediated editing of a stop codon positioned between the sensor and the effector, leading to expression of the downstream effector (here, mNeon). The binary system (right) uses a CellREADR virus (top right) to drive expression of tTA2 (Tet-Advanced transactivator) in the target cell. A reporter virus, designed to conditionally express a selected effector molecule (e.g. mNeon, ChIEF, GCamp7f, depending on experiment) under control of the Tet-Responsive Element (TRE), is co-applied.

(C) CellREADR targeting of diverse neuronal populations. Binary CellREADRs (mNeonGreen reporter) designed against Somatostatin (*SST*), Unc-5 netrin receptor B (*UNC5B),* Parvalbumin (*PVALB*), Forkhead box protein P2 (*FOXP2),* and Calretinin (*CALB2*) were applied to slices of human neocortex from at least 3 donors. Slices were immunostained for mNeon at 7 DIV. The CellREADR binary system led to robust expression of mNeon in differing laminar distributions and in cells of various morphologies. Dashed red boxes highlight the area depicted in the accompanying magnified image; pia and white matter (WM) are illustrated with dotted lines. Scale bars: 200 μm in large image and 50 μm in magnified image

(D) Histological analysis of binary *FOXP2* and *CALB2* CellREADR targeting. CellREADR mNeon distribution (green, left) and protein expression of target (purple, middle) are depicted at DIV 7. The fluorescence intensity profile to the far left of each image set reveals the distribution of cells targeted by the CellREADR. Dashed red boxes indicate areas depicted in (F); pia and white matter (WM) are illustrated with dotted lines. Scale bars: 200 μm

(E) Localization of CellREADR-targeted cells. CellREADR mNeon fluorescence intensity profiles measured across the depth of the cortex (0% depth = pial surface, 100% depth = white matter boundary) averaged from 6 slices (tissues from 3 donors). The solid line shows the mean and the shaded area shows the SEM. *FOXP2*-mNeon was relatively uniform throughout cortical layers, while *CALB2*-mNeon was more restricted to outer layers.

(F) Immunohistochemical characterization of CellREADR specificity. Representative images showing colocalization of CellREADR-mNeon with the corresponding target. Scale bars: 50um

(G) Quantification of CellREADR specificities, as measured by immunostaining. Each point denotes the specificity of labeling measured from an individual donor's tissue (*FOXP2* n=5 donors; Binary *CALB2* n=7, Singular *CALB2* n=3). Horizontal bars indicate mean specificity values for each CellREADR. Throughout figures, data are presented as mean ± SEM.

(H) Benchmarking of CellREADR efficiency with a human interneuron enhancer virus DLX2.0-YFP. *CALB2* CellREADR and DLX2.0-YFP viruses (rAAV2-retro) were applied to slices cut sequentially from the same neocortical tissue specimen. Tissue was fixed at 7 DIV and immunostained against mNeon or YFP. Slice boundaries are indicated by the solid white line, and the white matter is demarcated by a dashed white line; boxes mark the inset images below. Scale bars: upper images 1 mm, lower images 200 μm

(I) Localization of DLX2.0-YFP expression. Representative image illustrating the efficiency and distribution of cellular labeling achieved with a DLX2.0-YFP virus (left), with fluorescence intensity profile (right; averages are from tissues collected from 3 donors) illustrating distribution of cells labeled by the virus. As compared to *CALB2* CellREADR labeling (Figures 1D and 1H), DLX2.0-YFP+ cells were less concentrated in the outer cortex, as would be expected for pan-interneuron targeting by DLX2.0. Scale bar: 250 μm

(J) DLX2.0-YFP targeting of the *CALB2* population. Quantification of colocalization of DLX2.0-YFP expression and CALB2, as measured by immunostaining (images not shown); note inter-subject variability in targeting of CALB2 cells (n=7 donors).

514 (K) Efficiency of binary and singular *CALB2* CellREADRs compared to DLX2.0-YFP enhancer virus.
515 Quantification of cells labeled at DIV 7, per unit area (Binary *CALB2*, n=6 donors, Singular *CALB2*, n=3,
516 DLX2.0-YFP, n=3). Note, although the CellREADRs were designed to target only *CALB2* interneurons,
517 rather than multiple subclasses interneurons (as expected for DLX2.0), they labeled more cells (1-way
518 ANOVA with Tukey's post hoc).

Figure 2: PatchSeq validation of CALB2 CellREADR targeting

(A) Reference gene expression profiles of human MTG interneurons described in a prior study. *CALB2*
expression is largely restricted to *VIP* subclass interneurons, indicating that *CALB2* READR is expected
to primarily target *VIP* interneurons. Of the 19 *CALB2* READR-targeted cells studied by PatchSeq here,
all but 1 cell mapped transcriptomically to the *VIP* subclass (the transcriptomic types of the 19 cells are
listed at right; number indicates count of cells mapped to that type) according to the classification scheme
adapted from Tasic et al.[77] and Hodge et al.[7]

(B) Expression of selected genes in *CALB2* READR cells. *CALB2* transcripts were detected in all but 1
cell, which was mapped onto the *SST* subclass (*SST KLDHC8A*). READR-targeted cells rarely expressed
canonical markers of interneuron subclasses other than *VIP* (e.g., *LAMP5, SST, PVALB*). Interestingly, 5
of the 6 cells that mapped to the *PAX6 SYT6* type expressed the *SLC17A7* (vesicular glutamate
transporter 1), which is expressed widely across human MTG glutamatergic populations (Supplemental
Figure 2A); these cells also expressed the GABAergic marker, *GAD1*.

(C) PatchSeq electrophysiological measures of *CALB2* READR-targeted neurons. Input-output curves
illustrate the variation in rheobase, gain, and maximum firing frequency across the sampled population.
The interrelationship between gene expression and physiological properties is further illustrated in
Supplemental Figure 3.

(D) Variation in electrophysiological properties across and within interneuron transcriptomic types.
Membrane responses to depolarizing and hyperpolarizing currents from cells that were mapped to the
same transcriptomic types. Cells that mapped to the *PAX6 SYT6* type exhibited variation in the frequency
and accommodation of action potential firing and also exhibited varying responses to hyperpolarization.
Scale bars: 20 mV, 200 ms

(E) Morphologies of *CALB2* READR-targeted cells from the PatchSeq subset. The positions (within
cortical mantle) of reconstructed neurons are illustrated at top. Blue circles indicate locations of the cell
somata (not to scale), black profiles indicate the vertical and horizontal densities of cell dendrites, and red
profiles indicate the vertical and horizontal densities of cell axons. Cortical depth is indicated on the left
vertical axis; average cortical layer boundaries are shown on the right vertical axis. Complete cell tracings
are shown at bottom. Supplemental Figure 5 shows the morphologies of additional *CALB2*- and *FOXP2*-
targeted cells. Scale bars: 200 μm

Figure 3: CALB2- and FOXP2-targeted neurons exhibit distinct physiological profiles and spontaneous activities

(A) Passive membrane properties of cells targeted by *CALB2* (n=60 cells) and *FOXP2* CellREADRs
(n=43). Membrane capacitance and input resistance, but not hyperpolarizing sag ratio or resting
membrane potential, differed statistically between the two target populations. Black bars indicate mean
and SEM (Mann-Whitney U-test for non-normal distributions, t-test for normal distributions).

(B) Active properties of cells targeted by *CALB2* (n=60 cells) and *FOXP2* CellREADRs (n=43). Firing
responses of cells in response to current injections, with frequency plotted against current step. Insets
show action potentials elicited by current injections at 2X rheobase; these examples were selected to
represent the mean input-output curves for each cell type (black). The rheobase (current that initiated
firing), gain (initial slope), and max firing frequency all differed statistically between *CALB2* and *FOXP2*-
targeted populations (Mann-Whitney U-test for non-normal distributions, t-test for normal distributions).

(C) Action potential (AP) properties of cells targeted by *CALB2* (n=60 cells) and *FOXP2* CellREADRs
(n=43 cells). Action potentials were elicited with a 2ms current step. Action potential properties from
different stimulation regimes are presented in Supplemental Figure 4. Action potential half-width was
measured at 50% of peak voltage (height) and was found to be smaller in *CALB2*-targeted cells. Action
potential height and threshold did not vary between target cell groups. The fast afterhyperpolarization
(fAHP) following the AP was larger in *CALB2*-targeted cells. Action potentials were aligned on their rising
phase to compare AP waveforms between target cell groups (lower panel). Group means are illustrated
by the black lines. *CALB2*-targeted cells exhibited a shorter AP duration and a smaller afterdepolarization
than *FOXP2*-targeted cells (Mann-Whitney U-test for non-normal distributions, t-test for normal
distributions). Scales bars: left 20 mV, 2 ms; center 20 mV, 2 ms; right 5 mV, 20 ms

(D) Subthreshold synaptic activity and suprathreshold firing were observed in targeted cells (held at
resting membrane potential with 0 pA current). Excitatory and inhibitory postsynaptic potentials (EPSPs
and IPSPs, respectively) were detected in both *CALB2*- and *FOXP2*-targeted populations. Representative
traces demonstrating EPSPs and IPSPs in cells of both populations are shown. Scale bars: 1 mV, 200 ms

(E) Recorded cells exhibited spontaneous firing from their resting potentials. Examples of spontaneous
firing activity observed in *CALB2*- (red, n=3 cells) and *FOXP2*-targeted cells (blue, n=3) are illustrated,
with average firing frequencies (observed over a 3-minute period) indicated to the right of the traces.
Scale bars: 20 mV, 1000 ms

(F) *CALB2*-targeted cells (n= 66 cells) were more likely than *FOXP2*-targeted cells (n=50) to exhibit
spontaneous firing in the slice preparation (Chi-squared test).

(G) Spontaneous activity, measured as the frequency of spontaneous action potential firing, differed
between *CALB2- and FOXP2*-targeted populations (Mann-Whitney U test).

(H) Traces demonstrating spontaneous action potential firing from *CALB2*- (red) and *FOXP2*-targeted
cells (blue) in either baseline conditions (left) or in the presence of glutamatergic blockers (right). Scale
bars: 20 mV, 1000 ms

(I) Glutamatergic blockers (CNQX/AP-5) reduced spontaneous firing frequency in most cells (*CALB2*, n=6
cells; *FOXP2*, n=2), but did not eliminate firing in every cell.

Figure 4: Cell type-specific optical manipulation of human neurons with CellREADR
(A) CellREADR expression of an optogenetic effector, ChIEF, enables optogenetic control of action
potential firing in *CALB2*- but not *FOXP2*-targeted cells. *CALB2*-ChIEF neurons fired action potentials
(APs) in response to 100 ms, 642 nm light pulses (left) or trains of 2 ms pulses (6/6 cells fired action
potentials in response to both stimuli, with some failures at the 4th stimulus). Light stimuli depolarized
FOXP2-ChIEF, but did not drive AP firing (0/10 cells fired action potentials in response to stimuli; grey
traces, individual trials from a single cells, black traces, averages). Scale bars: 20mV, 20ms, 50ms

(B) Representative targeting scheme for optical interrogation of *CALB2* synaptic connectivity. Putative
postsynaptic cells were labeled with a DLX2.0-YFP virus (left top, green). A binary CellREADR vector
driving *CALB2*-ChIEF-tdTomato was used to activate presynaptic cells (right top, red). During optical
stimulation of the ChIEF-tdTomato$^+$ population, patch clamp recordings were made from YFP$^+$/ChIEF-
tdTomato$^-$ cells; at the end of recordings, biocytin-filled patched cells were recovered (right bottom,
purple, SAV). Scale bar: 50 μm

(C) Postsynaptic currents elicited by CALB2-ChIEF activation. Left, voltage clamp recordings (average of
10 trials in all cases) made from a DLX2.0-YFP$^+$/ChIEF-tdTomato$^-$ neuron during optical stimulation of the
CALB2-ChIEF population. Stimulation elicited optical postsynaptic currents (oPSCs), which were present
at the glutamate reversal potential ($V_{rev(Glu)}$) but absent at the GABA reversal potential ($V_{rev(GABA)}$),
indicating that they were mediated by synaptically released GABA. Bicuculline was applied to verify
observed oIPSCs were GABAergic. In DLX2.0-YFP$^+$/ChIEF-tdTomato$^-$ cells, oPSCs were never observed

at $V_{rev(GABA)}$. Right, averaged traces recorded from a hSyn-mCherry$^+$ pyramidal neuron during optical
stimulation of the *CALB2*-ChIEF population. In the recording shown here, oPSCs were detected at
$V_{rev(GABA)}$, but not $V_{rev(Glu)}$, indicating the cell received glutamatergic inputs from *CALB2*-ChIEF cells. These
inputs were not blocked by bicuculline. No cell exhibited oPSCs at both $V_{rev(GABA)}$ and $V_{rev(Glu)}$. Scale bar:
left 0.2 pA, 50 ms; 2 pA, 50ms

(D) Failure rate of optically triggered synaptic currents. To illustrate trial-to-trial variability in oPSC
responses, 10 trials from a DLX2.0-YFP$^+$/ChIEF-tdTomato$^-$ cell are shown at $V_{rev(Glu)}$. The first light pulse
had a failure rate of 10% (1/10), while the fourth light pulse had a failure rate of 40% (4/10; averaged
response, black). Scale bar: 2 pA, 50 ms

(E) Waveforms of optically-evoked postsynaptic responses. Optical PSCs measured at $V_{rev(Glu)}$ from
DLX2.0-YFP$^+$/ChIEF-tdTomato$^-$ interneurons and hSyn-mCherry+ pyramidal neurons were aligned to the
light pulse and overlaid to illustrate the variability in amplitudes (left). All oPSCs were normalized to the
peak current (right). The monophasic rise time and small jitter relative to the 2 ms light stimulus indicate
that PSCs were monosynaptic events directly triggered by the optical pulse (averaged response, black).
Scale bar: left 2 pA, 10 ms; 0.2, 5 ms.

(F) Quantification of postsynaptic responses to *CALB2*-ChIEF activation. Recordings were made from
DLX2.0-YFP$^+$/*CALB2*-ChIEF$^-$ interneurons during *CALB2*-ChIEF activation. Optical inhibitory PSCs
(oIPSCs, oPSCs detected at $V_{rev(Glu)}$) were observed in 6/16 cells (37.5%), and optical excitatory PSCs
(oEPSCs, oPSCs detected at $V_{rev(GABA)}$) were not detected. Recordings were also made from hSyn-
tdTomato$^+$ pyramidal neurons during *CALB2*-ChIEF activation. Optical IPSCs were detected in 3/14 cells
(21.4%), and oEPSCs were detected in 2/14 cells (14.3%).

Figure 5: Cell type-specific monitoring of population activity with CellREADR
(A) Correlation between action potential firing and GCamp7f signals. Action potentials (black) were
elicited by current injections in single cells expressing either *CALB2*-GCamp7f (n=4 cells) or *FOXP2*-
GCamp7f (n=4; top left, current injection is represented by gray box; the upper inset shows the voltage
trace at an expanded time scale to show the individual action potentials (APs)). Calcium transients
(green) were concurrently imaged at the cell soma and proximal dendritic compartment. Traces show 6
APs and resultant GCaMP ΔF/F signal measured from a *CALB2*-GCamp7f cell. The relationship between
the baseline-subtracted change in fluorescence and AP firing (right) was established by varying the
current injection, counting APs, and fitting the ΔF/F to action potential counts with a linear function
(*CALB2*: R^2 = 0.84; *FOXP2*: R^2 = 0.93). Scale bar: inset 20 mV, 100 ms; 20 mV, 2000 ms

(B) Calcium imaging detects subthreshold events as well as action potentials. Recording of a *CALB2*-
GCaMP7f demonstrating three evoked action potentials (black trace, at grey box) followed by a
spontaneous subthreshold depolarization then a single spontaneous action potential. The calcium
transient for these three events (green) is depicted; the rise times for each event were as follows: Tau_{3AP}
= 390 ms, $Tau_{Depolarization}$ = 1760 ms, Tau_{1AP} = 202 ms. Scale bars: inset 20 mV, 100 ms; 20 mV, 2000 ms

(C) Maximum projection of *CALB2*-GCamp7f (left, red) and *FOXP2*-GCamp7f (right, blue) neurons
measured in baseline conditions (standard artificial cerebrospinal fluid). Background-subtracted
fluorescent profiles of cell somata were collected from 90 s video stacks (red and blue traces).
CellREADR typically labeled 10-50 cells within a field of view. Scale bars: 50 μm, 20 s

(D) Color maps depicting population calcium signal dynamics. Calcium signals from slices labeled with
CALB2-GCamp7f (left, red; n=6 slices) or *FOXP2*-GCamp7f cells (right, blue n=7) are represented by a
color map (intensity scale at bottom; each map represents a recording performed on a single slice, slices
were collected from 4 donors). Spontaneous activity levels varied between slices. Calcium signal traces
from selected individual cells are depicted to the right of the color maps; most transients were small-
amplitude, fast-rising events, similar to the transients produced by APs in (A). Large-amplitude, slow-

rising transients (marked with asterisks) similar to the subthreshold events in (B), were also observed.
Scale bars: population maps 20 s, individual traces 10 s

(E) Correlated calcium events in slice populations. The correlation of cellular *CALB2*-GCaMP7f (n=6
slices) or *FOXP2*-GCaMP7f (n=7) calcium signals was measured within each slice and then compared
between groups (*CALB2*- and *FOXP2*-targeted populations). Calcium signals were more strongly
correlated in the *FOXP2*-targeted population (Unpaired t-test).

(F) Calcium signal responses to nicotinic acetylcholine receptor activation. Nicotine (300 µM) was applied
to a subset of slices following imaging of baseline activity. Each panel represents a ROI in a single slice
labeled with either *CALB2*-GCaMP7f (left, red; n=3 slices) and *FOXP2*-GCaMP7f (right, blue; n=3). Scale
bar: 20 s

(G) Nicotine application did not change the proportion of active cells within the labeled populations
(*CALB2*, p=0.08; *FOXP2*, p=0.999 by Fisher exact test). However, nicotine application increased the area
of measured calcium events, normalized to baseline areas in *CALB2*-targeted cells (p = 0.045 by 2-way
ANOVA with Fisher's LSD post-hoc). Dashed line on both graphs represents unity.

TABLES

Table 1. Donor demographics and specimen utilization

Donor ID	Age	Sex	Region of neocortex	Data collected
TE22	8	M	R parietal	*FOXP2* physiology
TE23	60	M	L temporal	*FOXP2* physiology
TE29	43	M	L temporal	*CALB2* physiology & morphology
TE30	52	M	R temporal	*CALB2* morphology
TE31	28	M	R temporal	*CALB2* morphology
TE32	12	F	L temporal	*SST* expression
TE33	45	M	L temporal	*UNC5B* expression, *CALB2* physiology & morphology
TE34	19	F	R temporal	*CALB2* physiology & morphology
TE35	30	M	L temporal	*PVALB* expression
TE36	35	F	R temporal	*CALB2* expression
TE37	23	M	R temporal	*CALB2* physiology
TE41	42	M	R temporal	*CALB2* physiology
TE44	14	M	L temporal	*CALB2* physiology
TE45	41	M	L temporal	*CALB2* physiology
TE46	43	F	L temporal	*CALB2* physiology
TE47	28	F	L temporal	*FOXP2* expression & physiology, Ca imaging
TE48	40	F	R temporal	*FOXP2* physiology
TE50	35	F	R temporal	*FOXP2* physiology, Ca imaging
TE51	29	F	R temporal	*FOXP2* physiology
TE52	19	M	L temporal	*FOXP2* physiology, Ca imaging
TE53	33	M	R temporal	*FOXP2* physiology, Ca imaging
TE57	34	F	L temporal	Ca imaging
TE65	47	F	L temporal	*CALB2* physiology
TE66	50	F	R temporal	*FOXP2* physiology

REFERENCES

1. Zeng, H., and Sanes, J.R. (2017). Neuronal cell-type classification: challenges, opportunities and the path forward. Nat. Rev. Neurosci. *18*, 530–546.

2. Konopka, G., and Bhaduri, A. (2023). Functional genomics and systems biology in human neuroscience. Nature *623*, 274–282.

3. Jorstad, N.L., Song, J.H.T., Exposito-Alonso, D., Suresh, H., Castro-Pacheco, N., Krienen, F.M., Yanny, A.M., Close, J., Gelfand, E., Long, B., et al. (2023). Comparative transcriptomics reveals human-specific cortical features. Science *382*, eade9516.

4. Siletti, K., Hodge, R., Mossi Albiach, A., Lee, K.W., Ding, S.-L., Hu, L., Lönnerberg, P., Bakken, T., Casper, T., Clark, M., et al. (2023). Transcriptomic diversity of cell types across the adult human brain. Science *382*, eadd7046.

5. Jorstad, N.L., Close, J., Johansen, N., Yanny, A.M., Barkan, E.R., Travaglini, K.J., Bertagnolli, D., Campos, J., Casper, T., Crichton, K., et al. (2023). Transcriptomic cytoarchitecture reveals principles of human neocortex organization. Science *382*, eadf6812.

6. Bakken, T.E., Jorstad, N.L., Hu, Q., Lake, B.B., Tian, W., Kalmbach, B.E., Crow, M., Hodge, R.D., Krienen, F.M., Sorensen, S.A., et al. (2021). Comparative cellular analysis of motor cortex in human, marmoset and mouse. Nature *598*, 111–119.

7. Hodge, R.D., Bakken, T.E., Miller, J.A., Smith, K.A., Barkan, E.R., Graybuck, L.T., Close, J.L., Long, B., Johansen, N., Penn, O., et al. (2019). Conserved cell types with divergent features in human versus mouse cortex. Nature *573*, 61–68.

8. Hawrylycz, M.J., Lein, E.S., Guillozet-Bongaarts, A.L., Shen, E.H., Ng, L., Miller, J.A., van de Lagemaat, L.N., Smith, K.A., Ebbert, A., Riley, Z.L., et al. (2012). An anatomically comprehensive atlas of the adult human brain transcriptome. Nature *489*, 391–399.

9. Krienen, F.M., Goldman, M., Zhang, Q., C H Del Rosario, R., Florio, M., Machold, R., Saunders, A., Levandowski, K., Zaniewski, H., Schuman, B., et al. (2020). Innovations present in the primate interneuron repertoire. Nature *586*, 262–269.

10. Yao, Z., van Velthoven, C.T.J., Kunst, M., Zhang, M., McMillen, D., Lee, C., Jung, W., Goldy, J., Abdelhak, A., Aitken, M., et al. (2023). A high-resolution transcriptomic and spatial atlas of cell types in the whole mouse brain. Nature *624*, 317–332.

11. BRAIN Initiative Cell Census Network (BICCN) (2021). A multimodal cell census and atlas of the mammalian primary motor cortex. Nature *598*, 86–102.

12. Tasic, B., Menon, V., Nguyen, T.N., Kim, T.K., Jarsky, T., Yao, Z., Levi, B., Gray, L.T., Sorensen, S.A., Dolbeare, T., et al. (2016). Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. Nat. Neurosci. *19*, 335–346.

13. Shi, Y., Wang, M., Mi, D., Lu, T., Wang, B., Dong, H., Zhong, S., Chen, Y., Sun, L., Zhou, X., et al. (2021). Mouse and human share conserved transcriptional programs for interneuron development. Science *374*, eabj6641.

14. Lee, A.T., Chang, E.F., Paredes, M.F., and Nowakowski, T.J. (2024). Large-scale neurophysiology and single-cell profiling in human neuroscience. Nature *630*, 587–595.

15. He, M., and Huang, Z.J. (2018). Genetic approaches to access cell types in mammalian nervous systems. Curr. Opin. Neurobiol. *50*, 109–118.

16. Feng, G., Jensen, F.E., Greely, H.T., Okano, H., Treue, S., Roberts, A.C., Fox, J.G., Caddick, S.,
Poo, M.-M., Newsome, W.T., et al. (2020). Opportunities and limitations of genetically modified
nonhuman primate models for neuroscience research. Proc. Natl. Acad. Sci. U. S. A. *117*, 24022–
24031.

17. Taniguchi, H., He, M., Wu, P., Kim, S., Paik, R., Sugino, K., Kvitsiani, D., Fu, Y., Lu, J., Lin, Y., et al.
(2011). A resource of Cre driver lines for genetic targeting of GABAergic neurons in cerebral cortex.
Neuron *71*, 995–1013.

18. Huang, Z.J., and Zeng, H. (2013). Genetic approaches to neural circuits in the mouse. Annu. Rev.
Neurosci. *36*, 183–215.

19. Vormstein-Schneider, D., Lin, J.D., Pelkey, K.A., Chittajallu, R., Guo, B., Arias-Garcia, M.A., Allaway,
K., Sakopoulos, S., Schneider, G., Stevenson, O., et al. (2020). Viral manipulation of functionally
distinct interneurons in mice, non-human primates and humans. Nat. Neurosci. *23*, 1629–1636.

20. Mich, J.K., Graybuck, L.T., Hess, E.E., Mahoney, J.T., Kojima, Y., Ding, Y., Somasundaram, S.,
Miller, J.A., Kalmbach, B.E., Radaelli, C., et al. (2021). Functional enhancer elements drive
subclass-selective expression from mouse to primate neocortex. Cell Rep. *34*, 108754.

21. Graybuck, L.T., Daigle, T.L., Sedeño-Cortés, A.E., Walker, M., Kalmbach, B., Lenz, G.H., Morin, E.,
Nguyen, T.N., Garren, E., Bendrick, J.L., et al. (2021). Enhancer viruses for combinatorial cell-
subclass-specific labeling. Neuron *109*, 1449-1464.e13.

22. Dimidschstein, J., Chen, Q., Tremblay, R., Rogers, S.L., Saldi, G.-A., Guo, L., Xu, Q., Liu, R., Lu, C.,
Chu, J., et al. (2016). A viral strategy for targeting and manipulating interneurons across vertebrate
species. Nat. Neurosci. *19*, 1743–1749.

23. Heinz, S., Romanoski, C.E., Benner, C., and Glass, C.K. (2015). The selection and function of cell
type-specific enhancers. Nat. Rev. Mol. Cell Biol. *16*, 144–154.

24. Hardison, R.C., and Taylor, J. (2012). Genomic approaches towards finding cis-regulatory modules
in animals. Nat. Rev. Genet. *13*, 469–483.

25. Challis, R.C., Ravindra Kumar, S., Chen, X., Goertsen, D., Coughlin, G.M., Hori, A.M., Chuapoco,
M.R., Otis, T.S., Miles, T.F., and Gradinaru, V. (2022). Adeno-Associated Virus Toolkit to Target
Diverse Brain Cells. Annu. Rev. Neurosci. *45*, 447–469.

26. Hoffmann, M.D., Gallant, J.P., LeBeau, A.M., and Schmidt, D. (2024). Unlocking precision gene
therapy: harnessing AAV tropism with nanobody swapping at capsid hotspots. NAR Mol Med *1*,
ugae008.

27. Castle, M.J., Turunen, H.T., Vandenberghe, L.H., and Wolfe, J.H. (2016). Controlling AAV tropism in
the nervous system with natural and engineered capsids. Methods Mol. Biol. *1382*, 133–149.

28. Matsuzaki, Y., Fukai, Y., Konno, A., and Hirai, H. (2024). Optimal AAV capsid/promoter
combinations to target specific cell types in the common marmoset cerebral cortex. bioRxiv.
https://doi.org/10.1101/2024.05.09.593444.

29. Sun, S., and Schaffer, D.V. (2018). Engineered viral vectors for functional interrogation,
deconvolution, and manipulation of neural circuits. Curr. Opin. Neurobiol. *50*, 163–170.

30. Campos, L.J., Arokiaraj, C.M., Chuapoco, M.R., Chen, X., Goeden, N., Gradinaru, V., and Fox, A.S.
(2023). Advances in AAV technology for delivering genetically encoded cargo to the nonhuman
primate nervous system. Curr. Res. Neurobiol. *4*, 100086.

31. Eugène, E., Cluzeaud, F., Cifuentes-Diaz, C., Fricker, D., Le Duigou, C., Clemenceau, S., Baulac,
M., Poncer, J.-C., and Miles, R. (2014). An organotypic brain slice preparation from adult patients
with temporal lobe epilepsy. J. Neurosci. Methods *235*, 234–244.

32. Schwarz, N., Uysal, B., Welzer, M., Bahr, J.C., Layer, N., Löffler, H., Stanaitis, K., Pa, H., Weber,
Y.G., Hedrich, U.B., et al. (2019). Long-term adult human brain slice cultures as a model system to
study human CNS circuitry and disease. Elife *8*. https://doi.org/10.7554/eLife.48417.

33. Lee, B.R., Dalley, R., Miller, J.A., Chartrand, T., Close, J., Mann, R., Mukora, A., Ng, L., Alfiler, L.,
Baker, K., et al. (2023). Signature morphoelectric properties of diverse GABAergic interneurons in
the human neocortex. Science *382*, eadf6484.

34. Cohen, I., Navarro, V., Clemenceau, S., Baulac, M., and Miles, R. (2002). On the origin of interictal
activity in human temporal lobe epilepsy in vitro. Science *298*, 1418–1421.

35. Gonzalez-Ramos, A., Waloschková, E., Mikroulis, A., Kokaia, Z., Bengzon, J., Ledri, M., Andersson,
M., and Kokaia, M. (2021). Human stem cell-derived GABAergic neurons functionally integrate into
human neuronal networks. Sci. Rep. *11*, 22050.

36. Dossi, E., Blauwblomme, T., Moulard, J., Chever, O., Vasile, F., Guinard, E., Le Bert, M., Couillin, I.,
Pallud, J., Capelle, L., et al. (2018). Pannexin-1 channels contribute to seizure generation in human
epileptic brain tissue and in a mouse model of epilepsy. Sci. Transl. Med. *10*.
https://doi.org/10.1126/scitranslmed.aar3796.

37. Andersson, M., Avaliani, N., Svensson, A., Wickham, J., Pinborg, L.H., Jespersen, B., Christiansen,
S.H., Bengzon, J., Woldbye, D.P.D., and Kokaia, M. (2016). Optogenetic control of human neurons
in organotypic brain cultures. Sci. Rep. *6*, 24818.

38. Kalmbach, B.E., Buchin, A., Long, B., Close, J., Nandi, A., Miller, J.A., Bakken, T.E., Hodge, R.D.,
Chong, P., de Frates, R., et al. (2018). h-Channels Contribute to Divergent Intrinsic Membrane
Properties of Supragranular Pyramidal Neurons in Human versus Mouse Cerebral Cortex. Neuron
100, 1194-1208.e5.

39. Ting, J.T., Kalmbach, B., Chong, P., de Frates, R., Keene, C.D., Gwinn, R.P., Cobbs, C., Ko, A.L.,
Ojemann, J.G., Ellenbogen, R.G., et al. (2018). A robust ex vivo experimental platform for molecular-
genetic dissection of adult human neocortical cell types and circuits. Sci. Rep. *8*, 8407.

40. Beaulieu-Laroche, L., Brown, N.J., Hansen, M., Toloza, E.H.S., Sharma, J., Williams, Z.M., Frosch,
M.P., Cosgrove, G.R., Cash, S.S., and Harnett, M.T. (2021). Allometric rules for mammalian cortical
layer 5 neuron biophysics. Nature *600*, 274–278.

41. Moradi Chameh, H., Rich, S., Wang, L., Chen, F.-D., Zhang, L., Carlen, P.L., Tripathy, S.J., and
Valiante, T.A. (2021). Diversity amongst human cortical pyramidal neurons revealed via their sag
currents and frequency preferences. Nat. Commun. *12*, 2497.

42. Wilbers, R., Galakhova, A.A., Driessens, S.L.W., Heistek, T.S., Metodieva, V.D., Hagemann, J.,
Heyer, D.B., Mertens, E.J., Deng, S., Idema, S., et al. (2023). Structural and functional
specializations of human fast-spiking neurons support fast cortical signaling. Sci. Adv. *9*, eadf0708.

43. Chartrand, T., Dalley, R., Close, J., Goriounova, N.A., Lee, B.R., Mann, R., Miller, J.A., Molnar, G.,
Mukora, A., Alfiler, L., et al. (2023). Morphoelectric and transcriptomic divergence of the layer 1
interneuron repertoire in human versus mouse neocortex. Science *382*, eadf0805.

44. Berg, J., Sorensen, S.A., Ting, J.T., Miller, J.A., Chartrand, T., Buchin, A., Bakken, T.E., Budzillo, A.,
Dee, N., Ding, S.-L., et al. (2021). Human neocortical expansion involves glutamatergic neuron
diversification. Nature *598*, 151–158.

45. Buchin, A., de Frates, R., Nandi, A., Mann, R., Chong, P., Ng, L., Miller, J., Hodge, R., Kalmbach, B.,
Bose, S., et al. (2022). Multi-modal characterization and simulation of human epileptic circuitry. Cell
Rep. *41*, 111873.

46. Boldog, E., Bakken, T.E., Hodge, R.D., Novotny, M., Aevermann, B.D., Baka, J., Bordé, S., Close,
J.L., Diez-Fuertes, F., Ding, S.-L., et al. (2018). Transcriptomic and morphophysiological evidence
for a specialized human cortical GABAergic cell type. Nat. Neurosci. *21*, 1185–1195.

47. Oláh, S., Komlósi, G., Szabadics, J., Varga, C., Tóth, E., Barzó, P., and Tamás, G. (2007). Output of
neurogliaform cells to various neuron types in the human and rat cerebral cortex. Front. Neural
Circuits *1*, 4.

48. Qian, Y., Li, J., Zhao, S., Matthews, E.A., Adoff, M., Zhong, W., An, X., Yeo, M., Park, C., Yang, X.,
et al. (2022). Programmable RNA sensing for cell monitoring and manipulation. Nature *610*, 713–
721.

49. Jiang, K., Koob, J., Chen, X.D., Krajeski, R.N., Zhang, Y., Volf, V., Zhou, W., Sgrizzi, S.R., Villiger,
L., Gootenberg, J.S., et al. (2023). Programmable eukaryotic protein synthesis with RNA sensors by
harnessing ADAR. Nat. Biotechnol. *41*, 698–707.

50. Kaseniit, K.E., Katz, N., Kolber, N.S., Call, C.C., Wengier, D.L., Cody, W.B., Sattely, E.S., and Gao,
X.J. (2023). Modular, programmable RNA sensing using ADAR editing in living cells. Nat.
Biotechnol. *41*, 482–487.

51. MacDermot, K.D., Bonora, E., Sykes, N., Coupe, A.-M., Lai, C.S.L., Vernes, S.C., Vargha-Khadem,
F., McKenzie, F., Smith, R.L., Monaco, A.P., et al. (2005). Identification of FOXP2 truncation as a
novel cause of developmental speech and language deficits. Am. J. Hum. Genet. *76*, 1074–1080.

52. Lai, C.S., Fisher, S.E., Hurst, J.A., Vargha-Khadem, F., and Monaco, A.P. (2001). A forkhead-
domain gene is mutated in a severe speech and language disorder. Nature *413*, 519–523.

53. Haesler, S., Rochefort, C., Georgi, B., Licznerski, P., Osten, P., and Scharff, C. (2007). Incomplete
and inaccurate vocal imitation after knockdown of FoxP2 in songbird basal ganglia nucleus Area X.
PLoS Biol. *5*, e321.

54. Groszer, M., Keays, D.A., Deacon, R.M.J., de Bono, J.P., Prasad-Mulcare, S., Gaub, S., Baum,
M.G., French, C.A., Nicod, J., Coventry, J.A., et al. (2008). Impaired synaptic plasticity and motor
learning in mice with a point mutation implicated in human speech deficits. Curr. Biol. *18*, 354–362.

55. Ferland, R.J., Cherry, T.J., Preware, P.O., Morrisey, E.E., and Walsh, C.A. (2003). Characterization
of Foxp2 and Foxp1 mRNA and protein in the developing and mature brain. J. Comp. Neurol. *460*,
266–279.

56. Allen Brain Atlas - ISH Data - Cortex Study https://human.brain-
map.org/ish/search/?page_num=0&page_size=400&no_paging=false&search_term=&search_type=
cortex https://human.brain-
map.org/ish/search/?page_num=0&page_size=400&no_paging=false&search_term=&search_type=
cortex.

57. Medalla, M., Mo, B., Nasar, R., Zhou, Y., Park, J., and Luebke, J.I. (2023). Comparative features of
calretinin, calbindin, and parvalbumin expressing interneurons in mouse and monkey primary visual
and frontal cortices. J. Comp. Neurol. *531*, 1934–1962.

58. Guet-McCreight, A., Skinner, F.K., and Topolnik, L. (2020). Common Principles in Functional
Organization of VIP/Calretinin Cell-Driven Disinhibitory Circuits Across Cortical Areas. Front. Neural
Circuits *14*, 32.

59. Allen Brain Atlas - Cell Types - Human MTG https://celltypes.brain-map.org/rnaseq/human/mtg
https://celltypes.brain-map.org/rnaseq/human/mtg.

60. Džaja, D., Hladnik, A., Bičanić, I., Baković, M., and Petanjek, Z. (2014). Neocortical calretinin
neurons in primates: increase in proportion and microcircuitry structure. Front. Neuroanat. *8*, 103.

61. Hladnik, A., Džaja, D., Darmopil, S., Jovanov-Milošević, N., and Petanjek, Z. (2014). Spatio-temporal
extension in site of origin for cortical calretinin neurons in primates. Front. Neuroanat. *8*, 50.

62. Allen Insitute for Brain Science scrattch: Single cell RNA-seq analysis for transcriptomic type
characterization. https://github.com/AllenInstitute/scrattch.

63. Allen Brain Map Documentation: Cell Types Database. https://community.brain-
map.org/t/documentation-cell-types-database/2845.

64. Shao, L.R., Halvorsrud, R., Borg-Graham, L., and Storm, J.F. (1999). The role of BK-type Ca2+-
dependent K+ channels in spike broadening during repetitive firing in rat hippocampal pyramidal
cells. J. Physiol. *521 Pt 1*, 135–146.

65. Lancaster, B., and Nicoll, R.A. (1987). Properties of two calcium-activated hyperpolarizations in rat
hippocampal neurones. J. Physiol. *389*, 187–203.

66. Gouwens, N.W., Sorensen, S.A., Berg, J., Lee, C., Jarsky, T., Ting, J., Sunkin, S.M., Feng, D.,
Anastassiou, C.A., Barkan, E., et al. (2019). Classification of electrophysiological and morphological
neuron types in the mouse visual cortex. Nat. Neurosci. *22*, 1182–1195.

67. Jonas, P., Bischofberger, J., Fricker, D., and Miles, R. (2004). Interneuron Diversity series: Fast in,
fast out--temporal and spatial signal processing in hippocampal interneurons. Trends Neurosci. *27*,
30–40.

68. Markram, H., Toledo-Rodriguez, M., Wang, Y., Gupta, A., Silberberg, G., and Wu, C. (2004).
Interneurons of the neocortical inhibitory system. Nat. Rev. Neurosci. *5*, 793–807.

69. Lin, J.Y., Lin, M.Z., Steinbach, P., and Tsien, R.Y. (2009). Characterization of engineered
channelrhodopsin variants with improved properties and kinetics. Biophys. J. *96*, 1803–1814.

70. Caputi, A., Rozov, A., Blatow, M., and Monyer, H. (2009). Two calretinin-positive GABAergic cell
types in layer 2/3 of the mouse neocortex provide different forms of inhibition. Cereb. Cortex *19*,
1345–1359.

71. Gulyás, A.I., Hájos, N., and Freund, T.F. (1996). Interneurons containing calretinin are specialized to
control other interneurons in the rat hippocampus. J. Neurosci. *16*, 3397–3411.

72. Meskenaite, V. (1997). Calretinin-immunoreactive local circuit neurons in area 17 of the cynomolgus
monkey, Macaca fascicularis. J. Comp. Neurol. *379*, 113–132.

73. Paunovska, K., Loughrey, D., and Dahlman, J.E. (2022). Drug delivery systems for RNA
therapeutics. Nat. Rev. Genet. *23*, 265–280.

74. Hou, X., Zaks, T., Langer, R., and Dong, Y. (2021). Lipid nanoparticles for mRNA delivery. Nat Rev
Mater *6*, 1078–1094.

75. Kügler, S., Kilic, E., and Bähr, M. (2003). Human synapsin 1 gene promoter confers highly neuron-
specific long-term transgene expression from an adenoviral vector in the adult rat brain depending
on the transduced area. Gene Ther. *10*, 337–347.

76. Furlanis, E., Dai, M., Leyva Garcia, B., Vergara, J., Pereira, A., Pelkey, K., Tran, T., Gorissen, B.L.,
Vlachos, A., Hairston, A., et al. (2024). An enhancer-AAV toolbox to target and manipulate distinct
interneuron subtypes. bioRxivorg. https://doi.org/10.1101/2024.07.17.603924.

77. Tasic, B., Yao, Z., Graybuck, L.T., Smith, K.A., Nguyen, T.N., Bertagnolli, D., Goldy, J., Garren, E.,
Economo, M.N., Viswanathan, S., et al. (2018). Shared and distinct transcriptomic cell types across
neocortical areas. Nature *563*, 72–78.

78. Gautier, A., Juillerat, A., Heinis, C., Corrêa, I.R., Jr, Kindermann, M., Beaufils, F., and Johnsson, K.
(2008). An engineered protein tag for multiprotein labeling in living cells. Chem. Biol. *15*, 128–136.

79. Brian Lee, K.H. (2021). Patch-Seq Recording and Extraction Detailed Protocol (Allen Brain Institute)
https://doi.org/10.17504/protocols.io.bw6gphbw.

80. Peng, H., Bria, A., Zhou, Z., Iannello, G., and Long, F. (2014). Extensible visualization and analysis
for multidimensional images using Vaa3D. Nat. Protoc. *9*, 193–208.

STAR⬝METHODS

KEY RESOURCES TABLE

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Chicken anti-GFP	Aves	Cat# GFP-1020; RRID:AB_10000240; 1:1000
Chicken anti-mCherry	Origene	Cat# TA150127; 1:1000
Goat anti-GFP	Rockland	Cat# 600-101-215; RRID:AB_218182; 1:1000
Mouse anti-mNeon	Proteintech	Cat# 32f6; 1:1000
Mouse anti-NeuN	Cell Signaling	Cat# 94403S; 1:1000
Rabbit anti-Calretinin	Swant	Cat# CR7697; RRID:AB_2619710; 1:1000
Rabbit anti-FoxP2	Abcam	Cat# ab16046; RRID:AB_2107107; 1:1000
Virus strains		
AAVphp.eb-DLX2.0 -YFP	AddGene	Cat# 163505-PHPeb; RRID:Addgene_163505
AAV-DLX2.0-YFP plasmid	AddGene	Cat# 163505; RRID:Addgene_163505
AAVrg-hSyn-mCherry	AddGene	Cat # 50459-AAVrg; RRID:Addgene_50459
Chemicals		
ATP-triphosphate, magnesium salt	SigmaAldrich	Cat# A9187
Bicuculline methiodide	Tocris	Cat# 2503
Biocytin	SigmaAldrich	Cat# B4261
Calcium Chloride, dihydrate	SigmaAldrich	Cat# C7902
Cesium Chloride	SigmaAldrich	Cat# 289329
Cesium methylsulfonate	SigmaAldrich	Cat# C1426
CNQX	Tocris	Cat # 190
D-(+)-Glucose	SigmaAldrich	Cat# G7528
DL-AP-5	Tocris	Cat# 105
EGTA	SigmaAldrich	Cat# 03777
GTP-triphosphate disodium salt	SigmaAldrich	Cat# 371701
HEPES	SigmaAldrich	Cat# H3537

Magnesium Chloride	SigmaAldrich	Cat# M4880
Na2-ATP	SigmaAldrich	Cat# A6419
Nicotine ditartrate	SigmaAldrich	Cat# 481975
Potassium Chloride	SigmaAldrich	Cat# P5405
Potassium gluoconate	SigmaAldrich	Cat# G4500
QX-314 bromide	Tocris	Cat# 1014
RNase inhibitor	Takara	Cat# 2313A
Sodium bicarbonate	SigmaAldrich	Cat# S5761
Sodium Chloride	SigmaAldrich	Cat# 31434
Sodium Phosphate, monobasic	SigmaAldrich	Cat# S9638
Thiodiethanol	SigmaAldrich	Cat# 166782
Tris-Phosphocreatine	SigmaAldrich	Cat# P1937
Critical commercial assays		
Bioanalyzer High Sensitivity DNA Analysis	Agilent	Cat# 5067-4626
SMART-seq v4 ultra low input mRNA kit	Takara	Cat# 634772
Software and algorithms		
BioRender App	BioRender	https://www.biorender.com/
GraphPad Prism	GraphPad Software, Inc.	https://www.graphpad.com
IgorPro 9.0	Wavemetrics	https://www.wavemetrics.com
Neuromatic Plugin for IgorPro	Freeware	www.neuromatic.thinkrandom.com
PClamp 10	Molecular Devices	
RStudio	Posit	https://posit.co/products/open-source/rstudio
Scrattch package for R	Allen Institute	https://github.com/AllenInstitute/scrattch

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Human tissue procurement

892
893 Human cortical tissue was donated by patients who underwent neurosurgical treatments for epilepsy.
894 Non-diseased tissue, which was surgically removed to access areas of disease pathology, was used for
895 this study. All donors provided informed consent under an IRB Protocol (00103019) at Duke University.
896 Biological sex was recorded, but gender identity was not collected. Donor demographics are listed in
897 Table 1.

METHOD DETAILS

Plasmids

899
900 CellREADR sensors were generated using gBlock synthesis (Integrated DNA Technologies). Vector
901 backbones were linearized using restriction digestion and DNA fragment inserts were generated using
902 PCR or gBlock synthesis (Integrated DNA Technologies).

CellREADR sensor RNA (sesRNA) design

903
The sensors used in this study were designed as previously described[48]. In brief, sensors designed for
both coding sequences and 5' untranslated regions of each target gene, depending on the gene length.
Synthesized sensor sequences were cloned into CAG-BFP-sesRNA-GFP constructs. We also
synthesized the target gene (coding sequence and 5'UTR) using gBlock and cloned them into expression
plasmids with CAG promoter. We tested the efficiency and specificity of some sensors in HEK cells using
methods we have previously described[48], and, when performed, used those with the highest specificity for

further study. Plasmids contained sequences to promote stability and intended cleavage of the
transduced product (Clip-F, WPRE, W3SL)[78] .

Viruses
CellREADR viruses were designed and prepared using methods we have previously described[48]. In brief,
for production of READR and reporter viruses, HEK293T cells were transfected with READR or Reporter
plasmids, AAV serotype plasmids, and pHelper using PEI MAX (Polysciences, 24765). Seventy-two hours
after transfection, the cells were collected in cell culture medium, and centrifuged at 4,000 rpm for
15 min. The supernatant was discarded, the pellet was resuspended in cell lysis buffer, frozen, and
thawed three times using a dry ice/ethanol bath. Cell lysate was centrifuged at 4,000 rpm for 20 min.
Contaminating DNA in the supernatant was removed by benzonase treatment. The crude viral
preparation was loaded onto an iodixanol density gradient and spun at 60,000 rpm for 90 min using a
Beckman Ti70 rotor. After the centrifugation, 3–4 ml of crude viral preparation was collected from the
40–60% layer with an 18-gauge needle attached to a 10-ml syringe. The viral crude solution was
concentrated using Amicon Ultra-15 centrifugal filters (100 kDa), washed with 8 ml PBS once, and
concentrated to an appropriate volume.) AAVrg-DLX2.0-YFP (Addgene #163505) and AAVrg-hSyn-
mCherry (Addgene #50459) were used to target interneurons or pyramidal neurons, respectively.

Human organotypic sample preparation and culture
Organotypic slices from surgical samples were prepared according to published methods[31,32]. Dissection
solution contained (in mM): 75 sucrose, 87 NaCl, 25 glucose, 26 $NaHCO_3$, 2.5 KCl, 1.2 NaH_2PO_4, 10
$MgCl_2$, 0.5 $CaCl_2$, bubbled with 95/5% O_2/CO_2 (pH 7.4). Cortex was carefully dissected under ice cold
(~4C) oxygenated dissection solution to remove the pia. Single gyri were blocked for slicing and cut at
350µm thickness on a vibratome. Slices were rinsed in Hank's Buffered Saline and plated on PTFE
membranes (Millipore, cat. # PICMORG50) in 6 well plates. Tissue was cultured in a conditioned media[31].
Conditioned media also contained antibiotic/antimycotic (1X, Gibco by ThermoFisher) for the first 7 days
in vitro, and cyclosporine (5 µg/mL) for the entire culture period. HEPES (20mM) was added to the media
for the first hour after slice preparation. Slices were cultured in an incubator at 35 Celsius with 5% CO_2
and 5% humidity. Virus-mediated expression of fluorescent reporters was monitored beginning at 3 DIV
until they were used for immunohistochemistry (DIV 3-16), patch clamp electrophysiology (DIV 4-12), or
calcium imaging experiments (DIV 4-7).

Virus application
On the day of tissue donation, after slicing and plating the tissue on semi-permeable membranes and
incubating slices in HEPES containing media, viruses were directly applied to the surface of the slice.
Viruses were diluted in media (CellREADR virus, 1:8 dilution; TRE-effector viruses, 1:6 dilution; AAVrg-
DLX2.0-YFP and AAVrg-hSyn-mCherry 1:10 dilution), and 7µL of the virus solution was applied to slice
grey matter. Plates were carefully returned to the incubator and left undisturbed overnight.

Immunohistochemistry
After 7 of incubation, or upon completion of patch clamp recordings, cortical slices were fixed overnight in
4% PFA, then rinsed in PBS and stored for 1-7 days in PBS with azide. For biocytin recovery or
colocalization IHC, tissue was permeabilized in PBS with 10% TritonX100, 5% NGS, and 100µM glycine
2 hours at room temperature on a shaker plate.

Primary antibodies were incubated 48 hours at 4C; secondary antibodies were incubated for 24 hours at
4C. Biocytin was developed with either Alexa647 conjugated streptavidin, or with a silver stain. Large
overview images of native mNeon fluorescence were made on a Leica SP8 upright confocal microscope,
or a Keyence BZ-X800 with a 10X or 20X objective.

The following antibodies were used for immunohistochemical labeling of mNeon, mCherry, FoxP2,
Calretinin and NeuN: mouse anti-mNeon Ab 1:500, Proteintech 32f6; chicken anti-mCherry Ab 1:500,
Origene TA150127; rabbit anti-FoxP2 Ab 1:500, Abcam ab16046; rabbit anti-Calretinin Ab 1:500, Swant
CR7697; mouse anti-NeuN Ab 1:500, Cell Signaling #94403S. Secondary antibodies were Alexa488,
Alexa594, or Alexa647 against the appropriate species (Invitrogen). Slices were mounted using a medium
containing DAPI.

Some slices were selected for chemical clearing to enhance imaging deeper into the slice and reduce
lipofuscin autofluorescence. These slices were incubated in thiodiethanol (TDE) solution of 30% for 35

minutes, 60% for 35 minutes, then 75% for 10 minutes. Slices were mounted in a 300μm thick spacer ring
(SunJin Labs, iSpacer, IS307), then the remaining volume was filled with 75% TDE before coverslip
placement.

Image analysis for colocalization & cell location
Immunohistochemistry measurements of CellREADR specificity were performed by immunostaining
against either CALB2 or FOXP2 protein and CellREADR mNeonGreen. Maximum projection images of
small z-stacks (40-60 μm total depth) were used for manual counts of mNeon-labeled cells and CALB2 or
FOXP2 using the CellCounter Plugin in FIJI. Colocalization was expressed as the percentage of
mNeonGreen+ cells that also expressed the marker protein of interest (CALB2 or FOXP2).

Electrophysiology
Slices were transferred to a heated recording bath (32-34C) on an Olympus BX-50 upright microscope.
Recording solution contained (in mM): 118 NaCl, 3 KCl, 25 $NaHCO_3$, 1 NaH_2PO_4, 1 $MgCl_2$, 1.5 $CaCl_2$, 30
glucose fully saturated with 95/5% O_2/CO_2. Borosilicate patch pipettes were pulled with a resistance of
3.5-5.5 MΩ, and filled with an internal solution containing (in mM): 134 K-gluconate, 10 HEPES, 4 ATP-
triphosphate Mg salt, 3 GTP-triphosphate Na salt, 14 Phosphocreatine, 6 KCl, 4 NaCl, pH adjusted to 7.4
with KOH for current clamp recordings For voltage clamp recordings (optogenetic stimulation of PSCs),
the internal solution continued (in mM): 100 Cs-MeSO3, 40 CsCl, 4 MgCl2, 0.1 Na2-GTP, 4 Na2-ATP, 10
Tris-Phosphocreatine, 10 HEPES, and 4 QX-314. Biocytin (0.5%) was added to the internal solution to
allow for morphological identification after recording. Patched cells were held in whole-cell mode for a
minimum of 10 minutes to ensure complete filling with biocytin.

Labeled cells were targeted for patch clamp under a 60X objective. Cells were sampled from across
cortical layers, in manners that evenly reflected where the *CALB2* and *FOXP2* CellREADR constructs
were expressed (Figure 1). Recordings were performed using an Axopatch 700B amplifier (Molecular
Devices), and data were digitized (Digidata 1550B) and captured with pClamp 10 (Molecular Devices).
Membrane voltage was recorded at 100k Hz and low-pass filtered at 10k Hz. Liquid junction potential was
not corrected. Pipette capacitance and series resistance were compensated at the start of each recording
and checked periodically for stability of the recording configuration. Intrinsic membrane properties (input
resistance, capacitance) were measured with a -10 pA current step. Rheobase was measured with a
ramp current of 1s duration and 100-300 pA final amplitude. Input-output curves were generated from a
series of current steps starting at -100 pA and increasing in 10 pA increments until a maximum firing rate
was elicited. Sag ratio was calculated from the -100pA step, and was the difference between the negative
peak deflection from baseline and the steady-state voltage normalized to the peak deflection[63]. In a
subset of cells glutamatergic blockers (20 μM CNQX + 50 μM AP-5) were bath applied to block excitatory
synaptic inputs. Data were analyzed using NeuroMatic and custom scripts in Igor Pro (Wavemetrics).

Optical stimulation experiments were performed by expressing either AAVrg-DLX2.0-YFP, AAV9-hSyn-
GFP or AAV9-hSyn-mCherry in putative postsynaptic cells, and CALB2-ses1-tTA and TRE-ChIEF-YFP or
TRE-ChIEF-tdTom CellREADR constructs in putative pre-synaptic populations. Cells that 1) expressed
only the DLX2.0-YFP or hSyn-mCherry, and 2) were situated in areas of CellREADR labeling, were
targeted for voltage clamp recordings. Blue light pulses (625 nm, 5x 2ms-long pulses at either 10Hz or
50Hz) illuminating the whole field of view were triggered for 10 sweeps while recording membrane
currents at -30 mV (V_{rev} GABA) and at 0 mV (V_{rev} Glu). If an optical response was detected in the average
of the 10 sweeps, then a GABAergic blocker (50 μM bicuculline) was bath-applied to the slice, then
stimulation was repeated to identify the neurotransmitter that produced the PSC.

Processing of cellular nuclei for PatchSeq
PatchSeq electrophysiology was performed using an internal solution that included (in mM): 110 K-
gluconate, 4 KCl, 10 HEPES, 1 ATP-Mg, 0.3 GTP-Na,10 Na-phosphocreatine, 0.2 EGTA, 20 μg/mL
glycogen, 0.5% biocytin, and 0.5 U/μL RNase inhibitor (Takara). The internal solution was prepared with
RNase-free water and stored in daily aliquots at -80 C until use. Equipment, materials preparation and
nuclei extraction followed previously published protocols[79]. Clean nucleus extraction was confirmed
visually with both DIC and fluorescent images. Nuclei were stored at -80C in 10μL of lysis buffer with
RNase inhibitor (40 U/μL) and then processed for sequencing in batches (5-12 nuclei/batch). Nuclear
RNA was processed into cDNA libraries using the SMART-Seq v4 Ultra Low Input kit (Takara) using 19
PCR cycles. Quality checks of the resulting cDNA were made on a Bioanalyzer (Agilent, High Sensitivity

DNA kit). Nuclei that yielded >200 pg/µL cDNA were sequenced. Borderline cDNA (concentrations just
below 200 pg/µL, and fragment sizes >500 bp) were re-amplified with 3 rounds of PCR, rechecked for
cDNA quality, and included in the sequencing set if reamplification increased cDNA concentration above
threshold.

<u>RNA sequencing and transcriptomic analysis</u>
cDNA sequencing was performed at the Duke Sequencing and Genomics Technologies core facility on
an Illumina NovaSeq 6000 flow cell with a targeted sequencing depth of approximately 50 million reads
per sample. Data were aligned to the human genome (GRCh38, annotations from ENSEMBL v106). Raw
count matrices from aligned data were analyzed using the Allen Institute scrattch.hicat (version 1.0.0)
package (https://github.com/AllenInstitute/scrattch.hicat) in R (2022.07.1 Build 554). Normalized counts
were calculated as log2(counts-per-million reads per cell) for downstream analysis. Heatmaps for select
genes were generated using the gplots package (version 3.1.3) in R.

Predicted transcriptomic cell type annotations were derived from a reference dataset of 15,928 nuclei
from the human middle temporal gyrus (MTG) that underwent SMART-seq single nucleus RNA
sequencing[59]. Annotation predictions were generated using scrattch.hicat (version 1.0.0) according to a
standard data analysis pipeline (https://taxonomy.shinyapps.io/scrattch_tutorial/#section-overview). The
reference MTG dataset was limited to interneuron clusters (class = GABAergic); cell class and cluster (i.e.
Inh L1-3 PAX6 SYT6) were used to generate reference cluster names. The reference count matrix and
the PatchSeq test count matrix were both normalized (log2(counts-per-million(raw matrix) +1)) and key
cluster specific genes were selected using the "select_markers" function in scrattch.hicat. Cluster
predictions for PatchSeq cells were then generated using the "map_sampling" function in scrattch.hicat
(markers.perc=0.8, iter=100).

To investigate differential gene expression between *CALB2*+ and *FOXP2*+ neurons we utilized an
existing single-nucleus RNA sequencing atlas from human MTG[59] . The atlas dataset was reconstituted in
Seurat via CreateSeuratObject, and then the raw count matrix was normalized and scaled via
SCTransform with default parameters. The human MTG Seurat object was further subsetted for cells with
non-zero expression of *CALB2* or *FOXP2*, yielding *CALB2*+ GABAergic neurons and *FOXP2*+
Glutamatergic neurons. Differential gene expression was then analyzed between these two populations
via FindMarkers(object, ident.1 = "GABAergic, ident.2 = "Glutamatergic", only.pos = FALSE), yielding
log2-fold changes in expression, percent gene expression in ident.1 and ident.2, and a p-value adjusted
for multiple comparisons where p-adj <0.05 was considered significant differential expression. The list of
differentially expressed genes was then manually investigated for significantly differentially expressed
genes of interest related to neuromodulation (e.g.. nicotinic acetylcholine receptor subunits), for the
studies of calcium imaging.

<u>Morphology</u>
For cortical layer identification, 4 sections from 4 donors were labeled with DAPI and imaged from the pia
to the white matter border. The density of nuclei delineated layer boundaries; the distances between each
layer were calculated for the 4 samples and averaged to create a "typical" cortical layer map. Our cortical
layer distances were as follows: L1-L2 boundary, 221.8 µm, L2-L3 boundary, 383.9 µm, L3-L4 boundary,
812.2 µm, L4-L5 boundary, 1677.6 µm, L5-L6 boundary, 2206.9 µm, L6-WM boundary, 3549.5 µm. This
is a slightly larger average thickness than reported elsewhere (2963µm)[38]. When a biocytin-filled cell was
identified, the distance from the pia to the cell soma was measured, as was the underlying Layer 6-white
matter boundary (these were measured along a line perpendicular to the overlying pia). Soma position
was then calculated as a percent depth (0% is located at the pial surface and 100% is located at the
Layer 6-white matter boundary. This normalized the depth position of the soma across slices. DAPI
labeling was used as a secondary criterion to ascertain the soma laminar position.

For morphological tracings, biocytin-filled neurons were imaged using a 40x objective on a Leica SP8
upright confocal microscope (oil immersion, NA = 1.3). Tiled z-stacks containing cell dendrites and axons
were captured at 1024x1024 resolution, with a z-step size of 1.0-1.2 µm. Overview images were captured
using a 10x objective to provide a reference for cell orientation relative to the pial surface. Images were
traced using the automatic tracing plugin in Vaa3D[80] and converted to .SWC format. The automatic
tracing function sometimes produced erroneous branching points, and so all traces were inspected and
manually corrected using either the SNT plugin in FIJI, or neuTube, a plugin for Vaa3D. To orient the

traces relative to the pial surface, corrected .SWC files were aligned in the SNT Viewer using the 10X
overview images to position all cells vertically in the cortical column.

Calcium Imaging

Four to 7 days after virus application, slices with robust GCaMP7f fluorescence were selected for
experiments. Slices were placed in a bath chamber at 32 C under a 10x objective and a high-power LED
(Thor Labs) at 488 nm. Videos of 60-90 s duration were captured at a frame rate of 100 ms with a Retiga
Electro 1.4 M pixel CCD camera (Teledyne Photometrics) with no binning. Imaging was performed in
supragranular cortex, where the greatest density of cellular labeling was observed with both the *CALB2*
and *FOXP2* CellREADRs. FIJI was used to check the standard deviation projection of the video for
spontaneous activity. If activity was detected, several videos were then captured in the same region, then
a different area of the slice was imaged. Between 4 and 10 videos were captured from each field of view
over the course of 10-15 minutes. Nicotine (300 µM) was bath-applied to a subset of slices, then videos
were recaptured 5 minutes later from the same field of view. After allowing nicotine to wash out for 15
minutes, baseline imaging was then performed at a new location.

One representative video of each field of view was selected if it contained a relatively high density of
labeled cells and exhibited minimal motion artifacts. Data were analyzed with ImageJ (FIJI) and custom
scripts in Igor Pro. If necessary, motion artifacts were corrected with the Image Stabilizer plugin.
Maximum projection images were used to select all virally neurons in the field of view for analysis (non-
active cells were also analyzed). ROIs were assigned to each neuron, and intensity values over time were
extracted for each ROI. Calcium signals were background subtracted and normalized to a baseline period
that exhibited no activity. Co-activation was calculated by a pairwise cross-correlation (Pearson's) of all
the ROIs in the field of view. Pearson's correlation coefficients were converted to Fisher's z-values for
comparison across slices and between groups. Activity during the nicotine application was normalized to
baseline activity in standard recording solution.

QUANTIFICATION AND STATISTICAL ANALYSIS

All data values in the text are the mean +/- SEM; error bars in the figures indicated the SEM. Statistical
tests were performed in GraphPad Prism. Data were checked for normality; normally distributed data
were compared with t-test or ANOVA. Non-normal data distributions were compared with the Mann-
Whitney U-test. Differences in proportions were tested with the Fisher's exact test (small sample size) or
Chi-squared test.

ADDITIONAL RESOURCES

Creative Commons license for figure Panels created with the BioRender app:

Graphical Abstract: Created in BioRender. Matthews, E. (2025) https://BioRender.com/0pgjkzm

Figure 2, Supplemental Figure 2, & Supplemental Figure 5: Created in BioRender. Matthews, E. (2025)
https://BioRender.com/qhb53fv

1103

A)

B)

Singular Vector

hSyn ClipF sensor 2a mNeon W3SL

Binary Vectors

READR AAV

hSyn ClipF sensor 2a tTA2 W3SL

Reporter AAV

TRE3g effector WPRE pA

C)

Pia

SST-mNeon

UNC5B-mNeon

PVALB-mNeon

FOXP2-mNeon   WM

CALB2-mNeon

D)

FOXP2 - mNeon

mNeon   FOXP2   Merge   WM

E)

mNeon Intensity

Cortical Depth (%)

FOXP2   *CALB2*

CALB2 - mNeon

Pia

mNeon   CALB2   Merge

F)

FOXP2   *CALB2*

mNeon   mNeon

FOXP2   CALB2

Merge   Merge

G)

Colocalization (%)

○ Binary *FOXP2*
○ Binary *CALB2*
△ Singlular *CALB2*

H)

CALB2 - mNeon   DLX2.0 - YFP

I)

Pia

YFP   WM

YFP intensity

Cortical Depth (%)

DLX2.0

J)

DLX2.0-YFP Colocalization (%)

CALB2

K)

Cell Counts (cells/mm²)

p = 0.04

Binary   Singular   DLX2.0