GraphVelo allows for accurate inference of multimodal velocities and molecular mechanisms for single cells

Yuhao Chen[1,2,a], Yan Zhang[2,a], Jiaqi Gan[2,a], Ke Ni[2], , Ming Chen[1,3,b], Ivet Bahar[4,b], Jianhua Xing[2,5,6,b]

1. *Department of Bioinformatics, College of Life Sciences, Zhejiang University, Hangzhou, 310058, China*
2. *Department of Computational and Systems Biology, School of Medicine, University of Pittsburgh, Pittsburgh, PA, USA*
3. *Zhejiang Key Laboratory of Multi-omics Precision Diagnosis and Treatment of Liver Diseases, Department of General Surgery, Sir Run-Run Shaw Hospital, Zhejiang University School of Medicine, 310016, Hangzhou, China.*
4. *Laufer Center for Physical and Quantitative Biology, and Department of Biochemistry and Cell Biology, Renaissance School of Medicine, Stony Brook University, Stony Brook, NY, USA*
5. *University of Pittsburgh Hillman Cancer Center, Pittsburgh, PA, USA*
6. *Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, PA, USA*

[a] Equal contribution

[b] Corresponding authors: xing1@pitt.edu, bahar@laufercenter.org, mchen@zju.edu.cn.

Abstract

RNA velocities and generalizations emerge as powerful approaches for extracting time-resolved information from high-throughput snapshot single-cell data. Yet, several inherent limitations restrict applying the approaches to genes not suitable for RNA velocity inference due to complex transcriptional dynamics, low expression, or lacking splicing dynamics, or data of non-transcriptomic modality. Here, we present GraphVelo, a graph-based machine learning procedure that uses as input the RNA velocities inferred from existing methods and infers velocity vectors lying in the tangent space of the low-dimensional manifold formed by the single cell data. GraphVelo preserves vector magnitude and direction information during transformations across different data representations. Tests on multiple synthetic and experimental scRNA-seq data including viral-host interactome and multi-omics datasets demonstrate that GraphVelo, together with downstream generalized dynamo analyses, extends RNA velocities to multi-modal data and reveals quantitative nonlinear regulation relations between genes, virus and host cells, and different layers of gene regulation.

Introduction

Cells need to constantly detect and adapt to changes in extracellular and intracellular environment. Regulation of their gene transcription is a common mechanism of response. Multiple factors affect the transcriptional activity of eukaryotic genes, including *cis* and *trans* regulatory elements and chromatin structure. High throughput single cell sequencing data provide the landscape of cell genotype. These data lack, however, information on how the genome state changes over time Continuous efforts have been made to extract information about gene regulation and developing methods for connecting the cell states to temporal sequences of events captured by single-cell snapshot data. One group of methods that has received extensive attention is that based on RNA velocity[1] for predicting the changes in RNA expression states in the cell. The original RNA velocity method leverages the ratio between nascent and mature transcripts to estimate the rate of change in gene expression. This seminal study has inspired numerous methods for improved RNA velocity estimation based on information from splicing[2-6], metabolic labeling[7,8], lineage tracing[9], and transcriptional factor binding[10].

The RNA velocity framework has, however, its own inherent limitations. First, none of the RNA velocity estimation methods could be applied to any single cell transcriptomic data without restrictions. For example, the splicing-based method is not applicable prokaryotes or viruses, or organisms without introns. Erroneous inferences of RNA velocities have also been noticed for genes having complex splicing dynamics[11]. Furthermore, it is difficult to estimate the RNA velocities of genes with low expression, which excludes most transcription factors. Second, multi-omics sequencing technologies provide multifaceted information on cellular states alongside the transcriptome modality, and currently there are limited systematic methods to extend such velocity estimations to other modalities[12,13].

The single cell transcriptome state is usually defined by the instantaneous distribution of RNA levels, represented by a multidimensional vector of RNA levels for all (measurable) genes. The usual practice is to map such single cell data onto a reduced space, e.g., transform from a principal component (PC) space to a UMAP representation to facilitate the visualization of the time-evolution of cell state, each cell state being represented by a point in that space. Numerous dimension reduction and manifold learning algorithms have been developed for such representation transformation. In comparison, transforming a velocity vector between

representations is a nontrivial task not rigorously addressed in the single cell field. Even worse, a

visually correct vector field does not necessarily imply accurate high-dimensional velocity

estimation[14]. La Manno et al. proposed a cosine kernel method to address this challenge, which

has been adopted since then in most subsequent studies[1]. Li et al. mathematically proved that the

cosine kernel asymptotically gives the correct direction of a velocity vector in the large sampling

limit, but the magnitude information is completely lost due to a normalization procedure[15]. This

loss of information casts concerns when such quantitative information is needed.

In this study, we tackle the above challenges through a graph-theoretical representation of RNA

velocities, called GraphVelo, with dynamical systems underpinnings. GraphVelo takes an ansatz

that the measured single cell expression profiles and inferred RNA velocities collectively reflect a

dynamical process and are connected through a set of dynamical equations. It exploits such

additional constraints that couples high dimensional velocity field and single cell state manifold,

and enables the generalization of the approach in the context of multi-modal single cell data. While

the combined expression and velocity information has been widely used to infer cell state transition

trajectories[2,16], GraphVelo presents the advantage of enabling downstream analyses such as that

performed by dynamo[7] to extract quantitative information on causal gene-gene relations that

dictate the cell state transitions. Benchmarking of the proposed graph framework against simulated

and experimental single cell data lends support to its broad utility.

Results

GraphVelo infers manifold-consistent single cell velocity vectors through tangent space
projection and transforms between representations through local linear embedding

87
88

Consider that the internal state of a cell can be specified by an N-dimensional state vector x, with

$N \gg 1$ generally. Assume that the temporal evolution of the cell state follows a continuous and

smooth curve $x(t)$ (see Methods for more general mathematical formulation). The instant velocity

vector $v(x,t) = \mathrm{d}x(t)/\mathrm{d}t$ is always tangent to the curve of $x(t)$ (as a function of t) at x. One can

generalize to the situation that the trajectories of a swarm of cells form a M-dimensional manifold

$\mathcal{M}(x)$ embedded in the N-dimensional state space with $M \ll N$ typically, as revealed by high-

throughput single cell omics data. Then under the ansatz that a velocity vector $v(x,t)$ dictates the

evolution of a state vector $x(t)$, v must lie in the tangent space of $\mathcal{M}(x)$, denoted as $T_p \mathcal{M}$. In

practice, the RNA velocity vectors inferred from existing methods do not automatically satisfy this

tangent space requirement (but see[4]).

Taking various inferred single cell RNA velocity vectors, e.g. splicing-based, metabolic labeling-

based, or lineage tracing-based, as input, GraphVelo takes advantage of the nature of the low-

dimensional cell state manifold to: 1) refine the estimated RNA velocity to satisfy the tangent

space requirement; 2) infer the velocities of non-transcriptomic modalities using RNA velocities.

GraphVelo thus serves as a plugin that can be seamlessly integrated into existing RNA velocity

analysis pipelines, and help process single cell data for downstream cellular dynamics analyses

using methods such as dynamo (Fig. 1a).

GraphVelo approximates the tangent space at a cell state x by a k-nearest neighbor (kNN) graph

following the local linear embedding algorithm[17], and uses the more reliable data manifold $\mathcal{M}$ to

refine the velocity vectors by imposing the constraint that the N-dimensional velocity vector v

should lie in the tangent space (Fig. 1b&c). Consider a given point x_i on a manifold corresponding

to the expression state i of the single cell. Its infinitesimal neighborhood forms a Euclid space that

approximates the tangent space $T_p\mathcal{M}$. With sufficient sampling of the neighboring cell states j in

the state space, the incremental displacement vectors between cell state i and its neighboring cell

states, $\delta_{ij} = x_j - x_i$, form a set of complete albeit possibly redundant and nonorthogonal/non-

normalized basis vectors of the Euclidean space in the local region. Then the projection of the

measured velocity vector onto $T_p\mathcal{M}$ can be expressed as a linear combination (see also

Supplemental Text I),

117
$$v_{\parallel}(x_i) = \sum_{j \in \mathcal{N}_i} \phi_{ij} \delta_{ij,} \tag{1}$$

118 where $\mathcal{N}_i$ is the neighborhood of cell state i, defined by its k nearest neighbors in the feature space

119 determined by sequencing profiles. Direct application of eq. 1 to determine the coefficients ϕ_{ij} is

120 numerically unstable in real data (see Supplemental Text II for detailed discussion). Instead, we

121 performed the projection by optimizing the following tangent space projection (TSP) loss function

122 (Fig. 1c),

123
$$\mathcal{L}(\phi_i) = \|v_i - v_{\parallel i}\|^2 - b \cdot \cos(\phi_i, \phi_i^{corr}) + \lambda \|\phi_i\|^2, \tag{2}$$

where $\|\cdot\|$ refers to vector modulus. ϕ_i^{corr} is a heuristic "cosine kernel" widely used in the RNA

velocity analyses for projecting velocity vectors onto a reduce space, the elements of which are

ϕ_{ij} $(with\ j \in \mathcal{N}_i)$ (see also Supplemental Text III); the second term $\cos(\cdot,\cdot)$ denotes the cosine

similarity. The first term in the loss function learns the correctly-scaled velocity magnitudes, and

the second term retains the reliable direction information based on previous mathematical analyses

showing that $\boldsymbol{\phi}_i^{corr}$ asymptotically gives correct direction of the velocity vector[15]. The L-2

regularization is used to bound parameters ϕ_i . b and λ are two hyperparameters for retaining the

direction and for regularization, respectively.

With local linear embedding, it is straightforward to transform velocity between different

representations. Assuming a mapping function f exists connecting manifold $\mathcal{M}$ and $\aleph$ such that

for cell i with state vector $\boldsymbol{x}_i$ in $\mathcal{M}$, the coordinate of the same cell in $\aleph$ is given as $\boldsymbol{y}_i = f(\boldsymbol{x}_i)$.

Since a given local patch of continuous manifold is approximated by a Euclidean space, a locally

linear transformation connects the patch in the two representations. Consequently, for a vector

described by eq. 1 in $\mathcal{M}$, the velocity vector in $\aleph$ is,

$$v_{\parallel}(\boldsymbol{y}_i) = \sum_{j \in \mathcal{N}_i} \phi_{ij} \boldsymbol{\delta}_{ij}', \tag{3}$$

where $\boldsymbol{\delta}_{ij}' = \boldsymbol{y}_j - \boldsymbol{y}_i$.

Therefore, eq. 1-3 form the mathematical and computational foundation of GraphVelo. With eq. 3

one can extend velocity inference to datasets that velocity inference is not traditionally applicable

such as host-virus interactome and multiomics datasets based on the Whitney embedding

theorem[18]. Details of the mathematical foundation were given in Methods. With velocity vectors

refined with GraphVelo, one can readily perform downstream analyses, as exemplified in Fig.1d-

f, which will be further elaborated below in the context of specific applications.

Benchmark studies demonstrate effectiveness of GraphVelo across simulation datasets with diverse topology

To demonstrate the effectiveness of the geometry-constrained projection, we first benchmarked

our method on a 3D bifurcation system constrained on a 2D manifold (Methods). We added a

random component vertical to the tangent plane to mimic the noise. The resulting velocity vectors

inferred by GraphVelo through minimizing the TSP loss were consistent with the ground truth

vectors (Fig. 2a). Both GraphVelo and cosine kernel successfully removed the normal components

(Fig. 2b i) and maintained the directional information (Fig. 2b ii), but only GraphVelo kept the

velocity magnitude information (Fig. 2b iii).

Next, we performed multifaceted evaluations of the ability of GraphVelo to robustly recover the
transcriptional dynamics across a range of simulated datasets with different underlining phenotypic
structures. We used dyngen[19], a multi-modal scRNA-seq simulation engine, to generate gene-wise
dynamics defined by gold-standard transcriptional regulatory networks (Methods). We generated
simulated scRNA-seq data for networks with a variety of underlying linear, cyclic, and bifurcating
topological structures, and recovered the corresponding vector field using GraphVelo-corrected
velocity vectors (Fig. 2c-e). To comprehensively assess the outcome, we used three diverse
metrics, cosine similarity, root-mean-square error (RMSE), and accuracy, which evaluate the
correctness of velocity direction, magnitude, and sign, respectively. We presented (Fig. 2c-e) the
comparative results obtained with the cosine kernel and with GraphVelo (TSP with (i.e., eq. 2 with
$b \neq 0$) and without (eq. 2 with $b = 0$) the cosine regularization term). By minimizing TSP loss,
GraphVelo preserved both the direction and magnitude of the vector field (Fig. 2f-h). With an
increase of noise level by adding Gaussian noise to the ground truth vectors, GraphVelo refined
the distorted velocity and outperformed the cosine kernel projection consistently (Extended Data
Fig. 1a-c).

Then, we tested whether manifold constraints could preserve the speed of the cell progression
across different representations. GraphVelo was able to scale velocity vectors between the original
space and the PCA space, showing a high correlation with the ground truth, even as noise levels
increased, whereas the cosine kernel failed (Extended Data Fig. 1d). The results on UMAP showed
less agreement, which is not surprising. UMAP is a convenient representation for visualizing single
cell data but not designed for representing quantitative cell state transition dynamics since UMAP
is not a continuous transformation from the original gene space and cannot preserve local distances
after projection.

To further explore whether GraphVelo could correct the RNA velocity estimated by the splicing
kinetics, we took the velocity inferred using different packages (scVelo[2], dynamo[7] and VeloVI[5])
as input. The output from GraphVelo agreed significantly better with the ground truth compared
to the raw input (Extended Data Fig. 1e), highlighting the significant improvement achieved by
GraphVelo in evaluating both the direction and magnitude of the velocity vector fields across all
datasets.

GraphVelo infers quantitative whole genome RNA velocity from a subset of genes with manifold-consistent RNA turnover kinetics

Most RNA velocity methods are based on biophysical models of mRNA turnover dynamics with specific assumptions that may break down in certain cases[11]. For example, the splicing-based RNA velocity may have an erroneous sign for processes under active regulation on mRNA degradation or promotors switching between states with different transcription efficiency (Fig. 3a, Extended Data Fig. 2). GraphVelo first uses the RNA velocities inferred from any method as input, then uses the velocities of high-confidence genes to infer velocities of other genes. One can use several existing approaches on evaluating the confidence scores of inferred RNA velocity values of genes[7]. Alternatively, we identified a subset of Manifold-consistent Kinetics (MacK) Genes based on their agreement with prior knowledge or additional information acquired from other methods such as lineage tracing (Fig. 3b),

We first applied GraphVelo to a mouse erythroid maturation dataset[20]. This study provided a transcriptional landscape of the erythroid lineage with well-documented differentiation trajectory during mouse gastrulation. Previous analyses have shown that the dataset contains genes with multiple rate kinetics, leading to erroneous prediction of the cell state transition direction[20,21]. We selected the top 200 out of 450 velocity genes as MacK genes, representing those with robustly estimated velocities (see Methods for details). The projected vector field in UMAP showed consistency with prior knowledge in developmental biology (Fig. 3c). We then used the corrected RNA velocities for dynamo velocity field analyses. The vector field-based pseudotime accurately predicted the lineage with scRNA-seq data of temporal mouse embryos (Fig. 3d).

Previous studies identified multiple rate kinetics (MURK) genes showing transcription bursts in the middle of erythroid differentiation[21]. For example, *Smim1* and *Hba-x* are two MURK genes showing complex patterns of phase portrait (Fig. 3f i). Consequently, the RNA velocity of *Simi1* inferred with scVelo was negative along a major part of the developmental axis (Fig. 3f ii), contradicting the trend of increasing *Simi1* mRNA levels (Fig. 3f iv). For *Hba-x*, scVelo even failed to infer its RNA velocity. On the other hand, GraphVelo inferred velocities and predicted correct kinetic patterns of these genes (Fig. 3f iii). Similar performances have been observed in other MURK genes (Extended Data Fig. 3a). To examine the overall prediction of cell state transitions from transcription burst genes, we projected the MURK genes velocity inferred from

GraphVelo and scVelo to the predefined UMAP. The velocities from GraphVelo but not scVelo

correctly captured the directional flow of differentiation using only MURK genes (Extended Data

Fig. 3b).

Next, we examined the inferred RNA velocities of an entire set of 2,000 highly variable genes.

For each gene we calculated the cross-boundary correctness (CBC) score, which quantifies how

likely, following its current velocity, a cell can develop to a target cell state[22]. GraphVelo

outperformed several existing tools that only used hundreds of 'well-estimated' kinetics genes (Fig.

3g). Furthermore, we estimated the speed of cell state transition using the norm of velocity vector

in high-dimensional space and identified the transcriptional surge stage (Extended Data Fig. 3c).

We hypothesized that the MURK genes, which exhibited a sudden increase in transcription rate

during this stage, were responsible for the sharp acceleration in cell state transition speed. Using

dynamo, we estimated the acceleration derived from the GraphVelo vector field and found that the

acceleration value, as the derivative of the velocity vector, demonstrated its potential as a predictor

for transcription burst genes (Extended Data Fig. 3d).

With the velocity estimation extended to the whole gene space, we were able to perform

comprehensive mechanistic analyses on the entire genome spectrum. First, we calculated the

MacK score for each gene using the corrected RNA velocities. We hypothesized that a gene with

a higher MacK score indicated a better agreement between its RNA velocity vector and the

developmental axis, suggesting that the gene served as a potential lineage-driver gene. We ranked

genes based on their scores and performed GO biological process enrichment analyses for the top

genes. Indeed, the enriched processes were associated with erythropoiesis, including the heme

biosynthetic process and interleukin-12-mediated signaling pathway[23] (Fig. 3e).

Next, we applied dynamo to perform differential geometry analyses of the vector field and

mechanistically dissected the activation cascade of erythroid marker gene *Klf1* (Extended Data Fig.

4a&b). Jacobian analyses based on GraphVelo vector field revealed sequential activation of driver

transcription factors (TFs) *Gata2*, *Gata1*, and *Klf1* during erythroid lineage differentiation, with

Gata1 subsequently repressing the expression of *Gata2* (Fig. 3h, Extended Data Fig. 4c)[7]. To

further demonstrate the crucial role of transcriptional factor *Gata1* during erythropoiesis, we

performed *in silico* genetic perturbation across all cells. Results showed that both inhibiting *Gata1*

and upregulating the *Gata1* repressor *Spi1* lead to a reversal of normal developmental flow (Fig.

3i, Extended Data Fig. 4d). The above analyses collectively suggest that activation of *Gata1* in the

blood progenitors biased its differentiation to erythropoiesis, agreeing with experimental reports[21].

To further evaluate GraphVelo, we tested the method on another dataset of human bone marrow

development[24]. This developmental process has complex progressions from hematopoietic stem

cells (HSCs) to three distinct branches: erythroid, monocyte, and common lymphoid progenitor

(CLP). Again, we used the top 100 out of 454 velocity genes as MacK genes to predict the RNA

velocities of 2,000 highly variable genes. The GraphVelo velocity field accurately recovered the

fate of cells on the sophisticated transcriptional landscape in contrast to scVelo (Fig. 3j and k,

Extended Data Fig. 5a). By combining the likelihood estimated by scVelo with the MacK score,

we identified rapid degradation and transcription burst genes whose dynamics deviated from the

RNA velocity assumptions (Extended Data Fig. 5b&c). *ANGPT1* and *RBPMS* are two examples

which were overall highly expressed in the progenitors and decreased quickly along the trajectories

(Fig. 3l), reminiscent of what was shown in Fig. 3a. These genes misled RNA velocity inference

with scVelo assuming a constant degradation rate constant. GraphVelo revealed a cell context-

specific transcription rate $\alpha = u + \frac{du}{dt}$ and degradation constant $\gamma = (u - \frac{ds}{dt})/s$, thus a

degradation wave along the differentiation path with (Fig. 3l, Extended Data Fig. 5d), consistent

with simulation result and reports on regulation of *ANGPT1* mRNA by microRNAs such as

miRNA-153-3p[25] (Extended Data Fig. 2d).

GraphVelo reconstructs host–pathogen transcriptome dynamics from infection trajectory

Continuous battle between human immune surveillance and viral immune evasion takes place in

the host cell system after viral entry. scRNA-seq data provide a massive and parallel way of

assessing the time evolution of both host and viral transcripts, unraveling the delicate inherent

dynamic of a virus-host system[26,27]. While existing splicing-based methods can robustly estimate

RNA velocity of host coding genes, it is infeasible to infer the dynamics of the viral transcriptome

due to lack of intron in the viral genome. GraphVelo enables inference of the velocities of virus

RNA abundance based on the kinetics of host transcripts velocities, as illustrated next.

We analyzed a human cytomegalovirus (HCMV) viral infection dataset to learn viral

transcriptomic kinetics in monocyte-derived dendritic cells (moDCs)[28]. The result from

GraphVelo unraveled how viral infection progressed along the transcriptional space (Fig. 4a). The

velocity vectors pointed to directions consistent with an increasing trend of the percentage of viral RNAs in a single cell, which inherently served as an indicator of the infection time course[29]. Compared to the trend obtained with the raw RNA velocities from scVelo, the vector field-based pseudotime calculated using GraphVelo-corrected RNA velocities consistently showed higher correlation with the (pseudo)temporal progression of viral infection as reflected by viral RNA percentage (Fig. 4b).

Furthermore, the examination of individual genes revealed that the RNA velocities consistently predicted the trend of the mRNA expression level change with increasing virus load (Fig. 4c, Extended Data Fig. 6a&b). Most viral genes started with a fast-increase phase, and the expressions of some genes (e.g. *UL22A*) gradually saturated at high virus load, together with the corresponding RNA velocities approaching zero. One exception is *UL122*, whose expression profile increased first then decreased to a steady state level lower than the peak value. This overshooting is characteristic of a negative feedback network structure[30]. Indeed, a recent study reported that *UL122* negatively regulates its own promotor[31]. Furthermore, comparison of MacK scores across GraphVelo, the CellRank pseudotime kernel, and randomized prediction showed that GraphVelo-computed viral RNA velocities aligned best with the transcriptome gradient of viral load (Fig. 4d). Note that the MacK score also served as a reliable predictor for dynamics-driving factors, specifically viral genes in this case (Extended Data Fig. 6d).

We further quantified the velocity norm of all viral factors as infection speed, and observed that the transcription of viral factors was significantly restricted initially, then gradually increased along the trajectory (Extended Data Fig. 6e). Interestingly, most of the genes that exhibited positive correlation with the infection speed were related to viral DNA synthesis, while those negatively correlated to the infection speed were engaged in host viral defense response[32](Fig. 4e, Extended Data Fig. 6f).

GraphVelo identifies host genomic response modules and predicts host-virus gene interactions

With GraphVelo-inferred RNA velocities, we probed the time evolution of lytic infection and the complex interplay between host and viral functional genomes. By fitting the GraphVelo velocity trends along the viral load axis, we identified genes with similar kinetic patterns (Fig. 4c). Using the smoothened velocity trends to calculate the distances, we clustered the genes into seven major

modules and visualized them on the UMAP space (Fig. 4f). Not surprisingly, viral genes were

concentrated in several enclosed regions, indicating that they formed distinct functional genomic

modules during the lytic cycle[29]. The genes showed two major dynamical features: acceleration

and deceleration along the viral load axis (Fig. 4g).

To systematically investigate whether the dichotomy between host kinetics genes and viral genes

share similar dynamics, we performed gene functional enrichment analyses of host genes residing

close to the viral gene clusters. Genes located in the deceleration part were associated with

repressing viral genome replication, which includes negative regulation of viral life cycle and

known restriction factors in antiviral responses activated in DCs such as the induction of cytokine

and chemokine responses as well as interactions with neutrophils[33,34]. In parallel, the toll-like

receptor signaling pathway, required for antiviral defense of the host, was arrested[35]. Neutrophil

related processes, which typically cooperate closely with DCs to modulate adaptive immune

responses[36], were suppressed. Therefore, the deceleration part also showed how critical set of host

factors were silenced by viral entry to achieve immune evasion.

The acceleration groups, on the other hand, demonstrated how viruses hijacked the host cell

endogenous cellular programs for virus replication. Notably, the pathways related to viral genome

replication were triggered, promoting DNA replication and transcription, such as negative

regulation of G1/S transition of mitotic cell cycle[37], cellular response to DNA damage stimulus[38]

and regulation of transcription from RNA polymerase II promoter[39]. Cells showed a shift towards

a transcriptional signature resembling the G1 phase (Extended Data Fig. 6g), agreeing with

previous report on HCMV infection[40]. Along the infection process, antiviral interferon (IFN)-γ

response of moDC cells was first activated then suppressed. These results highlighted the

organized and antagonistic strategies adopted by both host cells and viruses during their tug-of-

war for survival and proliferation.

To investigate the crosstalk between host and viral factors systematically in depth, we performed

dynamo Jacobian analyses. We scanned the entire spectrum of viral genome and delineated how

the HCMV factors silence IFN and NFκB signaling (Fig. 4i). A large proportion of the identified

viral factors functioned in evading host cell immune responses, a finding supported by several

recent studies[41-43]. *In silico* virus-directed knock out experiments revealed altered accumulation

patterns of viral transcripts (Fig. 4j). Notably, inhibition of *UL112*, which ranked first with the

total viral RNA inhibition in our analyses, led to a qualitatively distinct trajectory. These results

highlight the multifunctional *UL112* locus in the viral genome as a potential target for antiviral

intervention[29,44]. The analyses demonstrated potential usage of GraphVelo-inferred velocities for

understanding the interactions between viral and host factors, assessing the effects of perturbations

on infection, and designing potential antiviral interventions[29].

GraphVelo permits multi-omics velocity inference and chromatin dynamics analyses

The molecular anatomy during cell development entails multiple layers, and how different layers

coordinate to regulate gene expression is a fundamental problem. For example, the anagen hair

follicle features distinct lineages branching from a central population of progenitor cells. Ma et

al[45]. used SHARE-seq to capture both the transcriptome and the epigenome data simultaneously

for the lineage commitment process from transit-amplifying cells (TACs) to the inner root sheath

(IRS), cuticle layer, and medulla. Upon robust selection of estimated genes following dynamo

criteria (Methods), we further refined the RNA velocities of these genes through tangent space

projection and obtained the chromatin open/close dynamics from the corresponding scATAC data

using GraphVelo. The resultant vector field in the combined transcriptome-epigenome space

proved to reconstruct the correct multilineages differentiation paths during the anagen phase (Fig.

5a).

To test the consistency of dynamics across different modalities, we performed CellRank terminate

stage analyses[16] from the refined velocity vectors. Using GraphVelo velocities of either the RNA

modality or the ATAC modality, we accurately estimated three diverse terminal stages (Fig. 5b).

For comparison, we also performed similar analyses using MultiVelo, scVelo with all velocity

genes or robustly estimated genes in above GraphVelo studies and pseudotime-based vector field

inferred by CellRank. The 2D projection of these vector field functions also exhibited seemingly

correct velocity flow direction (Extended Data Fig. 7a). However, none of them captured the cell

fate commitment based on coarse-grained transition matrix (Fig. 5b, Extended Data Fig. 7b).

Notably, the results from the RNA modality and the ATAC modality of MultiVelo gave

inconsistent results. GraphVelo-corrected velocities, on the other hand, helped identify the top-

correlating genes towards individual terminal populations which showed agreement with previous

study[46] (Fig. 5c, Extended Data Fig. 8).

Next, we conducted differential geometry analyses based on the composite GraphVelo vector field.

We identified novel root cells, which were also characterized by chromatin potential (Fig. 5d)[45].

These novel root cells expressed distinct marker genes compared to the expected root cells using

the Wilcoxon test (Fig. 5e). Moreover, we unraveled differentially expressed markers identified

by the original study[45], as well as new differentiation-potent genes and validated their initiation

properties in another transcriptome dataset (Extended Data Fig. 9a)[46]. To further investigate how

these two distinct groups of root cells convert to other cell types, we performed a least action path

(LAP) analysis between different cell phenotypes. The expected and novel root cells converted to

the IRS terminal state following two distinct least action paths in the vector field (Extended Data

Fig. 9b&c). The two paths revealed different temporal change patterns of transcription factor

expression profiles (Extended Data Fig. 9d). We calculated the mean-squared displacement (MSD)

for every transcription factor to explore the dynamics of TFs along the path from novel root to IRS.

The result demonstrated that the fate conversion by novel root was mediated by the Shh-Runx1

signaling axis (Fig. 5f, Extended Data Fig. 9d), which has been demonstrated in human embryonic

stem cells[47] and is crucial for hair development[48]. In summary, GraphVelo unraveled the multiple

molecular mechanisms that orchestrated hair follicle morphogenesis.

With available chromatin velocity and RNA velocity, we set to quantify the coupling/decoupling

relationships between chromatin structure and gene expression for each gene (see Methods). Here

chromatin structure refers to the extent of exposure/accessibility of the gene locus to the

environment as indicated by scATACseq data, shortly called open or closed state; and chromatin

velocities refer to changes between these states as inferred from scATACseq counts at the

examined gene locus. We used dynamic time warping (DTW) distances between the velocities

from different omics layers to quantify the similarity between temporal patterns of these two

modalities for each gene. A higher DTW value indicates higher similarity. Using the elbow of the

ranked distance curve as a cutoff we identified genes that showed decoupled transcription and

chromatin structure dynamics. These decoupled genes had an accumulation of cell cycle-

dependent (CCD) genes found in previous study[49] (Fig. 5g). This group of genes showed strong

involvement in cell cycle-related processes, as indicated by GO enrichment analyses (Fig. 5h).

Close examinations indicated that the transcription of cell cycle related genes decreased along the

differentiation path, while the chromatin structure at the corresponding loci remained open (Fig.

5i, Extended Data Fig. 10). To validate this hypothesis, we further applied GraphVelo to a recently

published 10x Multiome dataset from developing human cortex[50] (Extended Data Fig. 11a).

Following the same analyses, we identified decoupled genes and found out that most of these genes

were related to cell cycle (Extended Data Fig. 11b-f). which has also been reported in a previous

MultiVelo study[13].

We further performed dynamo differential geometry analyses on the composite transcriptome-

chromatin vector field. One intriguing phenomenon observed in lineage dynamics is that *Lef1* and

Hoxc13 are the driver TFs correlated with domains of regulatory chromatin (DORCs) of *Wnt3* [45].

Differential geometry analyses on the composite vector field can go beyond correlation analyses

and provide an underlying casual mechanism. As a prerequisite for such analyses, GraphVelo-

inferred RNA and chromatin velocities of the three genes correctly predicted the trend of change

of mRNA and ATAC-seq counts (Extended Data Fig. 12a&b), in contrast to the performance of

MultiVelo (Extended Data Fig. 12c). Then, Jacobian analyses on the GraphVelo vector field

confirmed that priming activation of *Lef1* subsequently activated the *Hoxc13* TF[51] (Extended Data

Fig. 12d&e). Both *Lef1* and *Hoxc13* were found to activate the *Wnt3* target gene, initiating lineage

commitment (Extended Data Fig. 12f &g). To quantitatively understand how the two TFs affect

Wnt3 chromatin structure and transcription, we plotted the response heatmap to reflect the

distributions of Jacobian elements versus the abundance of mature mRNA for each TF (Fig. 5j).

The two terms $\partial f_{Wnt3-\text{chrom}} / \partial x_{Lef1}$ and $\partial f_{Wnt3-\text{chrom}} / \partial x_{Hoxc13}$ started with positive values at

low concentrations of TF mRNA copy numbers then decreased to zero, indicating that increasing

the level of either TF lead to further opening of the *Wnt3* chromatin region, and the effect saturated

at high TF expression. The other two term $\partial f_{Wnt3} / \partial x_{Lef1}$ and $\partial f_{Wnt3} / \partial x_{Hoxc13}$ increased with the

TF levels, indicating that these two TFs also activated Wnt3 transcription. Upon integration of the

Jacobian elements over regulator expression changes, we obtained the effective dose-response

curves obtained (see Methods), which revealed more transparently the TF dose lag between the

opening of the target chromatin region and the initialization of transcription (Fig. 5k).

Our results therefore illustrated the sequential events that these two driver TFs, *Lef1 and Hoxc13*,

drove as pioneer transcription factors (PTFs) to initiate local chromatin opening and then activated

the transcription of *Wnt3*. Notably, computational methods and experimental research confirm that

Lef1 acts as a nucleosome binder and exhibits diverse binding patterns across various cell lines[52].

The *Hox* family of TFs has also been shown to have the capacity to bind their targets in an inaccessible chromatin context and trigger the switch to an accessible state[53], consistent with our analyses that *Hoxc13* revealing that it shared a regulation mechanism similar to that of PTF *Lef1*.

Discussion

In this work, we provided a general framework that extends the framework developed for RNA velocity and related approaches to various data modalities such as proteomics, spatial genomics, 3d genome organization, and imaging data, which were originally beyond reach of this framework. We validated GraphVelo using various *in vivo* cellular kinetics models, confirmed its efficacy and robustness in handling complex and noisy multimodal data. Upon application to various datasets, we unraveled gene regulation relations of an extended list of genes, host-virus gene regulations, and coupling between transcription and local chromatin structures. GraphVelo can be seamlessly integrated with broad downstream analyses, such as dynamo continuous vector field analyses, as well as Markovian analyses using graph dynamo or CellRank.

References

1. La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498 (2018).
2. Bergen, V., Lange, M., Peidli, S., Wolf, F.A. & Theis, F.J. Generalizing RNA velocity to transient cell states through dynamical modeling. *Nat Biotechnol* **38**, 1408-1414 (2020).
3. Li, S. *et al.* A relay velocity model infers cell-dependent RNA velocity. *Nat Biotechnol* **42**, 99-108 (2024).
4. Lederer, A.R. *et al.* Statistical inference with a manifold-constrained RNA velocity model uncovers cell cycle speed modulations. *Nat Methods* (2024).
5. Gayoso, A. *et al.* Deep generative modeling of transcriptional dynamics for RNA velocity analysis in single cells. *Nat Methods* **21**, 50-59 (2024).
6. Gao, C.F., Vaikuntanathan, S. & Riesenfeld, S.J. Dissection and integration of bursty transcriptional dynamics for complex systems. *Proc Natl Acad Sci U S A* **121**, e2306901121 (2024).
7. Qiu, X. *et al.* Mapping transcriptomic vector fields of single cells. *Cell* **185**, 690-711 e45 (2022).
8. Weiler, P., Lange, M., Klein, M., Pe'er, D. & Theis, F. CellRank 2: unified fate mapping in multiview single-cell data. *Nat Methods* **21**, 1196-1205 (2024).
9. Wang, K. *et al.* PhyloVelo enhances transcriptomic velocity field mapping using monotonically expressed genes. *Nat Biotechnol* **42**, 778-789 (2024).
10. Li, J., Pan, X., Yuan, Y. & Shen, H.B. TFvelo: gene regulation inspired RNA velocity estimation. *Nat Commun* **15**, 1387 (2024).
11. Bergen, V., Soldatov, R.A., Kharchenko, P.V. & Theis, F.J. RNA velocity-current challenges and future perspectives. *Mol Syst Biol* **17**, e10282 (2021).
12. Gorin, G., Svensson, V. & Pachter, L. Protein velocity and acceleration from single-cell multiomics experiments. *Genome Biol* **21**, 39 (2020).
13. Li, C., Virgilio, M.C., Collins, K.L. & Welch, J.D. Multi-omic single-cell velocity models epigenome-transcriptome interactions and improves cell fate prediction. *Nat Biotechnol* **41**, 387-398 (2023).
14. Zheng, S.C., Stein-O'Brien, G., Boukas, L., Goff, L.A. & Hansen, K.D. Pumping the brakes on RNA velocity by understanding and interpreting RNA velocity estimates. *Genome Biol* **24**, 246 (2023).
15. Tiejun Li, T.L., Jifan Shi, J.S., Yichong Wu, Y.W. & Peijie Zhou, P.Z. On the Mathematics of RNA Velocity I: Theoretical Analysis. *CSIAM Transactions on Applied Mathematics* **2**, 1-55 (2021).
16. Lange, M. *et al.* CellRank for directed single-cell fate mapping. *Nat Methods* **19**, 159-170 (2022).

17. Roweis, S.T. & Saul, L.K. Nonlinear Dimensionality Reduction by Locally Linear Embedding. *Science* **290**,
2323-2326 (2000).
18. Whitney, H. The Self-Intersections of a Smooth n-Manifold in 2n-Space. *Annals of Mathematics* **45**, 220-
246 (1944).
19. Cannoodt, R., Saelens, W., Deconinck, L. & Saeys, Y. Spearheading future omics analyses using dyngen, a
multi-modal simulator of single cells. *Nat Commun* **12**, 3942 (2021).
20. Pijuan-Sala, B. *et al.* A single-cell molecular map of mouse gastrulation and early organogenesis. *Nature*
566, 490-495 (2019).
21. Barile, M. *et al.* Coordinated changes in gene expression kinetics underlie both mouse and human erythroid
maturation. *Genome Biol* **22**, 197 (2021).
22. Qiao, C. & Huang, Y. Representation learning of RNA velocity reveals robust cell transitions. *Proc Natl*
Acad Sci U S A **118**(2021).
23. Dybedal, I., Larsen, S. & Jacobsen, S.E. IL-12 directly enhances in vitro murine erythropoiesis in
combination with IL-4 and stem cell factor. *J Immunol* **154**, 4950-5 (1995).
24. Setty, M. *et al.* Characterization of cell fate probabilities in single-cell data with Palantir. *Nat Biotechnol* **37**,
451-460 (2019).
25. Ning, W. *et al.* Blocking exosomal miRNA-153-3p derived from bone marrow mesenchymal stem cells
ameliorates hypoxia-induced myocardial and microvascular damage by targeting the ANGPT1-mediated
VEGF/PI3k/Akt/eNOS pathway. *Cell Signal* **77**, 109812 (2021).
26. Bost, P. *et al.* Host-Viral Infection Maps Reveal Signatures of Severe COVID-19 Patients. *Cell* **181**, 1475-
1488 e12 (2020).
27. Ratnasiri, K., Wilk, A.J., Lee, M.J., Khatri, P. & Blish, C.A. Single-cell RNA-seq methods to interrogate
virus-host interactions. *Semin Immunopathol* **45**, 71-89 (2023).
28. Costa, B. *et al.* Human cytomegalovirus exploits STING signaling and counteracts IFN/ISG induction to
facilitate infection of dendritic cells. *Nat Commun* **15**, 1745 (2024).
29. Hein, M.Y. & Weissman, J.S. Functional single-cell genomics of human cytomegalovirus infection. *Nat*
Biotechnol **40**, 391-401 (2022).
30. Alon, U. *An introduction to systems biology: design principles of biological circuits*, (Chapman & Hall/CRC,
2007).
31. Ball, C.B. *et al.* Human Cytomegalovirus IE2 Both Activates and Represses Initiation and Modulates
Elongation in a Context-Dependent Manner. *mBio* **13**, e00337-22 (2022).
32. Manska, S. & Rossetto, C.C. Identification of cellular proteins associated with human cytomegalovirus
(HCMV) DNA replication suggests novel cellular and viral interactions. *Virology* **566**, 26-41 (2022).
33. Gredmark-Russ, S. & Soderberg-Naucler, C. Dendritic cell biology in human cytomegalovirus infection and
the clinical consequences for host immunity and pathology. *Virulence* **3**, 621-34 (2012).
34. Paulus, C., Krauss, S. & Nevels, M. A human cytomegalovirus antagonist of type I IFN-dependent signal
transducer and activator of transcription signaling. *Proc Natl Acad Sci U S A* **103**, 3840-5 (2006).
35. Park, A. *et al.* HCMV-encoded US7 and US8 act as antagonists of innate immunity by distinctively targeting
TLR-signaling pathways. *Nat Commun* **10**, 4670 (2019).
36. Yang, D., de la Rosa, G., Tewary, P. & Oppenheim, J.J. Alarmins link neutrophils and dendritic cells. *Trends*
Immunol **30**, 531-7 (2009).
37. Bogdanow, B. *et al.* Human cytomegalovirus tegument protein pp150 acts as a cyclin A2-CDK-dependent
sensor of the host cell cycle and differentiation state. *Proc Natl Acad Sci U S A* **110**, 17510-5 (2013).
38. Luftig, M.A. Viruses and the DNA Damage Response: Activation and Antagonism. *Annu Rev Virol* **1**, 605-
25 (2014).
39. Ball, C.B. *et al.* Human Cytomegalovirus Infection Elicits Global Changes in Host Transcription by RNA
Polymerases I, II, and III. *Viruses* **14**(2022).
40. Bogdanow, B., Phan, Q.V. & Wiebusch, L. Emerging Mechanisms of G(1)/S Cell Cycle Control by Human
and Mouse Cytomegaloviruses. *mBio* **12**, e0293421 (2021).
41. Goodwin, C.M., Ciesla, J.H. & Munger, J. Who's Driving? Human Cytomegalovirus, Interferon, and
NFkappaB Signaling. *Viruses* **10**(2018).
42. Manandhar, T., Ho, G.T., Pump, W.C., Blasczyk, R. & Bade-Doeding, C. Battle between Host Immune
Cellular Responses and HCMV Immune Evasion. *Int J Mol Sci* **20**(2019).
43. Zeng, J. *et al.* Insights into the Transcriptome of Human Cytomegalovirus: A Comprehensive Review.
Viruses **15**(2023).

44.   Schommartz, T., Tang, J., Brost, R. & Brune, W. Differential Requirement of Human Cytomegalovirus
520   UL112-113 Protein Isoforms for Viral Replication. *J Virol* **91**(2017).
521   45.   Ma, S. *et al.* Chromatin Potential Identified by Shared Single-Cell Profiling of RNA and Chromatin. *Cell*
522   **183**, 1103-1116 e20 (2020).
523   46.   Joost, S. *et al.* The Molecular Anatomy of Mouse Skin during Hair Growth and Rest. *Cell Stem Cell* **26**, 441-
524   457 e7 (2020).
525   47.   Pethe, P., Noel, V.S. & Kale, V. Deterministic role of sonic hedgehog signalling pathway in specification of
526   hemogenic versus endocardiogenic endothelium from differentiated human embryonic stem cells. *Cells Dev*
527   **166**, 203685 (2021).
528   48.   St-Jacques, B. *et al.* Sonic hedgehog signaling is essential for hair development. *Curr Biol* **8**, 1058-68 (1998).
529   49.   Mahdessian, D. *et al.* Spatiotemporal dissection of the cell cycle with single-cell proteogenomics. *Nature*
530   **590**, 649-654 (2021).
531   50.   Trevino, A.E. *et al.* Chromatin and gene-regulatory dynamics of the developing human cerebral cortex at
532   single-cell resolution. *Cell* **184**, 5053-5069 e23 (2021).
533   51.   Wang, S. *et al.* The inconsistent regulation of HOXC13 on different keratins and the regulation mechanism
534   on HOXC13 in cashmere goat (Capra hircus). *BMC Genomics* **19**, 630 (2018).
535   52.   Peng, Y. *et al.* Detection of new pioneer transcription factors as cell-type-specific nucleosome binders. *Elife*
536   **12**(2024).
537   53.   Desanlis, I. *et al.* HOX13-dependent chromatin accessibility underlies the transition towards the digit
538   development program. *Nat Commun* **11**, 2491 (2020).
539   54.   Gillespie, D.T. The chemical Langevin Equation. *J. Chem. Phys.* **113**, 297-306 (2000).
540   55.   Zwanzig, R. Ensemble method in the theory of irreversibility. *J. Chem. Phys.* **33**, 1338-1341 (1960).
541   56.   Mori, H. Transport, collective motion, and Brownian motion. *Prog. Theor, Phys.* **33**, 423-455 (1965).
542   57.   Xing, J. & Kim, K.S. Application of the projection operator formalism to non-Hamiltonian dynamics. *Journal*
543   *of Chemical Physics* **134**(2011).
544   58.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e29 (2021).
545   59.   Hastie, T. & Tibshirani, R. *Generalized additive models*, (Chapman and Hall, London ; New York, 1990).
546   60.   Mitra, S. *et al.* Single-cell multi-ome regression models identify functional and disease-associated enhancers
547   and enable chromatin potential analysis. *Nat Genet* **56**, 627-636 (2024).

Methods

Dynamical systems theory formulation of cellular state transitions

Assume that a cell state can be represented by the cell volume V (or cellular compartment size) and the copy number of $L >> 1$ pairs of gene products ($\mathbf{n}_m$, $\mathbf{n}_p$), where m and p designate mRNA and protein, and the bold fonts indicate vectors. For simplicity here we only consider m and p. It is straightforward to generalize to finer cell state specifications, for example, with distinction of nuclear and cytosol localizations, posttranslational states of proteins, other species such as microRNAs, epigenetic states, etc.

The temporal evolution of the cell state is described by a set of chemical master equations. When the copy numbers of molecular species are not too small, and the chemical reactions are not strongly coupled, Gillespie showed that the chemical master equations can be approximated by a set of chemical Langevin equations[54]. With extrinsic noises also included, we assume the ansatz that the dynamics of cell state is described by a set of generic stochastic differential equations,

$$\frac{dx_m}{dt} = F(x_m, x_p) + \zeta(x_m, x_p, t),$$

$$\frac{dx_p}{dt} = G(x_m, x_p) + \eta(x_m, x_p, t), \tag{4}$$

where the L-dimensional vectors $\mathbf{x}_m$ and $\mathbf{x}_p$ are cellular concentrations of m and p, and the ζ and η are taken as white noises with zero mean.

The low-dimensional manifold assumption is central to machine learning approaches on data analyses. From dynamical systems theory perspective, after a transient time a multi-dimensional dynamical system often converges to a low-dimensional slow manifold. In practice such property has been exploited with techniques such as quasi-steady-state approximation, quasi-equilibrium approximation. For a rigorous formulation, assume that one can identify a set of variables $(\mathbf{z}, \mathbf{Z})$ with ($2L$-M) dimensional fast variables $\mathbf{z} = \mathbf{z}(x_m, x_p)$ and M-dimensional slower variables $\mathbf{Z} = \mathbf{Z}(x_m, x_p)$. Xing and Kim extended the celebrated Zwanzig-Mori projection[55,56] to a general dynamical system described by eq. 4[57]. The projection procedure results in a set of stochastic integral-differential equations of $\mathbf{Z}$ with colored noises, which are formally equivalent to eq. 4. Then if one assumes clear time scale separation between $\mathbf{z}$ and $\mathbf{Z}$, the equations reduce to a set of

577   Langevin equations with white noises, $\frac{dZ}{dt} = A\big(Z(x_m, x_p)\big) + \eta(Z, t)$, where (Z, t) are white

578   noises with zero mean. Through ensemble averaging over the vicinity of a given point Z, one has

$$v \equiv <\frac{dZ}{dt}>_Z = A\big(Z(x_m, x_p)\big). \qquad (5)$$

The equations define a M-dimensional manifold embedded in the (x_m, x_p) space. A scRNA-seq

data then measures the corresponding manifold projected to the transcriptomic subspace.

One should note that in practice the reported RNA velocity vector of a cell state i is typically

obtained through averaging the raw velocity vectors of cell states within its neighborhood $\mathcal{N}_i$ on

the manifold as a numerical approximation of the ensemble average, $v_{xi} =<\frac{dx_m}{dt}> \approx \sum_{j \in \mathcal{N}_i} v_{xj}$.

In this work, we used the k-nearest neighbor (kNN) algorithm to define the neighborhood in single

modality datasets. For multiomics data, the neighborhood of a cell state was defined using

weighted nearest neighbors (WNN)[58] in the composite cell state space.

Mathematical foundation on applying GraphVelo to multiomics datasets

Equation 3 in the main text applies to transformation between a manifold embedded in a state

space and in a subspace. According to the Whitney Embedding theorem[18], any smooth real M-

dimensional manifold can be embedded in a $2M$-dimensional real space provided that $M > 0$.

Consider a full set of genes versus a subset in a scRNAseq dataset, or a combined

scRNAseq/scATACseq multiomics dataset versus the scRNAseq subset. Assume that the full cell

state space has a dimensionality N, while a single cell data manifold is typically low-dimensional

with $M \ll N$. Then the Whitney Embedding theorem[18] suggests that with proper choice of the

subset the manifolds in the full space and the subspace are homeomorphic or at least piece-wise

homeomorphic (Fig. 1b, Supplemental Text IV), i.e., a one-to-one mapping exists between the two.

Then applying eq. 3 allows one to infer the velocity vectors for the full-space representation from

those of the subspace.

Reconstruction of extended dynamo vector field from multiomics data

Dynamo is a general framework of reconstructing dynamical models from scRNAseq data, and it

is straightforward to generalize to multi-modal data. The framework is based on specific

realizations of eq. 5, $v \equiv <\frac{dx}{dt}>_x = A(x)$, with state vector x being the transcript concentrations

for scRNAseq data, and combined transcript concentrations and continuous quantification of
locus-specific chromatin open-close state for multiomics scRNAseq/scATACseq data. The
variables $\boldsymbol{x}$ can be defined in various representations, e.g., the original gene space, principal
component subspace, latent space defined by variational autoencoder, etc. With GraphVelo it is
straightforward to transform $\boldsymbol{v_x}$ between different representations.

The vector field functions $\boldsymbol{A(x)}$ contain quantitative regulation relations between genes that are
learned from single cell data points of $(\boldsymbol{x}, \boldsymbol{v_x})$. Various algorithms can be used to learn the
analytical forms of $\boldsymbol{A(x)}$. The original dynamo paper illustrated a Reproducing Kernel Hilbert
Space (RKHS) representation method. The method expresses $\boldsymbol{A(x)}$ as a linear combination of pre-
selected basis functions, $\boldsymbol{v} = \boldsymbol{A(x)} = \sum_\alpha \boldsymbol{c_\alpha} \boldsymbol{\Gamma_\alpha(x)}$, similar to the more familiar Taylor expansion
that uses a linear combination of polynomial functions to represent a continuous analytical function.
It should be noted that the basis functions and so $\boldsymbol{A(x)}$ are generally *nonlinear* functions of $\boldsymbol{x}$.
Following Qiu et al.[7], we chose Gaussian functions centered at selected reference points $\widetilde{\boldsymbol{x}}_\alpha$,
$\boldsymbol{\Gamma_a(x, \widetilde{x}_\alpha)} = e^{-2w\|x - \widetilde{x}_\alpha\|^2}$, with default parameter value of w in the package dynamo. Then we
determined the coefficient vectors $\boldsymbol{c_\alpha}$ through minimizing the loss function $\Phi(\boldsymbol{c_1}, \boldsymbol{c_2}, \dots, \boldsymbol{c_m}) =$
$\sum_{i=1}^n \|\boldsymbol{v_i} - \sum_\alpha \Gamma(\boldsymbol{x}, \widetilde{\boldsymbol{x}}_\alpha)\boldsymbol{c_\alpha}\|^2 + \frac{\lambda}{2}\sum_{\alpha=1}^m \sum_{\beta=1}^m \boldsymbol{c_\alpha^T}\Gamma(\widetilde{\boldsymbol{x}}_\alpha, \widetilde{\boldsymbol{x}}_\beta)\boldsymbol{c_\beta}$, where the first sum was over all the
data points, and the second term was Tikhonov regularization weighted by λ. The superscript T
means matrix transpose. One can also use neural networks, e.g., variational autoencoders, to learn
an optimal set of basis functions, and other algorithms such as neural ODE to learn $\boldsymbol{B(x)}$. The
difference is merely algorithmic under the same framework of dynamical systems theories.

Denoise velocity vectors in the space of principal components (PCs). Learning coefficients $\boldsymbol{\phi}_i$
in gene space directly often fails due to thousands of gene profiles. To avoid the curse of high
dimensionality and learn parameters in a compact manifold, we designed a procedure to denoise
the velocities in a reduced PCA space. Specifically, we extrapolated the cell state i in the original
space using the infinitesimal propagation operator to extrapolate the future state:

$$\boldsymbol{x'_i} = \boldsymbol{x_i} + \boldsymbol{v_i} \cdot dt_i$$

Moreover, we estimated an optimal step size dt based on the local density to guarantee the cell
states are bound to the manifold:

$$dt_i = \text{median}\left(\frac{\frac{1}{k}\sum_{j=1}^{k}\|\boldsymbol{\delta}_{ij}\|}{\|\boldsymbol{v}_i\|}\right)$$

633 After utilizing the cell-dependent t_i to forcing the predictions inhabiting regions of the
634 phenotypical manifold, we applied dimension reduction to project both current and future status
635 from the gene space to the PCA space through linear transformation. Then we obtained the
636 projected velocity vectors as:

637
$$\boldsymbol{v}_i^{\text{PCA}} = \frac{\boldsymbol{x}_i'Q - \boldsymbol{x}_iQ}{dt_i}$$

where Q is the PC loading matrix estimated using x that serves as the coordinate transformation
matrix.

Manifold-consistent kinetic genes. The kinetic assumptions between nascent and mature RNA
fail when the underlying parameters shift along the developmental trajectory[11], which leads to
transcription burst and rapid degradation in the phase portraits (Fig. 3a). An internal clock exists
during cell proliferation and differentiation. Current methods rely on different criteria to select
confident estimated velocity genes (see Supplemental Text V for detailed discussion).. Here, we
presume that the velocity of robustly estimated genes should be consistent with the (pseudo)time
derivative estimated under the manifold assumption. We can utilize any available ts inferred from
data manifold by either pseudotime, velocity latent time or lineage tracing to approximate the
temporal information and use k nearest neighbors to define the locally linear plane. After ordering
cells within the local Euclidean space, we calculate the MacK score for any gene g as an indicator
of whether the sign of estimated velocity agrees with the dynamic cascades within manifold,

651
$$\text{MacK score} = \frac{1}{n}\sum_{j\in\mathcal{N}(i)}\mathbb{I}\left(sgn\left(\frac{\Delta x_{ij}(g)}{\Delta t_{ij}}\right) = sgn(v_i(g))\right)$$

Where $\mathcal{N}_i$ indicates the neighbor points of cell i, $\mathbb{I}$ represents the indicator function and sgn returns
the sign of the values. $\Delta x_{ij}(g)$, $v_i(g)$ are the difference in abundance of gene g between cell i and
j, and the velocity of gene g in cell i, respectively. We parallelize the calculation to scale efficiently
with the number of genes, which is important due to the number of highly variable genes.

Dynamo criteria. Dynamo offers a correction strategy by removing genes with low gene-wise
confidence in the phase plane. This allows us to identify genes that appear in incorrect phase
portrait positions and contribute to erroneous flow directions (illustrated in Fig3. a). To filter out
genes with misleading dynamic patterns, a gene's behavior during cell differentiation from
progenitor to terminal states is evaluated based on its expression pattern relative to steady state.

Post-GraphVelo analyses

Markovian analyses. Marius et al.[16] have developed a framework named CellRank to study
cellular dynamics based on Markov chain formulation. We use CellRank to identify cell state
transitions using velocity kernel and identify terminal states within datasets by GPCCA function
module. In addition, CellRank pseudotime kernel is used for methods comparison in real datasets.

Functional form vector field and differential geometry analyses. Dynamo first proposed to
learn nonlinear functional form of RNA velocity vector field, providing a physical-informed
framework which integrates mechanism modeling and single cell data analyses. We use dynamo
to learn continuous vector filed and perform differential geometry analyses such as gene
acceleration, vector field-based pseudotime, LAP, Jacobian analyses and in silico perturbation.

Pseudotemporal orderings. GraphVelo itself does not compute an ordering index of cells as we
are seeking for a more quantitative method to infer RNA velocity. With an accurate RNA velocity
as input, we can approximate the vector field precisely. Thus, we use the scalar potential estimated
from the functional form vector field with Hodge decomposition as a proxy of time, which is
implemented by dynamo package.

Approximating smooth velocity trends with generalized addictive model. The variation of
transcription rates contains the high order dynamic information of the cell system. To model the
dynamic patterns of RNA velocity along transition path from noisy data, we refine the velocity
vectors by local geometry via TSP and further fit GAM to velocity value of each gene that has
been refined by GraphVelo. For any gene g, we model the velocity trend for the temporal variable
t via

682
$$v_{gi} = \beta_0 + f(t_i)$$

Where v_{gi} indicates the velocity of gene g in cell i, f is built using penalized B-splines which allow

us to automatically model non-linear mapping while maintaining additivity[59]. To visualize the

velocity trends, we select 100 equally spaced testing points along transition path and predict gene

expression at each of them using the fitted model. The estimated velocity trends can be treated as

smoothed time series for further analyses.

Clustering velocity trends along infection trajectory. With manifold-constrained velocity

estimated by GraphVelo, we are able to cluster genes into different functional modules which are

involved in the same regulatory circuit. We recover transcription variation of both host and virus

factors along infection trajectory by fitting GAMs in temporal indicator, the percentage of viral

RNA. Next, we select 100 equally spaced time points and generate the GAM-smoothed velocity

trends. We compute a kNN graph and cluster the kNN graph using the leiden algorithm. We used

k = 15 for the velocity-trend kNN graph and the leiden algorithm with resolution parameter set to

0.3 to avoid over clustering the trends. For each recovered cluster, we compute its mean and

standard deviation (pointwise, for all generated points that were used for smoothing) and visualize

the smoothed trends per cluster.

Characterizing decoupling genes based on dynamic time warping distance between multi-

modality velocities. We perform the DTW distance calculation by dtaidistance package. To

eliminate the influence of scale in different modality, we maximum-normalize the chromatin/RNA

velocity to the same range of [0, 1]. Then we fit the velocity trends of both modalities along vector

field-based pseudotime to yield the smoothed velocity trends. We calculate the DTW distance

between velocity trends per gene. To distinguish the decoupling genes based on their multi-

modality velocities, we rank the genes based on the DTW distance and identify elbow point as

cutoff. Genes with distance metric larger than the cutoff are characterized as decoupling genes and

used for visualization and functional analysis.

Synthetic datasets

Generating two genes bifurcation process and mapping it to 3d. The bifurcation data (n=2,000

cells) for the toggle-switch system is simulated using the Gillespie algorithm. We use activation

and inhibition Hill functions to model the induction and suppression effects between the two genes:

$$711 \qquad \dot{x} = \frac{a_1 x^n}{S_1^n + x^n} + \frac{b_1 K_1^n}{K_1^n + y^n} - \gamma_1 x,$$

$$712 \qquad \dot{y} = \frac{a_2 y^n}{S_2^n + y^n} + \frac{b_2 K_2^n}{K_2^n + x^n} - \gamma_2 y,$$

713 We use the simulation backend implemented by dynamo with default parameters except the

714 timescale (reset $\tau = 1$) to generate the bifurcating process. We then map the synthetic dataset onto

715 a sphere (radius $r=70$) and yield the variable z as:

$$716 \qquad z = \sqrt{max(r^2 - x^2 - y^2, 0)}$$

717 Then we are able to calculate the correctly-scaled 3d vectors by infinitesimal propagation operator

718 with sufficient small step size ($dt=1$ in our case):

$$719 \qquad \dot{z} = \lim_{\Delta t \to 0} \frac{\sqrt{max(r^2 - (x + \dot{x}\Delta t)^2 - (y + \dot{y}\Delta t)^2, 0)} - z}{\Delta t}$$

Generating scRNA-seq synthetic data with dyngen. To generate high-dimensional single-cell

transcriptomic data in silico, we use a multi-modal simulation engine, dyngen, to account for

different developmental topologies. We constructe module networks to represent regulatory

cascades and feedback loops driving progressive changes in gene expression and influencing cell

fate decisions. We generate three datasets with 1,000 cells and 100 genes using the linear, cyclic,

and bifurcating loop backbones provided by dyngen, with all other parameters set to default values.

These datasets include simulated nascent and mature mRNA counts along with ground-truth RNA

velocities and known manifold structure.

Simulation of rapid degradation and transcription burst events on phase portrait. As for

genes with variable degradation rate, we present a minimal regulatory network with linear model

in which an external signal both inhibits transcription and promotes microRNA (miRNA). The

miRNA exerts a linear influence on the degradation rate of mRNA. We have miRNA's velocity as

$$732 \qquad \frac{dm}{dt} = \alpha_m - \gamma_m m$$

733 The nascent gene transcription rate and mature mRNA's degradation rate would change to

$$734 \qquad \alpha = \alpha_0 - k_\alpha t$$

735
$$\gamma = \gamma_0 + k_\gamma m$$

736 where α_0 and γ_0 represent the constant transcription rate and degradation rate without the effect

737 of miRNA, k_α and k_γ represent the magnitude of influence from miRNA, t indicates the

738 simulation time to mimic the cell-context change along trajectory.

739
740
$$\frac{du}{dt} = \alpha - \beta u = \alpha_0 - k_\alpha m - \beta u$$
$$\frac{ds}{dt} = \beta u - \gamma s = \beta u - (\gamma_0 + k_\gamma m)s$$

We set the initial condition to a steady state with $u_0 = \alpha_0/\beta$ and $s_0 = \alpha_0/\gamma_0$, while the miRNA

abundance $m_0 = 0$. We simulate u_t and s_t as the microRNA signal m_t gradually increases. The

aim is to evaluate whether the estimated RNA velocity consistently aligns in sign with the ground

truth RNA velocity.

To generate genes with transcription burst phase portrait, we set the initial condition to $u_0 = 0$,

$s_0 = 0$, together with γ as constant and α promotes to $\alpha' = 3\alpha$ when simulation reaches specific

time.

Processing of sequencing data

All sequencing data in this study are downloaded publicly (see details in the 'Data availability'

section).

Analyses of scRNA-seq datasets. For the erythroid lineage of the mouse gastrulation, we follow

the standard data pre-processing procedures implemented by scVelo and select 9,815 cells and

2,000 highly variable genes to construct the k-nearest neighbor graph using 30 nearest neighbors

for downstream calculation.

For the human bone marrow dataset, we follow the standard data pre-processing procedures

implemented by scVelo and selected 5,780 cells and 2,000 highly variable genes to construct the

k-nearest neighbor graph using 30 nearest neighbors for downstream calculation. To estimate the

variation of degradation rate γ along the differentiation lineages, we divide the cells into five

discrete time bins based on precomputed Palantir pseudotime. We then estimate cell-specific

degradation rates and visualized their distribution shifts along the hematopoiesis trajectory.

Analyses of HCMV dataset. We sample the cells from donor 1 to eliminate sample-specific variation and further filter out cells lacking immediate early *UL123* gene expression to focus on the viral infection trajectory. We adapt monocle preprocessing recipe implemented by dynamo and yield 1,454 cells and 2,000 highly variable genes for further analyses. The number of nearest neighbors were set to 30 as default.

We collect the pathway-related genes from MSigDB and perform the Jacobian analyses implemented by dynamo, using viral genes as regulators and host genes as effectors. We rank the regulation relationships based on the collections of Jacobian elements. We pick top 50 inhibited effectors of viral genome and select the common set between pathway genes and all the effectors for visualization.

In silico knock out experiments are performed via dynamo. We suppressed every single virus factors per time using *dynamo.pd.perturbation* function and calculated the change of total viral gene expression after perturbation.

Analyses of multi-modality datasets.

RNA velocity estimation for multi-modality datasets. In traditional scRNA-seq datasets, RNA velocity methods use smoothed spliced and unspliced RNA counts through nearest-neighbor pooling, based on the PCA space computed from transcripts alone. However, this approach is not suitable for multimodal scenarios, as it overlooks hidden variables by relying on a single modality. To construct a consistent manifold combining information from multi-modal genomics data, we utilized WNN as implemented in MultiVelo[13]. The WNN algorithm combines low-dimensional representations from RNA and ATAC omics data. Specifically, we use PCA results from scRNA-seq data and latent semantic indexing (LSI) from scATAC-seq as inputs. The nearest neighbors identified by WNN were then used to calculate the first moment, reducing noise in separate modalities and approximating a unified manifold for GraphVelo.

Mouse skin dataset preprocessing. The preprocessed SHARE-seq mouse skin dataset[45] is adopted directly from MultiVelo data resources. All the procedures are consistent with MultiVelo except we get the LSI representation processed by SCARlink[60]. We construct the WNN graph using 50 nearest neighbors for downstream calculation. We run scVelo with 'stochastic' mode to estimate the RNA velocity based on the WNN graph as we discussed above.

Human cortex dataset preprocessing. The preprocessed human cerebral cortex data is adopted
directly from MultiVelo data resources. All the procedures are consistent with MultiVelo except
we get the LSI representation processed by SCARlink. We construct the WNN graph using 50
nearest neighbors for downstream calculation. We run scVelo with 'stochastic' mode to estimate
the RNA velocity based on the WNN graph.

Data availability

All the sequencing raw data are publicly accessible. The mouse gastrulation subset to erythroid
lineage can be extracted using scVelo's CLI: *scvelo.datasets.gastrulation_erythroid()* or from the
original work under accession number E-MTAB-6967 of ArrayExpress. The human bone marrow
can be extracted using scVelo's CLI: *scvelo.datasets. bonemarrow()* or through the Human Cell
Atlas data portal at https://data.humancellatlas.org/explore/projects/091cf39b-01bc-42e5-9437-
f419a66c8a45. The HCMV infected moDC data can be accessed via Zenodo
https://doi.org/10.5281/zenodo. 10404879 (https://zenodo.org/records/10404879). The
preprocessed mouse skin development dataset can be accessed via
https://figshare.com/articles/dataset/Mouse_Hair_Follicle_RNA_Data/22575307 and
https://figshare.com/articles/dataset/Mouse_hair_follicle_ATAC_data/22575313. The
preprocessed human cortex dataset can be downloaded from
https://figshare.com/articles/dataset/Developing_Human_Cortex_RNA_Data/22575376 and
https://figshare.com/articles/dataset/Developing_Human_Cortex_ATAC_Data/22575370.

Code availability

Python package GraphVelo can be accessed from https://github.com/xing-lab-pitt/GraphVelo.
Reproducibility and tutorials can be found in https://github.com/xing-lab-
pitt/GraphVelo/notebook.

References

54.   Gillespie, D.T. The chemical Langevin Equation. *J. Chem. Phys.* **113**, 297-306 (2000).
55.   Zwanzig, R. Ensemble method in the theory of irreversibility. *J. Chem. Phys.* **33**, 1338-1341 (1960).
56.   Mori, H. Transport, collective motion, and Brownian motion. *Prog. Theor, Phys.* **33**, 423-455 (1965).

57.   Xing, J. & Kim, K.S. Application of the projection operator formalism to non-Hamiltonian dynamics. *Journal*
of Chemical Physics **134**(2011).
58.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e29 (2021).
59.   Hastie, T. & Tibshirani, R. *Generalized additive models*, (Chapman and Hall, London ; New York, 1990).
60.   Mitra, S. *et al.* Single-cell multi-ome regression models identify functional and disease-associated enhancers
and enable chromatin potential analysis. *Nat Genet* **56**, 627-636 (2024).

Author contributions

826   JX conceived and formulated the theoretical framework, YC, and JG performed tests on simulated

827   data, YC and JX analyzed scRNA-seq datasets, YC, YZ, JG, and KN developed the package

828   GraphVelo, YC and JX wrote the draft, and all authors edited the manuscript.

Acknowledgments

This work was partially supported by National Institute of General Medical Sciences

(1R01GM148525 to JX and 1R01GM139297 to IB), and National Science Foundation (2325149)

to JX.

This work was supported by the National Key Research and Development Program of China

(2023YFE0112300); National Natural Sciences Foundation of China (32261133526; 32270709;

32070677); the 151 talent project of Zhejiang Province (first level), the Science and Technology

Innovation Leading Scientist (2022R52035)

Figure 1. Refining RNA velocity by tangent space projection and transforming between
representations using GraphVelo.

(a) Workflow of RNA velocity-based analyses incoporating GraphVelo. Note GraphVelo takes
any form of RNA velocity (i.e., not just splicing-based velocity) as input, and the kNN
neighborhood is defined in the full state space (e.g., by both scRNAseq and scATACseq in
multiomics data).

(b) Schematic of tangent space projection and velocity transformation between homeomophic
manifolds. *Left*: RNA velocity vectors are projected onto the tangent space defined by the
discretized local manifold of neighborhood cell samples. *Right*: GraphVelo allows for
transformation of velocity vectors from a manifold embeded in a higher dimensional space ($\mathcal{M}$)
to that in a lower dimensional space ($\aleph$), and *vice versa*.

(c) The process of minimizing the loss function of tangent space projection. Noisy velocity vectors
(*left*) generated by adding random components orthogonal to those sampled from an analytical 2D
manifold were projected back onto the 2D manifold, resulting in smooth velocity vectors that lie
in the tangent space (*right*).

(d) GraphVelo allows whole genome velocity inference based on the robustly estimated MacK
genes (see also Figure 3). Velocities of genes undergoing variable kinetic rates, such as rapid
degradation or transcription burst, are difficult to be correctly inferred by other methods, but can
be inferred robustly with GraphVelo.

(e) Virus infection dynamics and underlying host-virus interaction mechanisms uncovered by
GraphVelo (see also Figure 4). *Upper*: pathways involved in host-virus interactions were identified
using GraphVelo. *Lower*: GraphVelo predicted reversed trajectory of viral infection in response to
in silico perturbations of viral factors.

(f) GraphVelo provides a consistent view of epigenetic and transcription dynamics (see also Figure
5). *Upper*: GraphVelo analyses on multiomics data revealed that most cell-cycle dependent genes
showed decoupling between transcription dynamics and chromatin accessibility change dynamics.
Lower: Effective dose-response curves reconstructed from multiomics data revealed pioneer
transcription factors increased chromatin accessibility then transcription of targe genes.

Figure 2. Testing graphVelo on simulated datasets.

(a) Velocity vectors of an analytical three variables bifurcating vector field constrained to a spherical surface. The data points were colored by simulation time.

(b) Violinplots of: i) normal component of velocity vectors, ii) cosine similarity and iii) root mean

square error (RMSE) between ground truth and velocity vectors projected by GraphVelo and

cosine kernel, respectively.

(c-e) Simulation of scRNA-seq data using dyngen under linear, cycling, and bifurcating

differentiation models (*left*), and velocity fields (*right*) reconstructed using GraphVelo-corrected

velocities, respectively. Each simulation consists of 1,000 cell states and 100 genes. The cells in

different states were colored by their simulation time along trajectory.

(f-h) Boxplots of cosine similarity, , and accuracy between the ground truth velocity vectors and

dyngen simulated velocities after projection using GraphVelo TSP loss without cosine

regularization, GraphVelo TSP loss with cosine regularization, cosine kernel, and random

predictor, respectively.

$$\text{MacK score} = \frac{1}{n}\sum_{j \in \mathscr{N}(i)} \mathbb{I}(sgn(\frac{\Delta x_{i,j}(g)}{\Delta t_j}) = sgn(v_i(g)))$$

Figure 3. Delineating transcriptome-wise progression with manifold-consistent kinetic genes using GraphVelo

(a) Schematitc of transcritional events mislead RNA velocity estimation in the phase portrait by standard approaches. *Left*: for genes exhibiting rapid degradation, the cells appear above the steady state line on the phase portrait, whereas the true velocity is negative. *Right*: For genes exhibiting transcription burst, the transcription rate abruptly increases at intermediate states, leading to a steady state line whose slope is overestimated.

(b) Schematic of manifold-consistent score calculation for robustly estimated velocity genes.

(c) The projected velocity field from GraphVelo are consistent with the erythroid differentiation by using all highly variable genes.

(d) The correlation between GraphVelo vector field-based pseudotime and embryo time for erythroid lineage cells. Spearman correlation coefficients are shown.

(e) GO enrichment analyses of top ranked MacK genes.

(f) Scatter plots of: i) phase portrait, ii) velocities estimated by scVelo, iii) refined velocities by GraphVelo, and iv) mature mRNA expression of transcription burst genes (*Smim1*, *Hba-x*). Cells were colored by cell type, corresponding velocity, and mature mRNA abundance, respectively, and visualized on the phase portrait and UMAP representation.

(g) Cross boundary correctness calculated by the velocity input from GraphVelo, CellRank pseudotime kernel, VeloVI and scVelo, respectively.

(h) Gene regulatory cascade unraveled by GraphVelo-based vector field analyses that drives cell lineage commitment.

(i) Activation of *Gata1* inhibitor TF *Spi1* lead to reversed velocity flows in gastrulation erythroid maturation investigated through in silico perturbation analyses on GraphVelo-based vector field.

(j) Velocities derived from GraphVelo for the branching lineage in the hematopoiesis development and projected onto a pre-defined TSNE embedding. Directions of the projected cell velocities on TSNE are in agreement with the reported differentiation directions.

(k) Terminal states identified by CellRank based on Markov chain formulation derived from GraphVelo velocities.

(l) Phase portrait, velocity estimated by scVelo, refined velocity by GraphVelo, and gene
expression of mature mRNA of identified rapid degradation genes (*NPR3, ANGPT1*). The cells
were colored by the palantir pseudotime[24] in the phase portrait. The box plots showed cell-specific
γ for cells divided into bins according to pseudotime ordering in the phase protrait.

Figure 4. Using GraphVelo velocities to infer host-virus infection trajectory and identify

host-pathogen interactions.

(a) Viral infection captured by the GraphVelo velocity field. Cells were colored by the percentage of viral RNA within a single cell.

(b) Correlation between viral RNA percentage and pseudotime inferred by scVelo or GraphVelo. Spearman correlation coefficients and P values were shown.

(c) Viral RNA velocities infered by GraphVelo along the viral RNA percentage axis. The black dot line highlights the zero velocity.

(d) Boxplot summarizing the MacK scores of all viral genes calculated by GraphVelo, CellRank pseudotime kernel and random predictor.

(e) Correlation between viral infection speed and RNA abundance. Genes were ranked by Spearman correlatioin coefficients. Host and viral genes that contribute to viral DNA synthesis were marked in the left side and those contribute to viral defense response were marked in the right side. Viral genes were highlighted in red.

(f) UMAP representation of host and viral genes with distances defined by their dynamic expression patterns along the viral RNA percentage axis.

(g) Example dynamic expression patterns within specific clusters (Leiden4, 5, 3, 6 from top to bottom) along the viral RNA percentage axis. Zero velocity was highlighted by black dot line.

(h) GO enrichment of each cluster in (g).

(i) Top host genes inhibited by each viral factor based on dynamo Jacobian analyses. Host effectors were organized by their involved pathways.

(j) Dynamo prediction of total viral RNA change in response to in silico viral factor knockout. Viral factors were ranked by the mean of total viral RNA changes.

(k) Vector field change resultant from infinitesmal inhibition of *UL122* during the viral infection process.

Figure 5. Inferring epigenome and transcriptome consistent dynamics in mouse hair follicle

development using GraphVelo multiomics velocities.

(a) GraphVelo velocity fields of mouse hair follicle development. Cells were colored by cell macrostates.

(b) Number of terminal states predicted by CellRank using velocities inferred with different methods.

(c) Driver genes along multiple lineages identified through CellRank.

(d) Topological analyses of GraphVelo vector field identified novel root cells and attractors residing in three terminal states(IRS, hair shaft-cuticle cortex, and medulla).

(e) Expression levels of marker genes in novel root cells and expected root cells. Markers identified by Ma et al.[45] were highlighted with stars and newly identified markers were highlighted in bold.

(f) Regression results of MSD values along the transition path from the expected root or novel root to IRS. Two genes *Runx1* and *Shh* genes with large MSD originating from the novel root were highlighted.

(g) DTW distance between RNA velocity and chromatin velocity of individual genes. CCD genes were colored in red. The dotted line indicates the elbow point separating the decoupled genes from the rest.

(h) GO enrichment of decoupled genes in (g).

(i) Line plot of nomarlized RNA and chromatin velocity along pseudotime for genes predicted by GraphVelo to have notable decoupling patterns. Chromatin velocity trends were colored as brown and RNA velocity trends were colored as green.

(j) Heatmaps of Jacobian element distribution along the axis of regulator RNA abundance of four regulator effector circuits: i) *Lef1* versus *Wnt3* chromatin accessibilities. ii) *Hoxc13* versus *Wnt3* chromatin accessibilities. iii) *Lef1* versus *Wnt3* transcription. iv) *Hoxc13* versus *Wnt3* transcription.

(k) Effective dose-response curves obtained from integrating the averaged Jacobian elements over the corresponding normalized regulator mature mRNA regulator level in (j).

Extended Data Figures:

Extended Data Fig. 1. Refining RNA velocity from noisy simulation data or tradition splicing-based methods.

(a-c) Evaluation of on simulated scRNA-seq data under linear, cycling and bifurcating
differentiation models with an increasing noise level in velocity vectors.

(d) Heatmap of the correlation of cell speed calculated as the norm of velocity vector with respect
to the full-dimensional RNA velocity and the norm of velocity vectors projected to PCA or UMAP
space using GraphVelo, GraphVelo with cosine regularization, cosine kernel.

(e) Boxplots of metric evaluations on GraphVelo correction to the original velocities estimated by
scVelo, dynamo, and VeloVI. Box colors indicate the statistical test results for improvement
relative to zero.

Extended Data Fig. 2. RNA velocity inference on simulation results of splicing kinetics with constant or changed rates.

(a) Standard splicing kinetics along an induction trajectory in phase portrait and the reactions define how abundance levels of molecules change along simulation time. Right panel shows

corresponding trajectories over time (same below).

(b) Transcription burst along an induction trajectory in phase portrait. The transcription rate
constant α was promoted at specific time point.

(c) Standard splicing kinetics along a repression trajectory.

(d) Rapid time-varying degradation kinetics along repression trajectory. External signal promotes
synthesis of microRNA, which enhances degradation of target mRNA resulting in a microRNA -
dependent varying degradation rate "constant" γ, and inhibits the target gene via a decreasing α.

Extended Data Fig. 3. GraphVelo extrapolation of RNA velocities of an extended list of gene

set in mouse erythroid maturation dataset.

(a) Phase portrait, velocity estimated by scVelo, refined velocity by GraphVelo, and gene expression of mature mRNA of a selected set of genes. Cells were colored by cell type, corresponding velocity, and mature mRNA abundance, respectively, and visualized on the phase portrait and UMAP, respectively.

(b) Velocities of MURK genes derived from GraphVelo and scVelo for gastrulation erythroid maturation cells projected to a predefined UMAP representation. Note that RNA velocity estimated by scVelo was in a reverse flow, possibly influenced by transcription burst events.

(c) Cell speed distribution along the vector field pseudotime axis. Cells were colored by local density and red line indicates the fitted curve.

(d) Receiver operating curve analyses of MURK gene prediction based on dynamo accelerations analysis using GraphVelo as input, in contrast to a random predictor. AUC, area under curve.

1009

Extended Data Fig. 4. Regulatory cascades of driver TFs in mouse erythroid data.

(a) Expression and corresponding velocities of *Gata2*, *Gata1*, *Klf1* along the vector field-based

pseudotime in mouse erythroid data.

(b) Gene expression of *Gata2*, *Gata1*, *Klf1* on the UMAP space.

(c) Molecular mechanisms of driver TFs underlying erythroid lineage commitment based on
Jacobian analyses. (i) *Gata2* activates *Gata1*. (ii) Repression of *Gata2* by *Gata1*. (iii) Self-
activation of *Gata1*. (iv) *Gata1* activates *Klf1*.

(d) In silico perturbation analyses on GraphVelo-based vector field to examine the role of *Gata1*
in gastrulation erythroid maturation.

1019

Extended Data Fig. 5. RNA velocity estimated by scVelo and refined by GraphVelo from a

hematopoiesis dataset.

(a) Velocities derived from scVelo in the hematopoiesis development and projected to a pre-

defined TSNE embedding.

(b) TSNE visualization of the RNA velocity estimated by scVelo of *ANGPT1* and *RBPMS* genes.

(c) Scatter plots of: i) phase portrait, ii) velocities estimated by scVelo, iii) refined velocities by

GraphVelo, and iv) mature mRNA expression of transcription burst genes (e.g. *CALR*, *KLF1* and

AHSP). Cells were colored by cell type, corresponding velocity, and mature mRNA abundance,

respectively, and visualized on the phase portrait and TSNE, respectively.

(d) Cell-specific transcription rate constant α and degradation rate constant γ of rapid degradation

genes *ANGPT1* and *RBPMS* visualized on the TSNE space, which is consistent with simulation

result in Extended Data Fig. 2d.

1032

Extended Data Fig. 6. Dynamic patterns of gene trajectories during HCMV infection.

(a) Gene expression of selected viral factors on the UMAP space.

(b) Corresponding velocities inferred by GraphVelo for viral factors in (a).

(c) Viral RNA velocities derived from GraphVelo for infected cells, projected to the UMAP
representation only using viral genes.

(d) Receiver operating curve (ROC) analyses of MacK scores when using all detected viral genes
as the gold standard.

(e) Viral infection speed versus viral load. Cells were colored by the percentage of viral RNA.

(f) GO enrichment analyses of top host genes correlated with infection speed.

(g) Shifts in cell proportions across different cell cycle phases along the viral phase trajectory.
Cells were divided into bins according to viral RNA percentage. The result indicates a decreasing
S phase population corresponding to infection-driven transcriptional alterations of the cell cycle.
1045

1046

Extended Data Fig. 7. GraphVelo estimation of modality-consistent vector field based on multi-omics velocities.

(a) Velocity field projected to the pre-defined UMAP representation with different methods or gene sets: i) scVelo RNA velocity based on all velocity genes; ii) scVelo RNA velocity based on confident velocity genes filterred by dynamo criteria; iii) MultiVelo RNA velocity based on all velocity genes; iv) CellRank pseudotime kernel based on Palantir pseudotime.

(b) Terminal states identified by CellRank using different representation and corresponding velocity: i) GraphVelo-corrected RNA velocity; ii) GraphVelo-computed chromatin velocity; iii) MultiVelo-computed RNA velocity; iv) MultiVelo-computed chromatin velocity; v) scVelo-computed RNA velocity; vi) Palantir-based pseudotime kernel.

1057

a

i) scVelo   ii) Confident genes

Hair Shaft
-cuticle.cortex
IRS
Medulla
TAC-1
TAC-2

iii) MultiVelo   iv) CellRank (Palantir-based)

b   **Terminal states identified by CellRank**

i) GraphVelo (RNA)   ii) GraphVelo (ATAC)

IRS   IRS

Hair Shaft-cuticle.cortex   Hair Shaft-cuticle.cortex
Medulla   Medulla

iii) MultiVelo (RNA)   iv) MultiVelo (ATAC)

Medulla 2

TAC-1

Hair Shaft-cuticle.cortex
Medulla 1

v) scVelo   vi) Palantir

TAC-2

TAC-1

Hair Shaft-cuticle.cortex
Medulla   Medulla

1059

Extended Data Fig. 8. Reconstructed lineage commitment during mouse hair follicle differentiation using GraphVelo velocities.

(a) Lineage commitment probablity of each terminal cell type on UMAP vector field.

(b-d) Gene expression distribution of markers for cuticle/cortex, medulla and IRS lineages on the UMAP vector field.

1065

Extended Data Fig. 9. Novel root identified with multi-omic vector field from GraphVelo velocities.

(a) *Dcn*, *Mt2* expression dynamics during anagen hair follicle keratinocytes[46].

(b) Regions identified by topological analysis. Insert are ummarized cell-state transition vectors
calculated by GraphVelo and MultiVelo along the path from the novel root to IRS and projected
onto the UMAP representation.

(c) Predicted developmental LAPs from expected root or novel root to to each of the terminal cell
types in the UMAP embedding. Color of the node along the paths indicates the LAP transition
time.

(d) TF expression profiles along the LAP from novel root to IRS. *Shh* decays, alongside the
induction of the *Runx1* gene.

1077

1078 **Extended Data Fig. 10. Genomic patterns of decoupled CCD genes.**

1079 **(a)** Chromatin activity of decoupled CCD genes.

1080 **(b)** Gene expression of decoupled CCD genes.

1081

Extended Data Fig. 11. GraphVelo inferrence on epigenome and transcriptome decoupling dynamics in fetal human brain.

(a) GraphVelo velocity field colored by cell macrostates.

(b) DTW distance between RNA velocity and chromatin velocity of individual genes as a measure of the coupling/decoupling status. CCD genes were colored in red. The dotted line indicates the elbow point, with the decoupled genes on its right.

(c) GO enrichment of decoupled genes in (c).

(d) Line plot of nomarlized RNA and chromatin velocity along dynamo vector field pseudotime for genes predicted by GraphVelo to have notable decoupling patterns. Chromatin velocity trends were colored in brown and RNA velocity trends were colored in green.

(e) Chromatin activity of decoupled CCD genes.

(f) Gene expression of decoupled CCD genes.

1094

Extended Data Fig. 12. GraphVelo prediction on the regulatory mechanisms of *Lef1-Hoxc13-Wnt3* circuit during mouse skin development.

(a) Distributions of *Lef1* expression, *Hoxc13* expression, *Wnt3* chromatin openess/accessibility and *Wnt3* expression, respectively, on the projected UMAP vector field.

(b) Velocities of *Lef1* RNA, *Hoxc13* RNA, *Wnt3* chromatin and *Wnt3* RNA inferred by GraphVelo, visualized on the projected UMAP vector field.

(c) Velocities of *Lef1* RNA, *Hoxc13* RNA, *Wnt3* chromatin and *Wnt3* RNA inferred by MultiVelo,

visualized on the projected UMAP vector field.

(d) Expression of *Hoxc13* gene versus *Lef1* expression. *Lef1* expression primes the activation of

Hoxc13.

(e) Cell-wise Jacobian analyses of *Lef1-Hoxc13* activation cascade.

(f) Jacobian analyses of regulatory interactions between potential PTF *Hoxc13* and i) *Wnt3*

chromatin accessibility or ii) transcription..

(g) Jacobian analyses of regulatory interactions between PTF *Lef1* and i) *Wnt3* chromatin

accessibility or ii) transcription.

Supplemental text

I.   Connection of the projection to local linear embedding (LLE)

In this work, we use the neighborhood of a data point to approximate the corresponding tangent

space of the data manifold. Such approximation has been used in other applications such as the

local linear embedding (LLE)[1]. Here we provide a derivation of representing velocity vectors

based on LLE.

Consider a data point $\mathbf{x}_i$ and its neighbors. Applying LLE one has $\sum_{j \neq i} w_{ij}(x_j - x_i) =$

$\sum_{j \neq i} w_{ij} \delta_{ij} = 0$, with $\sum_j w_{ij} = 1$. The LLE algorithm gives $w_{ij} = \frac{\sum_k C_{jk}^{-1}}{\sum_{lm} C_{lm}^{-1}}$, with $C_{ij} = (x_i - x_j) \cdot$

$(x_i - x_k)$. Now consider after $\delta t \to 0$ $\mathbf{x}_i$ moves to $\mathbf{x}'_i = \mathbf{x}_i + \boldsymbol{v}_{pi} \delta t$, which should be also within

the local linear embedding space. Again with LLE one has $\sum_{j \neq i} w'_{ij}(x_j - x'_i) = 0$, with

$\sum_j w'_{ij} = 1$. Then one has $\boldsymbol{v}_{pi} = \sum_{j \neq i} \frac{(w'_{ij} - w_{ij})}{\delta t} \delta_{ij} = \sum_{j \neq i} \phi_{ij} \delta_{ij}$, with a form similar to that in eq.

1 of the main text.

The above derivation suggests an iterative trajectory propagation algorithm for obtaining the

velocities in the new representation:

1) Use a trial $\boldsymbol{v}_{pi}$ to propagate the state and obtain $(\mathbf{x}_i, \mathbf{x}'_i)$. One can propagate backward and

forward by $\delta t / 2$ and use central difference to estimate the velocity vectors.

2) Apply LLE to get w_{ij} and w'_{ij}. I

3) Obtain a new $\boldsymbol{v}_{pi}$

4) Go back to step 1 until the result converges below a threshold.

We numerically tested the above algorithm, the one with the loss function given in eq. 2 of the

main text with the parameter $b = 0$ and $b \neq 0$. All the three algorithms worked well on synthetic

data. The first two actually outperformed the third one (used in the main text). However, the first

two sometime showed numerical instability once applied to real single cell data. Therefore, in

subsequent applications, we used the general loss function form in eq. 2, using the direction

information from the second term (with $b \neq 0$) to further regularize the projection.

II.   Numerical issues of using eq. 1 for tangent space projection

The relation in eq. 1 provides an algorithm for projecting a measured **v** onto $\mathcal{M}$ by minimizing the

following loss function,

$$\mathcal{L}(\boldsymbol{\phi}_i) = \left\| \boldsymbol{v}_i - \boldsymbol{v}_{\|i} \right\|^2 + \lambda \|\boldsymbol{\phi}_i\|^2,$$

which unfortunately is numerically unstable. The redundancy of the basis vectors leads to

coefficients that are not uniquely determined, and failure of the projection. To see the latter, notice

that in real data the subspace formed by the displacement vectors only approximates the tangent

space $T\mathcal{M}$ locally, and it likely contains small components in the orthogonal space. Then the

projection procedure tries to express both $\mathbf{v}_{\|}(\mathbf{x})$ and all or part of the remaining $(\mathbf{v}_i - \mathbf{v}_{\|})$ as a linear

combination of $\boldsymbol{\delta}_{ij}$ with some $|\phi_{ij}| \gg 0$, which can only be counterbalanced by a very large value

of λ. The latter would impose large weight on the regularization. While further data preprocessing

may be developed in the future to alleviate the problem, in this work we provided a strategy of

adding an additional cosine kernel term for the direction information.

We also experimented with the idea of removing the redundancy through dimension reduction

with PCA. The numerical results were not satisfactory, since it converged slowly with the sampling

size, which becomes impractical.

III. Reformulation of Cosine kernel used in the literature in the context of tangent space projection

In the original RNA velocity study, a Cosine kernel has been proposed to transform the RNA velocity vectors between different representations[2]. We re-casted the cosine kernel in the context of the tangent space projection with a form similar to that of eq. 1 in the main text:

$$\mathbf{v}_{\parallel}^{corr}(\mathbf{x}_i) = \sum_{j \in \mathcal{N}_i} \phi_{ij}^{corr} \boldsymbol{\delta}_{ij},$$

where $\phi_{ij}^{corr} = P_{ij} - \frac{1}{k}$, and $P_{ij} = \frac{\exp[\cos(\mathbf{v}_i, \boldsymbol{\delta}_{ij})/\sigma]}{\sum_m \exp[\cos(\mathbf{v}_i, \boldsymbol{\delta}_{im})/\sigma]}$ defined in the form of softmax functions, with $\cos(\cdot, \cdot)$ denoting the cosine similarity between two input vectors, σ an arbitrary bandwidth parameter, and k the number of neighbors for each cell. Here P_{ij} gives a heuristic transition probability from cell i to j. The term $(-1/k)$, called the "density correction", is designed to correct the potential sampling bias where the embedded velocity vectors tend to point towards the direction of regions with high cell density. Li et al. showed that mathematically the cosine kernel asymptotically converges to the correct direction of a velocity vector[3]. However, the correlation kernel loses information about the magnitude of the velocity vectors $\mathbf{v}_i$ (i.e., the *speed*), due to the normalization in the correlation functions. Intuitively, the correlation kernel is qualitatively guided by the physical intuition that a cell has a high tendency to move along the direction of its velocity vector. Here, we used the direction information of ϕ_{ij}^{corr} to help on constraining $\boldsymbol{\phi}_i$.

IV. Further discussions on applications of GraphVelo to manifolds formed by multi-modal single cell data.

GraphVelo is based on the following mathematical assumption: the manifolds of a given system embedded in two different spaces are homeomorphic so one can establish a one-to-one mapping between the two. Here we use one simple example of scRNAseq/scATACseq multiomic data to illustrate that one can still apply GraphVelo if this assumption is violated. Consider that a cellular system switches between two discrete epigenetic states, and for each epigenetic state there is a corresponding (quasi)continuous transcriptomic manifold. Label the two disjoint manifolds as $\mathcal{M}_1(\mathbf{x})$ and $\mathcal{M}_2(\mathbf{x})$, where $\mathbf{x}$ represents the transcriptomic state. Consider two cells having the same $\mathbf{x}$ but different chromatin state, c_1 on $\mathcal{M}_1$ and c_2 on $\mathcal{M}_2$. Multiomic data allows distinction between two cells. That is, the neighborhood of c_1 is composed of cells on $\mathcal{M}_1$, and the neighborhood of c_2 is composed of cells on $\mathcal{M}_2$. Therefore, GraphVelo analyses treat cells on $\mathcal{M}_1$ and $\mathcal{M}_2$ separately.

V.   **Feature selection for robustly estimated velocity genes.**

In datasets featuring non-differentiating cell types organized in a hierarchical manner, application of RNA velocity may show different kinetic regimes. Analyzing cell fate transition using unreliable velocity genes can lead to emergence of spurious cell state transitions. Different packages subset the well-estimated kinetic genes for further analyses in different scenarios. In the implementation of scVelo, one first regresses out the uncertain genes are based on the r-squared coefficient in the steady-state model and further selects trustful velocity genes according to the log-likelihood in the dynamical mode. PhyloVelo relies on monotonically expressed genes to infer transcriptomic velocity, which is a more stringent requirement. However, the monotonic assumptions are counter intuitive and oversimplify the gene expression kinetics along differentiation or disease trajectories due to the non-sequential nature of these cascades. Dynamo

provides a biological prior-guided method and filters out the unreliable genes based on the

knowledge of lineage information.

Reference

1. Roweis, S.T. & Saul, L.K. Nonlinear Dimensionality Reduction by Locally Linear Embedding. *Science* **290**, 2323-2326 (2000).
2. La Manno, G. *et al*. RNA velocity of single cells. *Nature* (2018).
3. Li, T., Shi, J., Wu, Y. & Zhou, P. On the Mathematics of RNA Velocity I: Theoretical Analysis. *CSIAM Transactions on Applied Mathematics* **2**, 1-55 (2021).