Comparing Intra- and Inter-individual Correlational Brain Connectivity from Functional and

Structural Neuroimaging Data

Xin Di[1*], Bharat B. Biswal [1*], Alzheimer's Disease Neuroimaging Initiative[†]

1. Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, NJ, 07102, USA

* Corresponding authors:

Xin Di, Ph.D.

604 Fenster Hall, University Height

Newark, NJ, 07102, USA

xin.di@njit.edu

Bharat B. Biswal, Ph.D.

607 Fenster Hall, University Height

Newark, NJ, 07102, USA

bbiswal@yahoo.com

† Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Abstract

Inferring brain connectivity from inter-individual correlations has been applied to various neuroimaging modalities, such as glucose metabolic activity measured by positron emission tomography (PET) and brain structures assessed using MRI. The variability that drives these inter-individual correlations is generally attributed to individual differences, potentially influenced by factors like genetics, life experiences, and biological sex. However, it remains unclear whether long-term within-individual effects, such as aging, and state-like effects also contribute to the correlated structures, and how intra-individual correlations are compared to inter-individual correlations. In this study, we analyzed longitudinal data spanning a wide age range, examining regional brain volumes using structural MRI, and regional brain functions using both regional homogeneity (ReHo) of resting-state functional MRI and glucose metabolic activity measured with Fludeoxyglucose (18F) FDG-PET. In a first dataset from a single individual scanned over 15 years, we found that intra-individual correlations in both ReHo and regional volumes resembled resting-state functional connectivity. In a second dataset, involving multiple longitudinal points and participants for FDG-PET and MRI, we replicated these findings, showing that both intra- and inter-individual correlations were strongly associated with resting-state functional connectivity. Correlations in functional measures (i.e., ReHo or FDG-PET) showed greater similarity with resting-state connectivity than structural measures. Moreover, matrices from the same modality showed higher similarity between the two datasets, indicating modality specific contributions. These results suggest that multiple factors may contribute to both inter- and intra-individual correlational measures of connectivity. Understanding or controlling for these factors could enhance the interpretability of the inter-individual connectivity measures.

Keywords: Brain connectivity; Covariance network; Functional Connectivity; Inter-individual; Molecular connectivity.

1. Backgrounds

Studies of brain connectivity are essential for advancing our understanding of functional interactions between brain regions and the organization of the whole brain. The development of neuroimaging techniques has provided an exciting opportunity to study brain function in humans in vivo. Early research frequently employed positron emission tomography (PET) to measure glucose metabolic activity (Phelps et al., 1981) and cerebral blood flow (Fox and Raichle, 1984). These studies primarily used inter-individual correlations of PET measures to quantify brain connectivity based on glucose metabolism (Horwitz et al., 1984; Metter et al., 1984) or cerebral blood flow (Zeki et al., 1991). However, due to the nature of PET measurements, which are static, these studies were generally limited to inter-individual correlations. While they often identified statistically significant connectivity patterns, the similarities between connectivity derived from PET measures and resting-state networks identified using functional MRI (fMRI) were relatively modest (Di et al., 2017; Di and Biswal, and Alzheimer's Disease Neu, 2012; Lizarraga et al., 2023).

Functional MRI (fMRI) has become a widely used tool for studying brain connectivity due to its superior spatial and temporal resolution (Biswal et al., 1995, 2010). Beyond capturing moment-to-moment dynamics, fMRI data can be summarized over brief periods, often during resting-state sessions, to derive measures such as the amplitude of low-frequency fluctuations (ALFF) (Zang et al., 2007) and regional homogeneity (ReHo) (Zang et al., 2004). These metrics have also been applied to examine inter-individual correlations of brain (Di et al., 2024a; Taylor et al., 2012; Zhang et al., 2011). Additionally, the flexibility of task designs in fMRI enables researchers to explore how task performance impacts inter-individual connectivity correlations. Studies indicate that while task conditions can induce slight changes in connectivity patterns, the overall connectivity structure tends to remain largely consistent across different tasks (Di et al., 2024a).

An intriguing extension of inter-individual correlation analysis is its application to brain structural data, which tends to reflect more trait-like characteristics associated with slow and long-term effects (He et al., 2007; Mechelli et al., 2005). Mechelli and colleagues were among the first to use a seed-based

approach to examine inter-individual correlations of regional brain volumes, discovering strong

correlations between regions within the same functional brain systems (Mechelli et al., 2005). Building on

this, He and colleagues constructed whole-brain networks based on inter-individual correlations of

cortical thickness. Their findings demonstrated that these structural networks exhibit small-world

properties, highlighting the efficiency and organization of the brain's structural connectivity (He et al.,

2007).

Despite its growing popularity, questions remain about whether and to what extent inter-

individual correlations reflect functional connectivity, which is traditionally assessed intra-individually,

typically through resting-state fMRI. One approach to validate inter-individual correlational measures is

to compare their similarity to other established connectivity measures, such as intra-subject moment-to-

moment functional connectivity during rest or anatomical connectivity derived from white matter

tracking. When using white matter tracking from diffusion-weighted imaging (DWI) as a reference,

studies have found that inter-individual correlations of structural measures show limited similarity to

white matter tracts (Gong et al., 2012; Lizarraga et al., 2023). In contrast, inter-individual correlations

based on functional measures of glucose metabolic activity exhibit higher similarity with white matter

connectivity (Lizarraga et al., 2023). A similar pattern emerges when comparing these measures to

resting-state functional connectivity. Inter-individual structural correlations display limited similarity to

resting-state functional connectivity (Alexander-Bloch et al., 2013b; Di et al., 2017). However, inter-

individual correlations of functional measures, such as glucose metabolic activity, show greater alignment

with resting-state connectivity patterns (Di et al., 2017).

A critical question remains regarding the factors driving inter-individual variability that lead to

correlations in functional or structural brain measures. Do these correlations primarily reflect individual

differences shaped by genetic factors or life experiences, or do intra-individual factors also play a role?

For example, inter-individual correlation analyses often include large sample sizes spanning wide age

ranges, prompting the question of whether age-related effects contribute significantly to these correlations

in brain structure. Exploring intra-individual correlations could provide valuable insights into the

underlying causes of inter-individual variability.

In the context of functional data, such as glucose metabolism measured by FDG-PET, neural

activity introduces an additional variable. This state-like factor may be influenced by participants' mental

states at the time of measurement (Di et al., 2024a). Long-term brain activity, as reflected by metrics like

FDG-PET or regional homogeneity (ReHo), typically persists over minutes to hours. However, it is

unclear to what extent variability in this sustained activity contributes to the observed inter-individual

correlations. Investigating intra-individual correlations in these slow functional activity patterns could

shed light on the role of intra-individual variability in shaping inter-individual correlations.

In the current study, we examined correlations in brain structural and functional measures

typically calculated in an inter-individual manner. We analyzed two unique datasets, allowing us to

compute correlations both inter- and intra-individually, and compared the correlation structures derived

from these two approaches. This comparison enabled us to estimate the contribution of intra-individual

factors to the overall correlation structure. Specifically, the first dataset consists of a single individual

scanned over 16 years (Duchesne et al., 2019), providing a unique opportunity to estimate gray matter and

ReHo correlations intra-individually. The second dataset comes from the Alzheimer's Disease

Neuroimaging Initiative (ADNI), focusing on healthy participants with more than five longitudinal FDG-

PET scans. We calculated correlations in two ways: first, by calculating correlation matrices within each

participant and then averaging these matrices across participants, which minimizes individual variability

and focuses on intra-individual variability; and second, by calculating inter-individual correlations at each

age point and averaging these matrices across ages, which focuses exclusively on inter-individual

variability while controlling for factors such as age. Lastly, we compared correlation matrices between the

two datasets and investigate whether different imaging modalities have their unique correlation structures.

2. Materials and Methods

2.1. Datasets

2.1.1. Simon dataset

The Simon dataset is available through the International Neuroimaging Data-Sharing Initiative (INDI)

website (http://fcon_1000.projects.nitrc.org/indi/retro/SIMON.html). It includes data from a single heathy

male who was scanned across 73 sessions over a 16-year period, from the age of 30 to 47. Figure 1A

illustrates the distribution of these scanning sessions over time. In total, 73 MRI sessions are available,

conducted using various scanners and parameters. For more details, refer to the original paper by

(Duchesne et al., 2019). Our analysis focused on T1-weighted anatomical images and resting-state fMRI

data, with 71 sessions providing T1-weighted images and 58 sessions containing resting-state fMRI data.

Figure 1 Illustration of scan sessions for the Simon dataset (A) and fludeoxyglucose-18 (FDG) positron

emission tomography (PET) (B) and structural MRI (C) datasets from Alzheimer's Disease Neuroimaging

Initiative (ADNI). For the Simon dataset, a single participant was scanned for 73 sessions over 16 years.

Each dot represents one session. For the ADNI dataset, each row in y axis represents one participant,

where each participant was scanned for multiple sessions.

2.1.2. ADNI dataset

The ADNI dataset was obtained from the project website (adni.loni.usc.edu). The ADNI was launched in

2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner, MD. The primary

goal of ADNI has been to test whether serial magnetic resonance imaging (MRI), positron emission

tomography (PET), other biological markers, and clinical and neuropsychological assessment can be

combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease

(AD). For up-to-date information, see www.adni-info.org.

In this analysis, we only included data from healthy participants. All selected individuals showed

no evidence of depression, mild cognitive impairment (MCI), or dementia, as indicated by Mini-Mental

State Examination (MMSE) scores ranging from 24 to 30 and a Clinical Dementia Rating (CDR) of 0.

We manually curated FDG-PET, MRI, and resting-state fMRI data for this study. For FDG-PET and

MRI, we included participants with at least five sessions to ensure the calculation of reliable intra-

individual correlations. For the resting-state fMRI data, we included only one session per individual and

focused on the averaged correlation matrix across participants.

The data search began with FDG-PET, resulting in the inclusion of 72 participants (25 females)

with a total of 432 PET scan sessions. The number of sessions per participant ranged from 5 to 9 (Figure

1B). The participants' average age at the first session was 75.8 years, with a range of 62 to 86 years. For

each session, either a mean image was calculated, or a single representative image was used.

From the 72 participants with qualified PET data, we identified those with at least five sessions of

structural MRI scans. A total of 65 participants met this criterion, with the number of MRI sessions

ranging from 5 to 13. Figure 1C provides an overview of the session count and corresponding ages for

these participants.

Finally, among the 65 participants with structural MRI data, 17 individuals also had resting-state

fMRI scans available. For these participants, one session per individual was included, focusing on a

single resting-state fMRI session per participant.

2.2. Data processing

Neuroimaging data processing and analysis were conducted using SPM12

(https://www.fil.ion.ucl.ac.uk/spm/) within MATLAB environment, following preprocessing and quality

control procedures detailed in a prior study (Di and Biswal, 2023).

For the FDG-PET data, dynamic images (i.e., multiple images per session) were processed by

realigning all images within a session to the first image, followed by generating a mean image for that

session. For static PET data, which contained only a single image per session, no realignment was

required. Next, the mean images (or static images) from all sessions for each participant were realigned to

the image from the first session. A cross-sessional mean image was then normalized directly to the PET

template in SPM, aligned to the standard Montreal Neurological Institute (MNI) space. Normalization to

MNI space was performed using consistent parameters across all images. Direct normalization was

chosen over MRI-mediated normalization due to the sufficient spatial resolution of PET images and the

methodological advantages of direct normalization (Calhoun et al., 2017). Finally, the normalized images

were spatially smoothed using an 8 mm full-width at half-maximum (FWHM) Gaussian kernel, and each

image was normalized by dividing its signal by the mean signal within an intracranial volume mask.

Each anatomical image was treated as independent data, segmented into gray matter, white

matter, cerebrospinal fluid, and other tissues, and normalized to standard MNI space. Spatial

normalization included modulation to ensure that the resulting gray matter images reflected gray matter

volume (GMV). Quality control was performed by manually inspecting the anatomical images before and

after segmentation. In the Simon dataset, one session was excluded due to segmentation failure, resulting

in a total of 70 sessions being included in the final analysis.

For the fMRI data, the functional images were first realigned to the first image of each session.

The mean functional image was then coregistered to the corresponding anatomical image. Next, the

functional images were normalized to MNI space using the deformation field maps derived from the

segmentation step and spatially smoothed with an 8 mm full-width at half-maximum (FWHM) Gaussian

kernel. A voxel-wise general linear model was then applied to regress out head motion parameters and

white matter/CSF signals. This model included 24 regressors based on Friston's head motion model,

along with the first five principal components of signals from the white matter and cerebrospinal fluid.

The residual images from this step were saved for further analysis.

For the Simon dataset, ReHo was calculated for each resting-state fMRI session using the REST

toolbox (Song et al., 2011).

2.3. Brain parcellation

Cortical regions were defined using Schaefer's 100-region parcellation (Schaefer et al., 2018), while

subcortical regions were identified based on the Automated Anatomical Labeling (AAL) atlas. The

subcortical regions included the bilateral hippocampus, parahippocampus, amygdala, caudate, putamen,

pallidum, and thalamus (Tzourio-Mazoyer et al., 2002).

For each participant and region, voxel values were averaged to compute measures of gray matter volume

(GMV), FDG-PET, and regional homogeneity (ReHo), producing a 114-dimensional vector per

participant. For resting-state fMRI data, the average time series for each region of interest (ROI) was

extracted, resulting in a $114 \times n$ matrix, where n represents the number of time points, which varied

across sessions and participants.

2.4. Calculation of intra- and inter-individual correlation matrices

In the Simon dataset, mean GMV values across 114 regions of interest (ROIs) for the 70 sessions were

arranged into a 114×70 matrix. Pearson's correlation coefficients were then calculated to construct a

within-individual GMV correlation matrix (114×114). Similarly, a ReHo correlation matrix was

generated using ReHo maps from 58 sessions. For each of these 58 resting-state sessions, a resting-state

connectivity matrix was also computed from the fMRI time series data. Finally, the correlation matrices

from all sessions were averaged to produce a mean correlation matrix.

For the ADNI dataset, correlation matrices for FDG-PET or GMV were calculated across

sessions for each participant. These matrices were then averaged across participants to produce a mean

correlation matrix, referred to as the intra-individual correlation matrix. To calculate inter-individual

correlations, participants' ages were controlled. Specifically, inter-individual correlations were computed

at each integer age point where data from more than nine participants were available. This process was

applied to participants aged between 70 and 89 years, and the resulting inter-individual correlation

matrices were averaged to generate a mean correlation matrix, referred to as the inter-individual

correlation matrix. Finally, for the fMRI data, resting-state functional connectivity matrices were

calculated for each participant. These matrices were averaged across participants to obtain a mean

connectivity matrix.

2.5. Statistical analysis

To investigate the associations among the correlation matrices, we extracted the upper diagonal of each

matrix and converted it into a 6,441-dimensional vector (114 x ($114 - 1$) / 2). Given the potential non-

Gaussian distribution of the correlation matrices, Spearman's rank correlation coefficient (ρ) was used to

quantify the relationships among the matrices.

3. Results

3.1. Simon dataset

We first analyzed a multi-session dataset from a single individual spanning over 16 years. The averaged

correlation matrix, shown in Figure 2A, reveals clear modular structures, evident as square-like patterns

along the diagonal and additional squares representing left-right homotopic networks. Subsequently, we

computed intra-individual correlations for regional brain volume (GMV) and ReHo across all available

sessions. Both matrices display square-like patterns, although their spatial configurations differ. Notably,

both intra-individual correlation matrices show moderate but significant correlations with the averaged

resting-state time series correlation matrix (Spearman's correlation coefficients: $\rho_{GMV} = 0.29$; $\rho_{ReHo} =$

0.39).

Figure 2 Correlations of regional brain volume (GMV) (A), regional homogeneity (ReHo) (B), and

averaged correlations of resting-state time series (C) across 114 regions of interest (ROIs). D and E show

the Spearman's correlations (ρ) between the time-series correlation matrix and GMV or ReHo

correlations, respectively.

3.2. ADNI dataset

Next, we analyzed the ADNI dataset, where there were multiple participants, but each participant only

have a few sessions. We calculated averaged resting-state time series correlation from the 17 participants

(Figure 3A), which turned out to be very similar to those from the Simon dataset. We then calculated

structural correlations both intra-individually and inter-individually. Both matrices demonstrated

functional network structures along the diagonal and between left and right corresponding regions.

Moreover, both correlation matrices were correlated with resting-state time series correlation ($\rho_{intra} = 0.41$;

$\rho_{inter} = 0.37$), which were slightly higher than that in the Simon dataset (0.29).

Figure 3 Correlations of regional brain volume (GMV) calculated intra-individually (A) and inter-

individually (B), and averaged correlations of resting-state time series (C) across 114 regions o interests

(ROIs). D and E show the Spearman's correlations (ρ) between the resting-state time series correlation

matrix and GMV correlation matrices, respectively.

We next calculated correlations of glucose metabolic activity intra-individually and inter-

individually, and correlated the correlation matrices with resting-state time series correlation (Figure 4).

The correlations matrices of glucose metabolic activity showed more obvious functional network

structures than those in structural correlations. And most importantly, the both correlation matrices

showed strong correlations with resting-state time series correlation (ρ_{intra} = 0.71; ρ_{inter} = 0.64).

Figure 4 Correlations of regional brain glucose metabolism measured using positron emission

tomography (PET) calculated intra-individually (A) and inter-individually (B), and averaged correlations

of resting-state time series (C) across. D and E show the Spearman's correlations (ρ) between the resting-

state time series correlation matrix with the two PET correlation matrices, respectiveely.

3.3. Relationships among all the matrices

Finally, we calculated the correlations among the correlation matrices (Figure 5A). Due to the large

number of ROI pairs, all correlations were statistically significant. However, it is more meaningful to

focus on the relative effects of these correlations rather than their statistical significance. Here, we

emphasize the correlations between the matrices of the two datasets, where the highest correlations were

observed between corresponding modalities (highlighted blue rectangle and diamond markers). For

instance, the strongest correlation with the mean time-series correlations in the Simon dataset was found

with the time-series correlations in the ADNI dataset ($\rho = 0.67$). Similarly, the highest correlation with

GMV correlations in the Simon dataset occurred with intra-individual GMV correlations in the ADNI

dataset ($\rho = 0.46$), rather than inter-individual correlations. Notably, the highest correlation with ReHo

correlations in the Simon dataset was observed with inter-individual PET correlations in the ADNI dataset

($\rho = 0.51$).

Figure 5 A. Spearman's correlation matrix among the correlation matrices. The blue rectangle and

diamond markers highlight the effects of interest. B. The first principal component of the correlation

matrices. C. Loadings of the correlation matrices on the first and second principal components. ADNI,

Alzheimer's Disease Neuroimaging Initiative; GMV, gray matter volume; PET, positron emission

288   tomography; ROI region of interest.

We then conducted a PCA on the eight correlation matrices, with the first principal component

(PC) accounting for 59.0% of the variance (Figure 5B). Next, we visualized the loadings of the eight

matrices on the first two PCs (Figure 5C). The matrices formed distinct clusters in the plot. For instance,

the GMV correlation matrices were located at the bottom, while the time-series and PET correlation

matrices clustered in the top-left corner.

4. Discussion

Using two unique datasets encompassing both intra- and inter-individual effects, the current analysis

demonstrated how these effects contribute to correlation structures across brain regions. First, long-term

structural brain changes revealed correlation patterns that were small but significantly associated with

resting-state time-series correlations. Second, long-term functional activity, as measured by ReHo or

glucose metabolism, exhibited stronger correlation structures and greater alignment with resting-state

time-series correlations compared to structural measures. Finally, correlation matrices from the two

datasets showed greater similarity within the same modality than between different modalities, suggesting

that each modality provides distinct insights into interregional relationships.

This study analyzed a unique dataset from a single individual scanned over 16 years, revealing

that intra-individual structural correlations were modestly associated with resting-state connectivity (ρ =

0.29), the lowest correlation observed among the analyses. This finding suggests that structural brain

development and aging within an individual are partially constrained by the brain's functional

organization. These results were further validated using the ADNI dataset, which, despite fewer

longitudinal data points, showed a stronger correlation between intra-individual structural correlations

and resting-state connectivity (ρ = 0.41). To our knowledge, this is the first study to demonstrate

structural correlations within an individual, underscoring the influence of age as a single factor shaping

these correlations.

Notably, inter-individual structural correlations, when controlling for chronological age, showed

a similar association with resting-state connectivity (ρ = 0.37), indicating that individual differences

contribute comparably. The variability that give rise to the inter-individual correlation may be related to

genetics, life experience, and plasticity (Alexander-Bloch et al., 2013a; Evans, 2013). In previous studies

of structural "covariance", these factors could not be distinguished from development and aging effects.

In fact, the current findings suggest that multiple factors could give rise to the correlation structure. This

is also in line with studies showing that the structural "covariance" were modulated by factors such as age

(Vijayakumar et al., 2021). To enhance interpretability, researchers should consider restricting or

controlling for such factors in their analyses.

To the best of our knowledge, this study is the first to demonstrate intra-individual correlations in

long-term brain activity using either ReHo or FDG-PET measures. Notably, the correlations of ReHo and

metabolic activity were generally higher than those of regional brain volumes. This suggests that

summary measures of brain function may capture state-dependent activity, such as mood, thoughts, or

other transient mental states during scanning (Di et al., 2024b). The data revealed that even within a

single individual, long-term correlations showed strong similarity with resting-state connectivity ($\rho =$

0.39).

However, it is important to note that neural activity measured across different temporal scales

may not reflect identical processes. Thus, it remains unclear whether correlation structures derived from

distinct temporal scales are directly comparable. Prior studies on fMRI time series have shown that

connectivity structures within a single scanning session can vary depending on the frequency bands

analyzed (Gohel and Biswal, 2015; Kajimura et al., 2023; Yuen et al., 2019). Future research could

explore the comparison of correlation structures between slow and fast neural activity patterns to better

understand their relationship.

Compared to earlier studies, the current research reports slightly higher correlations with resting-

state connectivity. This improvement may stem from larger sample sizes, averaging data across multiple

sessions, or advancements in the preprocessing pipeline. These findings imply that the smaller

correlations observed in previous studies might partially result from noise, and that improvements in data

acquisition and processing can enhance the observed similarities. Nonetheless, the analysis also highlights

that each modality makes a unique contribution to the correlation structure, indicating an inherent limit to

the similarities that can be achieved between different modalities.

This analysis validates the use of structural and functional brain measures to investigate

interregional relationships, often referred to as functional connectivity when using functional data. Given

the complex factors influencing these measures, controlling for certain variables can enhance the

interpretability of correlation results. Our findings demonstrated that intra-individual correlations, which

account for individual differences, tend to exhibit stronger associations with resting-state functional

connectivity compared to inter-individual correlations, supporting the value of such controls. However, it

is important to note that multiple measures from the same individual are not always available, and in

some cases, inter-individual correlations may be the only feasible approach.

5. Conclusion

The current results to some extent validated the usage of inter-individual correlations as an estimate of

brain connectivity. The results also highlighted that multiple factors could contribute to the correlation

structure. Those factors may need to control or taken care of to boost interpretability of the results.

Acknowledgement

This study was supported by grants from (US) National Institutes of Health for Xin Di (R15MH125332)
and Bharat B. Biswal (5R01MH131335).
Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging
Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of
Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the
National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from
the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon
Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.;
Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its
affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer
Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research &
Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx
Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging;
Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of
Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions
are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee
organization is the Northern California Institute for Research and Education, and the study is coordinated
by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data
are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.
378

380   **References:**

381   Alexander-Bloch, A., Giedd, J.N., Bullmore, E., 2013a. Imaging structural co-variance between human
382   brain regions. Nat Rev Neurosci 14, 322–336. https://doi.org/10.1038/nrn3465

Alexander-Bloch, A., Raznahan, A., Bullmore, E., Giedd, J., 2013b. The convergence of maturational change and structural covariance in human cortical networks. The Journal of neuroscience : the official journal of the Society for Neuroscience 33, 2889–99. https://doi.org/10.1523/JNEUROSCI.3554-12.2013

Biswal, B., Yetkin, F.Z., Haughton, V.M., Hyde, J.S., 1995. Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. Magnetic resonance in medicine : official journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine 34, 537–41. https://doi.org/10.1002/mrm.1910340409

Biswal, B.B., Mennes, M., Zuo, X.-N., Gohel, S., Kelly, C., Smith, S.M., Beckmann, C.F., Adelstein, J.S., Buckner, R.L., Colcombe, S., Dogonowski, A.-M., Ernst, M., Fair, D., Hampson, M., Hoptman, M.J., Hyde, J.S., Kiviniemi, V.J., Kötter, R., Li, S.-J., Lin, C.-P., Lowe, M.J., Mackay, C., Madden, D.J., Madsen, K.H., Margulies, D.S., Mayberg, H.S., McMahon, K., Monk, C.S., Mostofsky, S.H., Nagel, B.J., Pekar, J.J., Peltier, S.J., Petersen, S.E., Riedl, V., Rombouts, S.A.R.B., Rypma, B., Schlaggar, B.L., Schmidt, S., Seidler, R.D., Siegle, G.J., Sorg, C., Teng, G.-J., Veijola, J., Villringer, A., Walter, M., Wang, L., Weng, X.-C., Whitfield-Gabrieli, S., Williamson, P., Windischberger, C., Zang, Y.-F., Zhang, H.-Y., Castellanos, F.X., Milham, M.P., 2010. Toward discovery science of human brain function. Proceedings of the National Academy of Sciences of the United States of America 107, 4734–9. https://doi.org/10.1073/pnas.0911855107

Di, X., Biswal, and Alzheimer's Disease Neu, B.B., 2012. Metabolic Brain Covariant Networks as Revealed by FDG-PET with Reference to Resting-State fMRI Networks. Brain Connectivity 2, 275–283. https://doi.org/10.1089/brain.2012.0086

Di, X., Biswal, B.B., 2023. A functional MRI pre-processing and quality control protocol based on statistical parametric mapping (SPM) and MATLAB. Frontiers in Neuroimaging 1, 1070151.

Di, X., Gohel, S., Thielcke, A., Wehrl, H.F., Biswal, B.B., Alzheimer's Disease Neuroimaging Initiative, 2017. Do all roads lead to Rome? A comparison of brain networks derived from inter-subject volumetric and metabolic covariance and moment-to-moment hemodynamic correlations in old individuals. Brain structure & function 222, 3833–3845. https://doi.org/10.1007/s00429-017-1438-7

Di, X., Jain, P., Biswal, B.B., 2024a. Effects of Tasks on Functional Brain Connectivity Derived from Inter-Individual Correlations: Insights from Regional Homogeneity of Functional MRI Data. https://doi.org/10.1101/2024.06.02.597063

Di, X., Jain, P., Biswal, B.B., 2024b. Effects of Tasks on Functional Brain Connectivity Derived from Inter-Individual Correlations: Insights from Regional Homogeneity of Functional MRI Data. https://doi.org/10.1101/2024.06.02.597063

Duchesne, S., Dieumegarde, L., Chouinard, I., Farokhian, F., Badhwar, A., Bellec, P., Tétreault, P., Descoteaux, M., Boré, A., Houde, J.-C., Beaulieu, C., Potvin, O., 2019. Structural and functional multi-platform MRI series of a single human volunteer over more than fifteen years. Sci Data 6, 245. https://doi.org/10.1038/s41597-019-0262-8

Evans, A.C., 2013. Networks of anatomical covariance. NeuroImage 80, 489–504. https://doi.org/10.1016/j.neuroimage.2013.05.054

Fox, P.T., Raichle, M.E., 1984. Stimulus rate dependence of regional cerebral blood flow in human striate cortex, demonstrated by positron emission tomography. Journal of neurophysiology 51, 1109–20.

Gohel, S.R., Biswal, B.B., 2015. Functional Integration Between Brain Regions at Rest Occurs in Multiple-Frequency Bands. Brain Connectivity 5, 23–34. https://doi.org/10.1089/brain.2013.0210

Gong, G., He, Y., Chen, Z.J., Evans, A.C., 2012. Convergence and divergence of thickness correlations with diffusion connections across the human cerebral cortex. NeuroImage 59, 1239–1248. https://doi.org/10.1016/j.neuroimage.2011.08.017

He, Y., Chen, Z.J., Evans, A.C., 2007. Small-world anatomical networks in the human brain revealed by cortical thickness from MRI. Cerebral cortex (New York, N.Y. : 1991) 17, 2407–19. https://doi.org/10.1093/cercor/bhl149

Horwitz, B., Duara, R., Rapoport, S.I., 1984. Intercorrelations of Glucose Metabolic Rates Between Brain
Regions: Application to Healthy Males in a State of Reduced Sensory Input. Journal of Cerebral
Blood Flow & Metabolism 4, 484–499. https://doi.org/10.1038/jcbfm.1984.73
Kajimura, S., Margulies, D., Smallwood, J., 2023. Frequency-specific brain network architecture in
resting-state fMRI. Sci Rep 13, 2964. https://doi.org/10.1038/s41598-023-29321-5
Lizarraga, A., Ripp, I., Sala, A., Shi, K., Düring, M., Koch, K., Yakushev, I., 2023. Similarity between
structural and proxy estimates of brain connectivity. J Cereb Blood Flow Metab
0271678X231204769. https://doi.org/10.1177/0271678X231204769
Mechelli, A., Friston, K.J., Frackowiak, R.S., Price, C.J., 2005. Structural covariance in the human cortex.
The Journal of neuroscience : the official journal of the Society for Neuroscience 25, 8303–10.
https://doi.org/10.1523/JNEUROSCI.0357-05.2005
Metter, E.J., Riege, W.H., Kuhl, D.E., Phelps, M.E., 1984. Cerebral Metabolic Relationships for Selected
Brain Regions in Healthy Adults. Journal of Cerebral Blood Flow & Metabolism 4, 1–7.
https://doi.org/10.1038/jcbfm.1984.1
Phelps, M.E., Kuhl, D.E., Mazziota, J.C., 1981. Metabolic Mapping of the Brain's Response to Visual
Stimulation: Studies in Humans. Science 211, 1445–1448.
https://doi.org/10.1126/science.6970412
Schaefer, A., Kong, R., Gordon, E.M., Laumann, T.O., Zuo, X.-N., Holmes, A.J., Eickhoff, S.B., Yeo,
B.T.T., 2018. Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional
Connectivity MRI. Cerebral cortex (New York, N.Y. : 1991) 28, 3095–3114.
https://doi.org/10.1093/cercor/bhx179
Song, X.-W., Dong, Z.-Y., Long, X.-Y., Li, S.-F., Zuo, X.-N., Zhu, C.-Z., He, Y., Yan, C.-G., Zang, Y.-
F., 2011. REST: a toolkit for resting-state functional magnetic resonance imaging data
processing. PloS one 6, e25031. https://doi.org/10.1371/journal.pone.0025031
Taylor, P.A., Gohel, S., Di, X., Walter, M., Biswal, B.B., 2012. Functional covariance networks:
obtaining resting-state networks from intersubject variability. Brain connectivity 2, 203–17.
https://doi.org/10.1089/brain.2012.0095
Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B.,
Joliot, M., 2002. Automated anatomical labeling of activations in SPM using a macroscopic
anatomical parcellation of the MNI MRI single-subject brain. NeuroImage 15, 273–89.
https://doi.org/10.1006/nimg.2001.0978
Vijayakumar, N., Ball, G., Seal, M.L., Mundy, L., Whittle, S., Silk, T., 2021. The development of
structural covariance networks during the transition from childhood to adolescence. Sci Rep 11,
9451. https://doi.org/10.1038/s41598-021-88918-w
Yuen, N.H., Osachoff, N., Chen, J.J., 2019. Intrinsic Frequencies of the Resting-State fMRI Signal: The
Frequency Dependence of Functional Connectivity and the Effect of Mode Mixing. Front
Neurosci 13, 900. https://doi.org/10.3389/fnins.2019.00900
Zang, Y., Jiang, T., Lu, Y., He, Y., Tian, L., 2004. Regional homogeneity approach to fMRI data
analysis. NeuroImage 22, 394–400. https://doi.org/10.1016/j.neuroimage.2003.12.030
Zang, Y.-F., He, Y., Zhu, C.-Z., Cao, Q.-J., Sui, M.-Q., Liang, M., Tian, L.-X., Jiang, T.-Z., Wang, Y.-F.,
2007. Altered baseline brain activity in children with ADHD revealed by resting-state functional
MRI. Brain & development 29, 83–91. https://doi.org/10.1016/j.braindev.2006.07.002
Zeki, S., Watson, J., Lueck, C., Friston, K., Kennard, C., Frackowiak, R., 1991. A direct demonstration of
functional specialization in human visual cortex. J. Neurosci. 11, 641–649.
Zhang, Z., Liao, W., Zuo, X.-N., Wang, Z., Yuan, C., Jiao, Q., Chen, H., Biswal, B.B., Lu, G., Liu, Y.,
2011. Resting-state brain organization revealed by functional covariance networks. PloS one 6,
e28817. https://doi.org/10.1371/journal.pone.0028817
481
482