Comparing Intra- and Inter-individual Correlational Brain Connectivity from Functional and

Structural Neuroimaging Data

Xin Di[1*], Bharat B. Biswal [1*], Alzheimer's Disease Neuroimaging Initiative[†]

1. Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, NJ, 07102, USA

* Corresponding authors:

Xin Di, Ph.D.

604 Fenster Hall, University Height

Newark, NJ, 07102, USA

xin.di@njit.edu

Bharat B. Biswal, Ph.D.

607 Fenster Hall, University Height

Newark, NJ, 07102, USA

bbiswal@yahoo.com

† Part of data used in preparation of this article were obtained from the Alzheimer's Disease
Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI
contributed to the design and implementation of ADNI and/or provided data but did not participate in
analysis or writing of this report. A complete listing of ADNI investigators can be found at:
http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Abstract

Inferring brain connectivity from inter-individual correlations has been applied across various

neuroimaging modalities, including positron emission tomography (PET) and MRI. The variability

underlying these inter-individual correlations is generally attributed to factors such as genetics, life

experiences, and long-term influences like aging. This study leveraged two unique longitudinal datasets to

examine intra-individual correlations of structural and functional brain measures across an extended time

span. By focusing on intra-individual correlations, we aimed to minimize individual differences and

investigate how aging and state-like effects contribute to brain connectivity patterns. Additionally, we

compared intra-individual correlations with inter-individual correlations to better understand their

relationship. In the first dataset, which included repeated scans from a single individual over 15 years, we

found that intra-individual correlations in both regional homogeneity (ReHo) during resting-state and gray

matter volumes (GMV) from structural MRI closely resembled resting-state functional connectivity.

However, ReHo correlations were primarily driven by state-like variability, whereas GMV correlations

were mainly influenced by aging. The second dataset, comprising multiple participants with longitudinal

Fludeoxyglucose (18F) FDG-PET and MRI scans, replicated these findings. Both intra- and inter-

individual correlations were strongly associated with resting-state functional connectivity, with functional

measures (i.e., ReHo and FDG-PET) exhibiting greater similarity to resting-state connectivity than

structural measures. This study demonstrated that controlling for various factors can enhance the

interpretability of brain correlation structures. While inter- and intra-individual correlation patterns

showed similarities, accounting for additional variables may improve our understanding of inter-

individual connectivity measures.

Keywords: Brain connectivity; Covariance network; Functional Connectivity; Inter-individual; Molecular

connectivity.

1. Backgrounds

Studies of brain connectivity are essential for advancing our understanding of functional interactions between brain regions and the organization of the whole brain. The development of neuroimaging techniques has provided an exciting opportunity to study brain function in humans in vivo. Early research frequently employed positron emission tomography (PET) to measure glucose metabolic activity (Phelps et al., 1981) and cerebral blood flow (Fox and Raichle, 1984). These studies primarily used inter-individual correlations of PET measures to quantify brain connectivity based on glucose metabolism (Horwitz et al., 1984; Metter et al., 1984) or cerebral blood flow (Zeki et al., 1991). However, due to the nature of PET measurements, which are static, these studies were generally limited to inter-individual correlations. While they often identified statistically significant connectivity patterns, the similarities between connectivity derived from PET measures and resting-state networks identified using functional MRI (fMRI) were relatively modest (Di et al., 2017; Di and Biswal, and Alzheimer's Disease Neu, 2012; Lizarraga et al., 2023).

Functional MRI (fMRI) has become a widely used tool for studying brain connectivity due to its superior spatial and temporal resolution (Biswal et al., 1995, 2010). Beyond capturing moment-to-moment dynamics, fMRI data can be summarized over brief periods, often during resting-state runs, to derive measures such as the amplitude of low-frequency fluctuations (ALFF) (Zang et al., 2007) and regional homogeneity (ReHo) (Zang et al., 2004). These metrics have also been applied to examine inter-individual correlations of brain (Di et al., 2024; Taylor et al., 2012; Zhang et al., 2011). Additionally, the flexibility of task designs in fMRI enables researchers to explore how task performance impacts inter-individual connectivity correlations. Studies indicate that while task conditions can induce slight changes in connectivity patterns, the overall connectivity structure tends to remain largely consistent across different tasks (Di et al., 2024).

An intriguing extension of inter-individual correlation analysis is its application to brain structural data, which tends to reflect more trait-like characteristics associated with slow and long-term effects (He et al., 2007; Mechelli et al., 2005). Mechelli and colleagues were among the first to use a seed-based

approach to examine inter-individual correlations of regional brain volumes, discovering strong

correlations between regions within the same functional brain systems (Mechelli et al., 2005). Building on

this, He and colleagues constructed whole-brain networks based on inter-individual correlations of

cortical thickness. Their findings demonstrated that these structural networks exhibit small-world

properties, highlighting the efficiency and organization of the brain's structural connectivity (He et al.,

2007).

Despite its growing popularity, questions remain about whether and to what extent inter-

individual correlations reflect functional connectivity, which is traditionally assessed intra-individually,

typically through resting-state fMRI. One approach to validate inter-individual correlational measures is

to compare their similarity to other established connectivity measures, such as intra-subject moment-to-

moment functional connectivity during rest or anatomical connectivity derived from white matter

tracking. When using white matter tracking from diffusion-weighted imaging (DWI) as a reference,

studies have found that inter-individual correlations of structural measures show limited similarity to

white matter tracts (Gong et al., 2012; Lizarraga et al., 2023). In contrast, inter-individual correlations

based on functional measures of glucose metabolic activity exhibit higher similarity with white matter

connectivity (Lizarraga et al., 2023). A similar pattern emerges when comparing these measures to

resting-state functional connectivity. Inter-individual structural correlations display limited similarity to

resting-state functional connectivity (Alexander-Bloch et al., 2013b; Di et al., 2017). However, inter-

individual correlations of functional measures, such as glucose metabolic activity, show greater similarity

with resting-state connectivity patterns compared with structural measures (Di et al., 2017).

A critical question remains regarding the factors driving inter-individual variability that lead to

correlations in functional or structural brain measures. Do these correlations primarily reflect individual

differences shaped by genetic factors or life experiences, or do intra-individual factors also play a role?

For example, inter-individual correlation analyses often include large sample sizes spanning wide age

ranges, prompting the question of whether age-related effects contribute significantly to these correlations

in brain structure. Exploring intra-individual correlations could provide valuable insights into the

underlying causes of inter-individual variability.

In the context of functional data, such as glucose metabolism measured by FDG-PET, neural

activity introduces an additional variable. This state-like factor may be influenced by participants' mental

states at the time of measurement (Di et al., 2024). Long-term brain activity, as reflected by metrics like

FDG-PET or regional homogeneity (ReHo), typically persists over minutes to hours. However, it is

unclear to what extent variability in this sustained activity contributes to the observed inter-individual

correlations. By comparing intra-individual correlations in these slow-varying functional patterns with

those from structural measures, which do not exhibit short-term fluctuations, we can gain deeper insights

into the impact of state-like functional dynamics on inter-individual correlations.

In the current study, we examined correlations in brain structural and functional measures

typically calculated in an inter-individual manner. We analyzed two unique datasets, allowing us to

compute correlations both inter- and intra-individually, and compared the correlation structures derived

from these two approaches. This comparison enabled us to estimate the contribution of intra-individual

factors to the overall correlation structure. Specifically, the first dataset consists of a single individual

scanned over 16 years (Duchesne et al., 2019), providing a unique opportunity to estimate gray matter and

ReHo correlations intra-individually. The second dataset comes from the Alzheimer's Disease

Neuroimaging Initiative (ADNI), focusing on healthy participants with more than five longitudinal FDG-

PET scans. We calculated correlations in two ways: first, by calculating correlation matrices within each

participant and then averaging these matrices across participants, which minimizes individual variability

and focuses on intra-individual variability; and second, by calculating inter-individual correlations at each

age point and averaging these matrices across ages, which focuses exclusively on inter-individual

variability while controlling for factors such as age. Lastly, we compared correlation matrices between the

two datasets and investigate whether different imaging modalities have their unique correlation structures.

2. Materials and Methods

2.1. Datasets

2.1.1. Simon dataset

The Simon dataset is available through the International Neuroimaging Data-Sharing Initiative (INDI)

website (http://fcon_1000.projects.nitrc.org/indi/retro/SIMON.html). It includes data from a single heathy

male who was scanned across 73 sessions over a 16-year period, from the age of 30 to 47. Figure 1A

illustrates the distribution of these scanning sessions over time. In total, 73 MRI sessions are available,

conducted using various scanners and parameters. For more details, refer to the original paper by

(Duchesne et al., 2019). Our analysis focused on T1-weighted anatomical images and resting-state fMRI

data, with 71 sessions providing T1-weighted images and 58 sessions containing resting-state fMRI data.

Figure 1 Illustration of scan sessions for the Simon dataset (A) and fludeoxyglucose-18 (FDG) positron

emission tomography (PET) (B) and structural MRI (C) datasets from Alzheimer's Disease Neuroimaging

Initiative (ADNI). For the Simon dataset, a single participant was scanned for 73 sessions over 16 years.

Each dot represents one session. For the ADNI dataset, each row in y axis represents one participant,

where each participant was scanned for multiple sessions.

2.1.2. ADNI dataset

The ADNI dataset was obtained from the project website (adni.loni.usc.edu). The ADNI was launched in

2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner, MD. The primary

goal of ADNI has been to test whether serial magnetic resonance imaging (MRI), positron emission

tomography (PET), other biological markers, and clinical and neuropsychological assessment can be

combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease

(AD). For up-to-date information, see www.adni-info.org.

In this analysis, we only included data from healthy participants. All selected individuals showed

no evidence of depression, mild cognitive impairment (MCI), or dementia, as indicated by Mini-Mental

State Examination (MMSE) scores ranging from 24 to 30 and a Clinical Dementia Rating (CDR) of 0.

We manually curated FDG-PET, MRI, and resting-state fMRI data for this study. For FDG-PET and

MRI, we included participants with at least five sessions to ensure the calculation of reliable intra-

individual correlations. For the resting-state fMRI data, we included only one session per individual and

focused on the averaged correlation matrix across participants.

The data search began with FDG-PET, resulting in the inclusion of 72 participants (25 females)

with a total of 432 PET scan sessions. The number of sessions per participant ranged from 5 to 9 (Figure

1B). The participants' average age at the first session was 75.8 years, with a range of 62 to 86 years. For

each session, either a mean image was calculated, or a single representative image was used.

From the 72 participants with qualified PET data, we identified those with at least five sessions of

structural MRI scans. A total of 65 participants met this criterion, with the number of MRI sessions

ranging from 5 to 13. Figure 1C provides an overview of the session count and corresponding ages for

these participants.

Finally, among the 65 participants with structural MRI data, 17 individuals also had resting-state

fMRI scans available. For these participants, one session per individual was included, focusing on a

single resting-state fMRI session per participant. fMRI data typically lack a sufficient number of

longitudinal sessions, making it impossible to conduct inter- and intra-individual correlation analyses

comparable to those performed with anatomical MRI and PET data using ReHo from resting-state fMRI.

2.2. Data processing

Neuroimaging data processing and analysis were conducted using SPM12

(https://www.fil.ion.ucl.ac.uk/spm/) within MATLAB environment, following preprocessing and quality

control procedures detailed in a prior study (Di and Biswal, 2023).

For the FDG-PET data, dynamic images (i.e., multiple images per session) were processed by

realigning all images within a session to the first image, followed by generating a mean image for that

session. For static PET data, which contained only a single image per session, no realignment was

required. Next, the mean images (or static images) from all sessions for each participant were realigned to

the image from the first session. A cross-sessional mean image was then normalized directly to the PET

template in SPM, aligned to the standard Montreal Neurological Institute (MNI) space. Normalization to

MNI space was performed using consistent parameters across all images. Direct normalization was

chosen over MRI-mediated normalization due to the sufficient spatial resolution of PET images and the

methodological advantages of direct normalization (Calhoun et al., 2017). Finally, the normalized images

were spatially smoothed using an 8 mm full-width at half-maximum (FWHM) Gaussian kernel, and each

image was normalized by dividing its signal by the mean signal within an intracranial volume mask.

Each anatomical image was treated as independent data, segmented into gray matter, white

matter, cerebrospinal fluid, and other tissues, and normalized to standard MNI space. Spatial

normalization included modulation to ensure that the resulting gray matter images reflected gray matter

volume (GMV). Quality control was performed by manually inspecting the anatomical images before and

after segmentation. In the Simon dataset, one session was excluded due to segmentation failure, resulting

in a total of 70 sessions being included in the final analysis.

For the fMRI data, the functional images were first realigned to the first image of each session.

The mean functional image was then coregistered to the corresponding anatomical image. Next, the

functional images were normalized to MNI space using the deformation field maps derived from the

segmentation step and spatially smoothed with an 8 mm full-width at half-maximum (FWHM) Gaussian

kernel. A voxel-wise general linear model was then applied to regress out head motion parameters and

white matter/CSF signals. This model included 24 regressors based on Friston's head motion model,

along with the first five principal components of signals from the white matter and cerebrospinal fluid.

The residual images from this step were saved for further analysis.

For the Simon dataset, ReHo was calculated for each resting-state fMRI session using the REST

toolbox (Song et al., 2011).

2.3. Brain parcellation

Cortical regions were defined using Schaefer's 100-region parcellation (Schaefer et al., 2018), while

subcortical regions were identified based on the Automated Anatomical Labeling (AAL) atlas. The

cortical regions were further grouped into seven networks (Yeo et al., 2011) and organized by

hemisphere. The subcortical regions included the bilateral hippocampus, parahippocampus, amygdala,

caudate, putamen, pallidum, and thalamus (Tzourio-Mazoyer et al., 2002).

For each participant and region, voxel values were averaged to compute measures of gray matter

volume (GMV), FDG-PET, and regional homogeneity (ReHo), producing a 114-dimensional vector per

participant. For resting-state fMRI data, the average time series for each region of interest (ROI) was

extracted, resulting in a $114 \times n$ matrix, where n represents the number of time points, which varied

across sessions and participants.

2.4. Calculation of intra- and inter-individual correlation matrices

In the Simon dataset, mean GMV values across 114 regions of interest (ROIs) for the 70 sessions were

arranged into a 114×70 matrix. Pearson's correlation coefficients were then calculated to construct a

within-individual GMV correlation matrix (114×114). Similarly, a ReHo correlation matrix was

generated using ReHo maps from 58 sessions. For each of these 58 resting-state sessions, a resting-state

connectivity matrix was also computed from the fMRI time series data. Finally, the correlation matrices

from all sessions were averaged to produce a mean correlation matrix.

For the ADNI dataset, correlation matrices for FDG-PET or GMV were calculated across

sessions for each participant. These matrices were then averaged across participants to produce a mean

correlation matrix, referred to as the intra-individual correlation matrix. To calculate inter-individual

correlations, participants' ages were controlled. Specifically, inter-individual correlations were computed

at each integer age point where data from more than nine participants were available. This process was

applied to participants aged between 70 and 89 years, and the resulting inter-individual correlation

matrices were averaged to generate a mean correlation matrix, referred to as the inter-individual

correlation matrix. Finally, for the fMRI data, resting-state functional connectivity matrices were

calculated for each participant. These matrices were averaged across participants to obtain a mean

connectivity matrix.

2.5. Fitting age models for the Simon dataset

For each region, we fitted the mean GMV or ReHo value series across sessions using cubic age models.

We therefore obtained fitted age effects series and residual series across sessions. We calculated

correlations of the fitted age effects and residual effects among the 114 ROIs, resulting in 114 x 114

matrices.

2.6. Similarities among correlation matrices

To investigate the associations among the correlation matrices, we extracted the upper diagonal of each

matrix and converted it into a 6,441-dimensional vector (114 x (114 - 1) / 2). Given the potential non-

Gaussian distribution of the correlation matrices, Spearman's rank correlation coefficient (ρ) was used to

quantify the relationships among the matrices.

Next, we examined the similarities between the correlation matrices from the Simon and ADNI

datasets. The Simon dataset included the averaged resting-state correlation matrix, GMV correlation

matrix, and ReHo correlation matrices, while the ADNI dataset contained the averaged resting-state

correlation matrix, intra- and inter-individual correlations of GMV, and intra- and inter-individual

correlations of PET glucose metabolic activity. To analyze these relationships, we extracted the upper

diagonal of the eight matrices, forming a $6,441 \times 8$ matrix. We first computed Spearman's correlation

coefficients among the eight matrices, followed by a principal component analysis (PCA). Before

applying PCA, the $6,441 \times 8$ matrix was z-transformed. We then obtained the scores of the first principal

component (PC) and reconstructed them into a 114×114 matrix, representing a shared correlation

structure across the eight matrices. Finally, we plotted the loadings of the first two PCs for the eight

matrices to examine their relative positions in the low-dimensional space.

3. Results

3.1. Simon dataset

We first analyzed a multi-session dataset from a single individual spanning over 16 years. The averaged correlation matrix of resting-state time series reveals clear modular structures, evident as square-like patterns along the diagonal and additional squares representing left-right homotopic networks (Figure 2C). Subsequently, we computed intra-individual correlations for regional GMV (Figure 2A) and ReHo (Figure 2B) across all available sessions. Both matrices display square-like patterns, although their spatial configurations differ. Notably, both intra-individual correlation matrices show moderate but significant correlations with the averaged resting-state time series correlation matrix (Spearman's correlation coefficients: $\rho_{GMV} = 0.29$; $\rho_{ReHo} = 0.39$).

Figure 2 Correlations of regional brain volume (GMV) (A), regional homogeneity (ReHo) (B), and averaged correlations of resting-state time series (C) across 114 regions of interest (ROIs). Black lines delineate Yeo's seven networks. The first 50 ROIs correspond to the left hemisphere, ROIs 51 to 100 correspond to the right hemisphere, and the remaining ROIs represent subcortical regions. D and E show the Spearman's correlations (ρ) between the time-series correlation matrix and GMV or ReHo correlations, respectively.

To examine how long-term age effects contribute to the observed correlations, we fitted a cubic

age model to the GMV time series for each ROI. Figures 3A–3C illustrate the raw time series, the fitted

cubic age effects, and the residuals for all 114 ROIs, respectively. The correlation matrices for these three

data types are presented in the bottom row of Figure 3. These matrices exhibited strong similarities, with

Spearman's correlation coefficients of 0.83 between the raw data and cubic age effects, 0.91 between the

raw data and residuals, and 0.56 between the cubic age effects and residuals (all p < 0.001).

Figure 3 Normalized gray matter volume (GMV) for the 114 regions of interest (ROIs) (A), their fitted

effects with a cubic age model (B) and the residuals (C). For illustration purpose, the time series in each

ROI was converted into z values. Panels D through F, correlation matrices for the raw, fitted cubic age

effects, and residuals across the 114 ROIs. Black lines delineate Yeo's seven networks. The first 50 ROIs

correspond to the left hemisphere, ROIs 51 to 100 correspond to the right hemisphere, and the remaining

ROIs represent subcortical regions.

Similarly, we fitted a cubic age model to the time series of normalized ReHo values for each ROI.

Figures 4A–4C display the raw time series, the fitted cubic age effects, and the residuals, respectively.

The correlation matrices for these three time series are shown in the bottom row of Figure 4. All matrices

were statistically significantly correlated with each other (all p < 0.001); however, the Spearman's

correlation coefficients varied considerably. The highest correlation was nearly perfect (0.99) between the

raw data and residuals, while the correlations between the cubic age effects and the other two (raw data

and residuals) were 0.34 and 0.21, respectively.

Figure 4 Regional Homogeneity (ReHo) for the 114 regions of interest (ROIs) (A), their fitted effects

with a cubic age model (B) and the residuals (C). For illustration purpose, the time series in each ROI was

converted into z values. Panels D through F, correlation matrices for the raw, fitted cubic age effects, and

residuals across the 114 ROIs. Black lines delineate Yeo's seven networks. The first 50 ROIs correspond

to the left hemisphere, ROIs 51 to 100 correspond to the right hemisphere, and the remaining ROIs

represent subcortical regions.

3.2. ADNI dataset

Next, we analyzed the ADNI dataset, where there were multiple participants, but each participant only

have a few sessions. We calculated averaged resting-state time series correlation from the 17 participants

(Figure 5C), which turned out to be very similar to those from the Simon dataset. We then calculated

structural correlations both intra-individually (Figure 5A) and inter-individually (Figure 5B). Both

matrices demonstrated functional network structures along the diagonal and between left and right

corresponding regions. Moreover, both correlation matrices were correlated with resting-state time series

correlation (ρ_{intra} = 0.41; ρ_{inter} = 0.37), which were slightly higher than that in the Simon dataset (0.29).

Figure 5 Correlations of regional brain volume (GMV) calculated intra-individually (A) and inter-

individually (B), and averaged correlations of resting-state time series (C) across 114 regions o interests

(ROIs). Black lines delineate Yeo's seven networks. The first 50 ROIs correspond to the left hemisphere,

ROIs 51 to 100 correspond to the right hemisphere, and the remaining ROIs represent subcortical regions.

D and E show the Spearman's correlations (ρ) between the resting-state time series correlation matrix and

GMV correlation matrices, respectively.

We next calculated correlations of glucose metabolic activity intra-individually and inter-

individually, and correlated the correlation matrices with resting-state time series correlation (Figure 6).

The correlations matrices of glucose metabolic activity showed more obvious functional network

structures than those in structural correlations. And most importantly, the both correlation matrices

showed strong correlations with resting-state time series correlation (ρ_{intra} = 0.71; ρ_{inter} = 0.64).

Figure 6 Correlations of regional brain glucose metabolism measured using positron emission tomography (PET) calculated intra-individually (A) and inter-individually (B), and averaged correlations of resting-state time series (C) across. Black lines delineate Yeo's seven networks. The first 50 ROIs correspond to the left hemisphere, ROIs 51 to 100 correspond to the right hemisphere, and the remaining ROIs represent subcortical regions. D and E show the Spearman's correlations (ρ) between the resting-state time series correlation matrix with the two PET correlation matrices, respectively.

3.3. Relationships among all the matrices

Finally, we calculated the correlations among the correlation matrices (Figure 7A). Due to the large number of ROI pairs, all correlations were statistically significant. However, it is more meaningful to focus on the relative effects of these correlations rather than their statistical significance. Here, we emphasize the correlations between the matrices of the two datasets, where the highest correlations were observed between corresponding modalities (highlighted blue rectangle and diamond markers). For instance, the strongest correlation with the mean time-series correlations in the Simon dataset was found with the time-series correlations in the ADNI dataset ($\rho = 0.67$). Similarly, the highest correlation with GMV correlations in the Simon dataset occurred with intra-individual GMV correlations in the ADNI dataset ($\rho = 0.46$), rather than inter-individual correlations. Notably, the highest correlation with ReHo

336   correlations in the Simon dataset was observed with inter-individual PET correlations in the ADNI dataset

337   (ρ = 0.51).

Figure 7 A. Spearman's correlation matrix among the correlation matrices. The blue rectangle and
diamond markers highlight the effects of interest. B. The first principal component of the correlation
matrices. Black lines delineate Yeo's seven networks. The first 50 ROIs correspond to the left
hemisphere, ROIs 51 to 100 correspond to the right hemisphere, and the remaining ROIs represent
subcortical regions. C. Loadings of the correlation matrices on the first and second principal components.
ADNI, Alzheimer's Disease Neuroimaging Initiative; GMV, gray matter volume; PET, positron emission
tomography; ROI region of interest.

We then conducted a PCA on the eight correlation matrices, with the first principal component
(PC) accounting for 59.0% of the variance (Figure 7B). Next, we visualized the loadings of the eight
matrices on the first two PCs (Figure 7C). The matrices formed distinct clusters in the plot. For instance,
the GMV correlation matrices were located at the bottom, while the time-series and PET correlation
matrices clustered in the top-right corner.

4. Discussion

Using two unique datasets encompassing both intra- and inter-individual effects, the current analysis
demonstrated how these effects contribute to correlation structures across brain regions. First, long-term

structural brain changes revealed correlation patterns that were small but significantly associated with

resting-state time-series correlations. Second, correlations in GMV were mainly driven by slow age

effects, white correlations in ReHo were mainly driven by state-like brain activity. Third, long-term

functional activity, as measured by ReHo or glucose metabolism, exhibited stronger correlation structures

and greater alignment with resting-state time-series correlations compared to structural measures. Finally,

correlation matrices from the two datasets showed greater similarity within the same modality than

between different modalities, suggesting that each modality provides distinct insights into interregional

relationships.

4.1. Structural correlations

This study analyzed a unique dataset from a single individual scanned over 16 years, revealing that intra-

individual structural correlations were modestly associated with resting-state connectivity (ρ = 0.29), the

lowest correlation observed among the analyses. These results were further validated using the ADNI

dataset, which, despite fewer longitudinal data points, showed a stronger correlation between intra-

individual structural correlations and resting-state connectivity (ρ = 0.41). These findings suggests that

structural brain development and aging within an individual are partially constrained by the brain's

functional organization. Moreover, by fitting cubic age models, the fitted curve yielded correlation

structures that were strongly correlated with the original GMV correlations (ρ = 0.83), further supported

that slow aging effects could contribute to the correlation structures.

However, it is noteworthy that the variability that could not be accounted for by cubic age

effects could still generate correlation matrix that were also even higher correlated with original GMV

correlations (ρ = 0.91). Although some studies reported evidence of changes of regional brain volumes

over short terms (Draganski et al., 2006, 2004), it is unlikely brain structures fluctuated with a large

amount over a short term. A more plausible explanation is that the variability is due to the variability of

MRI scanning settings (Di et al., 2022). For example, the homogeneity of magnetic fields may introduce

spatial signal variations in a scanner, which will in turn affect spatial variations in GMV estimates. The

spatial variability across scanner may introduce some correlation structures that somehow overlap with

brain functional networks. This calls for proper null models for the correlations between correlation

matrices (Váša and Mišić, 2022).

Notably, inter-individual structural correlations, when controlling for chronological age, showed

a similar association with resting-state connectivity ($\rho = 0.37$), indicating that individual differences

contribute comparably. The variability that give rise to the inter-individual correlation may be related to

genetics, life experience, and plasticity (Alexander-Bloch et al., 2013a; Evans, 2013). In previous studies

of structural "covariance", these factors could not be distinguished from development and aging effects.

In fact, the current findings suggest that multiple factors could give rise to the correlation structure. This

is also in line with studies showing that the structural "covariance" were modulated by factors such as age

(Vijayakumar et al., 2021). To enhance interpretability, researchers should consider restricting or

controlling for such factors in their analyses.

4.2. Functional correlations

Both ReHo and FDG-PET functional correlations differed from their structural counterparts. In the Simon

dataset, after removing cubic age effects, the residuals nearly perfectly mirrored the raw ReHo

correlations. This indicates that ReHo correlations are primarily driven by transient, state-dependent

fluctuations rather than long-term aging processes. Since ReHo summarizes neural activity during a scan

session and varies with different tasks (Di et al., 2024), these fluctuations likely capture changes in mood,

mental state, and daily experiences during scanning.

Additionally, correlations between ReHo or metabolic activity and resting-state connectivity were

generally stronger than those for regional brain volumes. This is expected because functional measures

are more directly linked to neural activity than structural ones. However, it's important to note that neural

activity measured over different temporal scales may represent distinct processes, making direct

comparisons of correlation structures challenging. Previous fMRI studies have shown that connectivity

patterns within a single session can vary depending on the frequency bands analyzed (Gohel and Biswal,

2015; Kajimura et al., 2023; Yuen et al., 2019). Future research should explore the characteristics of slow

functional fluctuations in summary measures like ReHo and glucose metabolic activity.

4.3. Implications for inter-individual correlation analysis

Compared to earlier studies, the current research reports slightly higher correlations with resting-state

connectivity. This improvement may stem from larger sample sizes, averaging data across multiple

sessions, or advancements in the preprocessing pipeline. These findings imply that the smaller

correlations observed in previous studies might partially result from noise, and that improvements in data

acquisition and processing can enhance the observed similarities. Nonetheless, the analysis also highlights

that each modality makes a unique contribution to the correlation structure, indicating an inherent limit to

the similarities that can be achieved between different modalities.

This analysis validates the use of structural and functional brain measures to investigate

interregional relationships, often referred to as functional connectivity when using functional data. Given

the complex factors influencing these measures, controlling for certain variables can enhance the

interpretability of correlation results. Our findings demonstrated that intra-individual correlations, which

account for individual differences, tend to exhibit stronger associations with resting-state functional

connectivity compared to inter-individual correlations, supporting the value of such controls. However, it

is important to note that multiple measures from the same individual are not always available, and in

some cases, inter-individual correlations may be the only feasible approach.

4.4. Limitations and future directions

One limitation of the current analysis is the narrow age range covered, making it unclear whether the

findings can be generalized to other age groups. However, due to the nature of long-term longitudinal

studies, obtaining suitable data across a broader age range is challenging, particularly with sufficient

within-individual runs and follow-up duration. Despite this limitation, the present findings offer novel

insights into the correlation structures underlying long-term effects on brain structure and function.

Further studies are needed to replicate and extend these results. Another limitation is that we relied solely

on resting-state functional connectivity to assess brain connectivity. Structural connectivity is typically

evaluated using diffusion-weighted imaging, and previous studies have established relationships between

inter-individual correlation measures and white matter connections (Gong et al., 2012; Lizarraga et al.,

2023). Future studies should incorporate diffusion-based measures to create a more comprehensive

picture of brain connectivity.

5. Conclusions

The current results to some extent validated the usage of inter-individual correlations as an estimate of

brain connectivity. The results also highlighted that multiple factors could contribute to the correlation

structure. Those factors may need to control or taken care of to boost interpretability of the results.

Acknowledgement

This study was supported by grants from (US) National Institutes of Health for Xin Di (R15MH125332)
and Bharat B. Biswal (5R01MH131335 and 1R01AG085665) and by (NJ) Governor's Council for
Medical Research and Treatment of Autism for Xin Di (CAUT25BRP005).
Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging
Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of
Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the
National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from
the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon
Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.;
Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its
affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer
Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research &
Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx
Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging;
Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of
Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions
are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee
organization is the Northern California Institute for Research and Education, and the study is coordinated
by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data
are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.
463

465 **References:**

466 Alexander-Bloch, A., Giedd, J.N., Bullmore, E., 2013a. Imaging structural co-variance between human
467   brain regions. Nat Rev Neurosci 14, 322–336. https://doi.org/10.1038/nrn3465

Alexander-Bloch, A., Raznahan, A., Bullmore, E., Giedd, J., 2013b. The convergence of maturational change and structural covariance in human cortical networks. The Journal of neuroscience : the official journal of the Society for Neuroscience 33, 2889–99. https://doi.org/10.1523/JNEUROSCI.3554-12.2013

Biswal, B., Yetkin, F.Z., Haughton, V.M., Hyde, J.S., 1995. Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. Magnetic resonance in medicine : official journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine 34, 537–41. https://doi.org/10.1002/mrm.1910340409

Biswal, B.B., Mennes, M., Zuo, X.-N., Gohel, S., Kelly, C., Smith, S.M., Beckmann, C.F., Adelstein, J.S., Buckner, R.L., Colcombe, S., Dogonowski, A.-M., Ernst, M., Fair, D., Hampson, M., Hoptman, M.J., Hyde, J.S., Kiviniemi, V.J., Kötter, R., Li, S.-J., Lin, C.-P., Lowe, M.J., Mackay, C., Madden, D.J., Madsen, K.H., Margulies, D.S., Mayberg, H.S., McMahon, K., Monk, C.S., Mostofsky, S.H., Nagel, B.J., Pekar, J.J., Peltier, S.J., Petersen, S.E., Riedl, V., Rombouts, S.A.R.B., Rypma, B., Schlaggar, B.L., Schmidt, S., Seidler, R.D., Siegle, G.J., Sorg, C., Teng, G.-J., Veijola, J., Villringer, A., Walter, M., Wang, L., Weng, X.-C., Whitfield-Gabrieli, S., Williamson, P., Windischberger, C., Zang, Y.-F., Zhang, H.-Y., Castellanos, F.X., Milham, M.P., 2010. Toward discovery science of human brain function. Proceedings of the National Academy of Sciences of the United States of America 107, 4734–9. https://doi.org/10.1073/pnas.0911855107

Di, X., Biswal, and Alzheimer's Disease Neu, B.B., 2012. Metabolic Brain Covariant Networks as Revealed by FDG-PET with Reference to Resting-State fMRI Networks. Brain Connectivity 2, 275–283. https://doi.org/10.1089/brain.2012.0086

Di, X., Biswal, B.B., 2023. A functional MRI pre-processing and quality control protocol based on statistical parametric mapping (SPM) and MATLAB. Frontiers in Neuroimaging 1, 1070151.

Di, X., Gohel, S., Thielcke, A., Wehrl, H.F., Biswal, B.B., Alzheimer's Disease Neuroimaging Initiative, 2017. Do all roads lead to Rome? A comparison of brain networks derived from inter-subject volumetric and metabolic covariance and moment-to-moment hemodynamic correlations in old individuals. Brain structure & function 222, 3833–3845. https://doi.org/10.1007/s00429-017-1438-7

Di, X., Jain, P., Biswal, B.B., 2024. Effects of Tasks on Functional Brain Connectivity Derived from Inter-Individual Correlations: Insights from Regional Homogeneity of Functional MRI Data. https://doi.org/10.1101/2024.06.02.597063

Di, X., Woelfer, M., Kühn, S., Zhang, Z., Biswal, B.B., 2022. Estimations of the weather effects on brain functions using functional MRI: A cautionary note. Human Brain Mapping 43, 3346–3356. https://doi.org/10.1002/hbm.25576

Draganski, B., Gaser, C., Busch, V., Schuierer, G., Bogdahn, U., May, A., 2004. Neuroplasticity: changes in grey matter induced by training. Nature 427, 311–2. https://doi.org/10.1038/427311a

Draganski, B., Gaser, C., Kempermann, G., Kuhn, H.G., Winkler, J., Büchel, C., May, A., 2006. Temporal and Spatial Dynamics of Brain Structure Changes during Extensive Learning. J. Neurosci. 26, 6314–6317. https://doi.org/10.1523/JNEUROSCI.4628-05.2006

Duchesne, S., Dieumegarde, L., Chouinard, I., Farokhian, F., Badhwar, A., Bellec, P., Tétreault, P., Descoteaux, M., Boré, A., Houde, J.-C., Beaulieu, C., Potvin, O., 2019. Structural and functional multi-platform MRI series of a single human volunteer over more than fifteen years. Sci Data 6, 245. https://doi.org/10.1038/s41597-019-0262-8

Evans, A.C., 2013. Networks of anatomical covariance. NeuroImage 80, 489–504. https://doi.org/10.1016/j.neuroimage.2013.05.054

Fox, P.T., Raichle, M.E., 1984. Stimulus rate dependence of regional cerebral blood flow in human striate cortex, demonstrated by positron emission tomography. Journal of neurophysiology 51, 1109–20.

Gohel, S.R., Biswal, B.B., 2015. Functional Integration Between Brain Regions at Rest Occurs in Multiple-Frequency Bands. Brain Connectivity 5, 23–34. https://doi.org/10.1089/brain.2013.0210

Gong, G., He, Y., Chen, Z.J., Evans, A.C., 2012. Convergence and divergence of thickness correlations with diffusion connections across the human cerebral cortex. NeuroImage 59, 1239–1248. https://doi.org/10.1016/j.neuroimage.2011.08.017

He, Y., Chen, Z.J., Evans, A.C., 2007. Small-world anatomical networks in the human brain revealed by cortical thickness from MRI. Cerebral cortex (New York, N.Y. : 1991) 17, 2407–19. https://doi.org/10.1093/cercor/bhl149

Horwitz, B., Duara, R., Rapoport, S.I., 1984. Intercorrelations of Glucose Metabolic Rates Between Brain Regions: Application to Healthy Males in a State of Reduced Sensory Input. Journal of Cerebral Blood Flow & Metabolism 4, 484–499. https://doi.org/10.1038/jcbfm.1984.73

Kajimura, S., Margulies, D., Smallwood, J., 2023. Frequency-specific brain network architecture in resting-state fMRI. Sci Rep 13, 2964. https://doi.org/10.1038/s41598-023-29321-5

Lizarraga, A., Ripp, I., Sala, A., Shi, K., Düring, M., Koch, K., Yakushev, I., 2023. Similarity between structural and proxy estimates of brain connectivity. J Cereb Blood Flow Metab 0271678X231204769. https://doi.org/10.1177/0271678X231204769

Mechelli, A., Friston, K.J., Frackowiak, R.S., Price, C.J., 2005. Structural covariance in the human cortex. The Journal of neuroscience : the official journal of the Society for Neuroscience 25, 8303–10. https://doi.org/10.1523/JNEUROSCI.0357-05.2005

Metter, E.J., Riege, W.H., Kuhl, D.E., Phelps, M.E., 1984. Cerebral Metabolic Relationships for Selected Brain Regions in Healthy Adults. Journal of Cerebral Blood Flow & Metabolism 4, 1–7. https://doi.org/10.1038/jcbfm.1984.1

Phelps, M.E., Kuhl, D.E., Mazziota, J.C., 1981. Metabolic Mapping of the Brain's Response to Visual Stimulation: Studies in Humans. Science 211, 1445–1448. https://doi.org/10.1126/science.6970412

Schaefer, A., Kong, R., Gordon, E.M., Laumann, T.O., Zuo, X.-N., Holmes, A.J., Eickhoff, S.B., Yeo, B.T.T., 2018. Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. Cerebral cortex (New York, N.Y. : 1991) 28, 3095–3114. https://doi.org/10.1093/cercor/bhx179

Song, X.-W., Dong, Z.-Y., Long, X.-Y., Li, S.-F., Zuo, X.-N., Zhu, C.-Z., He, Y., Yan, C.-G., Zang, Y.-F., 2011. REST: a toolkit for resting-state functional magnetic resonance imaging data processing. PloS one 6, e25031. https://doi.org/10.1371/journal.pone.0025031

Taylor, P.A., Gohel, S., Di, X., Walter, M., Biswal, B.B., 2012. Functional covariance networks: obtaining resting-state networks from intersubject variability. Brain connectivity 2, 203–17. https://doi.org/10.1089/brain.2012.0095

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B., Joliot, M., 2002. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. NeuroImage 15, 273–89. https://doi.org/10.1006/nimg.2001.0978

Váša, F., Mišić, B., 2022. Null models in network neuroscience. Nat Rev Neurosci 23, 493–504. https://doi.org/10.1038/s41583-022-00601-9

Vijayakumar, N., Ball, G., Seal, M.L., Mundy, L., Whittle, S., Silk, T., 2021. The development of structural covariance networks during the transition from childhood to adolescence. Sci Rep 11, 9451. https://doi.org/10.1038/s41598-021-88918-w

Yeo, B.T.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D., Hollinshead, M., Roffman, J.L., Smoller, J.W., Zöllei, L., Polimeni, J.R., Fischl, B., Liu, H., Buckner, R.L., 2011. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. Journal of neurophysiology 106, 1125–65. https://doi.org/10.1152/jn.00338.2011

Yuen, N.H., Osachoff, N., Chen, J.J., 2019. Intrinsic Frequencies of the Resting-State fMRI Signal: The Frequency Dependence of Functional Connectivity and the Effect of Mode Mixing. Front Neurosci 13, 900. https://doi.org/10.3389/fnins.2019.00900

Zang, Y., Jiang, T., Lu, Y., He, Y., Tian, L., 2004. Regional homogeneity approach to fMRI data analysis. NeuroImage 22, 394–400. https://doi.org/10.1016/j.neuroimage.2003.12.030

Zang, Y.-F., He, Y., Zhu, C.-Z., Cao, Q.-J., Sui, M.-Q., Liang, M., Tian, L.-X., Jiang, T.-Z., Wang, Y.-F.,
2007. Altered baseline brain activity in children with ADHD revealed by resting-state functional
MRI. Brain & development 29, 83–91. https://doi.org/10.1016/j.braindev.2006.07.002
Zeki, S., Watson, J., Lueck, C., Friston, K., Kennard, C., Frackowiak, R., 1991. A direct demonstration of
functional specialization in human visual cortex. J. Neurosci. 11, 641–649.
Zhang, Z., Liao, W., Zuo, X.-N., Wang, Z., Yuan, C., Jiao, Q., Chen, H., Biswal, B.B., Lu, G., Liu, Y.,
2011. Resting-state brain organization revealed by functional covariance networks. PloS one 6,
e28817. https://doi.org/10.1371/journal.pone.0028817
577