1 **Diverging transposon activity among polar bear sub-populations**
2 **inhabiting different climate zones**
3
4

5 **Authors:**
6
7 Alice M. Godden[1], Benjamin T. Rix[1] & Simone Immler[1]
8
9
10 **Affiliations**:
11 [1] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich NR4
12 7TJ, United Kingdom
13
14 Corresponding author: Alice M. Godden, Email: alice.godden@uea.ac.uk

15
16

Abstract

A new subpopulation of polar bears (*Ursus maritimus*) was recently discovered in the South-East

of Greenland. This isolated colony inhabits a warmer climate zone, akin to the predicted future

environments of polar bears with vastly reduced sea ice habitats, rendering this population of

bears particularly important. Over two-thirds of polar bears will be extinct by 2050 with total

extinction predicted by the end of this century, therefore understanding possible mechanisms of

adaptation via genomic analyses and preservation are critical. Transposable elements (TEs) are

parasitic mobile elements that may play a role in an adaptive response to environmental

challenges. We analysed transcriptome data from polar bear sub-populations in cooler North-

East (NEG) and warmer South-East Greenland (SEG) to compare TE activity between the two

populations and its correlation with temperature and associated changes in gene expression.

We identified activity hotspots in the genome of regions with significantly differentially expressed

TEs. LINE family TEs were the most abundant, and most differentially expressed and divergent

in the SEG population compared to reference TEs. We report a significant shift in TE activity and

age, with younger more abundant TEs in the SEG populations. Differentially expressed genes in

SEG populations were linked to *Foxo* signalling, ageing and metabolic pathways. Our results

provide insights into how a genomic response at the TE level may allow the SEG subpopulations

to adapt and survive to climate change and provides a useful resource for conservation in polar

bears.

Keywords: Transposable elements, thermal stress, adaptive response, Ursus, polar bear
37
38

Background

Global average temperatures in 2024 were more than 1°C above the pre-industrial era [1]. The

Arctic current is the warmest it has been in the last 125,000 years and temperatures continue to

rise [2,3]. In the South East of Greenland (SEG), the ice sheet margin is rapidly receding, causing

vast ice and habitat loss [4]. North-East Greenland (NEG) is an Arctic tundra, whereas SEG is

covered by forest tundra [5]. The SEG climate has high levels of precipitation, wind, and steep

coastal mountains [6]. This challenging habitat has led to severe warnings for the survival of the

polar bear *Ursus maritimus*, with the probability of a reduction in population size by over 90%

within 40 years set at 0.71. The IUCN have listed the polar bear as a vulnerable species [7], and

understanding possible ways of adaptation to climate change in such an enigmatic animal at the

genome level is key.

Environmental stress is reported to have a significant impact on genome structure, including the

expression and mobilisation of transposable elements (TEs) [8]. TEs are mobile genetic elements

that can shape evolution through self-replication and insertion events in the genome, which in

turn may change gene expression patterns [9,10]. TEs inserted into or near a promoter region may

alter the expression of genes and generate novel regulatory networks [11]. TEs make up 38.1% of

the genome in polar bears [12], which is similar to the giant panda *Ailuropoda melanoleuca* at

37%, and 36.1% of the dog *Canis familiaris* genome [13], and hence TEs may play an important

part in the process of adaptation. Current data show that over 150,000 genome insertion

mutations have been generated by TEs, driving variation in *ursine* and *tremarctine* bears [12]. The

aim of this study was to test if TE divergence between SEG and NEG polar bear populations

may explain genetic differences and variation in gene expression between these sub-

populations using a published transcriptome dataset [14]. We included information on temperature

variation into our analyses to test for a possible association between temperature and TE

activity.

Methods

Meteorological analysis

We collected and analysed meteorological data on the climate in NEG and SEG from the Danish

Meteorological Institute [15]. SEG locations included: Aputiteeq, Tasiilaq, Mittarfik, Kulusuk, Ikermiit

and Qaqortoq. NEG locations included: Ittoqqortoormiit, Daneborg, Port of Denmark and Station

Nord (see Supplementary File 1 for latitude and longitude co-ordinates). Temperature data

covering the highest, and lowest recorded temperatures each year from 1958-2024 were

included and the data was plotted with Tableau Public version 2024.1.

RNA-seq data

Transcriptome data from blood samples of adult polar bears were downloaded from the

European Nucleotide Archive under accession no. PRJNA669153 [14] (see Suppl. File 2 for

sample accession numbers and metadata) and aligned to the published reference polar bear

genome ASM1731132v1. As described in the original publication [14], RNA-sequencing (RNA-seq)

was based on total RNA and haemoglobin depletion step by CRISPR-Cas9 was used during

library preparation. The libraries were sequenced on Illumina HiSeqX to generate 151 bp paired

end reads [14]. We selected adult samples with an average coverage of 28.4 million aligned reads,

and included samples from five female NEG and seven female SEG bears and two male NEG

and three male SEG bears in our downstream analyses. The average number of mapped reads

across our 17 samples was 71 million. NF-core RNA-Seq pipeline version 1.4.2 was used to

analyse the RNA-seq data (https://nf-co.re/rnaseq/1.4.2/) [16]. Genome and annotation files were

used   and   accessed   as   follows: ASM1731132v1   (GCF_017311325.1)

GCF_017311325.1_ASM1731132v1_genomic.fna,   and   annotation

GCF_017311325.1_ASM1731132v1_genomic.gtf

https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_017311325.1/   ).   ASM1731132v1   is   the

most recent genome assembly and was used here as the reference genome. All differential

expression (DE) analyses are comparing intercepts? of NEG versus SEG population.

Differential TE expression analysis

To identify locus-specific expression of TEs and DE TE species, we used *Telescope* v1.0.3, a

software previously tested on blood sample data [17]. In addition to differential expression of TEs,

we used *RepeatMasker* [18] 4.1.1 and *RepeatModeler* 2.0.3 [19] for estimation of Kimura distances

to assess evolutionary divergence of nucleotide substitutions, TE activity and TE age: TEs with

Kimura distance <10 are younger and more recently active. For this pipeline, we merged the

bam files of individual samples by treatment group, then indexed and converted to fasta format

with *Samtools* [20]. *RepeatMasker* was used as follows: *RepeatMasker -species 29073 -par 5 -s -a*

-nolow -no_is, with fasta inputs split into 200 chunks for parallelisation. Aligned outputs were

merged and calcDivergenceFromAlign.pl was used for analysis. Genome sizes in the converted

fasta files were calculated using calcGenomesize.pl. This version of *RepeatMasker* utilised Dfam

v3.2, with genome version UrsMar1.0 TE annotations. The RepeatMasker default library for TEs

in polar bears utilises an UrsMar1.0 based annotation, therefore a custom library, "-lib", was

generated for the ASM1731132v1 reference genome assembly. To generate a custom library "-

lib" for use with RepeatMasker with the GCF_017311325.1_ASM1731132v1_genomic.fna as

reference genome for the ASM1731132v1 assembly. To create a *RepeatModeler* database*:*

BuildDatabase -name custom_repeats_db_ASM1731132v1 -engine ncbi "$GENOME_FILE".

Followed by: *RepeatModeler -database custom_repeats_db_ASM1731132v1 -pa 40 -engine*

ncbi. The output was used as the -lib input instead of species in the RepeatMasker pipeline.

RepeatModeler 2.0.3, ncbi-blast 2.10.1 and rmblast 2.10.0 [21], were used to analyse TE

divergence. To produce the TE gtf annotation file for Telescope pipeline, the Perl scripts from

TETranscripts [22] were used as follows: *perl makeTEgtf.pl -c 1 -s 2 -e 3 -o 6 -t 4*

ASM1731132v1_rmsk > ASM1731132v1_rmsk_TE.gtf. To test for a significant shift in the age of

Kimura substitution values, R scripts for general linear modelling were used with R packages

MASS v7.3-60.0.1 [23] *pscl* v1.5.9 [24]., with a negative binomial model (Count ~ Condition + Div;

where Count: the TE abundance at the Kimura substitution level; Condition: enrichment in SEG

population; Div: the Kimura substation levels) showing the best fit due to overdispersion in the

TE count data (See Suppl. Table 1 for model comparisons). To analyse genomic features

(coding sequence CDS, exon, gene, lncRNA, mRNA, pseudogene, snRNA and transcript) that

overlap with significantly DE TEs, the loci of TEs were extracted from the Telescope results and

mapped against gtf and bed file annotations to find genes and genomic features that overlap

with bedtools intersect [25]. All significantly DE TEs were tested for overlap with genomic features,

and significantly enriched DE genes overlapping with DE TEs in the SEG population were used

for GO term analysis.

Differential gene expression analysis

We used *DESeq2* [26] in *RStudio* v 1.4.1717 with R v4.1.1 [27] to run a differential expression

analyses between SEG and NEG bears. To account for spurious reads, any gene or TE

alignment with less than ten reads was discarded before running *DESeq2*. The basic model in

DESeq2 to generate a dds object was *design = ~ sex + population* (population referring to NEG

or SEG) to account for potential sex differences in expression levels. To test for a temperature

effect on TE expression, the average temperature over the year preceding blood sample

collection recorded in the town nearest to the bear's location was included in the analyses (mean

temperature over 2016, bears sampled in 2017) in an extended model: *design = ~ sex +*

temperature + population. Results were extracted as follows: *res_temperature <- results(dds,*

name="temperature"). Because location and temperature were colinear, the additive model was

used without an interaction term to avoid redundancy. The significance threshold for DE genes

and TEs was $\log_2$ fold change > 1 and a $p_{adj} < 0.05$. For GO term analyses, significantly we used

ShinyGO v0.77 [28] on DE genes with a background list of all genes found to be expressed in this

study, and figures were generated using custom Python scripts.

Software and analysis

All R scripts and Python scripts used to generate figures can be found at:

https://github.com/alicegodden/polarbear/. The following packages were used in the analysis

and presentation of the data in this project: *DESeq2* 1.34.1 [26], *Tidyverse* 2.0.0 [29],

EnhancedVolcano 1.12.0 [30], *ggplot2* 3.4.2 [31], *Readr* 2.1.4 [32], *tidyr* v1.3.0 [33], *viridis* v0.6.4 [34],

reshape [35], *hrbrthemes* v0.8.7, [36] and *gridExtra* v2.3, [37], *MASS* v7.3-60.0.1 [23] *pscl* v1.5.9 [24].

Results

Meteorological analysis

The meteorological observation data from Danish Meteorological Institute (DMI), revealed vast

variation in average lowest temperature measured from 1958-2024 between NEG and SEG (Fig.

1A), as denoted by the 95th percentile size of dots representing more variation. The variance in

observed lowest average temperatures was higher in SEG compared to NEG (Fig. 1B). Tasiilaq

(SEG) showed the highest reported temperature, and Station Nord (NEG) showed the lowest

recorded temperatures (Fig. 1A).

Differential TE expression analysis

We initially identified 179 significantly DE TEs (p_{adj} ≤ 0.05) between SEG and NEG bears when

including both sex and population into the model (Fig. Suppl. Fig. 2B). A Principal Components

Analysis showed modest clustering of samples based on population (Suppl. Fig. 2A). Most of the

significantly DE TE species belongs to the LINE family (Suppl. Fig. 2B-C). The most significantly

downregulated TE was DNA TE hAT-Charlie_dup1276 from the Charlie 8 subfamily. The most

significantly enriched TE was LINE TE L1_dup3552 from the L1 Carn1 subfamily (Suppl. Fig.

2B).

When adding temperature as a covariate into our analyses, samples clustered based on

temperature (Fig. 2B), and we identified 1,534 significantly DE TEs (Fig. 2A,C). The most

significantly enriched TE was L2b LINE L2_dup45185, and the most significantly downregulated

TE was L2a LINE/L2 dup_44480 (Fig. 2A). The most abundant TE class were again LINEs (Fig.

2C). To find genomic regions where TE activity may originate from, the genomic positions of

significantly DE TE species were plotted (Fig. 2D), revealing some clustering of DE TEs on

chromosomes NW_024423453.1, NW_024424119.1 and NW_02442563.1. No significantly DE

TE species were identified on the Y chromosome (NW_024423336.1), but a few LINE TE

species were located on the X chromosome (NW_024423319.1).

Age estimations of DE TEs showed significant divergence in TEs between the SEG and

NEG bear populations (Fig. 3), with both populations displaying divergence most strongly in

LINE family TEs, followed by DNA and LTR TEs. In the SEG population, we saw an overall

increase in TE divergence and a notable peak of younger divergent LINE TEs (Fig. 3). To

examine if the significantly DE TEs are overlapping with genic regions, we mapped the TE loci

against the bear genome. We found a significant enrichment of TEs in genic regions across the

whole genome and not restricted to DE genes (Suppl. Fig. 1A). We profiled the TE-overlapping

genes enriched in the SEG population with GO enrichment and found several significantly

enriched KEGG pathways, biological processes and molecular functions in SEG population. Of

relevance to a changed environment for the SEG bears, there was enrichment in KEGG

pathways (Suppl. Fig. 1B) for *Foxo* signalling, metabolism changes (ovarian steroidogenesis,

Thyroid hormone signalling) and immune response pathways including (human papillomavirus

infection, chemokine signalling and cancer pathways). For biological processes there was

significant enrichment of cardiac processes (SA node cell signalling pathways), metabolic

processes (phosphorous metabolic processes), and a range of developmental processes

including nervous system development (Suppl. Fig. 1C) and binding (Suppl. Fig. 1D).

Differential gene expression analysis

When running the model including *sex + temperature + population* as fixed variables,

clustering analysis revealed grouping of samples on PC1 with up to 29% variance explained by

temperature and latitude of sampling and 15% in expression (Fig. 4). From this DE analysis, we

identified 27 significantly DE genes, with LOC103677657 most downregulated and

LOC103668192 most enriched (Fig. 4B). We found no overlap in position of significantly DE

genes and TEs, but when using sex + population covariates, but addition of temperature

covariate for ASM1731132v1 revealed 10 overlapping genes with TEs (Fig. 4D). These genes

were: LOC121104823, LOC103667593, LOC103678552, LOC103677657, LOC121100717,

LOC121101012, BBS12 (ENSUMAG00000004608- Bardet-Biedl syndrome 12), LOC121104992,

SHANK1 and LOC103660844.

Differential gene expression analysis for the model with *sex + population* as fixed variables

revealed 13 significantly DE genes (Suppl. Fig. 3B). RSAD2 was most significantly

downregulated and LOC10367587 was most significantly enriched in SEG compared to NEG

bears. Of note, HSPH1 and HSPA4L were significantly enriched in SEG bears (Suppl. Fig. 3B).

GO terms analyses of biological processes, molecular function, and cellular function for the

significantly DE genes overlapping with DE TEs when aligned to ASM1731132v1 showed

enrichment for the following terms: internal cellular dynamics and movements: chromosome

movement towards spindle pole, PRC1 complex, organic acid:sodium symporter activity,

negative regulation of protein localization to nucleus, and immune responses: negative

regulation of viral genome replication, defence response to virus, defence response to symbiont,

(Suppl. Fig. 3C).

222
Discussion

The average lowest temperatures were highest in SEG towns in Greenland (Fig.1) suggesting

bears in this region to be exposed to a significantly warmer climate, which can trigger an

increase in TE activity. In accordance with this hypothesis, we found significantly DE TEs and

genes between SEG and NEG bears (Figs. 2-4). The most abundant DE TEs between the two

populations were LINE family TEs, and LTR TEs were significantly enriched in the SEG bears

(Fig. 2C). We also found that the loci of significantly DE TEs overlapped with genic regions of the

bear genome (Suppl. Fig. 1A). Kimura substitution analyses further showed an increased

abundance of younger DNA, LINE and LTR TE s in SEG bears (Fig. 3). Significantly DE genes

enriched in SEG polars included heat-shock protein genes HSPA4L and HSPH1 (Suppl. Fig.

3B). Taken together, our results suggest that TE activity in SEG bears is increased and may

contribute to the differential expression of genes between the two populations. Below, we

discuss the implications of our findings in a more general context and how they may help

understanding adaptation in wild animal populations more generally.

Meteorological analysis

Polar bears are dependent on sea ice, and the warmer and more variable SEG climate (Fig. 1A),

is forecast to be the norm for all polar bears by the end of the 21st century [14]. The warmest

temperatures in this study were observed in Tasiilaq (Fig. 1B), ranging between -30^{o}C and

+26^{o}C, which is characterised by a deep fjord and mountainous habitat with some drifting sea

pack ice as evidence of how temperature is isolating the SEG bears [6,38].

Genetic divergence between SEG and NEG polar bears

When comparing the genetic divergence between SEG and NEG bears, a previous study

reported an F_{ST} = 0.059 and concluded that SEG bears are more genetically diverged than other

polar bear populations in Greenland [14]. This genetic isolation is likely due to the limited gene flow

caused by the East Greenland coastal current [14], restricting bear movements and thus promoting

genetic drift over several hundred years of isolation. Despite this divergence, a previous study

using the same dataset reported no significantly DE genes, in contrast to our findings of 27 DE

genes. This discrepancy may be attributed to differences in model design, such as the inclusion

of intercept terms and the specific covariates considered during analysis. The previous analyses

were based on data from 16 adult samples, whereas we used 17 adult samples and used more

stringent DESeq2 analysis, filtering out all genes with less than 10 reads, compared to the 1

read threshold previously analysed. The PCA showed similar clustering patterns between our

study and the previous study.

Differential TE expression analysis

The main clusters of DE TEs across all samples identified in our analyses are LINEs, and they

were also the most abundant and significantly enriched in SEG bears (Fig. 2). LINEs and SINEs

are the most abundant family of TEs in mammalian genomes in general, with limited LTR and

DNA TE retention [11]. The DNA TEs detected in this RNA-seq analysis are likely being detected

through read-through of genes, this is unsurprising given the significant number of TEs in this

study that overlap with genic and coding regions of the genome (Suppl. Fig. 1A). LINE and SINE

TEs are also the most mobile and active TEs in bear genomes, and are most active in brown

bears, followed by polar bears [12]. To see if climate was impacting their expression, we included

impact of temperature in our analysis. This identified 1,534 significantly DE TEs (Fig. 2A, C).

This linking of temperature with genome mobility in polar bear aligns with similar studies in

model systems including *Arabidopsis* [39], *C. elegans* [40] and *Drosophila* [41], where environmental

stress was reported to increase TE mobility. Future work could include the use of DNA-

sequencing (DNA-seq) and small RNA-seq data to analyse for TE insertion events and

differential expression analysis of piRNAs.

Our finding of significant DE TEs (Fig.3) is in line with our idea of a putative role for TEs

in the adaptation of SEG polar bears and offers new avenues to understanding rapid adaptation

to changing environmental conditions. Divergent TEs between populations have also been

described in great pandas, where LINE and SINE TEs were generally younger in age, indicative

of recent mobilisation events [13]. We found a high abundance of LINE TEs in both populations,

potentially highlighting recent genome expansion of RNA class 1 TEs (Fig. 2C). LINE have been

previously shown to be important in the evolution of regulatory elements in the mammalian

genome [42]. This is interesting as we observed a peak of younger, divergent LINEs in the SEG

population of polar bears (Fig. 3). Older TEs included a modest amount of DNA TEs and LTRs in

older genome expansion events, although these divergent events constitute a smaller fraction of

the genome. This expansion of young LINEs, SINEs and LTRs was statistically significant (Fig.

3). Nevertheless, further DNA-sequencing analysis are needed to proof integration into the

genome. Overall, the observed significant enrichment of young TEs in SEG bears could be in

response to their challenging habitat and climate, but we need further evidence to fully

appreciate their integration back into the genome.

290   To test if TEs are being enriched in expression due to their localisation within a gene or

associated promoter region, we mapped the loci of the significantly DE TEs loci against the

reference genome and tested for overlap with genic regions. Due to limitations in the polar bear

genome assemblies, we could not perform intronic mapping, which is a technical as limitation

TEs are expected to fall into intronic regions [43]. However, we did manage to map TE loci to a

significant number of genes as well as to lncRNAs and pseudogenes (Suppl. Fig. 1A). We

observed significant enrichment of KEGG pathways and GO terms associated to these

overlapping genes (Suppl. Fig. 1). One of the key pathways, *Foxo* signalling, plays an important

role in adaptation to environmental stress and in ageing [44,45]. The significant enrichment of *Foxo*

signalling could be associated with increased TE activity in response to environmentally induced

DNA damage and repair [45]. The KEGG pathway *ovarian steroidogenesis* was also enriched,

genes in this process included: INSR and ALOX5. This is interesting as warmer environments

can affect polar bear diet, and if polar bears have lower fat reserves they are more likely to have

a failed pregnancy, with data previously showing a 28% reduction in the number of pregnant

female bears due to warmer environments [46]. Despite this, the upregulation of ovarian

steroidogenesis and lipid binding may be a way to compensate for hormonal imbalances caused

by temperature rise and nutritional deficits in the SEG population [47].

Increased TE activity has been shown to lead to immune responses in the host due to

detection of DNA damage and subsequent inflammatory responses [48]. This may explain the

enrichment of processes for immune signalling pathways associated with significantly DE genes

(Fig. 4B-C), and the genes that overlap with loci of significantly DE TEs (Suppl. Fig. 1). We

estimate that the recent spike in TE activity, as demonstrated by the significant peak of younger

TEs (Fig. 3) could be representative of at least a few hundred years, this would be enough to

cover several generations of polar bears and is a long time to allow for introduction of new

genetic variability due to TE activity to favour rapid adaptation [8,49].

While our study is assessing TE activity in a somatic tissue (blood), adaptation would rely on

germline TE activity. Some TEs are tissue specific, and their activity is generally less tightly

regulated in somatic tissues [50], providing a direct profile of a response to environmental stress

and possible resulting genetic divergence that may in time lead to genomic adaptation [8].

Nevertheless, the finding of differential TEs in a somatic tissue and the age difference in TEs

between the two populations, suggests an overall difference in TE activity across several

generations of bears in these two populations, possibly in response to differential environmental

factors.

Differential gene expression analysis

The evolutionary origins of bears as a taxonomic group are complicated, and they are thought to

have evolved over the past 5 million years in harsh and rapidly changing environments [51]. To

better understand polar bear genomes, we therefore profiled the differential gene expression of

SEG versus NEG polar bears, whilst using *sex* as a covariate to account for differences in

sample numbers and sex biases in gene expression. Given the differences in habitat over the

recent decades of extreme warming and variable weather patterns in SEG [6], it is unsurprising

that SEG bears are starting to show significant differences in gene expression compared to NEG

bears.

Linking gene and TE profiles in response to temperature

Our re-analyses of gene expression using a more conservative filtering steps of the data

revealed a number of significantly DE genes between the two populations when taking sex and

location into account, which contrasts with earlier results on the same dataset reporting no

significant DE genes [14]. Upregulated DE genes in SEG bears included RSAD2 (downregulated),

ACTR2, HSPA4L and HSPH1 (enriched) (Suppl. Fig. 3B), where RSAD2 (also known as Viperin)

is an evolutionarily conserved antiviral protein-coding gene [52]. ACTR2 (actin-related protein 2) in

house mice *Mus musculus* is involved in thrombosis, and mutations in the gene have been found

to be linked to thrombo-suppression [53] whereas HSPA4L (Hsp70) and HSPH1 (Hsp110) are

highly conserved and related families of proteins known as heat-shock proteins universal to all

eukaryotic species so far, acting as molecular chaperones [54]. This is interesting as other Hsp

chaperones, including *Hsp90*, regulate protein folding, unfolding and degradation [55]. *Hsp90* and

Hsp70 are also important in piRNA biogenesis and silencing of TEs, and function in elevated

temperature environments [56]. The most significant differential expression in TEs was observed in

the analysis with the addition of temperature as a covariate (Fig. 2A). This is supported by

evidence in *Drosophila* where TEs are activated in response to heat shock due to *Hsp70* and

other chaperones having a role in the collapse of piRNA production, and thus further TE action

[57]. Combined, this is evidence that the host genome is potentially responding to the changes in

environmental temperature.

When taking temperature as covariate into account (Fig. 4B), out of our 27 significantly

DE genes, 10 of these gene loci overlapped with significantly DE TEs, these included:

LOC121104823, LOC103678552, LOC121100717, and LOC121104992 (all enriched),

LOC121101012 (downregulated) are all long non-coding RNAs (lncRNAs) (Fig. 4B, D). Long

non-coding RNAs are highly abundant in mammals, with over 80% of transcribed products

generating lncRNAs that have roles in chromatin organization and gene regulation [58]. The

accessibility of chromatin, and methylation is significant as this can lead to de-repression of TEs

[59].

The most significantly downregulated gene in the SEG population was LOC103677657, a

pseudogene with unknown function. BBS12, an mRNA for Bardet-Biedl syndrome 12 protein,

involved in chaperone protein complex assemblies, and is orthologous with human *BBS12,* was

another downregulated in the SEG bears. Mutations or loss of BBS12 in humans leads to cilia

abnormalities [60]. Significantly enriched genes in SEG bears with reported functions included

LOC103667593 which is a predicted mRNA for filaggrin-2-like protein linked to inflammatory skin

diseases in human [61], and LOC121104992, a predicted mRNA for keratin-associated 4-8-like

protein. SHANK1 was also significantly upregulated in SEG bears and is an mRNA for SH3 and

multiple ankyrin repeat domains protein 1 and orthologous to human *SHANK1*. *SHANK1*

encodes a scaffold protein and can inhibit the activation of integrins [62], which can lead to

changes in cell signalling. This links to the GO term processes observed for phosphatidylinositol

activity highlighting cellular activity and signalling (Fig. 4C). Other GO terms in response to the

warmer SEG climate showed enrichment for terms related to immune responses including

defence response to virus, and negative regulation of viral genome replication (Suppl. Fig. 3C).

TE invasion is known to induce an immune response in mammals, activating the interferon and

innate immune pathways [63]. Polar bears in regions of reduced sea ice habitat are at higher risk

of pathogen exposure [64], which may explain the changes in SEG bears. However, whether SEG

bears are indeed exposed to higher pathogen loads needs further careful investigations.

Conclusions

Our work revealed key significant differences in the genomes of SEG bears compared to the

NEG bears, most strikingly in the mobile genome, potentially because of more variable and

higher temperatures in SEG. The main differences we found were the significantly DE TEs and

recent divergent genome expansion events exhibited by LINEs. Our study provides a basis for

further work into TEs in a wider subset of polar bear subpopulations and highlights potential

candidate TE species for further investigation as potential markers of environmental stress.

Future work to revise and further annotate the *Ursus maritimus* genome, at all levels not just TEs

will prove highly valuable in the efforts to preserve this vulnerable species through genomic

analyses. Larger scale analysis across all other subpopulations of polar bear, and comparative

analyses across different bear species in a range of climate zones may prove highly valuable for

the planning of conservation efforts and species management.

Abbreviations:

DE – differentially expressed

DMI - Danish Meteorological Institute

DNA-seq – DNA sequencing

lncRNAs – Long non-coding RNAs

NEG North-East Greenland

piRNA- piwi-interacting RNA

RNA-seq – RNA sequencing

SEG – South-East Greenland

TEs – Transposable elements

References

Masson-Delmotte, V. *et al. Global Warming of 1.5 C: IPCC special report on impacts of global warming of 1.5 C above pre-industrial levels in context of strengthening response to climate change, sustainable development, and efforts to eradicate poverty.* (Cambridge University Press, 2022).

Stendel, M., Christensen, J. H. & Petersen, D. Arctic climate and climate change with a focus on Greenland. *Advances in Ecological Research* **40**, 13-43 (2008).

Horhold, M. *et al.* Modern temperatures in central-north Greenland warmest in past millennium. *Nature* **613**, 503-507 (2023). https://doi.org/10.1038/s41586-022-05517-z

Bjørk, A. A. *et al.* An aerial view of 80 years of climate-related glacier fluctuations in southeast Greenland. *Nature Geoscience* **5**, 427-432 (2012).

Normand, S. *et al.* A greener Greenland? Climatic potential and long-term constraints on future expansions of trees and shrubs. *Philos Trans R Soc Lond B Biol Sci* **368**, 20120479 (2013). https://doi.org/10.1098/rstb.2012.0479

Berdahl, M. *et al.* Southeast Greenland winter precipitation strongly linked to the Icelandic low position. *Journal of Climate* **31**, 4483-4500 (2018).

Regehr, E. V. *et al.* Conservation status of polar bears (Ursus maritimus) in relation to projected sea-ice declines. *Biol Lett* **12** (2016). https://doi.org/10.1098/rsbl.2016.0556

Godden, A. M. & Immler, S. The potential role of the mobile and non-coding genomes in adaptive response. *Trends Genet* (2022). https://doi.org/10.1016/j.tig.2022.08.006

Bourque, G. *et al.* Ten things you should know about transposable elements. *Genome Biol* **19**, 199 (2018). https://doi.org/10.1186/s13059-018-1577-z

Chang, N. C., Rovira, Q., Wells, J., Feschotte, C. & Vaquerizas, J. M. Zebrafish transposable elements show extensive diversification in age, genomic distribution, and developmental expression. *Genome Res* **32**, 1408-1423 (2022). https://doi.org/10.1101/gr.275655.121

Platt, R. N., 2nd, Vandewege, M. W. & Ray, D. A. Mammalian transposable elements and their impacts on genome evolution. *Chromosome Res* **26**, 25-43 (2018). https://doi.org/10.1007/s10577-017-9570-z

Lammers, F., Gallus, S., Janke, A. & Nilsson, M. A. Phylogenetic Conflict in Bears Identified by Automated Discovery of Transposable Element Insertions in Low-Coverage Genomes. *Genome Biol Evol* **9**, 2862-2878 (2017). https://doi.org/10.1093/gbe/evx170

Peng, C. *et al.* Can-SINE dynamics in the giant panda and three other Caniformia genomes. *Mob DNA* **9**, 32 (2018). https://doi.org/10.1186/s13100-018-0137-0

Laidre, K. L. *et al.* Glacial ice supports a distinct and undocumented polar bear subpopulation persisting in late 21st-century sea-ice conditions. *Science* **376**, 1333-1338 (2022). https://doi.org/10.1126/science.abk2793

DMI.   (Danish Meteorological Institute, https://www.dmi.dk/vejrarkiv, 2024).

Ewels, P. A. *et al.* The nf-core framework for community-curated bioinformatics pipelines. *Nat Biotechnol* **38**, 276-278 (2020). https://doi.org/10.1038/s41587-020-0439-x

Bendall, M. L. *et al*. Telescope: Characterization of the retrotranscriptome by accurate estimation of transposable element expression. *PLoS Comput Biol* **15**, e1006453 (2019). https://doi.org/10.1371/journal.pcbi.1006453

Smit, A. F., Hubley, R. & Green, P. RepeatMasker software (2013). https://doi.org/http://www.repeatmasker.org

Smit, A. F. & Hubley, R. RepeatModeler Open-1.0. (2015). https://doi.org/http://www.repeatmasker.org

Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754-1760 (2009). https://doi.org/10.1093/bioinformatics/btp324

Camacho, C. *et al*. BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009). https://doi.org/10.1186/1471-2105-10-421

Jin, Y., Tam, O. H., Paniagua, E. & Hammell, M. TEtranscripts: a package for including transposable elements in differential expression analysis of RNA-seq datasets. *Bioinformatics* **31**, 3593-3599 (2015). https://doi.org/10.1093/bioinformatics/btv422

Venables, W. & Ripley, B. *Modern Applied Statistics with S, Springer, New York: ISBN 0-387-95457-0*. (2002).

Jackman, S. pscl: Classes and methods for r. developed in the political science computational laboratory, stanford university. department of political science, stanford university, stanford, ca. r package version 1.03. 5. *http://www*. *pscl. stanford. edu/* (2010).

Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841-842 (2010). https://doi.org/10.1093/bioinformatics/btq033

Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014). https://doi.org/10.1186/s13059-014-0550-8

R Core Team, R. R Core Team R: a language and environment for statistical computing. *Foundation for Statistical Computing* (2020).

Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals and plants. *Bioinformatics* **36**, 2628-2629 (2020). https://doi.org/10.1093/bioinformatics/btz931

Wickham, H. *et al*. Welcome to the tidyverse. *Journal of Open Source Software* **4**, 1-6 (2019). https://doi.org/https://doi.org/10.21105/joss.01686)

Blighe, K., Rana, S. & Lewis, M. EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling. https://github.com/kevinblighe/EnhancedVolcano (2021).

Ginestet, C. ggplot2: Elegant Graphics for Data Analysis. *J Roy Stat Soc A* **174**, 245-245 (2011). https://doi.org/DOI 10.1111/j.1467-985X.2010.00676_9.x

Wickham, H., Hester, J., Bryan, J.,. readr: Read Rectangular Text Data., https://CRAN.R-project.org/package=readr (2023). https://doi.org/ https://CRAN.R-project.org/package=readr

Wickham H, V. D., Girlich M,. tidyr: Tidy Messy Data. R package version 1.3.0. https://github.com/tidyverse/tidyr (2023).

Garnier, S., Ross, N., Rudis, B., Sciaini, M. & Scherer, C. viridis(Lite) - Colorblind-Friendly Color Maps for R., 10.5281/zenodo.4679423 (2023). https://doi.org/10.5281/zenodo.4679423

Wickham, H. reshape: Flexibly Reshape Data. (2022). https://doi.org/https://CRAN.R-project.org/package=reshape

Rudis, B. *et al*. hrbrthemes: Additional Themes, Theme Components and Utilities for 'ggplot2'. (2024). https://doi.org/ https://CRAN.R-project.org/package=hrbrthemes

Auguie, B. & Antonov, A. gridExtra: Miscellaneous Functions for "Grid" Graphics. *cran* (2017). https://doi.org/https://CRAN.R-project.org/package=gridExtra

Wiig, Ø. *et al*. Variation in non-metrical skull traits of polar bears (Ursus maritimus) and relationships across East Greenland and adjacent subpopulations (1830−2013). *Polar Biology* **42**, 461-474 (2019).

Roquis, D. *et al*. Genomic impact of stress-induced transposable element mobility in Arabidopsis. *Nucleic Acids Res* **49**, 10431-10447 (2021). https://doi.org/10.1093/nar/gkab828

Garrigues, J. M., Tsu, B. V., Daugherty, M. D. & Pasquinelli, A. E. Diversification of the Caenorhabditis heat shock response by Helitron transposable elements. *Elife* **8** (2019). https://doi.org/10.7554/eLife.51139

Garcia Guerreiro, M. P. What makes transposable elements move in the Drosophila genome? *Heredity (Edinb)* **108**, 461-468 (2012). https://doi.org/10.1038/hdy.2011.89

Roller, M. *et al*. LINE retrotransposons characterize mammalian tissue-specific and evolutionarily dynamic regulatory regions. *Genome Biol* **22**, 62 (2021). https://doi.org/10.1186/s13059-021-02260-y

Zhang, Y., Romanish, M. T. & Mager, D. L. Distributions of transposable elements reveal hazardous zones in mammalian introns. *PLoS Comput Biol* **7**, e1002046 (2011). https://doi.org/10.1371/journal.pcbi.1002046

Rodriguez-Colman, M. J., Dansen, T. B. & Burgering, B. M. T. FOXO transcription factors as mediators of stress adaptation. *Nat Rev Mol Cell Biol* **25**, 46-64 (2024). https://doi.org/10.1038/s41580-023-00649-0

Song, J. *et al*. FOXO-regulated OSER1 reduces oxidative stress and extends lifespan in multiple species. *Nat Commun* **15**, 7144 (2024). https://doi.org/10.1038/s41467-024-51542-z

Molnar, P. K., Derocher, A. E., Klanjscek, T. & Lewis, M. A. Predicting climate change impacts on polar bear litter size. *Nat Commun* **2**, 186 (2011). https://doi.org/10.1038/ncomms1183

Knott, K. K., Mastromonaco, G. F., Owen, M. A. & Kouba, A. J. Urinary profiles of progestin and androgen metabolites in female polar bears during parturient and non-parturient cycles. *Conserv Physiol* **5**, cox023 (2017). https://doi.org/10.1093/conphys/cox023

Kong, Y. *et al.* Transposable element expression in tumors is associated with immune infiltration and increased antigenicity. *Nat Commun* **10**, 5228 (2019). https://doi.org/10.1038/s41467-019-13035-2

Capy, P., Gasperi, G., Biemont, C. & Bazin, C. Stress and transposable elements: co-evolution or useful parasites? *Heredity (Edinb)* **85 (Pt 2)**, 101-106 (2000). https://doi.org/10.1046/j.1365-2540.2000.00751.x

Teixeira, F. K. *et al.* piRNA-mediated regulation of transposon alternative splicing in the soma and germ line. *Nature* **552**, 268-272 (2017). https://doi.org/10.1038/nature25018

Kumar, V. *et al.* The evolutionary history of bears is characterized by gene flow across species. *Sci Rep* **7**, 46487 (2017). https://doi.org/10.1038/srep46487

Upadhyay, A. S. *et al.* Viperin is an iron-sulfur protein that inhibits genome synthesis of tick-borne encephalitis virus via radical SAM domain activity. *Cell Microbiol* **16**, 834-848 (2014). https://doi.org/10.1111/cmi.12241

Westrick, R. J. *et al.* Sensitized mutagenesis screen in Factor V Leiden mice identifies thrombosis suppressor loci. *Proc Natl Acad Sci U S A* **114**, 9659-9664 (2017). https://doi.org/10.1073/pnas.1705762114

Easton, D. P., Kaneko, Y. & Subjeck, J. R. The hsp110 and Grp1 70 stress proteins: newly recognized relatives of the Hsp70s. *Cell Stress Chaperones* **5**, 276-290 (2000).

Rosenzweig, R., Nillegoda, N. B., Mayer, M. P. & Bukau, B. The Hsp70 chaperone network. *Nat Rev Mol Cell Biol* **20**, 665-680 (2019). https://doi.org/10.1038/s41580-019-0133-3

Karam, J. A., Parikh, R. Y., Nayak, D., Rosenkranz, D. & Gangaraju, V. K. Co-chaperone Hsp70/Hsp90-organizing protein (Hop) is required for transposon silencing and Piwi-interacting RNA (piRNA) biogenesis. *J Biol Chem* **292**, 6039-6046 (2017). https://doi.org/10.1074/jbc.C117.777730

Cappucci, U. *et al.* The Hsp70 chaperone is a major player in stress-induced transposable element activation. *Proc Natl Acad Sci U S A* **116**, 17743-17950 (2019). https://doi.org/10.1073/pnas.1903936116

Karlsson, O. & Baccarelli, A. A. Environmental Health and Long Non-coding RNAs. *Curr Environ Health Rep* **3**, 178-187 (2016). https://doi.org/10.1007/s40572-016-0092-1

Zhou, W., Liang, G., Molloy, P. L. & Jones, P. A. DNA methylation enables transposable element-driven genome expansion. *Proc Natl Acad Sci U S A* **117**, 19359-19366 (2020). https://doi.org/10.1073/pnas.1921719117

Seo, S. *et al.* BBS6, BBS10, and BBS12 form a complex with CCT/TRiC family chaperonins and mediate BBSome assembly. *Proc Natl Acad Sci U S A* **107**, 1488-1493 (2010). https://doi.org/10.1073/pnas.0910268107

Zeitvogel, J. *et al.* GATA3 regulates FLG and FLG2 expression in human primary keratinocytes. *Sci Rep* **7**, 11847 (2017). https://doi.org/10.1038/s41598-017-10252-x

Lilja, J. *et al.* SHANK proteins limit integrin activation by directly interacting with Rap1 and R-Ras. *Nat Cell Biol* **19**, 292-305 (2017). https://doi.org/10.1038/ncb3487

Gazquez-Gutierrez, A., Witteveldt, J., S, R. H. & Macias, S. Sensing of transposable elements by the antiviral innate immune system. *RNA* **27**, 735-752 (2021). https://doi.org/10.1261/rna.078721.121

Rode, K. D. *et al.* Increased pathogen exposure of a marine apex predator over three decades. *PLoS One* **19**, e0310973 (2024). https://doi.org/10.1371/journal.pone.0310973

Acknowledgements

We thank Dr. Oliver Tam (Hammel lab), for TE annotation file generation support. This study was funded by grants from the Natural Environment Research Council (NE/S011188/1) and the European Research Council (SELECTHAPLOID - 101001341) to SI.

Declarations

Authors declare no conflict or competing interests.

Author contributions

AMG conceived the idea and conducted the bioinformatic analyses, and wrote the manuscript.

BR contributed to data analysis and conducted metereological analyses. SI supported

manuscript preparation and funding.

Data and code availability

RNA-seq data was deposited by [14] under ENA accession PRJNA669153. All R and python

scripts used to analyse this data can be found in the following Github repository:

https://github.com/alicegodden/polarbear/. Metadata used for the files accessed can be found in

Supplementary file 2. The datasets supporting this article have been uploaded as part of the

supplementary material in files 1-8. Supplementary files include all raw counts data from RNA-

seq analyses, DESeq2 outputs for differential gene and TE expression with different models

can be found here: https://github.com/alicegodden/polarbear/tree/main/supplementary_data .

Diverging transposon activity among polar bear sub-populations
inhabiting different climate zones
3
4
5 **Authors:**
6

7 Alice M. Godden[1], Benjamin T. Rix[1] & Simone Immler[1]
8
9
10 **Affiliations**:
11 [1] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich NR4
12 7TJ, United Kingdom
13
14 Corresponding author: Alice M. Godden, Email: alice.godden@uea.ac.uk

15
16

FIGURES

Figure 1- Meteorological observation data of temperature from the DMI. **(A)** Highest and lowest observed temperatures from 1958-2024, listed from Northernmost to most Southern. **(B)** Colour of points indicates the average lowest observed temperature, with the size of point showing the variance at the 95th percentile. SEG locations included: Aputiteeq, Tasiilaq, Mittarfik, Kulusuk, Ikermit and Qaqortoq. NEG locations included: Ittoqqortoormiit, Daneborg, Port of Denmark and Station Nord. In this study north of 64° was considered NEG. The bear icons are indicative of the latitude where they were sampled, red meaning SEG and blue NEG. The Arctic circle is denoted by a dashed yellow line at 66°.

Figure 2- Comparison of expression and activity of TEs in NEG versus SEG bear taking temperature into account, profiling impact in SEG population. **(A)** Differential expression analysis from DESeq2 analysis of TE species identified 173 significantly differentially expressed TEs (p_{adj} ≤ 0.05) by the Telescope pipeline, with p cut off for significance p_{adj} ≥ 0.05 on y axis and log$_2$ fold change more than 0.5 (1). **(B)** Principal Component Analysis of in SEG and NEG polar bear populations including temperature and location. **(C)** Count data of significantly differentially expressed TEs at the family level as tested by hypergeometric testing (*** = p ≤0.001). **(D)** Genomic position of DE TE species and co-located DEGs.

74
Figure 3 – Repeat landscape of TE age and activity in NEG v SEG populations. TE age and
divergence estimation by Kimura substitution analysis with RepeatMasker. The estimated
divergence for SEG bears was greater than that in NEG bears. Analysis of negative binomial
testing with general linear modelling results to assay the model *Count ~ Condition + Div*, for a shift
in TE age with negative binomial modelling showed significant increase in younger TEs in SEG
populations at *p* ≤ 0.001 for all TE families. See Suppl. Table 1 for full statistical results.
81

82
83

85
Figure 4 – Analysis of differential gene expression to assay the main impact of temperature on
the genomes in DESeq2 outputs looking at changes in SEG population. **(A)** Principal Component
Analysis of differentially expressed genes in NEG and SEG polar bears taking temperature and
sex into account shows clustering according to geographical location. **(B)** Volcano plot of
differentially expressed genes, with p cut off for significance $p_{adj} \geq 0.05$ on y axis and $\log_2$ fold
change more than 0.5 (1), showing 27 significantly DE genes. **(C)** GO terms analysis of
significantly differentially expressed genes between SEG and NEG bears. **(D)** Loci of significantly
differentially expressed genes with and without temperature covariate in *DESeq2* analysis.

Diverging transposon activity among polar bear sub-populations
inhabiting different climate zones
3
4
5   **Authors:**
6
7   Alice M. Godden[1], Benjamin T. Rix[1] & Simone Immler[1]
8
9
10   **Affiliations**:
11   [1] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich
12   NR4 7TJ, United Kingdom
13
14   Corresponding author: Alice M. Godden, Email: alice.godden@uea.ac.uk
15
16

Supplementary Figure 1- Significantly DE TE loci and their overlap with genomic features and genes in NEG v SEG populations. **(A)** Abundance of TEs overlapping genomic features, statistical enrichment analysed if this overlap was more significant than to be expected by chance (χ^2 test of independence), $p \leq 0.001$, as denoted by ***. SINEs were most significantly enriched in gene regions ($\chi^2 = 1036.94$) and LINEs were most significantly depleted in transcript regions ($\chi^2 = 9.12$). Stars in red show significant enrichment, stars in blue denote significant depletion. **(B)** KEGG pathways of overlapping genes. GO terms analysis of genes overlapping with significantly enriched TEs for **(C)** biological processes and **(D)** molecular functions.

Supplementary Figure 2- Comparison of expression and activity of TEs between SEG and NEG bears modelled on *sex + population* **(A)** PCA analysis of NEG v SEG bears show clustering based on geographical location of sample for TE expression. **(B)** Differential expression in DESeq2 analysis of TE species identified 173 significantly differentially expressed TEs ($p_{adj} \leq 0.05$). **(C)** Count data of significantly differentially expressed TEs at the family level. **(D)** Genomic position of TE species that were significantly differentially expressed.

98
Supplementary Figure 3 – Differential expression analysis in *DESeq2* using RNA-
sequencing data from samples in PRJNA669153 aligned to the reference genome
ASM1731132v1 to compare NEG and SEG bears and observe impact on SEG population. **(A)**
Principal Component Analysis shows variation in gene expression correlates to geographical
location of the sample. **(B)** Volcano plot of the differentially expressed genes observed
following DESeq2 analysis, with 13 significantly differentially expressed genes in total, with
cut off for significance $p_{adj} \geq 0.05$ on y axis and $\log_2$ fold change more than 0.5 (1). **(C)** GO
terms analysis, examining biological processes (BP), Molecular function (MF) and cellular
function (CF). All genes expressed in the study were used as a background list, and those
with $p \leq 0.1$ processed for GO terms analysis with ShinyGO.

Supplementary Table 1 – Statistical analysis of models to examine a shift in TE family age
from Kimura Substitution analysis data from RepeatMasker outputs. General linear modelling
results to assay the model *Count ~ Condition + Div*, based on Akaike Information Criterion
(AIC) and overdispersion values. Lower AIC values indicate better model fit. The Negative
Binomial model is preferred over the Poisson model due to significant overdispersion in TE
counts. The lower section of the table presents the results of a shift analysis, showing the
estimated shift change, statistical significance (*p*-value), and the inferred shift direction in SEG
polar bears Negative shift values and significant indicate that younger TEs are enriched in
SEG regions.

TE_Family	Model_Type	AIC		Overdispersion
DNA	Poisson	244884562		NA
DNA	Negative Binomial	244599505		457503.742
LINE	Poisson	4401719973		NA
LINE	Negative Binomial	4396475509		11897277.4
LTR	Poisson	66546483.4		NA
LTR	Negative Binomial	66503766.4		137545.593
TE_Family	Model_Type	Shift change	P value	Shift direction in SEG
DNA	Negative binomial	-0.0461	<0.001 / ***	Younger
LINE	Negative binomial	-0.0469	<0.001/***	Younger
LTR	Negative binomial	-0.0230	<0.001/***	Younger

120
121

122   **SUPPLEMENTARY FILES**

123   All supplementary files can be found here:
124   https://github.com/alicegodden/polarbear/tree/main/supplementary_data
125
126   **Supplementary File 1-** Meteorological data accessed from DMI **–** "Suppl. File. 1-
127   Temperature_data - DMI.csv"
128
129   **Supplementary File 2**- Metadata of samples used for the RNA-seq analysis "Suppl. File. 2.
130   bear_metadata_adult.csv"
131
132   **Supplementary File 3 –** Raw counts RNA-seq ASM1731132v1: "Suppl. File 3-Raw counts
133   RNA-seq ASM1731132v1.tsv"
134
135   **Supplementary File 4 –** DESeq2 data RNA-seq ASM1731132v1: "Suppl. File 4-DESeq2 data
136   RNA-seq ASM1731132v1.csv"
137
138   **Supplementary File 5 –** Telescope raw counts RNA-seq- TEs ASM1731132v1: "Suppl. File
139   5-Telescope raw counts RNA-seq- TEs ASM1731132v1.zip" compressed csv file.
140
141   **Supplementary File 6 –** Telescope DESeq2 data RNA-seq ASM1731132v1: "Suppl. File 6-
142   Telescope DESeq2 data RNA-seq ASM1731132v1.csv"
143
144   **Supplementary File 7-** DESeq2 data RNA-seq ASM1731132v1 Temperature effect: "Suppl.
145   File. 7- Temperature_effect_results_newgen_bear.csv"
146
147   **Supplementary File 8-** DESeq2 data RNA-seq ASM1731132v1 Temperature effect on TE
148   expression: "Supp. File. 8- Temperature_effect_results_newgen_bear_TelescopeTEs.txt"
149
150