Spike sorting biases and information loss in a detailed cortical model

Steeve Laquitaine[1*], Milo Imbeni[1], Joseph Tharayil[1], James B. Isbister[1],
Michael W. Reimann[1]

[1]Blue Brain Project, The Swiss Federal Institute of Technology, 9 Chemin des Mines, Geneva, 1202, Switzerland.

*Corresponding author(s). E-mail(s): laquitainesteeve@gmail.com;
Contributing authors: milo.imbeni@maastrichtuniversity.nl; tharayiljoe@gmail.com;
isbisterjb@gmail.com; mwr@reimann.science;

Abstract

Sorting electrical signals (spikes) from extracellular recordings of large groups of connected neurons is essential to understanding brain function. Despite transformative advances in dense extracellular recordings, the activity of most cortical neurons remains undetected. Small simulations with known neuron spike times offer critical ground truth data to improve spike sorting. Yet, current simulations underestimate neuronal heterogeneity and connectivity, which can potentially make spike sorting more challenging. We simulated recordings in a detailed large-scale cortical microcircuit model to link spike sorting accuracy to neuronal heterogeneity, evaluate the performance of state-of-the-art spike sorters and examine how spike sorting impacts the retrieval of information encoded in the cortical circuit. We found that modern spike sorters accurately isolated about 15% of neurons within $50\,\mu m$ of the electrode shank, which contrasts with previous simulated yields but agrees with experiments. Neurons were unresolved because their spike trains were either missed (undersampling) or, when detected, incomplete or merged with other units' (assignment biases). Neuron isolation quality was influenced by both anatomical and physiological factors (selection bias), improving with increased neuron firing rate, spike spatial extent, for neurons in layer 5, and excitatory neurons. We exposed the network to various stimuli to dissociate the impact of these biases on its stimulus discrimination ability. Surprisingly, undersampling did not affect discrimination capacity, but selection and assignment biases nearly reduced it by half. These findings posit realistic models as a complementary method to evaluate and improve spike sorting and, hence, brain activity representations.

Keywords: spike sorting quality, Neuropixels, biophysical model, neocortex, neural manifold, information geometry

1 Introduction

Neurons convey and process information by emitting sequences of electric signals called spikes. These can be recorded as extracellular spikes with electrodes positioned in the space surrounding the neurons, with the amplitude and shape of the extracellular spikes unique to each neuron. It is established that brain functions emerge from sequences of spikes generated by large groups of interconnected neurons, called neuronal ensembles [1, 2]. Recent advances in large-scale extracellular recordings with densely packed electrodes have made it possible to capture the activity of neuronal ensembles with unprecedented spatiotemporal resolution and scale. These progresses provide a unique opportunity to deepen our understanding of the principles governing information processing by neuronal ensembles. In the last three decades, the field has advanced from arrays of four electrodes which captured only a few dozen neurons [3] to Neuropixels probes with 384 electrodes [4–6], increasing the size of the recorded region and the density of recording sites.

Spike sorting, the process of detecting and assigning spikes to their source neurons based on amplitudes and shapes of extracellular spikes is challenging, particularly in large scale recordings with dense electrode arrays where the signals from multiple neurons close to the electrodes frequently overlap, a problem called spike collision. The rapidly increasing volume of data has compelled researchers to develop and improve automated

spike-sorting algorithms, achieving state-of-the-art spike detection and neuron isolation with the Kilosort suite of computational algorithms [7]. To address the spike collision problem, the Kilosort sorters rely on template matching: detected spikes are iteratively subtracted from the raw signal after detection based on templates. This helps detect underlying spikes masked by the original spike. The combined hardware and software advances have elicited significant insights into how behavior is controlled by the activity of dynamic neural ensembles [2].

Studies using the Kilosort suite now typically report neuron yields of approximately 200 neurons for a Neuropixels 1.0 probe spanning the entire cortical thickness [1, 4, 8] in mice [4, 6], rats [8] and humans [9, 10] over periods of several days [11]. However, based on the density of cortical neurons and the rate of decay of spike amplitudes with distance to the electrodes one might expect 10 times more neurons to be isolated [1, 12]. For example, the most prominent neurons in the cortex, pyramidal cells, display sufficiently large spike amplitudes within 50 µm of an electrode site to be clearly detected, and so should in theory be easy to isolate using modern clustering methods [1]. It follows that spike sorting should accurately isolate about 800, 1300 and 1800 neurons in humans, rats, and mice respectively, given cortical thickness, neuron densities [13] and the geometry of a Neuropixels 1.0 probe [4]. The reported yields therefore correspond to only 25%, 16% and 11% of the theoretical maximum, highlighting substantial undersampling.

Improvement in spike sorting has been powered by small simulations of extracellular recordings, which provide an objective comparison with "ground truth" spike times of the recorded neurons. These synthetic simulations enabled an increase in the quality and number of isolated spike sorted units (i.e., the neuron yield). To improve the quality and yield of spike sorting algorithms, objective evaluations against "ground truth" have also been conducted using *paired extracellular and intracellular recordings* [8], where fractions of neurons not being isolated, i.e., error rates, as low as 5% have been reported for tetrodes [14]. However, this method is challenging for large dense electrode arrays, and may be limited to a small biased sample of neurons that exhibit the largest spike amplitudes [12]. Hybrid datasets combining spike sorted spike waveforms from ~ 1000 neurons with simulated noise offer another option, again yielding an estimate of errors on the order of 5-10% [7, 15]. However, since the dataset is based on spikes detected from spike sorting and is manually curated [7], it may lack the full diversity and complexity of extracellular spikes *in vivo*, limiting the accuracy of the evaluation. In contrast, full synthetic simulations from a subset of 250 biophysically-detailed cortical neuron types, with independently generated spike trains [16, 17] do not rely on spikes obtained from spike sorting. Both hybrid synthetic and full synthetic simulations are flexible, allowing

spiking features and noise levels to be easily tailored to test specific experimental hypotheses. However, despite allowing for greater control, these models may lack biophysical detail and do not capture the scale of real microcircuits, the diversity of spike waveforms arising from neuronal morphological heterogeneity [1] and the variability in spike firing patterns. Additionally, they do not incorporate synaptic connectivity, potentially failing to accurately replicate the statistics of temporally overlapping spikes [18]. These simplifications may result in inflated sorting accuracy and yield in synthetic extracellular recordings, which could explain the disparity between the experimental yield and the neuron yield predicted by the synthetic datasets.

More specifically, the discrepancy in neuron yield could be attributed to systematic biases in spike sorting algorithms. Indeed, the evaluation of spike sorters on synthetic ground truth datasets has revealed systematic biases, which can compromise the quality of isolated spike trains and how well they reflect true neural activity [16, 19]. For example, overmerging the spike trains of multiple neurons into a sorted unit or oversplitting a neuron's spike train across multiple sorted units [19] can obscure the information content in both individual neuronal and collective population responses [20, 21]. Additionally, background noise may produce spurious sorted units [7, 16, 20]. These biases could be explained by the sensitivity of spike sorters to certain anatomical and physiological neuron features, which, if insufficiently characterized due to lack of a realistic ground truth, may reduce yield. Such biases may preferentially sample certain neuronal populations and overlook others, such as sparsely firing neurons [6, 21, 22]. Such sampling biases may distort critical information about neural activity [23], and obscure the true nature of the neural code at both single neuron and population levels [20, 21, 23]. Furthermore, mean firing rates estimated from sorted spike trains are often used for the calibration of models of brain circuitry [24–27]. Biases may degrade these estimates, leading to less realistic models. Without realistic ground truth datasets to better characterize how spike sorters handle anatomical and physiological neuron features, these limitations remain particularly pronounced in large-scale, dense recordings where manual curation of millions of spikes from hundreds of neurons is impractical. A detailed network model with high biological validity could support automated curation aimed at minimizing spike sorting errors and enable comparisons between ground truth neuronal population representations and those derived from spike sorting.

Here, we tested whether modern spike sorting algorithms are biased towards certain anatomical and physiological neuron features in a biophysically-detailed model of an entire cortical column of the rat primary somatosensory cortex [28, 29], providing a new ground truth for improving spike sorting algorithms. The biophysically-detailed model was constructed in

a bottom-up manner, i.e., by carefully modeling and independently validating the individual components in separate in depth peer reviewed studies before combining them into a single model [30–36]. As such, extracellular activity emerges from biophysical principles, rather than being generated from a biased sample of waveforms. Neurons of diverse morphological and electrical classes were accurately distributed in accordance with anatomical laminar density profiles and modeled as fully extended multi-compartmental models with their electrical models fit and validated against single cell protocols. The simulated extracellular activity emerges from a generative model that captures the full heterogeneity of cortical tissue. During calibration of background input and validation of emerging population activity, potential overestimates of mean firing rates from sorted spike trains were taken into account. This enabled layer-wise responses of excitatory and inhibitory populations to single whisker deflection stimuli to closely match *in vivo* references on a millisecond time scale. Here we validated that the model produced realistic *in vivo* extracellular recording traces, spikes, and sorted neuronal firing statistics.

We evaluated the yield, accuracy, and sensitivity to neuronal features of the complete suite of Kilosort algorithms: Kilosort (1, 2, 2.5, 3, 4) [7]. As a baseline reference we also assessed herdingspikes, an alternative spike sorter also designed for large-scale multi-electrode arrays [37]. We found that the number of units sorted from the simulated data was comparable to yields reported *in vivo* for all tested spike sorters, but also that it represented only 15% of the theoretical yields, i.e., of neurons within 50 µm of the electrode shank. Most isolated neurons were either incomplete or merged with multiple units. Spike sorting accuracy was ten times lower in the biophysical models than in the less detailed full synthetic simulation [16], demonstrating a substantial effect of neuronal anatomical and physiological features on spike sorting. Regressing sorting accuracy against these features revealed a positive influence of small electrode distance, high firing rate, large spike spatial extent, excitatory neuron types, and cortical layer 5. This was associated with a pronounced over-representation of layer 5 pyramidal cells compared to other layers across all spike sorters, while interneurons were occasionally over-represented with older spike sorting methods.

We demonstrated that some of these biases significantly affected the recovered representations of stimuli. Biased sampling reduced a linear classifier's ability to distinguish between eight simulated stimulus inputs to the microcircuit based on population responses by nearly 50%, while spike assignment biases lowered performance by 30%. Surprisingly, the significant undersampling observed did not impact the circuit's stimulus discrimination ability. Overall, these findings suggest that while spike sorting preserves a substantial amount of stimulus information, it also produces an imperfect representation of neural activity. Based on the generated ground

truth data, we further provide a model that automates the curation of high-quality single neurons based on their spike features, achieving nearly 90% precision and 60% recall. Our openly available dataset, models, and analyses provide a biophysically realistic foundation for improving and automating spike sorting as datasets continue to expand.

2 Results

2.1 Simulation setup

To address the lack of biological detail of current synthetic simulations [7, 16], we used a detailed and heterogeneous model of cortical tissue incorporating diverse neuronal morphologies, firing patterns, and synaptic connections [28, 29]. The model extends the work of [30] to an atlas-based model of the full non-barrel primary somatosensory cortex. We simulated a 400 µm diameter sub-volume comprising 30,190 neurons distributed over all six cortical layers. Neurons were connected by a total of 36.7 million synaptic connections, with each neuron associated with a reconstructed morphology. The model captures a significant degree of biological variability in 60 morphological types (18 excitatory and 42 inhibitory, Fig. 1a), 11 firing classes (Fig. 1b), 30,190 neurons with unique reconstructed morphologies (Fig. 1a) and morpho-electric combinations (Fig. 1b).

To create realistic electrical signals in the extracellular space surrounding neurons, we used a well established approach, which divides neurons into smaller sections ($\sim$ 500 compartments per neuron on average) and calculating the contribution of each section to the electric signals in extracellular space [38] (Fig. 1c). This allowed us to describe in detail how different parts of the neuron - like the cell body (somata), the branches (dendrites), and the starting point of the axon - shape the electric field, that is, the electric signals in extracellular space. In these detailed simulations, we calculated the extracellular electrical signals by modeling the dendrites and the axon initial segment as lines emitting currents into the extracellular space (Fig. 1c, line-source equation [39, 40]), and the soma as a single point (see point-source equation in Methods). This generative model allowed us to simulate extracellular recordings at the sites of electrodes on different electrode probe designs used in real-world recordings (*in vivo*).

A recent study found that improving the spatial resolution of recorded electric signals around neurons can enhance spike sorting in mice by detecting small neurons with small electric fields (small spatial footprints) [6]. To explore the effect of electrode density on spike sorting accuracy, we tested two types of recording devices (probes): one with wide coverage and electrodes closely packed together, the Neuropixels 1.0 probe [4] (Fig. 1d), and another with more closely spaced electrodes but a smaller overall area, the dense probe introduced in Horvath et al., [41] (Fig. 1e). In this study, we did not use

the probe configuration as in [6], with both full coverage of the cortical thickness and high electrode density because of its high computational requirement. The Neuropixels 1.0 probe was made of a single shank with 384 square electrodes arranged in 4 staggered columns and distanced at $26\mu m$ (bottom, Fig. 1d), as described in [4]. For this probe, we simulated recordings and compared them against Neuropixels data from the anesthetized rat from [8]. The denser probe (bottom, Fig. 1e) had more restricted coverage of the cortical column. For this probe, we simulated recordings at three depths (top, Fig. 1e), replicating the setup used in the anesthetized rat study by [41], reconstructing their electrode locations as closely as possible (Supplementary Fig. 1c). The probe had 128 electrodes arranged in 4 columns, horizontally and vertically separated by $22.5\,\mu m$. To our knowledge, these were the only two publicly available dense recording datasets from the rat primary somatosensory cortex. Using identical probe configurations as *in vivo* enabled us to directly compare/validate the model's extracellular activity against experimental data and assess how spike sorting performance depends on electrode layout and spatial resolution. However, it should be noted that we did not attempt to simulate the exact experimental conditions of those references (such as anesthesia), hence differences in the dynamic states were expected.

We simulated both spontaneous and stimulus-evoked neural activity (Fig. 1f) to assess the robustness and accuracy of spike sorting across different activity regimes (Fig. 1g). The model included thalamic input fibers, which synapse onto cortical neurons with realistic layer-wise innervation profiles and were used here to apply whisker-like stimuli by sending realistic spike trains down a subset of fibres (Fig. 1g-d). In the spontaneous condition, the thalamic input fibers remained unused, and neurons fired at resting firing rates. In the evoked condition, stimuli were repeatedly delivered through the thalamic input fibers to mimic higher and more structured, activity-driven firing. We used 360 distinct *stimulus patterns*, each associated with a subset of the thalamic fibers. Each pattern was activated 50 times every 200 ms before moving onto the next pattern (red circles in schematic).

To evaluate whether using a more detailed and heterogeneous representation of cortical tissue had an impact on spike sorting performance, we compared sorting accuracies and yields between the biophysical simulations and the simpler full synthetic ground truth simulation described in [16], referred to here as the "synthetic model".

2.2 Model validation

Spike sorters' ability to detect and isolate a neuron's spike depends on several features of the recorded trace. It increases when the spike amplitude is large relative to other spikes of smaller amplitude produced by neurons at about $150\,\mu m$ (multi-unit activity) from the

electrode [1]. It also increases when spike amplitude is large relative to the amplitude of the background electrical signal produced by neurons located beyond $200\,\mu m$ (the background noise) [1]; i.e. when the spike has high signal-to-noise ratio. We thus assessed the validity of the Neuropixels and dense recordings simulations by comparing these features of the generated traces, the shape of its extracellular spikes, and the firing statistics of the sorted spikes to those described in the Marques-Smith Neuropixels data [8] and the Horvath dense probe data [41] respectively. However, we did not aim to replicate the dynamic states from those studies, so an exact match was not anticipated.

2.2.1 Validation of recording traces

As *in vivo*, the preprocessed traces contained spikes with high signal-to-noise ratio, multi-unit activity, and background noise (Fig. 2a-f). Single-unit waveforms were apparent at nearby sites (Fig. 2g-l) and exhibited correlated spatiotemporal dynamics similar to *in vivo*. As expected, the denser probe (Fig. 2k, l) sampled the electric field with higher spatial resolution than the Neuropixels probe (Fig. 2g-j).

Spikes with low signal-to-noise ratio (SNR) pose significant challenges for spike sorters, as their amplitude and shape are often obscured by background noise. To enhance the fidelity of the biophysical models, a model of external sources of noise was fitted layer by layer, and the trace extrema were adjusted to match *in vivo* (see Methods). We then assessed whether their SNR distributions aligned with *in vivo* ranges and reflected layer-specific variations across cortical tissue. Spike SNR is typically defined as the peak amplitude of the sorted spike divided by the amplitude of the background noise. To obtain a measure of SNR that is not biased by spike sorting, we compared the distribution of peak amplitudes of the electrode traces. In both Marques-Smith and Horvath's data, median peak amplitude increased with cortical depth (Fig. 2m, o, $p < 0.001$, Kruskall-Wallis). While the models' median peak amplitudes differed from *in vivo* (most $p < 0.01$, Mann-Whitney U test), all captured the *in vivo* trend ($p < 0.001$, Kruskal-Wallis). *In vivo* background noise also increased with cortical depth ($p < 0.001$, Kruskall Wallis, Fig. 2n, p), a trend reproduced by the biophysical models ($p < 0.001$, Kruskall Wallis). We also compared the amplitude-to-noise ratio (ANR), calculated by dividing the trace amplitudes by the median absolute deviation (MAD). The distributions of ANRs observed *in vivo* and in the models showed a wide dynamic range, with alignment of data points between -10 and 10 times the noise level (Fig. 2q), particularly in layers 1, 5, and 6. Despite deviations from *in vivo* due to difference in peak amplitudes, simulated ANRs remained within the *in vivo* range. These findings demonstrate that the biophysical models effectively capture key aspects of ANR distributions, including their

dynamic range and layer-specific trends, despite minor deviations in peak amplitudes.

A large number of spikes characteristic of multi-unit and single-unit activity should increase the number of temporally overlapping spikes making it harder for spike sorting to isolate spikes from each other. Importantly, we ensured that the level of spiking activity was within the range observed in Marques-Smith and Horvath's data. Particularly, power and frequency scaling provide two indirect measures of spiking activity that do not depend on spike sorting, offering an unbiased estimate of how many spikes should theoretically be present in the data. We decomposed the voltage trace into its constituent frequencies and compared the amount of frequencies (power) associated with single spikes (AP power) and multi-unit activity in all simulations to the real data. We also measured the scaling slope α of the distribution of frequencies as it quantifies the relative contributions of frequencies associated with multi-unit activity and single spikes to the voltage trace. In both the simulated and real traces, power decreased with increasing frequency in the frequency band of spiking activity (0.3 - 6 KHz) and followed a power law in all layers, as reported in [42](Fig. 2r). We defined the AP power [43] as the power of frequencies associated with spikes within the 0.3 - 2KHz frequency band. If all spikes had the same shape and amplitude, and did not overlap throughout the recording, AP power would be directly proportional to the number of spikes [43]. The biophysical models' median power in the frequency band of a spike waveform and the scaling slope in the spiking band also increased with cortical depth as *in vivo* (2s-v, all $p < 0.001$, Kruskall Wallis). The biophysical models reproduced *in vivo*'s AP power and frequency scaling for the Neuropixels probe better than it did for the dense probe, which displayed much weaker power and scaling slope than *in vivo* (2u, v). Nonetheless the dense probe's AP power and frequency scaling remained within *in vivo* range.

In Horvath's data but not in Marques-Smith's, the traces were dominated by high-amplitude slow (0.5 - 1.5Hz) and δ waves (1 - 4Hz) characteristic of slow-wave sleep [44] and visible to the naked eye across layers (Supplementary Fig. 2a and d). The biophysical models did not replicate these oscillations (Supplementary Fig. 2b, c, e) under the current parametrization but are able to do so with alternative meta-parameter combinations (results not shown). We assumed that these differences had a negligible effect on spike sorting, as the traces are typically high-pass filtered above 300 Hz to eliminate slow oscillations prior to spike sorting.

2.2.2 Validation of single unit spike waveforms

Crucially, spike waveform variations - such as reduced amplitude [1], broader spikes [45], and increased distortion with distance from the soma [1, 45] - can significantly affect the ability of sorting algorithms to detect spikes. If these sources of variability are not adequately represented in synthetic models, they may contribute to the yield discrepancy and systematic biases observed in spike sorting. Past experimental studies have identified five spike temporal profiles on single electrodes with shapes that are consistent across species (in humans, see panel Fig. 3 copied from [9], and in cats, see panel Fig. 3b copied from [46]). Although the most recently released Kilosort4 sorters has been shown to outperform all other sorters [7], we adopted the same approach as in [9] for direct comparison: both the simulated data and Marques-Smith's data were sorted with Kilosort3, and spike waveforms were extracted using spike-triggered averaging. The simulated temporal waveforms qualitatively matched the five shapes reported in cats [46] and humans [9] (Fig. 3c vs. a-b) and their variation with electrode location aligned with previous simulations [45]. Spike waveform variation exhibited three features with distance to the somata: redistribution from negative to positive components, decreased peak-to-peak amplitude, and widening of spike amplitudes (Fig. 3d). The peak-to-peak amplitude of the spikes rapidly decayed between nearby electrodes, dropping by more than half within 75 µm (Fig. 3e), therefore making the summed extracellular potentials difficult to isolate at 75 µm from the somata. The extracellular traces was strongly distorted by noise beyond 200 µm (Fig. 3f) in agreement with [1].

We further validated that the biophysical models not only replicated the general features of spike waveform variability but also captured the diversity of spatiotemporal profiles observed in *in vivo* data. We used *WaveMap*, a new machine learning tool designed to group sorted spikes based on the shapes of their waveforms across different electrodes [9, 47]. We then compared the spike profiles identified to those discovered in the Marques-Smith's dataset (Fig. 3f). Consistent with the experimental data, the algorithm identified six distinct profiles from the simulations (Fig. 3f, g). These profiles were visible on the isolated extracellular traces of both model excitatory (Supplementary Fig. 3a) and inhibitory neurons (b), and the shape of the waveforms varied with electrode locations.

2.2.3 Validation of firing rates

Firing rate is a key factor influencing spike sorting accuracy [7, 16]. To assess its variation across cortical layers and how well simulations replicate *in vivo* firing rate distributions, we used Kilosort4, the most advanced spike sorter [7] to ensure the comparisons reflected the highest achievable match. To improve unit isolation, Kilosort4 leverages the fact that neurons have a refractory period — a brief interval (1–5 ms [7]) following a spike during which they cannot fire again. It objectively differentiates 'single-units' from 'multi-units' by classifying sorted units with minimal violations of the neuronal refractory period as 'single-units' (a "contamination rate" less than 0.1, see Method). We compared the firing

rate distributions of 'single-units' across three datasets: Marques-Smith Neuropixels recordings [8], biophysical simulations of Neuropixels recordings, and Buccino's full synthetic simulations of Neuropixels recordings [16] (Fig. 4). We excluded comparisons involving the denser probe simulations with *in vivo* data because their lower peak amplitudes (Fig. 2O), spiking activity levels (U), and composition (V), although within the *in vivo* range, could skew sorting yield and quality. We analyzed the first ten minutes of each recording, which was the longest common recording duration. Both simulated and *in vivo* data showed lower single-unit yields in superficial layers, with significantly higher yields deeper layers, prompting a focus on layer 5 and 6. *In vivo*, the yields in these layers were at least 20 times that in layer 4, and no units were found in layers 1 and 2/3, a trend also reflected in the simulated data (Fig. 4a, see unit count).

The biophysical simulations successfully captured the major trends in firing rate statistics, despite minor differences. Like Marques-Smith's sorted unit firing rates, the firing rates from the spontaneous (red dots, Fig. 4a) and evoked (orange dots) biophysical simulations were distributed log-normally across layers (red and orange lines). The overall median firing rate averaged over layer 5 and 6 differed but only by a small margin (Marques-Smith: 2.8 $\pm 1\%$ (95% CI) spikes/secs, spontaneous biophysical simulation: 1.2 $\pm 0.2\%$, evoked simulation: 0.7 $\pm 0.1\%$ and Buccino's full synthetic simulation: 5 $\pm 0.4\%$ (95% CI)). Specifically, the biophysical simulations had lower median firing rates (Fig. 4b, $p < 0.001$, Mann-Whitney) and distributions were narrower (c, $p < 0.001$) than *in vivo* across layers. For reference, the synthetic model distribution also had a lower variance in layer 5 (all $p < 0.01$) but it displayed higher median firing rate ($p < 0.01$).

Older spike sorters also tended to produce firing rate distributions with lower variances compared to *in vivo* data, indicating less variability in the level of activity of the isolated units. While newer sorters like Kilosort3 produced log-normal distributions (Supplementary Fig. 4a) that better matched *in vivo* data (b, $p > 0.15$), older sorters often generated bimodal distributions with peaks aligned between *in vivo* and all three simulations. Across both spontaneous and evoked conditions, firing rate medians (b, $p < 0.05$) and variances (c, $p < 0.01$) in simulations tended to be reduced, highlighting a consistent but marginal gap in capturing the full range of neural activity seen in *in vivo* recordings.

Overall, these shared trends demonstrate that the biophysical simulations closely capture key aspects of *in vivo* variability, despite the absence of biological inputs from non-simulated regions that may account for lower overall activity.

2.3 Model predicts low yield, quality and biases

High-density probes typically allow the isolation of approximately 200 neurons per probe [1, 4, 6, 8–10]. After validating the biophysical simulations, we tested whether spike sorting produced comparable yields to those observed *in vivo*. To enable a fair comparison with the *in vivo* yield, we discarded signals from electrodes located in the white matter: counting only 'single-units' isolated by Kilosort4 on electrodes in the cortex in Horvath and Marques-Smith's data. Having access to ground truth data also permits evaluating the quality of the isolated units, to characterize systematic spike sorting biases and to improve the isolation of high-quality units.

The spontaneous biophysical simulations produced a yield of 'single-units' that more closely matched *in vivo* data compared to the synthetic model, despite producing more sorted 'single-units' than *in vivo*. We found that for spontaneous activity recorded with Neuropixels and sorted by Kilosort4, on average 41% of sorted units were 'single-units'. This proportion differed between experiments (Fig. 5a, $\chi^2(4, N = 5,241) = 43.9, p < 0.001$). At least twice more 'single-units' were detected from the simulations than *in vivo*, with the largest difference being 206 vs. 93 for the dense probe (Fig. 5a, 182 and 193 vs. 96 for Neuropixels). Yet the fractions of 'single-units' were similar ($\chi^2(2, N = 1,212) = 2.0, p = 0.37$ for Neuropixels, $\chi^2(1, N = 920) = 1.8, p = 0.18$ for the dense probe). In contrast with the *in vivo* yield, the majority of units sorted from the synthetic traces were 'single-units', four times more than *in vivo* (S: 377 vs. 163, $\chi^2(1, N = 740) = 4.0, p = 0.042$). The greater similarity of the biophysical model to *in vivo* yield could be attributed to differences in spike timing statistics. In the synthetic model, spike timing was generated with a independent Poisson process while they result from synaptic connectivity in the biophysical models, which will produce more spike collisions.

The model predicts that the number of sorted units should increase at least three times in a stimulus-evoked regime. Although the ratio should strongly shift to 'multi-units', a key prediction is that the single-unit yield should double.

To understand whether higher-resolution sampling of the electric field and the level of spiking activity influence the isolation of 'single-units', we compared the number and types of units detected by Kilosort4 for the two simulated probe designs and activity regime. Kilosort4 detected more units from the dense probe than the Neuropixels probe (358 vs 200 units and 562 vs. 504 units respectively). These were mostly 'multi-units', which were twice more frequent for the dense probe *in vivo* (M:104, H:265, $\chi^2(1, N = 558) = 6.6, p = 0.01$). The model did not reproduce this effect (NS:182, DS:206, $\chi^2(1, N = 1066) = 0.004, p = 0.9$). Despite a higher-resolution sampling of the electric field, the dense probe did not produce a higher single-unit yield

Fig. 1 Biophysical models and experiments a. Example of modeled inhibitory (left) and excitatory neuron morphologies (right) in per cortical layer [28]. Axon in blue, dendrites in red. **b**. 11 different firing patterns, **c**. Generative model of the extracellular recordings [39], **d**. Bouton distribution across cortical depth (line ascending from panel **g** [28, 30, 48]) assuming one afferent fiber from ventral posteromedial (VPM) thalamic nucleus, overlaid on randomly chosen neurons. Simulated **d**) Neuropixels 1.0 and **e**) Horvath probe [41]. Electrode widths but not inter-electrode distances, are multiplied by 3 compared to the column's scale. In bottom panels, distances are multiplied by 3. **f**. Spontaneous activity regime in which thalamic fibers remained unused and cortical neurons fired at resting state, **g**. Evoked regime in which 360 groups of fibers were activated to simulate neural responses to whisker deflections in 360 directions. Morphologies: DAC (HAC, LAC or SAC): Descending (Horizontal or Large or Small) Axon Cell, NGC-DA (or NGC-SA): Neurogliaform Cell with dense (or slender) axonal arborization, BP: Bipolar Cell, BTC: Bitufted Cell, ChC: Chandelier Cell, DBC: Double Bouquet Cell, LBC (NBC or SBC): Large (Nest or Small) Basket Cell, MC: Martinotti Cell, NGC: Neurogliaform Cell, IPC: Inverted Pyramidal Cell, TPC (or UTPC): Tufted (or Untufted) Pyramidal Cell, TPC:A (or TPC:C): Large (or Small) Tufted Pyramidal Cell, TPC:B: Early Bifurcating Pyramidal Cell, SSC: Spiny Stellate Cell, BPC: Bipolar Pyramidal Cell, HPC: Horizontal Cell, see [28].

Fig. 2 Validation of simulated traces. **a-f**. Voltage traces from a single electrode of a **a**. Neuropixels 1.0 probe in anesthetized rats [8], and **b**. of the biophysical model of spontaneous recordings, both in layer 5 **c**. 10 ms after simulated whisker deflections on the same site), **d**. from a synthetic simulation [16], **e**. recorded with the denser probe [41] (insertion depth 2, Fig. 1a) and **f**. simulated with the biophysical model with the denser probe in spontaneous condition. **g-l**. Multi-site traces, color-matched to **a-f**, are shown for one column of each probe (50 among 96 sites are shown for Neuropixels). Peak amplitude distributions by layer for the **m**) Neuropixels and **o** Denser probes. Distributions of noise levels for the **n**) Neuropixels and **p**) denser probes. **q**. Median distribution of the trace amplitude-to-noise ratios across sites and upward 95% confidence interval (ANR) for 100 bins. Voltage amplitudes were divided by the mean absolute deviation at each site. **r**. Median power spectral densities divided by the total power (PSDs) for the preprocessed traces are shown fitted with power-law functions ($1/f^\alpha$) in the 0-90 Hz (Local field potentials) and 0.3-6 KHz frequency band (spiking activity, black lines). Median power in spike waveform's frequency band (0.3-2 KHz) for the **s**) Neuropixels and **u**) the dense probes. Median α in the 0.3-6 KHz band (spiking) per layer and experiment for the **t**) Neuropixels and **v**) dense probes. Metrics for the denser probe were calculated at depth 1 for L1 and 2/3, at depth 2 for L4 and 5, and depth 3 for L6. Measurements reported in M-V are calculated across electrode sites from the full duration of the recordings. ***, p<0.001, Kruskall Wallis.

Fig. 3 Validation of spike spatiotemporal profiles. a. Five types of spike temporal profiles sorted with Kilosort3 from Neuropixels single-electrodes in the Human neocortex (panel adapted from [49]), **b**. Five similar types found in the cat visual cortex (panel adapted from [46]) and sorted with Kilosort3 in the **c**) biophysically simulated Neuropixels recording in the spontaneous regime. **d**. Isolated extracellular recording of the most active biophysically simulated pyramidal cell in layer 5 at 128 electrode locations organized according to a grid layout. The electrodes and spikes are spatially aligned with the cell morphology (grey) **e**. Extracellular recording of the cell in **d**, including the contribution of all the microcircuit cells. **f**. Clustered *Wavemap* [47] embedding for Marques-Smith's data [8] and for **g**) the biophysical simulation. The average spike waveform of each neuron (lines) corresponding to each identified cluster (colors) are shown in **f** and **g** for the three nearest electrodes with the largest average spike amplitude. n indicates the number of neurons per cluster.

Fig. 4 **Validation of the distribution of firing rates sorted in the simulated Neuropixels recordings with Kilosort4 against Marques-Smith's data**. **a**. Distribution of the firing rates of the single-units sorted with Kilosort4 in Marques-Smith's data (grey dots), the spontaneous (red dots) and evoked (orange dots) biophysical simulations and Buccino's full synthetic simulation in layer 5 (blue dots). The distributions were fitted with lognormal distributions (lines with associated colors), which provided a good qualitative match to the data. **b**. Boxplots of log-transformed firing rates. Firing rates were log-transformed with the decimal logarithm to facilitate the visualization of lognormally distributed data. **c**. Boxplots of the variances of distributions of the firing rates generated from 100 bootstraps of the decimal logarithm of the firing rates. The boxplot notches represent the 95% confidence intervals of the median and variances respectively. Statistical significance of the null hypothesis that the simulated median firing rates and variances are the same as *in vivo* data are reported with ns, *,**,***, which indicate $p <= 0.05$, $p > 0.05$, $p < 0.01$, $p < 0.001$ using the Mann-Whitney U test.

than Neuropixels, which cannot be attributed to a lower amplitude-to-noise (ANR) ratio since the *in vivo* dense recording had the highest ANR (Horvath, Fig. 2e, q).

A recent study has suggested that the longer duration of biophysically-simulated extracellular spikes compared to *in vivo* spikes reduces sorting accuracy by generating more false positives [7]. To address this, they proposed downsampling the simulated traces by a factor of two. We applied this approach but observed only a marginal 6% improvement in single-unit yield (experiment NS-2X in Fig. 5a).

The observed single-unit yields are significantly lower than the number of neurons with sufficiently large spikes within 50 μm of the probes [1, 12]. We quantified the gap between the observed and the theoretical yield. For the biophysical models, we counted the number of units within 50 μm of the probes that fired at least once (theoretical yield). To enable comparison between the two probes, which have different numbers of electrodes per

layer, we report the median yield per electrode averaged over layers. For Buccino's model the theoretical yield was simply 0.65 units per site (250 model units and 384 electrodes). For the biophysical models, the raw single-unit yields were an order of magnitude lower than the theoretical yields and were not significantly different between probes or spike sorters (5b). In Buccino's full synthetic model, Kilosort4 produced more 'single-units' than the number of modeled neurons, indicating the production of redundant units (b, blue). These observations were true for most tested sorters (Supplementary Fig. 5a), except for Kilosort2, which detected a large number of units [16]. The original Kilosort also apparently achieved maximal yield but because it does not curate 'single-units' from 'multi-units', it is possible that it also has low single-unit yield.

Since the simulator provided the actual spike times for each simulated neuron, we were able to assess the

quality of the isolated 'single-units' and identify significant issues with their accuracy. We matched isolated 'single-units' to corresponding *true units* by finding the simulated neuron that maximizes the agreement score of their spike trains [16] (for details, see methods). The agreement score is a similarity metric for pairs of spike trains that reach a value of 1.0 for a perfect match. How close the maximizing neuron gets to an agreement score of 1.0 can also serve as a measure of sorting quality for a given sorted unit. This also allowed us to classify sorted units based on the types of errors that occurred during sorting (Fig. 5c): If, for a single spike sorted unit for example, the second-best matching ground truth neuron has an agreement score close to the best ground truth neuron, it is likely that the unit is a 'merged unit' (blue in c), i.e., representing the union of the spike trains of both neurons. The majority of the isolated 'single-units' were less than 80% accurate and degraded across layers (Fig. 5d, white area) and spike sorters (Supplementary Fig. 5b). They contained spikes from multiple true units ('merged unit') and missed at least 20% of the matched true units spikes (Fig. 5c, green). Kilosort4 did not produce false positives (Fig. 5c,d, black area) in contrast with previous reports [7, 16].

To improve quality, the output of spike sorters is typically manually curated. To automate this arbitrary [50] and laborious process from Kilosort4 outputs, we developed a basic logistic regression model to classify units based on eleven spike waveform features which provide an comprehensive description of a unit's firing statistics and spike waveforms [16]. These features or 'quality metrics' [20] were chosen for their computational efficiency and broad availability across the majority of units (n=182 over 184 single-units for the spontaneous Neuropixels simulation). The model achieved 89% ±2% (95% CI) and 93% ±2% precision and 61% ±3% and 55% ±2% recall in the spontaneous (Fig. 5e) and evoked simulations (f) respectively. These high precision scores demonstrate that the features of the single-units isolated with Kilosort4 still hold valuable information for improved curation. We also introduced a new metric, which we called the "mad_ratio" (see Methods), which better reflected contamination by multiple units, displaying the largest correlation with unit sorting accuracy ($R = -0.46$). The model offered the additional insight that two key predictive features of a good unit are its firing rate statistics (Fig. 5g, including firing rate and range, $p < 0.001$) and the variability of its spike troughs compared to background noise variability (mad_ratio and sd_ratio, $p < 0.001$). These features influenced performance by an order of magnitude more than other quality metrics.

We tested whether we could improve curation by using the full waveform in addition to the unit quality metrics. We reduced the dimensionality of the waveform with CEBRA [51] and trained a k-Nearest Neighbor classifier to classify good and degraded units from unit's

embedding (see Methods). CEBRA is an autoencoder with robust generalization performance conferred by contrastive learning optimization of the InfoNCE loss. It allows for supervised label constraints on the produced embedding, makes no assumptions about the dataset statistics and outperforms most dimensionality reduction techniques [51]. Waveforms were reduced to three dimensions, using a binary vector of 'good' and 'degraded' unit labels as supervised auxiliary variables to shape the embedding. A k-nearest neighbor algorithm was then trained to classify units based on this embedding and their labels. The model achieved better recall (82%, ±3% and 90% ±2% for the spontaneous (Supplementary Fig. 5c) and evoked (d) regimes respectively) but lower precision (61% ±3% and 55% ±3%) than the logistic regression. This lower score cannot be explained by insufficient training (Supplementary Fig. 5e), indicating that spike quality metrics may capture all, or even more, discriminative information than the full waveforms.

2.4 Biological factors of sorting failures

The biophysical models successfully replicate several characteristics of dense recordings, including the low yields observed in vivo relative to theoretical expectations [1, 12], as well as previously documented sorting biases [16]. The biological validity of the models allowed us to explore how factors such as sorting algorithms, probe design, and the anatomical and physiological properties of model neurons impact spike sorting accuracy and contribute to the observed low yields and biases.

Comparing the sorting accuracy of spike sorters between the synthetic and biophysical models showed that increased biophysical validity leads to decreased accuracy. As previously reported, for the synthetic simulation [16] most spike sorters detected most neurons with accuracies ranging from 50% for herdingspikes to 99% for Kilosort2. These values also align with those reported for other similar hybrid datasets [7]. However, the sorters produced similarly low proportions of well-sorted ground truth neurons for the biophysical models (Fig. 6a), on average 15% of the neurons within 50 μm of the electrodes across the simulations (b, $\chi^2(5, N = 3,190) = 4.9, p = 0.43$).

Although the dense probe produced similar single-unit yield as the Neuropixels probe (Fig. 5a-b), it produced much lower sorting accuracy. The dense probe did not cover the entire thickness of the cortical column like the Neuropixels probe, missing portions of layers 5 and 6 (Fig. 1e), so we normalized the number of well sorted ground truth neurons by the number of neurons within the same volume of 50 μm from the electrodes, to obtain a sorting accuracy which avoids layer sampling biases. We found that the dense probe produced five times fewer neurons sorted with above 80% accuracy across spike sorters than the Neuropixels probe (on average 3% vs. 15%). Therefore, we chose to focus on the

Fig. 5 Single-unit yield and quality predicted with Kilosort4. **a**. Ratio of 'single-units' and 'multi-units' sorted on Marques-Smith's data (M), Horvath's data (H), spontaneous (NS) and evoked (E) Neuropixels biophysical simulations, Buccino's synthetic simulation [16] (S), and dense probe biophysical simulation (DS). The effect of shortening spike waveforms in NS was tested by down-sampling traces to match *in vivo* spike duration, maintaining sampling frequency at 40 kHz as in [7]. **b**. Single-unit count per electrode per experiment. Bars: median yield per electrode, averaged across layers; error bars: 95% CI; Colored lines: median theoretical yield per site (NS: 6.9, NE: 8.4, S: 0.65, DS: 5.3), calculated as number of active ground truth neurons within 50 µm per site and averaged across layers. **c**. Single-unit categorization. Good unit: unit with an agreement score (score, see Method) with a ground truth neuron above 80%; Merged unit: unit with an agreement score below 80% but above chance (see Methods) with multiple ground truth neurons; Partial unit or 'redundant' [16]: unit with a score between chance and 80% with a true unit also matched above chance by another ground truth neuron; False positive: unit with a score at chance. **d**. Ratio and count of single-units by quality type (colors), layer (x-axis), and simulation (panels). **e-f**. Precision and recall of a lasso-regularized logistic regression model classifying good vs. degraded units using spiking and waveform quality metrics [16]. Tested on **e**) NS and **f**) NE simulations, cross-validated on 100 bootstrapped train-test samples. Black dots: precision/recall for 25% test units; orange dots: median values. **g**. Regression model weights for the neuropixels spontaneous biophysical simulation. Pink indicates weights that are significantly different from 0; metrics were z-scored to enable the comparison of the weights.

Neuropixels probe, which had high sorting accuracy and had full coverage of the cortical column.

We leveraged the detailed morphological, anatomical, and physiological descriptions of our ground truth model neurons to characterize the impact of neuronal features on sorting accuracy. We used a simple lasso-regularized fractional logistic regression model (see Methods) to predict sorting accuracy. Unlike in the previous section, this predictor used features other than the spike waveform, which are only available in the model. The model provided a good fit to the data (Fig. 6, $R^2 = 0.42$ and $R^2 = 0.34$ in the spontaneous (c) and evoked (d) regimes respectively) and identified seven significantly predictive features out of 21 in both regimes. Accuracy improved as neuronal firing rates increased and spike spatial extent (or 'footprint' [6]) expanded. It was also higher for pyramidal cells compared to interneurons and in layer 5 compared to other cortical layers. Sorting accuracy decreased when neuron's distance to the nearest electrode increased (Fig. 6e,f) in both regimes), was larger for delayed non-accomodating neurons ('dNAC', evoked regime in f) and in layer 6 (spontaneous regime in e). These results demonstrate that, in both regimes, Kilosort4's output is biased toward very active layer 5 pyramidal neurons with large spike spatial extent. Notably, the bias for high firing units led to mean firing rates in most layers being overestimated by a factor of three to four from sorted spike trains compared to the ground truth (Supplementary Fig. 6). These results demonstrate that the low yield is due to neuronal feature sensitivity, which can also distort the circuit's firing rate distribution [12] (Fig. 4 and Supplementary Fig. 6).

We investigated whether spike sorters exhibited selection biases, by over-representing certain model neuron types while undersampling others (Fig. 6g,h). To quantify these biases, we assigned each sorted 'single-unit' the biophysical features of its best-matched ground truth neuron and assessed whether neuron types were more or less likely to appear after spike sorting compared to their representation in the ground truth. Excitatory pyramidal neurons, which made up 91% of the ground truth population (Fig. 6g), were consistently underrepresented across all layers by approximately a factor of 2, except in layer 5, where they were over-represented by the same factor (Fig. 6h, ratios in blue; Fisher's exact test, two-tailed, p < 0.05). In contrast, most interneurons were proportionally represented relative to the ground truth, except for rare interneuron types, which were over-represented by factors ranging from 2.9 to 8.1 by some spike sorters, particularly Kilosort (1 to 2.5).

2.5 Spike sorting degrades stimulus representation

After identifying the nature, sources, and extent of spike sorting failures, we investigated their functional consequences by assessing how these inaccuracies affect the readout of stimulus encoding. We determined whether

and how much the identified biases degraded the information contained in the sorted compared to the true spike trains. We simulated the model's responses to simulated whisker deflections in 8 of the 360 directions, separated by 45 degrees. For each stimulus, responses were defined as the spike count of each unit during the presentation of the stimulus. We reduced the dimensionality of the responses with PCA to enable the visualization of the ground truth population responses in three dimensions (neural latents) and plotted the transformed responses (neural manifolds, Fig. 7a) to the stimulus directions (colors) for each trial (dots).

The ground truth neural manifold exhibited a ring-like geometry (Fig. 7a), which continuously reflected changes in the stimulus direction across trials. Spike sorting distorted the manifold's geometry for all spike sorters, but the most recent sorters still preserved stimulus mapping (from Kilosort2 to 4).

We quantified stimulus information loss by measuring the manifold's information capacity, an information theoretical metric that captures a linear classifier's ability to distinguish between stimulus directions based on population responses [52]. Information capacity measures how much information about stimulus direction is stored per unit dimension. We expressed it as a ratio to the population's lower bound capacity, determined by shuffling the mapping between neural responses and stimulus directions. Since information capacity is sensitive to population size, we applied dimensionality reduction as in [52] to match the size of the ground truth and sorted unit populations. Information capacity remained robust across different numbers of dimensions (Fig. 6), so we used 200 unit dimensions (the default parameter from [52]) across all conditions to ensure a fair comparison. Distortions in manifold geometry were associated with significant reductions in information capacity, averaging 50% decrease across spike sorters, from 25% above the lower bound (Fig. 7b, t(199) ranging from -68 to -24, all $p < 0.001$, t-test).

The observed loss in discrimination capacity was partially explained by selection biases for unit types, but not by random undersampling of model neurons. Undersampling ground truth neurons to match sorting yields neither disrupted the manifold's structure nor its information capacity (example for Kilosort4, Fig. 7d, and all sorters in f, red boxes vs. red line, t(199) ranging from -0.71 to -0.02, all $p > 0.47$, t-test). However, also matching neuron type distribution to that of sorted units visibly degraded the manifold structure and its information capacity by an averaged 21% across sorters (Fig. 7f, white boxes, t(199) ranging from -17 to -8, all $p < 0.001$, t-test).

We evaluated the effect of assignment biases on the information capacity of sorted populations by comparing degraded sorted units ('merged' and 'partial') with good sorted units. Unit type distributions differed significantly between biased and good units across spike sorters

Fig. 6 Sorting accuracy and sensitivity to neuron anatomical and physiological features. **a**. Ground truth neuron sorting accuracy, i.e., the proportion of ground truth neurons (see annotations) sorted with above 80% accuracy within 50 μm of the electrodes, **b**. The total sorting accuracy summed across datasets, i.e., the number of well-detected neurons across ground truths divided by the number of neurons within 50 μm of the electrodes across simulations. For the dense probe, numbers were pooled across depths. Biophysical ground truth neurons' sorting accuracies against best-fitted regularized fractional regression models (red line) for Neuropixels simulations of the **c**) spontaneous and **d**) evoked regimes, sorted with Kilosort4 and their weights in **e** and **f**. The weights in pink are significantly different from 0 and numbers indicate weight amplitudes. **g**. Probability distribution of neuron types defined by their synaptic class, layer and electrical type, in the evoked biophysical ground truth. Number annotations indicates the count of each model neuron type **h**. Ratios of the probability of occurrence of model neuron types in the sorted data to their probability of occurrence in the ground truth. Each sorted single-unit was first assigned the biophysical features of its best ground truth match and we calculated the ratio of ground truth neurons' probability of occurrence after sorting and in the ground truth. The 10-minute evoked simulation at 40 kHz was used in panels **a** and **b**. To link neuron selection biases to stimulus representation quality (Fig. 7), the 1-hour evoked simulation at 20 kHz was analyzed in panels **d**, **f**, and **g-h**. *: $p > 0.05$, **: $p < 0.01$, ***: $p < 0.001$

.

($p < 0.001$, Fisher exact test with Monte Carlo simulations). To disambiguate the impact of the two biases, we resampled degraded units to match the distributions of good units. On average, the information capacity of good units was 37% lower than the ground truth across spike sorters (Fig. 7g, red boxes vs. red line, t(199) ranging from -0.31 to -15, all $p < 0.001$, t-test). Degraded units exhibited even lower capacity, averaging a 12% decrease from good units' capacity across most sorters (Kilosort4, 2.5, 1, and herdingspikes). These findings demonstrate that neuron selection and spike assignment biases strongly reduce information capacity (Fig. 7g, red vs. white).

These reductions in information capacity were primarily explained by increased direction manifold radii, dimensions and more correlated centroids. Except for Kilosort which showed a 3% decrease, direction manifolds' radii increased by an averaged 6% across sorters (Fig. 7h, t(199) ranging from -87 to 0.18, all $p < 0.001$, t-test). Except for Kilosort and Kilosort2 which displayed a 19% decrease in direction manifold's dimension, dimension also increased by an averaged 6% across sorters (i, t(199) ranging from -59 to 234, all $p < 0.001$). A marginal but significant increase in center correlation of 1% was observed for Kilosort3, 2, 1, and Herdingspikes (Fig. 7j). These changes indicate an increasing overlap in neural responses to distinct stimuli, reducing the linear separability of stimulus-specific responses.

3 Methods

3.1 Datasets and simulations

Marques-Smith et al.,'s dataset. We used a 26 minute recording performed with the Neuropixels 1.0 Phase3A probe [4] in the primary somatosensory cortex of anaesthetized adult rats aged between six weeks and eight months [8]. The probe, consisting of four columns with 96 electrode sites each, and rows spaced 20 µm apart, was aligned with a cortical column and spanned all six layers. The first electrode was positioned at the Pial surface. We used the cortical layer border depths from the Pial surface identified in P23-36 adult rats in [8, 53] to determine that layer 5 contained the highest number of isolated units. Traces were acquired at a sampling rate of 30 kHz, and analyses were performed on file c26, accessible at google-drive. The publicly shared recording underwent minimal processing, limited to offset subtraction [8].

Horvath et al.,'s dataset[41]. We utilized a dataset from a single rat, which included recordings at three cortical depths with a single probe insertion. Electric field potentials were recorded at a sampling rate of 20 kHz using an 8 mm long and 100 µm wide silicon probe with 128 dense recording sites, arranged in a 32 x 4 grid of square electrodes. Like Neuropixels 1.0, the electrodes were square-shaped and made from titanium nitride, with low impedance (50 kOhms) in saline, measured with 1 nA sinusoidal currents at 1 kHz, compared to Neuropixels 1.0's 149 pm6kOhms [4]. Unlike Neuropixels 1.0's staggered electrode layout, Horvath's probe features equidistant electrodes. The electrodes are also larger (20 µm per side, compared to Neuropixels 1.0's 12 µm), and electrode and positioned closer together, with a center-to-center distance of 22.5 µm for Horvath versus 25 µm for Neuropixels 1.0. We accurately reproduced the site locations from the experiment in [41] (Supplementary Fig 1). The dense probe captured activity from layers 1 and 2/3 at insertion depth 1, layers 2 to 5 at depth 2, and layer 6 at depth 3.

Synthetic model. We analyzed simulated Neuropixels recordings from [16], sampled at 32 kHz. The simulation included spiking activity from 250 units in layer 5, comprising 200 excitatory and 50 inhibitory biophysically simulated units with spike activity based on data from [30]. Importantly, this model lacks the cortical microcircuit connectivity that drives spiking dynamics and correlation statistics, whereas our proposed biophysical model includes these connections.

Biophysical model. We utilized a model of the rat primary somatosensory cortex, at age P14 [28]. Of the 4,234,929 neurons in the circuit, we simulated the voltage activity of a microcircuit column comprising 30,190 neurons using the NEURON software [54], via Blue Brain Project's Neurodamus interface, [55] under spontaneous and stimulus-evoked conditions. We detected spikes from 1,388 model neurons during the spontaneous activity condition, 1,836 model neurons during the 1-hour evoked condition sampled at 20 kHz, and 1,552 neurons during the 10 minute evoked condition sampled at 40 kHz, within 50 µm of the Neuropixels probe. The network was calibrated to emulate an *in vivo* state of spontaneous firing with low synaptic release probability and asynchronous firing at low rates [29]. All model neurons fired at least once in the first 10 minutes, making detection theoretically possible. Meta-parameters included extracellular calcium concentration (modulating synaptic reliability), noise in external input compensation, and the overall activity level, described as a proportion of layer-specific reference firing rates. The model was configured with parameters $P_{fr} = 0.3$, [Ca] = 1.05, and ROU = 0.4 [29].

Probe reconstruction. We simulated extracellular recordings from 384 recording sites of a Neuropixels 1.0 probe, without accounting for electrode impedance, capacitance, or the effects of electrode shape and size. Instead, the electrodes were modeled as infinitesimal points [39]. For the Horvath setup, extracellular recordings were simulated from 128 electrodes, and the probe was positioned at three different depths in the cortical microcircuit to replicate the three in vivo locations [41]. However, the geometry of our microcircuit prevented us from perfectly matching the electrode distribution across

Fig. 7 Impact of spike sorting biases on stimulus discriminability. **a**. Neural manifolds and **b**) information capacity of ground truth and sorted populations. Neural manifolds for **c**) ground truth, for **d**) a random sample of the ground truth neurons with Kilosort4's yield, **e**. A sample with distribution of model neuron types matched to Kilosort4 sorted population's, **f**. Information capacity for the undersampling **d** (red box) and biased sampling in **e** (white box). **g**. Information capacity for ground truth and good units and degraded units ('merged' and 'partial') in Fig. 5c,d. The distribution of unit types were matched between the two. **h**. Radius and **i**) dimension of the manifold evoked by each stimulus direction, **j**. Correlations of the centroids of these manifolds. Distributions of normalized capacities, radii, dimensions and center correlations were obtained with 200 bootstraps of the populations. All analyses were performed on the 1-hour biophysical simulation of the Neuropixels evoked condition with 20 kHz sampling frequency.

.

layers observed in vivo (Supplementary Fig. 1, Depth 1, Depth 3 had 68 sites in L6, 60 outside the cortex).

Generative model of the electric field. Neurons are modeled as multicompartment cables using the NEURON simulation environment [56]. Extracellular signals are calculated at simulation-time using the BlueRecording pipeline [39]. Briefly, we assume that the extracellular medium can be treated as homogeneous, with a conductivity σ, and effectively infinite relative to the size of the neural circuit and recording electrodes. We also assume that the recording electrode contacts can be treated as infinitesimally small.

At each time step, the simulator calculates the extracellular signal $S(t) = \Sigma_{j=1}^{n} C_j \texttt{i\_membrane}_j(t)$, where `i_membrane` is the total compartmental transmembrane current, j is the index of each neural compartment in the circuit, and C is a precomputed scaling factor.

For simulations of the probes used by Horvath et al., we calculate the scaling factors C_j using the line-source approximation [40]. We assume that each neural compartment can be treated as a 1-dimensional line. For each neural segment, the scaling factor $C_j = \frac{1}{4\pi\sigma_e \Delta_s} \int_{-\Delta_s}^{0} \frac{ds}{\sqrt{(h-s)^2+r^2}}$, where Δ_s is the length of the segment, h is the distance from the near end of the segment to the recording electrode in the axial direction, and r is the absolute value of the distance from the near end of the electrode to the electrode in the perpendicular direction.

For simulations of Neuropixels probes, we calculate the scaling factors C_j using the point-source approximation. We assume that each neural compartment is infinitesimally small. The scaling factors $C_j = \frac{|\vec{r}_j - \vec{r}_e|}{4\pi\sigma_e}$, where $\vec{r}_e$ is the position of the electrode and $\vec{r}_j$ is the position of the compartment.

Sampling frequency. The simulated Neuropixels traces were sampled at 40 kHz in the spontaneous regime and at 20 and 40 kHz in the 1-hour long and 10 minute evoked regimes respectively which are close to sampling frequencies used in past studies [9, 16, 41, 57]. The traces collected with the reconstructed Horvath probe were sampled at a frequency of 20 kHz as in [41].

Fitting amplitude and background noise. The biophysically simulated traces missed noise from external sources and voltage amplitudes ranged from $-1.26\,\mu$V to $0.6\,\mu$V, a narrow range near zero that reduced spike resolution, causing their omission after Kilosort processing, as it assumes 16-bit integers. To address these, we scaled the traces and modeled external background noise as an additive Gaussian distribution with zero mean across all electrodes, following [7, 16]. We ensured biological validity by fitting the trace extrema and background noise to Marques-Smith's data in three steps: applying a gain derived from dividing the extrema of preprocessed *in vivo* and simulated traces, adjusting noise levels measured by the trace mean absolute deviation (MAD), for each layer, and manually fine-tuning the gain and noise after preprocessing. All fits were performed using the Nelder-Mead algorithm. We used the same gain and noise parameters for both spontaneous and evoked regimes. The best-fit gain was 677, with missing noise root mean square values of $2.8\,\mu$V, $3.4\,\mu$V, $3.7\,\mu$V, $3.6\,\mu$V, and $3.8\,\mu$V for layers 1, 2/3, 4, 5, and 6. The spontaneous regime's maximal amplitude ($361\,\mu$V) closely matched Marques-Smith's ($362\,\mu$V), while the evoked regime produced a slightly higher value ($387\,\mu$V). The gain to fit the synthetic traces' extrema to Marques-Smith's data was 0.41.

For the denser probe, separate gains were derived for each depth, and background noise was fitted to Horvath's data [41] by matching depth and layer. Gains were 12,463, 1,949, and 20,778 for depths 1, 2, and 3, respectively. Missing noise RMS values were $27.3\,\mu$V and $30.3\,\mu$V for layers 1 and 2/3 at depth 1, $37.4\,\mu$V and $37.9\,\mu$V for layers 4 and 5 at depth 2, and $35.6\,\mu$V for layer 6 at depth 3. Maximal amplitudes closely matched Horvath's data, with errors below 3.5%.

Preprocessing. For all analyses, the traces were high-pass filtered above 300 Hz as in [4, 6, 7, 9, 16], using *SpikeInterface* [16] to attenuate their low-frequency noise component while preserving spikes' frequency band. The common median voltage across electrodes was then subtracted to remove the global noise component shared across electrodes (common median referencing [58]). Whitening was applied by all Kilosort sorters to reduce local spatial correlations across channels [7].

3.2 Validation

Amplitude-to-noise ratio. To minimize computational cost, we analyzed the first 10 minutes of all recordings from cortical electrode sites. The amplitude-to-noise ratio (ANR) was calculated by dividing the trace amplitudes by the median absolute deviation (MAD). We compared ANR distributions across experiments by using the largest ANR range, divided into 100 bins, and calculated the probability of each ANR bin per electrode and experiment. The ANR was then averaged across electrodes and plotted with confidence intervals across electrodes for each experiment (Figure 2Q).

Spectral composition. Power spectral densities (PSD) were computed with the Welch method [59], at 1 Hz resolution with the Scipy package. Filtering window sizes matched recording sampling frequencies, and PSDs were denoised with a Hann window with a one-second overlap.

Power law fits were performed by fitting a linear regression to the log-transformed PSDs, whose coefficients were the α parameters of the $1/f^\alpha$ power laws.

WaveMap. Single-unit spikes were extracted using spike-triggered averaging. The six nearest electrode sites, which exhibited the largest average spike amplitudes, were selected and ranked in descending order based on the first positive peak, trough (maximum negative amplitude), or second positive peak—depending on which had the largest absolute amplitude. Spikes were aligned at

the trough over a 12 ms window centered on the spike timestamp, later reduced to 6 ms. Waveform amplitudes were normalized to a range of 0 to 1, and UMAP (umap-learn v0.5.3) was applied to project the waveforms into a two-dimensional latent space. The resulting embedding was clustered using the Louvain algorithm (python-louvain v0.16) as in [9, 47].

Spike-triggered average of isolated intracellular recordings were calculated for the active morpho-electric types that were found within 50 μm of the Neuropixels electrodes (Supplementary Fig. 3).

Firing rate. Since firing rate distributions followed a lognormal pattern, we compared simulated and *in vivo* firing rates using their decimal logarithm (log) transformations. A Mann-Whitney U test was conducted to assess whether the medians and variances of the log-transformed firing rates of the sorted units were similar (Figure 4B). To calculate the median and confidence intervals of the variance shown in Figure 4B, we bootstrapped the firing rates 100 times to generate a distribution of variances.

3.3 Spike sorting

Spike sorting was performed on the probe-wired raw recordings using SpikeInterface version 0.100.5 [16]. Sorting generally succeeded with default parameters. For all sorters, we used the default settings, with a few key adjustments: the minimum firing rates for unit detection (*minFR*) and site curation (*minfr_goodchannels*) were both set to 0 to maximize unit detection. This adjustment enabled a fair comparison with sorters like Kilosort 4 and Herdingspikes, which do not filter units based on firing rates. Additionally, batch sizes were adjusted to accommodate GPU memory constraints.

We report results from Kilosort2 for 4 minutes of evoked recordings, as it failed to process longer durations. Kilosort2.5 only functioned without drift correction (with the *nblocks* parameter set to 0). Mountainsort 4 was not used due to its long sorting time, which exceeded eight hours. Klusta and HDSort were excluded because they are not maintained to be compatible with Python versions later than 3.9.

All tested sorters, except for the original Kilosort version and Herdingspikes, isolated 'single-units' and labeled them as either 'good' or 'multi-unit' in the *KSLabel* metadata. For Kilosort and Herdingspikes, all sorted units were reported.

3.4 Sorted unit quality

Unit isolation ratio. Kilosort (2, 2.5, 3, 4) isolate 'single-units' as clusters of spikes with minimal violations of the neuronal refractory period — a brief interval following a spike (1–5 ms [7]) during which neurons cannot generate another spike. Units that do not satisfy this criterion are classified as 'multi-units'. To distinguish between these categories, these sorters calculate the "contamination rate", which measures the proportion of refractory period violations. This metric is defined as the ratio of the frequency of spikes occurring within the refractory period to the baseline frequency of spikes outside this period [60]. A contamination rate below 0.1 is used as the threshold for identifying 'single-units' [7]. The unit isolation ratio was calculated as the proportion of 'single-unit' to the 'multi-unit' produced by the spike sorters. To investigate the impact of waveform duration on unit isolation, we ran Kilosort4 on a biophysical simulation of the spontaneous regime, processing the data at twice the normal speed. This was achieved by downsampling the trace to every second sample while maintaining the original sampling frequency of 40 kHz, as described in [7].

Unit yield. The sorting yield refers to the number of 'single-units' isolated by Kilosort versions 2.0, 2.5, 3.0, and 4.0. This measure of sorting quality is accessible to experimentalists who do not have access to ground truth data. To enable comparisons with *in vivo* studies and across experiments with varying cortical column coverage, we employed the yield per site metric.

Agreement Score. We utilized the agreement score defined in [16], which is the ratio of correct spike coincidences between a ground truth neurons and a sorted unit to the total number of spikes.

$$A = \frac{N_{tp}}{N_{tp} + N_{fp} + N_{fn}} \tag{1}$$

where N_{tp} is the number of true positives (detected spikes), N_{fp} is the number of false positives (spurious spikes), and N_{fn} is the number of false negatives (missed spikes). A score of one hundred percent can only be achieved when all spikes from the ground truth and the sorted units coincide.

Coincidences were detected within a time window of Δ_t = 1.3 ms around each ground truth spike. This duration maximized sorting accuracy for Kilosort4 (Supplementary Fig. 3A). To ensure that a Δ_t of 1.3 ms was not overly lenient—potentially assigning high scores for random matches, we compared the distribution of ground truth neuron sorting accuracies for zero shift versus long constant shifts of ground truth spike timestamps. When Δ_t becomes too long, the accuracy distribution will show a single peak near 0, indicating chance-level performance. For meaningful Δ_t, the distribution will display two peaks: one for correctly detected units and another near chance level for missed units.

In our simulation, a spike waveform is timestamped when the voltage reaches $-10\,\mu V$, while spike sorters may timestamp the spike at a different point in the spike waveform. As we expected, for small shifts, the accuracy distribution for ground truth neurons showed two distinct peaks. However, with larger shifts (e.g., 30 ms), the accuracies degraded and peaked at 0, indicating chance-level performance (Supplementary Fig. 3B). Based on this analysis, we finalized $\Delta_t = 1.3$ ms with no shift.

Coincidences between two independent spike trains can occur by chance, resulting in non-zero agreement scores. We assumed that the sorted unit's spikes occurred independently and stochastically with a constant firing rate ν. Under this assumption, the probability of observing at least one coincidence (a true positive tp) within the interval surrounding the ground truth spike ($T = 2 * \Delta_t$) is given by the homogeneous Poisson equation:

$$p_{chance}^{tp} = 1 - \frac{\lambda^k e^{-\lambda}}{k!} \qquad (2)$$

where $k = 0$ and the term $\frac{\lambda^k e^{-\lambda}}{k!}$ is the probability of observing no coincidence. λ is the expected number of coincidences within the interval T and for a firing rate $\nu = min(N_{true}, N_{sorted})$ taken as the firing rate of the least active unit between the ground truth and the sorted unit:

$$\lambda = \nu T \qquad (3)$$

Chance agreement scores A_{chance} were calculated following equation 1 for each pair of ground truth neuron and sorted units by setting the number of true positives to $N_{tp} = \nu * p_{chance}^{tp}$, the number of false positives to $N_{tp} = N_{sorted} - N_{tp}$ and the number of false negatives to $N_{true} - N_{tp}$.

Sorting accuracy. We defined the sorting accuracy of a ground truth neuron as the highest agreement score achieved with its best-matching sorted unit.

3.5 Model selection

Logistic regression model for classification. The logistic regression model was a generalized linear model for the Binomial family and the logit link (statsmodel v0.14.3). The models' precisions and recalls for more than 80% accurate sorted single-units (high-quality units) were calculated by splitting the dataset into an 80% train set and a 20% test set randomly sampled a hundred times (number of folds). We selected the unit quality metrics that were the most efficient to compute and that could be computed for the largest number of units (n=182 units) to maximize the size of the training dataset. The quality metrics were z-scored before training and testing, which allowed to compare their weights from the fractional logistic model. Z-scoring did not noticeably change the precision and recalls of the models.

cebraspike model. We used 40 iterations which were sufficient for the convergence of CEBRA's loss functions for all cross-validation groups of samples (the folds). We used the smallest possible number of iterations to minimize the large computational cost of running 100-fold cross-validation. We manually searched for the best set of hyperparameters, by changing one parameter at a time. We changed the number of spike waveforms (25 and 50) in the dataset, CEBRA's output dimensions (3 and 10) and the k-NN's number of neighbors (1, 2 and 10, 20).

Regularized fractional regression model. We trained a generalized linear model for the Binomial family and a logit link (statsmodel v0.14.2) to predict ground-truth unit sorting accuracy based on 21 available descriptive features. The model weights were regularized with Lasso (L1-norm) which enables feature selection by setting the weights of linearly dependent unit features with high variance, which effect on sorting accuracy can not be disambiguated, to zero. We ensured model convergence by increasing the number of iterations (maxiter parameter) and reducing the loss function convergence tolerance (cnvrg_tol) until the models' cross-validated explained variances (R^2) became stable. The models converged for with 100 iterations and a convergence tolerance of 1e-10.

3.6 Manifold information geometry

We used information geometric metrics to quantify the impact of the identified sorting biases on the capacity of the sorted populations to perform a coarse stimulus discrimination task: the metrics were calculated for 8 stimulus directions separated by 45 degrees out of the tested 360, to manage computational cost. We used the neural_manifolds_replicaMFT package to calculate the average mean-field theoretic manifold capacity, radius and dimensionality for each category of unit bias (available on Github, [52]). Because manifold capacity changes with the dataset size, the number of ground truth, and sorted units in each sorted unit class, dimensionality reduction was applied to all datasets to enforce the same number of unit features, set to 200 (the default parameter n_t) and capacity was divided by the lower bound capacity of each dataset; the lower bound can be empirically calculated from a shuffled label dataset [52]. Medians and confidence intervals were obtained by sampling each dataset 1000 times. The model's number of samples n_t was set to 200 (its default value), the margin κ to zero.

Software

We used Spikeinterface 0.100.5, ProbeInterface, Kilosort3, Python and Matlab. The Fractional regression model was implemented with Statsmodel in Python. All Kilosort sorters were run with Matlab R2022b with the Parallel Computing Toolbox (Version 7.7), Signal Processing Toolbox (Version 9.1) and the Statistics and Machine Learning Toolbox (version 12.4). Extracellular signals were calculated using an experimental version of the BlueRecording [39] pipeline. Source code and instructions for precomputing the scaling factors for each compartment are available on GitHub. The codebase for the analyses is available on GitHub.

3.7 Hardware

Biophysical simulations. We run simulations parallelized on 120 cpu clx nodes (384 GB of memory: 12

DIMMs (6 per CPU) of each 32GB at 2933MT/s 2 Intel Xeon Gold 6248 CPUs at 2.50GHz with max speed of 3.9Ghz). Each CPU has 20 physical cores. Due to hyperthreading, slurm provides 80 cores per node scheduled with Slurm. With this setup a simulation of 10 minute of extracellular recordings typically took 6 days.

Spike sorting. Spike sorting was supported by GPUs (4 volta v100 skl each with 16GB of RAM) on nodes with 768 GB of memory: 12 DIMMs (6 per CPU) of each 64GB at 2666MT/s, 2 Intel Xeon Gold 6140 CPUs at 2.30GHz with max speed of 3.7Ghz). Each CPU has 18 physical cores => 36 physical cores in total. Due to hyperthreading, slurm provides 72 cores per node.

Parallelized computing. Biophysical simulations were performed on a computer cluster with resource allocation managed by SLURM and Python multiprocessing framework was used to speed up preprocessing and data analysis.

4 Discussion

We simulated dense recordings from a large-scale, detailed model of the rat cortical microcircuit designed to capture neuron heterogeneity in a data-driven and biophysically-detailed manner. Our goal was to test if spike sorting sensitivity to neuronal biological features could explain the large gap between the yield of modern spike sorters and the theoretical yield in a data-driven and biophysically-principled manner and characterize the theoretical impact of this. We found that a fractional regression model, where several model neuron features strongly impacted spike sorting accuracy, effectively accounted for the proportion of spikes correctly detected and assigned to isolated units. The model accounts for the yield gap by predicting reduced detection and isolation of neurons within isolable distances, influenced by the combination of electrode distance, layer-specific location, morphology, synaptic and electrical properties. The model further predicts that most presumably isolated units are sorted with accuracies below 80%, include spikes from multiple neurons and/or are incomplete and that all spike sorters display selection biases for particular unit types. Yet despite these biases, the most recent spike sorters preserved the overall low-dimensional shape of stimulus-evoked responses as well as most stimulus information content.

In agreement with evaluations reporting sorting accuracies above 80% on synthetic datasets [7, 16, 21], most tested spike sorters achieved high yields, accurately assigning spikes to over 85% of the neurons in synthetic datasets. However, in contrast to this predictions but in agreement with reports from dense extracellular recording experiments [4, 8–11], the spike sorters isolated about 200 neurons on the biophysical models — only a tenth of the achievable yield predicted by theoretical models [1, 12]. This comparison between synthetic model yield and theoretical yield was enabled by the scale and high level of biophysical detail of the model. It closely reproduces the average neuron density distribution across the full thickness of the rat primary somatosensory cortex [30]. The cortical microcircuit comprises 30,000 morphologically detailed neurons, captures cortical neuron diversity in 60 morphological types, grouped into inhibitory and excitatory synaptic types and 11 electrical neuron types, spanning all six cortical layers and receiving inputs from simulated thalamic fibers, which have been extensively validated against *in vivo* data [28–30]. It also features realistic synaptic connectivity and short-term plasticity which constrain firing synchrony, critical for spike sorting [18], and are not modeled by current synthetic datasets. The model is also grounded in a biophysically principled description of spike amplitude decay with distance [38, 39].

We demonstrate that the extracellular traces simulated from our detailed model replicate key characteristics of two published experimental datasets from the rat primary somatosensory cortex [8, 41]: they exhibited spiking components, spectral composition, amplitude-to-noise ratios that were similar and/or in the *in vivo* range of these datasets. They contained spikes with five temporal profiles that matched those reported in these [8] and other studies in humans [9] and cats [46] and the dependencies on electrode distance and location reported in yet another set of studies [1, 12, 45]. Extracellular spike amplitudes remained resolvable within 75 microns and disappeared by 200 microns, in line with prior research [1] and ruling out the possibility that the low yield is attributed to a non-realistic rapid decay of spike amplitude with distance. The spike power's tendency to increase with cortical depth may come to further validate the model's accurate description of the distribution of neuron density and activity level across layers, as spike power has been shown to reflect neuronal density and activity across cortical thickness [43].

The biophysical models did not replicate the slow oscillations observed under anesthesia with the current parametrization; however, they are capable of generating activity resembling other brain states under alternative meta-parameter combinations. This flexibility makes them suitable for quantifying the impact of different brain states on the quality of spike sorting in the cortex [61].

Although the biophysical simulations produced spike waveforms that were longer than those observed *in vivo* — a factor known to potentially degrade spike sorting accuracy and increase false positives [7] — the resulting accuracy losses were minimal. We further demonstrated that reducing the waveform length by half through subsampling the traces, as suggested by Pachitariu et al. [7], did not significantly alter the number or proportion of isolated 'single-units'. Additionally, for the non-subsampled traces the unit yield was already statistically consistent with the Marques-Smith's dataset [8]. Finally, Pachitariu et al. [7] reported false positives as the main

error resulting from longer waveforms, while we observed very few false positives in our results.

We developed a fractional regression model that succinctly describes and quantifies the relationship between cortical neuron location, biological features, and the proportion of correctly isolated spikes. Our findings reveal that, aside from distance, a sparse firing rate is the strongest determinant of low sorting accuracy, corroborating other studies that highlight the dependence of sorting accuracy on firing rate [7, 16]. This also explains previous experimental evidence showing that spike activity levels estimated from extracellular recordings can sometimes be significantly higher than the true average level of spiking activity measured with intracellular recordings [12] and predicted by modeling studies [62], as extracellular methods shift firing rate distributions toward higher values (Fig. 6). This has an immediate impact on modeling work that often uses sorted firing rates as targets for their simulations [24, 25, 63]. It could be argued that the presence of low-firing rate neurons in our simulations, and hence this conclusion, is artificial. We note that, if the low-firing units were spiking at higher frequencies they would be more likely to be detected by the sorters, increasing the yield. The yield observed for our simulations is already higher than for the *in vivo* traces.

Additionally, our model identifies the second largest factor impacting sorting accuracy as the spatial footprint of spikes, aligning with analyses on spike spatial extent [7]. This finding directly supports recent demonstrations that denser probes than Neuropixels could enhance spike sorting yield threefold in cortical recordings by capturing small-footprint neurons more effectively [6] and is consistent with the observation that approximately half of isolated units with Neuropixels probes span multiple electrodes [9]. The fractional regression model offers a concise framework that disentangles and quantifies the effects of the key factors hypothesized to influence spike sorting accuracy [1, 12].

A novel and important prediction of the model is that a significant portion of the units isolated by the tested sorters are merged and incomplete, with sorting accuracies below 80%. The primary quality metric used to distinguish 'single-units' from 'multi-units' is the interspike interval (ISI, [7]). This metric is based on the reasoning that all neurons exhibit an absolute refractory period; thus, any cluster with a high proportion of ISIs less than 1–2 ms cannot represent a well-isolated unit. However, the opposite does not hold true: a clear refractory period does not necessarily indicate good isolation quality. For instance, a cluster containing intermixed spikes from two different neurons firing at distinct times may still show a clear refractory period.

To address this issue, we propose a new, straightforward model based on a linear combination of six quality metrics [16]. We demonstrate that this model effectively differentiate high-quality 'single-units' from merged and incomplete units among those produced by the Kilosort

sorters. The model's principles are intuitive: merged and incomplete units typically exhibit lower firing rates, a wider range of firing rates [16], greater shape variation (quantified by the median absolute deviation ratio, or 'mad_ratio' which we introduce here), and clusters that are more cohesive and better separated from others, as indicated by a silhouette score close to 1 [64].

In addition to these spike assignment biases, spike sorters displayed similar selection biases, particularly for excitatory pyramidal cells. Pyramidal cells were underrepresented in most layers, and overrepresented in layer 5 across sorters. Increased activity evoked by the model stimuli in Layer 5, which is an important input layer, receiving inputs from our model thalamus may contribute to the over-representation of excitatory cells in layer 5 compared to other layers. Why layer 2/3, which is also an input layer, did not show such an increase in excitatory cells could be explained by different morphologies of the cells in these two layers. Further analyses will also be needed to determine why a few rare interneurons were over-represented by some spike sorters while most were accurately represented. Altogether, these findings suggest that most selection biases may stem from mechanisms shared among sorters, while others are influenced by processes unique to each spike sorter. Understanding these characterized selection biases is crucial, as recording from statistically representative samples of isolated neurons is a high-priority objective in systems neuroscience [1]. As combined Neuropixels and optogenetic techniques soon become available, it will be possible to validate these predictions.

5 Conclusion

This study underscores the critical limitations of current spike sorting methods in capturing the full complexity of dense cortical activity, revealing significant undersampling, assignment, and selection biases that impair accurate neural representation. By leveraging a biophysically-detailed model of the rat cortical column, we have demonstrated that current algorithms capture only a fraction of isolable neurons and preferentially sort certain neuron types, thus distorting stimulus encoding. Despite undersampling not directly diminishing information capacity, biases in neuron selection and assignment notably reduced stimulus discriminability, affecting the interpretation of cortical responses. These findings highlight the value of realistic, high-resolution models as complementary tools for improving spike sorting algorithms, setting a path toward more accurate large-scale neural activity representations that can deepen our understanding of cortical function and the neural basis of behavior.

References

[1] Buzsáki, G.: Large-scale recording of neuronal ensembles. Nature Neuroscience **7**(5), 446–451

(2004) https://doi.org/10.1038/nn1233

[2] Urai, A.E., Doiron, B., Leifer, A.M., Churchland, A.K.: Large-scale neural recordings call for new insights to link brain and behavior. Nature Neuroscience **25**(1), 11–19 (2022) https://doi.org/10.1038/s41593-021-00980-9

[3] Gray, C.M., Maldonado, P.E., Wilson, M., McNaughton, B.: Tetrodes markedly improve the reliability and yield of multiple single-unit isolation from multi-unit recordings in cat striate cortex. Journal of Neuroscience Methods **63**(1-2), 43–54 (1995) https://doi.org/10.1016/0165-0270(95)00085-2

[4] Jun, J.J., Steinmetz, N.A., Siegle, J.H., Denman, D.J., Bauza, M., Barbarits, B., Lee, A.K., Anastassiou, C.A., Andrei, A., Aydın, C., Barbic, M., Blanche, T.J., Bonin, V., Couto, J., Dutta, B., Gratiy, S.L., Gutnisky, D.A., Häusser, M., Karsh, B., Ledochowitsch, P., Lopez, C.M., Mitelut, C., Musa, S., Okun, M., Pachitariu, M., Putzeys, J., Rich, P.D., Rossant, C., Sun, W.-l., Svoboda, K., Carandini, M., Harris, K.D., Koch, C., O'Keefe, J., Harris, T.D.: Fully integrated silicon probes for high-density recording of neural activity. Nature **551**(7679), 232–236 (2017) https://doi.org/10.1038/nature24636

[5] Steinmetz, N.A., Aydin, C., Lebedeva, A., Okun, M., Pachitariu, M., Bauza, M., Beau, M., Bhagat, J., Böhm, C., Broux, M., Chen, S., Colonell, J., Gardner, R.J., Karsh, B., Kloosterman, F., Kostadinov, D., Mora-Lopez, C., O'Callaghan, J., Park, J., Putzeys, J., Sauerbrei, B., Daal, R.J.J., Vollan, A.Z., Wang, S., Welkenhuysen, M., Ye, Z., Dudman, J.T., Dutta, B., Hantman, A.W., Harris, K.D., Lee, A.K., Moser, E.I., O'Keefe, J., Renart, A., Svoboda, K., Häusser, M., Haesler, S., Carandini, M., Harris, T.D.: Neuropixels 2.0: A miniaturized high-density probe for stable, long-term brain recordings. Science **372**(6539), 4588 (2021) https://doi.org/10.1126/science.abf4588

[6] Ye, Z., Shelton, A.M., Shaker, J.R., Boussard, J., Colonell, J., Manavi, S., Chen, S., Windolf, C., Hurwitz, C., Namima, T., Pedraja, F., Weiss, S., Raducanu, B., Ness, T.V., Einevoll, G.T., Laurent, G., Sawtell, N.B., Bair, W., Pasupathy, A., Mora Lopez, C., Dutta, B., Paninski, L., Siegle, J.H., Koch, C., Olsen, S.R., Harris, T.D., Steinmetz, N.A.: Ultra-high density electrodes improve detection, yield, and cell type specificity of brain recordings. bioRxiv (2023) https://doi.org/10.1101/2023.08.23.554527

[7] Pachitariu, M., Sridhar, S., Pennington, J., Stringer, C.: Spike sorting with Kilosort4. Nature Methods **21**(5), 914–921 (2024) https://doi.org/10.1038/s41592-024-02232-7

[8] Marques-Smith, A., Neto, J.P., Lopes, G., Nogueira, J., Calcaterra, L., Frazão, J., Kim, D., Phillips, M.G., Dimitriadis, G., Kampff, A.R.: Recording from the same neuron with high-density cmos probes and patch-clamp: a ground-truth dataset and an experiment in collaboration. bioRxiv (2020) https://doi.org/10.1101/370080

[9] Paulk, A.C., Kfir, Y., Khanna, A.R., Mustroph, M.L., Trautmann, E.M., Soper, D.J., Stavisky, S.D., Welkenhuysen, M., Dutta, B., Shenoy, K.V., Hochberg, L.R., Richardson, R.M., Williams, Z.M., Cash, S.S.: Large-scale neural recordings with single neuron resolution using Neuropixels probes in human cortex. Nature Neuroscience **25**(2), 252–263 (2022) https://doi.org/10.1038/s41593-021-00997-0

[10] Chung, J.E., Sellers, K.K., Leonard, M.K., Gwilliams, L., Xu, D., Dougherty, M.E., Kharazia, V., Metzger, S.L., Welkenhuysen, M., Dutta, B., Chang, E.F.: High-density single-unit human cortical recordings using the Neuropixels probe. Neuron **110**(15), 2409–24213 (2022) https://doi.org/10.1016/j.neuron.2022.05.007

[11] Beest, E.H., Bimbard, C., Fabre, J.M., Dodgson, S.W., Takács, F., Coen, P., Lebedeva, A., Harris, K.D., Carandini, M.: Tracking neurons across days with high-density probes. Nature Methods, 1–10 (2024)

[12] Shoham, S., O'Connor, D.H., Segev, R.: How silent is the brain: is there a "dark matter" problem in neuroscience? Journal of Comparative Physiology A **192**(8), 777–784 (2006) https://doi.org/10.1007/s00359-006-0117-6

[13] DeFelipe, J., Alonso-Nanclares, L., Arellano, J.I.: Microstructure of the neocortex: comparative aspects. Journal of neurocytology **31**, 299–316 (2002)

[14] Harris, K.D., Henze, D.A., Csicsvari, J., Hirase, H., Buzsáki, G.: Accuracy of Tetrode Spike Separation as Determined by Simultaneous Intracellular and Extracellular Measurements. Journal of Neurophysiology **84**(1), 401–414 (2000) https://doi.org/10.1152/jn.2000.84.1.401

[15] Rossant, C., Kadir, S.N., Goodman, D.F.M., Schulman, J., Hunter, M.L.D., Saleem, A.B., Grosmark, A., Belluscio, M., Denfield, G.H., Ecker, A.S., Tolias, A.S., Solomon, S., Buzsáki, G., Carandini,

M., Harris, K.D.: Spike sorting for large, dense electrode arrays. Nature Neuroscience **19**(4), 634–641 (2016) https://doi.org/10.1038/nn.4268

[16] Buccino, A.P., Hurwitz, C.L., Garcia, S., Magland, J., Siegle, J.H., Hurwitz, R., Hennig, M.H.: SpikeInterface, a unified framework for spike sorting. eLife **9**, 61834 (2020) https://doi.org/10.7554/eLife.61834

[17] Martinez, J., Pedreira, C., Ison, M.J., Quian Quiroga, R.: Realistic simulation of extracellular recordings. Journal of Neuroscience Methods **184**(2), 285–293 (2009) https://doi.org/10.1016/j.jneumeth.2009.08.017

[18] Garcia, S., Buccino, A.P., Yger, P.: How Do Spike Collisions Affect Spike Sorting Performance? eneuro **9**(5), 0105–222022 (2022) https://doi.org/10.1523/ENEURO.0105-22.2022

[19] Chung, Jason E., Magland, Jeremy F., Barnett, Alex H., Tolosa, Vanessa M., Tooker, Angela C., Lee, Kye Y., Shah, Kedar G., Felix, Sarah H., Frank, Loren M., Greengard, Leslie F.: MountainSort - A Fully Automated Approach to Spike Sorting. Neuron **95**(6), 1381–13946 (2017) https://doi.org/10.1016/j.neuron.2017.08.030

[20] Hill, D.N., Mehta, S.B., Kleinfeld, D.: Quality Metrics to Accompany Spike Sorting of Extracellular Signals. Journal of Neuroscience **31**(24), 8699–8705 (2011) https://doi.org/10.1523/JNEUROSCI.0971-11.2011

[21] Harris, K.D., Quiroga, R.Q., Freeman, J., Smith, S.L.: Improving data quality in neuronal population recordings. Nature Neuroscience **19**(9), 1165–1174 (2016) https://doi.org/10.1038/nn.4365

[22] Olshausen, B.A., Field, D.J.: How Close Are We to Understanding V1? Neural Computation **17**(8), 1665–1699 (2005) https://doi.org/10.1162/0899766054026639

[23] Quirk, M.C., Wilson, M.A.: Interaction between spike waveform classification and temporal sequence detection. Journal of neuroscience methods **94**(1), 41–52 (1999)

[24] Pronold, J., Meegen, A., Shimoura, R.O., Vollenbröker, H., Senden, M., Hilgetag, C.C., Bakker, R., Albada, S.J.: Multi-scale spiking network model of human cerebral cortex. Cerebral Cortex **34**(10), 409 (2024). Publisher: Oxford University Press

[25] Schmidt, M., Bakker, R., Shen, K., Bezgin, G., Diesmann, M., Albada, S.J.: A multi-scale layer-resolved spiking network model of resting-state dynamics in macaque visual cortical areas. PLOS Computational Biology **14**(10), 1006359 (2018). Publisher: Public Library of Science San Francisco, CA USA

[26] Dura-Bernal, S., Griffith, E.Y., Barczak, A., O'Connell, M.N., McGinnis, T., Moreira, J.V., Schroeder, C.E., Lytton, W.W., Lakatos, P., Neymotin, S.A.: Data-driven multiscale model of macaque auditory thalamocortical circuits reproduces in vivo dynamics. Cell reports **42**(11) (2023)

[27] Dura-Bernal, S., Herrera, B., Lupascu, C., Marsh, B.M., Gandolfi, D., Marasco, A., Neymotin, S., Romani, A., Solinas, S., Bazhenov, M., et al.: Large-scale mechanistic models of brain circuits with biophysically and morphologically detailed neurons. Journal of Neuroscience **44**(40) (2024)

[28] Reimann, M.W., Bolanõs-Puchet, S., Courcol, J.-D., Egas Santander, D., Arnaudon, A., Coste, B., Delalondre, F., Delemontex, T., Devresse, A., Dictus, H., Dietz, A., Ecker, A., Favreau, C., Ficarelli, G., Gevaert, M., Herttuainen, J., Isbister, J.B., Kanari, L., Keller, D., King, J., Kumbhar, P., Lapere, S., Lazovskis, J., Lu, H., Ninin, N., Pereira, F., Planas, J., Pokorny, C., Riquelme, J.L., Romani, A., Shi, Y., Smith, J.P., Sood, V., Srivastava, M., Van Geit, W., Vanherpe, L., Wolf, M., Levi, R., Hess, K., Schürmann, F., Muller, E.B., Markram, H., Ramaswamy, S.: Modeling and Simulation of Neocortical Micro- and Mesocircuitry. Part I: Anatomy. eLife **13**, 99688 (2024) https://doi.org/10.7554/eLife.99688.2

[29] Isbister, J.B., Ecker, A., Pokorny, C., Bolaños-Puchet, S., Santander, D.E., Arnaudon, A., Awile, O., Barros-Zulaica, N., Alonso, J.B., Boci, E., Chindemi, G., Courcol, J.-D., Damart, T., Delemontex, T., Dietz, A., Ficarelli, G., Gevaert, M., Herttuainen, J., Ivaska, G., Ji, W., Keller, D., King, J., Kumbhar, P., Lapere, S., Litvak, P., Mandge, D., Muller, E.B., Pereira, F., Planas, J., Ranjan, R., Reva, M., Romani, A., Rössert, C., Schürmann, F., Sood, V., Teska, A., Tuncel, A., Van Geit, W., Wolf, M., Markram, H., Ramaswamy, S., Reimann, M.W.: Modeling and simulation of neocortical micro- and mesocircuitry. part ii: Physiology and experimentation. eLife **13**, 99693 (2024) https://doi.org/10.7554/elife.99693.1

[30] Markram, H., Muller, E., Ramaswamy, S., Reimann, M.W., Abdellah, M., Sanchez, C.A., Ailamaki, A., Alonso-Nanclares, L., Antille, N., Arsever, S., Kahou, G.A.A., Berger, T.K., Bilgili, A., Buncic, N., Chalimourda, A., Chindemi, G., Courcol, J.-D., Delalondre, F., Delattre, V., Druckmann, S., Dumusc, R., Dynes, J., Eilemann, S., Gal, E.,

Gevaert, M.E., Ghobril, J.-P., Gidon, A., Graham, J.W., Gupta, A., Haenel, V., Hay, E., Heinis, T., Hernando, J.B., Hines, M., Kanari, L., Keller, D., Kenyon, J., Khazen, G., Kim, Y., King, J.G., Kisvarday, Z., Kumbhar, P., Lasserre, S., Le Bé, J.-V., Magalhães, B.R.C., Merchán-Pérez, A., Meystre, J., Morrice, B.R., Muller, J., Muñoz-Céspedes, A., Muralidhar, S., Muthurasa, K., Nachbaur, D., Newton, T.H., Nolte, M., Ovcharenko, A., Palacios, J., Pastor, L., Perin, R., Ranjan, R., Riachi, I., Rodríguez, J.-R., Riquelme, J.L., Rössert, C., Sfyrakis, K., Shi, Y., Shillcock, J.C., Silberberg, G., Silva, R., Tauheed, F., Telefont, M., Toledo-Rodriguez, M., Tränkler, T., Van Geit, W., Díaz, J.V., Walker, R., Wang, Y., Zaninetta, S.M., DeFelipe, J., Hill, S.L., Segev, I., Schürmann, F.: Reconstruction and Simulation of Neocortical Microcircuitry. Cell **163**(2), 456–492 (2015) https://doi.org/10.1016/j.cell.2015.09.029

[31] Reimann, M.W., King, J.G., Muller, E.B., Ramaswamy, S., Markram, H.: An algorithm to predict the connectome of neural microcircuits. Frontiers in computational neuroscience **9**(120) (2015) https://doi.org/10.3389/fncom.2015.00120

[32] Reimann, M.W., Gevaert, M., Shi, Y., Lu, H., Markram, H., Muller, E.: A null model of the mouse whole-neocortex micro-connectome. Nature Communications **10**(3903) (2019) https://doi.org/10.1038/s41467-019-11630-x

[33] Keller, D., Meystre, J., Veettil, R.V., Burri, O., Guiet, R., Schürmann, F., Markram, H.: A Derived Positional Mapping of Inhibitory Subtypes in the Somatosensory Cortex. Frontiers in Neuroanatomy **13**(78) (2019) https://doi.org/10.3389/fnana.2019.00078

[34] Kanari, L., Ramaswamy, S., Shi, Y., Morand, S., Meystre, J., Perin, R., Abdellah, M., Wang, Y., Hess, K., Markram, H.: Objective Morphological Classification of Neocortical Pyramidal Cells. Cerebral Cortex **29**(4), 1719–1735 (2019) https://doi.org/10.1093/cercor/bhy339

[35] Reva, M., Rössert, C., Arnaudon, A., Damart, T., Mandge, D., Tuncel, A., Ramaswamy, S., Markram, H., Geit, W.V.: A universal workflow for creation, validation and generalization of detailed neuronal models. Patterns **4**(100855) (2023) https://doi.org/10.1016/j.patter.2023.100855

[36] Barros-Zulaica, N., Rahmon, J., Chindemi, G., Perin, R., Markram, H., Muller, E., Ramaswamy, S.: Estimating the Readily-Releasable Vesicle Pool Size at Synaptic Connections in the Neocortex. Frontiers in Synaptic Neuroscience **11**(29) (2019) https://doi.org/10.3389/fnsyn.2019.00029

[37] Hilgen, Gerrit, Sorbaro, Martino, Pirmoradian, Sahar, Muthmann, Jens-Oliver, Kepiro, Ibolya Edit, Ullo, Simona, Ramirez, Cesar Juarez, Encinas, Albert Puente, Maccione, Alessandro, Berdondini, Luca, Murino, Vittorio, Sona, Diego, Zanacchi, Francesca Cella, Sernagor, Evelyne, Hennig, Matthias Helge: HerdingSpikes - Unsupervised Spike Sorting for Large-Scale, High-Density Multielectrode Arrays. Cell Reports **18**(10), 2521–2532 (2017) https://doi.org/10.1016/j.celrep.2017.02.038

[38] Lindén, H., Hagen, E., Łęski, S., Norheim, E.S., Pettersen, K.H., Einevoll, G.T.: LFPy: a tool for biophysical simulation of extracellular potentials generated by detailed model neurons. Frontiers in Neuroinformatics **7** (2014) https://doi.org/10.3389/fninf.2013.00041

[39] Tharayil, J., Blanco Alonso, J., Farcito, S., Lloyd, B., Cassara, A., Schürmann, F., Neufeld, E., Kuster, N., Reimann, M.: Bluerecording: A pipeline for the efficient calculation of extracellular recordings in large-scale neural circuit models. bioRxiv (2024) https://doi.org/10.1101/2024.05.14.591849

[40] Holt, G.R.: A critical reexamination of some assumptions and implications of cable theory in neurobiology. California Institute of Technology (1998)

[41] Horváth, C., Tóth, L.F., Ulbert, I., Fiáth, R.: Dataset of cortical activity recorded with high spatial resolution from anesthetized rats. Scientific Data **8**(1), 180 (2021) https://doi.org/10.1038/s41597-021-00970-3

[42] Buzsáki, G., Anastassiou, C.A., Koch, C.: The origin of extracellular fields and currents — EEG, ECoG, LFP and spikes. Nature Reviews Neuroscience **13**(6), 407–420 (2012) https://doi.org/10.1038/nrn3241

[43] Zhang, L.A., Li, P., Callaway, E.M.: High-resolution laminar identification in macaque primary visual cortex using neuropixels probes. eLife **13**, 97290 (2024) https://doi.org/10.7554/elife.97290.1

[44] Destexhe, A., Contreras, D., Steriade, M.: Spatiotemporal Analysis of Local Field Potentials and Unit Discharges in Cat Cerebral Cortex during Natural Wake and Sleep States. The Journal of Neuroscience **19**(11), 4595–4608 (1999) https://doi.org/10.1523/JNEUROSCI.19-11-04595.1999

[45] Schomburg, E.W., Anastassiou, C.A., Buzsáki,

G., Koch, C.: The spiking component of oscillatory extracellular potentials in the rat hippocampus. Journal of Neuroscience **32**(34), 11798–11811 (2012)

[46] Sun, S.H., Almasi, A., Yunzab, M., Zehra, S., Hicks, D.G., Kameneva, T., Ibbotson, M.R., Meffin, H.: Analysis of extracellular spike waveforms and associated receptive fields of neurons in cat primary visual cortex. The Journal of Physiology **599**(8), 2211–2238 (2021) https://doi.org/10.1113/JP280844

[47] Lee, E.K., Balasubramanian, H., Tsolias, A., Anakwe, S.U., Medalla, M., Shenoy, K.V., Chandrasekaran, C.: Non-linear dimensionality reduction on extracellular waveforms reveals cell type diversity in premotor cortex. eLife **10**, 67490 (2021) https://doi.org/10.7554/eLife.67490

[48] Meyer, H.S., Wimmer, V.C., Hemberger, M., Bruno, R.M., De Kock, C.P.J., Frick, A., Sakmann, B., Helmstaedter, M.: Cell Type–Specific Thalamic Innervation in a Column of Rat Vibrissal Cortex. Cerebral Cortex **20**(10), 2287–2303 (2010) https://doi.org/10.1093/cercor/bhq069

[49] Paulk, A.C., Kfir, Y., Khanna, A., Mustroph, M., Trautmann, E.M., Soper, D.J., Stavisky, S.D., Welkenhuysen, M., Dutta, B., Shenoy, K.V., Hochberg, L.R., Richardson, R.M., Williams, Z.M., Cash, S.S.: Large-scale neural recordings with single-cell resolution in human cortex using high-density neuropixels probes. bioRxiv (2021) https://doi.org/10.1101/2021.06.20.449152

[50] Wood, F., Black, M.J., Vargas-Irwin, C., Fellows, M., Donoghue, J.P.: On the variability of manual spike sorting. IEEE Transactions on Biomedical Engineering **51**(6), 912–918 (2004) https://doi.org/10.1109/TBME.2004.826677

[51] Schneider, S., Lee, J.H., Mathis, M.W.: Learnable latent embeddings for joint behavioural and neural analysis. Nature **617**(7960), 360–368 (2023) https://doi.org/10.1038/s41586-023-06031-6

[52] Stephenson, C., Feather, J., Padhy, S., Elibol, O.H., Tang, H., McDermott, J.H., Chung, S.: Untangling in invariant speech recognition. CoRR **abs/2003.01787** (2020) 2003.01787

[53] Meyer, H.S., Schwarz, D., Wimmer, V.C., Schmitt, A.C., Kerr, J.N.D., Sakmann, B., Helmstaedter, M.: Inhibitory interneurons in a cortical column form hot zones of inhibition in layers 2 and 5A. Proceedings of the National Academy of Sciences **108**(40), 16807–16812 (2011) https://doi.org/10.1073/pnas.1113648108

[54] Hines, M.: Efficient computation of branched nerve equations. International journal of bio-medical computing **15**(1), 69–76 (1984)

[55] King, J.G., Hines, M., Hill, S.L., Goodman, P.H., Markram, H., Schürmann, F.: A component-based extension framework for large-scale parallel simulations in NEURON. Frontiers in Neuroinformatics **3**, 456 (2009)

[56] Hines, M.L., Carnevale, N.T.: The NEURON simulation environment. Neural computation **9**(6), 1179–1209 (1997). Publisher: MIT Press

[57] Trautmann, E.M., Hesse, J.K., Stine, G.M., Xia, R., Zhu, S., O'Shea, D.J., Karsh, B., Colonell, J., Lanfranchi, F.F., Vyas, S., Zimnik, A., Steinmann, N.A., Wagenaar, D.A., Andrei, A., Lopez, C.M., O'Callaghan, J., Putzeys, J., Raducanu, B.C., Welkenhuysen, M., Churchland, M., Moore, T., Shadlen, M., Shenoy, K., Tsao, D., Dutta, B., Harris, T.: Large-scale brain-wide neural recording in nonhuman primates. bioRxiv (2023) https://doi.org/10.1101/2023.02.01.526664

[58] Rolston, J.D., Gross, R.E., Potter, S.M.: Common median referencing for improved action potential detection with multielectrode arrays. In: 2009 Annual International Conference of the IEEE Engineering in Medicine and Biology Society, pp. 1604–1607. IEEE, Minneapolis, MN (2009). https://doi.org/10.1109/IEMBS.2009.5333230

[59] Welch, P.: The use of fast Fourier transform for the estimation of power spectra: A method based on time averaging over short, modified periodograms. IEEE Transactions on Audio and Electroacoustics **15**(2), 70–73 (1967) https://doi.org/10.1109/TAU.1967.1161901

[60] Pillow, J.W., Shlens, J., Chichilnisky, E., Simoncelli, E.P.: A model-based spike sorting algorithm for removing correlation artifacts in multi-neuron recordings. PloS one **8**(5), 62123 (2013)

[61] Hildebrandt, K.J., Sahani, M., Linden, J.F.: The Impact of Anesthetic State on Spike-Sorting Success in the Cortex: A Comparison of Ketamine and Urethane Anesthesia. Frontiers in Neural Circuits **11**, 95 (2017) https://doi.org/10.3389/fncir.2017.00095

[62] Lennie, P.: The Cost of Cortical Computation. Current Biology **13**(6), 493–497 (2003) https://doi.org/10.1016/S0960-9822(03)00135-0

[63] Rimehaug, A.E., Stasik, A.J., Hagen, E., Billeh, Y.N., Siegle, J.H., Dai, K., Olsen, S.R., Koch, C., Einevoll, G.T., Arkhipov, A.: Uncovering circuit

mechanisms of current sinks and sources with bio-physical simulations of primary visual cortex. elife **12**, 87169 (2023)

[64] Rousseeuw, P.J.: Silhouettes: a graphical aid to the interpretation and validation of cluster analysis. Journal of computational and applied mathematics **20**, 53–65 (1987)

Acknowledgements

The authors would like to thank Enny Van Beest, Jennifer Colonell, Alessio Buccino and Marius Pachitariu for very insightful and critical feedback; Enny Van Beest, Ilias Rentzeperis and Cameron Mckenzie for helpful discussions and paper review suggestion; Alessio Buccino, Daniela Egas Santander, Christoph Pokorny, Andras Ecker, Natali Barros Zulaica for helpful discussions and support in identifying and accessing data analytical resources; the BBP Core Services team for responding to IT requests and services surrounding the research; Karin Holm for support of manuscript development and helpful discussions.

Funding

This study was supported by funding to the Blue Brain Project, a research center of the Ecole polytechnique fédérale de Lausanne (EPFL), from the Swiss government's ETH Board of the Swiss Federal Institutes of Technology.

Declarations

The authors declare no competing interests.

Supplementary figures

Fig. 1 Model neuron combinations of morphological and electrical properties and coverage of the cortical column by the dense probe. a. morpho-electric types used in this study, i.e., the combinations of neuron morphologies, electrical firing patterns, layers and their proportions (heatmap colors). Inhibitory parvalbumin positive (PV+) cell class (blue bar), somatostatin positive cell class (Sst+, green) and 5HT3aR+ cell class (orange) are indicated by the vertical colored bars on the side. The cACint electrical type belongs to the Sst+ class (green boxes). The panel is adapted from [29]. **b.** The number of electrodes per layer of the Horvath setup [41] (filled grey bars) is plotted against the number of electrodes per layer of the simulated dense probe (empty red bars).

Fig. 2 Validation of raw simulated traces. First 4 seconds of raw *in vivo* traces for **a**) Marques-Smith's Neuropixels [8], **b**) biophysically simulated recordings in the **b**) spontaneous and **c**) evoked regime, the **d**) Horvath's dense recordings [41] and **e**) biophysical simulated recordings with Horvath's dense probe, from the most superficial recording site of each cortical layers. The Horvath traces display slow waves typically observed during slow wave sleep. **f**. Average power spectral densities (PSDs) of **f**) the Neuropixels and **g**) the dense probe raw recordings, calculated over sites. Means are shown with their 95% confidence interval (areas). Each trace's mean was subtracted beforehand to remove the signal's DC components, then PSDs were computed with the Welch method on the first 10 minutes of recording, compressed to int16 and divided by the total power, **h**. PSDs on log-linear plots with power law fits in the LFP band (0 - 90Hz) and spiking band (300 - 6000Hz), and the mean and 95% confidence intervals (text on the right), **i**. Boxplots of the parameter α of the power law fits in the LFP band.

Fig. 3 Spike shape change with electrode location. Spike-triggered average of simulated recording traces from 16 excitatory (A) and 20 inhibitory (B) cells located within 50 µm of the Neuropixels probe and sorted by morphological type. The spike waveforms were recorded with two electrodes, one 20 µm above the cell somata (red in A, blue in B) and the other 20 µm below (orange in A, purple in B). For each morphological type, we display the cell with the highest firing rate.

Fig. 4 Validation of firing rate distributions. a. Statistics of sorted single-unit firing rates. Units were sorted with all Kilosort sorters applied to *in vivo* data and simulations per cortical layer. Statistics are reported for the decimal logarithm of the firing rate. All units are reported for Kilosort, which does not classify single and 'multi-unit', **b**. Statistics of the distribution variances, calculated from 100 bootstraps of the decimal logarithm of the sorted firing rates. Boxplot notches represent the 95% confidence intervals. Statistical significance of the null hypothesis that the model sorted median firing rate and variance are the same as *in vivo* data are reported with ns, *,**,***, which indicate p<=0.05, p>0.05, p<0.01, p<0.001 using the Mann-Whitney U test, **c**. Lines represent lognormal distribution fits to the firing rate distributions.

Fig. 5 Single-unit yield and quality predicted by the different ground truth simulations for all tested spike sorters. **a**. Number of isolated single-units per electrode for each experiment. The bar heights indicate the median yields, averaged over layers and the error bars indicate its 95% confidence interval. The colored horizontal lines indicate the median theoretical yield per site averaged over layers. It is the number of active ground truth neurons within 50 μm divided by the number of sites, and averaged over layers, **b**. Ratio of isolated single-units of each quality by layer and spike sorter for each ground truth simulation (rows). The overlaid numbers indicate the number of single-units. Precision and recall performance of a CEBRA-KNN model that classifies good units and degraded units in the **c**) spontaneous and **d**) evoked biophysical simulations of Neuropixels recordings. The Cebra-KNN model transforms single-unit quality metrics and their full spike waveforms into a lower dimensional embedding which was used to train a binary k-nearest neighbor classifier to predict good units from other units, **e**. The model's InfoNCE loss functions for all bootstrapped samples. Loss functions converged within 60 iterations.

.

Fig. 6 Sorting accuracy. A) Effect of the *delta_time* parameter on sorting accuracy, for 10 min recording of the simulation of Neuropixels recordings sorted with Kilosort3.

Fig. 7 Ground truth firing rate distribution. **a**. Violinplots of the distributions of sorted (orange) and true (blue) firing rates in individual layers for the spontaneous biophysical simulations of Neuropixels recording. True firing rates are based on all neurons within $50\mu m$ of the recording electrode. Widths of the violins normalized for each class. **b**. as **a**, but with non-normalized widths, i.e. the widths are directly proportional to the number of neurons in each bin. **c**. Ratio of sorted divided by true neuron count in each logarithmic firing rate bin. Colors indicate results for individual layers. **d**. Ratio of true divided by sorted mean firing rates in each layer. Grey area indicates a 90% confidence interval based on bootstrapping. **e**. as D, but instead the difference between the logarithm of the means is indicated.

Fig. 8 Information capacity is robust to change in dimensionality. Information capacity of randomly sampled cortical populations or sampled to reproduce Kilosort4's unit type distribution, using the same probability distribution over unit types as observed in Kilosort4's sorted population.