Mesoscale simulations of membrane-tethered reactions to parameterize cell-scale models of signaling**.**

Kelvin J. Peterson, Boris M. Slepchenko and Leslie M. Loew*

R. D. Berlin Center for Cell Analysis and Modeling, University of Connecticut School of Medicine, Farmington, CT USA

*Correspondence to les@uchc.edu

ABSTRACT. Biochemical interactions at membranes are the starting points for cell signaling networks. But bimolecular reaction kinetics are difficult to experimentally measure on 2-dimensional membranes and are usually measured in volumetric *in vitro* assays. Membrane tethering produces confinement and steric effects that will significantly impact binding rates in ways that are not readily estimated from volumetric measurements. Also, there are situations when 2D reactions do not conform to simple mass action kinetics. Here we show how highly coarse-grained molecular simulations using the SpringSaLaD software can be used to estimate membrane-tethered rate constants from experimentally determined volumetric kinetics. The approach is validated using an analytical solution for dimerization of binding sites anchored via stiff linkers. This approach can provide 2-dimensional bimolecular rate constants to parameterize cell-scale models of receptor-mediated signaling. We explore how factors such as molecular reach, steric effects, disordered domains, local concentration and diffusion affect the kinetics of binding. We also develop a general scheme to assess whether simple mass action rate constants can be applied for a given scenario, taking into account the diffusivity of the membrane anchors and tethered binding sites, the initial membrane densities of the reactants and the desired level of completion for the fitted rate constant. We then apply our approach to epidermal growth factor receptor (EGFR) mediated activation of the membrane-bound small GTPase Ras. The analysis reveals how binding of Ras to the allosteric site of SOS, a guanine nucleotide exchange factor (GEF) that is recruited to EGFR, significantly accelerates Ras binding to the SOS catalytic site. A small biochemical network model parametrized with the derived 2D rate constants shows how recruitment of SOS via EGFR can significantly enhance Ras activation.

SIGNIFICANCE STATEMENT. In cell signaling, the activation of a surface receptor leads to a cascade of intracellular biochemical events. Many protein interactions occur near the inner plasma membrane surface. However, accurate rate parameters for these steps in models of signaling are rarely available because membrane-tethered reaction kinetics are difficult to experimentally measure. Here, we use a highly coarse-grained molecular simulator to model the kinetics of reactions between binding sites that are tethered to a membrane. We can fit these simulation outputs with 2-dimensional rate laws to obtain rate constants that can be used to build complex models of cell signaling. The derived rate constants can also be analyzed to understand the key biophysical features controlling the kinetics of bimolecular membrane reactions.

Introduction.

The cell membrane responds to and integrates electrical, mechanical and chemical signals from the extracellular environment. For chemical signals, the initial step is binding of a ligand to an external binding site on a membrane receptor protein. This triggers a chain of events that typically involves a change of state of the cytoplasmic receptor domain and subsequent recruitment of adapter proteins, enzymes and/or cytoskeletal regulators to further evoke a cell biological response. Mathematical modeling of signaling pathways is a powerful tool to systematically organize the experimental knowledge we have about these complex systems and then develop predictions, through simulations, to inspire new experiments (1-3).

A challenge in developing cell signaling models is the acquisition of the appropriate experimentally-grounded input parameters. Often, kinetic data is available from *in vitro* biochemistry and this has served the mature field of metabolic modeling very well. However, rate parameters are less available and more difficult to measure for signaling pathways and networks. Among the key challenges is that many of the essential steps are associated with the plasma membrane, where multiple molecules are recruited before a messenger ultimately diffuses to an intracellular target (e.g. the nucleus). It is experimentally difficult to measure reaction rates on membranes because complex reconstituted bilayer methods must be used (4), so available data is commonly derived from volumetric measurements. While such quantitative data is useful, it can be challenging to translate rate parameters derived from 3D solution to the very different biophysical environment of a 2D membrane.

Indeed, the biophysics of membrane associated reactions has a long scientific history. An early focus of investigation, initiated with a classic paper by Adam and Delbruck (5), was the difference between 2D and 3D diffusion limited reactions; they argued that the 2-step process of absorbing a cytosolic molecule to the membrane and subsequent 2D search for an enzyme or binding partner might offer a kinetic advantage over a fully 3D search. This argument emerges from the fundamental difference between 2D and 3D diffusion: a diffusing molecule in 2D is guaranteed to eventually find its target, while no such guarantee can be made in 3D (6). In practice, Adam and Delbruck's assertion that reactions in membranes have an inherent advantage has been disputed using subsequent experimentally determined realistic parameters and biological scenarios; this is because of the generally low diffusivity of membrane embedded molecules compared to the same sized molecule diffusing in the cytosol (7).

However, regardless of whether 2D reactions are advantageous over 3D, a consequence of the Adam and Delbruck (5) analysis is that there is a fundamental difference between 2D and 3D bimolecular reactions. Reactions in 3D can generally be described by mass action kinetics, where rate constants are truly constant, i.e independent of time and reactant concentrations. However, bimolecular reactions on a surface cannot be fundamentally described by mass action kinetics (8-14). With regard to cell signaling, theoretical analyses of reactions at membranes have been extended and elaborated to consider both diffusion-limited and reaction-limited bimolecular kinetics (1,8,10,11). Intuitively, for reaction-limited cases (where diffusion is fast compared to the intrinsic reaction rate upon encounter) the Adam and Delbruck idea is not pertinent and the reaction can be well described by simple mass action kinetics. In particular, Yogurtcu and Johnson (10) developed the theory to determine where the transition from reaction-limited to diffusion limited 2D kinetics occurs – i.e. when mass action in 2D is an adequate description or when a more complex time-dependent parameterization of the rate is required.

These pioneering studies treated membrane reactions as strictly two-dimensional surface
events. However, most biological membrane associated reactions actually occur in the
immediately adjacent cytosol, with interacting sites tethered to the membrane through lipid or
protein anchors. A well-known feature of anchoring bimolecular reactions to a membrane is the
effect of locally increased effective concentration: compared to the same reaction by the same
number of molecules within the cell volume, anchoring the reaction to the membrane generally
increases concentration by confining the reaction volume to a thin layer above the membrane
(15). The thickness of that layer is often parametrized as *h,* sometimes called the "confinement
length" (16,17). Roughly, *h* is related to the distance the binding sites can sample above the
membrane surface. The smaller *h*, the greater the effective concentrations of binding sites and
the faster the membrane associated bimolecular kinetics. In principle, *h*, can be determined by
analyzing detailed molecular dynamics simulations on the flexibility and motions of binding
domains tethered to the membrane (17,18). Recently, the concept of *molecular reach* was
introduced as a more general framework for assessing how molecular structure influences the
steady state phosphorylated fraction of a membrane bound substrate interacting with a tethered
kinase (19). The *reach* is defined as the distance of the kinase site from the membrane anchor
and is directly related to *h* when the anchor diffuses freely in the membrane. However, when
lateral diffusion is restricted (e.g. within large signaling clusters such as the immune synapse),
binding sites with longer *reach* may have an advantage in being able to find more binding
partners; for such diffusion-limited scenarios, this increase in reach can outweigh the decrease
in local concentration associated with increased *h* (19).
118
Thus, it is clear from these foundational studies that the membrane environment and the
structural features of interacting membrane-bound molecules must be considered when
converting a measured 3D on-rate to a 2D on-rate suitable for cell-scale continuum models
based on ordinary or partial differential equations (ODEs or PDEs). In this work, we show how
this can be done using simulations from SpringSaLaD (20) to derive 2D rate constants.
Importantly, we are treating tethered binding sites. This is in fact the more general situation in
cell signaling where a more-or-less flexible protein domain containing the binding site extends
into the cytosol a significant distance from the actual membrane – typically much longer than the
actual thickness of the lipid bilayer. Because our method accounts for the structure of the
protein (albeit very coarsely), the distance of the binding sites from the membrane as well as
other biophysical features are accounted for in the estimation of the 2D on-rate constant.
130
SpringSaLaD uses a series of variously sized spherical sites linked together with stiff springs to
coarsely model the key structural features of macromolecules such as flexibility, excluded
volume and binding site localization. Each sphere within the molecule can be assigned its own
diffusion coefficient and Brownian diffusion is simulated via a Langevin dynamics algorithm. The
molecule can be tethered to a surface, representing a membrane, via a specialized anchoring
sphere that has a lateral diffusion coefficient; the rest of the molecule, including the spheres
designated as binding sites, are free to explore the volume above the membrane within their
reach. Naturally therefore (and particularly advantageous for the purpose of this work),
volumetric rate constants are used for bimolecular rate expressions even for membrane-bound
molecules. In previous work, this feature was used to show how multivalent clustering is
enhanced when one of the interacting molecules is tethered to a membrane (21).
142
To determine 2D rate constants, we fit the stochastic kinetics simulated with SpringSaLaD to a
deterministic mass-action rate law based on the corresponding surface densities of the binding
partners. Using idealized structures, we validate this procedure against theoretical calculations.
We then explore how molecular structural features (e.g. tether length and stiffness, steric
access), surface density and lateral diffusion affect binding kinetics. Additionally, we develop a

theory for membrane-tethered reactions, similar to that of Yogurtcu and Johnson for purely 2D
reactions (10), to estimate the applicability of the mass action rate law for varying tether lengths,
anchor diffusion coefficients, reaction rates, initial surface densities and the desired level of
completion of the reaction. Then, as a biologically relevant example, we apply this approach to
recruitment of SOS to the epidermal growth factor receptor (EGFR). SOS is the G-protein
exchange factor (GEF) for Ras (22) and a better understanding of SOS activation of Ras
emerges from this analysis. Thus, we offer a procedure for parameterizing membrane kinetics
that serves to bridge molecular to cell scale simulations of cell signaling.
156
157 **Methods**
158
All simulations were performed with SpringSaLaD v. 2.3.4 (https://vcell.org/ssalad). While the
full physics and math behind the calculations performed by SpringSaLaD can be found in the
original publication (20), we provide a brief overview of its functionality. A molecule in
SpringSaLaD is approximated as a series of variably sized hard spheres, representing protein
domains, connected by stiff springs. Random forces (chosen from a normal distribution with a
variance determined from the diffusion coefficient assigned to each sphere by the user) impinge
on each sphere at each time step and are transmitted to neighboring spheres through the stiff
springs. The software models membranes as a planar surface within the volume of the 3D
simulation domain. The membrane can have embedded spheres that serve as diffusing anchors
(with user-assigned 2D diffusion coefficients) linked to spheres that extend into the volumetric
space. Each sphere may also be designated as the site of a reaction. The user can input
macroscopic rate constants for reactions and the software converts these to reaction
probabilities within each time step; for bimolecular reactions, this conversion also depends on
the radii of the spherical binding sites and the sum of their diffusion coefficients. SpringSaLaD
allows users to manually edit molecules to fully customize structure and molecule flexibility.
Orientational approaches between spherical binding sites can be restricted by the appropriate
placement of neighboring spheres adjacent to the respective binding sites within the reacting
molecules. To build coarse grained molecular structures used in SpringSaLaD for Fig. 3, we
used AlphaFold2 (23,24) to generate PDB file estimates of protein structures for EGFR, Grb2,
SOS, and Ras via input of entire amino acid sequences. These PDB files are converted to
highly coarse-grained molecular models via the mol2sphere (25) utility embedded in
SpringSaLaD. In some cases, we manually edited the structures to capture their essential
features from measurements on the PDB structures, as visualized in PyMol (Schrödinger, Inc.).
In particular, for SOS we subdivide the CDC25 and REM domains into multiple smaller spherical
sites with only two sites (the allosteric and catalytic sites for RAS) capable of participating in a
binding reaction. This process maintains the structural characteristics of these domains, while
ensuring that the binding radius of the domain is not artificially inflated. Modeling disordered
regions, such as the PRM region of SOS, can be challenging due to low confidence in the
AlphaFold2-generated geometry of these regions. To model disordered domains, we use
PyMOL to measure the length of entire straight chain amino acid sequences, then model this
sequence in SpringSaLaD using 1.0 nm diameter sites connected by 3.1 nm linkers (in
SpringSaLaD, a string of relatively small spheres connected by links that are larger than the
sphere diameters can be used to model flexible (i.e. disordered) domains). Binding reactions in
all simulations have rates input in terms of $\mu M^{-1}s^{-1}$; in our simulations, successful binding results
in 1nm (Tables 1 and 2, Fig. 4) distances between the surfaces of the spherical sites. The
default simulation time step of 10ns was sufficiently accurate for all the SpringSaLaD
simulations in this work (as checked by testing 2 ns timesteps and noting no significant
differences). Simulations for Figs. 1 and 2 were run with 40 molecules, while Figure 4 used 200
molecules of Ras and 20 complexes of EGFR-Grb2-SOS to proportionally reflect the difference
in concentrations between Ras and SOS molecules *in-vivo* (26,27). The 3D problem domain

had a z coordinate of 100nm, with the membrane placed in the XY plane at Z=10nm; the
boundaries at the edges of the domain have reflective boundary conditions. X and Y
coordinates were set to provide the desired initial surface densities. Initial placement of the
molecules was random and different for each trajectory. A set of 100 trajectories are simulated
in parallel using the Center for Cell Analysis and Modeling High Performance Compute Cluster
(https://health.uconn.edu/high-performance-computing/resources/). One run for Tables 1 and 2
required approximately 3 hours and one run for Fig. 4 required approximately 30 hours.
SpringSaLaD input files are in Supporting Information and provide all the geometric details for
the molecules in each computational experiment.

The mean time dependence of dimer surface density $[dimer]_t$, based on 100 SpringSaLaD
runs for each of the parameter sets in Tables 1 and 2 and Fig. 4, were fitted with a deterministic
2D mass action rate law to obtain $k_{on}^{(mem)}$. For irrreversible dimerization (Table 1, first 8 rows
and Table 2), a fit to the analytical mass action expression for the appearance of dimer (Eq. 1)
used the Excel solver.

$$[dimer]_t = [dimer]_\infty - \frac{[dimer]_\infty}{(4*k_{on}^{(mem)}*[dimer]_\infty*t)+1}$$ Eq. 1

For reversible dimerization (Table 1, rows 9 and 10) and for the fits in Fig. 4, we used the
COPASI (28) parameter estimation tool within Virtual Cell (VCell) (29,30). The latter can be
accessed in the VCell published BioModel "Peterson Figure 4: Ras-
SOS_Binding_fit_to_SpringSaLaD". All these results with some further analysis can also be
found in the spreadsheets included in the Supporting Information. The VCell model related to
Fig. 5 can be found in the VCell database with the name "SOS_Recruitment_Ras_Binding"

Results

<u>The general approach.</u>

For bimolecular reactions, SpringSaLaD determines the microscopic probability of 2 binding
sites forming a bond as they diffuse within a reaction radius that is slightly larger than the sum of
their physical radii. The input to the algorithm is simply the macroscopic volumetric on rate
constant ($k_{on}^{(vol)}$, units of μM^{-1}s^{-1}) and the diffusion coefficient of the individual spheres. Full
details on the derivation of the reaction probability and a thorough validation of its accuracy can
be found in the original paper describing SpringSaLaD (20). Importantly for the purposes of this
work, the rate of binding for sites that happen to be tethered to a membrane are still treated as
volumetric, because the spherical sites are located in the volume compartment even while they
are constrained with links to the 2D membrane surface.

Step 1: Run 100 dimerization Simulations in SpringSaLaD

$k_{on}^{(vol)} = 0.47 \mu M^{-1}s^{-1}$

Step 2: Fit mean of the 100 SpringSaLaD trajectories to find $k_{on}^{(mem)}$ in a deterministic dimerization model using surface densities

Fit $k_{on}^{(mem)} = 0.109 \ \mu m^2 molecule^{-1}s^{-1}$

Figure 1. Workflow for finding membrane bimolecular binding rate constants, $k_{on}^{(mem)}$, in terms of surface densities. Step 1 is to run 100 SpringSaLaD trajectories with the coarse-grained model of the binding partners. Illustrated are a pair of simple monomers (top) consisting of 2nm diameter spheres (yellow) tethered to a 1nm membrane anchor sphere (gray) with a 5nm link; both the anchor and binding spheres are assigned the diffusion coefficient of 1 μm^2/s. At the bottom, the product dimer is depicted. Step 2 consists of fitting the average of 100 outputs from stochastic SpringSaLaD (SS) simulations with a deterministic (ODE) non-spatial model of surface-bound dimerization.

Figure 1 illustrates this for a simple dimerization reaction where the two yellow binding sites are
tethered to the membrane anchor (gray sphere) by a 5nm link; in these simulations both the
anchor and tether sites are given identical diffusion coefficients of 1μm^2/s. We ran 100
SpringSaLaD simulations each with 40 dimerizing molecules (Fig. 1 only shows two molecules
for clarity). The mean trajectories for these 100 runs are then fitted with a deterministic mass
action membrane binding model in terms of surface densities using either COPASI or Virtual
Cell (although an analytical solution can be also fit for simple dimerization) (Step 2 in Fig. 1).
The output binding rate constant, $k_{on}^{(mem)}$ in units of $\mu m^2 molecule^{-1}s^{-1}$, can then be used to
parameterize larger deterministic or stochastic models (that utilize mass action rate constants
as inputs) with molecule numbers (>1000) or timescales (>10 s) that would be too large for even
highly coarse-grained molecular simulators like SpringSaLaD. Also, we emphasize that binding
rates are typically determined experimentally using in vitro volumetric measurements; that
SpringSaLaD uses volumetric rate constant inputs, makes the procedure in Fig. 1 especially
appropriate and convenient. The results in Fig. 1 show that a 2nm diameter binding site with an
on rate for dimerization of 0.47 $\mu M^{-1}s^{-1}$ and tethered to a membrane surface through a 5nm link,
can be modeled as a 2D surface reaction with an on rate, $k_{on}^{(mem)}$, of 0.109 $\mu m^2 molecule^{-1}s^{-1}$.
Using idealized models, we now explore how various structural and biophysical parameters

control $k_{on}^{(mem)}$ and when mass action rate constants may not be appropriate to describe
membrane-associated binding kinetics.
258
259 <u>Effects of volumetric on-rate, 2D diffusion rate, surface density and structural features on</u>
260 <u>dimerization kinetics in idealized systems</u>
261
We start by analyzing the case of a 5nm stiff tether between a 1nm diameter binding sphere and
the membrane. The diffusion coefficient for the binder sphere is set to 1µm²/s. Table 1 gives
results for all combinations of two volumetric on rate constants, two surface densities and two
membrane diffusion coefficients. All the dimerization rate laws are irreversible except for the last
two rows, where the off-rate constant is indicated. The 2D on-rate constant derived by fitting the
SpringSaLaD simulation, $k_{on}^{(mem)}$, is in the fifth column; for consistency, all these fits are
performed for kinetics at 80% completion (the impact of the choice of the level of completion is
discussed in the next section). For comparison we also provide the 2D on-rate constant, $k_{on}^{(h)} =$
$k_{on}^{(vol)}/h$, where h is the linker length plus the radius of the binder sphere. If h is in units of µm,
dividing by a unit conversion factor of 602.2 converts $k_{on}^{(h)}$ from units of µM⁻¹s⁻¹ to units of
µm²molecules⁻¹ s⁻¹; $k_{on}^{(h)}$ is an equivalent 2D binding rate constant of freely diffusing monomers
confined within a thin volume with a height h adjacent to the membrane. The Relative Sq Dev
column provides the ratio of the sum of the squared deviations to the sum of the squared
SpringSaLaD mean values; this ratio provides a measure of the goodness of fit to the
bimolecular mass action rate law, with anything less than ~10⁻³ representing a good fit.
Examples of the fits are shown in Figure 2. The last column, δ, is a measure of how applicable
mass action kinetics are for these combinations of parameters; this will be further discussed in
the next Section.

Table 1. Results for single stiff link, binding site D_{vol}=1.0µm²/s, h=0.0055µm

$k_{on}^{(vol)}$ (µM⁻¹s⁻¹)	Surface density (molecules/µm²)	D_{mem} (µm²/s)	$k_{on}^{(h)} = k_{on}^{(vol)}/h/602.2$ (µm²*molecules⁻¹*s⁻¹)	$k_{on}^{(mem)}$, SpringSaLaD fit (µm²*molecules⁻¹*s⁻¹)	Relative Sq Dev	δ
0.01	2500	0.01	3.0 X10⁻³	2.9 x 10⁻³	3.0 x 10⁻⁵	0.09
0.01	2500	1.0	3.0 X10⁻³	3.1 x 10⁻³	3.1 x 10⁻⁵	0.001
0.01	25	0.01	3.0 X10⁻³	2.4 x 10⁻³	2.9 x 10⁻⁵	0.13
0.01	25	1.0	3.0 X10⁻³	3.0 x 10⁻³	2.3 x 10⁻⁵	0.002
1.0	2500	0.01	0.30	0.08	7.4 x 10⁻³	0.78
1.0	2500	1.0	0.30	0.26	2.6 x 10⁻⁴	0.04
1.0	25	0.01	0.30	0.02	1.5 x 10⁻³	0.9
1.0	25	1.0	0.30	0.20	4.9 x 10⁻⁴	0.09
1.0 (k_{off}= 10³s⁻¹)	2500	0.01	0.30	0.18	1.3 x 10⁻³	
1.0 (k_{off}= 10s⁻¹)	25	0.01	0.30	0.1	6.2 x 10⁻²	

280
We looked at two anchor diffusion coefficients corresponding to that of a large transmembrane
protein domain (D_{mem}= 0.01 µm²/s) and a lipid anchor (D_{mem}=1 µm²/s). The first row of Table 1
considers a case where diffusion of the binder (D_{vol}=1 µm²/s) is 100 times that of the anchor site
in the membrane. Because the linker is stiff, the anchor acts as a pivot and the binder rapidly

moves within a hemispherical shell to effectively create a reaction region with a thickness
slightly larger than the 1-nm diameter of the binder sphere. Because the region of spatial
overlap of the two shells where the binding may occur is restricted, it might seem surprising that

Figure 2. Examples of SpringSaLaD simulation data and their fit to surface-confined mass action dimerization kinetics. The initial surface density of monomers in each case is 2500 molecule/μm^2. The diffusion coefficient of the anchor and the volumetric binding rate constant of the binding sites are indicated above each graph, corresponding respectively to rows 1, 6 and 5 of Table 1. Each of the SpringSaLaD simulations were run to 80% completion.

$k_{on}^{(mem)}$ is so closely approximated by $k_{on}^{(h)}$. However, as shown in the Supporting Text I, the
reaction probability is enhanced because of the effectively higher density of binding sites within
this restricted region, which compensates for the smaller spatial overlap. Thus, the calculations
in the Supporting Text I both explain and validate the SpringSaLaD results. The second row of
Table 1 corresponds to the case where the anchor and the binder have the same fast diffusion.
In this scenario, that $k_{on}^{(mem)} \cong k_{on}^{(h)}$ is intuitive, because the effect of the tether in this case
essentially reduces to confining the binders within the layer adjacent to the membrane. The
solutions in the Supporting Information assume that the molecular distributions are spatially
uniform at any time (i.e., well-mixed), which pertains best to the cases where the reaction is
slow on the timescale of <u>anchor</u> diffusion (reaction-limited kinetics).
298
It has been shown that the rates of membrane reactions may be susceptible to deviations from
a simple mass action rate law (8,10,14,31-35), which manifests themselves as significant
changes of the apparent rate constant with initial surface density. These changes need to be
assessed before using the rate constants we obtain by the procedure of Fig. 1 in large cell-scale
models. We start by a quantitative exploration of how variations of some of the key parameters
in the test cases of Table 1 might expose deviations from mass action. We probed for this by
decreasing the initial density by a factor of 100. The combinations of D_{mem} and $k_{on}^{(vol)}$ in the third
and fourth rows of Table 1 resulted in relatively small changes in $k_{on}^{(mem)}$, indicating that the
mass action rate law applies to these cases. We further tested this by increasing the $k_{on}^{(vol)}$ by a
factor of 100 in the lower half of Table 1. Clearly, for the case of D_{mem}= 0.01 μm^2/s, there is a
strong dependence on surface density. Furthermore, the system under these conditions
develops depletion zones, as discussed below. Thus, the combination of $k_{on}^{(vol)}$= 1.0 $\mu M^{-1}s^{-1}$ and
D_{mem}= 0.01 μm^2/s present cases where mass action rate laws would not apply. Of course, the
appropriateness of the mass action rate law can also be judged by the goodness of fit when
determining $k_{on}^{(mem)}$ from the SpringSaLaD simulation (last column of Table 1); as demonstrated
in Fig. 2, the case of $k_{on}^{(vol)}$= 1.0 $\mu M^{-1}s^{-1}$ and D_{mem}= 0.01 μm^2/s is not approximated very well by
a mass-action dimerization rate. The simulation results are initially faster and then ultimately
slower than the best fit that assumes mass action kinetics. This is because the monomers

whose binding sites are initially close to each other (effectively within the "reach" of the tether)
will react, but leave behind depletion zones where monomers are too far away from potential
binding partners (33,34). These isolated monomers can be discerned toward the end of Movie
1, which presents an example trajectory for this case. A discussion of how to more generally
assess the appropriateness of mass action kinetics is provided in the next Section.
322
Consistent with the idea of depletion zones, the last 2 rows of Table 1 show that when
reversibility is introduced, the fitted on-rate constant increases (compare, respectively, to rows 5
and 7 of Table 1). This is because when dimerization is reversible, free monomers can reappear
to fill in depletion zones, thereby countering the slow diffusion. While the mass action kinetics
approximations are still poor, especially for the low density case, the respective mean steady-
state densities of dimers in the SpringSaLaD simulations, 570 and 5.7 molecules/μm^2, are
consistent with the *thermodynamic* law of mass action (9). According to thermodynamics, the
ratio of the squared monomer surface density and the dimer surface density is equal at steady
state to the equilibrium dissociation constant K_d, which is determined by the strength of the
interaction between the monomers and thus is independent of reaction kinetics. When kinetics
obey mass action, K_d is equal to the ratio of off and on rate constants. In our case, $K_d = k_{off}/$
$k_{on}^{(h)}$, because the interactions of the binding sites in the 3D volume above the membrane obey
mass-action kinetics and $k_{on}^{(h)}$ is the 3D association rate constant adjusted for the restricted
volume above the membrane. It is gratifying that the SpringSaLaD simulations produce these
correct steady state surface densities even when, as in the last 2 rows of Table 1, mass action
kinetics are not obeyed.

Table 2. Dimerization of idealized structure types.

D_{mem}=0.01μm^2/s, binding site D_{vol}=1.0μm^2/s, $k_{on}^{(vol)} = 1$ $\mu M^{-1}s^{-1}$, h=0.0205μm, surface density = 2500 molecules/μm^2

Structure	$k_{on}^{(h)} = k_{on}^{(vol)}/h/602.2$ (μm^2*molecules^{-1}*s-1)	$k_{on}^{(mem)}$, SpringSaLaD fit (μm^2*molecules^{-1}*s-1)
Single site, stiff	0.081	0.078
Steric sites, stiff	0.081	0.061
Steric sites, flexible	0.081	0.064

339
Table 2 provides results for three computational experiments in which structural features of the
SpringSaLaD molecules are varied. These are all for dimerization reactions where the maximum
distance between the membrane and the binding site is four times longer than in Table 1: $h =$
0.0205 μm (linker length of 20nm and binding site radius of 0.5nm). The $k_{on}^{(h)}$ is therefore a
factor of ~4 slower than that in Table 1 for the same $k_{on}^{(vol)}$ of 1$\mu M^{-1}s^{-1}$. The SpringSaLaD
simulations were carried out with a slow membrane-anchor diffusion coefficient, D_{mem}=
0.01μm^2/s, to model a transmembrane protein domain. The first row can be directly compared to
the fifth row of Table 1, where the only difference is the length of the linker. As would be
expected from the increase in *h*, the membrane on rate constant is decreased; however,
importantly, this constant is now closer to the diffusion rate and therefore deviations from mass
action are significantly reduced. The second row in Table 2 shows results for a structure where
additional spherical sites are introduced between the membrane anchor and the binding site to
model the space occupied by a cytosolic protein sequence; this steric effect results in a small

decrease in $k_{on}^{(mem)}$. This decrease is reversed when flexibility is introduced by allowing the
spherical sites to be pivot points in the third row of the Table; this is how disordered domains
may be modeled in SpringSaLaD. Overall, for these idealized structures and simple
dimerization, the fitted on-rate constants in Table 2 are relatively close to $k_{on}^{(h)}$.
357
358 <u>The appropriateness of mass action kinetics for reactions of binding sites tethered to a</u>
359 <u>membrane.</u>
360
Qualitatively, one can consider the overall rate coefficient, k_{on}, to be affected by the intrinsic
binding occurring once the sites are close enough to bind and characterized by the rate
constant k_0, and by diffusion-influenced reactant encounters characterized by their
corresponding rate coefficient k_D. In 3D, k_D is a time-independent constant (12), and so is k_{on},
which in this case is computed as $k_{on} = \frac{k_D k_0}{k_D + k_0}$ (13). As discussed in the Introduction and in the
previous section in connection with the third panel of Figure 2, simple mass action kinetics are
not always appropriate for bimolecular reactions within a surface because k_D becomes a
function of time (8,12,13). Indeed, in the diffusion-limited regimes with $k_D(t) \ll k_0$, the k_{on} will
be largely determined by $k_D(t)$ and thus will depend on time, so a mass action rate law will be a
poor fit to the kinetics. We now wish to analyze our tethered binder reactions for their
conformance to mass action, using the criteria proposed by Yogurtcu and Johnson (10). Full
mathematical details of our approach are found in Supporting Text II; we will provide a summary
here.
374
For the dimerization of binding sites anchored to the membrane with a single simple tether, the
intrinsic binding rate constant is $k_0 = k_{on}^{(h)}$. For cases of short tethers with lengths that are
significantly less than the average initial distance between the anchors (exemplified by Table 1),
the effective reaction radii are determined by tether lengths h, and the effective diffusion
coefficient is determined by the diffusivity of anchors D_{mem}. Also among the factors affecting
possible deviations from simple mass action are the reactant densities (or the degrees of
reaction completion), as discussed above in relation to the development of depletion zones.
Thus, for a given combination of tether length, diffusion coefficient, effective 2D on-rate
constant, initial surface density and desired degree of completion for a fit, we need to define a
metric that describes how well k_{on} may be approximated as a constant (i.e. by $k_{on}^{(mem)}$).
385
As a first step, we needed to determine the
general functional form of $k_D(t)$, which is the
source of the time dependence of k_{on}. To do
this, we solved the Smoluchowski model (12)
numerically in 2D to produce the relation
shown in Fig. 3. The results are plotted using
dimensionless variables to allow us to
determine k_D as a function of time for any
combination of D' (the sum of the D_{mem} for the
two reactants) and r_{AB} (the sum of the radii of
the two reactants). Full details of how Fig. 3
was calculated are found in Supporting Text II
in relation to Fig. S7, where the calculations
are also validated against short-time and long-
time asymptotic analytical solutions.
401

Figure 3. Time dependence of k_D in 2D.

With $k_D(t)$ in hand, we can now ask how much it contributes to k_{on} for a desired level of

completion, p, (e.g. 80% in Tables 1 and 2.); $p = 1 - \frac{\sigma(t_c)}{\sigma_0}$, where $\sigma(t)$ is the monomer surface

density and t_c is the time to the desired level of completion, which can be read off the kinetic

data for σ vs. t. (It is important to appreciate that t_c is smaller for higher initial surface density,

all other parameters being equal; therefore, the range of $k_D(t)$ that pertains is higher up the

initial segment of the plot in Fig. 3, increasing the contribution of k_0 to k_{on}). In general, we can

establish the relative importance of $k_D(t)$ by defining the metric δ as

$$\delta = (1 + \bar{k}_D(t_c)/k_0)^{-1} \ , \hspace{4cm} \text{Eq. 2}$$

where $\bar{k}_D(t_c) = \frac{1}{t_c}\int_0^\infty k_D(t) \exp\left(-\frac{t}{t_c}\right) dt$ is, in effect, a weighted average of $k_D(t)$ over the time

interval between $t = 0$ and $t = t_c$. Thus, Eq 2 is indeed a measure of deviation from mass-

action kinetics k_0: if δ is close to 1, $\bar{k}_D(t_c) \ll k_0$, i.e. the reaction is diffusion-limited and k_{on} will

not be constant; conversely, if δ is small, the kinetics will be dominated by k_0 and will be well

approximated by mass action. Eq. 2 is fully derived in Supporting Text II, where it is Eq. S11.

Values of δ are included in Table 1 for all the irreversible reactions; these were calculated using

$k_0 = k_{on}^{(h)}$ and $r_{AB} = 2h$, as discussed above (see also Supporting Text I). As illustrated in Figure

2, the parameters associated with row #5 of Table 1 produce kinetics in SpringSaLaD that are

not approximated well by the mass-action rate law; the corresponding value of δ, 0.78 is indeed

close to 1. We judge a value of $\delta < 0.3$ to allow for the application of 2D mass action as a

reasonable approximate rate law, but individual judgment should be made on how critically the

overall system behavior might depend on conformity to mass action kinetics.

In Supporting Text II, we also show that a weighted average of the overall on-rate constant,

$\bar{k}_{on}(t_c) = \frac{1}{t_c}\int_0^\infty k_{on}(t) \exp\left(-\frac{t}{t_c}\right) dt,$ can be computed as

$$\bar{k}_{on}(t_c) = \bar{k}_D(t_c)k_0/(\bar{k}_D(t_c) + k_0), \hspace{3cm} \text{Eq. 3}$$

which is structurally similar to the equation for k_{on} in 3D. Using Eq. 3, we obtained a gratifyingly

good match for $k_{on}^{(mem)}$, derived by fitting the SpringSaLaD simulations.

In the case of long tethers, $h > (\pi\sigma_0)^{-\frac{1}{2}}$, exemplified by Table 2, geometric constraints imposed

by tethers and slow anchor diffusion are less limiting in two respects. First, multiple binding sites

may now encounter each other under the hemisphere of radius h. As a result, the tether length

no longer represents the effective reaction radius, which should now be determined by the

assigned radii of the binding sites $\frac{d}{2}$. Second, the main mixing mechanism is now volumetric

diffusion. Because the area of the hemisphere is twice the area of its projection on the

membrane, and both are covered by diffusion within the same time, the corresponding effective

2D diffusivity is $\frac{D_{vol}}{2}$. Thus, to compute δ from Eq. 2 for the case of long tethers, we should

compute $k_D(t)$ from the relation of Fig. 3 with $r_{AA} = d$ and $D' = D_{vol}$. The resultant δ for the first

row of Table 2 is 0.03, indicating that mass action is a good approximation. The other rows of

Table 2 also are clearly well fitted by mass action (see the fits in the spreadsheets in Supporting

Information), but we did not calculate δ because $k_{on}^{(h)}$ is not as well defined when the structures

become more complex than a single tethered sphere. This is because both steric effects and
excluded volume effects will impact binding kinetics in these cases.

In general, metrics like δ will not be easy or possible to calculate for complex structures based
on real molecules. Thus, in general the best way to assess the applicability of mass action as a
reasonable approximation for membrane-tethered reactions is to judge the fit of the
SpringSaLaD simulation to the integrated mass-action rate law. As can be seen by comparing
the plots in Fig. 2 and the corresponding entries in Table 1, it is clear that the fits for rows 1 and
6 are good, while the fit for row 5 is poor, consistent with both the entries for δ and the relative
squared deviation of the fit. But even a poor fit to mass action may be an adequate
approximation for some situations, especially where the uncertainty in other parameters of a
large signaling system can be large. Additionally, for reversible binding, the steady state
distribution is not dependent on the accuracy of mass action kinetics. We now illustrate our
method by applying it to a real signaling module that has been widely investigated because of
its importance in cancer biology.

<u>Application of the method to interaction of receptor-bound SOS with Ras.</u>

Until now, we have employed idealized molecular structures to validate our method and to learn
more about biophysical principles that control the on-rate constants of binding sites tethered to
membranes. We now illustrate the application of this approach to a biologically relevant
example, namely the interaction of the G-protein exchange factor (GEF) SOS with the lipid
anchored small G-protein Ras (22).

SOS has two binding sites for Ras: an allosteric site and a catalytic site. When a Ras molecule
binds to the allosteric site it increases the GEF activity of the catalytic site (36). Additionally,
before SOS binds to Ras it is first recruited to an active receptor tyrosine kinase (RTK) through
an adapter protein; the adaptor binds to a proline rich motif (PRM) on SOS via a SH3 domain
and to a phosphorylated tyrosine via a SH2 domain. One such RTK is the Epidermal Growth
Factor Receptor (EGFR) and one such adaptor protein is Grb2 (37). Once SOS is bound to
Grb2, it becomes membrane-tethered and its interaction with Ras is facilitated (36). However
the complex mechanistic details of this system are still emerging (38).

We asked the limited question of how binding of Ras with the receptor-associated SOS catalytic
site might depend on whether SOS is prebound to Ras at the allosteric site. Ras is a lipid-
anchored protein, so we reasoned that binding of Ras to the allosteric site of SOS would bring
the SOS catalytic site closer to the membrane to enhance binding to a second Ras and
subsequent exchange of GDP for GTP. Just how large an effect this is, may be estimated by the
procedure developed above, with the results shown in Fig. 4.

Figure 4. Membrane binding of Ras to the catalytic site of Receptor-bound SOS. Top: direct binding. Bottom: after pre-binding of Ras to the allosteric site. The molecular structures are approximated by using the SpringSaLaD 3D editing utility based on atomic structures derived from AlphaFold2. The top left structure is an EGFR cytoplasmic domain anchored to the membrane (red kinase domain, followed by yellow disordered tail capped by a phosphotyrosine in green); the latter is linked to a cyan SH2 domain in Grb2; one of its magenta SH3 domains is linked to an olive PRM on the end of the disordered region of SOS; the violet SOS binding site for the catalytic domain of Ras is indicated with an asterisk (*). The bottom left structure is identical, except that the pink allosteric site on SOS is prebound to Ras. The Ras structures are shown in the center with the yellow binding sites indicated by an asterisk. The input rate constants for the SpringSaLaD simulations are shown at the top, corresponding to the volumetric on rate for Ras binding to the catalytic site of SOS. The EGFR anchor diffusion coefficient is 0.01µm^2/s. All other site diffusion coefficients are 1.0µm^2/s. For each condition, 20 EGFR-Grb2-SOS molecules react with 200 Ras molecules on a 250nmX250nm membrane surface to generate 100 SpringSaLaD trajectories. Their means were fitted to a deterministic 2D rate law to derive $k_{on}^{(mem)}$, fixing k_{off} at 4.0 s^{-1}; results for the 2 conditions are shown on the right.

480
We developed molecular models with the aid of the mol2sphere (25) utility within SpringSaLaD
and were guided by AlphaFold 2 atomic structure predictions (23,24); all the site diameters and
linker lengths are available in the SpringSaLaD input file included in the Supporting Information;
snapshots of the structure are shown in Fig. 4. The top of Fig. 4 displays results for binding of
SOS-Grb2-EGFR to Ras at the SOS catalytic site; the bottom shows results for the same
reaction, except SOS-Grb2-EGFR had been first bound to a Ras molecule at the SOS allosteric
site. The input rates shown at the top of Fig. 4 are based on experimentally measured data (22)
and are applied to both of the reactions. Special considerations were applied to ensure
accuracy of the on rate, which was parameterized via an initial cytosolic reaction where the
output rate constant was matched to the experimentally measured rate data (22). This important

step allows us to calibrate the coarse-grained protein structure used in our model to real protein
structures, which inherently impact binding *in-vivo* and are therefore reflected in experimentally
measured reaction rates. The desired output rate corresponding to an experimentally observed
rate constant of 0.27 μM^{-1}s^{-1} (22), was achieved with an input rate constant of 0.35 μM^{-1}s^{-1},
which was then used in subsequent simulations of membrane bound interactions in Fig. 4. In
these models, the EGFR membrane anchor site is assigned a diffusion coefficient of 0.01 μm^2/s
to represent a large transmembrane protein, while the Ras membrane anchor is assigned a
diffusion coefficient of 1.0 μm^2/s to represent a lipid anchor; all the sites that are dangling in the
cytosol volume are given D_{vol} of 1.0 μm^2/s, but since the binding reaction is not close to
diffusion-limited, the precise values are not critical. Consistent with these being reaction-limited
on-rates, the SpringSaLaD simulation outputs (averages of 100 runs) are very well fitted to
reversible mass action kinetic law, as shown in the plots on the right of Fig. 4 (relative squared
deviations are, respectively, 4.1 X 10^{-4} and 2.0 X 10^{-4}). Importantly, the 2D on-rate constants
($k_{on}^{(mem)}$) derived from these fits are, respectively, 5.0 X 10^{-3} μm^{2}molecules^{-1}s^{-1} and 1.9 X 10^{-2}
μm^{2}molecules^{-1}s^{-1}. Likewise, the affinity of the catalytic site is increased by allosteric site pre-
association: K_d = 800 molecules/μm^2 for the top of Fig. 3 and 210 molecules/μm^2 for the bottom
pre-association case. Thus, SOS allosteric site association with Ras is estimated to accelerate
its catalytic site binding and affinity by a factor of ~4 – even when SOS is already confined to the
membrane through Grb2-mediated association with EGFR.
510
511   <u>A simple ODE model of receptor-mediated signaling</u>
512
To illustrate the application of our approach, we built a small signaling model based on the
values of $k_{on}^{(mem)}$ determined in the previous section. The ODE model was built in VCell (29,30)
and total concentrations of the various species were from measurements on HeLa cells (26,27).
In addition to the two reactions from Fig. 4, we also modeled recruitment of Grb_SOS from the
cytosol to the membrane by phosphorylated EGFR (EGFRPTyr) and direct stepwise binding of
two Ras molecules to Grb_SOS; the model assumes that almost all the cytosolic SOS is
prebound to Grb2, which is present in excess (39). The VCell reaction diagram in Fig. 5A
displays the connectivity of the network and Table 3 provides the corresponding mass action
rate constants.
522

Figure 5. A simple VCell model and simulation results for EGFR signaling to Ras. A. Reaction network identifying all the variables (see also the corresponding reactions in Table 3.). B. The kinetics reach steady state within 30s. The primary plot shows all the membrane species and the inset shows Grb_SOS in the cytosol. Steady state surface densities of the 4 different species of 1 or 2 Ras bound to Grb_SOS or EGFRPtyr_Grb_SOS under 3 conditions: "Basal", "Partial activation" and "Full activation" corresponding respectively to 0, 200 and 840 molecules/μm^2 initial EGFRPtyr.

We used the data for EGFR measured in HeLa cells (26) to derive an initial surface density for
EGFPtyr of 840 molecules/μm^2, which we took to be the fully activated receptor (we did not
attempt to account for kinetics of inactivation, which would lower this number, or for the multiple
phosphorylation sites available on the cytoplasmic domain of EGFR, which would raise this
number). Fig. 5B shows the approach to steady state for this "Full activation" condition. It shows
that free EGFPtyr and free Ras are only partially depleted at steady state, while ~90%
Grb_SOS is removed from the cytosol to become membrane associated. Figure 5C provides a
more detailed look at the association of Grb_SOS with Ras. We chose three levels of initial
EGFRPTyr, 0, 200 and 840 molecules/μm^2, to model, respectively, basal unstimulated cells,
partial activation of EGFR and full activation. The species composed of doubly bound Ras,
Grb_SOS_Ras_Ras and EGFR_Grb_SOS_Ras_Ras (GSRR and EGSRR in Fig. 5), are always
a factor of ~3.5 higher than the corresponding species with one Ras. This is important because
the binding two Ras molecules is required for SOS to optimally express its GEF activity (40); the

second binding event is, of course, significantly enhanced by the proximity effect quantified in
Fig. 4; this enhancement at the membrane has been reported based on experiments for the
binding of Ras to SOS alone (41). Another conclusion from Fig. 4C is that the overall levels of
Ras doubly bound to SOS are significantly increased by EGFR activation; this is consistent with
the high affinity of the SH2 domains on Grb2 for phosphotyrosine (42), creating a higher local
concentration of SOS near the membrane surface. Indeed, under the "Basal" condition without
participation of EGFRPtyr, 80% of the Grb_SOS remains unbound in the cytosol (data not
shown but available in the public VCell Model "SOS_Recruitment_Ras_Binding").
544
545

Table 3. Rate constants for the VCell model			
Reaction	$\textbf{\textit{k}}_{\textit{on}}$	$\textbf{\textit{k}}_{\textit{off}}$	**Source**
EGFRPtyr + Grb_SOS -> EGFR_Grb_SOS	1.0 $\mu M^{-1}s^{-1}$	0.3 s^{-1}	(42)
Grb_SOS + Ras -> Grb_SOS_Ras	0.27 $\mu M^{-1}s^{-1}$	4.0 s^{-1}	(22)
EGFR_Grb_SOS + Ras -> EGFR_Grb_SOS_Ras	0.005 $\mu m^2 molecule^{-1}s^{-1}$	4.0 s^{-1}	(22) with Fig. 4
Ras + EGFR_Grb_SOS_Ras -> EGFR_Grb_SOS_Ras_Ras	0.019 $\mu m^2 molecule^{-1}s^{-1}$	4.0 s^{-1}	(22) with Fig. 4
Ras + Grb_SOS_Ras -> Grb_SOS_Ras_Ras	0.019 $\mu m^2 molecule^{-1}s^{-1}$	4.0 s^{-1}	(22) with Fig. 4

546
547
548   **Discussion**
549
The kinetics of reactions at membranes have long fascinated biophysicists
(1,5,7,8,10,11,17,19,32). These studies have produced theoretical insights to illuminate how
surface-associated reactions have distinct properties compared with reactions occurring in 3D
solution. Which of these special properties are most pertinent to any given membrane-bound
molecular interaction is difficult to ascertain *a priori*. Furthermore, experiments to measure
bimolecular kinetics on membrane surfaces are complex (4), so often only on-rate constants
measured in 3D are accessible. Fundamentally, however, the kinetics of key membrane-
associated reactions depend on the 2D surface densities and 2D rate constants, not on the bulk
cellular concentrations and 3D rate constants. Indeed, because surface to volume ratios of
different cell types vary tremendously, volumetric rate constants cannot be readily used to model
and simulate cell signaling systems. To address these theoretical and practical problems, we
describe a procedure (Fig. 1) using experimentally accessible volumetric on-rate constants,
$k_{on}^{(vol)}$ within the SpringSaLaD simulation software to estimate the $k_{on}^{(mem)}$, the 2-dimensional rate
constant for a membrane-confined bimolecular reaction.
564
To validate the method, we applied it to the dimerization of a single binding site tethered to a
surface through a 5nm stiff linker, where the membrane anchor acts as a pivot (Table 1). For the
situation where the reaction is rate limiting, this system can be solved theoretically (see
Supporting Text I); gratifyingly, $k_{on}^{(mem)}$ determined by our method is well reproduced by this
solution. Interestingly, for these cases, $k_{on}^{(mem)}$ is well approximated by $k_{on}^{(h)} = k_{on}^{(vol)}/h/602.2$
($\mu m^2 molecules^{-1}s^{-1}$), where h is the distance of the binding site from the membrane anchor (in
μm) and 602.2 is a units conversion factor. The parameter h has also been referred to as the

"confinement length" (17), defining a thin volume above the membrane that concentrates the
binding sites and directly producing the relationship between $k_{on}^{(h)}$ and $k_{on}^{(vol)}$.
574
While mass action kinetics are generally applicable for both encounter-limited and reaction-
limited kinetics in 3D solution (but see ((12)15)), it has long been appreciated that the situation
may be more complex for 2D kinetics (5,8,10,11,32,43). This is demonstrated by the results in
Table 1 for situations where the diffusion coefficient of the anchor is slow, but the volumetric on-
rate constant is fast. For these cases, different estimates of $k_{on}^{(mem)}$ are obtained at different
initial surface density – clearly incompatible with the mass action kinetics. Indeed, the third
panel of Figure 2 shows that the SpringSaLaD kinetic data is not well fit by a 2D mass action
rate law. A video of one of these trajectories (Movie 1) nicely illustrates how the initial rate is
fast, while the binding sites are within "reach" (19), but falls off as binding sites are left orphaned
outside the reach of the remaining slowly diffusing monomers. Because of this, a mass action
rate law applies better to high initial concentrations and to shorter time durations, before
depletion zones develop. To address the applicability of mass action theoretically, we solved the
Smoluchowski model (12) in 2D using VCell (Supporting Text II) to generate the relation plotted
in Fig. 3 for the time-dependence of the diffusion limited reaction coefficient. This allowed us to
implement a pipeline for computing a weighted average of the on-rate coefficient over the time
to a particular level of reaction completion, for any intrinsic on-rate, k_0, membrane diffusion
coefficient, initial surface density, and desired level of completion for a fit. We used this to see
how well the idealized scenarios in the first 8 rows of Table 1 and the first row of Table 2 could
be fitted by simple mass action (i.e. where the rate coefficients can be reasonably well
approximated as independent of time). We were able to derive a metric, δ, that measures how
significantly the overall rate is influenced by diffusion. This is similar to the detailed analysis of
Yogurtcu and Johnson (10) for 2D reactions, except their analysis focused on pure 2D (i.e. $h =$
0). Our results allow us to generalize that mass action applies as long as $k_{on}^{(mem)}$ is close to
$k_{on}^{(h)}$. Importantly, there was a clear relationship between how well the SpringSaLaD simulation
results can be approximated with a mass action rate law, as measured by the relative squared
deviations and δ in Table 1. Thus, our analysis provides an approach to determine whether 2D
binding might be well approximated by mass action kinetics, and if so, to estimate the 2D mass
action on-rate constant.
603
To explore how other molecular structural features might affect dimerization of the monomers
tethered to the membrane, we looked at three additional idealized systems in Table 2. In all
these, *h* was 20.5nm (as opposed to 5.5nm in Table 1). As expected, the longer confinement
length decreased the estimated $k_{on}^{(mem)}$ by a factor of ~4. The insertion of steric sites between
the anchor and the binding site or allowing for flexibility of the linker region have minor effects
on $k_{on}^{(mem)}$, which is relatively well approximated by $k_{on}^{(h)}$. Importantly, the lower value of $k_{on}^{(h)}$ for
this system was closer to D_{mem}, resulting in a better fit by the mass action kinetics than for the
similar case with the 5nm linker (fifth row of Table 1).
612
For membrane binding of real biological molecules, the average location of binding sites relative
to the membrane surface (i.e. *h*) will generally be not well defined, since it is influenced by the
steric effects and the variable flexibility of neighboring protein domains. Also, the two binding
sites may be parts of very different structures with different distances from the membrane
surface. In situations like this, our approach has the potential to provide good estimates of rate
constants that can be applied to larger cell signaling systems. Indeed, there may be direct
insights that can be realized just by considering the structural details of the interacting
membrane molecules. We have illustrated this in relation to adaptor-mediated protein kinase

receptor signaling mechanisms, specifically for the interaction of the GEF SOS with its effector Ras (Fig. 4); the fit of the SpringSaLaD simulations to a mass action rate law is excellent in both scenarios of Fig. 4, indicating that mass action is applicable. It has been shown that direct catalysis by the SOS catalytic domain of Ras conversion from the GDP to the GTP states is relatively slow. However, binding of Ras to SOS at a site that is not catalytic (termed the "allosteric" site on SOS) significantly accelerates the catalytic activity(40,41), where the catalysis becomes processive (22,36,38). The results in Fig. 4 suggest that at least part of this acceleration may be due to the close proximity of the SOS catalytic site to the membrane once it is bound to Ras at its allosteric site. Even though SOS is already localized to the membrane by initially binding to EGFR via Grb2 in our computational experiment, pre-binding of the SOS allosteric site to Ras brings it even closer to the membrane. Of course, there could be additional effects such as a direct allosteric enhancement through a conformational change or release of self-inhibition (22,36,38), but here we focus on the significance of constricting the binding zone through membrane tethers of varying length and flexibility.

To illustrate the application of our approach to cell signaling, we constructed a simple model of EGFR signaling to Ras via SOS in HeLa cells (Fig. 5). The results show that the proximity effect established in Fig. 4 for the binding of SOS to the second Ras molecule results in the dominance of doubly bound Ras species. This no doubt contributes to the efficiency of the GEF activity associated with SOS. Our simple model also demonstrates the effectiveness of active EGFR in boosting the recruitment of Grb_SOS to the membrane where it presents an effectively high local concentration to Ras. The model, which is available in the VCell database, can be further explored for the effect of varying Ras or SOS concentrations or the behavior of different cell types with different levels of the key signaling models; it could also be used as a starting point for more elaborate models that incorporate more detailed mechanisms and additional downstream signaling events.

Thus, our approach toward estimating membrane on-rates will aid in the parametrization of ODE and PDE models that could help elucidate the full kinetic and mechanistic details of cell signaling systems. In ODE models, diffusion is assumed to be fast on the timescale of reaction rates; however, our approach is necessary to capture the special local membrane effects that allow conversion of a bulk measurement of the on-rate to $k_{on}^{(mem)}$. In cell-scale PDE (or spatial stochastic) models, where diffusion is considered explicitly, the membrane molecules are treated as points and the distances are very large on the scale of molecular dimensions. So the most appropriate diffusion coefficient would be that of the membrane anchor. These are typically measured experimentally using fluorescence recovery after photobleaching (FRAP) or fluorescence correlation spectroscopy (FCS). It is also possible to estimate membrane diffusion coefficients based on the size of the membrane embedded region and the known viscosity of biological membranes.

Author contributions
KJP created the computational models, performed the simulations, analyzed the data and wrote the paper; BMS performed the calculations in the Supporting Text I and II and wrote the paper; LML conceived the research, analyzed the data and wrote the paper.

Declaration of interests
The authors declare no competing interests.

Acknowledgments

This work was supported by NIH grants R24 GM137787 and R01 GM132859. We are pleased to acknowledge the advice of Aniruddha Chattaraj with some of the data analysis.

1. Haugh, J. M. 2002. A Unified Model for Signal Transduction Reactions in Cellular Membranes. *Biophys. J.* 82(2):591-604, http://www.biophysj.org/cgi/content/abstract/82/2/591.

2. Slepchenko, B. M., J. C. Schaff, I. Macara, and L. M. Loew. 2003. Quantitative cell biology with the Virtual Cell. *Trends Cell Biol.* 13(11):570-576, doi: S0962892403002150 [pii], http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14573350.

3. Eungdamrong, N. J., and R. Iyengar. 2004. Modeling Cell Signaling Networks. *Biology of the Cell*. 96(5):355-362, doi: https://doi.org/10.1016/j.biolcel.2004.03.004, https://www.sciencedirect.com/science/article/pii/S0248490004000772.

4. Groves, J. T., and M. L. Dustin. 2003. Supported planar bilayers in studies on immune cell adhesion and communication. *Journal of Immunological Methods*. 278(1):19-32, doi: https://doi.org/10.1016/S0022-1759(03)00193-5, https://www.sciencedirect.com/science/article/pii/S0022175903001935.

5. Adam, G., and M. Delbrück. 1968. Reduction of Dimensionality in Biological Diffusion. In Structural Chemistry and Molecular Biology. A. Rich, and N. Davidson, editors. W. H. Freeman and Co., San Francisco, pp. 198-215.

6. Pólya, G. 1921. Über eine Aufgabe der Wahrscheinlichkeitsrechnung betreffend die Irrfahrt im Straßennetz. *Mathematische Annalen*. 84(1):149-160, doi: 10.1007/BF01458701, https://doi.org/10.1007/BF01458701.

7. McCloskey, M. A., and M. Poo. 1986. Rates of Membrane-associated Reactions: Reduction of Dimentionality Revisited. *Journal of Cell Biology*. 102:88-96.

8. Torney, D. C., H. M. McConnell, and G. R. Porter. 1983. Diffusion-limited reaction rate theory for two-dimensional systems. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*. 387(1792):147-170, doi: doi:10.1098/rspa.1983.0055, https://royalsocietypublishing.org/doi/abs/10.1098/rspa.1983.0055.

9. Koudriavtsev, A. B., R. F. Jameson, W. Linert, and SpringerLink. 2001. The Law of Mass Action. Springer Berlin Heidelberg : Imprint: Springer, Berlin, Heidelberg.

10. Yogurtcu, O. N., and M. E. Johnson. 2015. Theory of bi-molecular association dynamics in 2D for accurate model and experimental parameterization of binding rates. *The Journal of Chemical Physics*. 143(8), doi: 10.1063/1.4929390, https://doi.org/10.1063/1.4929390.

11. Axelrod, D., and M. D. Wang. 1994. Reduction-of-dimensionality kinetics at reaction-limited cell surface receptors. *Biophys J*. 66(3 Pt 1):588-600, doi: 10.1016/s0006-3495(94)80834-3, https://www.ncbi.nlm.nih.gov/pubmed/8011892.

12. Keizer, J. 1987. Diffusion effects on rapid bimolecular chemical reactions. *Chemical Reviews*. 87(1):167-180, doi: 10.1021/cr00077a009, https://doi.org/10.1021/cr00077a009.

13. Collins, F. C., and G. E. Kimball. 1949. Diffusion-controlled reaction rates. *Journal of Colloid Science*. 4(4):425-437, doi: https://doi.org/10.1016/0095-8522(49)90023-9, https://www.sciencedirect.com/science/article/pii/0095852249900239.

14. Szabo, A. 1989. Theory of diffusion-influenced fluorescence quenching. *The Journal of Physical Chemistry*. 93(19):6929-6939, doi: 10.1021/j100356a011, https://doi.org/10.1021/j100356a011.

15. Windisch, B., D. Bray, and T. Duke. 2006. Balls and Chains—A Mesoscopic Approach to Tethered Protein Domains. *Biophysical Journal*. 91(7):2383-2392, doi: https://doi.org/10.1529/biophysj.105.078543, https://www.sciencedirect.com/science/article/pii/S0006349506719531.

16. Bell, G. I., M. Dembo, and P. Bongrand. 1984. Cell adhesion. Competition between nonspecific repulsion and specific bonding. *Biophys J*. 45(6):1051-1064, doi: 10.1016/S0006-3495(84)84252-6, https://www.ncbi.nlm.nih.gov/pubmed/6743742.

17. Wu, Y., J. Vendome, L. Shapiro, A. Ben-Shaul, and B. Honig. 2011. Transforming binding affinities from three dimensions to two with application to cadherin clustering. *Nature*. 475(7357):510-513, doi: 10.1038/nature10183, https://doi.org/10.1038/nature10183.

18. Xie, Z.-R., J. Chen, and Y. Wu. 2014. Linking 3D and 2D binding kinetics of membrane proteins by multiscale simulations. *Protein Science*. 23(12):1789-1799, doi: https://doi.org/10.1002/pro.2574, https://onlinelibrary.wiley.com/doi/abs/10.1002/pro.2574.

19. Zhang, Y., L. Clemens, J. Goyette, J. Allard, O. Dushek, and S. A. Isaacson. 2019. The Influence of Molecular Reach and Diffusivity on the Efficacy of Membrane-Confined Reactions. *Biophys J*. 117(7):1189-1201, doi: 10.1016/j.bpj.2019.08.023.

20. Michalski, P. J., and L. M. Loew. 2016. SpringSaLaD: A Spatial, Particle-Based Biochemical Simulation Platform with Excluded Volume. *Biophys J*. 110(3):523-529, doi: 10.1016/j.bpj.2015.12.026, http://www.ncbi.nlm.nih.gov/pubmed/26840718.

21. Chattaraj, A., M. Youngstrom, and L. M. Loew. 2018. The interplay of structural and cellular biophysics controls clustering of multivalent molecules. *bioRxiv (accepted in Biophys. J.)*.373084, doi: 10.1101/373084, https://www.biorxiv.org/content/biorxiv/early/2018/12/26/373084.full.pdf.

22. Iversen, L., H.-L. Tu, W.-C. Lin, S. M. Christensen, S. M. Abel, J. Iwig, H.-J. Wu, J. Gureasko, C. Rhodes, R. S. Petit, S. D. Hansen, P. Thill, C.-H. Yu, D. Stamou, A. K. Chakraborty, J. Kuriyan, and J. T. Groves. 2014. Ras activation by SOS: Allosteric regulation by altered fluctuation dynamics. *Science*. 345(6192):50-54, doi: 10.1126/science.1250373, https://science.sciencemag.org/content/sci/345/6192/50.full.pdf.

23. Abramson, J., J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L. Willmore, A. J. Ballard, J. Bambrick, S. W. Bodenstein, D. A. Evans, C.-C. Hung, M. O'Neill, D. Reiman, K. Tunyasuvunakool, Z. Wu, A. Žemgulytė, E. Arvaniti, C. Beattie, O. Bertolli, A. Bridgland, A. Cherepanov, M. Congreve, A. I. Cowen-Rivers, A. Cowie, M. Figurnov, F. B. Fuchs, H. Gladman, R. Jain, Y. A. Khan, C. M. R. Low, K. Perlin, A. Potapenko, P. Savy, S. Singh, A. Stecula, A. Thillaisundaram, C. Tong, S. Yakneen, E. D. Zhong, M. Zielinski, A. Žídek, V. Bapst, P. Kohli, M. Jaderberg, D. Hassabis, and J. M. Jumper. 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature*. 630(8016):493-500, doi: 10.1038/s41586-024-07487-w, https://doi.org/10.1038/s41586-024-07487-w.

24. Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Žídek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A. Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P. Kohli, and D. Hassabis. 2021. Highly accurate protein structure prediction with AlphaFold. *Nature*. 596(7873):583-589, doi: 10.1038/s41586-021-03819-2, https://doi.org/10.1038/s41586-021-03819-2.

25. Masison, J., P. J. Michalski, L. M. Loew, and A. D. Schuyler. 2018. mol2sphere: spherical decomposition of multi-domain molecules for visualization and coarse grained spatial

modeling. *Bioinformatics*. 34(22):3948-3950, doi: 10.1093/bioinformatics/bty487,
https://www.ncbi.nlm.nih.gov/pubmed/29931043.

26.   Kamioka, Y., S. Yasuda, Y. Fujita, K. Aoki, and M. Matsuda. 2010. Multiple decisive
phosphorylation sites for the negative feedback regulation of SOS1 via ERK. *J Biol*
Chem. 285(43):33540-33548, doi: 10.1074/jbc.M110.135517,
https://www.ncbi.nlm.nih.gov/pubmed/20724475.

27.   Fujioka, A., K. Terai, R. E. Itoh, K. Aoki, T. Nakamura, S. Kuroda, E. Nishida, and M.
Matsuda. 2006. Dynamics of the Ras/ERK MAPK Cascade as Monitored by Fluorescent
Probes*. *Journal of Biological Chemistry*. 281(13):8917-8926, doi:
https://doi.org/10.1074/jbc.M509344200,
https://www.sciencedirect.com/science/article/pii/S0021925819565978.

28.   Hoops, S., S. Sahle, R. Guages, C. Lee, J. Pahle, N. Simus, M. Singhal, L. Xu, P.
Mendes, and U. Kummer. 2006. COPASI: a COmplex PAthway SImulator.
Bioinformatics. 22:3067-3074, doi: 10.1093/bioinformatics/btl485,
https://doi.org/10.1093/bioinformatics/btl485.

29.   Schaff, J., C. C. Fink, B. Slepchenko, J. H. Carson, and L. M. Loew. 1997. A general
computational framework for modeling cellular structure and function. *Biophys J*.
73(3):1135-1146, doi: 10.1016/S0006-3495(97)78146-3,
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
&list_uids=9284281.

30.   Cowan, A. E., Moraru, II, J. C. Schaff, B. M. Slepchenko, and L. M. Loew. 2012. Spatial
modeling of cell signaling networks. *Methods Cell Biol*. 110:195-221, doi: 10.1016/B978-
0-12-388403-9.00008-4, http://www.ncbi.nlm.nih.gov/pubmed/22482950.

31.   Kopelman, R. 1986. Rate processes on fractals: Theory, simulations, and experiments.
Journal of Statistical Physics. 42(1-2):185-200, doi: 10.1007/BF01010846,
https://www.scopus.com/inward/record.uri?eid=2-s2.0-
30244438013&doi=10.1007%2fBF01010846&partnerID=40&md5=8220d250ae1bd74cb
14dad318176663e (Article).

32.   Berry, H. 2002. Monte Carlo Simulations of Enzyme Reactions in Two Dimensions:
Fractal Kinetics and Spatial Segregation. *Biophysical Journal*. 83(4):1891-1901, doi:
https://doi.org/10.1016/S0006-3495(02)73953-2,
https://www.sciencedirect.com/science/article/pii/S0006349502739532.

33.   Ovchinnikov, A. A., and Y. B. Zeldovich. 1977. Role of density fluctuations in bimolecular
reaction kinetics. *Chemical Physics*. 28:215-218.

34.   Toussaint, D., and F. Wilczek. 1983. Particle–antiparticle annihilation in diffusive motion.
The Journal of Chemical Physics. 78(5):2642-2647, doi: 10.1063/1.445022,
https://doi.org/10.1063/1.445022.

35.   Tachiya, M. 1983. Theory of diffusion-controlled reactions: Formulation of the bulk
reaction rate in terms of the pair probability. *Radiation Physics and Chemistry (1977)*.
21(1):167-175, doi: https://doi.org/10.1016/0146-5724(83)90143-7,
https://www.sciencedirect.com/science/article/pii/0146572483901437.

36.   Bandaru, P., Y. Kondo, and J. Kuriyan. 2019. The Interdependent Activation of Son-of-
Sevenless and Ras. *Cold Spring Harb Perspect Med*. 9(2), doi:
10.1101/cshperspect.a031534, https://www.ncbi.nlm.nih.gov/pubmed/29610148.

37.   Jorissen, R. N., F. Walker, N. Pouliot, T. P. J. Garrett, C. W. Ward, and A. W. Burgess.
2003. Epidermal growth factor receptor: mechanisms of activation and signalling.
Experimental Cell Research. 284(1):31-53, doi: https://doi.org/10.1016/S0014-
4827(02)00098-8, http://www.sciencedirect.com/science/article/pii/S0014482702000988.

38.   Ren, H., A. A. Lee, L. J. N. Lew, J. B. DeGrandchamp, and J. T. Groves. 2024. Positive
feedback in Ras activation by full-length SOS arises from autoinhibition release

mechanism. *Biophysical Journal*. 123(19):3295-3303, doi: 10.1016/j.bpj.2024.07.014,
https://doi.org/10.1016/j.bpj.2024.07.014.
39. Giubellino, A., T. R. Burke, and D. P. Bottaro. 2008. Grb2 signaling in cell motility and
cancer. *Expert Opinion on Therapeutic Targets*. 12(8):1021-1033, doi:
10.1517/14728222.12.8.1021, https://doi.org/10.1517/14728222.12.8.1021.
40. Margarit, S. M., H. Sondermann, B. E. Hall, B. Nagar, A. Hoelz, M. Pirruccello, D. Bar-
Sagi, and J. Kuriyan. 2003. Structural Evidence for Feedback Activation by Ras·GTP of
the Ras-Specific Nucleotide Exchange Factor SOS. *Cell*. 112(5):685-695, doi:
https://doi.org/10.1016/S0092-8674(03)00149-1,
https://www.sciencedirect.com/science/article/pii/S0092867403001491.
41. Gureasko, J., W. J. Galush, S. Boykevisch, H. Sondermann, D. Bar-Sagi, J. T. Groves,
and J. Kuriyan. 2008. Membrane-dependent signal integration by the Ras activator Son
of sevenless. *Nature Structural & Molecular Biology*. 15(5):452-461, doi:
10.1038/nsmb.1418, https://doi.org/10.1038/nsmb.1418.
42. Chook, Y. M., G. D. Gish, C. M. Kay, E. F. Pai, and T. Pawson. 1996. The Grb2-mSos1
complex binds phosphopeptides with higher affinity than Grb2. *J Biol Chem*.
271(48):30472-30478, doi: 10.1074/jbc.271.48.30472,
https://www.ncbi.nlm.nih.gov/pubmed/8940013.
43. Bell, G. I. 1978. Models for the Specific Adhesion of Cells to Cells. *Science*.
200(4342):618-627, doi: doi:10.1126/science.347575,
https://www.science.org/doi/abs/10.1126/science.347575.

841
842
843
844
845   **Figure Captions**
846
Figure 1. Workflow for finding membrane bimolecular binding rate constants ($k_{on}^{(mem)}$) in
terms of surface densities. Step 1 is to run 100 SpringSaLaD trajectories with the coarse
grained model of the binding partners. Illustrated are a pair of simple monomers (top) consisting
of 2nm diameter spheres (yellow) tethered to a 1nm membrane anchor sphere (gray) with a
5nm link; at the bottom, the product dimer is depicted. Step 2 consists of fitting the average of
100 outputs from stochastic SpringSaLaD simulations to a deterministic (ODE) non-spatial
model of surface-bound dimerization.
854
855   **Figure 2. Examples of SpringSaLaD simulation data and their fit to surface-confined**
856   **mass action dimerization kinetics.** The initial surface density of monomers in each case is
857   2500 molecule/μm^2. The diffusion coefficient of the anchor and the volumetric binding rate
858   constant of the binding sites are indicated above each graph, corresponding respectively to
859   rows 1, 6 and 5 of Table 1.
860
861   **Figure 3. Time dependence of k_D in 2D.**
862
863
Figure 4. Membrane binding of Ras to the catalytic site of Receptor-bound SOS. Top:
direct binding. Bottom: after pre-binding of Ras to the allosteric site. The molecular
structures are approximated by using the SpringSaLaD 3D editing utility based on atomic
structures derived from AlphaFold2. The top left structure is an EGFR cytoplasmic domain
anchored to the membrane (red kinase domain, followed by yellow disordered tail capped by a
phosphotyrosine in green); the latter is linked to a cyan SH2 domain in Grb2; one of its magenta

SH3 domains is linked to an olive PRM on the end of the disordered region of SOS; the violet
SOS binding site for the catalytic domain of Ras is indicated with an asterisk (*). The bottom left
structure is identical, except that the pink allosteric site on SOS is prebound to Ras. The Ras
structures are shown in the center with the yellow binding sites indicated by an asterisk. The
input rate constants for the SpringSaLaD simulations are shown at the top, corresponding to the
volumetric on rate for Ras binding to the catalytic site of SOS. The EGFR anchor diffusion
coefficient is 0.01µm^2/s. All other site diffusion coefficients are 1.0µm^2/s. For each condition, 20
EGFR-Grb2-SOS molecules react with 200 Ras molecules on a 250nmX250nm membrane
surface to generate 100 SpringSaLaD trajectories. Their means were fitted to a deterministic 2D
rate law to derive $k_{on}^{(mem)}$, fixing k_{off} at 4.0 s^{-1}; results for the 2 conditions are shown on the
right.
881
Figure 5. A simple VCell model and simulation results for EGFR signaling to Ras. A.
Reaction network identifying all the variables (see also the corresponding reactions in Table 3.).
B. The kinetics reach steady state within 30s. The primary plot shows all the membrane species
and the inset shows Grb_SOS in the cytosol. Steady state surface densities of the 4 different
species of 1 or 2 Ras bound to Grb_SOS or EGFRPtyr_Grb_SOS under 3 conditions: "Basal",
"Partial activation" and "Full activation" corresponding respectively to 0, 200 and 840
molecules/µm^2 initial EGFRPtyr.
889
890
List of Supporting Data and Information

Derivations for limiting cases related to Table 1 (Supporting Text I) and Derivations related to
Applicability of mass-action kinetics to dimerization of molecules with binding sites tethered to
the membrane. Filename: Supporting Text_revised.docx

SpringSaLaD input files for the models in Tables 1 and 2 and Figure 4. Filenames: Table 1 and 2
SIMS.zip; EGFR_Grb_SOS binding to Ras.txt; EGFR_Grb_SOS_prebound at allo binding to
Ras.txt

Spreadsheets related to Tables 1 and 2 and Figure 4 containing the SpringSaLaD simulation
results and the fits to determine $k_{on}^{(mem)}$. Filenames: Table 1 5nm_stiff_fits.xlsx; Table2
20nmT_fits.xlsx; Data for Figure 4.xlsx

Movie of an example trajectory for the model of Table 1 Row 5 (and also Fig. 2, 3rd panel).
Filename: Movie kon=1 Danchor=membrane protein.mp4