Evolution in response to prophage activation attenuates the virulence of culturable *Serratia symbiotica* relatives of aphid endosymbionts

Anthony J. VanDieren[1] and Jeffrey E. Barrick[1#]

[1]Department of Molecular Biosciences, Center for Systems and Synthetic Biology,

The University of Texas at Austin, Austin, Texas, USA

[#]Correspondence: jbarrick@cm.utexas.edu

Running title: *S. symbiotica* evolution following prophage activation

ABSTRACT

Serratia symbiotica bacteria exhibit a range of relationships with aphids. They may be co-obligate mutualists, commensals, or even pathogens depending on the strain, aphid host species, and environment. *Serratia symbiotica* CWBI-2.3^T (CWBI), a culturable member of this group, is transmitted to embryos transovarially when it is injected into pea aphids (*Acyrthosiphon pisum*), the same route used by *S. symbiotica* strains that are vertically inherited endosymbionts. Yet, aphids colonized with CWBI die before they give birth to infected offspring. We evolved laboratory populations of CWBI through 15-30 serial passages at two different temperatures in rich media. These conditions mimic aspects of the nutritional environment in aphid hosts that lead to the evolution of reduced endosymbiont genomes. Unexpectedly, all *S. symbiotica* populations propagated at one temperature appeared to evolve slower growth after only a few days due to reactivation of a lytic prophage from the CWBI genome. Though these populations continued to reach saturating cell densities slower than cultures of the ancestor throughout the experiment, most bacteria in them had mutations affecting lipopolysaccharide biosynthesis and were resistant to the phage. Some evolved strains exhibited less virulence when injected into aphids, and we observed instances of gene inactivation and loss mediated by insertion elements. Our results illustrate how transposons and prophages can dominate laboratory evolution of newly cultured bacteria, particularly those that are host-associated in nature and have genomes rife with selfish DNA elements. They also suggest that bacteria-phage coevolution can catalyze evolutionary paths that contribute to converting pathogens into stably inherited endosymbionts.

IMPORTANCE

Laboratory experiments can be used to explore evolutionary innovations in how microbes associate with animal hosts. *Serratia symbiotica* bacteria exhibit a variety of interactions with aphids. Some strains are obligate endosymbionts. Others have facultative associations with benefits or costs depending on the environmental context. *S. symbiotica* CWBI-2.3^T (CWBI) resembles aphid endosymbionts in how it can be transovarially transmitted to aphid embryos. However, adults injected with CWBI do not survive long enough to give birth to infected offspring. We evolved this aphid protosymbiont in rich media to see if this would attenuate its virulence and recapitulate genome reduction observed in endosymbionts. We observed large deletions and gene inactivation, but reactivation of a prophage from the CWBI genome and then evolution of phage resistance dominated. Some evolved strains became less virulent to aphids, suggesting that evolution driven by selfish DNA elements can contribute to the emergence of new endosymbionts from pathogen ancestors.

INTRODUCTION

Microbes have evolved a wide variety of associations with insects, including living as endosymbionts within the cells and tissues of their hosts (1–3). Insect symbionts can be broadly classified as obligate or facultative in terms of their relationships with their hosts. Obligate symbionts are required for host survival. They typically colonize specialized host cells or organs, are reliably vertically transmitted to offspring, and evolve severely reduced genomes that make it difficult or impossible to axenically culture them. Facultative symbionts are not required for host survival and can be vertically or horizontally transmitted. They often display complex relationships with their hosts. For example, a facultative symbiont may impose a general fitness cost on its host, but provide a benefit, such as protection against parasitoids, that can outweigh this cost under certain conditions (4). Facultative symbionts often have larger genomes than obligate symbionts, and some can be cultured outside of their insect hosts.

Pea aphids (*Acyrthosiphon pisum*) have both obligate and facultative endosymbionts. The obligate symbiont (*Buchnera aphidicola*) is required for aphid survival because it synthesizes essential amino acids that are not present in sufficient quantities in their diet (5, 6). *Buchnera* has a greatly reduced genome and has not been cultured. Facultative symbionts of pea aphids—including *Hamiltonella defensa, Regiella insecticola*, and *Serratia symbiotica*—colonize bacteriocytes and can also be found in the hemolymph and various tissues (6). Different strains of the bacterium *Serratia symbiotica* exhibit a range of relationships with aphids. Some *S. symbiotica*, like those found in the aphid *Cinara cedri*, are co-obligate symbionts with reduced genomes and characteristics similar to *Buchnera* (7–10). Others, like *S. symbiotica* Tucson in *A. pisum*, are facultative, providing a series of context dependent benefits, such as protection against heat stress and parasitoid wasps (4, 11, 12).

Recently, the first culturable strains of *S. symbiotica* have been isolated from aphids. These culturable *S. symbiotica*, which include strain CWBI-2.3T (CWBI), primarily colonize the

aphid gut and may be horizontally transmitted between aphids feeding on the same plant (7, 13–16). They have genomes that are intermediate in size between those of other culturable *Serratia* species, such as *Serratia marcescens*, and uncultured facultative symbionts, like *S. symbiotica* Tucson (10, 17, 18). When injected into aphids, CWBI is transovarially transmitted to developing aphid embryos in the same manner as *Buchnera* and *S. symbiotica* Tucson (10). However, CWBI is unable to establish stable vertical transmission because infected adults to not survive long enough to give birth to infected offspring (10). Because of its latent capacity for transmission, CWBI and similar culturable strains like *S. symbiotica* HB1, may represent aphid protosymbionts that are poised to evolve into vertically inherited endosymbionts (10, 18, 19).

It has been hypothesized that endosymbionts arise from pathogenic ancestors and that evolving reduced virulence is necessary for transitioning to this new lifestyle (20–22). If we could evolve or engineer CWBI to be less virulent, this might complete its transformation into a vertically inherited aphid endosymbiont. Passaging microbes *in vitro* is a classical method for attenuating pathogens. Key virulence factors are often genetically unstable and rapidly lost in an environment in which they provide no benefits. Laboratory passaging has been used to develop live attenuated bacterial vaccines (23–26), and unintentional losses of virulence have occurred in pathogens when they were cultured in the laboratory (27–30). Mutations that reduce virulence may evolve due to random genetic drift if populations are subjected to single-cell bottlenecks as they are passaged, but this process is slow because bacterial mutation rates are low. Alternatively, serially passaging large populations of microbes can rapidly select for mutants with increased growth rates *in vitro* and possibly, thereby, reduce virulence *in vivo*. Both types of experiments can lead to chromosomal deletions and gene loss, potentially mimicking mechanisms of genome evolution experienced by transitional symbionts in nature.

We performed serial transfer evolution experiments with CWBI to examine what mutations accumulated in its genome during adaptation to laboratory culture conditions. Reactivation of a prophage occurred in all populations evolved at one temperature followed by the evolution of

resistance to this phage through mutations in genes involved in lipopolysaccharide biosynthesis. Insertion sequence activity led to gene inactivation and loss in evolved strains, but not at as high of a rate as might be expected from their prevalence in the CWBI genome. Aphids survived for longer after they were injected with some evolved strains compared to the ancestor. This attenuated virulence, if compounded by further evolution, might eventually lead to stable vertical transmission. Our results illustrate how mobile DNA elements, such as transposons and prophages, which may proliferate in the genomes of host-associated bacteria (17, 31, 32), shape how they evolve. They further suggest that coevolution with these elements may be important for the emergence of new endosymbionts from pathogen ancestors.

RESULTS

Serratia symbiotica laboratory evolution experiment

CWBI-GFP is a genetically modified variant of *Serratia symbiotica* CWBI-2.3^T with a cassette expressing GFP integrated into its chromosome (10). We evolved CWBI-GFP by serially propagating twenty-four populations through cycles of 1:1000 dilution and regrowth in a rich medium (tryptic soy broth) (**Fig. 1A**). Twelve of the populations (LT-01 through LT-12) were cultured at a lower temperature, the ambient temperature of 22–25°C in our laboratory. The other twelve (HT-01 through HT-12) were cultured at a higher temperature, in an incubator set to 30°C. The CWBI-GFP ancestor grew to saturation within 24 hours at both temperatures, and all populations were initially transferred daily. Two of the 30°C populations (HT-05 and HT-12) were eventually discontinued due to suspected contamination.

Delays in population growth always evolved in the LT treatment

For every LT population, we observed a pronounced delay in how long it took the cultures to reach normal saturating turbidity levels beginning between 4 and 10 passages (**Fig. 1B**). These delayed cultures were considerably less turbid than the CWBI-GFP ancestor after 24 hours and

sometimes contained stringy debris resembling cobwebs. All eventually reached an apparently saturating turbidity after additional days of incubation, at which point they were transferred. After further variability in the delays observed on subsequent days, all LT populations began to consistently reach full turbidity between 24 and 48 hours of incubation. We propagated all LT populations for a total of 15 transfers, which took from 21 to 33 days.

Most HT populations continued to reach saturation within 24 hours for the duration of the evolution experiment and were propagated through a total of 30 transfers (**Fig. 1C**). The one exception was population HT-11, which exhibited a growth delay on day 19, but then returned to reaching saturation each day. On day 20 and thereafter, cells in this population rapidly settled to the bottom of the test tube when it was removed from the shaking incubator. To further investigate this new phenotype, we randomly selected ten colonies from the HT-11 population for characterization at the end of the experiment. Saturated cultures grown from all ten of these clonal isolates exhibited the same rapid settling as cultures of the full population.

Fig. 1. *Serratia symbiotica* **CWBI-3.2^T evolution experiment.** (**A**) Procedures used to begin the

independent populations of the evolution experiment, carry out cycles of serial transfer and regrowth, and

isolate endpoint clones for characterization. (**B**) Transfer histories of lower temperature (LT) populations

evolved at ambient laboratory temperature (22–25°C). (**C**) Transfer histories of higher temperature (HT)

populations evolved in an incubator at 30°C. In panels B and C, darker shading indicates that a

population was transferred to fresh medium with a 1:1000 dilution on that day. Lighter shading is used for

days when a population was not transferred because it had not yet reached the normal turbidity expected

for a saturated culture of the ancestor strain.

Growth rates of evolved LT populations and clonal isolates are poorly correlated

To characterize CWBI evolution, we picked a single clonal isolate from each endpoint population

(**Fig. 1A**). We recorded growth curves for all populations and these clonal isolates in microwell

plates using a plate reader (**Fig. 2**). Nine of the LT populations evolved at 22–25°C had

significantly slower exponential growth rates than the ancestor at 25°C (**Fig. 2A**). The evolved

LT clones exhibited more variability in their growth rates compared to their populations of origin

(**Fig. 2B**). Some grew more rapidly, while others grew much more slowly. The net effect was

that, on average, population growth rates were not significantly different from clone growth rates

(*F*-test, *p* = 0.283), even though the growth rates of individual clones relatively poorly correlated

with the growth rates measured for their populations of origin (r_{xy} = 0.465, sample Pearson's

correlation coefficient).

Fig. 2. Growth rates of evolved populations and strains. Exponential growth rates estimated from

microplate assays for (A) endpoint LT populations at ambient room temperature, (B) clonal isolates from

the LT populations at ambient room temperature, (C) endpoint HT populations at 30°C, and (D) clonal

isolates from the HT populations at 30°C. Horizontal dashed lines are reference lines at the growth rate of

the CWBI-GFP ancestor strain (Anc). Error bars are standard deviations. Asterisks above or below points

indicate that there is a significant difference between the measured growth rate of a population or clonal isolate and the ancestor in the same microplate assay (two-tailed t-tests with Bonferroni correction, *$p <$ 0.05 and **$p < 0.01$).

There were fewer changes in the evolved samples and less variability between them in the 30°C treatment. A majority of the evolved HT population had growth rates that were indistinguishable from that of the ancestor strain, with just four exhibiting significantly reduced growth rates (**Fig. 2C**). Evolved HT clone growth rates (**Fig. 2D**) also correlated better with their populations of origin (r_{xy} = 0.766) than was the case for the LT samples. The main outlier was population HT-11 and its clonal isolate, which both appeared to grow more slowly, likely due to the rapid settling phenotype interfering with measuring growth from changes in culture turbidity. Though there was no difference, on average, in the growth rates of the HT clones and populations (F-test, p = 0.605), the HT-03 and HT-10 clones individually exhibited significantly increased growth rates compared to the ancestor (**Fig. 2D**).

Phages derived from a CWBI lysogen are present in evolved LT populations

The final LT populations took more than 24 h to reach saturation and accumulated stringy debris. We hypothesized that this debris was derived from cells lysed by phages, and that partial lysis of the cell populations was causing the apparent growth delays. When we spotted phage lysates that we made from LT endpoint populations on top agar containing the CWBI-GFP ancestor, we observed turbid zones of clearing after 48 h of incubation that supported this hypothesis (**Fig. 3A**). Transmission electron microscopy (TEM) imaging of lysates showed that they contained tailed phages (**Fig. 3B**). These phages have a capsid with a diameter of ~70 nm and a tail with a length of ~150 nm. The diameter of the tail is ~10 nm except for a ~50 nm segment near the end where it widens to ~20 nm.

Fig. 3. Activation of *S. symbiotica* CBWI prophages. (**A**) Turbid plaques formed by lysate containing

the Laboratory-Activated Serratia Phage (LASφ) on the *S. symbiotica* CWBI-GFP ancestor. The lysate

was made from a revived sample of the LT-02 endpoint population. Ten-fold serial dilutions of lysate,

beginning on the left, were spotted on top agar containing host cells. This plate was incubated at ambient

room temperature for 48 hours before imaging. (B) Representative TEM images of phage particles in

lysate from the LT-02 endpoint population. Scale bars are 50 nm. (C) Coverage of DNA sequencing reads

mapped to the CWBI-GFP genome from a pooled lysate of all 12 endpoint LT populations which

contained LASφ phage. (D) Coverage of DNA sequencing reads mapped to the CWBI-GFP genome from

a sample of the supernatant of ancestor cells treated with mitomycin C (MC).

To identify the reactivated phage and ascertain its origin, we pooled lysates made from

all LT populations and extracted and sequenced their DNA. The CWBI-GFP genome contains

21 putative prophage regions, 10 of which are predicted to encode intact prophages (**Table 1**).

Sequencing reads from the pooled phage lysates mapped to coordinates that matched a

prediction of a 36.7-kb prophage at coordinate 1,323,010 (**Fig. 3C**). This genomic region is

predicted to encode a complete copy of a P2-like prophage in the family Peduoviridae. We

named this phage, which recurrently became lytic during our evolution experiments at the lower

temperature, the Laboratory-Activated Serratia Phage (LASφ).

Table 1. Prophage regions in the *S. symbiotica* ancestor chromosome predicted by PHASTER

Designation	Position	Size (kb)	Proteins	Completeness	Score	Annotation
	33,885	36.6	49	intact	100	Entero_phiT5282H_NC_049429
	224,112	32.3	28	intact	100	Vibrio_CTX_NC_015209
	269,576	4.5	11	questionable	80	Vibrio_Thalassa_NC_042095
	441,049	35.8	51	intact	150	Haemop_SuMu_NC_019455
	545,061	9.7	14	incomplete	60	Bacill_G_NC_023719
	656,411	7.6	13	questionable	70	Mycoba_Tonenili_NC_031080
	706,563	6.6	11	incomplete	50	Shigel_Sf6_NC_005344
	705,025	29.6	25	intact	120	Shigel_Sf6_NC_005344
MC	771,733	48	74	intact	150	Salmon_118970_sal3_NC_031940
	914,879	35.7	54	intact	150	Haemop_SuMu_NC_019455
	1,295,865	19.8	21	incomplete	40	Mannhe_vB_MhM_1127AP1_NC_047750
LASφ	1,323,010	36.7	52	intact	120	Escher_500465_1_NC_049342
	1,722,414	8	9	incomplete	30	Bacill_SP_15_NC_031245
	1,885,653	17.4	27	intact	100	Cronob_ENT39118_NC_019934
	1,921,934	89.6	102	intact	150	Salmon_118970_sal3_NC_031940
	2,111,306	44.2	38	intact	103	Entero_phiT5282H_NC_049429
	2,236,529	16.8	32	questionable	90	Burkho_KS5_NC_015265
	2,267,340	20.8	12	incomplete	50	Escher_500465_1_NC_049342
	2,465,690	17.8	22	questionable	70	Shigel_SfII_NC_021857
	2,632,536	33.6	11	incomplete	40	Shigel_Sf6_NC_005344
	3,329,468	5.2	8	incomplete	40	Escher_SH2026Stx1_NC_049919

LASφ is the prophage region with nearly all DNA sequencing coverage in a pooled phage lysate from

evolved samples. MC is the prophage region with DNA sequencing coverage in the supernatant of

cultures of the ancestor treated with mitomycin C.

Next, we next tested whether LASφ replication was activated by DNA damage by sequencing DNA from the supernatant of a CWBI-GFP culture exposed to mitomycin C. DNA in this sample was enriched in sequences that mapped to the approximate coordinates 772,000–819,000 of the CWBI-GFP genome (**Fig. 3D**). This region of the CWBI genome encodes genes associated with a different prophage that is also predicted to be complete (**Table 1**). Therefore, we conclude that LASφ is not activated by the SOS response and that the CWBI genome encodes at least this one other intact prophage that is capable of entering the lytic cycle.

Evolved LT clones are resistant to LASφ

Given the cell death and growth delays we observed in the LT populations, we expected there to be strong selection for the evolution of resistance to LASφ. To test whether this was the case, made phage lysate from the 15-passage sample of population LT-02. We tested the effect of adding this lysate or the same lysate after it was heated to 90°C, on the growth of cultures of the ancestor and the LT clones (**Fig. 4**). The ancestor showed a normal growth curve when exposed to the heat-inactivated lysate but did not develop turbidity when subjected to phage lysate. By contrast, there was no difference in the growth curves of most evolved clones whether they were exposed to active versus heated lysate. The exception was the LT-07 clone, which was also inhibited by the active phage. In conclusion, all but one of the evolved LT clones evolved resistance to LASφ.

Fig. 4. Evolved clones are resistant to LASφ. Treatment with phage lysate from a sample of the endpoint LT-02 population inhibits growth of the CWBI-GFP ancestor and the LT-07 clone but not any of the other characterized LT clones. Inactivating the phage lysate by heating abolishes this inhibition.

Genome evolution

Next, we sequenced the genomes of all LT and HT endpoint clones to understand how they adapted to the laboratory environment and determine the genetic basis of resistance to LASφ. There were 1.1 and 0.8 mutations on average in each LT and HT clone, respectively (**Table 2**). Most genomes clones had exactly one mutation, except two had two mutations each (LT-05 and HT-11) and three—all in the high-temperature treatment—had no mutations (HT-01, HT-02, and HT-08). Base substitutions were the most abundant type of mutation, with 12 total across both temperature conditions, accounting for 57% of the 21 total mutations. All base substitutions were nonsynonymous mutations in protein-coding genes. There were also four small insertions and deletions of ≤ 50-bp, two of which were in homopolymer repeats. Lastly, we observed three IS element insertions, a 54-bp deletion, and a 27.4-kb deletion.

Table 2. Mutations in evolved clonal isolates

Clone	Episome	Position	Mutation	Annotation	Gene	Description
LT-01	Chromosome	1,046,686	C→T	S209F (TCT→TTT)	mshA_2	D-inositol-3-phosphate glycosyltransferase
LT-02	Chromosome	1,047,886	G→T	G229C (GGC→TGC)	mshA_3	D-inositol-3-phosphate glycosyltransferase
LT-03	Chromosome	1,047,886	G→C	G229R (GGC→CGC)	mshA_3	D-inositol-3-phosphate glycosyltransferase
LT-04	Chromosome	1,040,286	IS3_176 (+) +3 bp	coding (497-499/1206 nt)	01118	FkbM family methyltransferase
LT-05	Chromosome	782,372	Δ54 bp	coding (803-805/1413 nt)	ydhT_1	Putative protein YdhT
	Chromosome	2,985,053	C→T	R137H (CGT→CAT)	dam_4	DNA adenine methylase
LT-06	Plasmid 1	111,666	(A)$_{7\rightarrow8}$	intergenic (+331/+60)	03511 / rep_2	IS481 family transposase ISErsp1 / ATP-dependent DNA helicase Rep
LT-07	Chromosome	1,314,304	+16 bp	coding (189/195 nt)	01374	IS481 family transposase ISFw3
LT-08	Chromosome	1,040,768	IS3_176 (+) +3 bp	coding (979-981/1206 nt)	01118	FkbM family methyltransferase
LT-09	Chromosome	1,037,303	G→A	G302S (GGT→AGT)	algC	Phosphomannomutase/phosphoglucomutase
LT-10	Chromosome	77,271	Δ12 bp	coding (398-409/1098 nt)	pglJ	N-acetylgalactosamine-N, N'-diacetylbacillosaminyl-diphospho-undecaprenol 4-α-N-acetylgalactosaminyltransferase
LT-11	Chromosome	1,035,798	T→C	S276P (TCT→CCT)	rfbM	Mannose-1-phosphate guanylyltransferase RfbM
LT-12	Chromosome	1,654,473	C→T	R124C (CGC→TGC)	manA	Mannose-6-phosphate isomerase
HT-03	Chromosome	2,338,798	G→T	Q455H (CAG→CAT)	glnD	Bifunctional uridylyltransferase/uridylyl-removing enzyme
HT-04	Chromosome	2,727,639	Δ27,462 bp	IS3_176-mediated	cra–csbX	21 genes
HT-06	Chromosome	2,728,332	IS3_176 (+) +3 bp	intergenic (−11/+1203)	cra / leuO	Catabolite repressor/activator / HTH-type transcriptional regulator LeuO
HT-07	Chromosome	3,170,649	G→A	R265H (CGC→CAC)	glnL	Sensory histidine kinase/phosphatase NtrB
HT-09	Chromosome	3,170,202	G→A	R116H (CGC→CAC)	glnL	Sensory histidine kinase/phosphatase NtrB
HT-10	Chromosome	1,743,150	(G)$_{10\rightarrow9}$	intergenic (+108/+172)	ansA / 01771	L-asparaginase 1/hypothetical protein
HT-11	Chromosome	1,044,558	C→A	T144K (ACG→AAG)	gmd	GDP-mannose 4,6-dehydratase
	Chromosome	3,002,184	G→A	R234C (CGC→TGC)	envZ	Osmolarity sensor protein EnvZ

The mutation column shows the base change or summarizes more complex mutations. For IS element insertions the name of the element is followed by its orientation and the number of bases duplicated at the target site. The annotation column gives the amino acid and codon changes for base substitutions in protein-coding genes and the coordinates relative to genes that they are within or between for other types of mutations in genes or intergenic regions. Five-digit numbers are locus tags for genes that are not annotated with a gene name in the reference genome. For intergenic mutations, information about the two neighboring genes is shown separated by a slash with the annotation column displaying the distance and orientation of the mutation relative to these genes (+, upstream; –downstream).

Mutations affecting LPS biosynthesis are associated with phage resistance

Most mutations in the LT evolved clones appear to be associated with synthesis or modification of lipopolysaccharide (LPS) or other extracellular polysaccharides (**Table 2**). LT-09, LT-11, and LT-12 had mutations in genes that are involved in the biosynthesis of GDP-mannose, a key LPS precursor. LT-01, LT-02, LT-03, and LT-10 had mutations in glycosyltransferases that could link monosaccharide units or attach them to lipids. LT-04 and LT-08 had mutations in the same FkbM family methyltransferase that could modify sugars displayed on the cell surface. As these

were often the sole mutations in these clones and they are broadly convergent in terms of their possible functions, it seems likely that these mutations confer resistance to LASφ by altering polysaccharides on the bacterial cell surface. Clone LT-07 was the only endpoint clone at RT that was not resistant to LASφ (**Fig. 4**), and it did not have mutations in any of these pathways.

Clone HT-11, which exhibited the settling phenotype observed in this population, had two mutations. One was in a mannose biosynthesis gene *gmd*, and one was in the osmolarity sensor *envZ*. It is possible that stochastic reactivation and the evolution of resistance to LASφ also caused the evolution of the unique phenotype displayed by this population, though the way in which it altered its cell surface. None of the other evolved HT clones (HT-03) had mutations in genes with a direct connection to polysaccharide biosynthesis or modification. Instead, nearly all mutations in HT clones are in genes involved in regulation of nitrogen metabolism or central carbon metabolism, including convergent mutations affecting *glnL* and *cra*.

IS*3* elements mediated gene loss in CWBI evolved at both temperatures

The CWBI genome is predicted to have 232 insertion sequence (IS) elements from fifteen families, though 124 of these (53.8%) appear to be incomplete or have inactivating mutations. The most frequent families are IS*3* (100 total, 49 complete), IS*256* (37 total, 12 complete), and IS*4* (23 total, 21 complete). In endpoint clones from the evolution experiment, we only observed activity of one specific IS*3* subfamily with 48 copies of which 40 appear to be active. We observed a large deletion that led to the loss of 21 genes in HT-04 mediated by insertion of a new copy of this IS element in the *csrA* gene followed by recombination with a preexisting copy overlapping the end of the *csbX* gene. The same IS*3* element inserted into genes in LT-04 and LT-08 and into an intergenic region in HT-06. No other IS element families mediated deletions or were observed to insert new copies during the evolution experiment.

Evolved LT clones are less virulent than the ancestor when injected into aphids

To determine if mutations that accumulated during the evolution experiment attenuated CWBI virulence, we injected cohorts of 4th instar aphids with each of the evolved LT clones and compared their survival to aphids injected with the ancestor strain. Under these conditions aphids rapidly become systemically infected with the *S. symbiotica* CBWI-GFP ancestor (**Fig. 5A**). A majority of aphids die 3-4 days after injection with the CWBI-GFP ancestor (**Fig. 5B**) whereas a majority survive for approximately 12 days after mock injections with buffer (**Fig. 5C**). We observed significantly longer aphid survival after injecting 5 of the 12 evolved LT clones compared to paired injections with the ancestor strain ($p < 0.05$, log-rank tests) (**Fig. 5B**). The most dramatic changes were for LT-10 and LT-05. These clones killed aphids an average of 2.19 and 1.48 days more slowly than the ancestor, respectively. One clone, LT-07, appeared to kill aphids more quickly ($p = 0.028$), though the difference was only 0.55 days on average.

Fig. 5. Some LT-evolved clones exhibit slower killing of aphids. (**A**) The CWBI-GFP ancestor rapidly and systemically infects when ~10^4 bacterial cells are microinjected into aphids. GFP fluorescence in living aphids was imaged on a blue-light transilluminator with an orange filter at 4 days post-injection. (**B**)

Survival curves for cohorts of 40-50 fourth-instar aphids injected with ~10^4 *S. symbiotica* cells beginning at 2 days post-injection. Paired ancestor controls were tested at the same time as each of the 12 endpoint clones from the LT populations. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ for log-rank tests comparing Kaplan-Meier survival curves for each clone and its paired ancestor control. (**C**) Corresponding survival curves for a cohort of aphids that was mock injected with buffer that did not contain any *S. symbiotica* cells.

A faster *in vitro* growth rate for a given evolved strain might generally lead to trade-offs that made it less virulent to aphids. Alternatively, a faster growth rate in the rich media used in the evolution experiment might also track with faster *in vivo* growth within an aphid's body and therefore increase virulence. We did not find support for either of these possibilities. There was no significant correlation between the *in vitro* growth rates of the evolved clones (**Fig. 2B**) and how much longer aphids survived on average when they were injected with the evolved clones versus ancestor (the differences in the areas under the survival curves in **Fig. 5B**) ($p = 0.16$, *t*-test on slope in linear regression). There was also no pattern with respect to the presence of mutations in certain genes or pathways in the evolved clones that exhibited significantly reduced virulence versus the others. Finally, we did not observe vertical transmission of ancestor or evolved *S. symbiotica* to aphid offspring in these experiments, which is in line with the delay of 10 days post-injection before infected offspring are observed in prior experiments with *S. symbiotica* (10). Nevertheless, the substantial changes in survival for certain clones indicate that evolution in this short laboratory experiment was able to substantially attenuate the virulence of this *S. symbiotica* protosymbiont in some cases.

DISCUSSION

Our evolution experiments with culturable *S. symbiotica* strains began with two major goals related to investigating routes by which this protosymbiont strain could mutate to become a stably vertically inherited endosymbiont of aphids. First, we wanted to understand whether

adaptation to a nutrient-rich laboratory environment will attenuate the growth and virulence of CWBI when it is injected into aphids. Second, we wanted to characterize the activities of different families of mobile elements in the CWBI genome to understand their relative contributions to inactivating and deleting genes, which could also attenuate CWBI virulence. In general, one can imagine distinct routes by which mutations resulting from either non-adaptive genetic drift or adaptive decay due to trade-offs between growth in different environments transform a pathogen into an endosymbiont (17, 18, 20, 21, 31). Recapitulating this process could inform our understanding of how and when each pathway contributes to this transition.

Insertion sequence (IS) elements are simple bacterial transposons that can inactivate genes and mediate large chromosomal rearrangements (33, 34). Given the prevalence and diversity of IS elements in the *S. symbiotica* CWBI genome, we expected that IS-mediated mutations would dominate during our evolution experiment. However, only one subfamily of IS*3* elements was active. Of 21 total mutations, just three were new insertions of this transposon and one was a 27-kb deletion mediated by this transposon. The IS-mediated fraction of total mutations in our experiment with CWBI (19%) is not as high as that seen in evolution experiments with laboratory strains of *E. coli* (30%) (35) and *Acinetobacter baylyi* (41%) (36), even though these bacteria have fewer transposon copies than CWBI in their genomes. Thus, perhaps surprisingly, the genome of CWBI does not appear to be unusually unstable. Nevertheless, we would expect IS elements—both existing copies and new ones—to have an outsize role in catalyzing large-scale deletions if our evolution experiment continued.

Recently, within-host evolution experiments showed that a single mutation in the *crp* master regulator of central carbon metabolism can make *E. coli* an effective substitute for the native *Pantoea* gut symbiont of the stinkbug *Plautia stali* in supporting its nutrition and development (37). Similar global regulatory mutations affecting different growth switches, particularly stationary phase to exponential phase transitions and biofilm versus planktonic lifestyles, could greatly alter CWBI interactions with aphid hosts. In this vein, it is interesting that

we observed a mutation in the upstream region of *cra*, another master regulator of central carbon metabolism, and an IS-mediated deletion overlapping this gene in two of the evolved clones from the higher-temperature treatment. This condition also led to the evolution of *glnD* and *glnL* mutations that are expected to affect global regulation of nitrogen metabolism. How these mutations affect *S. symbiotica* growth in aphids and host survival remains to be tested.

Emergence of the Laboratory-Activated *Serratia* Phage (LASφ) complicated the initial stages of our evolution experiment, particularly in the lower temperature treatment. Reactivation of this prophage created a strong selective pressure for CWBI to evolve resistance, which occurred through mutations in lipopolysaccharide biosynthesis genes. Mutations in bacterial pathogens that confer resistance to phage often affect similar pathways and result in trade-offs that reduce fitness *in vitro* and *in vivo* when phage is not present (38, 39). For example, *Listeria monocytogenes* that evolved mutations in teichoic acid glycosylation genes that conferred resistance to a phage was less virulent in mice and exhibited a reduced ability to invade cells (40). Similar trade-offs have been observed in phage-resistant mutants of *Vibrio cholerae* (41), *Shigella flexneri* (42), and *Staphylococcus aureus* (43), among others. Some of the clonal endpoint isolates from our experiment that were resistant to LASφ grew more slowly than the ancestor, suggesting that they exhibit fitness trade-offs for phage resistance *in vitro*.

The APSE prophage in the genome of the endosymbiont *Hamiltonella defensa* encodes a toxin that has been shown to protect aphids from parasitoid wasps, thereby contributing to persistence of this facultative symbiont in some aphid populations (44–46). Genes expressed from WO prophage regions in the genomes of *Wolbachia* mediate how this endosymbiont manipulates reproduction in other types of insects (47, 48). We found that some of our evolved *S. symbiotica* clones with LASφ-resistance mutations are less virulent when injected into aphids, which suggests additional dimensions to how bacteria-phage coevolution might shape the emergence of new endosymbionts. Culturable *S. symbiotica* that are naturally acquired by aphids from their diet sometimes escape the gut and systemically infect their bodies (8).

Prophage activation, phage predation, or the evolution of phage resistance could reduce the titer or growth rates of these bacteria, thereby increasing the chances that they are able to evolve and establish as stably vertically inherited symbionts before they kill their hosts. It is also possible that phage-resistant mutations could prevent the establishment of endosymbionts. For instance, changes in LPS or loss of cell-surface proteins that act as phage receptors in resistant mutants could affect interactions with aphid embryos that lead to endocytosis of some but not other bacteria (10, 49). Additional evolution or engineering will be needed to understand whether the mutations we observed are potential steppingstones to aphid endosymbiosis.

MATERIALS AND METHODS

Culture conditions

We used CWBI-GFP, a strain of *S. symbiotica* CWBI-2.3^T with genes for GFP and kanamycin resistance integrated into its chromosome (10). CWBI-GFP was cultured in trypticase soy broth (TSB) or on trypticase soy agar (TSA) (BD Biosciences) with 50 µg/mL kanamycin (Kan). Unless specified otherwise, cultures were grown at in an incubator at 30°C or at the ambient temperature in our laboratory (22-25°C) with orbital shaking at 200 r.p.m. over a one-inch diameter in 5 mL of TSB in 150 × 18 mm test tubes. Samples were cryopreserved by adding glycerol to a final concentration of 16% (v/v) and stored at −80°C.

Evolution experiment

Twenty-four colonies of CWBI-GFP from a TSA+Kan plate were used to inoculate separate cultures of TSB+Kan to begin the separate populations of the evolution experiment. Half were grown at ambient temperature (LT populations). Half were grown at 30°C (HT populations). Upon achieving saturated growth, as judged by visual inspection, 5 µL of each culture was transferred to a new test tube containing fresh TSB+Kan. LT and HT populations were passaged 15 and 30 times, respectively. Samples of these cultures were cryopreserved every five

passages. To isolate endpoint clones, glycerol stocks of the final populations were struck out on TSA+Kan plates. Then, single large colonies were picked from each plate into TSB+Kan, and these cultures were grown to saturation and cryopreserved.

Growth curves

Cultures of evolved clones or populations revived from cryopreserved samples and grown to saturation were diluted 1:100 into TSB+Kan. From each diluted culture, 300 μL was pipetted into a clear, flat-bottomed 96-well plate (Corning). The final layout of the plate included three replicates of a TSB+Kan blank, nine replicates of the ancestor, and three replicates of each evolved clone. Growth curves were measured in a Tecan Infinite M200 PRO plate reader. Measurements of optical density at 600 nm (OD600) were recorded every 15 minutes with 14 minutes of orbital shaking at 200 r.p.m. between each reading for 48 hours. The temperature control in the instrument was set to 23ºC for LT samples and 30ºC for HT samples, but actual temperatures varied to as high as 27ºC or 32ºC, respectively, during runs due to heating from operation of the instrument. Growth curves were performed separately for each set of populations and clones evolved at different temperatures. Statistical analysis and plotting were performed in R using the *tidyverse, ggplot2*, *statix*, and *growthcurver* packages (50).

Phage lysate preparations

For each lysate, 5-25 mL of saturated culture was mixed with an equal volume of chloroform in a centrifuge tube. After inverting vigorously for 15 seconds to mix, tubes were centrifuged for 5 minutes at 3,220 × g at ambient temperature. The supernatant was transferred to a fresh tube and 100 μL of chloroform mixed in by inverting the tube. The resulting lysate was stored at 4°C.

Phage plaque assay

CWBI-GFP was grown to an OD600 of ~0.4 in TSB+Kan at ambient temperature. Phage lysates were serially diluted 10-fold using TSB in a 96-well plate up to a maximum dilution of 10^5. TSA top agar was made by halving the concentration of agar used to make TSA base media. Top

agar was melted, then 25 mL was poured into a 100 mL flask and left to cool until warm. Kanamycin was added along with 1.5 mL of culture. The mix was swirled briefly to mix, then poured onto the top of a TSA+Kan plate. Upon cooling, 2 μL of the lysate dilutions were pipetted onto the cooled top agar. After allowing the spots to dry, the plates were incubated at room temperature. Phage plaques became visible after 24 h and were more pronounced after 48 h.

Transmission electron microscopy imaging

To isolate phage particles, 4 mL of lysate from population LT-02 was mixed with 1 mL of phage precipitation solution (20% w/v PEG 8000, 2.5 M NaCl) and mixed by inverting the tube for 15 seconds. The solution was stored at 4°C overnight then centrifuged for 30 minutes at 10,000 × g at 4°C. Supernatant was removed, and the phage pellet was resuspended and diluted in 1× phage resuspension buffer (1M NaCl, 10 mM Tris•HCl pH 7.5, 0.1 mM EDTA). The phage suspension was added to a TEM grid, washed with water, and stained with 2% uranyl acetate. Images were obtained using a FEI Technai transmission electron microscope at the University of Texas at Austin Center for Biomedical Research Support (CBRS) microscopy facility.

Mitomycin C treatment

The CWBI ancestor strain was grown in TSB+Kan at room temperature to saturation from a cryopreserved stock. In a 50 mL Erlenmeyer flask, 110 μL of saturated culture was added to 11 mL of fresh TSB+Kan. Using a Genesys 150 UV-Vis spectrophotometer (ThermoFisher Scientific), initial OD600 values were measured using 1 mL of the culture. After 3 hours at room temperature under standard shaking conditions, the culture was split by pipetting 5 mL into two test tubes. In one test tube, 5 μL of a 0.6 mM solution of mitomycin C (MilliporeSigma) dissolved in water was added to one of the tubes (0.6 μM final concentration). After 24 hours of additional growth, 1 mL of the culture treated with mitomycin C was removed and centrifuged for 10 minutes at 4,000 × g at 4ºC. From the supernatant, 857 μL was removed, added to 6 mL of DNA binding buffer from the DNA Clean & Concentrator kit (Zymo Research). DNA purified with this

kit and eluted in 25 µL of DNAse/RNAse free water. The DNA concentration was measured using an Invitrogen Qubit 4 Fluorometer (ThermoFisher Scientific).

Phage susceptibility testing

Phage lysate was prepared from 25 mL of culture as described above. Approximately 2 mL of lysate was inactivated by heating to 90°C for 10 minutes. For each growth curve replicate, 100 µL of lysate was added to 400 µL of TSB+Kan in a 1.7 mL Eppendorf tube, and then 5 µL of saturated ancestor strain was added to the tube. After mixing by inverting, 300 µL from the tube was added to a single well on a 96-well plate. Nine replicate wells of ancestor and three replicates well of each evolved clone were each tested both with lysate and with heated lysate. Nine blank wells of TSK+Kan were included. Growth curves were carried out and analyzed as described above.

Phage DNA purification

Isolation and precipitation of phage particles was performed as described for TEM imaging. To eliminate carryover of nucleic acids from host cells, 40 µL of 10× DNase I buffer (New England Biolabs) was added to 360 µL of phage sample in resuspension buffer. Then, 1 µL DNase I (2000 U/mL, New England Biolabs) and 1 µL RNase A (20 mg/mL, Invitrogen PureLink™ RNase A) were added. Following incubation at 37°C for 30 min, 10 µL of 0.5 M EDTA (pH 8.0) was added to halt digestion. To remove enzymes and purify DNA from phage particles, the digested sample was phenol extracted and ethanol precipitated. The final DNA pellet was resuspended in 10 mM Tris•Cl (pH 7.5) and stored at −20°C.

Genome sequencing

TSB+Kan cultures of the ancestor strain and evolved clones were grown to saturation at their respective temperatures. For Illumina sequencing, genomic DNA (gDNA) was isolated from 1 mL of culture using a PureLink Genomic DNA Mini Kit (Invitrogen). A Qubit 2.0 Fluorometer with the broad range assay was used to measure DNA concentrations. Up to 10 ng of gDNA was

used as input to the IDT xGen DNA Lib Prep EZ using the xGen Deceleration Module to isolate

larger fragment sizes according to the manufacturer's instructions except that all reactions took

place at 20% reaction volume. Libraries were sequenced in paired-end 150-bp mode using

unique 6-bp dual index sample barcodes. Bacterial genomes were sequenced on a HiSeq X Ten

at Psomagen (Rockville, MD). Phage genomes were sequenced on an iSeq 100.

For nanopore sequencing, high molecular weight genomic DNA was isolated using the

20/G Genomic-tips (Qiagen) or the Quick-DNA HMW MagBead Kit (Zymo research). For each

sample, 325-400 ng of this gDNA was prepared for sequencing using a Rapid Barcoding kit

(Oxford Nanopore Technologies). Samples were sequenced on a MinION Mk1C using V9.4.1

flow cells. Onboard basecalling using Guppy (V 5.0.13) was done in fast mode.

Ancestor genome assembly and annotation

CWBI-GFP nanopore reads were trimmed of adaptors using Porechop and filtered based on

quality and length using Filtlong (51). Illumina reads were trimmed using fastp (52). We used

Trycycler (53) to generate a consensus assembly from Flye (54), Miniasm+Minipolish (55),

Canu (56), NECAT (57), Raven (58), and Unicycler (59) assemblies of long read subsets. The

final assembly was iteratively polished, first using Medaka (Oxford Nanopore Technologies) with

nanopore reads and then using Polypolish (60), POLCA (61), and *breseq* (62) with Illumina

reads. Finally, we used Prokka (64) and ISEscan (65) to annotate genes and IS elements in the

final genome assembly (**Data File S1**). Putative prophages were identified with PHASTER (66).

Read mapping and mutation calling

We predicted mutations in evolved clones relative to the annotated CWBI-GFP genome using

breseq (v0.38.1) (62). This pipeline uses bowtie2 (67) for read alignment. We also mapped

reads from phage and mitomycin C samples to the reference genome and examined read

coverage using *breseq*. In these samples, some regions of the LASφ prophage and the

prophage activated by mitomycin C share short stretches of exact homology with other

prophage regions that initially resulted in ambiguous mapping of some reads. To create the final coverage graphs, we used an artificial reference sequence that avoided these artifacts by including just the first half of the genome. Coverage plots zoomed to the prophage regions were created using the bam2cov post-processing command from *breseq*.

Aphid Injection

Ancestor and evolved clones were inoculated into TSB+Kan from glycerol stocks and grown at ambient temperature. Cells were pelleted by centrifugation at 6800 × g for 3 min and then washed three times in 1× PBS with 1 min of centrifugation each time. Cell pellets were resuspended in buffer A (25 mM KCl, 10 mM $MgCl_2$, 250 mM sucrose, 35 mM Tris-HCl, pH 7.5) at an OD600 of 1.0. For injection, 40-50 fourth-instar aphids per treatment were chilled overnight at 4°C, then injected at the base of the hindleg with <0.1 μL of resuspended cells using pulled 5 μL calibrated micropipettes (Drummond) loaded into a IM-400 electric microinjector (Narishige) (MPa = 0.037, 0.2 seconds). To estimate the number of cells injected into each aphid, the microinjector was used to dispense <0.1 μL of resuspended cells into tubes containing 50 μL of buffer A, 1:10 serial dilutions were spot-plated, and CFUs were counted. The number of cells injected per aphid was approximately ~10^4 cells. Immediately post-injection, aphids were placed onto 5- to 5.5-inch plants and allowed to recover for 24 h. Counting of surviving aphids began 48 hours post-injection and continued daily. For counting, aphids were gently removed from the plants and placed under a blue-light illuminator to also monitor them for colonization. Aphids were placed back on plants immediately after counting. Statistical tests and plotting of survival curves were conducted in R using the *survminer* and *survival* packages (68).

DATA AVAILABILITY

Whole-genome sequencing reads have been deposited in NCBI Sequence Read Archive (PRJNA1111250).

COMPETING INTERESTS

The authors declare no competing interests.

ACKNOWLEDGMENTS

We thank Daniel Deatherage for assistance with DNA sequencing; Victor Li for assistance with phage experimental techniques; and Michelle Mikesh of the UT Microscopy Core for assistance with TEM sample preparation and operation.

FUNDING

This research was supported by the U.S. Army Research Office (W911NF-20-1-0195 to J.E.B). A.J.V. acknowledges support from a UT Austin Graduate School summer fellowship. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

REFERENCES

1. Oliver KM, Degnan PH, Burke GR, Moran NA. 2010. Facultative symbionts in aphids and the horizontal transfer of ecologically important traits. Annu Rev Entomol 55:247–266.
2. Perlmutter JI, Bordenstein SR. 2020. Microorganisms in the reproductive tissues of arthropods. Nat Rev Microbiol 18:97–111.
3. Elston KM, Leonard SP, Geng P, Bialik SB, Robinson E, Barrick JE. 2022. Engineering insects from the endosymbiont out. Trends Microbiol 30:79–96.
4. Oliver KM, Russell JA, Moran NA, Hunter MS. 2003. Facultative bacterial symbionts in aphids confer resistance to parasitic wasps. Proc Natl Acad Sci U S A 100:1803–1807.
5. Douglas AE, Prosser WA. 1992. Synthesis of the essential amino acid tryptophan in the pea aphid (*Acyrthosiphon pisum*) symbiosis. J Insect Physiol 38:565–568.

6. Douglas AE. 1998. Nutritional interactions in insect-microbial symbioses: aphids and their symbiotic bacteria *Buchnera*. Annu Rev Entomol 43:17–37.

7. Sabri A, Leroy P, Haubruge E, Hance T, Frère I, Destain J, Thonart P. 2011. Isolation, pure culture and characterization of *Serratia symbiotica* sp. nov., the R-type of secondary endosymbiont of the black bean aphid *Aphis fabae*. Int J Syst Evol Microbiol 61:2081–2088.

8. Elston KM, Perreau J, Maeda GP, Moran NA, Barrick JE. 2021. Engineering a culturable *Serratia symbiotica* strain for aphid paratransgenesis. Appl Environ Microbiol 87:e02245-20.

9. Manzano-Marín A, Lamelas A, Moya A, Latorre A. 2012. Comparative genomics of *Serratia* spp.: two paths towards endosymbiotic life. PloS One 7:e47274.

10. Perreau J, Patel DJ, Anderson H, Maeda GP, Elston KM, Moran NA, Barrick JE. 2021. Vertical transmission at the pathogen-symbiont interface: *Serratia symbiotica* and aphids. mBio 12:e00359-21.

11. Burke G, Fiehn O, Moran N. 2010. Effects of facultative symbionts and heat stress on the metabolome of pea aphids. ISME J 4:242–252.

12. Burke GR, Moran NA. 2011. Massive genomic decay in *Serratia symbiotica*, a recently evolved symbiont of aphids. Genome Biol Evol 3:195–208.

13. Pons I, Renoz F, Noël C, Hance T. 2019. Circulation of the cultivable symbiont *Serratia symbiotica* in aphids is mediated by plants. Front Microbiol 10:764–764.

14. Pons I, Renoz F, Noël C, Hance T. 2019. New insights into the nature of symbiotic associations in aphids: infection process, biological effects, and transmission mode of cultivable *Serratia symbiotica* bacteria. Appl Environ Microbiol 85:e02445-18.

15. Renoz F, Noël C, Errachid A, Foray V, Hance T. 2015. Infection dynamic of symbiotic bacteria in the pea aphid *Acyrthosiphon pisum* gut and host immune response at the early steps in the infection process. PLOS One 10:e0122099–e0122099.

16. Pons I, Scieur N, Dhondt L, Renard M-E, Renoz F, Hance T. 2022. Pervasiveness of the symbiont *Serratia symbiotica* in the aphid natural environment: distribution, diversity and evolution at a multitrophic level. FEMS Microbiol Ecol 98:fiac012.

17. Manzano-Marín A, Latorre A. 2016. Snapshots of a shrinking partner: genome reduction in *Serratia symbiotica*. Sci Rep 6:32590–32590.

18. Renoz F, Foray V, Ambroise J, Baa-Puyoulet P, Bearzatto B, Mendez GL, Grigorescu AS, Mahillon J, Mardulyn P, Gala J-L, Calevro F, Hance T. 2021. At the gate of mutualism: identification of genomic traits predisposing to Insect-bacterial symbiosis in pathogenic strains of the aphid symbiont *Serratia symbiotica*. Front Cell Infect Microbiol 11:660007.

19. Renoz F, Pons I, Vanderpoorten A, Bataille G, Noël C, Foray V, Pierson V, Hance T. 2019. Evidence for gut-associated *Serratia symbiotica* in wild aphids and ants provides new perspectives on the evolution of bacterial mutualism in insects. Microb Ecol 78:159–169.

20. Husník F, Chrudimský T, Hypša V. 2011. Multiple origins of endosymbiosis within the Enterobacteriaceae (γ-Proteobacteria): convergence of complex phylogenetic approaches. BMC Biol 9:87.

21. McCutcheon JP, Boyd BM, Dale C. 2019. The life of an insect endosymbiont from the cradle to the grave. Curr Biol 29:R485–R495.

22. Clayton AL, Oakeson KF, Gutin M, Pontes A, Dunn DM, von Niederhausern AC, Weiss RB, Fisher M, Dale C. 2012. A novel human-infection-derived bacterium provides insights into the evolutionary origins of mutualistic insect-bacterial symbioses. PLoS Genet 8:e1002990.

23. Oettinger T, Jørgensen M, Ladefoged A, Hasløv K, Andersen P. 1999. Development of the *Mycobacterium bovis* BCG vaccine: review of the historical and biochemical evidence for a genealogical tree. Tuber Lung Dis 79:243–250.

24. Luca S, Mihaescu T. 2013. History of BCG Vaccine. Maedica 8:53–58.

25. Hanley KA. 2011. The double-edged sword: how evolution can make or break a live-attenuated virus vaccine. Evol Educ Outreach 4:635–643.

26. Kenner J, Cameron F, Empig C, Jobes DV, Gurwith M. 2006. LC16m8: an attenuated smallpox vaccine. Vaccine 24:7009–7022.

27. Somerville GA, Beres SB, Fitzgerald JR, DeLeo FR, Cole RL, Hoff JS, Musser JM. 2002. In vitro serial passage of *Staphylococcus aureus*: changes in physiology, virulence factor production, and agr nucleotide sequence. J Bacteriol 184:1430–1437.

28. Sher AA, Jerome JP, Bell JA, Yu J, Kim HY, Barrick JE, Mansfield LS. 2020. Experimental evolution of *Campylobacter jejuni* leads to loss of motility, *rpoN* (σ54) deletion and genome reduction. Front Microbiol 11:579989.

29. Evans DG, Silver RP, Evans DJ, Chase DG, Gorbach SL. 1975. Plasmid-controlled colonization factor associated with virulence in *Esherichia coli* enterotoxigenic for humans. Infect Immun 12:656–667.

30. Duangurai T, Reamtong O, Rungruengkitkun A, Srinon V, Boonyuen U, Limmathurotsakul D, Chantratita N, Pumirat P. 2020. *In vitro* passage alters virulence, immune activation and proteomic profiles of *Burkholderia pseudomallei*. Sci Rep 10:8320.

31. Song H, Hwang J, Yi H, Ulrich RL, Yu Y, Nierman WC, Kim HS. 2010. The early stage of bacterial genome-reductive evolution in the host. PLoS Pathog 6:e1000922.

32. Haudiquet M, de Sousa JM, Touchon M, Rocha EPC. 2022. Selfish, promiscuous and sometimes useful: how mobile genetic elements drive horizontal gene transfer in microbial populations. Philos Trans R Soc Lond B Biol Sci 377:20210234.

33. Raeside C, Gaffé J, Deatherage DE, Tenaillon O, Briska AM, Ptashkin RN, Cruveiller S, Médigue C, Lenski RE, Barrick JE, Schneider D. 2014. Large chromosomal rearrangements during a long-term evolution experiment with *Escherichia coli*. mBio 5:e01377-14.

34. Lee H, Doak TG, Popodi E, Foster PL, Tang H. 2016. Insertion sequence-caused large-scale rearrangements in the genome of *Escherichia coli*. Nucleic Acids Res 44:7109–7119.

35. Consuegra J, Gaffé J, Lenski RE, Hindré T, Barrick JE, Tenaillon O, Schneider D. 2021. Insertion-sequence-mediated mutations both promote and constrain evolvability during a long-term experiment with bacteria. Nat Commun 12:980–980.

36. Renda BA, Dasgupta A, Leon D, Barrick JE. 2015. Genome instability mediates the loss of key traits by *Acinetobacter baylyi* ADP1 during laboratory evolution. J Bacteriol 197:872–881.

37. Koga R, Moriyama M, Onodera-Tanifuji N, Ishii Y, Takai H, Mizutani M, Oguchi K, Okura R, Suzuki S, Gotoh Y, Hayashi T, Seki M, Suzuki Y, Nishide Y, Hosokawa T, Wakamoto Y, Furusawa C, Fukatsu T. 2022. Single mutation makes *Escherichia coli* an insect mutualist. Nat Microbiol 7:1141–1150.

38. Hasan M, Ahn J. 2022. Evolutionary dynamics between phages and bacteria as a possible approach for designing effective phage therapies against antibiotic-resistant bacteria. Antibiot Basel Switz 11:915.

39. Mangalea MR, Duerkop BA. 2020. Fitness Trade-offs resulting from bacteriophage resistance potentiate synergistic antibacterial strategies. Infect Immun 88:e00926-19.

40. Wendlinger G, Loessner MJ, Scherer S. 1996. Bacteriophage receptors on *Listeria monocytogenes* cells are the *N*-acetylglucosamine and rhamnose substituents of teichoic acids or the peptidoglycan itself. Microbiol Read Engl 142 (Pt 4):985–992.

41. Yen M, Cairns LS, Camilli A. 2017. A cocktail of three virulent bacteriophages prevents *Vibrio cholerae* infection in animal models. Nat Commun 8:14187.

42. Kortright KE, Done RE, Chan BK, Souza V, Turner PE. 2022. Selection for phage resistance reduces virulence of *Shigella flexneri*. Appl Environ Microbiol 88:e0151421.

43. Capparelli R, Nocerino N, Lanzetta R, Silipo A, Amoresano A, Giangrande C, Becker K, Blaiotta G, Evidente A, Cimmino A, Iannaccone M, Parlato M, Medaglia C, Roperto S, Roperto F, Ramunno L, Iannelli D. 2010. Bacteriophage-resistant *Staphylococcus aureus*

mutant confers broad immunity against staphylococcal infection in mice. PloS One 5:e11720.

44. Degnan PH, Moran NA. 2008. Diverse phage-encoded toxins in a protective insect endosymbiont. Appl Environ Microbiol 74:6782–6791.

45. Oliver KM, Degnan PH, Hunter MS, Moran NA. 2009. Bacteriophages encode factors required for protection in a symbiotic mutualism. Science 325:992–994.

46. Weldon SR, Strand MR, Oliver KM. 2013. Phage loss and the breakdown of a defensive symbiosis in aphids. Proc R Soc B Biol Sci 280:20122103.

47. LePage DP, Metcalf JA, Bordenstein SR, On J, Perlmutter JI, Shropshire JD, Layton EM, Funkhouser-Jones LJ, Beckmann JF, Bordenstein SR. 2017. Prophage WO genes recapitulate and enhance *Wolbachia*-induced cytoplasmic incompatibility. Nature 543:243–247.

48. Bordenstein SR, Bordenstein SR. 2022. Widespread phages of endosymbionts: Phage WO genomics and the proposed taxonomic classification of Symbioviridae. PLOS Genet 18:e1010227.

49. Koga R, Meng X-Y, Tsuchida T, Fukatsu T. 2012. Cellular mechanism for selective vertical transmission of an obligate insect symbiont at the bacteriocyte–embryo interface. Proc Natl Acad Sci U S A 109:E1230–E1237.

50. Sprouffske K, Wagner A. 2016. Growthcurver: an R package for obtaining interpretable metrics from microbial growth curves. BMC Bioinformatics 17:172.

51. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Completing bacterial genome assemblies with multiplex MinION sequencing. Microb Genomics 3:e000132.

52. Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics 34:i884–i890.

53. Wick RR, Judd LM, Cerdeira LT, Hawkey J, Méric G, Vezina B, Wyres KL, Holt KE. 2021. Trycycler: consensus long-read assemblies for bacterial genomes. Genome Biol 22:266.

54. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using repeat graphs. 5. Nat Biotechnol 37:540–546.

55. Li H. 2016. Minimap and miniasm: fast mapping and de novo assembly for noisy long sequences. Bioinformatics 32:2103–2110.

56. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res 27:722–736.

57. Chen Y, Nie F, Xie S-Q, Zheng Y-F, Dai Q, Bray T, Wang Y-X, Xing J-F, Huang Z-J, Wang D-P, He L-J, Luo F, Wang J-X, Liu Y-Z, Xiao C-L. 2021. Efficient assembly of nanopore reads via highly accurate and intact error correction. Nat Commun 12:60.

58. Vaser R, Šikić M. 2021. Time- and memory-efficient genome assembly with Raven. Nat Comput Sci 1:332–336.

59. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. PLoS Comput Biol 13:e1005595.

60. Wick RR, Holt KE. 2022. Polypolish: Short-read polishing of long-read bacterial genome assemblies. PLoS Comput Biol 18:e1009802.

61. Zimin AV, Salzberg SL. 2020. The genome polishing tool POLCA makes fast and accurate corrections in genome assemblies. PLoS Comput Biol 16:e1007981.

62. Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using *breseq*. Methods Mol Biol 1151:165–188.

63. Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory evolved microbes from next-generation sequencing data using breseq. Methods Mol Biol Clifton NJ 1151:165–188.

64. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069.

65. Xie Z, Tang H. 2017. ISEScan: automated identification of insertion sequence elements in prokaryotic genomes. Bioinformatics 33:3340–3347.

66. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. Nucleic Acids Res 44:W16–W21.

67. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9:357–360.

68. Terry M. Therneau, Patricia M. Grambsch. Modeling Survival Data: Extending the Cox Model. Springer, New York.