1  **Article Title:** Sublethal pesticide exposure decreases mating and disrupts chemical signaling in a
2  beneficial pollinator
3

4  **Author Names:** Nathan Derstine, Cameron Murray, Frederick Purnell, Etya Amsalem
5
**Affiliations:** Department of Entomology, Center for Chemical Ecology, Center for Pollinator

Research, Huck Institutes of the Life Sciences, Pennsylvania State University, University Park,

PA 16802 U.S.A.

Nathan Derstine (corresponding author) – ntd34@psu.edu
Cameron Murray - csm5580@psu.edu
Frederick Purnell - frederickspurnell@gmail.com
Etya Amsalem – eua6@psu.edu
13

**Acknowledgements**

The authors would like to acknowledge the Huck Metabolomics Core Facility

(RRID:SCR_023864) for use of the Waters Acquity UPLC and Xevo TQ-S mass spectrometer and

Dr. Ashley Shay for collecting the data shown in Figure 2. We also acknowledge the Huck

Microscopy Core Facility (RRID:SCR_024457) for use of the Olympus Fluoview 10i-LIV

confocal microscope. This research was funded through a USDA-NIFA Predoctoral Fellowship

(2021-67034-35047) to ND.

**Declaration of Interests**

The authors have nothing to declare.

**Author Contributions**

N.D. designed and carried out the experiments, analyzed the data and prepared the figures, and

wrote the main manuscript text. E.A. designed experiments, wrote and revised the manuscript.

Under the supervision of N.D., C.M. and F.P collected callow bees, carried out mating

experiments, administered pesticide treatments, and took size measurements of bees (weight and

head width).

**Abstract**

Pesticides provide vital protection against insect pests and the diseases they vector but are
simultaneously implicated in the drastic worldwide decline of beneficial insect populations.
Convincing evidence suggests that even sublethal pesticide exposure has detrimental effects on
both individual- and colony-level traits, but the mechanisms mediating these effects remained
poorly understood. Here, we use bumble bees to examine how sublethal exposure to pesticides
affects mating, a key life history event shared by nearly all insects, and whether these impacts are
mediated via impaired sexual communication. In insects, mate location and copulation are
primarily regulated through chemical signals and rely on both the production and perception of
semiochemicals. We show through behavioral bioassays that mating success is reduced in bumble
bee gynes after exposure to field-relevant sublethal doses of imidacloprid, and that this effect is
likely mediated through a disruption of both the production and perception of semiochemicals.
Semiochemical production was altered in gyne and male cuticular hydrocarbons (CHCs), but not
in exocrine glands where sex pheromones are presumably produced (i.e., gyne mandibular glands
and male labial glands). Male responsiveness to gyne mandibular gland secretion was reduced, but
not the queen responsiveness to the male labial secretion. In addition, pesticide exposure reduced
queen fat body lipid stores and male sperm quality. Overall, the exposure to imidacloprid affected
the fitness and CHCs of both sexes and the antennal responses of males to gynes. Together, our
findings identify disruption of chemical signaling as the mechanism through which sublethal
pesticide exposure reduces mating success.

**Introduction**

Pesticide exposure is a major contributing factor to insect decline, including beneficial insects such as pollinators (Potts *et al.* 2010; Cameron *et al.* 2011; Goulson *et al.* 2015; Rundlöf *et al.* 2015; Brühl & Zaller 2019; Forister, Pelton & Black 2019; Sánchez-Bayo & Wyckhuys 2019). Negative impacts on pollinators are of particular concern because of their integral role in food production and wild ecosystems (Klein *et al.* 2007; Aizen *et al.* 2009; Ollerton, Winfree & Tarrant 2011; Potts *et al.* 2016; Jordan *et al.* 2021). While the effects of exposure are dependent on the dose, active ingredient, route of exposure, clearance rate (Cresswell *et al.* 2014), and species (Cresswell *et al.* 2012; Gradish *et al.* 2018; Sgolastra *et al.* 2018; Franklin & Raine 2019; Rondeau *et al.* 2022), continually mounting evidence supports the idea that even sublethal doses of pesticides negatively impact insect behavior and physiology (Blacquière *et al.* 2012; Bantz *et al.* 2018; Müller 2018; Tosi *et al.* 2022).

Globally, one of the most widely used class of pesticides are neonicotinoids (Douglas & Tooker 2015), which are applied as foliar sprays, soil drenches, and seed treatments (Jeschke *et al.* 2011) to suppress agricultural pests. After application, neonicotinoids act systemically in plants and persist in the soil, pollen and nectar, thereby creating multiple routes of exposure for pollinators (Goulson *et al.* 2015; Raine & Rundlöf 2024). Neonicotinoids disrupt central nervous system function in insects by binding to nicotinic acetylcholine receptors (Jeschke *et al.* 2011; Moffat *et al.* 2016), leading to a vast array of detrimental impacts on behavior and physiology. In bees, the negative impacts of sublethal neonicotinoid exposure include, but are not limited to: flight ability (Tosi, Burgio & Nieh 2017), foraging (Gill & Raine 2014; Colin *et al.* 2019), hygienic behavior (Tsvetkov *et al.* 2017), thermoregulation (Crall *et al.* 2018), sleep (Tasman, Rands & Hodge 2020), learning and memory (Stanley, Smith & Raine 2015; Muth, Francis & Leonard 2019; Siviter & Muth 2022), mating (Chen *et al.* 2024) and sperm viability (Chaimanee *et al.* 2016; Ciereszko *et al.* 2016; Straub *et al.* 2016). Many of these impacts likely occur simultaneously, therefore it is not surprising that sublethal neonicotinoid exposure ultimately impacts fitness by reducing reproductive output (Laycock *et al.* 2012; Whitehorn *et al.* 2012; Sandrock *et al.* 2014; Baron *et al.* 2017; Wu-Smart & Spivak 2018; Siviter *et al.* 2020; Stuligross & Williams 2020; Willis Chan & Raine 2021). However, despite the intensive efforts to describe the impacts of sublethal pesticide

exposure in insects, the mechanisms mediating these effects remained poorly understood. Here,
we examine how pesticide exposure affects mating, a key life-history bottleneck shared by the
majority of insects, and whether the impacts are mediated through disruption of sexual
communication.

Mate location and copulation in insects are primarily regulated through chemical signals called
pheromones (Ayasse, Paxton & Tengö 2001; Keeling, Plettner & Slessor 2004; Yew & Chung
2015). Pheromones are often produced in exocrine glands, secreted by either males, females or
both (Keeling, Plettner & Slessor 2004; Bruschini, Cervo & Turilazzi 2010) and elicit a response
in the recipient. In bumble bees, males mark mating sites with secretions from their labial glands
to attract gynes (newly-emerged queens) (Bergman & Bergström 1997; Ayasse & Jarau 2014).
When gynes find males, short-range attraction is likely mediated by both cuticular hydrocarbons
and pheromones from the head of gynes (Krieger *et al.* 2006), possibly from their mandibular
glands. Previous studies suggest that this type of sexual communication can be impaired by
sublethal doses of pesticide, a phenomenon known as info-disruption (Lürling & Scheffer 2007;
Tricoire-Leignel *et al.* 2012). For example, in the parasitoid *Nasonia vitripennis*, imidacloprid
impaired olfaction, which reduced the recognition of sexual signals and ultimately successful
mating (Tappert *et al.* 2017). In a study of three tortricid moth species, sublethal exposure to
thiacloprid altered female calling behavior in all three species but affected the chemical blend of
the pheromone in only one species (Navarro-Roldán & Gemeno 2017). Because females of many
insect species have low mating frequencies and are often singly mated (Ridley 1990; Schmid-
Hempel & Schmid-Hempel 2000; Ayasse, Paxton & Tengö 2001; Bird *et al.* 2024), any reduction
in their ability to find mates and successfully copulate could reduce their fitness to zero (Van
Wilgenburg, Driessen & Beukeboom 2006; Heimpel & de Boer 2008). This disruption can result
in large consequences on insect fitness.

Here, we examine the sublethal effects of the neonicotinoid imidacloprid on mating behavior and
their underlying chemical and physiological mediators. We used *Bombus impatiens*, a native,
social, and commercially available bumble bee species. *B. impatiens* is an excellent model to
address these questions because like many other pollinators (mostly solitary bees), bumble bee
gynes have a low mating-frequency (Schmid-Hempel & Schmid-Hempel 2000; Bird *et al.* 2024),

typically mating once prior to entering a winter diapause, after which they start a social colony

(Plowright & Laverty 1984). Single mating places greater importance on mate choice for gynes.

Bumble bee males employ one of three mate-finding strategies: flying a patrol route between

locations marked with their cephalic labial gland secretion, waiting at the nest entrance of a colony

for gynes to leave, and perching and pouncing on passing gynes (Valterová *et al.* 2019). Patrolling

behavior is the most common, and has been observed in many members of the *Pyrobombus*

subgenus to which *B. impatiens* belongs (Svensson & Bergstrom 1977). The labial gland scent

marks are attractive to gynes (Lecocq *et al.* 2015; Kubo, Harano & Ono 2017). Gynes are also

assumed to produce sex pheromones in glands located in their head (Krieger *et al.* 2006),

potentially the mandibular glands. These compounds presumably mediate the short-range

attraction between gynes and males. We hypothesize that sublethal pesticide exposure will have

detrimental effects on mating by either 1) interfering with chemical signal production or

perception, and/or 2) by reducing fitness such that the insects communicate their decreased quality

through honest chemical signals. To test these hypotheses, we first examine the effect of sublethal

exposure on mating behavior by exposing either gynes or males to one of two sublethal doses of

imidacloprid (6 ppb or 60 ppb). Next, we examine the impact of this exposure on signal production

and perception through chemical and electrophysiological analyses. Finally, we measure potential

fitness outcomes of imidacloprid exposure by measuring lipid storage in gynes, and sperm viability

in males. Gynes rely on lipid reserves to complete diapause (Hahn & Denlinger 2007), which are

accumulated in the pre-diapause period (Treanore & Amsalem 2020).

**Methods**

*Bee colonies and sampling.* Bees were collected from *B. impatiens* colonies obtained from Koppert

Biological Systems (Howell, Michigan, USA) and housed in walk-in environmental chambers

maintained at a constant temperature of 28–30°C, 60% relative humidity, and constant darkness.

These colonies provided gynes (unmated, pre-diapause queens) and males, which were sampled

on the day of emergence and kept separately in small plastic cages (11 cm diameter × 7 cm height).

Males and gynes were grouped by colony and age cohort, to ensure they were unmated prior to

experimentation. Colonies and cages were supplied with *ad libitum* 60% sucrose solution and

honeybee-collected pollen. With some exceptions, the head width and wet mass of the bees used

in experiments were taken as a proxy for size (Amsalem *et al.* 2009), which was used as a covariate

in statistical models where applicable.

*Pesticide treatment*

Imidacloprid was provided to gynes and males in all experiments listed below via chronic, *ad*

*libitum* feeding in sugar solution. Imidacloprid (Pestanal, Sigma-Aldrich) was first diluted in

deionized water to an initial concentration 100 ng/µL. This solution was further diluted in 60%

sucrose solution to achieve the desired concentration. Bees were collected upon emergence to

control for their age and ensure they were unmated, then were housed in small microcolony cages

of 2-4 gynes and 4-6 males per cage, and access to *ad libitum* sucrose solution. On day 3, gynes

were divided among treatment groups and fed either 0, 6 or 60 ppb imidacloprid for the next 3

days. Males were treated the same, except their 3 day exposure started on day 7 after emergence.

Gynes were then used in mating bioassays on day 6, and males on day 10 after emergence. This

feeding regime was designed to mimic chronic environmental exposure and keep the pesticide

exposure consistent between gynes and males while enabling their use in mating bioassays at

optimal ages of sexual maturity (i.e., males sexually mature later than gynes; (Treanore *et al.*

2021)). This served a second purpose of eliminating the variability associated with age. Six and

60 ppb concentrations represent sublethal doses that span a common range of field realistic

exposure encountered in pollen and nectar, with 60 ppb representing the higher end of reported

values (Schmuck *et al.* 2001; Chauzat *et al.* 2006; Krischik, Landmark & Heimpel 2007; Krupke

*et al.* 2012; Stoner & Eitzer 2012; Zioga *et al.* 2020).

*Determination of imidacloprid concentration in bees*

To assess the amount of imidacloprid in bees of different pesticide treatments, imidacloprid was

extracted and analyzed by UPLC-MS as in (Crall *et al.* 2018). Gynes (n = 29, day 6) and males (n

= 19, day 10) were homogenized in a Fastprep-bead beater with 1 mL 25% methanol, 2 metal

beads, and 20 µL of 500 nM d4-imidacloprid as an internal standard. After centrifugation at 16,000

rpm for 10 min, 900 µL of supernatant was transferred to a clean 2 mL tube containing 900 µL of

25% acetic acid. Solid phase extraction columns (SPE, DSC-18, 1mL, 50 mg, Supelco) were then

used to purify the samples. After conditioning the cartridges with 1 mL methanol, washing with 1

mL water, and drying, 600 µL of sample was loaded on each cartridge. The cartridges were again

washed with 1 mL water, dried and eluted with 800 µL methanol into a glass vial. This was dried,

resuspended in 40 µL of 50% methanol, and 5 µL injected on a Waters Acquity UPLC connected

to a Waters Xevo TQ-S mass spectrometer with a BEH C18 column (1.7 µm, 2.1*100mm) at the

Huck Metabolomics Core Facility. The mobile phases were A: methanol/water 2:98 v/v with 5

mM ammonium formate and 0.1% formic acid and B: methanol/water 98:2 v/v with 5 mM

ammonium formate and 0.1% formic acid. The gradient started with 0% B, then increased to 20%

B at 1.1 min, 90% B at 8 min, returned to 20% by 8.1 min, where it was held until 10 min. The

flow rate was kept at 0.3 mL/min and the column was heated to 40°C.

*Mating bioassays*

To examine the effect of sublethal imidacloprid exposure on mating, gynes and males were video

recorded in paired mating bioassays. To examine how males respond to sublethally-exposed gynes,

we placed one exposed and one unexposed sister gyne in a mating arena with 6 unexposed 10-day

old males, and video recorded all interactions with a smartphone (iPhone 6s) or GoPro Hero 6 for

15 minutes, or until both gynes had mated (Fig 1A). The males were unrelated to the gynes but

siblings with each other. This was done twice, once with one of the gynes exposed to 6 ppb

imidacloprid, and once with one of the gynes exposed to 60 ppb imidacloprid. To evaluate how

male pesticide exposure affects mating, we used the same setup, but inverted the exposure, with

the two gynes remaining unexposed and half of the males (3/6) receiving imidacloprid (Fig 1B).

The 1:3 ratio of gynes to males was maintained to increase the probability of successful mating

(Treanore *et al.* 2021). Gynes and males were taken from different colonies to remove possible

effects of inbreeding avoidance and nestmate recognition (Gerloff & Schmid-Hempel 2005;

Whitehorn, Tinsley & Goulson 2009). Bees were paint marked with red or white on the dorsal

thorax according to treatment. The color designation for treatment and control was alternated

between trials. Videos were scored for the duration of male mounting behavior, and any successful

mating was recorded to calculate mating frequency between groups. 'Mounting' was defined as

the male being attached to the gynes' abdomen in a position that precedes copulation. 'Mating'

was defined as the stereotypical posture of the male having fallen backwards so that he was

attached to the gyne only with his genitalia, with his legs twitching in the air.

*Chemical extracts and analysis*

All bees were frozen at -80° C prior to dissection or extraction of their cuticular hydrocarbons
(CHCs) and resulting samples were stored at -20° C until analysis. Gyne mandibular glands (n =
15 per treatment) were dissected by bisecting the head with razor blade and then using a micro
scissors to cut through the cuticle towards the mandible, exposing the gland. A clean, fine forceps
was then used to remove the gland into a glass insert in a 2 mL glass vial containing 100 µL of
dichloromethane and left to extract overnight. After removing the solvent with passive evaporation
in a fume hood, 5 µL of BSTFA with 1% TCMS (Millipore Sigma) was added to make
trimethylsilyl derivatives of the hydroxy acids typically found in the mandibular gland, which
greatly improves their chromatography. After 4 hours at room temperature the reaction was
quenched by adding 50 µL of hexane containing 1 µg of eicosane (as an internal standard) to each
sample.
218
Male labial glands (n = 20 per treatment) were dissected by pinning the head face down in a
Sylgard dissection dish and using a micro scissors to make an incision from the occipital foramen
towards the vertex of the head. This incision was used as an entry point to pry up and break off
pieces of cuticle and expose the labial glands, which were removed with fine forceps into a 2 mL
glass vial containing 500 µL of hexane with 10 µg eicosane as an internal standard. After overnight
extraction, a 100 µL aliquot was transferred to a clean glass vial with an insert and treated with
BSTFA as above. 100 µL of hexane was added to quench the reaction.
226
227 CHCs of gynes and males were extracted by immersing the bees in hexane for 30 seconds. In
228 correspondence with their body size, gynes were washed in 1.5 mL and males in 1 mL of hexane
229 containing either 30 µg (gynes) or 20 µg (males) eicosane as internal standard.
230
To identify compounds, samples were analyzed on an Agilent 7890A GC equipped with a HP-5
MS column (0.25 mm id × 30 m × 0.25 µm film thickness, Agilent) and interfaced to an Agilent
5975C mass spectrometer. Compounds were tentatively identified based on retention indices and
diagnostic ions, and when possible, identifications were confirmed by matching retention times
and mass spectra with those of authentic standards. The double bond positions of monounsaturated
cuticular hydrocarbons were determined by GC-MS analysis of their dimethyldisulfide (DMDS)
adducts (Dunkelblum et al. 1985), described in Table S6 and Figure S1. DMDS derivatizations

were carried out by adding 100 µl DMDS and 1 drop of 60 µg/µL $I_2$ in diethyl ether to the
evaporated CHC residue extracted from a single bee. The reaction was kept in a 2 mL screw-cap
vial overnight at 40° C, after which 200 µl of hexane was added, and the mixture washed with an
equal volume of 5% aqueous sodium thiosulfate. The organic layer was removed and dried over
anhydrous sodium sulfate for GC-MS analysis.
243
To quantify relative and absolute amounts of compounds, 1 µL of extracts was injected on a Trace
1310 GC (Thermo Fisher, Waltham, MA USA) equipped with a flame-ionization detector (FID)
and a TG-5MS column (0.25 mm id × 30 m × 0.25 µm film thick- ness; Thermo Fisher). The
temperature program was as follows: 60˚C to 120˚C at 15˚C/min followed by 120˚C to 300˚C (5
m hold) at 4˚C/min. The injector port and FID were held constant at 250˚C and 320˚C, respectively.
Analyses were conducted with Chromeleon version 7.2 SR4.
250
*Electroantennography*
To examine the impact of imidacloprid exposure on signal perception, antennal responses to gland
extracts (i.e., gyne antenna vs. male labial glands, and male antenna vs. gyne mandibular glands)
were recorded using electroantennography (EAG) as previously described (Derstine *et al.* 2021),
with slight modifications. The antennae of gynes (total sample size = 60; control - 30, 6 ppb – 15,
60 ppb - 15) and males (total sample size = 54; n = 18 per treatment group) fed with 0, 6, and 60
ppb imidacloprid were exposed to a control stimulus and a gland extract from the opposite sex.
Mandibular glands from 70 gynes and labial glands from 50 males were dissected into hexane to
create pooled extracts that ensured consistent stimuli across replicates. The pooled extracts were
evaporated to a concentration of 1 gland equivalent per 10 µL. Odor cartridges were created by
pipetting 1-bee equivalent of either mandibular or labial extract, or 10 µL hexane as a blank
control, onto strips of Whatman filter paper. These strips were inserted into glass pipettes after the
solvent evaporated and connected to the air delivery tube. For EAG preparations, a single antenna
was excised, and the tip snipped with fine scissors prior to mounting between the paddles of a
Syntech EAG probe with Spectra 360 electroconductive gel and placed in the air stream. Each
antenna was allowed to reach a steady baseline before starting the experiment, at which point the
test odorants were delivered into a steady airstream by using a stimulus flow controller (Syntech,
Hilversum, Netherlands) to pass air through the odor cartridges. Responses to each stimulus were

amplified with a 10 × pre-amplifier (Syntech, Netherlands) and further amplified 10 × before being

recorded and analyzed on a desktop computer using Syntech EAG software.

Each antenna was exposed to two control, and two gland extract puffs, all 15 seconds apart. For

each antenna, responses to test odorants were averaged and normalized by subtracting the average

response to the control odorants and the corrected values were used in analyses.

*Lipid measurement*

Total lipids per mass of gynes' fat body was measured with the vanillin-phosphoric acid bioassay

as in (Treanore & Amsalem 2020). Eviscerated abdomens (control n = 30, 6 ppb n = 27, 60 ppb n

= 32) were weighed and then homogenized in 1 mL 2% sodium sulfated solution. For each sample,

a 200 µL aliquot of the homogenate was added to 2.8 mL of 1:1 chloroform:methanol (v:v),

vortexed for 30 seconds, and centrifuged to precipitate glycogen (3000 rpm for 5 min). The

supernatant was transferred to a clean glass vial and mixed with an additional 2 mL distilled water.

The upper fraction was discarded while the lower fraction containing lipids was reserved. After

removing the solvent by evaporation, 200 µL sulfuric acid was added to the lipid residue and

heated for 10 min at 100°C. 5 mL of vanillin-phosphoric acid reagent was added, and after cooling,

300 µL was transferred to a 96-well plate in triplicate. Absorbance was measured at 525 nm using

a BioTek Microplate Spectrophotometer (BioTek Instruments Inc., Winooski, VT). For each

sample, resulting absorbance values measured against a standard curve which consisted of five

different concentrations of 0.1% vegetable oil diluted in chloroform. Absorbance values were

converted to micrograms based on a formula calculated from the regression line derived from the

standard curve. Lipid quantity is presented as lipids in µg per mg of fat body to correct for body

size.

*Sperm viability*

Sperm from 10-day old males exposed to 0 ppb, 6 ppb, and 60 ppb imidacloprid (n = 17 per

treatment group) were visualized using an Olympus Fluoview 10i-LIV confocal microscope after

staining with the Live/Dead Sperm Viability Kit (L-7011). This kit uses SYBR 14 and propidium

iodide to fluorescently stain live and dead sperm, respectively. The dyes were diluted according to

manufacturer's instructions in Kiev buffer and used on the day of preparation. Males treated with

0, 6, or 60 ppb imidacloprid dosed sucrose solution were dissected at 10 days old after cold

anesthesia. Accessory testes were removed into a 25 µL drop of Kiev buffer, and gently torn open

with forceps to allow sperm to flow out into the liquid. The liquid was transferred to a 2 mL

Eppendorf tube containing 25 µL Live/Dead dye working solution and incubated at 27° C for 10

minutes. 5 µL of stained sperm solution was pipetted onto a clean glass slide, covered with a glass

coverslip, and scanned in green and red wavelengths. From the resulting image, 10 subsections

were randomly selected, and all live and dead sperm were counted within each subsection to

calculate the proportion of live sperm. These subsections were also averaged to give the mean total

sperm per sampling area.

*Statistics*

Statistical analyses were performed using R (version 4.1.3) in RStudio. In mating trials, male

mounting behavior was analyzed with a paired Wilcoxon Rank Sum test, whereas the proportion

of successful mating between control and treatment groups was analyzed using a two-proportion

Z-test. Permutational analysis of variance (PERMANOVA) was used to examine overall chemical

differences in extracts of gyne mandibular glands, male labial glands, and CHCs between treatment

groups of gynes and males. Only peaks > 0.5% of the total peak area were used. These selected

peaks were then re-standardized to 100%. Individual components were compared using linear

mixed effects models with colony as a random factor and Tukey's Honest Significant Difference

post-hoc test to obtain pairwise contrasts of treatment groups. In the analyses of EAG, total lipids,

and sperm viability, we used linear mixed effects models with colony as a random effect to test

the impact of treatment on the response variable. Pairwise contrasts between treatment groups were

subsequently evaluated with Tukey's Honest Significant Difference post-hoc test.

**Results**

*Imidacloprid body concentration*

Gynes and males consuming sucrose solution containing imidacloprid for 3 days had comparable

amounts of imidacloprid in their body when given 6 ppb (gynes - 0.093 ng $\pm$ 0.02, males - 0.091

ng $\pm$ 0.01, mean $\pm$ SE, n = 10) and 60 ppb (gynes - 1.005 ng $\pm$ 0.31, males - 1.102 ng $\pm$ 0.18, mean

$\pm$ SE, n = 10), while imidacloprid was not found in any of the control gynes (Fig 2). The statistics

of all pairwise comparisons are given in Table S1.

*Effect of imidacloprid on mating behavior*

Treatment with 6 ppb imidacloprid reduced mating attempts by males toward exposed gynes (Fig 3). Gynes which consumed 6 ppb imidacloprid received less male mounting (p = 0.008, Wilcoxon Rank Sum test; $1.2 \pm 0.6$ and $3 \pm 0.7$ minutes, mean $\pm$ SE in 6 ppb vs. control groups, respectively, Fig 3A) and mated less frequently than unexposed gynes (p = 0.05 two-proportion Z-test, n = 30, 20% to 43% in 6 ppb vs. control groups, respectively, Fig 3B). In the 60 ppb treatment group, the overall mating success was significantly lower in both the treatment and control groups and there were no significant differences between treatment groups in male mounting (p = 0.97, Wilcoxon Rank Sum test, $0.27 \pm 0.2$ min and $0.45 \pm 0.2$ min in 6 ppb vs control groups, respectively, Fig 3C) or mating frequency (p = 0.09 two-proportion Z-test, n = 30, 3% to 14% in 6 ppb vs. control groups, respectively, Fig 3D).

In mating trials where only the males were exposed to imidacloprid, there was no effect of treatment on male mounting (p = 0.19, Wilcoxon Rank Sum test; $1.8 \pm 0.7$ and $3.4 \pm 1.0$ minutes, mean $\pm$ SE in 6 ppb vs. control groups, respectively, Fig 3E) or successful mating (p = 0.68 two-proportion Z-test, n = 27, 11% to 14% in 6 ppb vs control groups, respectively, Fig 3F).

*Effect of imidacloprid on chemical signaling*

To evaluate if altered chemical signaling was responsible for reduced mating success in bees exposed to imidacloprid, we examined CHCs in gynes and males, gyne mandibular glands and male labial glands in both control and treatments. Gyne and male CHCs were comprised primarily of saturated and unsaturated aliphatic alkanes and alkenes, with gynes having a much higher proportion of unsaturated compounds than males. These desaturations occurred predominately at the 9 and 7 positions for all chain lengths encountered. Gyne mandibular glands contained a series of 3-hydroxy acids in addition to hydrocarbons and several compounds which did not vary across treatments and thus their identification was not pursued. Male labial glands contained predominantly two compounds, 2,3-dihydrofarnesol (previously reported in (Luxová *et al.* 2004)), and hexadecenoic acid, in addition to a smaller proportion of hydrocarbons and farnesyl esters. As in the mandibular glands, because these compounds were not affected by the pesticide treatment, their complete identification was not pursued. The amounts of unidentified

compounds were included in all analyses, provided they met the 0.5% relative peak area cutoff.

The identity, retention indices, and average proportion of compounds used in statistical analyses

are given in Table S2 (Gyne CHCs), Table S3 (Male CHCs), Table S4 (Gyne mandibular gland),

and Table S5 (Male labial gland).

Gyne CHCs were altered by 60 ppb but not 6 ppb imidacloprid exposure. An NMDS plot of gyne

CHCs (Fig 4A) shows separation between the 60 ppb imidacloprid group and the two other

groups (0 and 6 ppb) which cluster together. This visualization is supported by pairwise

PERMANOVA which finds CHCs in the 60 ppb group significantly different from the 6 ppb (p

= 0.006) and the control (p = 0.05) groups. One compound in particular seems to drive this

separation, 9-pentacosene (Fig 4B). Simultaneous pairwise comparisons using a linear mixed

effects model with Tukey's HSD test indicated 9-pentacosene was significantly increased in the

60 ppb group, representing $18.1 \pm 0.5\%$ of the total peak area in the 60 ppb treatment group,

compared with $15.6 \pm 0.6\%$ in the 6 ppb treatment group and $15.3 \pm 0.4\%$ in the control group

(mean $\pm$ SE, n = 15; 60 ppb vs. 6 ppb p = 0.001, 60 ppb vs. control p = 0.0001, 6 ppb vs. control

p = 0.88).

Similarly, exposure to 60 ppb, but not 6 ppb imidacloprid, altered the overall male CHC profile

(Fig 4C, pairwise PERMANOVA, 60 ppb vs. 6 ppb p = 0.01, 60 ppb vs. control p = 0.02, 6 ppb

vs. control p = 0.36). However, this difference was not obviously explained by individual

compounds, and while the entire profile differed between the groups, no single individual

compound was significantly different between treatment groups (Fig 4D).

In contrast, imidacloprid exposure had no effect on the chemical composition of the gyne

mandibular gland (p = 0.81, PERMANOVA, Fig 4E-F) or the male labial gland (p = 0.45,

PERMANOVA, Fig 4G-H), at either 6 ppb or 60 ppb doses.

*Effect of imidacloprid on signal perception*

To evaluate if altered perception of chemical signaling was responsible for reduced mating success

in bees exposed to imidacloprid, we examined gyne perception of male labial gland secretion and

male perception of gyne mandibular gland secretion. EAG recordings of gynes responding to male

labial gland extract showed that imidacloprid did not affect signal perception at the tested doses

(Fig. 5A, B). A linear mixed effects model found no significant differences of 6 ppb treatment ($\chi^2$

= 0.56, $df$ = 1, p = 0.45), or 60 ppb treatment ($\chi^2$ = 0.004, $df$ = 1, p = 0.98) compared to their

corresponding controls. However, signal perception of males responding to gyne mandibular gland

extract was affected by the imidacloprid treatment (Fig 5C, $\chi^2$ = 7.7, $df$ = 2, p = 0.02). Post-hoc

comparisons between treatment groups showed that males in the 60 ppb, but not in the 6 ppb group,

had significantly lower EAG responses than controls (Tukey HSD, 60 ppb vs. control p = 0.02, 60

ppb vs. 6 ppb p = 0.11, 6 ppb vs. control p = 0.93).

*Effect of imidacloprid on gyne fat body lipid stores*

Imidacloprid exposure at the 60 ppb, but not 6 ppb dose, reduced gyne fat body lipid stores

compared to controls (Fig 6, $\chi^2$ = 15.5, $df$ = 2, p < 0.001; 133.4 ± 8.2, 115.1 ± 7.4 and 95.5 ± 4.3

µg lipid/mg fat body ± SE, n = 30/27/32 for 0, 6 and 60 ppb, respectively). Post-hoc comparisons

between treatment groups showed that the 60 ppb, but not the 6 ppb group, had significantly lower

fat body lipids than controls (Tukey HSD, 60 ppb vs. control p < 0.001, 60 ppb vs. 6 ppb p = 0.19,

6 ppb vs. control p = 0.09).

*The effect of imidacloprid on male sperm viability*

Imidacloprid exposure in males affected sperm viability in the 60 ppb but not in the 6 ppb group

(Fig 7). The mean total number of sperm was different between groups ($\chi^2$ = 9.4, $df$ = 2, p =

0.01), of which the 60 ppb group was reduced compared to controls (Tukey HSD: 60 ppb vs. 6

ppb p = 0.01, 60 ppb vs. 6 ppb p = 0.07, 6 ppb vs. control p = 0.71). The mean proportion of live

sperm was not significantly different between treatment groups ($\chi^2$ = 5.4, $df$ = 2, p = 0.07; Fig

7B).

**Discussion**

Our results show that chronic sublethal exposure to imidacloprid negatively affects mating

behavior in *B. impatiens* at doses as low as 6 ppb and identifies altered semiochemical production

and perception as a likely mechanism mediating this effect. The effect on mating was sex specific

and the gynes were more vulnerable to the same dose compared to the males, despite being 2-3

times larger than the males. Imidacloprid exposure altered the cuticular hydrocarbon profile of

both sexes and reduced male antennal sensitivity to the gyne mandibular glands at the 60 ppb dose.

Furthermore, we found negative effects on sperm viability in males and lipid storage in gynes

when exposed to 60 ppb imidacloprid. Given that mate location and copulation are primarily

regulated by chemical signaling in insects, the effects of pesticides on chemical communication

adds to growing concerns about the impact of widespread pesticide use on insect populations

(Forister, Pelton & Black 2019; Sánchez-Bayo & Wyckhuys 2019; Main *et al.* 2020; Minnameyer

*et al.* 2021; Willis Chan & Raine 2021).

While the 60 ppb dose negatively affected nearly all the chemical and physiological variables we

tested, including chemical production and perception, total lipids, and sperm viability, it did not

influence mating behavior. This outcome contrasts with the negative impact on mating behavior

observed at the 6 ppb dose. We suspect that the lack of behavioral impact at the 60 ppb dose is an

artifact of the overall lower mating frequency in that experiment, which likely reduced statistical

power and increased the risk of false negative (Type II error). Overall, bees clearly detected and

responded to changes in the 6 ppb treatment that were below the sensitivity of our instruments but

became evident with the higher 60 ppb dose.

Our data further indicate a stronger effect of the same dose on gynes compared to males. This is

evident in two key findings: first, gynes had comparable pesticide levels in their bodies to males,

despite being 2–3 times larger; and second, the treatment had an asymmetrical impact on mating

behavior, with males avoiding treated gynes, while gynes showed no such avoidance of treated

males. Sex differences in response to pesticide exposure have several possible explanations. Males

and females of many insect species have clear differences in overall body size and the amount of

fat body tissue, and in bees, also in the number of chromosomes. All of these could interact to

affect the processing and physiological response to pesticide exposure. In *B. impatiens,* gynes can

weigh 2-5x more than males (see Fig S2), which could impact both the ultimate body concentration

of pesticide and how much nectar is consumed. Once ingested, imidacloprid is primarily

metabolized rather than excreted (Suchail *et al.* 2004; Suchail, Debrauwer & Belzunces 2004).

This is significant because detoxification enzymes are predominantly produced in fat body tissue

(Li, Yu & Feng 2019), which is much more abundant in gynes than males (yet, was not more

effective in gynes vs. males in our data). Additionally, gynes are diploid. Haploid individuals can

be more susceptible to neonicotinoids than their diploid sisters, likely because of reduced variation

in detoxification genes (Friedli *et al.* 2020). The combined effects of these interactions argue that

males should suffer greater effects from pesticide exposure than females. However, in our

experiments and others (Boff & Ayasse 2023), sublethal pesticide exposure did not significantly

reduce mating when only males were exposed, as it did when only females were exposed. In

addition, the concentration of pesticides provided and the total amount found in the body of males

and gynes was the same, despite differences in body size. Thus, it is more likely that the differences

in behavior of pesticide exposed males and gynes found in this study are a result of differences in

the level of choosiness by gynes and males and the range at which mate choice takes place. This

is in line with the assumption that bumble bee gynes choose males from a long range by attracting

to their labial gland secretion and mate choice in the short range is likely determined by the males.

Differences in pesticide levels in the body could be influenced not only by body size but also by

consumption, as the pesticides were administered chronically. However, sugar consumption was

not measured in this study, so we cannot determine whether males and gynes arrived at similar

amounts of imidacloprid in the body because they regulated their consumption differentially, or if

they had different clearance rates. This would be a worthwhile subject of future investigation.

Given the mating system of *B. impatiens*, where males can mate multiply but gynes rarely do (Starr

1984), gynes should be more incentivized to choose high quality mates than males (Boomsma,

Kronauer & Pedersen 2009). In this study, gynes were less selective than males. Generally, it is

assumed that long distance mate-finding in *Pyrobombus* bumble bees is mediated by female

attraction to male labial gland pheromones (Valterová *et al.* 2019), and that less volatile CHCs

may play a role once contact is initiated. Thus, females have an opportunity to assess and choose

males both in the initial long-range attraction phase, and in close pre-copulatory behavior, while

male choice can only be exercised in the later. One possible explanation for the data presented here

is that by placing males and gynes together in a mating arena, the long-distance mate finding step

mediated by male labial gland pheromones was circumvented, preventing one avenue of mate

discrimination or assessment (by gynes), which translates into negative impacts on mating in that

sex.

Under honest signaling theory, chemical signals are expected to convey true and reliable

information in order to be evolutionarily stable (Smith & Harper 2003). In moths, for example, sex

pheromone signals have been shown to covary with fitness and convey information about

individual quality (Foster & Johnson 2011; Blankers *et al.* 2021). In this context, pheromones

should convey accurate information about mate quality. However, the putative volatile sex

pheromone compounds of both sexes were unaffected by imidacloprid treatment at doses which

impacted clear components of what would make a quality mate - sperm viability in males and lipid

stores in gynes. This would suggest that these signals are not honest, or at least are not tied to these

measures of quality and warrants more research to identify the specific compounds responsible for

sex attraction in *B. impatiens.* Instead, the profile of cuticular hydrocarbons of both males and

females was altered when bees ingested imidacloprid, potentially acting as honest signals.

It is important to note that while the male labial glands were shown to contain a sex pheromone

across multiple species of bumble bees (Valterová *et al.* 2019), the production of a sex pheromone

in the gyne mandibular gland is not a well-established phenomenon. One study relied on a highly

invasive gland excision surgery with low replication and variable results in different species (van

Honk, Velthuis & Röseler 1978). Another demonstrated that while a freshly frozen gyne could

elicit copulation from males, extracts from both the cuticular surface and whole head (but not

specifically the mandibular glands) were successful in eliciting mounting, but not copulation, from

males (Krieger *et al.* 2006). These extracts contained GC-EAD active compounds but were not

localized to a particular gland. Urbanová et al subsequently found many of the

electrophysiologically active compounds, including the 3-hydroxy acids, in the mandibular glands

(Urbanová *et al.* 2008). This indicates that the male response may require multi-modal signals to

copulate, and that a whole bee extract is more likely to contain the sex pheromone, given that

compounds from multiple glands may be spread onto the cuticle to elicit copulation.

Pesticides' effects on mating and sexual communication are often elusive and hard to detect since

they manifest in species- and sex-specific ways and do not always affect fitness immediately. For

example, in the mason bee *Osmia cornuta,* fungicide exposure alters male vibrational and cuticular

signals which decreases mating (Boff *et al.* 2022), while in another megachilid, *Heriades*

*truncorum,* pesticide exposure affects precopulatory behaviors in males and cuticular signals

females (Boff & Ayasse 2023). Similar to our findings, pesticide exposure did not affect the

proportion of successful mating in *H. truncorum* in males but did in females.

Overall, our data show reduced mating success in gynes exposed to imidacloprid accompanied by

changes in their cuticular profiles and the responsiveness of males to their chemical secretion. We

further demonstrate fitness consequences to imidacloprid exposure in both gynes and males and

together, propose a plausible mechanism explaining the impact of imidacloprid on mating with

disrupt sexual communication. Such disruption to communication systems is not likely to be

immediately apparent in insects and its consequences may be extended over several generations.

These data add up to the growing concerns of the use of sublethal, field relevant, doses of

neonicotinoids and their impacts on pollinator health. Finally, to control as many variables as

possible, we chose an exposure scenario of 3 days of adult exposure, which is conservative with

regards to exposure scenarios of wild bees. In a natural setting, bees might contend with pesticide

exposure during development (Kozii *et al.* 2021) and potentially continuously as adults, both of

which have been shown to have detrimental effects on queens.

Figure 1 – Diagram of reciprocal mating experiments. In "A" one of two gynes was fed on
imidacloprid dosed sugar solution and then placed in a mating arena with six unexposed males.
In "B", three of the six males were fed imidacloprid, while the gynes were unexposed.

Figure 2 - LCMS analysis of imidacloprid amount in individual gynes and males after 3 days of
feeding on either control, 6 ppb, or 60 ppb imidacloprid-dosed sugar solution. Letters above the
boxplots indicate statistical differences determined by pairwise contrasts using the Kruskal-
Wallis test followed by Dunn's post hoc test with Benjamini and Hochberg adjusted p-values ($\alpha$
= 0.05).

Figure 3 – The impact of imidacloprid exposure on mounting behavior by males (A,C,E) and mating
behavior (B,D,F) when either gynes or males were fed imidacloprid. Panels A-D correspond to mating
trials where half of the gynes were fed imidacloprid and, panels E-F correspond to mating trials where
half of the males were fed imidacloprid. Asterisks denote significant differences at p < 0.05.

Figure 4 – The effect of pesticide exposure on signal production. Panels A, C, E, G show NMDS plots
based on the Bray-Curtis dissimilarity matrix of relative amounts of chemical components in gland
extracts or cuticular washes. Panels B, D, F, H show relative amounts of each chemical component
quantified from GC-FID and GC-MS analyses. Points are means ± 2SE (to approximate 95%
confidence intervals). Panels A,B and C,D describe gyne and male cuticular hydrocarbon data,
respectively. Panels E,F and G,H describe gyne mandibular gland and male labial gland extracts,
respectively.

Figure 5 – The effect of imidacloprid exposure on signal perception. Panels show antennal responses of gynes (A, B) and males (C) exposed to either control, 6 ppb, or 60 ppb imidacloprid after receiving puffs of male labial gland extract. As the 6 ppb and 60 ppb treatments were run separately for gynes, they are presented separately and next to their concurrently run controls. Letters above boxplots denote statistical differences of pairwise contrasts at $p < 0.05$ (calculated from a linear mixed effects model with colony as a random factor, using Tukey's HSD post-hoc test for pairwise comparisons).

560      Figure 6 – Lipid stores of 6-days old gynes fed either 0, 6, or 60 ppb imidacloprid in sugar solution.
561      Values are presented in micrograms of lipid per milligram of fat body tissue. Letters above boxplots
562      indicate significant differences (linear mixed effects model followed by Tukey HSD, $\alpha = 0.05$).
563

565  Figure 7 – A – The mean number of sperm counted per subsection, per male. B – The proportion of live
566  sperm in each treatment group. Letters above boxplots indicate significant differences (linear mixed
567  effects model followed by Tukey HSD, α = 0.05). C – Example image of fluorescently dyed *B. impatiens*
568  sperm (green indicates living and red indicates dead sperm).
569

570
571

References

Aizen, M.A., Garibaldi, L.A., Cunningham, S.A. & Klein, A.M. (2009) How much does
agriculture depend on pollinators? Lessons from long-term trends in crop production.
*Annals of Botany,* **103,** 1579-1588.
Amsalem, E., Twele, R., Francke, W. & Hefetz, A. (2009) Reproductive competition in the
bumble-bee *Bombus terrestris*: do workers advertise sterility? *Proc Biol Sci,* **276,** 1295-
1304.
Ayasse, M. & Jarau, S. (2014) Chemical ecology of bumble bees. *Annu Rev Entomol,* **59,** 299-
319.
Ayasse, M., Paxton, R.J. & Tengö, J. (2001) Mating behavior and chemical communication in
the order Hymenoptera. *Annual review of entomology,* **46,** 31-78.
Bantz, A., Camon, J., Froger, J.-A., Goven, D. & Raymond, V. (2018) Exposure to sublethal
doses of insecticide and their effects on insects at cellular and physiological levels.
*Current Opinion in Insect Science,* **30,** 73-78.
Baron, G.L., Jansen, V.A.A., Brown, M.J.F. & Raine, N.E. (2017) Pesticide reduces bumblebee
colony initiation and increases probability of population extinction. *Nat Ecol Evol,* **1,**
1308-1316.
Bergman, P. & Bergström, G. (1997) Scent marking, scent origin, and species specificity in male
premating behavior of two Scandinavian bumblebees. *Journal of Chemical Ecology,* **23,**
1235-1251.
Bird, S.A., Pope, N.S., McGrady, C.M., Fleischer, S.J. & López-Uribe, M.M. (2024) Mating
frequency estimation and its importance for colony abundance analyses in eusocial
pollinators: a case study of *Bombus impatiens* (Hymenoptera: Apidae). *Journal of*
*economic entomology.*
Blacquière, T., Smagghe, G., van Gestel, C.A.M. & Mommaerts, V. (2012) Neonicotinoids in
bees: a review on concentrations, side-effects and risk assessment. *Ecotoxicology,* **21,**
973-992.
Blankers, T., Lievers, R., Plata, C., van Wijk, M., van Veldhuizen, D. & Groot, A.T. (2021) Sex
pheromone signal and stability covary with fitness. *Royal Society open science,* **8,**
210180.
Boff, S. & Ayasse, M. (2023) Exposure to sublethal concentration of flupyradifurone alters
sexual behavior and cuticular hydrocarbon profile in *Heriades truncorum*, an oligolectic
solitary bee. *Insect Science.*
Boff, S., Conrad, T., Raizer, J., Wehrhahn, M., Bayer, M., Friedel, A., Theodorou, P., Schmitt, T.
& Lupi, D. (2022) Low toxicity crop fungicide (fenbuconazole) impacts reproductive
male quality signals leading to a reduction of mating success in a wild solitary bee.
*Journal of Applied Ecology,* **59,** 1596-1607.
Boomsma, J.J., Kronauer, D.J. & Pedersen, J. (2009) The evolution of social insect mating
systems. *Organization of insect societies*, 3-25.
Brühl, C.A. & Zaller, J.G. (2019) Biodiversity decline as a consequence of an inappropriate
environmental risk assessment of pesticides. *Frontiers in Environmental Science*, 177.
Bruschini, C., Cervo, R. & Turilazzi, S. (2010) Pheromones in Social Wasps. *Pheromones*, pp.
447-492.

Cameron, S.A., Lozier, J.D., Strange, J.P., Koch, J.B., Cordes, N., Solter, L.F. & Griswold, T.L. (2011) Patterns of widespread decline in North American bumble bees. *Proc Natl Acad Sci U S A,* **108,** 662-667.

Chaimanee, V., Evans, J.D., Chen, Y., Jackson, C. & Pettis, J.S. (2016) Sperm viability and gene expression in honey bee queens (*Apis mellifera*) following exposure to the neonicotinoid insecticide imidacloprid and the organophosphate acaricide coumaphos. *J Insect Physiol,* **89,** 1-8.

Chauzat, M.-P., Faucon, J.-P., Martel, A.-C., Lachaize, J., Cougoule, N. & Aubert, M. (2006) A survey of pesticide residues in pollen loads collected by honey bees in France. *Journal of economic entomology,* **99,** 253-262.

Chen, X., Wang, Y., Zhou, Y., Wang, F., Wang, J., Yao, X., Imran, M. & Luo, S. (2024) Imidacloprid reduces the mating success of males in bumblebees. *Science of The Total Environment,* **928,** 172525.

Ciereszko, A., Wilde, J., Dietrich, G.J., Siuda, M., Bąk, B., Judycka, S. & Karol, H. (2016) Sperm parameters of honeybee drones exposed to imidacloprid. *Apidologie,* **48,** 211-222.

Colin, T., Meikle, W.G., Wu, X. & Barron, A.B. (2019) Traces of a Neonicotinoid Induce Precocious Foraging and Reduce Foraging Performance in Honey Bees. *Environmental science & technology,* **53,** 8252-8261.

Crall, J.D., Switzer, C.M., Oppenheimer, R.L., Ford Versypt, A.N., Dey, B., Brown, A., Eyster, M., Guerin, C., Pierce, N.E., Combes, S.A. & de Bivort, B.L. (2018) Neonicotinoid exposure disrupts bumblebee nest behavior, social networks, and thermoregulation. *Science,* **362,** 683-686.

Cresswell, J.E., Page, C.J., Uygun, M.B., Holmbergh, M., Li, Y., Wheeler, J.G., Laycock, I., Pook, C.J., de Ibarra, N.H., Smirnoff, N. & Tyler, C.R. (2012) Differential sensitivity of honey bees and bumble bees to a dietary insecticide (imidacloprid). *Zoology,* **115,** 365-371.

Cresswell, J.E., Robert, F.X., Florance, H. & Smirnoff, N. (2014) Clearance of ingested neonicotinoid pesticide (imidacloprid) in honey bees (*Apis mellifera*) and bumblebees (*Bombus terrestris*). *Pest Manag Sci,* **70,** 332-337.

Derstine, N., Villar, G., Orlova, M., Hefetz, A., Millar, J. & Amsalem, E. (2021) Dufour's gland analysis reveals caste and physiology specific signals in *Bombus impatiens. Sci Rep,* **11,** 2821.

Douglas, M.R. & Tooker, J.F. (2015) Large-Scale Deployment of Seed Treatments Has Driven Rapid Increase in Use of Neonicotinoid Insecticides and Preemptive Pest Management in U.S. Field Crops. *Environmental science & technology,* **49,** 5088-5097.

Forister, M.L., Pelton, E.M. & Black, S.H. (2019) Declines in insect abundance and diversity: We know enough to act now. *Conservation Science and Practice,* **1**.

Foster, S.P. & Johnson, C.P. (2011) Signal Honesty through Differential Quantity in the Female-Produced Sex Pheromone of the Moth *Heliothis virescens. Journal of Chemical Ecology,* **37,** 717-723.

Franklin, E.L. & Raine, N.E. (2019) Moving beyond honeybee-centric pesticide risk assessments to protect all pollinators. *Nature Ecology & Evolution,* **3,** 1373-1375.

Friedli, A., Williams, G.R., Bruckner, S., Neumann, P. & Straub, L. (2020) The weakest link: Haploid honey bees are more susceptible to neonicotinoid insecticides. *Chemosphere,* **242,** 125145.

661    Gerloff, C.U. & Schmid-Hempel, P. (2005) Inbreeding depression and family variation in a
662             social insect, *Bombus terrestris* (Hymenoptera: Apidae). *Oikos,* **111,** 67-80.
663    Gill, R.J. & Raine, N.E. (2014) Chronic impairment of bumblebee natural foraging behaviour
664             induced by sublethal pesticide exposure. *Functional Ecology,* **28,** 1459-1471.
665    Goulson, D., Nicholls, E., Botias, C. & Rotheray, E.L. (2015) Bee declines driven by combined
666             stress from parasites, pesticides, and lack of flowers. *Science,* **347,** 1255957.
667    Gradish, A.E., Steen, J.v.d., Scott-Dupree, C.D., Cabrera, A.R., Cutler, G.C., Goulson, D., Klein,
668             O., Lehmann, D.M., Lückmann, J., O'Neill, B., Raine, N.E., Sharma, B. & Thompson, H.
669             (2018) Comparison of Pesticide Exposure in Honey Bees (Hymenoptera: Apidae) and
670             Bumble Bees (Hymenoptera: Apidae): Implications for Risk Assessments. *Environmental*
671             *Entomology,* **48,** 12-21, 10.
672    Hahn, D.A. & Denlinger, D.L. (2007) Meeting the energetic demands of insect diapause: nutrient
673             storage and utilization. *J Insect Physiol,* **53,** 760-773.
674    Heimpel, G.E. & de Boer, J.G. (2008) Sex determination in the hymenoptera. *Annu Rev Entomol,*
675             **53,** 209-230.
676    Jeschke, P., Nauen, R., Schindler, M. & Elbert, A. (2011) Overview of the status and global
677             strategy for neonicotinoids. *Journal of agricultural and food chemistry,* **59,** 2897-2908.
678    Jordan, A., Patch, H.M., Grozinger, C.M. & Khanna, V. (2021) Economic dependence and
679             vulnerability of United States agricultural sector on insect-mediated pollination service.
680             *Environmental science & technology,* **55,** 2243-2253.
681    Keeling, C.I., Plettner, E. & Slessor, K.N. (2004) Hymenopteran semiochemicals. *Top Curr*
682             *Chem,* **239,** 133-177.
683    Klein, A.M., Vaissiere, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C. &
684             Tscharntke, T. (2007) Importance of pollinators in changing landscapes for world crops.
685             *Proc Biol Sci,* **274,** 303-313.
686    Kozii, I.V., Barnsley, S., Silva, M.C.B.d., Wood, S.C., Klein, C.D., de Mattos, I.M., Zabrodski,
687             M.W., Silva, R.d.C.M., Fabela, C.I.O., Guillemin, L., Dvylyuk, I., Ferrari, M.C.O. &
688             Simko, E. (2021) Reproductive fitness of honey bee queens exposed to thiamethoxam
689             during development. *Veterinary Pathology,* **58,** 1107-1118.
690    Krieger, G.M., Duchateau, M.J., Van Doorn, A., Ibarra, F., Francke, W. & Ayasse, M. (2006)
691             Identification of queen sex pheromone components of the bumblebee *Bombus terrestris.*
692             *J Chem Ecol,* **32,** 453-471.
693    Krischik, V.A., Landmark, A.L. & Heimpel, G.E. (2007) Soil-Applied Imidacloprid Is
694             Translocated to Nectar and Kills Nectar-Feeding *Anagyrus pseudococci* (Girault)
695             (Hymenoptera: Encyrtidae). *Environmental Entomology,* **36,** 1238-1245.
696    Krupke, C.H., Hunt, G.J., Eitzer, B.D., Andino, G. & Given, K. (2012) Multiple routes of
697             pesticide exposure for honey bees living near agricultural fields. *PLoS One,* **7,** e29268.
698    Kubo, R., Harano, K.-i. & Ono, M. (2017) Male scent-marking pheromone of *Bombus ardens*
699             *ardens* (Hymenoptera; Apidae) attracts both conspecific queens and males. *The Science*
700             *of Nature,* **104,** 1-5.
701    Laycock, I., Lenthall, K.M., Barratt, A.T. & Cresswell, J.E. (2012) Effects of imidacloprid, a
702             neonicotinoid pesticide, on reproduction in worker bumble bees (*Bombus terrestris*).
703             *Ecotoxicology,* **21,** 1937-1945.
704    Lecocq, T., Coppée, A., Mathy, T., Lhomme, P., Cammaerts-Tricot, M.-C., Urbanová, K.,
705             Valterová, I. & Rasmont, P. (2015) Subspecific differentiation in male reproductive traits
706             and virgin queen preferences, in *Bombus terrestris. Apidologie,* **46,** 595-605.

Li, S., Yu, X. & Feng, Q. (2019) Fat Body Biology in the Last Decade. *Annual review of*
*entomology,* **64,** 315-333.
Lürling, M. & Scheffer, M. (2007) Info-disruption: pollution and the transfer of chemical
information between organisms. *Trends in Ecology & Evolution,* **22,** 374-379.
Luxová, A., Urbanová, K., Valterová, I., Terzo, M. & Borg-Karlson, A.K. (2004) Absolute
configuration of chiral terpenes in marking pheromones of bumblebees and cuckoo
bumblebees. *Chirality: The Pharmacological, Biological, and Chemical Consequences of*
*Molecular Asymmetry,* **16,** 228-233.
Main, A.R., Webb, E.B., Goyne, K.W. & Mengel, D. (2020) Reduced species richness of native
bees in field margins associated with neonicotinoid concentrations in non-target soils.
*Agriculture, Ecosystems & Environment,* **287,** 106693.
Minnameyer, A., Strobl, V., Bruckner, S., Camenzind, D.W., Van Oystaeyen, A., Wäckers, F.,
Williams, G.R., Yañez, O., Neumann, P. & Straub, L. (2021) Eusocial insect declines:
Insecticide impairs sperm and feeding glands in bumblebees. *Science of The Total*
*Environment,* **785,** 146955.
Moffat, C., Buckland, S.T., Samson, A.J., McArthur, R., Chamosa Pino, V., Bollan, K.A.,
Huang, J.T.-J. & Connolly, C.N. (2016) Neonicotinoids target distinct nicotinic
acetylcholine receptors and neurons, leading to differential risks to bumblebees. *Scientific*
*reports,* **6,** 1-10.
Müller, C. (2018) Impacts of sublethal insecticide exposure on insects — Facts and knowledge
gaps. *Basic and Applied Ecology,* **30,** 1-10.
Muth, F., Francis, J.S. & Leonard, A.S. (2019) Modality-specific impairment of learning by a
neonicotinoid pesticide. *Biology letters,* **15,** 20190359.
Navarro-Roldán, M.A. & Gemeno, C. (2017) Sublethal Effects of Neonicotinoid Insecticide on
Calling Behavior and Pheromone Production of Tortricid Moths. *Journal of Chemical*
*Ecology,* **43,** 881-890.
Ollerton, J., Winfree, R. & Tarrant, S. (2011) How many flowering plants are pollinated by
animals? *Oikos,* **120,** 321-326.
Plowright, R. & Laverty, T. (1984) The ecology and sociobiology of bumble bees. *Annual*
*review of entomology,* **29,** 175-199.
Potts, S.G., Biesmeijer, J.C., Kremen, C., Neumann, P., Schweiger, O. & Kunin, W.E. (2010)
Global pollinator declines: trends, impacts and drivers. *Trends Ecol Evol,* **25,** 345-353.
Potts, S.G., Imperatriz-Fonseca, V., Ngo, H.T., Aizen, M.A., Biesmeijer, J.C., Breeze, T.D.,
Dicks, L.V., Garibaldi, L.A., Hill, R. & Settele, J. (2016) Safeguarding pollinators and
their values to human well-being. *Nature,* **540,** 220-229.
Raine, N.E. & Rundlöf, M. (2024) Pesticide Exposure and Effects on Non-Apis Bees. *Annual*
*review of entomology,* **69,** 551-576.
Ridley, M. (1990) The control and frequency of mating in insects. *Functional Ecology***,** 75-84.
Rondeau, S., Baert, N., McArt, S. & Raine, N.E. (2022) Quantifying exposure of bumblebee
(*Bombus spp.*) queens to pesticide residues when hibernating in agricultural soils.
*Environmental Pollution,* **309,** 119722.
Rundlöf, M., Andersson, G.K., Bommarco, R., Fries, I., Hederström, V., Herbertsson, L.,
Jonsson, O., Klatt, B.K., Pedersen, T.R. & Yourstone, J. (2015) Seed coating with a
neonicotinoid insecticide negatively affects wild bees. *Nature,* **521,** 77-80.
Sánchez-Bayo, F. & Wyckhuys, K.A.G. (2019) Worldwide decline of the entomofauna: A
review of its drivers. *Biological Conservation,* **232,** 8-27.

Sandrock, C., Tanadini, L.G., Pettis, J.S., Biesmeijer, J.C., Potts, S.G. & Neumann, P. (2014) Sublethal neonicotinoid insecticide exposure reduces solitary bee reproductive success. *Agricultural and Forest Entomology,* **16,** 119-128.

Schmid-Hempel, R. & Schmid-Hempel, P. (2000) Female mating frequencies in *Bombus spp.* from Central Europe. *Insectes Sociaux,* **47,** 36-41.

Schmuck, R., Schöning, R., Stork, A. & Schramel, O. (2001) Risk posed to honeybees (*Apis mellifera* L, Hymenoptera) by an imidacloprid seed dressing of sunflowers. *Pest Management Science: formerly Pesticide Science,* **57,** 225-238.

Sgolastra, F., Hinarejos, S., Pitts-Singer, T.L., Boyle, N.K., Joseph, T., Lūckmann, J., Raine, N.E., Singh, R., Williams, N.M. & Bosch, J. (2018) Pesticide Exposure Assessment Paradigm for Solitary Bees. *Environmental Entomology,* **48,** 22-35.

Siviter, H., Horner, J., Brown, M.J. & Leadbeater, E. (2020) Sulfoxaflor exposure reduces egg laying in bumblebees *Bombus terrestris*. *Journal of Applied Ecology,* **57,** 160-169.

Siviter, H. & Muth, F. (2022) Exposure to the novel insecticide flupyradifurone impairs bumblebee feeding motivation, learning, and memory retention. *Environmental Pollution,* **307,** 119575.

Smith, J.M. & Harper, D. (2003) *Animal signals*. Oxford University Press.

Stanley, D.A., Smith, K.E. & Raine, N.E. (2015) Bumblebee learning and memory is impaired by chronic exposure to a neonicotinoid pesticide. *Sci Rep,* **5,** 16508.

Starr, C.K. (1984) Sperm competition, kinship, and sociality in the aculeate Hymenoptera. *Sperm competition and the evolution of animal mating systems,* **428,** 459.

Stoner, K.A. & Eitzer, B.D. (2012) Movement of soil-applied imidacloprid and thiamethoxam into nectar and pollen of squash (Cucurbita pepo). *PLoS One,* **7,** e39114.

Straub, L., Villamar-Bouza, L., Bruckner, S., Chantawannakul, P., Gauthier, L., Khongphinitbunjong, K., Retschnig, G., Troxler, A., Vidondo, B., Neumann, P. & Williams, G.R. (2016) Neonicotinoid insecticides can serve as inadvertent insect contraceptives. *Proc Biol Sci,* **283**.

Stuligross, C. & Williams, N.M. (2020) Pesticide and resource stressors additively impair wild bee reproduction. *Proceedings of the Royal Society B,* **287,** 20201390.

Suchail, S., De Sousa, G., Rahmani, R. & Belzunces, L.P. (2004) In vivo distribution and metabolisation of 14C-imidacloprid in different compartments of *Apis mellifera* L. *Pest Management Science: formerly Pesticide Science,* **60,** 1056-1062.

Suchail, S., Debrauwer, L. & Belzunces, L.P. (2004) Metabolism of imidacloprid in *Apis mellifera*. *Pest Management Science: formerly Pesticide Science,* **60,** 291-296.

Svensson, B. & Bergstrom, G. (1977) Volatile marking secretions from the labial gland of North European *Pyrobombus* D.T. males (Hymenoptera, Apidae).

Tappert, L., Pokorny, T., Hofferberth, J. & Ruther, J. (2017) Sublethal doses of imidacloprid disrupt sexual communication and host finding in a parasitoid wasp. *Scientific reports,* **7,** 1-9.

Tasman, K., Rands, S.A. & Hodge, J.J.L. (2020) The Neonicotinoid Insecticide Imidacloprid Disrupts Bumblebee Foraging Rhythms and Sleep. *iScience,* **23,** 101827.

Tosi, S., Burgio, G. & Nieh, J.C. (2017) A common neonicotinoid pesticide, thiamethoxam, impairs honey bee flight ability. *Sci Rep,* **7,** 1201.

Tosi, S., Sfeir, C., Carnesecchi, E., vanEngelsdorp, D. & Chauzat, M.-P. (2022) Lethal, sublethal, and combined effects of pesticides on bees: A meta-analysis and new risk assessment tools. *Science of The Total Environment,* **844,** 156857.

Treanore, E. & Amsalem, E. (2020) The effect of intrinsic physiological traits on diapause
survival and their underlying mechanisms in an annual bee species *Bombus impatiens*.
*Conservation Physiology,* **8,** coaa103.
Treanore, E., Barie, K., Derstine, N., Gadebusch, K., Orlova, M., Porter, M., Purnell, F. &
Amsalem, E. (2021) Optimizing Laboratory Rearing of a Key Pollinator, *Bombus
impatiens*. *Insects,* **12,** 673.
Tricoire-Leignel, H., Thany, S., Gadenne, C. & Anton, S. (2012) Pest Insect Olfaction in an
Insecticide-Contaminated Environment: Info-Disruption or Hormesis Effect. *Frontiers in
Physiology,* **3**.
Tsvetkov, N., Samson-Robert, O., Sood, K., Patel, H., Malena, D., Gajiwala, P., Maciukiewicz,
P., Fournier, V. & Zayed, A. (2017) Chronic exposure to neonicotinoids reduces honey
bee health near corn crops. *Science,* **356,** 1395-1397.
Urbanová, K., Cahlíková, L., Hovorka, O., Ptáček, V. & Valterová, I. (2008) Age-dependent
changes in the chemistry of exocrine glands of *Bombus terrestris* queens. *Journal of
Chemical Ecology,* **34,** 458-466.
Valterová, I., Martinet, B., Michez, D., Rasmont, P. & Brasero, N. (2019) Sexual attraction: a
review of bumblebee male pheromones. *Z Naturforsch C J Biosci,* **74,** 233-250.
van Honk, C.G.J., Velthuis, H.H.W. & Röseler, P.F. (1978) A sex pheromone from the
mandibular glands in bumblebee queens. *Experientia,* **34,** 838-839.
Van Wilgenburg, E., Driessen, G. & Beukeboom, L.W. (2006) Single locus complementary sex
determination in Hymenoptera: an" unintelligent" design? *Frontiers in Zoology,* **3,** 1-15.
Whitehorn, P.R., O'Connor, S., Wackers, F.L. & Goulson, D. (2012) Neonicotinoid pesticide
reduces bumble bee colony growth and queen production. *Science,* **336,** 351-352.
Whitehorn, P.R., Tinsley, M.C. & Goulson, D. (2009) Kin recognition and inbreeding reluctance
in bumblebees. *Apidologie,* **40,** 627-633.
Willis Chan, D.S. & Raine, N.E. (2021) Population decline in a ground-nesting solitary squash
bee (*Eucera pruinosa*) following exposure to a neonicotinoid insecticide treated crop
(*Cucurbita pepo*). *Scientific reports,* **11,** 4241.
Wu-Smart, J. & Spivak, M. (2018) Effects of neonicotinoid imidacloprid exposure on bumble
bee (Hymenoptera: Apidae) queen survival and nest initiation. *Environmental
Entomology,* **47,** 55-62.
Yew, J.Y. & Chung, H. (2015) Insect pheromones: An overview of function, form, and
discovery. *Progress in Lipid Research,* **59,** 88-105.
Zioga, E., Kelly, R., White, B. & Stout, J.C. (2020) Plant protection product residues in plant
pollen and nectar: A review of current knowledge. *Environmental Research,* **189,**
109873.
835