Title: Caliber of zebrafish touch-sensory axons is dynamic in vivo

Abbreviated title: Somatosensory axon caliber

Authors: Kaitlin Ching[1*], Alvaro Sagasti[1†]

Affiliations: [1] Department of Cell, Molecular, and Developmental Biology, University of California,

Los Angeles, Los Angeles, California 90095, USA

*Corresponding author: Kaitlin Ching (kaitlinching@ucla.edu)

[†]Corresponding author: Alvaro Sagasti (sagasti@mcdb.ucla.edu)

number of pages: 36

number of figures: 8, videos: 4

number of words in abstract: 230, intro: 417, discussion: 1665

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: The studies were supported by National Institutes of Health grant

R01AR064582. The content is solely the responsibility of the authors and does not necessarily

represent the official views of the National Institutes of Health. We thank Van Savage for

valuable discussions and suggestions regarding branch analysis, Melissa Emami for providing

the Isl1[ss]:lexA, lexAop:mRuby-caax fish, D'Juan Farmer and the UCLA Broad Stem Cell

Research Center Microscopy Core for microscope access, and the UCLA Zebrafish Core

Facility for animal care. We thank the Sagasti, Farmer, and Koehler labs for helpful discussion

and suggestions.

Abstract

Cell shape is crucial to cell function, particularly in neurons. The cross-sectional diameter, also

known as caliber, of axons and dendrites is an important parameter of neuron shape, best

appreciated for its influence on the speed of action potential propagation. Many studies of axon

caliber focus on cell-wide regulation and assume that caliber is static. Here, we have

characterized local variation and dynamics of axon caliber in vivo using the peripheral axons of

zebrafish touch-sensing neurons at embryonic stages, prior to sex determination. To obtain

absolute measurements of caliber in vivo, we paired sparse membrane labeling with super-

resolution microscopy of neurons in live fish. We observed that axon segments had varicose or

"pearled" morphologies, and thus vary in caliber along their length, consistent with reports from

mammalian systems. Sister axon segments originating from the most proximal branch point in

the axon arbor had average calibers that were uncorrelated with each other. Axon caliber also

tapered across the branch point. Varicosities and caliber, overall, were dynamic on the

timescale of minutes, and dynamicity changed over the course of development. By measuring

the caliber of axons adjacent to dividing epithelial cells, we found that skin cell division is one

aspect of the cellular microenvironment that may drive local differences and dynamics in axon

caliber. Our findings support the possibility that spatial and temporal variation in axon caliber

could significantly influence neuronal physiology.

Significance Statement

Axon caliber directly influences how quickly neurons send messages to other cells and likely

plays a role in neurons' overall health. In the peripheral nervous system, where neurons cover

particularly long distances, cell shape can determine whether an animal successfully executes

behaviors such as escape responses. We found that axon caliber can vary between locations

within the same cell and is highly dynamic. Taking these variations into account may allow

neuroscientists to better estimate transmission speeds for cells in neural circuits. We observed

that axon caliber is distorted when nearby skin cells change shape. Thus, cells not classically

considered part of the nervous system can also contribute to caliber dynamics, broadening our

view of axon caliber determinants.

Introduction

The shape of a neuron's axons or dendrites is central to its function. A key characteristic

of axon and dendrite shape is their cross-sectional diameter, also known as caliber. Axon

caliber intrinsically influences the speed of action potential propagation (Kriz et al., 2000;

Sakaguchi et al., 1993) and may also play a role in cell health and structural stability (Dollé et

al., 2018; Perge et al., 2012).

Rohon-Beard (RB) neurons are a type of touch-sensing neuron that develops in early

embryonic zebrafish and project skin-innervating peripheral arbors with branched architecture

(Fig 1A) (Katz et al., 2021; Shorey et al., 2021; Wang et al., 2013). RB peripheral arbors are

afferent axons that detect somatosensory stimuli in the skin and relay messages to interneurons

via central axons in the spinal cord, thus promoting behavioral responses.

The significance of axon caliber in cell function has been described for about a century

(Hursh, 1939), and it is widely regarded as a cell-wide, static trait despite reports of variations in

axon or dendrite caliber over space and time. Axon caliber can vary along the length of axons

due to inherent biophysical properties (Griswold et al., 2024), myelination (Seidl, 2014), and

mechanical insult (Pan et al., 2024). Dendrite caliber is also known to taper across branch

points (Craig and Banker, 1994). Axon caliber has been observed to change over development

(Bomont, 2021; Seidl and Rubel, 2016), aging (Metzner et al., 2022), and in response to target

size (Voyvodic, 1989a, 1989b) or activity (Nabel et al., 2024; Sinclair et al., 2017). These

studies of caliber changes have primarily focused on changes to axon caliber over many days

and assume that caliber undergoes little to no change on the timescale of hours. However, the

cytoskeletal structures (Costa et al., 2020; Taylor et al., 2012; Wang et al., 2020) and cargoes

(Costa et al., 2020; Greenberg et al., 1990; Wang et al., 2020) that can impact axon caliber are

dynamic on the timescale of seconds to minutes, prompting us to ask how axon caliber and

morphology change on shorter timescales.

Here, we characterized axon caliber in zebrafish RB neurons. We found that each single

arbor, in vivo, can have a variety of axon calibers, varying within a segment and between sister

segments. Axon calibers were dynamic on the timescale of minutes, and these dynamics

changed over the developmental timescale of days. Finally, we found that the behavior of

epithelial cells in the skin impacts sensory axon caliber, highlighting the complexities of the in

vivo axon environment.

Materials and Methods

Fish care

Zebrafish (Danio rerio) adults were housed at 28.5°C with 13.5/10.5-hour light/dark cycles.

Embryos were raised at 28.5°C in water containing 0.06 g/L Instant Ocean salt mix (Spectrum

Brands, AA1-160P) and 0.05% methylene blue (ThermoFisher, 042771.AP).

Injection for sparse labeling

To sparsely label RB neurons for observation of development and axon caliber, adult fish

expressing Isl1[ss]:Gal4, UAS:dsRed (Sagasti et al., 2005) were crossed, and resulting embryos

were allowed to develop for approximately one hour after fertilization. When embryos reached

the 4-cell or 8-cell stage, a single cell in each embryo was injected with approximately 1-3 nL of

plasmid containing the Gal4 effector, UAS:egfp-caax, diluted to 10-15 ng/μL in water, with or

without phenol red (Sigma, P4758). After injection, embryos were raised as described above.

Mounting embryos for microscopy

Unless otherwise indicated, embryos were immobilized on coverslips in 1.2% low-melt agarose

dissolved in water (Fisher, BP165-25) and mounted with an equal volume of water containing

0.12 g/L Instant Ocean (Spectrum Brands, AA1-160P) and 0.4 mg/mL tricaine (MS-222,

Western Chemical Inc.).

Developmental time lapse imaging

Embryos injected for sparse labeling were sorted at approximately 20 hours post-fertilization

(hpf) to find animals with a single cell expressing EGFP-CAAX in the tail. Selected embryos

were immediately dechorionated using forceps and anesthetized in approximately 0.2 mg/mL

tricaine (MS-222, Western Chemical Inc.). Embryos were mounted as described above. A cut

was made in the agarose posterior to the tail using a fly pin probe to allow room for growth.

Embryos were imaged at 5-minute intervals on a ZEISS laser scanning microscope (LSM) 800

with a stage heater at 28°C. Magnification was increased as needed by changing objectives to

see branching. The most proximal branch point at approximately 28 hpf was identified and

visually tracked back to the point of formation within the time lapse for scoring.

Line scan and symmetry analysis

Embryos were injected for sparse labeling, as described above. Embryos with a distinct EGFP-

CAAX-positive neuron in the tail were dechorionated using forceps and anesthetized in

approximately 0.2 mg/mL tricaine (MS-222, Western Chemical Inc.). For this experiment,

embryos were immobilized in 1.1 – 1.2% low-melt agarose on No.1 coverslips (Epredia,

12460S) to improve available working distance, and water used in initial anesthetization was

added before sealing with a glass slide. The primary axon segment was identified as the one

that exited the spinal cord and entered the skin. The most proximal branch point was imaged on

a ZEISS LSM 880 with Airyscan deconvolution at 27 – 31 hpf. If two neurons or two arbors were

present in the same animal or cell, the more posterior one was scored. Line scan analysis was

performed using the Line and Plot Profile tools in ImageJ (Schindelin et al., 2012) to find

intensity values along a line perpendicular to the center axis of the axon. Axon caliber was

measured as the distance between the two peaks on the plot that corresponded to the plasma

membrane. For symmetry analysis, axons were measured 3, 4, and 5 µm away from the branch

point and averaged to find the caliber of the given segment. The caliber of secondary branches

was normalized by dividing by the caliber of the corresponding primary branch. We defined

symmetry as S2/S1, which is equivalent to the normalized S2 caliber divided by the normalized

S1 caliber.

Caliber dynamics time lapse imaging

Embryos were injected for sparse labeling, as described above. Around 24 hpf, embryos were

mounted as described above. The proximal region of the axon arbor was imaged on a ZEISS

LSM 980, heated to approximately 28°C, and processed with Airyscan Joint Deconvolution at a

maximum of five iterations. Time lapses with 5-minute intervals were started at 28 - 31 hpf and

acquired until the axon drifted out of the field of view, typically one to two hours. After imaging,

embryos were recovered from the mounting agarose using forceps and a probe and grown an

additional 24 hours under normal conditions (described above), with the addition of 0.2 mM PTU

(1-phenyl 2-thiourea, Sigma, P7629) in the water. At 52 - 55.5 hpf, embryos were mounted and

imaged in the same manner as the first day. For scoring, axon segment calibers were measured

3 µm from a branch point at every frame of the time lapse, and segments used were each

separated by at least one branch point. If the same segments could be identified on the second

day, caliber was scored at the same locations for every frame of the time lapse.

Dual color time lapse imaging of cell division

To observe axons near dividing cells, transgenic embryos expressing Isl1[ss]:lexA,

lexAop:mRuby-caax and tp63:Gal4VP16; UAS:egfp-plcδ-ph (Rasmussen et al., 2015) were

mounted as described above and imaged on a ZEISS LSM 980, heated to approximately 28°C,

and processed with Airyscan Joint Deconvolution at a maximum of five iterations. Basal skin

cells in a rounded state were identified by eye when embryos were 28 - 31 hpf and immediately

imaged by time lapse with 5-minute intervals, until daughter basal cells appeared flattened for at

least two time points. Axon caliber was measured by line scan analysis, as described above, at

1-μm intervals, excluding regions less than 3 μm from a branch point. Border to border length of

axons was measured using the Line tool in ImageJ (Schindelin et al., 2012). Orthogonal

projections were generated in ZEN Blue (ZEISS).

Plotting

All dot plots were created in R. Code can be found at github.com/katieching/CaliberAnalysis.

Histograms, dynamics, and pearling plots were created using Microsoft Excel.

Experimental design and statistical analysis

For descriptive analyses, an appropriate size for the data set was chosen in advance, and

images were collected in experimental batches until the predetermined size was exceeded. For

statistical comparisons, a small data set was initially collected, and a power analysis was

performed to determine the data set size needed to reach a p-value < 0.05. The data set was

collected and analyzed before performing final statistical analyses. Data were excluded as

necessary for technical reasons, namely image quality that was too poor for line scan analysis,

segments where branch points were closer together than the distance indicated for the

experiment (e.g. < 5 μm for symmetry analysis), images in which the primary branch caliber was

below the limit of resolution for the system (Fig 3, 4), and axons in which > 20% of

measurements at the 1 day post-fertilization (dpf) time point were below the limit of resolution

for the system (Fig 5, 6). Significance of differences in paired data was evaluated by paired

permutation test in R. Differences between measurements treated as populations were

evaluated by Mann-Whitney test. Analyses performed in R can be found at

github.com/katieching/CaliberAnalysis. The sex of animals included in the study was not

considered because experiments occurred prior to sex determination.

Results

Rohon-Beard (RB) neurons display a variety of calibers.

RB neurons are touch-sensing neurons that differentiate and become functional within

two days of development in embryonic zebrafish and other anamniotes (Katz et al., 2021). The

peripheral axons of RB neurons can be identified by transmission electron microscopy (TEM) as

membrane-bound ovals between the outermost (periderm) skin cells and lower (basal) skin cells

in cross section (O'Brien et al., 2012). Using existing TEM data sets (Faas et al., 2012; O'Brien

et al., 2012), we identified peripheral axons of RB neurons and trigeminal neurons, closely-

related sensory neurons that perform the same touch-sensing function in the head. By

measuring the short axis of these axon cross-sections, we found that calibers ranged from

approximately 0.084 to 1.459 µm. This wide range prompted us to ask if axon caliber varies

within each cell or only between cells, perhaps by sensory subtype (Gau et al., 2013; Palanca et

al., 2013; Tuttle et al., 2024).

To determine if axon caliber can vary within an axon arbor, we sought to measure

caliber at consistent locations across many intact cells. To enable absolute measurements of

axon caliber, we developed an imaging protocol that paired sparse labeling with super-

resolution microscopy. Using transient transgenesis, we generated embryos that expressed a

membrane-localized fluorescent protein (EGFP-CAAX) in a single RB neuron in the tail (Fig 1B).

We performed time lapse confocal microscopy starting shortly after RB neuron differentiation

(20 hours post-fertilization, hpf) and confirmed that branches of the peripheral axon arbor form

by growth cone bifurcation as well as collateral sprouting (Haynes et al., 2022; Sagasti et al.,

2005), resulting in a highly branched axon arbor within hours of skin entry. To identify a

consistent area of measurement, we observed the most proximal branch point in the peripheral

arbor at 27 - 30 hpf and found that it typically forms by collateral sprouting (10 out of 10 cells,

Fig 1C-C', Video 1). In these images, some branches of each arbor appeared to be thick while

others appeared to be thin (Fig 1D). This observation suggested that axon caliber can vary

within each cell and supports the use of this location and technique for studying axon caliber

variation.

Caliber varies along the length of an axon segment.

Previous studies have shown that axons are not cylindrical but vary in caliber along their

lengths (Bar-Ziv et al., 1999; Greenberg et al., 1990; Griswold et al., 2024; Pullarkat et al.,

2006). This phenomenon is sometimes referred to as pearling or varicose morphology. To

determine if pearling is observed in live animals, we labeled individual RB neurons and

measured axon caliber at 1-μm increments along the length of a segment, shortly after arbor

formation began (29 hpf, Fig 2A-B). Variations in axon caliber were visible along the length of all

resolvable axon segments, as has been described in mammalian neurons (Greenberg et al.,

1990; Griswold et al., 2024). In some instances, axons had distinct pearls (Fig 2A') and in

others, more subtle thick and thin stretches without distinct inflection points (Fig 2C). These

observations of lengthwise caliber variation confirm that fish and mammalian neurons share

similarities in local axon morphology in vivo.

S*ister axon branch calibers are largely uncorrelated with each other.*

Next, we asked how axon caliber varies between different axon segments within a single

cell. We labeled individual RB neurons and measured axon caliber near the branch point most

proximal to the cell body in 27 - 31 hpf embryos (Fig 3A). The axon segment coming from the

spinal cord was designated the primary axon (P), and the axon segments distal to the branch

point were designated secondary axons (sister branches designated S1 and S2, for the thicker

and thinner segments, respectively). Each segment was measured at three locations (3, 4, and

5 µm from the branchpoint), which were averaged to estimate the caliber of that axon segment.

By measuring these locations, we observed that axon segments of different calibers exist within

the same RB neuron and that secondary branches were typically thinner than the primary

branch from which they arose (Fig 3B, P = 0.43 µm, S1 = 0.32 µm, S2 = 0.20 µm). These data

confirmed that the average caliber of axons is not uniform across the cell.

Because sister segments, S1 and S2, often have different calibers, we sought to assess

the relationship between them. We considered two hypotheses: 1) despite variations, each

neuron has characteristically thick or thin calibers, and thus S1 and S2 have positively

correlated calibers, or 2) S1 and S2 compete for cellular material, and thus their calibers are

negatively correlated. To distinguish between these two hypotheses, we compared the S1 and

S2 caliber for each neuron. Direct comparison showed a weak relationship between S1 and S2

caliber (Fig 3C, R^2 = 0.13). This weak correlation suggests that neither hypothesis is strongly

supported.

In keeping with this interpretation, symmetry across the branch point was weakly

correlated with normalized S1 caliber (Fig 3D, R^2 = 0.04), which we would expect to be strongly

correlated if branches compete for cellular material to increase caliber. On the other hand,

symmetry is more strongly correlated with normalized S2 caliber (Fig 3E, R^2 = 0.54), suggesting

that symmetry is achieved by the thinner branch being thicker, unrelated to the caliber of the S1

branch.

RB axons taper.

For an RB neuron to function properly, the cell needs electrical signals to travel

efficiently. At the time points measured in Fig 3, the arbor was just a few hours old, functional,

and continuing to expand. Already, optimal RB neuron function of transmitting sensory signals

from the periphery requires efficient propagation of electrical signals from distal to proximal axon

segments (toward the soma). Although the peripheral arbor of RB neurons transmits these

afferent signals, as dendrites typically do, we refer to the peripheral arbor as an axon due to its

uniformly plus-end-out microtubule organization (Shorey et al., 2021). Tapering is often

considered a characteristic unique to dendrites (Craig and Banker, 1994; Jan and Jan, 2010),

though more recent analyses suggest that branched axons may taper as well (Desai-Chowdhry

et al., 2022).

To facilitate comparison of RB axon branching to other systems, we assessed tapering.

First, we calculated the cross-sectional area of each primary and secondary branch ($area =$

$\pi * (\frac{1}{2}caliber)^2$). We compared the primary cross-sectional area to the sum of the secondary

cross-sectional areas (Fig 4A). On average, the ratio of the combined secondary cross-sectional

areas to the primary cross-sectional area was less than 1 (mean of $\frac{area_{S1} + area_{S2}}{area_P} = 0.888$, Fig

4B), meaning that the RB peripheral arbor tapers in area.

Tapering can influence the health and function of a neuron in many ways, some of which

scale with area, and some of which scale with radius (Craig and Banker, 1994; Desai-Chowdhry

et al., 2022). Hence, we also compared the radius scaling ratio between secondary branches

and their corresponding primary branches ($\frac{r_S}{r_P}$, Fig 4C). The average ratio for the thicker branch

was much larger than for the thinner sister branch (mean $\frac{S1}{P} = 0.77 \pm 0.033\ SEM,$ and mean

$\frac{S2}{P} = 0.49 \pm 0.030\ SEM$, Fig 4D). In evaluating a branched system, some scaling rules hold for

the population average, even when branches are asymmetric (Brummer et al., 2017). Hence, to

evaluate the function of the arbor more generally, we combined the measurements for the two

branches and found that the average ratio (mean $\frac{S}{P} = 0.629 \pm 0.0282\ SEM$) was similar to what

is predicted for a system that optimizes for power dissipation rather than time delay (Desai-

Chowdhry et al., 2022).

Here, we refer to the RB peripheral arbor as an axon due to its plus-end-out microtubule

organization (Shorey et al., 2021). Therefore, our observations demonstrate that it is possible

for neuronal processes with an axon-like cytoskeletal organization to taper.

Axon caliber is highly dynamic.

Having documented variability in axon caliber across locations within each RB neuron,

we wondered how calibers change with time. To investigate caliber dynamics, we performed

time-lapse imaging in the proximal region of the tail-innervating axon arbors at 28 - 31 hpf

(Video 2). Axon caliber changed in several qualitatively different ways over time. We

categorized these dynamic behaviors into four categories (Video 3): (1) a traveling pearl,

instances of bubble-like regions moving along an axon (Fig 5A), (2) focal inflation and deflation,

in which a pearl appears and disappears over the course of multiple time points (Fig 5B), (3)

segment widening and narrowing, in which an entire segment thickens and thins between time

points (Fig 5C), and (4) a constriction point that appears and disappears, sometimes repeatedly

(Fig 5D). This variety of dynamic behaviors suggests that several distinct processes likely

contribute to axon caliber dynamics.

Axon caliber is known to be altered by changes in target cells (Nabel et al., 2024;

Sinclair et al., 2017; Voyvodic, 1989a) or myelination (Ciocanel et al., 2020; Seidl, 2014; Walker

et al., 2019), but these developmental processes occur on different timescales from the

dynamics we observed with intervals of 5 minutes (Fig 5A-D). To assess caliber dynamics

quantitatively, we selected several locations within the proximal region of each neuron, each 3

μm away from a branch point, and measured axon caliber at that location over the course of

imaging (Fig 5E). Axon caliber was highly dynamic (Fig 5F), with standard deviations ranging

from 0.045 μm to 0.095 μm, and could vary with a range of up to four-fold at the same location

in less than two hours. Hence, axon caliber and local morphology may be more dynamic than

has been widely appreciated.

Next, we asked if thin or thick axon segments are more dynamic. First, we assessed

absolute dynamicity by comparing average caliber to the standard deviation (SD) or

measurements observed during the time lapse (Fig 5G). We found that SD and axon caliber

correlated positively, suggesting that thicker axons experienced larger or more fluctuations in

caliber. A possible caveat for our measurements is that the lower dynamicity could be due to

thin axons spending more time below the limit of resolution. To determine if this was the case,

we also assessed relative dynamicity. We calculated relative standard deviation (RSD) for each

location by dividing the standard deviation by the average caliber across the movie. Comparing

dynamicity across segments revealed that caliber and RSD were negatively correlated, meaning

that thick axons were less dynamic than thin axons (Fig 5H). If differences in absolute

dynamicity were due to resolution limitations, we would expect to see that relative dynamicity of

the thinnest axons is also low. Instead, thin axons had high relative dynamicity, indicating that, if

anything, we may have underestimated the dynamicity of thin axons. Together, these results

suggest that thick axons have more or larger absolute fluctuations in caliber, but the fluctuations

experienced by thin axons are more dramatic when compared to their average caliber.

Caliber dynamicity changes over development.

Because we observed caliber to be highly dynamic early in the life and function of RB

neurons, we sought to determine if caliber continues to be dynamic later in development. To do

this, we recovered embryos after imaging at 28 - 31 hpf and allowed them to continue

developing for an additional 24 hours. These fish were re-mounted at 52 - 55.5 hpf and imaged

by time-lapse microscopy again. Although all cells had changed shape and axon arbors had

grown, often the locations imaged on the first day could be identified, imaged, and measured

again on the second day (Fig 6A-B). We found that all resolvable axons continued to be

dynamic. However, dynamicity measured as %RSD decreased from the first to the second day

in most axon segments (10 out of 11 axon segments decreased. Mean %RSD = 28% at 1 dpf

and 20% at 2 dpf, p = 0.036, Fig 6C). Because some axon segments got thicker (n = 6 out of

11) and some got thinner (n = 5 out of 11), this decline in relative dynamicity is not simply a

product of axons becoming universally thicker. Instead, this dynamicity likely reflects

developmental changes that occur in the cell and tissue.

Cellular microenvironment can impact axon caliber.

Based on the qualitative variety of caliber dynamics we observed on the minutes

timescale (Fig 5A-D), we hypothesized that there are multiple contributors to caliber dynamics.

Cell-intrinsic effectors that may cause axon caliber to be dynamic on short timescales, such as

cargo transport and contraction of the membrane periodic skeleton, have been described by

others (Costa et al., 2020; Wang et al., 2020). However, another contributor is the cellular

microenvironment. Mechanical force at the soma has been shown to influence axon and

dendrite morphology (Pan et al., 2024). We hypothesized that, in RB peripheral axons, caliber

changes could be influenced locally, including by contact with surrounding epithelial cells.

Because RB axons are embedded in actively developing skin, they are subject to local

deformation caused by morphological changes in surrounding epithelial cells. Specifically, RB

peripheral axons grow between two epithelial cell layers, the periderm and basal cells, each of

which undergoes rapid expansion at these developmental stages (Fig 1A) (O'Brien et al., 2012).

To determine if changes to the shape or adhesion of adjacent epithelial cells can impact

the caliber of axons, we performed simultaneous time lapse imaging of axons and the

underlying basal cells starting at 28 - 31 hpf (Video 4). We imaged basal cells using a marker of

lipid microdomains under the control of the ΔNp63 promoter (Rasmussen et al., 2015; Rosa et

al., 2023). One of the most dramatic and frequent perturbations to the RB axon environment

occurs when basal cells round up for mitosis, during which the lipid reporter clearly highlights

rounded cells against the field of surrounding, flat basal cells (Fig 7A). We measured caliber at

1-μm intervals along the length of the axon, from a flat neighbor cell, onto the rounded cell, and

onto the next flat neighbor cell (Fig 7B). We found that segments of axon on the rounded cell

were significantly thicker than segments of the same axon on the non-dividing neighbor cells

(caliber = 0.21 µm on rounded cell, 0.16 and 0.14 µm on non-dividing neighbors, p-value =

0.025 and 0.00035, respectively, n = 114 measurement locations, N = 4 axons/fish, Fig 7C).

This difference contrasts with the two non-dividing neighbor groups, which did not have

significantly different calibers, as expected (p-value = 0.059). Although this observation does not

distinguish between mechanical or signaling interactions as the driving factor for caliber

changes, these data align with a model in which dynamic properties of the dividing cell may

impact local axon caliber.

Forces from surrounding cells may impact axon morphology.

One reason why axons might be thicker on rounded cells could be due to changes in

tension (Griswold et al., 2024; Pan et al., 2024). As a basal cell enters mitosis, it pulls its

borders inward, becoming shorter in planar distance, and rounds up, creating a greater vertical

distance (Fig 7D). We used images with the clearest orthogonal projections to obtain estimates

of how the dimensions of a dividing basal cell differs from its non-dividing neighbors. Using

these estimates, we calculated theoretical changes in axon length as the cell rounds up (Fig

7E). To simplify these calculations, we assumed that the axon is fixed at the cell borders and

runs in a straight path across the basal cell surface. We found that the rounding of basal cells

resulted in a shorter axon path during division than when cells were flat (path length when round

≈ 0.78 * path length when flat). Assuming that axons are adhered to the non-dividing

neighboring cells, these estimates are consistent with the axon being under lower tension on the

rounded cell than on a non-dividing neighbor cell.

Based on our observations of rounded cells and our estimates of length changes, we

expected that axon caliber would decline as the daughter basal cells flatten. To test this

prediction, we performed time-lapse imaging with 5-minute intervals on the rounded cells,

including those in Fig 7C, to see if axon caliber changes as the rounded cell divides and flattens

into two daughter cells. For each time point, we measured the distance between the two points

where the axon crossed the border of the dividing cell or its daughter cells (Fig 8A). The two

time points between which the change in length was greatest were selected, and the caliber of

the axon on the rounded cell or its daughters was measured at 1-µm increments for those two

time points (Fig 8B-C). We paired measurements by location to assess if there was a significant

drop in axon caliber between these two time points. Although axon caliber declined from the

round to flat time point, on average, this subtle decrease did not reach statistical significance

(caliber = 0.28 µm when the basal cell was round and 0.25 µm when the basal cell was flat, p-

value = 0.17, n = 54 measurements, N = 7 axons/fish, Fig 8D). This contrasts with the portions

of the axon that were on non-dividing neighbor cells, where the difference could have easily

occurred by random chance (p-value = 0.94, n = 37 measurements). This result suggests that

the decrease in average axon caliber as the underlying basal cell flattened is small and likely

not meaningful.

Despite the lack of change in average caliber, visual assessment of the videos appeared

to show a change in morphology during basal cell flattening (Fig 7A, 8C). Tension and

mechanical force may be an important parameter in pearling (Bar-Ziv et al., 1999; Griswold et

al., 2024; Pan et al., 2024; Pullarkat et al., 2006), so we wondered if thick regions (i.e. pearls)

and thin regions (i.e. connectors) undergo different changes during basal cell division.

To assess morphological changes to axons during basal cell division, we calculated

pearling as the standard deviation of measurements taken along the length of each axon for the

rounded time point and for the flat time point. By pairing values from each movie, we found that

one subset of axons became more pearled (increased SD) and the rest became less pearled

(decreased SD), resulting in two groups of cellular responses to flattening (Fig 8E). This

heterogeneity might suggest that either 1) different responses arise from heterogeneity among

the RB neuron population, or 2) our 5-minute intervals do not provide sufficient time resolution

to capture cellular responses to tension, which may occur in less than 10 minutes (Sinha et al.,

2011; Sitarska and Diz-Muñoz, 2020). Regardless of the reason, these results demonstrate that

axon morphology changes with basal cell flattening, though additional complexities remain to be

explored.

Discussion

Tapered architecture in branched sensory axons

We have found that the calibers of axon segments in the same RB neuron can be largely

independent of one another, highly dynamic, and likely shaped by multiple determinants,

including contact with epithelial cells. Variation in axon caliber had previously been observed

along short stretches of fixed axons in rodent neurons (Greenberg et al., 1990; Griswold et al.,

2024), in response to developmental changes to the neuron's environment and signaling (Bin et

al., 2024; Ford et al., 2015; Kumar et al., 2005; Metzner et al., 2022; Nabel et al., 2024;

Sánchez et al., 1996; Seidl, 2014; Seidl et al., 2010; Seidl and Rubel, 2016; Sinclair et al.,

2017), and with forces on millimeters-length scales (Pan et al., 2024). Our results expand upon

these findings by characterizing caliber variations in vivo across a branched axon, with fine

timescale, and with local deformations caused by epithelial cell division.

The cutaneous axon endings of touch-sensing neurons, including trigeminal, dorsal root

ganglion (DRG), and RB neurons, share characteristics with both dendrites and axons. Like

dendrites, they detect stimuli from the periphery and relay signals towards the cell body, but, like

axons, their microtubules are oriented plus-end-out (Shorey et al., 2021). We found that the

branched axon arbors of RB neurons taper across the most proximal branch point, becoming

narrower in caliber and cross-sectional area further from the soma (Fig 4). This feature is often

regarded as a characteristic of dendrites, not of axons (Craig and Banker, 1994; Jan and Jan,

2010). Mechanisms of signal relay from periphery to soma in RB neurons have not been fully

elucidated. If RB neurons fire action potentials, then their tapering (Fig 5) would favor afferent

action potential conduction, as signals travel from the periphery toward the soma. In measuring

the radius scaling ratio, we found it to be consistent with neurons optimized for power

dissipation, which aligns well with a more graded, dendrite-like signal transduction (Desai-

Chowdhry et al., 2022). Together, our findings are consistent with a model in which tapering

may be most aligned with the direction of functional signal propagation rather than microtubule

polarity. Whether our findings hold true for other axons, especially afferent axons, remains to be

explored.

In comparing sister axon branches, we hypothesized that sister branches are related to

each other, resulting from either cell-wide effects or a competition for cellular materials. Instead,

we found that their calibers are largely uncorrelated with each other, despite being in close

proximity and sharing a cytoplasm. Interestingly, we observed varying degrees of symmetry,

ranging from pairs of sister branches with the same caliber to highly asymmetrical pairs, where

one secondary branch is much thicker than the other. One reasonable hypothesis to explain this

observation could be that branches of different symmetry form by different mechanisms. For

example, highly asymmetric branches might have formed by collateral sprouting and symmetric

branches by growth bifurcation. However, we found a full spectrum of symmetry values, not a

bimodal distribution, arguing against this hypothesis (Fig 3D-E). Moreover, though bifurcations

are common in RB arbor outgrowth (Haynes et al., 2022) the first branch point often, if not

always, forms as a collateral (Fig 1C). Thus, collateral sprouting can give rise to segments of a

variety of calibers within a few hours of formation, highlighting the malleability of axon caliber.

Mechanisms of caliber variation over time and space

461   We found that caliber dynamicity changed over development (Fig 6). Many

developmental changes could account for a shift in dynamicity, including changes within the cell

itself, such as cargo transport (Costa et al., 2020; Wang et al., 2020), or changes to the

extracellular environment, such as axon ensheathment (Jiang et al., 2019; O'Brien et al., 2012;

Rosa et al., 2023). Time lapse imaging of axon caliber also revealed multiple qualitatively

different types of dynamics (Fig 5A-D), which we refer to as traveling pearls, focal inflation and

deflation, segment widening and narrowing, and constriction points. This variety suggests that

there are likely multiple contributors to variations in axon caliber, including both cell-intrinsic and

cell-extrinsic mechanisms.

470   Cell-intrinsic determinants of axon caliber, such as the cytoskeleton, have been studied

to varying degrees. One of the best-known caliber determinants is neurofilaments, a type of

intermediate filament specific to the nervous system. Increased neurofilament abundance

increases axon caliber in mammalian and bird neurons (Elder et al., 1998; Eyer and Peterson,

1994; Kriz et al., 2000; Ohara et al., 1993; Sainio et al., 2021; Sakaguchi et al., 1993; Zhu et al.,

1997). Neurofilaments are also highly dynamic. Differences in transport, differential

modifications, and dynamics (e.g. folding, severing, annealing) of neurofilaments are all possible

ways in which these filaments might contribute to both local regulation and axon caliber

dynamics within a cell (Boyer et al., 2022; Çolakoğlu and Brown, 2009; Uchida et al., 2023).

479   Another cytoskeletal candidate for cell-intrinsic effectors of caliber variation is the more

recently discovered membrane periodic skeleton (MPS), regularly-spaced actin rings connected

by spectrin tetramers that run the length of axons and dendrites (He et al., 2016; Xu et al.,

2013). These rings expand when large cargo such as mitochondria and lysosomes pass

through, and altering myosin contractility changes caliber and retrograde trafficking (Costa et al.,

2020; Leite et al., 2016; Wang et al., 2020). Observations of cargo transport through the MPS

(Costa et al., 2020; Wang et al., 2020), as well as fixed observations that large cargo is often

found within pearls (Greenberg et al., 1990), suggest that intracellular cargo transport along

microtubules may also contribute to axon caliber dynamics. Work by Griswold et al.

demonstrated that myosin contractility can impact pearling (Griswold et al., 2024). Furthermore,

the MPS and the propensity of neurons to bead have been found to be protective against low-

level mechanical forces (Krieg et al., 2017; Pan et al., 2024), highlighting MPS contractility as

another potential source of caliber variations.

Other intracellular factors, such as importin beta (Bin 2024), PI3K/AKT signaling (Kumar

2005) and membrane composition (Griswold 2024) also regulate axon caliber. Although some of

these factors likely influence caliber by regulating the cytoskeleton (Bin 2024), membrane

composition is a non-cytoskeletal, cell-intrinsic candidate for causing caliber variation. Griswold

et al. found that membrane fluidity can change which size and shape of pearls are most

energetically favorable, thereby changing axon morphology (Griswold et al., 2024). Thus, given

the dynamic nature of caliber in vivo, changes to membrane composition could also change

caliber dynamics.

Damage can also influence axon morphology. Markedly pearled or varicose

morphologies, sometimes referred to as beading or swelling, often form in damaged axons

(Datar et al., 2019; Kerschensteiner et al., 2005; Shao et al., 2020; Williams et al., 2014).

Indeed, damaged trigeminal and RB axons, themselves, become dramatically more beaded just

before degeneration (Martin et al., 2010). Mechanical forces below the level necessary to break

axons have also been found to impact axon and dendrite pearling (Pan et al., 2024). Our

observations of variation in caliber in this study were made in anesthetized, healthy embryos

and are unlikely to result from axon damage, as neurons remained intact, even after time lapse

microscopy. Similarity between these phenomena in healthy and damaged axons may reflect

related processes if the healthy malleability of axon caliber is pushed to extremes upon insult.

Although much attention has focused on potential intrinsic determinants for axon caliber

variation, we considered the possibility that extrinsic influences from surrounding cells might

affect axon caliber locally. Cell-extrinsic factors that influence axon caliber include target size

(Voyvodic, 1989a) and distance (Seidl et al., 2010), neuronal function (Nabel et al., 2024;

Sinclair et al., 2017), and mechanical forces at millimeters-length scale (Pan et al., 2024). Each

of these factors likely impacts the entire cell and is, therefore, unlikely to regulate caliber

variation locally.

Our study focused on smaller deformations that occur on smaller length scales and

timescales. Although myelination is thought to influence local axon caliber (Rydmark, 1981;

Seidl, 2014; Swärd et al., 1995), questions remain about the direction and specific nature of this

intercellular relationship (Bin et al., 2025; Friede, 1972; Sánchez et al., 1996). RB axon arbors

are embedded in a developing epidermis, which undergoes growth and morphogenetic

changes. RB axons span tens or even hundreds of micrometers in length and branch to cover

large areas of the skin. Hence, each segment of an axon contacts different basal cells, which

may each impact the local axon caliber independently.

Basal cells can undergo many morphological changes to alter the axon's environment.

One important change is ensheathment (O'Brien et al., 2012) (Fig 1A), which increases

between 1 and 2 dpf (Jiang et al., 2019; Rosa et al., 2023), possibly contributing to differences

in dynamicity between these two developmental time points (Fig 6C). The idea that

ensheathment may alter axon caliber is further consistent with the observation that myelinated

portions of an axon are thicker than the nodes of Ranvier (Rydmark, 1981; Swärd et al., 1995).

Another morphological change in basal cells is rounding for division, which we found

changes axon caliber and morphology (Fig 7,8). We hypothesize that changes in tension

caused by the movement of cells adjacent to axons are candidates for cell-extrinsic regulators

of axon caliber. Division events likely vary in frequency throughout embryonic development,

which may further contribute to developmental differences in caliber dynamicity (Fig 6C).

Additionally, axons may experience end-to-end stretch as embryos lengthen. Finally, although

our studies were in embryonic zebrafish, when the environment surrounding these neurons is a

rapidly expanding bilayered epidermis, dynamics are unlikely to cease in adulthood, since the

stratified epidermis in both fish and mammals is constantly renewing as basal cells differentiate

into higher strata. How caliber variations and dynamics differ in this stratified system remains to

be explored. In conjunction with findings that myelination influences axon caliber (Ciocanel et

al., 2020; Rydmark, 1981; Seidl, 2014; Swärd et al., 1995; Walker et al., 2019), our results

support the model that lengthwise variation, local variations in average caliber, and caliber

dynamics may be influenced, in part, by the behavior of surrounding cells.

546
547
Figure 1: Axon branch development and caliber in sparsely-labeled Rohon-Beard (RB)
neurons
(A) Diagram depicting an RB neuron in an embryonic zebrafish. Light grey squares represent
periderm (outermost) epithelial cells. Dark grey triangles represent basal epithelial cells, which
ensheath some RB axons around 2 - 3 dpf. **(B)** Example of embryonic zebrafish tail with sparse
labeling showing one RB neuron expressing EGFP-CAAX among many RB neurons expressing
cytoplasmic DsRed. Scale bar: 50 μm. **(C)** Illustrative images taken from time lapse of RB
neuron development. Proximal branch point formed by collateral sprouting in 10 out of 10 time
lapses. Arrow: formation of main branch in peripheral axon arbor. 26.42 hpf panel shows
multiple new branches that formed within minutes of each other. Open arrowhead: growth cone
bifurcation event. Solid arrowheads: collateral sprouting events. Scale bars: 10 μm. **(C')**
Proximal region of the axon arbor shown in B. Magenta pseudo-colored axon shows the main
branch, which branched by growth cone bifurcation. Green pseudo-colored axons show
branches that formed by subsequent collateral sprouting. Scale bar: 5 μm. **(D)** Method used for
measuring caliber. RB neurons expressing EGFP-CAAX were measured by line scan (green
line). Caliber was scored as distance between intensity peaks, as illustrated in example plot. P:
primary branch, which extends from spinal cord to skin, S1: thicker secondary branch, S2:
thinner secondary branch. Scale bar: 2 μm.

Figure 2: Lengthwise variation in axon caliber
(A) Image of an RB axon arbor at 29.00 hpf. Blue box shows location of inset in A'. Green box shows location of inset in B. Scale bar: 5 μm. **(A')** Inset showing axon branch with pearled morphology. Arrowheads: Thicker regions of axon, resembling "pearls" previously described in mammalian neurons. Scale bar: 2 μm. **(B)** Long axon segment measured by line scan at 1-μm increments. Scale bar: 2 μm. **(C)** Plot of caliber showing variation along the length of axon segment in panel B.

576
577
Figure 3: Comparison of axon segments within each arbor
(A) Example image of proximal RB axon arbor. Arrowheads indicate approximate location of line
scans for average caliber measurement. P: primary branch, S1: thicker secondary branch, S2:
thinner secondary branch. Scale bar: 5 μm. **(B)** Plot of average axon calibers at the proximal
branch point. Grey lines connect data points from the same neuron. Horizontal lines: mean
values with colors corresponding to branch categories shown in panel A. n = 31 neurons. **(C)**
Plot comparing caliber of sister axon segments shown in panel A. **(D)** Plot of branch symmetry
versus normalized S1 caliber (S1/P). **(E)** Plot of branch symmetry versus normalized S2 caliber
(S2/P). Blue lines in C, D, and E show linear regression.
587

588
589
Figure 4: Analysis of tapering across the most proximal branch point
(A) Diagram of tapering analysis in panel B. Combined cross-sectional area of secondaries (S1
and S2) is divided by the cross-sectional area of the primary (P) to evaluate tapering. **(B)**
Histogram of cross-sectional area ratios for data shown in Figure 1. **(C)** Diagram of tapering
analysis in panel D. Caliber of each secondary (S1 or S2) is divided by the caliber of the primary
(P) to estimate the impact of tapering on action potential conduction velocity. **(D)** Plot of caliber
ratios for data shown in Figure 1. Grey lines connect data points from the same neuron. In
panels B and D, dotted black lines mark reference value 1.0 (no tapering), and magenta lines
mark mean values.
599

Figure 5: Qualitative and quantitative analysis of caliber dynamics
(A) Example of possible traveling pearl. Short distance traveled allowed for acquisition despite 5-minute intervals. Arrowhead: initial location of pearl. **(B)** Example of focal inflation and deflation event. Puncta containing EGFP-CAAX accumulate during inflation and are seen exiting during deflation. Arrowhead: location of event. **(C)** Example of segment widening and narrowing. Arrowhead pairs denote width at t = 5 minutes. **(D)** Example of constriction point. Arrowhead: location of event. In panels A-D, scale bars: 2 µm. **(E)** Image of axon arbor in the proximal region. Different segments are labeled from proximal to distal: i-iv. Scale bar: 5 µm. **(F)** Plot of axon caliber dynamics, n = 11 axon segments, N = 5 fish. Blue lines: axon segments shown in panel E. Grey lines: all other axon segments measured. **(G)** Plot of mean caliber versus absolute dynamicity (SD) for all axons shown in panel F. Mean caliber is the average at the given location across all time points. **(H)** Plot of mean caliber versus relative dynamicity (%RSD

613   = (SD / mean caliber) * 100) for all axons shown in panel F. In panels G and H, blue lines: linear
614   regression.
615

Figure 6: Tracking caliber dynamics over development

(A) Experimental workflow with example images (same neuron as Figure 2). Arrowheads: locations for line scan measurements. Each location is 3 µm from a branch point and separated from other locations by at least one branch point. Scale bars: 5 µm. **(B)** Representative example plots of caliber dynamics for the same segment at 1 dpf and 2 dpf. **(C)** Plot of relative dynamicity (%RSD) at 1 dpf and 2 dpf (same neurons as Figure 5) n = 11 neurons, N = 5 fish. Grey lines connect data points from the same neuron. Magenta lines: mean values. *p < 0.05.

625
626
Figure 7: Axon caliber on dividing basal epithelial cells
(A) Representative images of time lapse showing an RB axon adjacent to a dividing basal cell.
Basal cells express EGFP-PLCδ-PH. RB neurons express mRuby-CAAX. Scale bar: 10 μm. **(B)**
Diagram of an axon on a dividing basal cell that is rounded while undergoing mitosis. Green
line: axon segment in contact with dividing basal cell. Blue lines: axon segments on neighboring,
non-dividing basal cells. **(C)** Plot of axon caliber measurements grouped by location, n = 114
measurement locations, N = 4 axons/fish. Green dots: caliber of axon on a dividing basal cell.
Blue dots: caliber of axon on neighboring cells to one side (neighbor1) or the other (neighbor2)
of the dividing basal cell. Black triangles: mean value. *p-val < 0.05. **(D)** Orthogonal projection of
a dividing basal cell, which expresses EGFP-PLCδ-PH and is highlighted against non-dividing
basal cells due to a variegated expression pattern. An RB axon is in contact with its apical
surface, which is rounded upward. Vertical scale bar: 10 μm. **(E)** Diagram comparing axon path
length on a rounded, dividing basal cell to its flattened state as an approximation of how basal
cell shape changes may impact tension on the axon. Magenta: approximated, theoretical axon
length, estimated as the longest distance across the apical surface in an orthogonal projection.
Approximate arc length was calculated based on height and length estimates in orthogonal
projections. ℓ: original length of basal cell, h: original height of basal cell.

644
645
Figure 8: Changes to axon morphology as new basal epithelial cells flatten
(A) Diagram of how border-to-border length was measured for an axon on a dividing basal cell.
(B) Example plot of changes in border-to-border axon length during time lapse imaging for axon
shown in Fig 7A. Round and flat time points are defined as sequential points (5-min. interval)
during which border-to-border length increases the most. **(C)** Images from time points
highlighted in panel B. Arrowheads: five measurement locations. Scale bars: 5 µm. **(D)** Plot of
axon caliber for all locations that remained on the dividing basal cell or its daughter cells, n = 54
measurement locations, N = 7 axons/fish. Grey lines connect data points from the same
location. Black lines: mean values. **(E)** Plot of pearling, defined as SD across all locations, to
assess morphology change for axons shown in panel D, N = 7 axons/fish. Grey lines connect
data points from the same axon.
657
658

Video Legends

Video 1: Time-lapse microscopy of axon branch formation in RB neurons.
Confocal time lapse microscopy of RB neuron development in the embryonic zebrafish tail. The
main branch of the peripheral axon arbor grows out of the developing spinal cord and into the
skin. Shortly thereafter, many branches form by growth cone bifurcation and collateral sprouting.
Pseudocoloring in the final frames highlights proximal arbor, and arrows indicate branch points
that formed by collateral sprouting events at the times indicated. Scale bars: 5 µm.
666
667   **Video 2: Time-lapse microscopy of peripheral RB axon.**
668   Super-resolution time-lapse microscopy of the peripheral axon arbor showing the most proximal
669   branch points. Frequent and rapid changes in axon caliber occur within the approximately 1
670   hour captured prior to drift. Time intervals: 5 minutes. Scale bar: 5 µm.
671
672   **Video 3: Types of caliber fluctuations in RB axons.**
673   Super-resolution time-lapse microscopy of peripheral axon arbors showing examples of
674   qualitatively different fluctuations in axon caliber. Time intervals: 5 minutes. Scale bars: 2 µm.
675
676   **Video 4: Time-lapse microscopy of RB axon in contact with mitotic basal keratinocyte.**
677   Super-resolution time-lapse microscopy of an RB axon segment in contact with a skin cell that is
678   undergoing mitosis. Morphological changes in both cells were visualized via membrane
679   fluorescent membrane tags. Magenta: RB neuron labeled with mRuby-caax. Green: Basal
680   keratinocyte labeled with egfp-plcδ-ph. Time interval: 5 minutes. Scale bar: 5 µm.
681
682

References

Bar-Ziv R, Tlusty T, Moses E, Safran SA, Bershadsky A (1999) Pearling in cells: a clue to

understanding cell shape. Proc Natl Acad Sci U S A 96:10140–10145.

Bin JM, Emberley K, Buscham TJ, Eichel-Vogel MA, Doan RA, Steyer AM, Nolan MF, Möbius

W, Monk KR, Werner HB, Emery B, Lyons DA (2025) Developmental axon diameter growth

of central nervous system axons does not depend on ensheathment or myelination by

oligodendrocytes. bioRxivorg.

Bin JM, Suminaite D, Benito-Kwiecinski SK, Kegel L, Rubio-Brotons M, Early JJ, Soong D,

Livesey MR, Poole RJ, Lyons DA (2024) Importin 13-dependent axon diameter growth

regulates conduction speeds along myelinated CNS axons. Nat Commun 15:1790.

Bomont P (2021) The dazzling rise of neurofilaments: Physiological functions and roles as

biomarkers. Curr Opin Cell Biol 68:181–191.

Boyer NP, Julien J-P, Jung P, Brown A (2022) Neurofilament Transport Is Bidirectional In Vivo.

eNeuro 9.

Brummer AB, Savage VM, Enquist BJ (2017) A general model for metabolic scaling in self-

similar asymmetric networks. PLoS Comput Biol 13:e1005394.

Ciocanel M-V, Jung P, Brown A (2020) A mechanism for neurofilament transport acceleration

through nodes of Ranvier. Mol Biol Cell 31:640–654.

Çolakoğlu G, Brown A (2009) Intermediate filaments exchange subunits along their length and

elongate by end-to-end annealing. The Journal of Cell Biology.

Costa AR, Sousa SC, Pinto-Costa R, Mateus JC, Lopes CD, Costa AC, Rosa D, Machado D,

Pajuelo L, Wang X, Zhou F-Q, Pereira AJ, Sampaio P, Rubinstein BY, Mendes Pinto I,

Lampe M, Aguiar P, Sousa MM (2020) The membrane periodic skeleton is an actomyosin

network that regulates axonal diameter and conduction. Elife 9.

Craig AM, Banker G (1994) Neuronal polarity. Annu Rev Neurosci 17:267–310.

Datar A, Ameeramja J, Bhat A, Srivastava R, Mishra A, Bernal R, Prost J, Callan-Jones A,

Pullarkat PA (2019) The roles of microtubules and membrane tension in axonal beading,

retraction, and atrophy. Biophys J 117:880–891.

Desai-Chowdhry P, Brummer AB, Savage VM (2022) How axon and dendrite branching are

guided by time, energy, and spatial constraints. Sci Rep 12:20810.

Dollé J-P, Jaye A, Anderson SA, Ahmadzadeh H, Shenoy VB, Smith DH (2018) Newfound sex

differences in axonal structure underlie differential outcomes from in vitro traumatic axonal

injury. Exp Neurol 300:121–134.

Elder GA, Friedrich VL Jr, Kang C, Bosco P, Gourov A, Tu PH, Zhang B, Lee VM, Lazzarini RA

(1998) Requirement of heavy neurofilament subunit in the development of axons with large

calibers. J Cell Biol 143:195–205.

Eyer J, Peterson A (1994) Neurofilament-deficient axons and perikaryal aggregates in viable

transgenic mice expressing a neurofilament-beta-galactosidase fusion protein. Neuron

12:389–405.

Faas FGA, Avramut MC, van den Berg BM, Mommaas AM, Koster AJ, Ravelli RBG (2012)

Virtual nanoscopy: generation of ultra-large high resolution electron microscopy maps. J

Cell Biol 198:457–469.

Ford MC, Alexandrova O, Cossell L, Stange-Marten A, Sinclair J, Kopp-Scheinpflug C, Pecka

M, Attwell D, Grothe B (2015) Tuning of Ranvier node and internode properties in

myelinated axons to adjust action potential timing. Nat Commun 6:8073.

Friede RL (1972) Control of myelin formation by axon caliber (with a model of the control

mechanism). J Comp Neurol 144:233–252.

Gau P, Poon J, Ufret-Vincenty C, Snelson CD, Gordon SE, Raible DW, Dhaka A (2013) The

zebrafish ortholog of TRPV1 is required for heat-induced locomotion. J Neurosci 33:5249–

5260.

Greenberg MM, Leitao C, Trogadis J, Stevens JK (1990) Irregular geometries in normal

unmyelinated axons: a 3D serial EM analysis. J Neurocytol 19:978–988.

Griswold JM, Bonilla-Quintana M, Pepper R, Lee CT, Raychaudhuri S, Ma S, Gan Q, Syed S, Zhu C, Bell M, Suga M, Yamaguchi Y, Chéreau R, Valentin Nägerl U, Knott G, Rangamani P, Watanabe S (2024) Membrane mechanics dictate axonal pearls-on-a-string morphology and function. Nat Neurosci.

Haynes EM, Burnett KH, He J, Jean-Pierre MW, Jarzyna M, Eliceiri KW, Huisken J, Halloran MC (2022) KLC4 shapes axon arbors during development and mediates adult behavior. Elife 11.

He J, Zhou R, Wu Z, Carrasco MA, Kurshan PT, Farley JE, Simon DJ, Wang G, Han B, Hao J, Heller E, Freeman MR, Shen K, Maniatis T, Tessier-Lavigne M, Zhuang X (2016) Prevalent presence of periodic actin-spectrin-based membrane skeleton in a broad range of neuronal cell types and animal species. Proc Natl Acad Sci U S A 113:6029–6034.

Hursh JB (1939) Conduction velocity and diameter of nerve fibers. Am J Physiol 127:131–139.

Jan Y-N, Jan LY (2010) Branching out: mechanisms of dendritic arborization. Nat Rev Neurosci 11:316–328.

Jiang N, Rasmussen JP, Clanton JA, Rosenberg MF, Luedke KP, Cronan MR, Parker ED, Kim H-J, Vaughan JC, Sagasti A, Parrish JZ (2019) A conserved morphogenetic mechanism for epidermal ensheathment of nociceptive sensory neurites. Elife 8.

Katz HR, Menelaou E, Hale ME (2021) Morphological and physiological properties of Rohon-Beard neurons along the zebrafish spinal cord. J Comp Neurol 529:1499–1515.

Kerschensteiner M, Schwab ME, Lichtman JW, Misgeld T (2005) In vivo imaging of axonal degeneration and regeneration in the injured spinal cord. Nat Med 11:572–577.

Krieg M, Stühmer J, Cueva JG, Fetter R, Spilker K, Cremers D, Shen K, Dunn AR, Goodman MB (2017) Genetic defects in β-spectrin and tau sensitize C. elegans axons to movement-induced damage via torque-tension coupling. Elife 6.

Kriz J, Zhu Q, Julien JP, Padjen AL (2000) Electrophysiological properties of axons in mice

lacking neurofilament subunit genes: disparity between conduction velocity and axon

diameter in absence of NF-H. Brain Res 885:32–44.

Kumar V, Zhang M-X, Swank MW, Kunz J, Wu G-Y (2005) Regulation of dendritic

morphogenesis by Ras-PI3K-Akt-mTOR and Ras-MAPK signaling pathways. J Neurosci

25:11288–11299.

Leite SC, Sampaio P, Sousa VF, Nogueira-Rodrigues J, Pinto-Costa R, Peters LL, Brites P,

Sousa MM (2016) The Actin-Binding Protein α-Adducin Is Required for Maintaining Axon

Diameter. Cell Rep 15:490–498.

Martin SM, O'Brien GS, Portera-Cailliau C, Sagasti A (2010) Wallerian degeneration of

zebrafish trigeminal axons in the skin is required for regeneration and developmental

pruning. Development 137:3985–3994.

Metzner K, Darawsha O, Wang M, Gaur N, Cheng Y, Rödiger A, Frahm C, Witte OW, Perocchi

F, Axer H, Grosskreutz J, Brill MS (2022) Age-dependent increase of cytoskeletal

components in sensory axons in human skin. Front Cell Dev Biol 10:965382.

Nabel AL, Teich L, Wohlfrom H, Alexandrova O, Heß M, Pecka M, Grothe B (2024)

Development of myelination and axon diameter for fast and precise action potential

conductance. Glia 72:794–808.

O'Brien GS, Rieger S, Wang F, Smolen GA, Gonzalez RE, Buchanan J, Sagasti A (2012)

Coordinate development of skin cells and cutaneous sensory axons in zebrafish. J Comp

Neurol 520:816–831.

Ohara O, Gahara Y, Miyake T, Teraoka H, Kitamura T (1993) Neurofilament deficiency in quail

caused by nonsense mutation in neurofilament-L gene. J Cell Biol 121:387–395.

Palanca AMS, Lee S-L, Yee LE, Joe-Wong C, Trinh LA, Hiroyasu E, Husain M, Fraser SE,

Pellegrini M, Sagasti A (2013) New transgenic reporters identify somatosensory neuron

subtypes in larval zebrafish. Dev Neurobiol 73:152–167.

Pan X, Hu Y, Lei G, Wei Y, Li J, Luan T, Zhang Y, Chu Y, Feng Y, Zhan W, Zhao C, Meunier

FA, Liu Y, Li Y, Wang T (2024) Actomyosin-II protects axons from degeneration induced by

mild mechanical stress. J Cell Biol 223.

Perge JA, Niven JE, Mugnaini E, Balasubramanian V, Sterling P (2012) Why do axons differ in

caliber? J Neurosci 32:626–638.

Pullarkat PA, Dommersnes P, Fernández P, Joanny J-F, Ott A (2006) Osmotically driven shape

transformations in axons. Phys Rev Lett 96:048104.

Rasmussen JP, Sack GS, Martin SM, Sagasti A (2015) Vertebrate epidermal cells are broad-

specificity phagocytes that clear sensory axon debris. J Neurosci 35:559–570.

Rosa JB, Nassman KY, Sagasti A (2023) Sensory axons induce epithelial lipid microdomain

remodeling and determine the distribution of junctions in the epidermis. Mol Biol Cell

34:ar5.

Rydmark M (1981) Nodal axon diameter correlates linearly with internodal axon diameter in

spinal roots of the cat. Neurosci Lett 24:247–250.

Sagasti A, Guido MR, Raible DW, Schier AF (2005) Repulsive interactions shape the

morphologies and functional arrangement of zebrafish peripheral sensory arbors. Curr Biol

15:804–814.

Sainio MT, Rasila T, Molchanova SM, Järvilehto J, Torregrosa-Muñumer R, Harjuhaahto S,

Pennonen J, Huber N, Herukka S-K, Haapasalo A, Zetterberg H, Taira T, Palmio J, Ylikallio

E, Tyynismaa H (2021) Neurofilament Light Regulates Axon Caliber, Synaptic Activity, and

Organelle Trafficking in Cultured Human Motor Neurons. Front Cell Dev Biol 9:820105.

Sakaguchi T, Okada M, Kitamura T, Kawasaki K (1993) Reduced diameter and conduction

velocity of myelinated fibers in the sciatic nerve of a neurofilament-deficient mutant quail.

Neurosci Lett 153:65–68.

Sánchez I, Hassinger L, Paskevich PA, Shine HD, Nixon RA (1996) Oligodendroglia regulate

the regional expansion of axon caliber and local accumulation of neurofilaments during

development independently of myelin formation. J Neurosci 16:5095–5105.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,

Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A (2012) Fiji: an open-source platform for biological-image analysis.

Nat Methods 9:676–682.

Seidl AH (2014) Regulation of conduction time along axons. Neuroscience 276:126–134.

Seidl AH, Rubel EW (2016) Systematic and differential myelination of axon collaterals in the

mammalian auditory brainstem: Systematic and Differential Myelination of Axon Collaterals.

Glia 64:487–494.

Seidl AH, Rubel EW, Harris DM (2010) Mechanisms for adjusting interaural time differences to

achieve binaural coincidence detection. J Neurosci 30:70–80.

Shao X, Sørensen MH, Xia X, Fang C, Hui TH, Chang RCC, Chu Z, Lin Y (2020) Beading of

injured axons driven by tension- and adhesion-regulated membrane shape instability. J R

Soc Interface 17:20200331.

Shorey M, Rao K, Stone MC, Mattie FJ, Sagasti A, Rolls MM (2021) Microtubule organization of

vertebrate sensory neurons in vivo. Dev Biol.

Sinclair JL, Fischl MJ, Alexandrova O, Heβ M, Grothe B, Leibold C, Kopp-Scheinpflug C (2017)

Sound-evoked activity influences myelination of brainstem axons in the trapezoid body. J

Neurosci 37:8239–8255.

Sinha B, Köster D, Ruez R, Gonnord P, Bastiani M, Abankwa D, Stan RV, Butler-Browne G,

Vedie B, Johannes L, Morone N, Parton RG, Raposo G, Sens P, Lamaze C, Nassoy P

(2011) Cells respond to mechanical stress by rapid disassembly of caveolae. Cell 144:402–

413.

Sitarska E, Diz-Muñoz A (2020) Pay attention to membrane tension: Mechanobiology of the cell

surface. Curr Opin Cell Biol 66:11–18.

Swärd C, Berthold CH, Nilsson-Remahl I, Rydmark M (1995) Axonal constriction at Ranvier's

node increases during development. Neurosci Lett 190:159–162.

Taylor NJ, Wang L, Brown A (2012) Neurofilaments are flexible polymers that often fold and

unfold, but they move in a fully extended configuration. Cytoskeleton (Hoboken) 69:535–

544.

Tuttle AM, Miller LN, Royer LJ, Wen H, Kelly JJ, Calistri NL, Heiser LM, Nechiporuk AV (2024)

Single-Cell Analysis of Rohon-Beard Neurons Implicates Fgf Signaling in Axon

Maintenance and Cell Survival. J Neurosci 44.

Uchida A, Peng J, Brown A (2023) Regulation of neurofilament length and transport by a

dynamic cycle of phospho-dependent polymer severing and annealing. Mol Biol Cell

34:ar68.

Voyvodic JT (1989a) Target size regulates calibre and myelination of sympathetic axons. Nature

342:430–433.

Voyvodic JT (1989b) Peripheral target regulation of dendritic geometry in the rat superior

cervical ganglion. J Neurosci 9:1997–2010.

Walker CL, Uchida A, Li Y, Trivedi N, Fenn JD, Monsma PC, Lariviére RC, Julien J-P, Jung P,

Brown A (2019) Local acceleration of neurofilament transport at nodes of Ranvier. J

Neurosci 39:663–677.

Wang F, Julien DP, Sagasti A (2013) Journey to the skin: Somatosensory peripheral axon

guidance and morphogenesis. Cell Adh Migr 7:388–394.

Wang T, Li W, Martin S, Papadopulos A, Joensuu M, Liu C, Jiang A, Shamsollahi G, Amor R,

Lanoue V, Padmanabhan P, Meunier FA (2020) Radial contractility of actomyosin rings

facilitates axonal trafficking and structural stability. J Cell Biol 219.

Williams PR, Marincu B-N, Sorbara CD, Mahler CF, Schumacher A-M, Griesbeck O,

Kerschensteiner M, Misgeld T (2014) A recoverable state of axon injury persists for hours

after spinal cord contusion in vivo. Nat Commun 5:5683.

Xu K, Zhong G, Zhuang X (2013) Actin, spectrin, and associated proteins form a periodic

cytoskeletal structure in axons. Science 339:452–456.

Zhu Q, Couillard-Després S, Julien JP (1997) Delayed maturation of regenerating myelinated

axons in mice lacking neurofilaments. Exp Neurol 148:299–316.