Title: Avian telencephalon and cerebellum volumes can be accurately estimated from digital brain

endocasts

Aubrey R. Keirnan[1], Felipe Cunha[2], Sara Citron[2], Gavin Prideaux[1], Andrew N. Iwaniuk[2]*, Vera

Weisbecker[1]*

[1] College of Science and Engineering, Flinders University, Adelaide SA, Australia

[2] Department of Neuroscience, University of Lethbridge, Lethbridge, Alberta, Canada

ORCiD:

Aubrey Keirnan 0000-0002-4033-6217

Felipe Cunha 0000-0002-3656-732X

Sara Citron 0009-0006-8138-3905

Gavin Prideaux 0000-0002-9958-0265

Andrew N. Iwaniuk 0000-0001-9273-3655

Vera Weisbecker 0000-0003-2370-4046

ABSTRACT

For studies of the evolution of vertebrate brain anatomy and potentially associated behaviours,

reconstructions of digital brain endocasts from computed tomography scans have revolutionised our

capacity to collect neuroanatomical data. However, measurements from digital endocasts must be

validated as reflecting actual brain anatomy, which is difficult because the collection of soft tissue

information through histology is laborious and time consuming. In birds, the reliability of digital

endocast measurements as volume proxies for the two largest brain regions – the telencephalon and

cerebellum - remains to be validated despite their use as proxies e.g. of cognitive performance or

flight ability. We here use the largest dataset of histology and digital endocasts to date, including

136 species from 25 avian orders, to compare digital endocast surface area measurements with

actual brain volumes of the telencephalon, cerebellum, and whole-brain endocast. Using linear and

phylogenetically informed regression analyses, we demonstrate that endocast surfaces are strongly

correlated with their brain volume counterparts for both absolute and relative size. This provides

empirical support for using endocast-derived cerebellar and telencephalic surface areas in existing

and future studies of living and extinct birds, with potential to expand to the dinosaur-bird transition

in the future.

INTRODUCTION

Over the past two decades, digital endocasts have become an increasingly important tool in the

study of neural and behavioural evolution in vertebrates [1-6]. Using computed tomography (CT)

scanned skulls, the brain cavity is digitally "filled" to approximate the volume and shape of the

brain [7]. These methods are largely non-destructive, allowing research into the neuroanatomy

of rare [8] and extinct species [9], including fossil specimens [10, 11]. For preservation purposes,

such as museum specimens or fossils embedded in sedimentary matrix, CT scanning offers non-

destructive access to otherwise unattainable information [12, 13]. To date, endocasts have been

used to study brain evolution across all vertebrate classes, including fishes [5], amphibians [2],

birds [14], non-avian reptiles [15, 16], and mammals [6, 17, 18]. Therefore, our understanding of

the brain-to-endocast relationship is fundamental to the study of vertebrate brain evolution.

Our knowledge of evolutionary patterns in brain size [13, 19], shape [6, 20], and composition [21]

has advanced substantially over the past century through endocast comparisons [13, 22]. For

example, early endocast research on pterosaur fossils, an extinct clade of flying reptiles,

revealed similarities between their brains with those of modern birds [23]. These included the

positions of the optic lobes (or midbrain) and the cerebellum, distinguishing pterosaurs and birds

from other reptiles [23]. More recently, endocast studies have been used to infer brain–behaviour

relationships, including social behaviour in big cats [24], nocturnality in birds [8, 9, 25], and the

habitats of snakes [16]. However, there is substantial variation in the brain–endocast

relationships among vertebrates [26], yet the use of endocast measurements relies on the

assumption that they are indicative of the brain region of interest [2, 3, 12]. While some studies

have verified the brain–endocast correlations for particular brain regions [27, 28], measurement

modalities such as shape or volume are not necessarily interchangeable [4, 6], meaning that

both must be tested to establish the validity of these methods. For fishes, amphibians, and non-

avian reptiles, in which brains do not fill the cranial cavity, inferences are limited as endocasts

overestimate brain size [2, 4, 29]. This can be an issue particularly at the root of iconic radiations

such as dinosaurs, whose closest living non-avian relatives – the crocodiles – have brains that

typically occupy less than half of their endocranial space in adults [26, 30]. In contrast, the

endocranial volume in birds is strongly correlated with brain volume [31].

The brains of birds have been the subject of extensive efforts to establish the relationship

between the endocranial cavity and brain tissue [22, 28, 31-33]. Their endocasts are remarkably

faithful to the shape of the brain itself, and endocranial volume is a valid proxy for actual brain

volume and widely used in studies of brain size (e.g. [14, 31, 34, 35]). Further, several brain

regions are discernible in endocasts [10, 12, 22, 28]. The endocast surface areas of two of these

brain regions, the wulst and optic lobes, are reliable estimates of the underlying hyperpallium

and optic tectum regions, respectively, of the brain [28]. However, inferences have also been

drawn from endocast measurements of the telencephalon and cerebellum, even though the

reliability of measuring these brain regions from endocasts remains untested. Changes in the

size of both regions are important in the evolution of differences in avian cognition [36] and the

brain becoming "flight-ready" [37, 38]. For example, the expansion of the telencephalon is a key

anatomical change in the evolution of the avian brain [37, 39] and associated with higher rates of

innovative behaviour [40]. Similarly, changes in relative cerebellum size, and part of the

cerebellum, are associated with behavioural differences across avian clades [41, 42]. They also

comprise the majority of brain volume [43] and the endocast surface. Exploiting the potential of

these two brain regions in explaining major transitions in avian brain evolution therefore requires

validation of how faithfully their endocast imprints reflect their soft tissue volume.

Here, we determine the brain–endocast relationships of the telencephalon and cerebellum using

the largest, phylogenetically diverse sample of bird species to date. While brain regions are

typically measured as volumes, endocasts lack the internal anatomical landmarks needed to

accurately divide them into volumes. This is especially true of the avian cerebellum, because its

anterior surface is covered by the posterior telencephalon. While surface area may not be

interchangeable with volume [6], we hypothesised this comparison would show a typical

isometric relationship, close to or equal to a 2:3 ratio or surface area increasing at a rate of 0.67

to that of the volume. However, as the relative size of these brain regions is often important for

studying neurobehavioural evolution [3, 26, 32, 44, 45], we also asked if their relative size when

measured from endocasts was comparable to that of brains. We expected that, if the endocast

reliably estimates the relative size of these regions, they would produce comparable

relationships, which would result in similar slopes.

METHODS

Specimens

Data on whole brain, telencephalon and cerebellum volumes were compiled from the literature [41,

43, 46-56] and supplemented with previously unpublished data from a histological collection

maintained by ANI [57]. These specimens were provided to ANI dead and were not euthanised for the

purpose of this study. Further, all procedures were completed in accordance with the guidelines and

policies of the Canada Council on Animal Care. All of these brains were gelatin-embedded and serially

sectioned in the coronal plane at a thickness of 40μm. After mounting and staining with thionin

acetate, the volumes were measured using unbiased stereology in StereoInvestigator

(Microbrightfield Inc., VT, USA) by FC. For some species, the telencephalon and cerebellum volumes

were measured from different individuals. In such cases, we included the brain volumes for both

individuals to accurately calculate the relative size of brain regions. More details on the brain

processing and volume measurement methods are described in [46]. All data and their specific sources

are provided in the electronic supplementary material [58].

To compare the brain volumes with surface areas from endocasts, we collected existing micro-

computed tomography (μCT) scans as well as completing additional scans for this study. Existing scans

were sourced through the online repository Morphosource (Project ID: 000642669), our previous

studies [8, 25] as well as completing additional scans. The source, museum identification, and the

scanning parameters for each specimen are provided in Supplementary Table 1. In total, we had brain

volume measurements and CT scans from 136 species, representing 58 families and 25 orders.

However, for four species, the volumes for either the telencephalon or cerebellum was unavailable –

two species from each region - reducing the total to 134 species for these brain regions.

Endocast reconstruction and surface area data collection

All endocast reconstruction and measurements were completed by AK. We imported the CT scans into

the image processing software Materialise Mimics (v.24.0; released in 2021 by Materialise NV) and

exported reconstructed skulls. Meshes of the inner surface of the brain cavity – the endocast – were

derived using the endomaker function from the Arothron package [59] in R [60] with the meshes

decimated to 20000 but otherwise using the base settings for all reconstructions. The meshes were

incomplete due to natural openings or damage to the skull, and often included excess material. To

complete the endocasts, these were imported into Geomagic Wrap (v.2021.2.2; released in 2023 by

3D Systems) to remove excess material and patch openings using the "flat" fill setting. Bumps and

crevices were also removed and subsequently patched with the "flat" fill along with the removal and

patching of nerves and blood vessels, following best practices as outlined by Balanoff, Bever [33]

(example in electronic supplementary material figure S1). As is to be expected, there was some

variation in the amount removed and patching needed for each specimen. However, efforts were

made to keep the reconstructions consistent and maintain the surface details of all the endocasts for

accuracy.

We defined the borders for the telencephalon and cerebellum by consulting previous studies which

partitioned endocasts for volume, surface area or landmark based measurements [4, 8, 21, 28, 61].

The posterior border of the telencephalon fossa was defined as the junction where it meets the

cerebellar fossa and continues ventrally through the crista tentorialis above the fossa tecti

mesenchephali. These borders then meet rostrally above the optic chiasma. For the cerebellum fossa,

the dorsal border was defined by the junction that meets the telencephalon fossa and continues along

the caudal border of the fossa tecti mesencephali on either side. This border continues as the crista

present at the ventral region of the cerebellum fossa, dorsal to the brainstem fossa, meeting in the

posterior region of the foramen magnum. We used the surface areas within these borders, which

encompass the visible imprints of both regions to estimate their sizes from the endocasts (Figures 1a,

b). We collected all measurements using Geomagic Wrap by following the borders outlined above. In

Figure 1c, we provide examples of these selections from each of the 25 clades in this study.

Figure 1. The endocast of a Eurasian treecreeper (*Certhia familiaris*) in rostral (A) and caudal views

(B). Both images show osteological landmarks used to delineate the endocast telencephalon and

cerebellum for measurements. The telencephalon is blue, the cerebellum is red, and the rest of the

endocast is grey. (C) Endocasts of a representative species of each of the orders examined in this

study using the same colour coding as in A and B. Note that endocasts shown here are of reduced

resolution and are not to scale.

Statistical analysis

All data were natural log transformed for analyses. To assess if the absolute size of the endocast

surface areas is an appropriate proxy of the absolute size of their brain counterpart, we completed

ordinary least squares (OLS) regression analyses with brain region volume as predictor and endocast

surface area as response variables in R [60]. From these, we calculated confidence intervals (the

confidence of the fit), prediction intervals (the confidence with which an unknown specimen could be

given a value), and the line of best fit for each set of measurements: whole endocast, telencephalon

and cerebellum. We expected the endocast surface areas to scale isometrically with the brain volume,

with surface area scaling at two-thirds the volume (i.e., yielding slopes of 0.67). To test this, we

assessed the slope coefficients of the linear relationships and additionally used Wald tests in the *car*

package [62], which uses the model coefficients and coefficient-covariance matrix to determine if the

relationship significantly deviated from the isometric slope of 0.67 [63]. Note that the relationship

between brains and endocasts is not expected to be impacted by phylogenetic structure of the data.

For this reason, linear modelling was the preferred statistical procedure. However, phylogenetic

generalised least squares models were also computed using the *evomap* package [64] to assess if

phylogenetic relationships impact on the estimates of the slope [65].

As we have previously shown that the variation in voxel sizes of the CT scans may cause minor

discrepancies for endocast measurements [8] we tested if this may have had an artificial effect on

endocast size. We did this by calculating the residuals of OLS regression analyses using the whole

endocast surface area and whole brain volume as dependent variables and voxel size as the predictor

variable. We expected that if this factor does not influence the results that the residuals should show

a similar relationship to our comparisons of the endocast and brain measurements.

We also computed the relative size for both the telencephalon and cerebellum by scaling them with

the remainder of the brain as the dependent variable [66], both in the surface area (endocast-based)

data and the volume (histology-based) data. This was to confirm that the slopes of the scaling

relationships were comparable between the two data sets. Because phylogenetic relationships exert

a significant effect on the relative size of brain regions [43, 65, 67, 68], we used both linear modelling

and phylogenetic generalised least squares modelling to derive the slopes and intercepts.

We derived our phylogeny by pruning the family-level phylogeny by Stiller, Feng [69] and used

Mesquite [70] to place species on this tree using order specific phylogenetic studies [71-81]. We then

completed the PGLS regression analyses and calculated phylogenetic confidence intervals (CI) for each

of the regions against the remainder of either the endocast or brain using the *evomap* package [64].

RESULTS

Our comparisons of the surface area measurements for the whole endocast, telencephalon, and

cerebellum against their brain volume counterparts yielded highly significant associations (Figure 2;

Table 1) which are unlikely to be influenced by the voxel size (see electronic supplementary material,

figure S2). Furthermore, the resulting slopes indicate isometric relationships (Table 1) which are not

significantly different from the expected 0.67 according to the Wald test comparison (Table 1).While

this overall pattern is highly consistent, some species fell outside of the prediction intervals (PI) for

the whole sample, meaning that their surface area measurements were not expected given their

corresponding volumes. In total, nine species fell outside of the PIs for at least one of the regressions

with three outside of the PIs for all regressions: the satin bowerbird (*Ptilonorhynchus violaceus*;

Passeriformes), black-capped chickadee (*Poecile atricapillus*; Passeriformes), and the collared

sparrowhawk (*Accipiter cirrocephalus*; Accipitriformes) (Figure 2).

For the comparisons of the relative sizes of the telencephalon and cerebellum, the surface area for

these regions regressed against the endocast remainder yielded similar slopes to that of the region

volume regressed against the brain remainder, for both the OLS and PGLS (Figure 3; Table 2). Parrots

(Psittaciformes) appear to deviate more from the regression lines in Figure 3 for both regions. This is

because parrots have relatively large telencephala and small cerebella compared with other species

[43]. The surface area measurements show a somewhat larger deviation for the telencephalon, which

could arise from the relationship between surface area and volume or that the optic lobes are smaller

in parrots [43, 82], allowing the telencephalon to spread out more. One species, the Kākāpō (*Strigops*

habroptilus), is outside of the PGLS confidence interval for the expected range of the telencephalon

surface area scaled against the endocast remainder surface area (Figure 3a). This species is a uniquely

nocturnal and flightless parrot that is neuroanatomically distinct within its clade [54] which may then

result in an atypical relationship between endocast and brain measurements. Overall, these results

indicate that the scaling relationship for both brain regions is similar when measured from endocast

surface areas and brain volumes.

Figure 2. Log transformed measurements of the (a) whole endocast, (b) telencephalon, and (c)

cerebellum surface areas measured from endocasts regressed against their brain volume counterparts

for 136 species of birds. The orders from this data are represented by unique combinations of colours

with point shapes. The black line represents the ordinary least-squares linear model regression; dark

grey area represents 95% confidence intervals; light grey area represents 95% prediction intervals.

Table 1. Results from ordinary least squares (OLS) regression analyses and Wald tests. OLS results

show the significant (p=<0.05) relationships between the endocast surface area measurements and

the brain volume measurements. Wald test results show that slopes do not significantly ($p=>0.05$)

differ from the expected isometric relationship between surface area and volume measurements

(0.67).

Dependent	Predictor	*df*	F-statistic	*p*-value	*y* intercept	R^2	Slope	Wald test *p*-value
log Endo. SA	log Brain VOL	134	3969	<0.01	1.78	0.97	0.68	0.64
log Tele. SA	log Tele. VOL	132	2954	<0.01	1.62	0.96	0.67	0.99
log Cere. SA	log Cere. VOL	132	2324	<0.01	1.17	0.95	0.67	0.84

215   Abbreviations: SA, surface area; Vol., volume; Endo., endocast; Tele., telencephalon; Cere.,

216   cerebellum.

Figure 3. Log transformed measurements comparing the scaling relationships of the telencephalon for

(a) the surface area against endocast remainder, and (b) the volume against brain remainder, as well

as the cerebellum for (c) the surface area against endocast remainder, and (d) the volume against

brain remainder for 136 species of birds. The orders from this data are represented by unique

combinations of colours with point shapes. Ordinary least squares regression (LM slope) is shown by

the black line; dark grey for the 95% confidence intervals; light grey for the 95% prediction intervals.

Phylogenetic generalised least squares (PGLS) regression shown by the dashed red line; pink area

shows the 95% confidence intervals of the PGLS.

226 **Table 2.** Results from phylogenetic generalised least squares (PGLS) regression models used to

227 compare the scaling relationship of endocast surface area and brain volume measurements of the

228 telencephalon and cerebellum.

Dependent	Predictor	*df*	*t*-value	*p*-value	*y* intercept	Slope
Tele. SA.	Endo. Remainder	134	32.2	<0.01	0	1.04
Tele. VOL	Brain remainder	134	28.97	<0.01	0.2	1.04
Cere. SA	Endo. Remainder	134	29.35	<0.01	-1.65	0.97
Cere. VOL	Brain remainder	134	33.48	<0.01	-1.57	0.96

Abbreviations: SA, surface area; Vol., volume; Endo., endocast; Tele., telencephalon; Cere.,

cerebellum.

DISCUSSION

Our results reveal that avian endocast surface areas for the whole endocast, the telencephalon, and

the cerebellum exhibit tight isometric relationships with the volume of their brain counterparts. This

result is in spite of a sample size of one individual per species for the endocast measurements and a

reliance on different individuals for the endocast and brain measurements, both of which could have

reduced the strength of the association between the two datasets [31]. By including a broad range of

species, our validation supports the use of endocast measurements for these regions in future studies,

while also supporting the conclusions from previous studies which relied on such measurements [8,

9, 25, 36, 37, 83, 84]. This result is particularly beneficial for studies on rare species [8, 25], and

palaeontological specimens [9, 85] for which brain measurements are not attainable.

We also found that the relative sizes for both the telencephalon and cerebellum regions are consistent

between the endocast surface areas and the soft tissue brain volumes, supporting their use in

comparisons of brain composition [43]. This is in spite of the absence of the entire anterior lobe of the

cerebellum in endocasts which is generally hidden from external view by the telencephalon, and the

cerebellar surface often obscured by the occipital sinus [3, 32, 86, 87]. For other endocast validation

research, such as that of the optic lobes and wulst [28], the regions are typically clearly defined and

largely unobscured. Nonetheless, our findings strongly suggest that the portion of the cerebellum

visible in endocasts is sufficient for estimating its relative volume.

While there is a strong brain–endocast association despite our reliance on different individuals, some

species showed disparities between specimens, likely due to intraspecific variation in brain size and

composition arising from sexual dimorphism [88-90] or subspecific, clinal, and other regional variation

[91]. In fact, in a previous study, we found substantial skull and endocast size differences for two

collared sparrowhawk specimens, likely due to sexual dimorphism [25]. Within this study, we suspect

variation from sexual dimorphism in hawks [88, 90], and pheasants [56, 92],and regional variation for

the bowerbird [89], chickadee [93], burrowing owl [94] and cormorant [95]. Furthermore, as variation

occurred when comparing volumes and surface areas directly to one another but did not appear to be

present in the analyses of relative brain region size we suspect this variation stems from using different

specimens for both datasets. While variation is expected, this raises concerns regarding the sample

size needed to represent a species in comparative studies, particularly of narrow taxonomic scope.

We nevertheless note the high correspondence between the brain and endocast datasets, suggesting

that these measurements can generally be used with small sample sizes.

The telencephalon and cerebellum make up the majority of the brain and endocast surface, making

their inclusion necessary for studies on brain composition [43]. By using known brain–behaviour

associations, such as the telencephalon size with cognitive abilities [36, 40, 96], and the cerebellum

with flight abilities [37, 38, 42], some aspects of ecology and behaviours may be inferred from these

regions [40, 46, 97-99]. Furthermore, even though some behaviourally relevant brain regions (such as

the hippocampus [100]) will always remain hidden within the brain, holistic approaches to endocast

comparisons can improve the accuracy of inferences by accounting for the influences that changes in

brain regions sizes can have on one another [14, 43, 99, 101]. For example, the relative size of the

telencephalon in owls is comparable with that of parrots [96], but this is largely due to their

hypertrophied wulst in relation to vision [43] and not the regions of the telencephalon associated with

cognition [102].

Overall, this study has confirmed the strong relationships between the surface areas of endocasts and

brain volumes in birds for absolute and relative size, both wholly and when divided into the

telencephalon and cerebellum regions. This result is in concordance with other endocast validations

[28, 31], and we suggest that further expanding the use of avian endocasts is plausible. However, these

methods remain speculative beyond the crown group Aves, such as for non-avian dinosaurs [2, 4, 29].

Outside of validating bird endocasts, comparisons with other living archosaurs, the crocodilians, may

assist with resolving the brain–endocast correspondence [2] and provide further evidence for the

strengths and limitations of these methods [4, 30]. With the increased accessibility that CT scanning

and associated digital reconstruction methods present [1, 10, 103], expanding and refining endocast

uses has the potential to shed light on the lives of extinct species and further resolve our

understanding of vertebrate brain evolution.

REFERENCES

1. Degrange, F.J., et al., *Anatomy and Evolution of Avian Brain and Senses: What Endocasts Can Tell Us*, in *Paleoneurology of Amniotes : New Directions in the Study of Fossil Endocasts*, M.T. Dozo, et al., Editors. 2023, Springer International Publishing: Cham. p. 333-364.

2. Clement, A.M., et al., *Brain Reconstruction Across the Fish-Tetrapod Transition; Insights From Modern Amphibians.* Frontiers in Ecology and Evolution, 2021. **9**.

3. Balanoff, A. and G.S. Bever, *The Role of Endocasts in the Study of Brain Evolution*. 2017. p. 223-241.

4. Watanabe, A., et al., *Are endocasts good proxies for brain size and shape in archosaurs throughout ontogeny?* Journal of Anatomy, 2019. **234**(3): p. 291-305.

5. Clement, A.M., et al., *Morphometric analysis of lungfish endocasts elucidates early dipnoan palaeoneurological evolution.* eLife, 2022. **11**: p. e73461.

6. Weisbecker, V., et al., *Global elongation and high shape flexibility as an evolutionary hypothesis of accommodating mammalian brains into skulls.* Evolution, 2021. **75**(3): p. 625-640.

7. Kubo, D., et al., *Accuracy and precision of CT-based endocranial capacity estimations: a comparison with the conventional millet seed method and application to the Minatogawa 1 skull.* Anthropological Science, 2008. **116**(1): p. 77-85.

8. Iwaniuk, A.N., et al., *The endocast of the Night Parrot (Pezoporus occidentalis) reveals insights into its sensory ecology and the evolution of nocturnality in birds.* Scientific Reports, 2020. **10**(1): p. 9258.

9. Torres, C.R. and J.A. Clarke, *Nocturnal giants: evolution of the sensory ecology in elephant birds and other palaeognaths inferred from digital brain reconstructions.* Proceedings. Biological sciences, 2018. **285**(1890).

10. Knoll, F. and S. Kawabe, *Avian palaeoneurology: Reflections on the eve of its 200th anniversary.* Journal of Anatomy, 2020. **236**(6): p. 965-979.

11. Heingård, M., et al., *New Remains of Scandiavis mikkelseni Inform Avian Phylogenetic Relationships and Brain Evolution.* Diversity, 2021. **13**(12).

12. Walsh, S.A. and M.A. Knoll, *Directions in palaeoneurology.* Special Papers in Palaeontology, 2011. **86**(1228): p. 263-279.

13. Buchholtz, E.A. and E.-A. Seyfarth, *The Study of "Fossil Brains": Tilly Edinger (1897–1967) and the Beginnings of Paleoneurology.* BioScience, 2001. **51**(8): p. 674-682.

14. Ksepka, D.T., et al., *Tempo and Pattern of Avian Brain Size Evolution.* Curr Biol, 2020. **30**(11): p. 2026-2036.e3.

15. Beyrand, V., et al., *Multiphase progenetic development shaped the brain of flying archosaurs.* Scientific Reports, 2019. **9**(1): p. 10807.

16. Allemand, R., et al., *Comparative morphology of snake (Squamata) endocasts: evidence of phylogenetic and ecological signals.* Journal of Anatomy, 2017. **231**(6): p. 849-868.

17. Todorov, O.S., et al., *Down a Rabbit Hole: Burrowing Behaviour and Larger Home Ranges are Related to Larger Brains in Leporids.* Journal of Mammalian Evolution, 2022. **29**(4): p. 957-967.

18. Marino, L., et al., *Reconstructing cetacean brain evolution using computed tomography.* The anatomical record. Part B, New anatomist, 2003. **272B**(1): p. 107-117.

19. Montgomery, S.H., et al., *Reconstructing the ups and downs of primate brain evolution: implications for adaptive hypotheses and Homo floresiensis.* BMC Biology, 2010. **8**(1): p. 9.

20. Carril, J., et al., *Comparative brain morphology of Neotropical parrots (Aves, Psittaciformes) inferred from virtual 3D endocasts.* Journal of Anatomy, 2016. **229**(2): p. 239-251.

21. Balanoff, A.M., J.B. Smaers, and A.H. Turner, *Brain modularity across the theropod–bird transition: testing the influence of flight on neuroanatomical variation.* Journal of anatomy, 2016. **229**(2): p. 204-214.

22. Edinger, T., *Paleoneurology 1804-1966 : an annotated bibliography*. 1st 1975. ed. Advances
in Anatomy, Embryology and Cell Biology, 49. 1975, Berlin ;: Springer-Verlag.
23. Edinger, T., *The brain of Pterodactylus.* American journal of science: devoted to the
geological sciences and to related fields, 1941. **239**(9): p. 665.
24. Sakai, S.T., et al., *Big Cat Coalitions: A Comparative Analysis of Regional Brain Volumes in
Felidae.* Frontiers in Neuroanatomy, 2016. **10**.
25. Keirnan, A., et al., *Not like night and day: the nocturnal letter-winged kite does not differ
from diurnal congeners in orbit or endocast morphology.* Royal Society Open Science, 2022.
9(5): p. 220135.
26. Jerison, H.J., *Evolution of the Brain and Intelligence.* Current Anthropology, 1975. **16**(3): p.
403-426.
27. Sakai, S.T., et al., *Virtual endocasts: an application of computed tomography in the study of
brain variation among hyenas.* Annals of the New York Academy of Sciences, 2011. **1225**(S1):
p. E160-E170.
28. Early, C.M., et al., *Endocast structures are reliable proxies for the sizes of corresponding
regions of the brain in extant birds.* Journal of Anatomy, 2020. **237**(6): p. 1162-1176.
29. Morhardt, A.C., R.C. Ridgely, and L.M. Witmer. *From endocast to brain: assessing brain size
and structure in extinct archosaurs using gross anatomical brain region approximation
(GABRA)*. in *Journal of Vertebrate Paleontology*. 2012. TAYLOR & FRANCIS INC 325
CHESTNUT ST, SUITE 800, PHILADELPHIA, PA 19106 USA.
30. Jirak, D. and J. Janacek, *Volume of the crocodilian brain and endocast during ontogeny.* PLOS
ONE, 2017. **12**(6): p. e0178491.
31. Iwaniuk, A. and J. Nelson, *Can endocranial volume be used as an estimate of brain size in
birds? Can J Zool.* Canadian Journal of Zoology, 2002. **80**: p. 16-23.
32. Witmer, L.M., et al., *Using CT to Peer into the Past: 3D Visualization of the Brain and Ear
Regions of Birds, Crocodiles, and Nonavian Dinosaurs*, in *Anatomical Imaging: Towards a
New Morphology*, H. Endo and R. Frey, Editors. 2008, Springer Japan: Tokyo. p. 67-87.
33. Balanoff, A.M., et al., *Best practices for digitally constructing endocranial casts: examples
from birds and their dinosaurian relatives.* Journal of Anatomy, 2016. **229**(2): p. 173-190.
34. Smaers, J.B., et al., *The evolution of mammalian brain size.* Science Advances. **7**(18): p.
eabe2101.
35. Weisbecker, V. and A. Goswami, *Brain size, life history, and metabolism at the
marsupial/placental dichotomy.* Proceedings of the National Academy of Sciences, 2010.
107(37): p. 16216-16221.
36. Walsh, S.A. and F. Knoll, *The Evolution of Avian Intelligence and Sensory Capabilities: The
Fossil Evidence*. 2017, Tokyo: Springer Japan: Tokyo. p. 59-69.
37. Balanoff, A.M., et al., *Evolutionary origins of the avian brain.* Nature, 2013. **501**(7465): p. 93-
96.
38. Witmer, L.M., et al., *Neuroanatomy of flying reptiles and implications for flight, posture and
behaviour.* Nature, 2003. **425**(6961): p. 950-953.
39. Striedter, G.F. and R.G. Northcutt, *Brains through time: a natural history of vertebrates*.
2019: Oxford University Press.
40. Lefebvre, L., et al., *Feeding innovations and forebrain size in birds.* Animal behaviour, 1997.
53(3): p. 549-560.
41. Cunha, F., et al., *A quantitative analysis of cerebellar anatomy in birds.* Brain Structure and
Function, 2021. **226**(8): p. 2561-2583.
42. Iwaniuk, A.N., P.L. Hurd, and D.R.W. Wylie, *Comparative Morphology of the Avian
Cerebellum: II. Size of Folia.* Brain, behavior and evolution, 2007. **69**(3): p. 196-219.
43. Iwaniuk, A.N. and P.L. Hurd, *The evolution of cerebrotypes in birds.* Brain, Behavior and
Evolution, 2005. **65**(4): p. 215-230.

44.   Sol, D., et al., *Evolutionary Divergence in Brain Size between Migratory and Resident Birds.*
PLOS ONE, 2010. **5**(3): p. e9617.

45.   Sol, D., L. Lefebvre, and J.D. Rodríguez-Teijeiro, *Brain size, innovative propensity and*
migratory behaviour in temperate Palaearctic birds. Proceedings of the Royal Society B:
Biological Sciences, 2005. **272**(1571): p. 1433-1441.

46.   Barros da Cunha, F., *Allometric Relationships and the Evolution of the Avian Brain*. 2021,
ProQuest Dissertations Publishing.

47.   Iwaniuk, A.N. and J.E. Nelson, *Developmental differences are correlated with relative brain*
size in birds: a comparative analysis. Canadian journal of zoology, 2003. **81**(12): p. 1913-
1928.

48.   Portmann, A., *Etudes sur la cérébralisation des oiseaux. II. Les indices intra-cérébraux.*
Alauda, 1947. **15**: p. 1-15.

49.   Kalisińska, E., *Anseriform brain and its parts versus taxonomic and ecological categories.*
Brain, Behavior and Evolution, 2005. **65**(4): p. 244-261.

50.   Corfield, J.R., et al., *Morphometric Analysis of Telencephalic Structure in a Variety of*
Neognath and Paleognath Bird Species Reveals Regional Differences Associated with Specific
Behavioral Traits. Brain, Behavior and Evolution, 2012. **80**(3): p. 181-195.

51.   Alama, S. and N. Bee-dee-Speroni, *índices cerebrales y composición cuantitativa encefálica*
en Athene cunicularia y Tyto alba (Strigiformes: Strigidae y Tytonidae). Facena, 1992. **9**: p.
19-37.

52.   Boire, D. and G.J.J.o.B.R. Baron, *Allometric comparison of brain and main brain subdivisions*
in birds. 1994. **35**(1): p. 49-66.

53.   Carezzano, F. and N. Bee de Speroni, *Composicion volumetrica encefalica e indices cerebrales*
en tres aves de ambiente acuatico (Ardeidae, Podicipedidae, Rallidae). Facena, 1995. **11**: p.
75-83.

54.   Corfield, J.R., et al., *Anatomical Specializations for Nocturnality in a Critically Endangered*
Parrot, the Kakapo (Strigops habroptilus) (Visual System of the Kakapo). PLoS ONE, 2011.
6(8): p. e22945.

55.   Day, L.B., D.A. Westcott, and D.H. Olster, *Evolution of bower complexity and cerebellum size*
in bowerbirds. Brain behavior and evolution, 2005. **66**(1): p. 62-72.

56.   Krilow, J.M. and A.N. Iwaniuk, *Seasonal Variation in Forebrain Region Sizes in Male Ruffed*
Grouse (Bonasa umbellus). Brain, Behavior and Evolution, 2015. **85**(3): p. 189-202.

57.   Iwaniuk, A.N., *The importance of scientific collecting and natural history museums for*
comparative neuroanatomy. Annals of the New York Academy of Sciences, 2011. **1225**(S1):
p. E1-E19.

58.   Keirnan, A.R., et al., *Avian telencephalon and cerebellum volumes can be accurately*
estimated from digital brain endocasts 2024.

59.   Profico, A., et al., *Endomaker, a new algorithm for fully automatic extraction of cranial*
endocasts and the calculation of their volumes. American Journal of Physical Anthropology,
2020. **172**(3): p. 511-515.

60.   R Core Team, *R: A language and environment for statistical computing. R Foundation for*
Statistical Computing, Vienna, Austria. 2021, R Foundation for Statistical Computing,.

61.   Kawabe, S., et al., *Variation in avian brain shape: relationship with size and orbital shape.* J
Anat, 2013. **223**(5): p. 495-508.

62.   Fox, J. and S. Weisberg, *An R companion to applied regression*. 2018: Sage publications.

63.   Goldman, C.A., et al., *Principles of allometry.* Tested studies for laboratory teaching, 1990.
11: p. 43-72.

64.   Smaers, J. and C. Mongle, *evomap: R package for the evolutionary mapping of continuous*
traits. GitHub (https://github. com/JeroenSmaers/evomap), 2014.

65.   Garland, T., Jr. and A.R. Ives, *Using the Past to Predict the Present: Confidence Intervals for*
Regression Equations in Phylogenetic Comparative Methods. Am Nat, 2000. **155**(3): p. 346-
364.
66.   Deacon, T.W., *Fallacies of progression in theories of brain-size evolution.* International
Journal of Primatology, 1990. **11**(3): p. 193-236.
67.   Garland, T., Jr., A.F. Bennett, and E.L. Rezende, *Phylogenetic approaches in comparative*
physiology. Journal of Experimental Biology, 2005. **208**(16): p. 3015-3035.
68.   Corfield, J.R., et al., *Diversity in olfactory bulb size in birds reflects allometry, ecology, and*
phylogeny. Front Neuroanat, 2015. **9**: p. 102.
69.   Stiller, J., et al., *Complexity of avian evolution revealed by family-level genomes.* Nature,
2024.
70.   Maddison, W.P. and D.R. Maddison, *Mesquite: a modular system for evolutionary analysis.*
Version 3.81. 2023.
71.   Fuchs, J., J.A. Johnson, and D.P. Mindell, *Rapid diversification of falcons (Aves: Falconidae)*
due to expansion of open habitats in the Late Miocene. Molecular Phylogenetics and
Evolution, 2015. **82**: p. 166-182.
72.   Prum, R.O., et al., *A comprehensive phylogeny of birds (Aves) using targeted next-generation*
DNA sequencing. Nature, 2015. **526**(7574): p. 569-573.
73.   Andersen, M., et al., *A phylogeny of kingfishers reveals an Indomalayan origin and elevated*
rates of diversification on oceanic islands. Journal of Biogeography, 2017. **45**.
74.   Shakya, S.B., et al., *Tapping the woodpecker tree for evolutionary insight.* Molecular
Phylogenetics and Evolution, 2017. **116**: p. 182-191.
75.   Garcia-R, J.C., et al. *Phylogenomic Reconstruction Sheds Light on New Relationships and*
Timescale of Rails (Aves: Rallidae) Evolution. Diversity, 2020. **12**, DOI: 10.3390/d12020070.
76.   Kimball, R.T., P.A. Hosner, and E.L. Braun, *A phylogenomic supermatrix of Galliformes*
(Landfowl) reveals biased branch lengths. Mol Phylogenet Evol, 2021. **158**: p. 107091.
77.   Wink, M. and H. Sauer-Gurth, *Molecular taxonomy and systematics of owls (Strigiformes)—*
An update. Airo, 2021. **29**: p. 487-500.
78.   Černý, D. and R. Natale, *Comprehensive taxon sampling and vetted fossils help clarify the*
time tree of shorebirds (Aves, Charadriiformes). Molecular Phylogenetics and Evolution,
2022. **177**: p. 107620.
79.   Hruska, J.P., et al., *Ultraconserved elements resolve the phylogeny and corroborate patterns*
of molecular rate variation in herons (Aves: Ardeidae). Ornithology, 2023. **140**(2): p.
ukad005.
80.   Catanach, T.A., M.R. Halley, and S. Pirro, *Enigmas no longer: using ultraconserved elements*
to place several unusual hawk taxa and address the non-monophyly of the genus Accipiter
(Accipitriformes: Accipitridae). Biological journal of the Linnean Society, 2024.
81.   Smith, B.T., et al., *Phylogenomic Analysis of the Parrots of the World Distinguishes Artifactual*
from Biological Sources of Gene Tree Discordance. Systematic Biology, 2023. **72**(1): p. 228-
241.
82.   Iwaniuk, A.N., et al., *Allometric Scaling of the Tectofugal Pathway in Birds.* Brain, Behavior
and Evolution, 2010. **75**(2): p. 122-137.
83.   Early, C.M., R.C. Ridgely, and L.M. Witmer, *Beyond endocasts: Using predicted brain-*
structure volumes of extinct birds to assess neuroanatomical and behavioral inferences.
Diversity, 2020. **12**(1).
84.   Milner, A.C. and S.A. Walsh, *Avian brain evolution: new data from Palaeogene birds (Lower*
Eocene) from England. Zoological journal of the Linnean Society, 2009. **155**(1): p. 198-219.
85.   Smith, N.A. and J.A. Clarke, *Endocranial Anatomy of the Charadriiformes: Sensory System*
Variation and the Evolution of Wing-Propelled Diving. PLOS ONE, 2012. **7**(11): p. e49584.

86.   Ksepka, D.T., et al., *Evolution of the brain and sensory organs in Sphenisciformes: new data*
from the stem penguin Paraptenodytes antarcticus: PENGUIN BRAIN EVOLUTION. Zoological
journal of the Linnean Society, 2012: p. no-no.

87.   Walsh, S.A., et al., *Avian Cerebellar Floccular Fossa Size Is Not a Proxy for Flying Ability in*
Birds. PloS one, 2013. **8**(6): p. e67176-e67176.

88.   Debus, S.J.S., *Birds of prey of Australia: a field guide*. 2nd ed. ed, ed. J.N. Davies and A.
BirdLife. 2012, Collingwood, Vic.: Collingwood, Vic. : CSIRO Publishing in association with
Birdlife Australia.

89.   Menkhorst, P., et al., *The Australian bird guide*. 2017: Csiro Publishing Melbourne.

90.   Dunne, P. and K.T. Karlson, *Birds of Prey: Hawks, Eagles, Falcons, and Vultures of North*
America. 2017: Houghton Mifflin Harcourt.

91.   Baladrón, A.V., et al., *Body size and sexual dimorphism in the southernmost subspecies of the*
Burrowing Owl (Athene cunicularia cunicularia). Journal of Raptor Research, 2015. **49**(4): p.
479-485.

92.   McGowan, P.J.K. and G.M. Kirwan. *Silver Pheasant (Lophura nycthemera), version 1.0. In*
Birds of the World (J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, and E. de Juana, Editors).
2020 [cited 2024.

93.   Roth, T.C. and V.V. Pravosudov, *Hippocampal volumes and neuron numbers increase along a*
gradient of environmental harshness: a large-scale comparison. Proc Biol Sci, 2009.
276(1656): p. 401-5.

94.   Poulin, R.G., et al. *Burrowing Owl (Athene cunicularia), version 1.0. In Birds of the World (A.*
F. Poole, Editor). 2020.

95.   Hatch, J.J., et al. *Great Cormorant (Phalacrocorax carbo), version 1.0. In Birds of the World (S.*
M. Billerman, Editor). 2020 [cited 2024.

96.   Iwaniuk, A.N., K.M. Dean, and J.E. Nelson, *Interspecific Allometry of the Brain and Brain*
Regions in Parrots (Psittaciformes): Comparisons with Other Birds and Primates. Brain Behav
Evol, 2004. **65**(1): p. 40-59.

97.   Burish, M.J., H.Y. Kueh, and S.S.H. Wang, *Brain Architecture and Social Complexity in Modern*
and Ancient Birds. Brain, behavior and evolution, 2004. **63**(2): p. 107-124.

98.   Lefebvre, L., et al., *Feeding Innovations and Forebrain Size in Australasian Birds.* Behaviour,
1998. **135**(8/9): p. 1077-1097.

99.   Iwaniuk, A.N., K.M. Dean, and J.E. Nelson, *A mosaic pattern characterizes the evolution of*
the avian brain. Proceedings of the Royal Society B: Biological Sciences, 2004. **271**(Suppl 4):
p. S148-S151.

100.   Todorov, O.S., et al., *Primate hippocampus size and organization are predicted by sociality*
but not diet. Proceedings of the Royal Society B: Biological Sciences, 2019. **286**(1914): p.
20191712.

101.   Carlisle, A., et al., *Testing hypotheses of developmental constraints on mammalian brain*
partition evolution, using marsupials. Scientific Reports, 2017. **7**(1): p. 4241.

102.   Sayol, F., et al., *Environmental variation and the evolution of large brains in birds.* Nature
Communications, 2016. **7**(1): p. 13971.

103.   Davies, T.G., et al., *Open data and digital morphology.* Proceedings of the Royal Society B:
Biological Sciences, 2017. **284**(1852): p. 20170194.

526