1 **Midbrain dopamine neuronal activity modulates splenic immunity through a brain-to-body**
2 **circuit**
3

4 Adithya Gopinath[1,2]*, Alfonso junior Apicella[3], Alice Bertero[3], Leah T. Phan[1], Emily Miller[1], Jordan
5 Seibold[1], Shannon C. Wall[2], Jordan Follett[2], Aidan Smith[1], Elijah Freedman[1], Lauryn Hornstein[1],
6 Logan Wood[1], Hannah Ciupe[1], Diya Behal[1], Arianna Grillo[1], Delaney Booth[1], Sabrina Campos[1],
7 Nirvana Soundara[1], Daniel L. Purich[4], Pedro Trevizan-Baú[5], Leah R. Reznikov[5], Nikhil Urs[6], Olga
8 Guryanova[6], Mathew J. Farrer[2], Daniel W. Wesson[6], Habibeh Khoshbouei[1]*
9
10
1- Department of Neuroscience, University of Florida, College of Medicine, Gainesville, FL
USA
2- Department of Neurology, University of Florida, College of Medicine, Gainesville, FL USA
3- Department of Neuroscience, Developmental and Regenerative Biology, University of
Texas San Antonio, San Antonio, Texas USA
4- Department of Biochemistry and Molecular Biology, University of Florida, College of
Medicine, Gainesville, FL USA
5- Department of Physiological Sciences, University of Florida, College of Veterinary
Medicine, Gainesville, FL USA
6- Department of Pharmacology and Therapeutics, University of Florida, College of Medicine,
Gainesville, FL USA

22
Abstract

Midbrain dopamine neurons are well-known to shape central nervous system function, yet
there is growing evidence for their influence on the peripheral immune system. Here we
demonstrate that midbrain dopamine neurons form a circuit to the spleen via a multi-synaptic
pathway from the dorsal vagal complex (DVC) through the celiac ganglion. Midbrain dopamine
neurons modulate the activity of D1- and D2-like dopamine receptor-expressing DVC neurons. *In*
vivo activation of midbrain dopamine neurons induced dopamine release in the DVC and
increased immediate early gene expression in both the DVC and celiac ganglion. Activation of
this midbrain-to-spleen circuit reduced spleen weight and decreased naïve CD4$^+$ T-cell
populations without affecting total T-cell numbers. These findings unveil a functional midbrain-
DVC-celiac ganglion-spleen pathway, through which midbrain dopamine neurons modulate
splenic immunity. These novel insights into the neural regulation of the immune system have
important implications for diseases involving altered dopamine neurotransmission and highlight
potential targets for immunotherapeutic interventions.

37
Introduction
While substantial progress has been made in unraveling body-to-brain communication[1], the
reverse pathway, how the brain influences peripheral organs, particularly through dopamine
signaling, remains unclear. Emerging evidence highlights that midbrain dopamine neurons exert
significant influence on peripheral systems, particularly in the context of neuroimmune interactions,
but the circuits connecting them to peripheral immune organs remain largely undefined. Zhu and
colleagues[2], for instance, identified a functional link between central nervous system (CNS) pain
processing and splenic immunity, suggesting that neural pathways involved in immune regulation
extend well beyond the brain's immediate environment to influence key organs, such as the
spleen. Such findings imply that midbrain dopaminergic neurons may play a previously
uncharacterized role in orchestrating peripheral immune responses. Indeed, degeneration of
midbrain dopamine neurons is closely associated with changes in peripheral immune dysfunction
believed to contribute to disease progression[3-5]. Similarly, psychostimulant-induced activation of
midbrain dopamine neurons regulates peripheral immunity and influence wound healing[6-11].

Moreover, the well-characterized connection between peripheral inflammation and infection,
manifested symptomatically as anhedonia and reduced motivation[12-14], strongly suggest a link
between dopamine neurotransmission and peripheral immunity. Despite such findings, the
specific circuits through which midbrain dopamine neurons communicate with peripheral immune
organs remain unclear.
By connecting the brainstem to multiple peripheral organs including heart, lung, gut, and
spleen[15-18], the vagus nerve is one of the most characterized routes for brain-body communication.
While body-to-brain communication through vagal efferents has been extensively studied, the
corresponding brain-to-periphery circuits that use dopamine signaling to mediate the brain's
control over immune functions remain to be defined. The dorsal vagal complex (DVC) plays a
pivotal role in modulating immune function through the cholinergic anti-inflammatory pathway[19-21].
The DVC includes the dorsal motor nucleus of the vagus (DMV), the area postrema (AP), and the
nucleus of the solitary tract (NTS) and is optimally positioned to transmit signals from midbrain
dopamine neurons to peripheral organs. Prior work revealed that midbrain neurons innervate the
DVC and separately that midbrain neuron output appears to influence colon motility[22,23]. In
addition, vagal innervation of peripheral organs including the spleen contribute to both
inflammation and its' resolution[24-28]. While these advances have strongly supported the existence
of a brain-to-periphery system which can influence both peripheral organs and immune system
function, the specific role of dopamine in the modulation of splenic immunity is unclear.
The spleen is innervated by both cholinergic[29-31] and catecholaminergic[32-35] fibers and is
essential for survival, with roles in the production of blood cells and immune function. Both inputs
modulate splenic function, immune cell populations and the broader peripheral immune landscape.
Here we used a combination of viral approaches to map a midbrain dopamine neurons→DVC→
spleen circuit, focusing on how midbrain dopamine neuron activity modulates splenic immune
phenotype. We demonstrate that midbrain dopamine neurons project to the DVC, can activate
dopaminergic receptors expressed on DVC neurons, and that their activity is transmitted through
the celiac ganglia to reach the spleen. Furthermore, we show that the activation of this circuit
alters peripheral immune function, including spleen weight and splenic T-cell populations. Notably,
transection of the cervical vagus nerve abolishes the impact of midbrain dopamine neuron
activation on splenic immunity, confirming the essential role of vagal signaling in this brain-to-
spleen pathway. Collectively, our findings provide novel insights into the mechanisms underlying
the regulation of peripheral immunity by midbrain dopamine neurons, integrating prior
observations into a cohesive circuit model, casting new light on the interplay between central
dopamine signaling and peripheral immunity, and taking us closer to the root causes of disorders
characterized by dysregulated CNS dopamine signaling.
87
88
89 **Results and Discussion**
90
Anterograde mapping of midbrain dopaminergic neurons confirms dopaminergic
projections to the dorsal vagal complex.
Prior work established CNS circuity linking midbrain with the DVC[1,2,22,23,36-38]. Here we built
upon these reports and performed two initial viral-tracing based studies to identify 1) are midbrain
neuron inputs to the DVC dopaminergic and 2) are they monosynaptic?
To address the first question we utilized DAT-Cre mice, which express Cre recombinase under
the control of the dopamine transporter (DAT)[39], and injected AAV2-Ef1a-DIO-EYFP into the
midbrain (Figure 1A). This allowed for EYFP expression in tyrosine hydroxylase+ (TH[+]) dopamine
neuron cell bodies in the midbrain as well as their axons in the median forebrain bundle (Figures
1B & 1C and Supplementary Figure 1). As the gold standard for cholinergic neurons, choline
acetyltransferase (ChAT) co-staining was used to identify the DVC (Figure 1D). As expected,

EYFP[+] fibers were detected throughout the DVC (Figures 1D-E), including in the DMV and NTS,
showing midbrain dopaminergic neurons send projections to the DVC.
We next investigated if these midbrain dopaminergic neurons form synapses within the DVC.
Using a complementary viral tracing approach (Figure 2A), we injected DAT-Cre mice with AAV1-
hSyn-Flex-mGFP-2A-Synaptophysin-mRuby into the midbrain. This drives expression of the
Synaptophysin-mRuby fusion protein in dopaminergic synapses that originate form midbrain
neurons[40]. We next inspected the DVC for mRuby[+] puncta. mRuby[+] puncta were confirmed at the
injection site in the midbrain as well as in known projection areas, such as the striatum (Figure
2B & 2C, Supplemental Figure 2). Importantly, mRuby[+] puncta were observed throughout the
DVC, including in the DMV and NTS (Figure 2D-E), indicating that DAT[+] midbrain dopamine
neurons form synapses within the DVC and its subnuclei.
113
DVC neurons express D1- and D2-like receptors.
The above findings support a two-part hypothesis: 1) changes in midbrain dopamine neuron
activity and dopamine release modulate DVC neuronal activity; and 2) this modulation, in turn,
regulates peripheral systems downstream of the DVC, and may comprise a previously
unrecognized component of the midbrain dopamine neuron-to-peripheral immunity circuit. As
noted above, choline acetyltransferase (ChAT) is a reliable marker to help identify the DVC. To
establish a receptor-based foundation for testing our hypotheses, we performed whole-cell patch-
clamp recordings in ChAT-GFP mice, targeting GFP-expressing ChAT[+] neurons, the major
parasympathetic output population[20,41], as well as GFP-negative neurons in the DVC. This
experimental approach allows for an assessment the effects of dopamine on the intrinsic
excitability and firing properties of the neurons identified. Whole-cell patch-clamp recordings were
performed on coronal brainstem slices in the presence of synaptic blockers for GABAergic and
glutamatergic transmission, enabling isolation of the direct, postsynaptic actions of dopamine
(Figure 3A). Firstly, since GFP expression might highlight only a subset of ChAT neurons in the
DVC and cause misleading identification of DVC neurons' identity, we compared the intrinsic
properties of GFP-positive (N=22) and GFP-negative (N=19) neurons, to determine if they belong
to two separate classes of neurons. The vast majority of GFP-positive neurons (77%, 17/22)
showed spontaneous firing activity when recorded in cell-attached or current-clamp configuration
(Figure 3B), a common feature of ChAT-expressing neurons in other brain regions[42] and of DVC
neurons[43]. GFP-negative neurons, on the other hand, do not display such behavior. The two
populations have comparable passive properties, such as input resistance (Ri GFP[+] 639.6 ± 100.5
MΩ, Ri GFP[-] 878.9 ± 80.3 MΩ, p=0.076) and membrane time constant (tau, GFP[+] 0.46 ± 0.03
msec, GFP[-] 0.36 ± 0.03 msec) (Figure 3C), and their voltage threshold to fire an action potential
(V threshold, GFP[+] -32.1 ±1.5 mV, GFP[-] -31.9 ±1.6 mV, p=0.92, Figure 3C). However, they display
a wide array of differences in their active properties, such as current threshold (I threshold, GFP[+]
81.8 ± 8.4 pA, GFP[-] 50 ± 0 pA, p=0.001), action potential height (V height, GFP[+] 90.6 ± 2.3 mV,
GFP[-] 59.0 ± 2.2 mV, p=0.0008), action potential halfwidth (AP halfwidth, GFP[+] 0.73 ± 0.04 msec,
GFP[-] 0.55 ± 0.2 msec, p=0.0003), the relationship between injected current and AP frequency (f/I
slope, GFP[+] 0.08 ± 0.01 Hz/pA, GFP[-] 0.25 ± 0.05 Hz/pA, p=0.001) and spike frequency adaptation
ratio (SFA ratio GFP[+] 1.07 ± 0.011, GFP[-] 0.59 ±0.05, p=0.0006) (Figure 3D). This confirms that
GFP-positive and GFP-negative neurons in the DVC are two separate neuronal populations with
essentially different firing properties.
We next examined how dopamine modulates the intrinsic properties of GFP-positive (N=22)
and GFP-negative (N=19) DVC neurons. After determining their baseline intrinsic properties, we
applied 50 μM dopamine to the ACSF bath and reassessed the neurons' properties, for
comparison. Overall, dopamine did not change the passive membrane properties of GFP-positive
neurons (Figure 3E) but affected the input resistance of GFP-negative neurons, which showed a
significant decrease (Pre dopamine: 878.89 ± 80.26 MΩ; Post dopamine: 656.37 ± 63.43 MΩ;
p=0.036, Figure 3E). Based on the effect of the firing properties (Figure 3F, Supplementary Table

1), we could identify three main neuronal groups: GFP-expressing ChAT neurons (N=22) and
GFP-negative neurons (N=10/19) with an overall decrease in firing activity and increase in
rheobase, and a subset of GFP-negative neurons (N=9/19) unresponsive to dopamine, as shown
in three representative examples in Figure 3F, whose f/I curve, representing their changes in firing
pattern, demonstrate the criteria for classification (Figure 3G, Supplementary Table 1). The two
responsive DVC populations show significant changes in membrane and current thresholds
(Figure 3H, Supplementary Table 1) and V height, but not in AP halfwidth, f/I slope and SFA ratio
(Figure 3I-J, Supplementary Table 1). This behavior is compatible with D2 receptor expression
[44,45]. It is worth noting that a very small fraction of GFP-expressing ChAT-positive neurons showed
an increased F/I slope (Figure 3I, F/I slope, N=2/22, Supplementary Table 1) without changes in
rheobase. Such behavior suggests enhanced excitability following dopamine administration,
consistent with the expression of D1 receptors[23,46]. We also tested if dopamine influences the
synaptic integration of DVC neurons by injecting a train of 5 EPSP-like waveform at 20Hz (α-
EPSPs) into the soma via the patch pipette. We found that, after bath application of dopamine,
the three populations showed different alterations of synaptic integration, with responsive GFP-
negative neurons showing the most significant reduction in temporal summation (Figure 3K,
Supplementary Table 1). Overall, these findings indicate that both ChAT-positive and ChAT-
negative neurons in the DVC respond to dopamine, exhibiting changes in intrinsic excitability,
firing patterns, and synaptic integration properties.
172
Stimulation of midbrain dopamine neurons induces dopamine release in the DVC.
We next tested the hypothesis that stimulation of midbrain dopamine neurons causes
dopamine release in the DVC. We injected AAV2-hSyn-DIO-hM3D(Gq)-mCherry (encoding
excitatory Gq-coupled DREADDs) into the midbrain of DAT-Cre mice. Three weeks post-injection,
the Cre-mediated expression of Gq-DREADD-mCherry was verified in the somatic (midbrain) and
terminal regions (striatum and DVC) (Figure 4C, Supplementary Figure 3).
In a separate cohort, we injected the same Gq-DREADD-encoding AAV into the midbrain and
AAV9-hSyn-dLight1.3b into the DVC, leading to the expression of the high-affinity dopamine
sensor dLight1.3b[47] in the DVC (Figure 4A-B). Three weeks later, 250 µm coronal slices of the
DVC were imaged with epifluorescence microscopy. For each experimental condition, pre-drug
fluorescence signals (-30 seconds → 0 seconds) were subtracted from the signal peak after drug
application (30 seconds) (Figure 4D) (Materials and Methods). We observed significantly
increased dLight1.3b fluorescence after clozapine-N-oxide (CNO) application (Figure 4E),
indicating that dopamine is released from Gq-DREADD-expressing midbrain-originating
dopamine terminals in the DVC. DVC sections were also imaged at baseline and after perfusion
with either 10 µM methamphetamine (Figure 4F), a well-characterized stimulator of dopamine
release[48-50], or 0.1 µM dopamine (Figure 4G) - another positive control. Both treatments
significantly increased dLight1.3b fluorescence, confirming the specificity of dopamine release
from dopaminergic terminals in the DVC. These findings support the conclusion that the activation
of midbrain dopamine neurons induces dopamine release in the DVC, and suggests that
dopaminergic projections from the midbrain regulate DVC activity.
194
Anterograde multi-transsynaptic tracing reveals midbrain-to-spleen connectivity via celiac
ganglion.
The celiac ganglion, which receives projections from the DVC, acts as a relay to peripheral
organs, including the liver, bone marrow and spleen[51-55]. Vagal afferents via the celiac ganglion
regulate splenic immune functions[33,56-59]. Decades of research support the existence of the celiac
ganglia-DVC circuit[60,61], and with the midbrain dopamine neuron-to-DVC projections identified in
this study, we propose a multi-synaptic link between midbrain dopamine neurons and the spleen.
We used the Ai9 TdTomato Cre-reporter line[62] and performed midbrain injections of pENN-AAV1-
hSyn-Cre-WPRE-hGH to transduce Cre in an anterograde multi-transsynaptic manner[63] (Figure

5A). Three weeks later, TdTomato$^+$ cell bodies were observed in the DVC (Fig 5C). We also
observed TdTomato expression in both the celiac ganglion and spleen (Figure 5D-E,
Supplementary Figure 5). TdTomato$^+$ expression in the spleen surrounded T-cell zones, identified
by CD3$^+$ T-cells (Figure 5E, Supplementary Figure 5). As that the spleen contains few neuronal
cell bodies[64], the TdTomato detected in the spleen likely reflects neuronal fibers innervating the
spleen from the celiac ganglion, reinforcing the multi-synaptic midbrain-DVC-celiac ganglion-
spleen projection pathway. Although these results suggest a midbrain-DVC-celiac ganglion-
spleen pathway, they do not exclude the possibility that other direct or indirect connections exist
which may also influence peripheral immunity[65,66]. DVC to the celiac ganglion connections likely
reflect a composite of efferent (CNS-to-periphery) and afferent (periphery-to-CNS) activity, and
would involve sympathetic spinal inputs to the celiac ganglion. In this study, our experiments are
designed to interrogate the efferent arm of this pathway, namely the DVC → celiac ganglia →
spleen axis. In addition, the efferent modulation within a midbrain–DVC–celiac ganglion–spleen
circuit is likely to be accompanied by feedback through afferent pathways back to the CNS.
Nevertheless, the midbrain-DVC-celiac ganglion-spleen pathway offers one mechanism by which
midbrain dopamine neuron activity may influence splenic immune function. To investigate this
possibility, we developed a stepwise experimental approach to assess 1) whether there is
increased DVC neuronal activity when midbrain dopamine neurons are activated; 2) whether
celiac ganglion neurons exhibit increased activity under the same conditions; 3) whether splenic
immune cell populations are altered during dopamine neuron activation, and 4) whether
perturbation of the midbrain-DVC-celiac ganglion-spleen pathway via vagus nerve transection
abrogates these effects. Following this stepwise plan, we sought to investigate the functional
impact of midbrain dopamine neuron activation at each junction of this circuit.
227
Chemogenetic stimulation of midbrain dopamine neurons increases cFos expression in
the DVC and celiac ganglion.
Thus far, we have shown: 1) that midbrain dopamine neurons mono-synaptically project to
and release dopamine in the DVC; 2) that DVC neurons can respond to dopaminergic input, and
3) that multi-synaptic projections exist from the midbrain to the spleen via a midbrain-DVC-celiac
ganglion-spleen circuit. This led us to ask whether activation of midbrain dopamine neurons
increase neuronal activity along the midbrain-DVC-celiac ganglion axis. Immunostaining for cFos
expression, an immediate-early gene induced by neuronal activity that reaches peak expression
within 90 minutes after neuronal activation[67-70], is a reliable measure of increased neuronal activity
along the midbrain-DVC-celiac ganglion axis. Therefore, we next quantified cFos expression in
the DVC and celiac ganglion as a proxy indication of neuronal activity. We injected AAV2-hSyn-
DIO-hM3D(Gq)-mCherry or AAV2-hSyn-DIO-mCherry (control vector) into the midbrain of DAT-
Cre mice. The same animals also received either a cervical vagotomy, or sham surgery targeting
the cervical region. Four weeks post-injection, the mice received daily intraperitoneal injections of
CNO (1 mg/kg) for 14 days to activate the Gq-DREADD-expressing midbrain dopamine neurons
(Figure 6A). The choice of 1 mg/kg CNO was guided by prior work reporting that this dose does
not elicit reward-seeking behavior or toxicity[71]. In addition, to minimize ligand-related confounds,
all animals, both Gq-DREADD and empty-vector controls, received the same CNO regimen
throughout the experiment. Thus, any direct or metabolite-mediated effects of CNO should be
similar across all experimental and control groups.
Locomotor activity, a reliable readout of dopamine neuron activation, was assessed at baseline
and following CNO injections on Days 1, 7, and 13 (Supplementary Figure 6 and 7; Supplemental
Video 1 and 2). Mice injected with AAV2-hSyn-DIO-hM3D(Gq)-mCherry increased locomotor
activity in response to CNO, confirming activation of midbrain dopamine neurons (Supplementary
Figure 6B & 6C). Consistent with our hypothesis, we observed a robust increase in cFos
expression in the DVC, particularly within the DMV and NTS subregions (Figure 6); increased
DVC cFos expression was present in both sham surgery (Figure 6D) and vagotomized mice

(Figure 6E). This effect was absent in the empty-vector control groups (AAV2-hSyn-DIO-mCherry)
(Figure 6B-E). cFos expression in the DVC was detected in both ChAT⁺ and ChAT⁻ neurons,
consistent with the heterogeneous neuronal populations in this region and its role as a relay hub
that integrates and bidirectionally transmits information between the CNS and peripheral
organs[72,73 74,75]. We did not detect cFos expression in any of amygdala's subregions that are
known to receive inputs from, and send projections to, the midbrain and DVC (Supplemental
Figure 8). The observation that midbrain dopamine neuron activation increased neuronal activity
across multiple DVC subregions, including cholinergic neurons in the DMV and catecholaminergic
neurons in the NTS agrees with prior reports[72,73 74,75], supporting the overall interpretation that
activation of midbrain dopamine neurons influences neuronal activity along the midbrain-DVC
axis, as evidenced by increases in cFos expression.
To further investigate downstream effects of midbrain dopamine neuron activation, we
assessed cFos expression in the celiac ganglia of mice injected with Gq-DREADDs (excitatory
DREADD) or control vector, and either cervical vagotomy or sham cervical surgery, as described
above (Figure 7A). Following 14 days of CNO injection (1mg/kg, *i.p.*), intact celiac ganglia were
dissected, whole-mounted, post-fixed, and immunostained. Consistent with midbrain-DVC-celiac
ganglion circuitry, we observed increased neuronal activity, as evidenced by enhanced cFos
expression in Gq-DREADD-injected mice, compared to the empty vector control (AAV2-hSyn-
DIO-mCherry) (Figure 7B & 7C). Quantification of cells with robust cFos expression showed that,
compared to mice that received the control vector which showed only basal levels of cFos
expression, Gq-DREADD-treated mice receiving CNO revealed a marked increase in cFos+ cells
(Figure 7D).
Since the celiac ganglion receives diverse inputs[53,59,76] and innervates multiple peripheral
organs[53,64,77,78], we considered the possibility that the increased celiac ganglion activity observed
after chemogenetic activation of dopamine neurons could arise form a parallel or indirect pathway.
A straightforward way to test this was to disrupt the proposed DVC–celiac ganglion circuit by
transecting the vagus nerve. Thus we tested whether increased celiac ganglion activity (in the
context of midbrain dopamine neuron activation) persists following cervical vagotomy. An
additional cohort of DAT-Cre mice received midbrain injections of either Gq DREADDs or empty
vector control, followed by a sham surgery or a cervical vagotomy (Figure 7A). After 4 weeks to
allow vial expression and surgical recovery, animals received systemic CNO as described above.
Locomotor activity was assessed to verify dopamine neuron activation in both sham surgery and
vagotomized mice (Supplemental Figures 6 and 7; Supplemental Video 1 and 2). Intact celiac
ganglia were dissected, whole mounted, post-fixed and immunostained. Consistent with our
hypothesis, animals that received both midbrain Gq-DREADD AAV and cervical vagotomy exhibit
no change in celiac ganglion cFos expression relative to empty vector controls that received
cervical vagotomy (Figure 7E-G). These results strongly support the interpretation that midbrain
dopamine neuron activity functionally regulates splenic immunity via a midbrain-DVC-celiac
ganglion-spleen circuitry. Together, these findings demonstrate that midbrain dopamine neuron
activation profoundly influences neuronal activity along the midbrain-DVC axis and downstream
autonomic circuits, highlighting a functional role in regulating peripheral immune responses.
296
Chemogenetic stimulation of midbrain dopamine neurons reduces spleen weight and
naïve CD4⁺ T-cell populations without affecting total cell numbers. Vagotomy abolishes
changes in spleen weight and changes in naïve CD4⁺ T-cells.
As a key immune organ, the spleen plays a central role in orchestrating both innate and
adaptive immune responses, responding to pathogens, immune activation, and even cancer[79-86].
Disruptions in splenic immune cell populations or function due to altered CNS dopamine
neurotransmission could have profound immunological consequences. Such behavior is
particularly relevant in diseases characterized by dysregulated dopamine neuron activity and
existing peripheral immune disturbances, potentially influenced by CNS dopamine signaling. For

instance, Parkinson's disease (PD), conventionally defined by the loss of midbrain dopamine
neurons and motor deficits, also exhibits peripheral immune manifestations, including changes in
the phenotype of immune cell populations, gut dysfunction, and systemic inflammation[3-5,87], with
PD patients exhibiting impaired CD4$^+$ T-cell migration, accompanied by an increase total CD4$^+$ T-
cell populations[87-89]. Conversely, psychostimulant use, which elevates brain dopamine levels,
leads to opposing T-cell responses, including a reduction in total T-cells, and an increased
propensity towards inflammatory T-cell phenotypes[90-94]. These observations suggest that
changes in CNS dopamine transmission, whether through loss, as in PD, or through augmentation,
as seen with psychostimulants, can influence immune function[87-89, 90-94]. The interplay between
midbrain dopamine neuron activity and peripheral immune responses underscores a relationship
where CNS signaling impacts systemic immunity[6,95-98]. To probe whether the midbrain-DVC-celiac
ganglion-spleen circuit modulates splenic immunity, we assessed splenic immune cell
populations following the activation of midbrain dopamine neurons. The proposed midbrain-DVC-
celiac ganglion-spleen circuit relies on vagal projections from DVC to spleen via the celiac
ganglion[24,25]; therefore, we reasoned that the effects of midbrain dopamine neuron activation on
splenic immunity would be abolished if the vagus nerve were transected. Previous work has
shown that cervical, but not subdiaphragmatic, vagotomy preserves connections between the
celiac ganglion and spleen while disrupting vagal efferents to the celiac ganglion[99,100].
Therefore, we employed a four-pronged experimental design, in which DAT-cre mice
received an intracranial AAV injection targeting the midbrain to express either Cre-dependent Gq-
DREADDs or a control vector expressing a fluorescent reporter, followed by either a left cervical
vagus transection (vagotomy) or a sham surgery. After four weeks to allow viral expression, we
administered CNO daily (1 mg/kg, I.P.) for 14 days to stimulate DAT$^+$ midbrain neurons (Figure
8A, Figure 9A). Locomotor activity was assessed at baseline and following CNO injections on
Days 1, 7, and 13 to verify chemogenetic stimulation of dopamine neurons (Supplementary Figure
6 and 7; Supplemental Video 1 and 2). Chemogenetic stimulation of dopamine neurons did not
alter body weight in either sham surgery or vagotomy groups (Figure 8B and Figure 9B),
Firstly, examining the sham surgery group, we found that mice with chemogenetic stimulation
of midbrain dopamine neurons showed a significant reduction in spleen weight relative to empty
vector controls (Figure 8C), despite no change in body weight, reinforcing the idea that midbrain
dopamine signaling influences splenic immune function. These data are consistent with previous
studies in DAT-knockout (DAT-KO) mice, marked by elevated CNS dopamine, which is
associated with spleen hypoplasia, and reduced spleen weight[98,101]. Next, in the sham surgery
group, we performed flow cytometry to assess splenic adaptive and innate immune subsets.
These analyses included T-cell populations, including CD4$^+$, CD8$^+$, effector memory, Central
memory, and naïve T-cells (Figure 8C-H, Supplemental Figure 6); B-cell populations, including
total B-cells, immature and maturing B-cells (Supplemental Figures 7 and 12); NK-cell populations,
including total NK-cells, immature and mature NK-cells (Supplemental Figures 8 and 12); and
splenic macrophages (Supplemental Figures 10 and 14). While total numbers of CD4$^+$ T-cells,
effector memory T-cells (CD4$^+$ CD44$^+$ CD62L$^-$) and central memory T-cells (CD4$^+$ CD44$^+$ CD62L$^+$)
were unchanged between groups (Figure 8 F-G; J-L), naïve CD4$^+$ T-cells (CD4$^+$ CD44$^-$ CD62L$^+$)
were significantly reduced in mice that received chemogenetic stimulation of midbrain dopamine
neurons, compared to empty vector control (Figure 8M). No significant changes were observed
in CD8$^+$ T-cell populations (Figure 8 H-I; N-Q).
To determine the involvement of the vagus nerve in modulating splenic immunity in the
proposed midbrain-DVC-celiac ganglion-spleen when dopamine neurons are chemogenetically
activated, we examined the groups that received midbrain injection of Gq-DREADD AAV or empty
vector control followed by vagotomy (Figure 9). In contrast to the sham surgery groups, in
vagotomized mice, chemogenetic stimulation of midbrain dopamine neurons show no change in
spleen weight compared to empty vector controls (Figure 9C). Immunophenotyping of splenic
immune cells in vagotomized mice showed that total CD4+ T-cells, effector memory cells (CD4+

CD44$^+$ CD62L$^-$), and central memory cells are unchanged (CD4$^+$ CD44$^+$ CD62L$^+$) (Figure 9J-L),
similar to sham controls. Importantly, diverging from sham surgery groups, we found that
vagotomy abolishes the reduction in naïve CD4$^+$ T-cells (CD4$^+$ CD44$^-$ CD62L$^+$) in mice with
chemogenetic stimulation of midbrain dopamine neurons, compared to empty vector controls
(Figure 9 F-G, Figure 9M). No changes were observed in CD8$^+$ T-cell populations in both groups
of vagotomized mice (Figure 9H-I, Figure 9N-Q). Finally, in both sham surgery and vagotomy
groups, we observed no changes in B-cell subtypes (Supplementary Figure 9), NK-cell subtypes
(Supplementary Figure 10) or splenic macrophages (Supplementary Figure 11).
Collectively, these findings suggest a brain-body circuit for specialized modulation of specific
splenic immune cell populations by midbrain dopamine neurons, rather than global suppression,
shedding light on mechanisms by which altered dopamine neurotransmission influences
peripheral immune function. Importantly, our results align with previous studies demonstrating
that top-down regulation of splenic T-cells can affect peripheral immune manifestations such as
pain[96].
Through the celiac ganglia, the spleen is innervated by both cholinergic[29-31] and
catecholaminergic fibers[32-35], both of which can modulate spleen function[26], immune cell
populations, and, by extension, influence the peripheral immune landscape[102-104]. Therefore,
dopaminergic projections from midbrain regulate DVC activity, that via the vagus nerve and celiac
ganglion with downstream adrenergic and cholinergic fibers, could modulate splenic immunity[102-
104]. This pathway likely underpins the immunophenotypic changes observed following midbrain
dopamine neuron activation. These are consistent with the central finding of the current study that
activation of midbrain dopamine neurons induces functional immunological changes in the spleen.
It should be noted that the DVC contains multiple vagal nuclei upstream of the periphery and
projects to numerous peripheral organs via the vagal efferent pathway. This raises the possibility
that the changes in splenic immunity observed upon chemogenetic activation of midbrain
dopamine neurons may also reflect alterations in the activity of other peripheral organs outside
the proposed midbrain-DVC-celiac ganglion-spleen circuit[38,51,75,105].
384
Concluding remarks
This study provides strong evidence for a functional connection between the CNS and
peripheral immunity through midbrain dopamine signaling. Our findings suggest that midbrain
dopamine neuron activation not only affects splenic weight and T-cell populations but also
underscores a broader role for dopamine signaling in regulating peripheral immunity. Disruptions
in brain-body communication have been implicated in various diseases, including
neurodegenerative disorders[3], cardiovascular disorders[106-108], and metabolic syndromes[109,110].
Numerous studies have shown that altered dopamine neurotransmission is linked to changes in
peripheral immunity, such as during psychostimulant use and with degeneration of dopamine
neurons[6,87,97,98,111-113]. However, prior evidence for the circuitry between midbrain dopamine
neurons and the immune system was scant. In the present study, we now demonstrate the
existence of a multi-synaptic brain-to-periphery pathway, linking midbrain dopamine neurons to
the DVC and, via the celiac ganglia, to the spleen[26]. Vagal fibers are known to innervate the celiac
ganglion[25,99,100,114]; in turn, projections from the celiac ganglion innervate the spleen[114]. Given the
evidence linking the vagus to the spleen, prior studies have focused on the cholinergic anti-
inflammatory pathway that can modulate inflammation via vagal transmission[26-28], further
supporting the importance these of brain-body connection.
While our findings demonstrate that midbrain dopamine neuron activity, acting through a
midbrain–DVC–celiac ganglion pathway, modulates splenic immunity, our data do not exclude
contributions from parallel routes within the vagal–spleen axis. Indeed, the vagus nerve sends
and receives projections from the gastrointestinal tract[36,115-117], and gut–spleen communication
via autonomic and immune pathways are well documented[104,118,119]. These observations raise the
possibility that a midbrain–DVC–gut–spleen circuit could contribute to midbrain dopamine

neuron–dependent regulation of peripheral immunity. Supporting this idea, Chang et al. (2024)
identified a stress-sensitive circuit in which vagal projections regulate gut microbiome composition,
and altered gut microbiota influence both splenic immunity and brain function[116,120-122]. Thus, a
brain–gut–spleen axis represents a plausible complementary pathway for dopamine neuron
modulation of peripheral immunity. Finally, although the present study focuses on the efferent
limb of the midbrain–DVC–celiac ganglion–spleen pathway, our data do not preclude afferent
signaling from the periphery back to the CNS and the existence of other reciprocal, bidirectional
circuits between these compartments[123-125].
In the present study, we expand previous findings demonstrating that midbrain-originating
dopamine signaling emerges as a key modulator of peripheral immunity, with the observed
reduction in spleen weight and alterations in T-cell populations, particularly the decrease in naïve
CD4$^+$ T-cells, following chemogenetic activation of midbrain dopamine neurons. These results
emphasize the influence of midbrain dopamine neurons on adaptive immune functions, with broad
implications for conditions where immune dysregulation and chronic inflammation are central to
disease manifestation and progression. Furthermore, dopamine-driven modulation of splenic
immune cell populations suggests that midbrain dopamine neuron activity may extend regulatory
effects to immune responses throughout the body. A multitude of diseases, from autoimmunity to
cancer, are treated using immunomodulatory drugs, modulating the immune response as a
treatment method. Given that dopamine neuron activity affects splenic T-cell populations, we
pondered whether peripheral immunomodulation may also change CNS dopamine
neurotransmission, via the same circuit. It is well known that during infection and inflammation,
patients experience a lack of motivation, social withdrawal, and anhedonia[12,126-128]. Therefore, it
is reasonable to postulate that peripheral immunomodulation, via drugs that target the immune
system, may function in a feedback/feedforward loop to modulate CNS dopamine
neurotransmission. Future studies employing clinically relevant immunomodulatory drugs, or
experimental methods such as chemogenetic manipulation of the spleen, will help clarify this
compelling question.
Finally, there remains the fundamental question: why have mammals evolved a biological
circuit that links midbrain dopamine neurons to the peripheral immune system? Dopamine
neurotransmission is fundamentally involved in socially motivated behaviors, such as social
bonding[129-133]. Such behaviors are conserved among numerous species. Indeed, in contexts of
anhedonia, such as during depression, peripheral inflammation and immune dysfunction is
increased[134,135]. In this respect, midbrain dopamine regulation of the peripheral immune system
may facilitate immunity through socialization and bonding reward mechanisms. This idea is
supported by literature showing that activation of midbrain dopamine neurons helps to resolve
infection[95], to improve anti-tumor immunity[96], and to enhance stress resilience in individuals
experiencing chronic stress[136,137], emphasizing the function of dopaminergic signaling in
peripheral immunity along the midbrain-spleen axis. Our study also highlights a critical role for
midbrain dopamine neurons beyond CNS neurotransmission, positioning them as an important
regulator of peripheral immunity. Such considerations strengthen the concept of brain-body
communication, linking neuroimmune interactions to the onset and progression of systemic
diseases and offering promising leads for therapeutic interventions that target these circuits.
450
Limitations of the methodology and systems used in this study
While our approach yielded significant findings, several methodological limitations warrant
consideration. First, the viral constructs employed in our study reliably transduced the entire
dopaminergic midbrain, encompassing both the substantia nigra pars compacta (SNc) and the
ventral tegmental area (VTA), precluding efforts to distinguish the specific roles of SNc versus
VTA neurons in the regulation of the peripheral immune system. Future studies utilizing more
refined viral constructs or intersectional genetics will enable dissection these distinct neuronal
populations and likely would uncover even more robust outcomes. Second, our data suggest that

midbrain dopamine neurons send projections throughout the DVC, including AP, NTS, and DMV,
thus preventing us from determining whether a specific nucleus or neuronal subtype within the
DVC contributes predominantly to splenic immunoregulation by midbrain dopamine signaling.
Future studies employing targeted inhibition of individual DVC nuclei, during dopamine neuron
activation will help delineate the specific roles of each nucleus and individual cell types in splenic
immune modulation. Third, literature suggests that unilateral vagus nerve stimulation may induce
more pronounced changes in the peripheral immune system[138,139]; however, our bilateral
approach does not allow us to assess whether the left or right vagal pathway exerts greater
influence on midbrain dopamine neuron regulation of splenic immunity. Addressing this limitation
in future studies will involve investigating the laterality of midbrain-spleen immunoregulation,
which could provide deeper insights into the lateral specificity of this pathway. Such findings could
have important implications for experimental designs and therapeutic applications based on these
mechanisms. While we have taken a significant step toward understanding the role of midbrain
dopamine signaling in regulating peripheral immunity, addressing these constraints should
provide a fuller perspective on the neuroimmunological implications of the midbrain-DVC-celiac
ganglion-spleen pathway.
475
476
477 **Materials and Methods:**
478
Animals
We used the following strains of mice, both males and females, on a C57BL/6 background:
B6.Cg-*Gt(ROSA)26Sor*$^{tm9(CAG-tdTomato)Hze}$/J (Jax strain #007909)[140], *Slc6a3*$^{tm1(cre)Xz}$/J (Jax strain
#020080), B6.Cg-Tg(RP23-268L19-EGFP)2Mik/J (Jax strain#007902). Adult male and female
mice were between 8-16 weeks of age, on a 12-hour day/night cycle with ad libitum access to
food and water. Animals were bred in-house at the University of Florida or at the University of
Texas at San Antonio, following approved procedures by the University of Florida IACUC (IACUC
202400000184) or by the University of Texas at San Antonio IACUC (IACUC MU096).
487
Stereotactic surgeries and viral injections
Mice were anesthetized using gas anesthetic (3% isoflurane mixed with oxygen). After
head shaving, mice were head-fixed in a stereotactic frame (Kopf Instruments). Following
sterilization of the surgical site, an incision was made, skull exposed and leveled from bregma to
lambda, and burr holes drilled at the surgical coordinates (Bregma: +0.4ML, -3.4AP, -4.4DV).
Viruses were injected via a pulled glass capillary at a rate of 5nL every 5 seconds, until the
injection was complete (Drummond Nanoject III). The injector was left in place for 5 minutes to
allow spread prior to retraction. Scalp skin was sutured using 3 interrupted sutures (5.0
polypropylene sutures). The virus was allowed to express for several weeks prior to experiments
as indicated throughout the Results. Viral vectors are provided in **Table 1**.
498
Vagotomy or sham procedure
Mice were placed in a supine position on a sterile drape, maintained on isoflurane at 0.8-
1.2%. Following sterilization of the neck surgical site, a vertical incision was made 3mm caudal to
the sternal notch. The inferior border of the thyroid tissue, a distinct V-shaped line at the cervico-
sternal junction, was identified. By blunt dissection, submaxillary salivary glands were exposed
and separated through the midline fascial plane to expose the trachea. The incision was kept
open using a micro-retractor. The left cervical vagus nerve, located within a neurovascular bundle,
was readily identified by the pulsation of the carotid artery. The carotid sheath was isolated from
the connective tissue using pickup forceps. The nerve was gently dissected from the sheath,
taking care not to damage the vessels or tug on the nerve. Once isolated, a small section of sterile
suture was used to suspend the nerve from surrounding tissue and severed using fine surgical

scissors (vagotomy) or simply manipulated and then restored to the sheath (sham procedure).
The incision site was cleaned and sutured with 3-5 interrupted sutures.
Image analysis
Fluorescence intensity of EYFP+ processes in the DVC were calculated as total
fluorescence intensity on background-corrected images in the channel corresponding to EYFP in
NIS Elements software. Units are expressed as arbitrary fluorescence units per square micron.
mRuby+ puncta were quantified using open-source software Cell Profiler, using an analysis
pipeline adapted from Cell Profiler "Speckles Quantification". cFos+ cells in the DVC and celiac
ganglion were assessed by setting a threshold for cFos intensity of greater than or equal to 61%
of the fluorescence intensity of the nuclear counterstain (DAPI). Cells which were positive for cFos
were manually counted and are presented as number of cFos+ cells per square micron.
521
Electrophysiology
As described in previous studies[141-147], after 3-4 weeks from the viral injection, mice were
anesthetized with isoflurane and decapitated. Coronal slices (300 µm) containing the area of
interest were obtained on a vibratome (VT1200S, Leica) in a chilled cutting solution containing
100 mM choline chloride, 25 mM $NaHCO_3$, 25 mM D-glucose, 11.6 mM sodium ascorbate, 7 mM
$MgSO_4$ 3.1 mM sodium pyruvate, 2.5 mM KCl, 1.25 mM NaH_2PO_4, 0.5 mM $CaCl_2$. Slices were
then incubated in oxygenated artificial cerebrospinal fluid (ACSF) in a submerged chamber at 35–
37°C for 30 min and then at room temperature (21–25°C), until recordings were performed. ACSF
contained 126 mM NaCl, 26 mM $NaHCO_3$, 10 mM D-glucose, 2.5 mM KCl, 2 mM $CaCl_2$, 1.25 mM
NaH_2PO_4, 1 mM $MgCl_2$; osmolarity was ~290 Osm/L.
Whole-cell recordings were performed in 31–33°C ACSF. Thin-walled borosilicate glass
pipettes (Warner Instruments) were pulled on a vertical pipette puller (PC-10, Narishige), resulting
in a typical range of 3–5 MΩ resistance. Intrinsic properties were recorded in the current-clamp
configuration using a potassium-based intracellular solution which contained 120 mM potassium
gluconate, 20 mM KCl, 10 mM HEPES, 10 mM phosphocreatine, 4 mM ATP, 0.3 mM GTP, 0.2
mM EGTA, and 0.3–0.5% biocytin (mg/mL). Signals were sampled at 10 kHz and filtered (lowpass
filter) at 4 kHz, in the presence of excitation and inhibition blockers: CPP (5 µM; Tocris Bioscience),
NBQX (10 µM; Abcam), and gabazine (25 µM; Abcam). Hardware control and data acquisition
were performed with the Matlab-based Ephus package (Suter et al., 2010). The resting membrane
potential (Vm) was calculated in current clamp mode (I = 0) immediately after breaking in. Series
(R_s) and input resistance (R_{in}) were calculated in voltage-clamp mode (V_{hold} = −70 mV) by giving
a −5 mV step, which resulted in transient current responses. R_s was determined by dividing the
voltage step amplitude by the peak of the capacitive current generated by the voltage step. The
difference between baseline and steady-state hyperpolarized current (ΔI) was used to calculate
Rin using the following formula: R_{in} = −5 mV/ΔI − R_s. Subthreshold and suprathreshold membrane
responses in current-clamp configuration were elicited by injecting −100 to +500 pA in 50-pA
increments at V_{hold} = −70 mV. The first resulting AP at rheobase (the minimal current of infinite
duration - experimentally limited to 1 s, as required to generate an AP, from here on current
threshold, or I threshold) was analyzed for AP half-width, peak, height, Voltage threshold. The
adaptation ratio was measured at the current step that gave the closest APs firing rate to 20 Hz,
calculated dividing the second instantaneous frequency by the last. The f/I graphs were generated
by calculating the firing frequency at each positive current step administered to the cell while
holding it at -70 mV. Alpha synapse was calculated by measuring the peak of five consecutive 5
msec EPSP-like steps at 20Hz and normalizing on the amplitude of the first peak.
556
557 *Ex vivo* **dLight1.3b recording**
558 **Preparation of acute DVC slices:** The mice were perfused with ice cold sucrose-sodium-
559 replacement solution containing 250 mM sucrose, 26 mM $NaHCO_3$, 2 mM KCl, 1.2 mM NaH_2PO_4,
560 11 mM dextrose, 7 mM $MgCl_2$, and 0.5 mM $CaCl_2$. Mice were decapitated and their brains were

promptly removed. For slicing, the brain was cut rostrally to the cerebellum in the coronal plane to create a flat surface for attachment to the cutting stage. Coronal brainstem sections containing the DVC were sectioned at 250 μm (VT1200, Leica). The slices were transferred to a recovery chamber containing oxygenated ACSF at 34°C. The ACSF composition included: 126 mM NaCl, 2.5 mM KCl, 2 mM $CaCl_2$, 26 mM $NaHCO_3$, 1.25 mM NaH_2PO_4, 2 mM $MgSO_4$, and 10 dextrose, equilibrated with 95% O_2 and 5% CO_2. The osmolarity for both the sucrose-sodium-replacement and ACSF was adjusted to 300-310 mOsm and the pH was set to 7.35-7.4. Sections remained in the recovery chamber for at least 30 minutes at 34°C until used for subsequent ex-vivo dLight1.3b recording. Sections containing the DVC were identified using the AP and hypoglossal nucleus as anatomical landmarks.

Recording: The slices were then transferred to a low-profile open diamond bath imaging chamber (RC-26GLP, Warner Instruments, Hamden, CT, USA) and perfused with ACSF at 36-37°C. Sections were imaged using a 40x water-immersion objective on an Eclipse FN1 upright microscope (Nikon Instruments, Inc., Melville, NY, USA), at 25-30fps with no delay between frames. The microscope was equipped with a Spectra X light engine (Lumencor, Beaverton, OR, USA) and a 470/24 nm solid-state illumination source. Images were captured with a Hamamatsu BT Fusion camera (Hamamatsu). Gq-DREADD receptors were activated via bath perfusion of 10 μM clozapine-N-oxide (CNO) in ACSF.

Signal bleach correction: Raw fluorescent intensity values were extracted using ImageJ. A photobleaching curve was generated by fitting the baseline florescent signal using a non-linear least regression with a biexponential model. Traces were then corrected and normalized using MATLAB. Codes used for bleach correction and normalization were generated by Leah T. Phan (co-author), and are available at:
https://github.com/LTP08/Khosh_CommonCodes/blob/899d7403fb39b3d7d41978a11e6284de1787d55e/biexponential_photobleachcorrection_bulksignal.mlx

dF/F dlight1.3b signal calculation: Baseline values, the median value of the 30 second baseline recording (pre-drug) were subtracted from the peak signal 30 seconds following drug perfusion (~1 minute after recording start). The resulting fluorescence value was divided by the baseline fluorescence value to yield a normalized dF/F value for comparison across experiments.

Dissection of the sympathetic splanchnic-celiac-superior mesenteric ganglion complex (SCSMG)

Mice were deeply anesthetized with isoflurane, and transcardially perfused with PBS, followed by 4% paraformaldehyde (PFA). The sympathetic ganglion complex was dissected following Trevizan-Baú et al. (2024)[59]. Briefly, mice were positioned in dorsal recumbency, and the abdominal wall was opened (in the Linea Alba). With fine forceps and Noyes Iris scissors, the sympathetic ganglion complex was dissected and collected under a stereo microscope. The complex was then placed onto a Sylgard-coated dissection dish with PBS and pinned down on the dish through connective tissue and nerves, not through the ganglia. Then, after removing any additional connective tissue or fat, the sympathetic ganglion complex was post-fixed for 24 hours before using for immunostaining and later imaging.

Brain tissue and spleen immunofluorescence and imaging

Animals were terminally anesthetized with isoflurane, and transcardiac perfused with 1x PBS followed by 4% PFA dissolved in PBS (pH 7.4). Brains were removed and post-fixed in 4% PFA for 24 hours at 4°C. Spleens were removed prior to perfusion and post-fixed in 4% PFA for 24 hours at 4°C. Thereafter, sections were cut at 40um thickness at room temperature on a vibrating microtome (Leica VT1000). Free-floating sections were blocked at 37°C for 1 hour in 0.3% Triton X-100 in PBS, containing 5% normal goat serum or fetal bovine serum. The slices were incubated with primary antibody overnight in 0.3% TritonX-100 in PBS, containing 5% normal goat serum or fetal bovine serum. Following washes, slices were incubated with

612 corresponding fluorophore-conjugated secondaries for 1 hour at room temperature, washed and
613 mounted (Fluoromount, Invitrogen). Antibodies and concentrations are given in **Table 2**.
614
Flow cytometry
Following terminal isoflurane anesthesia, the spleen was removed and placed in PBS on
ice. Sterile tissue forceps were used to remove the spleen, place it on a slide and disrupt it to
produce a splenocyte suspension. Slides were washed using 3 volumes of PBS to collect
splenocyte suspension, followed by filtration through a 40 µm cell strainer placed over a 50mL
conical tube. Following centrifugation (300 x g, 5 min), the cell pellet was resuspended in 70%
Percoll density medium, and then underlaid below 37% Percoll. Centrifugation for 30 min at 500
x g with brakes and acceleration set to low results in a suspension of splenocytes at the interface
between 37% and 70% Percoll. Cells were collected, washed twice with excess PBS, and then
counted prior to the density being adjusted to 10,000 cells per microliter. Flow cytometry staining
was conducted using 1 million cells suspended in 100uL of PBS. Antibodies and concentrations
are given in **Table 3**.
627
628
629 **Table 1: Adeno-associated viruses**

AAV	Purpose	Vendor	Catalog #	Concentration	Volume injected
AAV2-Ef1a-DIO-EYFP	Tracing	Addgene	27056	1×10^{13} vg/mL	200nL
AAV1-hSyn-Flex-mGFP-2A-Synaptophysin-mRuby	Tracing	Addgene	71760	7×10^{12} vg/mL	100nL
pENN-AAV1-hSyn-Cre-WPRE-hGH	Tracing	Addgene	105553	7×10^{12} vg/mL	100nL
AAV9-syn-dLight1.3b	Dopamine sensor	Addgene	135762	1×10^{13} vg/mL	50nL
AAV2-hSyn-DIO-hM3D(Gq)-mCherry	Excitatory DREADD	Addgene	44361	6×10^{12} vg/mL	200nL
AAV2-hSyn-DIO-mCherry	Empty Vector control	Addgene	50459	6×10^{12} vg/mL	200nL

630
631 **Table 2: Primary and secondary antibodies for immunofluorescence**

Specificity	Clone/Species	Conjugate	Vendor	Catalog Number	Purpose	Dilution
DAT	Polyclonal/Rat	N/A	EMD Millipore Corp.	MAB369	IHC	1:500
TH	Polyclonal/Rabbit Polyclonal/Chicken	N/A	Encor	RPCA-TH CPCA-TH	IHC	1:500
CD3	Polyclonal/Rat	N/A	BioLegend	100201	IHC	1:1000

ChAT	Polyclonal/Goat	N/A	EMD Millipore Corp.	AB144P	IHC	1:200
GFP	Monoclonal/Mouse	N/A	Invitrogen	A11122	IHC	1:500
mCherry	Monoclonal/Mouse	N/A	Encor	MPCA-mCherry	IHC	1:500
cFos	Polyclonal/Rabbit	N/A	Encor	RPCA-c-FOS	IHC	1:500
B220	Polyclonal/Rat	N/A	BioLegend	103202	IHC	1:500
Rat	Polyclonal/Goat	Alexafluor-488 Alexafluor-647	Invitrogen	A11006 A21247	IHC	1:500
Rabbit	Polyclonal/Goat	Alexafluor-488 Alexafluor-568	Invitrogen	A11034 A11011	IHC	1:500
Guinea Pig	Polyclonal/Goat	Alexafluor-488	Invitrogen	A11073	IHC	1:500
Mouse	Polyclonal/Goat	Alexafluor-647	Invitrogen	A21236	IHC	1:500
Chicken	IgY	Alexafluor-647	Invitrogen	A21449	IHC	1:1000
Goat	Polyclonal/Donkey	Alexafluor-647	Invitrogen	A21447	IHC	1:500

632
633
634
635
636 **Table 3: Antibodies for flow cytometry**
637

Specificity	Clone/Species	Conjugate	Vendor	Catalog #	Purpose	Dilution
CD3	17A2/Rat	Pac Blue	BioLegend	100214	FC	1:50
CD4	GK1.5/Rat	AF700	BioLegend	100430	FC	1:100
CD8a	QA17A07/Mouse	Spark 538	BioLegend	155020	FC	1:50
CD127	S18006K/Rat	PE/Fire 640	BioLegend	158213	FC	1:100
KLRG1	2F1/KLRG1/Syrian Hamster	PE/Fire 810	BioLegend	138437	FC	1:200
CD19	6D5/Rat	BV605	BioLegend	115540	FC	1:50
CD45	30-F11/Rat	Spark Blue 574	BioLegend	103184	FC	1:200

NK1.1	S17016D/Mouse	Fire 700	BioLegend	156527	FC	1:100
CD335	29A1.4/Rat	PerCP Fire 806	BioLegend	137650	FC	1:50
CD11b	M1-70/Rat	FITC	BioLegend	101206	FC	1:200
F4/80	BM8/Rat	Spark NIR 685	BioLegend	123168	FC	1:50
CD44	IM7/Rat	Spark Yg 594	BioLegend	103078	FC	1:100
CD62L	MEL-14/Rat	BV711	BioLegend	104445	FC	1:200
IgM	RMM-1/Rat	PE	BioLegend	406507	FC	1:200
IgD	11-26c.2a/Rat	PerCP Cy5.5	BioLegend	405709	FC	1:200

638
639
640

Funding: This work was supported by NIH grants to Habibeh Khoshbouei (H.K.), Adithya
Gopinath (A.G), and Matthew J. Farrer (M.J.F.): R01NS071122-07A1 (to H.K.), R01DA026947-
10, National Institutes of Health Office of the Director Grant 1S10OD020026-01 (to H.K.)
R01DA058143-02 (to H.K.), R21NS133384-01 (to H.K.), K99DA063743 (to A.G.),
1R21NS136890-01A1 (to M.J.F.), Evelyn F. and William L. McKnight Brain Institute's Gator
Neuroscholars Program (to A.G. and P.T.B.), and the Karen Toffler Charitable Trust (to A.G.).
647
648 **Acknowledgements**: We acknowledge assistance from Kaitlyn N. Kiel and technical assistance
649 from the University of Florida Center for Immunology and Transplantation.

650
651

652
Figure 1. Anterograde mapping of midbrain dopaminergic neurons confirms dopaminergic
projections to the dorsal vagal complex. Panel A: Anatomic location of injection of AAV2-Ef1a-
DIO-EYFP into the midbrain of DAT[IRES]-Cre mice, followed by three-week expression period.
Panel B: EYFP expression in positive control region, the median forebrain bundle (MFB). **Panel**
C: EYFP expression at the injection site in the midbrain. Note also that tyrosine hydroxylase (TH)
co-staining confirmed the fidelity of expression in dopaminergic projections from the midbrain in
B and C. **Panel D:** EYFP+ projections surrounding ChAT+ DMV neurons in the DVC, suggesting
monosynaptic projections from the midbrain to DVC. **Panel E:** EYFP intensity per square
millimeter (expressed as arbitrary fluorescence units). *Data representative of n=6 independent*
experiments.

Figure 2. DAT+ midbrain neurons synapse throughout the dorsal vagal complex. Panel A:
Anatomic locations in brains of DAT-Cre mice receiving midbrain injections of AAV1-hSyn-Flex-
mGFP-2A-Synaptophysin-mRuby. Three weeks after midbrain injection, the resulting
fluorescence patterns indicate that midbrain dopamine neurons not only send projections to the
DVC but also make synapses in the DVC. **Panel B:** synaptophysin-mRuby+ puncta in control
regions of striatum. **Panel C:** synaptophysin-mRuby+ puncta at the injection site in the midbrain.
[Panels B and C also show tyrosine hydroxylase (TH) co-staining to confirm expression in
dopaminergic projections from the midbrain.] **Panel D:** Presence of synaptophysin-mRuby+
puncta throughout DVC, identified by presence of choline acetyltransferase+ (ChAT+) neurons.
Panel E: Quantification of synaptophysin-mRuby+ puncta. *Data representative of n=6*
independent experiments.

690
Figure 3. DVC neurons express D1-like and D2-like receptors. Panel A: DVC neurons of
adult Chat-GFP mice were recorded, using whole-cell patch clamp. Left: Anatomic drawing of
brain (sagittal), showing region of interest (green shading). Vertical line represents the coronal
plane(s) collected for patching, as shown in middle panel. Right: Position of the patching pipette
on a GFP-positive and a GFP-negative neuron in the DVC. **Panel B**: GFP-expressing neurons
are tonically active: they spontaneously fire action potentials in resting conditions, both in cell-
attach (top trace) and in whole-cell current-clamp configuration (bottom trace). **Panel C**: GFP-
expressing and GFP-negative neurons display similarities in intrinsic properties such as input
resistance (Ri), membrane time constant (τ). **Panel D**: active properties of GFP-expressing and
GFP-negative neurons in the DVC: $V_{threshold}$, the membrane voltage threshold to generate an
action potential; $I_{threshold}$, the minimal current step to elicit an action potential (rheobase); V_{Height} the
total amplitude of the action potential (AP); action potential halfwidth ($AP_{halfwidth}$ the width of the
first action potential at threshold, measured at half the peak amplitude); f/I slope, the relationship
between AP frequency and increasing depolarizing currents applied; SFA, spike frequency
adaptation, measured as the ratio between the second and last instantaneous frequency of a train

of action potentials upon depolarizing currents. *** = p<0.005, **** = p<0.001 (two tailed t test).
Panel E: Summary boxplots of passive properties of GFP-positive (GFP+, shades of green, N=22)
and GFP-negative (GFP-, shades of red, N=19) neurons in the DVC, before ("Pre", lighter circles)
and after ("Post", darker circles) bath administration of 50 µM dopamine. **Panel F:** Representative
traces from the three populations identified, showing the membrane responses to increasing
depolarizing current steps (from 0 pA to 500 pA in 50 pA steps). The lighter traces were recorded
in baseline conditions, holding the neurons at -70mV in absence of dopamine, while the darker
traces were recorded 4 minutes after bath-application of 50 µM dopamine. **Panel G**: Quantification
of the firing rate (frequency of action potentials, Hz) in response to depolarizing currents (pA) of
each cell type in Panel-E, same color coding: GFP+ neuron before (light green) and after
dopamine application (dark green); responsive GFP- neuron before (magenta) and after
dopamine application (red); unresponsive GFP- neuron before (light blue) and after dopamine
application (dark blue). Arrowheads indicate rheobases. **Panel H**: Quantification of V threshold
and I threshold for the three populations identified, pre (light colored circles) and post bath
application of dopamine (dark colored circles). **Panel I**: Quantification of V height and AP halfwidth
for the three populations identified, pre (light colored circles) and post bath application of
dopamine (dark colored circles). **Panel J:** Quantification of f/I slope and SFA ratio for the three
populations identified, pre (light colored circles) and post bath application of dopamine (dark
colored circles). **Panel K**: Left: three representative alpha-synapse responses from the
populations analyzed pre (lighter traces) and post (darker traces) bath application of dopamine.
Right: population analysis of synaptic integration, expressed as a fold increase on the first of five
EPSP-like inputs before (lighter traces) and after (darker traces) bath application of dopamine,
showing a significant effect on GFP+ and responsive GFP- neurons in the DVC. * = p <0.05; ** =
p<0.01 (two tailed t-test comparing before and after bath application of dopamine).
730

Figure 4. Midbrain dopamine neuron projections expressing excitatory Gq-DREADD receptors in the DVC release dopamine upon stimulation with CNO, and upon stimulation with methamphetamine, indicating functional projections from midbrain DAT+ neurons to DVC. Panel A: Anatomic midbrain locations of DAT-Cre mice receiving AAV2-DIO-hM3D(Gq)-DREADD-mCherry. In the same surgery, mice received AAV9-hSyn-dLight1.3b in the DVC, followed by expression for 3 weeks. **Panel B:** Coronal sections containing DVC (highlighted in orange) were obtained. **Panel C:** Co-staining for ChAT to identify DVC where DREADD expression in dopamine neuron terminals was verified (data representative of *n* = 3 biological replicates). **Panel D:** Live 250um coronal sections containing the DVC were collected and placed in a recovery chamber at 34 degrees C with carbogenation. After allowing for 30 minutes for rest and recovery, sections were transferred to an imaging chamber perfused with carbogenated ACSF. After collecting a fluorescence baseline (-30 sec) (Panel D- experimental design), drug perfusion was initiated (time zero). Peak signal (30 seconds) was assessed by subtracting baseline fluorescence from stimulated dLight1.3b fluorescence at the 30 second mark. **Panel E:** Stimulated release of dopamine from dopamine terminals originating in the midbrain upon bath perfusion of CNO (10 µM), resulting in a significant increase in dLight1.3b signal. **Panel F:** Additional incremental increase in DA release within the DVC, upon bath perfusion of methamphetamine (10 µM), as determined in separate experiments. **Panel G:** Positive control obtained with 100 µM dopamine (*n=4 independent biological replicates, with 2 technical replicates per experiment; ***p<0.005, ****p<0.0005, two-tailed unpaired Student's T-test*).

757
758
Figure 5. Anterograde multi-transsynaptic tracing reveals midbrain-to-spleen connectivity
via the celiac ganglion. Panel A: Sagittal view of the mouse brain, showing multi-transsynaptic
site of midbrain, where Ai9-Flox-stop-TdTomato mice were injected with pENN-AAV1-hSyn-Cre-
WPRE-hGH. **Panel B:** Diagram showing where, after three weeks of viral expression, the DVC,
celiac ganglia and spleen were assessed via confocal microscopy for expression of TdTomato.
Panel C: DVC neurons are positive for TdTomato, indicating midbrain-to-DVC projections, in a
manner that is consistent with our prior findings. **Panels D-E:** Assessment of celiac ganglia and
spleen show presence of TdTomato⁺ neurons in both organs, indicating both organs received
multi-transsynaptic projections from midbrain dopamine neurons. Taken together with our earlier
neural circuit tracing data, these findings indicate that multi-synaptic projections exist from
midbrain dopamine neurons, via DVC and celiac ganglia, to the spleen. *Data representative of*
n=4 independent experiments.
771

Figure 6. Stimulation with Gq-DREADD of midbrain dopamine neurons increases cFos expression in the DVC, indicating that dopamine neuron activation can modulate DVC activity. Panel A: Experiment timeline, showing (left) site of AAV2-hSyn-DIO-hM3D(Gq)-mCherry or empty vector (AAV2-hSyn-DIO-mCherry) injections into the midbrain DAT-Cre mice, (middle-left) four-week expression interval, (middle-right) daily I.P. injections of clozapine-*N*-oxide (1mg/kg), and (right) cFos immunolocalization in DVC as a proxy measure of neuronal activation. **Panel B:** Minimal/basal cFos immunostaining in the DVC of mice receiving empty vector (AAV2-hSyn-DIO-mCherry). **Panel C:** Robust cFos immunostaining in midbrain of mice injected with AAV encoded with Cre-dependent Gq-DREADD and treated with CNO. **Panel D:** Compared to empty vector control, mice receiving AAV2-hSyn-DIO-hM3D(Gq)-mCherry show significantly increased number of cFos+ cells in the DVC. **Panel E**: Following cervical vagotomy, animals that received AAV2-hSyn-DIO-hM3D(Gq)-mCherry continue to show increased DVC cFos expression compared to empty vector control. (*Data representative of n=6-7 independent biological replicates; ***p<0.005, ****p<0.0005, two-tailed unpaired Student's T-test*)

792
Figure 7. Gq-DREADD stimulation of midbrain dopamine neurons increases cFos expression in celiac ganglia, indicating that dopamine neuron activity can modulate celiac ganglia activity via midbrain-DVC-celiac ganglion pathway. Cervical vagotomy abolishes increased cFos expression in the celiac ganglia. Panel A: Cre-dependent excitatory Gq-DREADDs or empty vector control (AAV2-hSyn-DIO-mCherry) was injected in the midbrain of DAT-Cre mice. Following, mice received either a sham cervical surgery (top) or cervical vagotomy (bottom). After 4 weeks to allow viral expression, mice received 14 days of daily CNO (1mg/kg, I.P.). cFos immunoreactivity was examined as a proxy measure for increased celiac ganglion neuronal activity due to midbrain dopamine neuron activation. **Panel B-C**: Compared to empty

vector control, animals that received Gq-DREADDs and I.P. CNO show increased celiac ganglion
cFos expression. **Panel D**: Mice receiving Gq-DREADDs show significantly increased number of
cFos+ cells in the celiac ganglion. **Panel E-F**: In mice that received cervical vagotomy, Gq-
DREADD mediated stimulation of midbrain dopamine neurons does not significantly increase the
number of cFos+ cells in the celiac ganglion. **Panel G**: Mice that received cervical vagotomy do
not show increased (*Data representative of n=3-4 independent biological replicates; *p<0.05, two-*
tailed unpaired Student's T-test)

817
Figure 8. Spleen weight and flow cytometric analysis, showing that stimulation of midbrain
dopamine neurons by Gq-DREADD results in reduced spleen weight and reduced naïve
CD4$^+$ T-cell numbers in the spleen without effect on CD8$^+$ T-cell populations. Panel A:
Experiment timeline, showing site of AAV2-hSyn-DIO-hM3D(Gq)-mCherry or empty vector
(AAV2-hSyn-DIO-mCherry) injections into the midbrain DAT-Cre mice, followed by sham surgery
targeting the cervical region. After four-weeks for viral expression, followed by daily I.P. injections
of clozapine-*N*-oxide (1mg/kg) for two weeks, harvesting spleen for weighing cellular dissociation,
and (right) flow cytometry using antibodies against T-cell subtypes. **Panel B:** Animal body weight
was monitored throughout CNO administration; mice that received Gq-DREADDs vs empty vector
control did not exhibit decreases in body weight. **Panel C**: Reduced spleen weight (adjusted for
body weight) in mice receiving midbrain Gq-DREADD and systemic CNO, as compared to empty
vector control (p<0.05). **Panels D-E**: Single live cells, free of debris, were gated for CD3
expression, a pan-T-cell marker, and then gated for expression of CD4 and CD8. **Panels F-I:**

CD4$^+$ and CD8$^+$ T-cells were analyzed by expression of CD44 and/or CD62L to assess
populations of T-effector memory cells (TEMs), T-central memory cells (TCMs) or naïve T-cells.
Panels H-M: With no change in total CD4+ T-cell populations, compared to empty vector control,
animals that received Gq-DREADDs in the midbrain show significantly reduced naïve CD4+ T-
cells, marked by expression of CD62L and lacking expression of CD44. CD4$^+$ T-effector memory
cells and T-central memory cells were not significantly changed. **Panels N-Q:** CD8$^+$ T-cell subsets
are not significantly changed by Gq-DREADD-mediated midbrain dopamine neuron activation.
(n=8-10 independent biological replicates. Panel B: p>0.05, 2-way ANOVA, N.S. Panels J-Q:
*p<0.05, two-tailed unpaired Student's T-test)

845
Figure 9. Following vagotomy, effect on spleen weight and splenic immune cells is
abolished despite activation of midbrain dopamine neurons, indicating a specific pathway
from midbrain to spleen via vagal afferents. Panel A: Experimental timeline showing site of
AAV2-hSyn-DIO-hM3D(Gq)-mCherry or empty vector (AAV2-hSyn-DIO-mCherry) injections into
the midbrain DAT-Cre mice, followed by cervical surgery transecting the vagus nerve (cervical
vagotomy). After four-weeks for viral expression, followed by daily I.P. injections of clozapine-*N*-
oxide (1mg/kg) for two weeks, harvesting spleen for weighing and cellular dissociation, and
analysis flow cytometry using antibodies against T-cell subtypes. **Panel B**: Animal body weight
was monitored throughout CNO administration; mice that received Gq-DREADDs vs empty vector
control did not exhibit decreases in body weight. **Panel C**: In contrast to intact animals receiving
midbrain Gq-DREADDs, vagotomized animals that received Gq-DREADDs or Empty Vector
Control did not exhibit significantly different spleen weights adjusted for body weight. **Panels D-**
E: Single live cells, free of debris, were gated for CD3 expression, a pan-T-cell marker, and then

gated for expression of CD4 and CD8. **Panels F-I:** In vagotomized animals that received either empty vector control or Gq-DREADDs, CD4$^+$ and CD8$^+$ T-cells were analyzed by expression of CD44 and/or CD62L to assess populations of T-effector memory cells (TEMs), T-central memory cells (TCMs) or naïve T-cells. **Panel J-R**: Regardless of whether animals received empty vector control or Gq-DREADD AAV in the midbrain, cervical vagotomy abolishes the impact of midbrain dopamine neuron activation on splenic T-cell populations *(n=4-6 independent biological replicates. Panel B: p>0.05, 2-way ANOVA, N.S. Panels J-Q: p>0.05, two-tailed unpaired Student's T-test, N.S.).*

890
891 **Supplementary Figure 1. Single channel control images for EYFP-expressing midbrain and**
892 **median forebrain bundle verify injection site and expression in midbrain dopamine neuron**
893 **projections**. **Panel A:** Median forebrain bundle of DAT-Cre mice injected with AAV2-Ef1a-DIO-
894 EYFP shows EYFP+ fibers and overlap with TH immunoreactive fibers. **Panel B:** At the midbrain
895 site of injection, EYFP+ cell bodies overlap with TH+ midbrain dopamine neurons.
896
897

898
Supplementary Figure 2. Single channel control images for synaptophysin-expressing
midbrain and striatum verify injection site and expression in midbrain dopamine neuron
projections. **Panel A:** Striatum of DAT-Cre mice injected with AAV1-hSyn-Flex mGFP-2A-
Synaptophysin-mRuby shows synaptophysin-mRuby+ terminals and overlap with TH
immunoreactive terminals. **Panel B**: At the midbrain site of injection, synaptophysin-mRuby+ cell
bodies overlap with TH+ midbrain dopamine neurons.
905
906
907
908

909
Supplementary Figure 3. mCherry-tagged Cre-dependent Gq-DREADDs are expressed in
midbrain dopamine neurons and terminal regions including the DVC. DAT-Cre mice
received midbrain microinjections of Cre-dependent AAV-DIO-mCherry-hM3D (Gq-DREADD), or
AAV2-hSyn-DIO-mCherry (empty vector). mCherry fluorescence was assessed in midbrain and
in terminal regions. **Panel A**: Single channel control images show midbrain dopamine neurons,
identified by expression of TH, express mCherry-tagged Gq-DREADDs. **Panel B**: mCherry[+]
terminals are evident throughout the striatum, indicating this AAV construct expresses Gq-
DREADD receptors in terminal regions. **Panel C**: mCherry[+] Gq-DREADD-expressing terminals
are evident throughout the DVC.

Supplementary Figure 4: Gq-DREADD stimulation of midbrain dopamine neurons increases cFos expression in both midbrain and striatum. DAT-Cre mice received midbrain microinjections of AAV-hSyn-DIO-mCherry-hM3D (Gq-DREADD), or AAV2-hSyn-DIO-mCherry (empty vector). **Panel A-C:** Midbrain dopamine neurons of mice that received excitatory Gq-DREADDs, identified by expression of DAT, show robust cFos expression following systemic CNO administration. **Panel D-F:** Striatal terminal regions of dopamine neurons, identified by DAT expression, also exhibit robust cFos expression.

950
951
Supplementary Figure 5: Single channel control images for TdTomato+ signals in the
celiac ganglion and spleen. Panel A-C: Single channel images showing TdTomato (Panel A),
TH (Panel B) and a merge image (Panel C) showing TdTomato+ cells in proximity to TH+ cell
bodies in the celiac ganglion of Ai9 mice that received midbrain injections of pENN-AAV1-hSyn-
Cre-WPRE-hGH. **Panel D-G**: Single channel images showing TdTomato expression (Panel D),
TH-expressing fibers (Panel E) and CD3+ T-cell zones (Panel F), and a merged image (Panel G)
showing TdTomato+ fibers surrounding T-cell zones in the spleen of Ai9 mice that received
midbrain injections of pENN-AAV1-hSyn-Cre-WPRE-hGH.
960
961
962
963

978
Supplementary Figure 7: Following cervical vagotomy, Gq-DREADD stimulation results in
increased locomotor activity following intraperitoneal injection of 1mg/kg clozapine-*N*-
oxide. Panel A: Locomotor activity in the home cage was measured prior to drug injection (Day
0), and on Days 1, 7 and 13 during the CNO administration period. **Panel B:** On Day 0, total
distance traveled is not significantly different between animals receiving Gq-DREADD in midbrain
dopamine neurons relative to empty vector (AAV2-hSyn-DIO-mCherry). Consistent with the
literature in the field, CNO increases locomotor activity in animals with midbrain dopamine neuron
Gq-DREADDs compared to empty vector on Days 1, 7 and 13. **Panel C:** Average speed is also
significantly increased in animals receiving Gq-DREADDs relative to empty vector control (AAV2-
hSyn-DIO-mCherry) on Days 1, 7 and 13. *(n=8 independent biological replicates;*
p<0.005,*p<0.0005, two-tailed unpaired Student's T-test). See also **Supplementary Movie**
2.

991
992
993
994 **Supplementary Figure 8: Chemogenetic activation of midbrain dopamine neurons does**
995 **not increase cFos expression in amygdala. Panels A-B:** In the central amygdala, identified by
996 anatomical landmarks and presence of TH+ fibers, chemogenetic activation of midbrain dopamine
997 neurons does not increase cFos expression.

998
999
Supplementary Figure 9. B-cell populations are not significantly altered when midbrain
dopamine neurons are stimulated with Gq DREADDs. Panel A-B: Single, live cells, free of
debris were gated for expression of CD45. **Panel C-D**: Total CD19+ B-cells, and **Panels G-H**:
Immature and Maturing B-cells, identified by expression of IgM alone or co-expression of IgM and
IgD, were assessed to determine impact of midbrain dopamine neuron activation. **Panel E-F**:
Total CD19+ B-cells were not significantly altered when midbrain dopamine neurons were
activated. **Panel I-L**: Immature B-cells and Maturing B-cells were not significantly altered when
midbrain dopamine neurons were activated. *(n=6 independent biological replicates; two-tailed*
unpaired Student's T-test, p>0.05; N.S.)

1009
1010
1011

1012

1013
1014

Supplementary Figure 10. NK-cell populations are not significantly altered when midbrain dopamine neurons are stimulated with Gq DREADDs. Panel A-B: Single, live cells, free of debris were gated for expression of CD45 and lack of CD3 expression (CD45+/CD3-) to identify NK cells. **Panel C-D**: Total NK cells, identified by expression of CD335 and NK1.1, and **Panel G-H**: Immature and mature NK cells, identified by expression of KLGR1, were assessed to determine impact of midbrain dopamine neuron activation. **Panel E-F**: Total NK-cells were not significantly altered when midbrain dopamine neurons were activated. **Panel I-L**: Immature and Mature NK-cells were not significantly altered when midbrain dopamine neurons were activated. *(n=6 independent biological replicates; two-tailed unpaired Student's T-test, p>0.05; N.S.)*

1025
Supplementary Figure 11. Splenic macrophages are not significantly altered when
midbrain dopamine neurons are stimulated with Gq DREADDs. Panel A-B: Single, live cells,
free of debris were gated for CD45 expression. **Panel C-D**: Splenic macrophages were identified
by co-expression of CD11b and F4/80 (CD11b+/F480+). Panel E-F: Splenic macrophages were
not significantly altered when midbrain dopamine neurons were activated. *(n=6 independent*
biological replicates; two-tailed unpaired Student's T-test, p>0.05; N.S.).
1032
1033
1034

1035
1036
Supplementary Figure 12. Fluorescence-minus-one generated gating strategy to analyze
splenic T-cell populations. **Panel A**: Single, live cells free of debris were (**Panel B**) assessed
for expression of CD45. **Panel B**: Subsequently, cells expressing CD3 were assessed for CD4
and CD8 expression and then assessed for T-central memory (TCM), T-effector memory (TEM)
and Naïve T-cells. **Panels C-H**: Gates for analyses for each of the above populations were set by
fluorescence-minus-one gating strategy.

1043
1044
1045 **Supplementary Figure 13. Fluorescence-minus-one generated gating strategy to analyze**
1046 **splenic B-cell populations**. **Panel A**: Single, live cells free of debris were (**Panel B**) assessed
1047 for expression of CD45. **Panel B**: Subsequently, cells expressing CD19 were assessed for IgM
1048 and IgD expression. **Panels C-E**: Gates for analyses for each of the above populations were set
1049 by fluorescence-minus-one gating strategy.

1057
1058
Supplementary Figure 14. Fluorescence-minus-one generated gating strategy to analyze
splenic NK-cell populations. **Panel A**: Single, live cells free of debris were (**Panel B**) assessed
for expression of CD45 without CD3 expression (CD45+/CD3-). **Panel B**: Subsequently, cells co-
expressing CD335 and NK1.1 (total NK-cells) were assessed for KLGR1 expression. **Panels C-**
G: Gates for analyses for each of the above populations were set by fluorescence-minus-one
gating strategy.
1065
1066

1067
1068
1069

1070 **Supplementary Figure 15. Fluorescence-minus-one generated gating strategy to analyze**
1071 **splenic macrophages**. **Panel A**: Single, live cells free of debris were (**Panel B**) assessed for
1072 expression of CD45. **Panel B**: Subsequently, cells were assessed for co-expression of CD11b
1073 and F4/80 (splenic macrophages). **Panels C-D**: Gates for analyses for each of the above
1074 populations were set by fluorescence-minus-one gating strategy.

1100
1101 **Supplementary Video 1: Increased locomotor activity evident throughout 14 days of**
1102 **systemic CNO treatment in mice that received excitatory Gq-DREADD AAV injection in**
1103 **midbrain dopamine neurons and sham cervical surgery.**
1104
1105 **Supplementary Video 2: Increased locomotor activity evident throughout 14 days of**
1106 **systemic CNO treatment in mice that received excitatory Gq-DREADD AAV injection in**
1107 **midbrain dopamine neurons and cervical vagotomy.**

References

Jin, H., Li, M., Jeong, E., Castro-Martinez, F. & Zuker, C. S. A body–brain circuit that regulates body inflammatory responses. *Nature* **630**, 695-703, doi:10.1038/s41586-024-07469-y (2024).

Zhu, X. *et al.* Somatosensory cortex and central amygdala regulate neuropathic pain-mediated peripheral immune response via vagal projections to the spleen. *Nature Neuroscience* **27**, 471-483, doi:10.1038/s41593-023-01561-8 (2024).

Tansey, M. G. *et al.* Inflammation and immune dysfunction in Parkinson disease. *Nat Rev Immunol* **22**, 657-673, doi:10.1038/s41577-022-00684-6 (2022).

Cossu, D., Hatano, T. & Hattori, N. The Role of Immune Dysfunction in Parkinson's Disease Development. *Int J Mol Sci* **24**, doi:10.3390/ijms242316766 (2023).

Tan, E. K. *et al.* Parkinson disease and the immune system - associations, mechanisms and therapeutics. *Nat Rev Neurol* **16**, 303-318, doi:10.1038/s41582-020-0344-4 (2020).

Gopinath, A. *et al.* Methamphetamine induces a low dopamine transporter expressing state without altering the total number of peripheral immune cells. *Basic Clin Pharmacol Toxicol* **133**, 496-507, doi:10.1111/bcpt.13838 (2023).

Weaver, J. L. *et al.* Methamphetamine Use is Associated with Increased Surgical Site Infections after Trauma Laparotomy. *J Surg Res* **267**, 563-567, doi:10.1016/j.jss.2021.06.034 (2021).

Hulsebos, I. F. *et al.* Stimulant Abuse in Burn Patients Is Associated With an Increased Use of Hospital Resources. *J Burn Care Res* **41**, 921-925, doi:10.1093/jbcr/iraa087 (2020).

Salamanca, S. A., Sorrentino, E. E., Nosanchuk, J. D. & Martinez, L. R. Impact of methamphetamine on infection and immunity. *Front Neurosci* **8**, 445, doi:10.3389/fnins.2014.00445 (2014).

Miller, D. R., Bu, M., Gopinath, A., Martinez, L. R. & Khoshbouei, H. Methamphetamine Dysregulation of the Central Nervous System and Peripheral Immunity. *J Pharmacol Exp Ther* **379**, 372-385, doi:10.1124/jpet.121.000767 (2021).

Miller, E. J. & Khoshbouei, H. Immunity on ice: The impact of methamphetamine on peripheral immunity. *Adv Pharmacol* **99**, 217-250, doi:10.1016/bs.apha.2023.09.003 (2024).

Eisenberger, N. I. *et al.* Inflammation-induced anhedonia: endotoxin reduces ventral striatum responses to reward. *Biol Psychiatry* **68**, 748-754, doi:10.1016/j.biopsych.2010.06.010 (2010).

Lee, Y. *et al.* Anti-cytokine agents for anhedonia: targeting inflammation and the immune system to treat dimensional disturbances in depression. *Ther Adv Psychopharmacol* **8**, 337-348, doi:10.1177/2045125318791944 (2018).

Mehta, N. D. *et al.* Inflammation, reward circuitry and symptoms of anhedonia and PTSD in trauma-exposed women. *Soc Cogn Affect Neurosci* **15**, 1046-1055, doi:10.1093/scan/nsz100 (2020).

Cooper, C. M., Farrand, A. Q., Andresen, M. C. & Beaumont, E. Vagus nerve stimulation activates nucleus of solitary tract neurons via supramedullary pathways. *J Physiol* **599**, 5261-5279, doi:10.1113/jp282064 (2021).

Jayaprakash, N. *et al.* Organ- and function-specific anatomical organization of vagal fibers supports fascicular vagus nerve stimulation. *Brain Stimul* **16**, 484-506, doi:10.1016/j.brs.2023.02.003 (2023).

Matteoli, G. & Boeckxstaens, G. E. The vagal innervation of the gut and immune homeostasis. *Gut* **62**, 1214, doi:10.1136/gutjnl-2012-302550 (2013).

Tao, J. *et al.* Highly selective brain-to-gut communication via genetically defined vagus neurons. *Neuron* **109**, 2106-2115.e2104, doi:10.1016/j.neuron.2021.05.004 (2021).

Pavlov, V. A. & Tracey, K. J. The cholinergic anti-inflammatory pathway. *Brain Behav Immun* **19**, 493-499, doi:10.1016/j.bbi.2005.03.015 (2005).

Alen, N. V. The cholinergic anti-inflammatory pathway in humans: State-of-the-art review and future directions. *Neurosci Biobehav Rev* **136**, 104622, doi:10.1016/j.neubiorev.2022.104622 (2022).

Wu, S. J. *et al.* Activation of the Cholinergic Anti-inflammatory Pathway Attenuated Angiotension II-Dependent Hypertension and Renal Injury. *Front Pharmacol* **12**, 593682, doi:10.3389/fphar.2021.593682 (2021).

Holt, M. K. *et al.* Synaptic Inputs to the Mouse Dorsal Vagal Complex and Its Resident Preproglucagon Neurons. *The Journal of Neuroscience* **39**, 9767, doi:10.1523/JNEUROSCI.2145-19.2019 (2019).

Xing, T., Nanni, G., Burkholder, C. R., Browning, K. N. & Travagli, R. A. The substantia nigra modulates proximal colon tone and motility in a vagally-dependent manner in the rat. *J Physiol* **601**, 4751-4766, doi:10.1113/jp284238 (2023).

Kobori, N., Moore, A. N., Redell, J. B. & Dash, P. K. Caudal DMN neurons innervate the spleen and release CART peptide to regulate neuroimmune function. *J Neuroinflammation* **20**, 158, doi:10.1186/s12974-023-02838-2 (2023).

Bassi, G. S. *et al.* Anatomical and clinical implications of vagal modulation of the spleen. *Neuroscience & Biobehavioral Reviews* **112**, 363-373, doi:https://doi.org/10.1016/j.neubiorev.2020.02.011 (2020).

Tanaka, S. *et al.* Vagus nerve stimulation activates two distinct neuroimmune circuits converging in the spleen to protect mice from kidney injury. *Proceedings of the National Academy of Sciences* **118**, e2021758118, doi:10.1073/pnas.2021758118 (2021).

Hoeger, S. *et al.* Modulation of brain dead induced inflammation by vagus nerve stimulation. *Am J Transplant* **10**, 477-489, doi:10.1111/j.1600-6143.2009.02951.x (2010).

Inoue, T. *et al.* Vagus nerve stimulation mediates protection from kidney ischemia-reperfusion injury through α7nAChR+ splenocytes. *J Clin Invest* **126**, 1939-1952, doi:10.1172/jci83658 (2016).

Rosas-Ballina, M. *et al.* Splenic nerve is required for cholinergic antiinflammatory pathway control of TNF in endotoxemia. *Proc Natl Acad Sci U S A* **105**, 11008-11013, doi:10.1073/pnas.0803237105 (2008).

Huston, J. M. *et al.* Cholinergic neural signals to the spleen down-regulate leukocyte trafficking via CD11b. *J Immunol* **183**, 552-559, doi:10.4049/jimmunol.0802684 (2009).

Leaders, F. E. & Dayrit, C. THE CHOLINERGIC COMPONENT IN THE SYMPATHETIC INNERVATION TO THE SPLEEN. *J Pharmacol Exp Ther* **147**, 145-152 (1965).

Vida, G., Peña, G., Deitch, E. A. & Ulloa, L. α7-cholinergic receptor mediates vagal induction of splenic norepinephrine. *J Immunol* **186**, 4340-4346, doi:10.4049/jimmunol.1003722 (2011).

Mota, C. M. D. & Madden, C. J. Neural control of the spleen as an effector of immune responses to inflammation: mechanisms and treatments. *Am J Physiol Regul Integr Comp Physiol* **323**, R375-r384, doi:10.1152/ajpregu.00151.2022 (2022).

Bellinger, D. L., Felten, S. Y., Lorton, D. & Felten, D. L. Origin of noradrenergic innervation of the spleen in rats. *Brain Behav Immun* **3**, 291-311, doi:10.1016/0889-1591(89)90029-9 (1989).

Wu, M. *et al.* Innervation of nociceptor neurons in the spleen promotes germinal center responses and humoral immunity. *Cell* **187**, 2935-2951.e2919, doi:10.1016/j.cell.2024.04.027 (2024).

Han, W. *et al.* A Neural Circuit for Gut-Induced Reward. *Cell* **175**, 665-678.e623, doi:10.1016/j.cell.2018.08.049 (2018).

Marvel, F. A., Chen, C. C., Badr, N., Gaykema, R. P. & Goehler, L. E. Reversible inactivation of the dorsal vagal complex blocks lipopolysaccharide-induced social

withdrawal and c-Fos expression in central autonomic nuclei. *Brain Behav Immun* **18**, 123-134, doi:10.1016/j.bbi.2003.09.004 (2004).

Chavan, S. S., Pavlov, V. A. & Tracey, K. J. Mechanisms and Therapeutic Relevance of Neuro-immune Communication. *Immunity* **46**, 927-942, doi:10.1016/j.immuni.2017.06.008 (2017).

Zhuang, X., Masson, J., Gingrich, J. A., Rayport, S. & Hen, R. Targeted gene expression in dopamine and serotonin neurons of the mouse brain. *J Neurosci Methods* **143**, 27-32, doi:10.1016/j.jneumeth.2004.09.020 (2005).

Beier, K. T. *et al.* Circuit Architecture of VTA Dopamine Neurons Revealed by Systematic Input-Output Mapping. *Cell* **162**, 622-634, doi:10.1016/j.cell.2015.07.015 (2015).

Pavlov, V. A., Wang, H., Czura, C. J., Friedman, S. G. & Tracey, K. J. The cholinergic anti-inflammatory pathway: a missing link in neuroimmunomodulation. *Mol Med* **9**, 125-134 (2003).

Bennett, B. D., Callaway, J. C. & Wilson, C. J. Intrinsic membrane properties underlying spontaneous tonic firing in neostriatal cholinergic interneurons. *J Neurosci* **20**, 8493-8503, doi:10.1523/jneurosci.20-22-08493.2000 (2000).

Trapp, S. & Ballanyi, K. Vol. 73   1-59 (2012).

Anselmi, L., Toti, L., Bove, C. & Travagli, R. A. Vagally mediated effects of brain stem dopamine on gastric tone and phasic contractions of the rat. *Am J Physiol Gastrointest Liver Physiol* **313**, G434-g441, doi:10.1152/ajpgi.00180.2017 (2017).

Lawrence, A. J., Krstew, E. & Jarrott, B. Functional dopamine D2 receptors on rat vagal afferent neurones. *Br J Pharmacol* **114**, 1329-1334, doi:10.1111/j.1476-5381.1995.tb13352.x (1995).

Anselmi, L., Toti, L., Bove, C., Hampton, J. & Travagli, R. A. A Nigro−Vagal Pathway Controls Gastric Motility and Is Affected in a Rat Model of Parkinsonism. *Gastroenterology* **153**, 1581-1593, doi:https://doi.org/10.1053/j.gastro.2017.08.069 (2017).

Patriarchi, T. *et al.* An expanded palette of dopamine sensors for multiplex imaging in vivo. *Nat Methods* **17**, 1147-1155, doi:10.1038/s41592-020-0936-3 (2020).

Yorgason, J. T. *et al.* Methamphetamine increases dopamine release in the nucleus accumbens through calcium-dependent processes. *Psychopharmacology (Berl)* **237**, 1317-1330, doi:10.1007/s00213-020-05459-2 (2020).

Goodwin, J. S. *et al.* Amphetamine and methamphetamine differentially affect dopamine transporters in vitro and in vivo. *J Biol Chem* **284**, 2978-2989, doi:10.1074/jbc.M805298200 (2009).

Sambo, D. O. *et al.* The sigma-1 receptor modulates methamphetamine dysregulation of dopamine neurotransmission. *Nat Commun* **8**, 2228, doi:10.1038/s41467-017-02087-x (2017).

Pavlov, V. A., Chavan, S. S. & Tracey, K. J. Molecular and Functional Neuroscience in Immunity. *Annu Rev Immunol* **36**, 783-812, doi:10.1146/annurev-immunol-042617-053158 (2018).

Hoover, D. B. Cholinergic modulation of the immune system presents new approaches for treating inflammation. *Pharmacol Ther* **179**, 1-16, doi:10.1016/j.pharmthera.2017.05.002 (2017).

Kaestner, C. L., Smith, E. H., Peirce, S. G. & Hoover, D. B. Immunohistochemical analysis of the mouse celiac ganglion: An integrative relay station of the peripheral nervous system. *J Comp Neurol* **527**, 2742-2760, doi:10.1002/cne.24705 (2019).

Klein Wolterink, R. G. J., Wu, G. S., Chiu, I. M. & Veiga-Fernandes, H. Neuroimmune Interactions in Peripheral Organs. *Annu Rev Neurosci* **45**, 339-360, doi:10.1146/annurev-neuro-111020-105359 (2022).

55 Wang, T., Teng, B., Yao, D. R., Gao, W. & Oka, Y. Organ-specific Sympathetic Innervation
Defines Visceral Functions. *bioRxiv*, 2024.2009.2019.613934,
doi:10.1101/2024.09.19.613934 (2024).
56 Kressel, A. M. *et al.* Identification of a brainstem locus that inhibits tumor necrosis factor.
Proc Natl Acad Sci U S A **117**, 29803-29810, doi:10.1073/pnas.2008213117 (2020).
57 Carnevale, D. *et al.* A cholinergic-sympathetic pathway primes immunity in hypertension
and mediates brain-to-spleen communication. *Nat Commun* **7**, 13035,
doi:10.1038/ncomms13035 (2016).
58 Shinohara, K. Celiac ganglia: potential new targets in neuromodulation for hypertension.
Hypertension Research **46**, 2235-2236, doi:10.1038/s41440-023-01355-2 (2023).
59 Trevizan-Baú, P. *et al.* Protocol for the isolation of the mouse sympathetic splanchnic-
celiac-superior mesenteric ganglion complex. *STAR Protoc* **5**, 103036,
doi:10.1016/j.xpro.2024.103036 (2024).
60 Fernandez, R. *et al.* Neural reflex regulation of systemic inflammation: potential new
targets for sepsis therapy. *Front Physiol* **5**, 489, doi:10.3389/fphys.2014.00489 (2014).
61 Browning, K. N. & Travagli, R. A. Central nervous system control of gastrointestinal motility
and secretion and modulation of gastrointestinal functions. *Compr Physiol* **4**, 1339-1368,
doi:10.1002/cphy.c130055 (2014).
62 Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system
for the whole mouse brain. *Nat Neurosci* **13**, 133-140, doi:10.1038/nn.2467 (2010).
63 Zingg, B. *et al.* AAV-Mediated Anterograde Transsynaptic Tagging: Mapping
Corticocollicular Input-Defined Neural Pathways for Defense Behaviors. *Neuron* **93**, 33-
47, doi:https://doi.org/10.1016/j.neuron.2016.11.045 (2017).
64 Moura, M. *et al.* Macro- and Microanatomy of the Sympathetic Innervation of the Spleen
in Rodents. *J Comp Neurol* **533**, e70086, doi:10.1002/cne.70086 (2025).
65 Martelli, D., Yao, S. T., McKinley, M. J. & McAllen, R. M. Reflex control of inflammation by
sympathetic nerves, not the vagus. *J Physiol* **592**, 1677-1686,
doi:10.1113/jphysiol.2013.268573 (2014).
66 McAllen, R. M., McKinley, M. J. & Martelli, D. Reflex regulation of systemic inflammation
by the autonomic nervous system. *Autonomic Neuroscience* **237**, 102926,
doi:https://doi.org/10.1016/j.autneu.2021.102926 (2022).
67 Morgan, J. I., Cohen, D. R., Hempstead, J. L. & Curran, T. Mapping patterns of c-fos
expression in the central nervous system after seizure. *Science* **237**, 192-197,
doi:10.1126/science.3037702 (1987).
68 Barros, V. N. *et al.* The pattern of c-Fos expression and its refractory period in the brain
of rats and monkeys. *Front Cell Neurosci* **9**, 72, doi:10.3389/fncel.2015.00072 (2015).
69 Gao, Y. J. & Ji, R. R. c-Fos and pERK, which is a better marker for neuronal activation
and central sensitization after noxious stimulation and tissue injury? *Open Pain J* **2**, 11-
17, doi:10.2174/1876386300902010011 (2009).
70 Cullinan, W. E., Herman, J. P., Battaglia, D. F., Akil, H. & Watson, S. J. Pattern and time
course of immediate early gene expression in rat brain following acute stress.
Neuroscience **64**, 477-505, doi:10.1016/0306-4522(94)00355-9 (1995).
71 Lawson, K. A., Ruiz, C. M. & Mahler, S. V. A head-to-head comparison of two DREADD
agonists for suppressing operant behavior in rats via VTA dopamine neuron inhibition.
Psychopharmacology (Berl) **240**, 2101-2110, doi:10.1007/s00213-023-06429-0 (2023).
72 Ahern, J., Chrobok, Ł., Champneys, A. R. & Piggins, H. D. A new phase model of the
spatiotemporal relationships between three circadian oscillators in the brainstem. *Sci Rep*
13, 5480, doi:10.1038/s41598-023-32315-y (2023).
73 Espinoza, D., Seibold, F. & Stanley, S. Central and peripheral neural circuits regulating
glucose homeostasis. *npj Biomedical Innovations* **2**, 34, doi:10.1038/s44385-025-00036-
8 (2025).

Hes, C. *et al.* A unified rodent atlas reveals the cellular complexity and evolutionary divergence of the dorsal vagal complex. *eLife* **14**, RP106217, doi:10.7554/eLife.106217 (2025).

Ludwig, M. Q. *et al.* A genetic map of the mouse dorsal vagal complex and its role in obesity. *Nat Metab* **3**, 530-545, doi:10.1038/s42255-021-00363-1 (2021).

Jobling, P. & Gibbins, I. L. Electrophysiological and morphological diversity of mouse sympathetic neurons. *J Neurophysiol* **82**, 2747-2764, doi:10.1152/jn.1999.82.5.2747 (1999).

Hayakawa, T., Kuwahara, S., Maeda, S., Tanaka, K. & Seki, M. Morphology and ultrastructure of the sympathetic celiac ganglion neurons projecting to the cardia and pylorus of the rat stomach. *Auton Neurosci* **134**, 8-17, doi:10.1016/j.autneu.2007.01.011 (2007).

Wang, T., Teng, B., Yao, D. R., Gao, W. & Oka, Y. Organ-specific sympathetic innervation defines visceral functions. *Nature* **637**, 895-902, doi:10.1038/s41586-024-08269-0 (2025).

Lewis, S. M., Williams, A. & Eisenbarth, S. C. Structure and function of the immune system in the spleen. *Sci Immunol* **4**, doi:10.1126/sciimmunol.aau6085 (2019).

Wluka, A. & Olszewski, W. L. Innate and adaptive processes in the spleen. *Ann Transplant* **11**, 22-29 (2006).

Ugel, S. *et al.* Immune tolerance to tumor antigens occurs in a specialized environment of the spleen. *Cell Rep* **2**, 628-639, doi:10.1016/j.celrep.2012.08.006 (2012).

Bronte, V. & Pittet, M. J. The spleen in local and systemic regulation of immunity. *Immunity* **39**, 806-818, doi:10.1016/j.immuni.2013.10.010 (2013).

Mebius, R. E. & Kraal, G. Structure and function of the spleen. *Nat Rev Immunol* **5**, 606-616, doi:10.1038/nri1669 (2005).

Wu, C., Hua, Q. & Zheng, L. Generation of Myeloid Cells in Cancer: The Spleen Matters. *Front Immunol* **11**, 1126, doi:10.3389/fimmu.2020.01126 (2020).

Beheshti, A. *et al.* Tumor-host signaling interaction reveals a systemic, age-dependent splenic immune influence on tumor development. *Oncotarget* **6**, 35419-35432, doi:10.18632/oncotarget.6214 (2015).

Markel, J. E. *et al.* Using the Spleen as an In Vivo Systemic Immune Barometer Alongside Osteosarcoma Disease Progression and Immunotherapy with α-PD-L1. *Sarcoma* **2018**, 8694397, doi:10.1155/2018/8694397 (2018).

Chen, X. *et al.* Evidence for Peripheral Immune Activation in Parkinson's Disease. *Front Aging Neurosci* **13**, 617370, doi:10.3389/fnagi.2021.617370 (2021).

Ghanta, M. K. *et al.* Role of CD4+ T Cells in Parkinson's Disease. *Crit Rev Immunol* **42**, 17-25, doi:10.1615/CritRevImmunol.2023047372 (2022).

Mamula, D., Khosousi, S., He, Y., Lazarevic, V. & Svenningsson, P. Impaired migratory phenotype of CD4(+) T cells in Parkinson's disease. *NPJ Parkinsons Dis* **8**, 171, doi:10.1038/s41531-022-00438-0 (2022).

Bravo, J., Magalhães, C., Andrade, E. B., Magalhães, A. & Summavielle, T. The impact of psychostimulants on central and peripheral neuro-immune regulation: a scoping review of cytokine profiles and their implications for addiction. *Front Cell Neurosci* **17**, 1109611, doi:10.3389/fncel.2023.1109611 (2023).

Carrico, A. W. *et al.* Stimulant use is associated with immune activation and depleted tryptophan among HIV-positive persons on anti-retroviral therapy. *Brain Behav Immun* **22**, 1257-1262, doi:10.1016/j.bbi.2008.07.010 (2008).

Glac, W. *et al.* Prolonged Peripheral Immunosuppressive Responses as Consequences of Random Amphetamine Treatment, Amphetamine Withdrawal and Subsequent Amphetamine Challenges in Rats. *J Neuroimmune Pharmacol* **16**, 870-887, doi:10.1007/s11481-021-09988-1 (2021).

93 Wu, H. *et al.* Dynamic immune and exosome transcriptomic responses in patients
undergoing psychostimulant methamphetamine withdrawal. *Front Cell Neurosci* **16**,
961131, doi:10.3389/fncel.2022.961131 (2022).
94 Assis, M. A., Hansen, C., Lux-Lantos, V. & Cancela, L. M. Sensitization to amphetamine
occurs simultaneously at immune level and in met-enkephalin of the nucleus accumbens
and spleen: an involved NMDA glutamatergic mechanism. *Brain Behav Immun* **23**, 464-
473, doi:10.1016/j.bbi.2009.01.003 (2009).
95 Ben-Shaanan, T. L. *et al.* Activation of the reward system boosts innate and adaptive
immunity. *Nat Med* **22**, 940-944, doi:10.1038/nm.4133 (2016).
96 Ben-Shaanan, T. L. *et al.* Modulation of anti-tumor immunity by the brain's reward system.
Nat Commun **9**, 2723, doi:10.1038/s41467-018-05283-5 (2018).
97 Gopinath, A. *et al.* DAT and TH expression marks human Parkinson's disease in
peripheral immune cells. *NPJ Parkinsons Dis* **8**, 72, doi:10.1038/s41531-022-00333-8
(2022).
98 Gopinath, A. *et al.* Who Knew? Dopamine Transporter Activity Is Critical in Innate and
Adaptive Immune Responses. *Cells* **12**, doi:10.3390/cells12020269 (2023).
99 Berthoud, H. R. & Powley, T. L. Interaction between parasympathetic and sympathetic
nerves in prevertebral ganglia: morphological evidence for vagal efferent innervation of
ganglion cells in the rat. *Microscopy research and technique* **35**, 80-86 (1996).
100 Berthoud, H.-R. & Powley, T. L. Characterization of vagal innervation to the rat celiac,
suprarenal and mesenteric ganglia. *Journal of the autonomic nervous system* **42**, 153-169
(1993).
101 Kavelaars, A., Cobelens, P. M., Teunis, M. A. & Heijnen, C. J. Changes in innate and
acquired immune responses in mice with targeted deletion of the dopamine transporter
gene. *J Neuroimmunol* **161**, 162-168, doi:10.1016/j.jneuroim.2005.01.004 (2005).
102 Bencsics, A. *et al.* Dopamine, as well as norepinephrine, is a link between noradrenergic
nerve terminals and splenocytes. *Brain Res* **761**, 236-243, doi:10.1016/s0006-
8993(97)00313-2 (1997).
103 Cano, G., Sved, A. F., Rinaman, L., Rabin, B. S. & Card, J. P. Characterization of the
central nervous system innervation of the rat spleen using viral transneuronal tracing. *J
Comp Neurol* **439**, 1-18, doi:10.1002/cne.1331 (2001).
104 Murray, K. *et al.* Neuroanatomy of the spleen: Mapping the relationship between
sympathetic neurons and lymphocytes. *PLoS One* **12**, e0182416,
doi:10.1371/journal.pone.0182416 (2017).
105 Elenkov, I. J., Wilder, R. L., Chrousos, G. P. & Vizi, E. S. The sympathetic nerve--an
integrative interface between two supersystems: the brain and the immune system.
Pharmacol Rev **52**, 595-638 (2000).
106 Vaccarino, V. *et al.* Brain-heart connections in stress and cardiovascular disease:
Implications for the cardiac patient. *Atherosclerosis* **328**, 74-82,
doi:10.1016/j.atherosclerosis.2021.05.020 (2021).
107 Liu, W. *et al.* Brain-heart communication in health and diseases. *Brain Res Bull* **183**, 27-
37, doi:10.1016/j.brainresbull.2022.02.012 (2022).
108 Braun, J. *et al.* Central stress pathways in the development of cardiovascular disease. *Clin
Auton Res* **34**, 99-116, doi:10.1007/s10286-023-01008-x (2024).
109 Zouridis, S., Nasir, A. B., Aspichueta, P. & Syn, W. K. The Link between Metabolic
Syndrome and the Brain. *Digestion*, 1-9, doi:10.1159/000541696 (2024).
110 Petersen, M. *et al.* A latent clinical-anatomical dimension relating metabolic syndrome to
brain structure and cognition. *Elife* **12**, doi:10.7554/eLife.93246 (2024).
111 Buchanan, J. B., Sparkman, N. L. & Johnson, R. W. Methamphetamine sensitization
attenuates the febrile and neuroinflammatory response to a subsequent peripheral
immune stimulus. *Brain Behav Immun* **24**, 502-511, doi:10.1016/j.bbi.2009.12.008 (2010).

112 Kustrimovic, N., Marino, F. & Cosentino, M. Peripheral Immunity, Immunoaging and
Neuroinflammation in Parkinson's Disease. *Curr Med Chem* **26**, 3719-3753,
doi:10.2174/0929867325666181009161048 (2019).
113 Muñoz-Delgado, L. *et al.* Peripheral inflammatory immune response differs among
sporadic and familial Parkinson's disease. *NPJ Parkinsons Dis* **9**, 12, doi:10.1038/s41531-
023-00457-5 (2023).
114 Bratton, B. *et al.* Neural regulation of inflammation: no neural connection from the vagus
to splenic sympathetic neurons. *Experimental physiology* **97**, 1180-1185 (2012).
115 Ratcliffe, E. M., Farrar, N. R. & Fox, E. A. Development of the vagal innervation of the gut:
steering the wandering nerve. *Neurogastroenterol Motil* **23**, 898-911, doi:10.1111/j.1365-
2982.2011.01764.x (2011).
116 Chang, H. *et al.* Stress-sensitive neural circuits change the gut microbiome via duodenal
glands. *Cell* **187**, 5393-5412.e5330, doi:10.1016/j.cell.2024.07.019 (2024).
117 Powley, T. L. *et al.* Gastrointestinal projection maps of the vagus nerve are specified
permanently in the perinatal period. *Brain Res Dev Brain Res* **129**, 57-72,
doi:10.1016/s0165-3806(01)00183-3 (2001).
118 Murray, K., Barboza, M., Rude, K. M., Brust-Mascher, I. & Reardon, C. Functional circuitry
of neuro-immune communication in the mesenteric lymph node and spleen. *Brain Behav
Immun* **82**, 214-223, doi:10.1016/j.bbi.2019.08.188 (2019).
119 Dey, P. The Gut-Spleen Axis in Liver Disease: Mechanistic Insights Into Metabolism,
Metabolome, and Microbiome Interactions. *Faseb j* **39**, e71070,
doi:10.1096/fj.202502523RR (2025).
120 Wan, X. *et al.* Impact of broad-spectrum antibiotics on the gut-microbiota-spleen-brain axis.
Brain Behav Immun Health **27**, 100573, doi:10.1016/j.bbih.2022.100573 (2023).
121 Belkaid, Y. & Hand, T. W. Role of the microbiota in immunity and inflammation. *Cell* **157**,
121-141, doi:10.1016/j.cell.2014.03.011 (2014).
122 Fang, H. *et al.* Gut-Spleen Axis: Microbiota via Vascular and Immune Pathways Improve
Busulfan-Induced Spleen Disruption. *mSphere* **8**, e0058122,
doi:10.1128/msphere.00581-22 (2023).
123 Wei, Y. *et al.* Brain-spleen axis in health and diseases: A review and future perspective.
Brain Res Bull **182**, 130-140, doi:10.1016/j.brainresbull.2022.02.008 (2022).
124 Komegae, E. N. *et al.* Vagal afferent activation suppresses systemic inflammation via the
splanchnic anti-inflammatory pathway. *Brain Behav Immun* **73**, 441-449,
doi:10.1016/j.bbi.2018.06.005 (2018).
125 Buijs, R. M., van der Vliet, J., Garidou, M. L., Huitinga, I. & Escobar, C. Spleen vagal
denervation inhibits the production of antibodies to circulating antigens. *PLoS One* **3**,
e3152, doi:10.1371/journal.pone.0003152 (2008).
126 Lucido, M. J. *et al.* Aiding and Abetting Anhedonia: Impact of Inflammation on the Brain
and Pharmacological Implications. *Pharmacol Rev* **73**, 1084-1117,
doi:10.1124/pharmrev.120.000043 (2021).
127 Lalousis, P. A. *et al.* Anhedonia as a Potential Transdiagnostic Phenotype With Immune-
Related Changes in Recent-Onset Mental Health Disorders. *Biol Psychiatry* **96**, 615-622,
doi:10.1016/j.biopsych.2024.05.019 (2024).
128 Carpinelli, L. *et al.* Anhedonia in irritable bowel syndrome and in inflammatory bowel
diseases and its relationship with abdominal pain. *Neurogastroenterol Motil* **31**, e13531,
doi:10.1111/nmo.13531 (2019).
129 Krach, S., Paulus, F. M., Bodden, M. & Kircher, T. The rewarding nature of social
interactions. *Front Behav Neurosci* **4**, 22, doi:10.3389/fnbeh.2010.00022 (2010).
130 Torquet, N. *et al.* Social interactions impact on the dopaminergic system and drive
individuality. *Nat Commun* **9**, 3081, doi:10.1038/s41467-018-05526-5 (2018).

131 Manduca, A. *et al.* Dopaminergic Neurotransmission in the Nucleus Accumbens
Modulates Social Play Behavior in Rats. *Neuropsychopharmacology* **41**, 2215-2223,
doi:10.1038/npp.2016.22 (2016).
132 Han, M., Jiang, G., Luo, H. & Shao, Y. Neurobiological Bases of Social Networks. *Front*
Psychol **12**, 626337, doi:10.3389/fpsyg.2021.626337 (2021).
133 Yamaguchi, Y., Lee, Y. A. & Goto, Y. Dopamine in socioecological and evolutionary
perspectives: implications for psychiatric disorders. *Front Neurosci* **9**, 219,
doi:10.3389/fnins.2015.00219 (2015).
134 Milaneschi, Y. *et al.* Association of inflammation with depression and anxiety: evidence for
symptom-specificity and potential causality from UK Biobank and NESDA cohorts. *Mol*
Psychiatry **26**, 7393-7402, doi:10.1038/s41380-021-01188-w (2021).
135 Osimo, E. F. *et al.* Inflammatory markers in depression: A meta-analysis of mean
differences and variability in 5,166 patients and 5,083 controls. *Brain Behav Immun* **87**,
901-909, doi:10.1016/j.bbi.2020.02.010 (2020).
136 Isingrini, E. *et al.* Resilience to chronic stress is mediated by noradrenergic regulation of
dopamine neurons. *Nat Neurosci* **19**, 560-563, doi:10.1038/nn.4245 (2016).
137 Willmore, L., Cameron, C., Yang, J., Witten, I. B. & Falkner, A. L. Behavioural and
dopaminergic signatures of resilience. *Nature* **611**, 124-132, doi:10.1038/s41586-022-
05328-2 (2022).
138 Falvey, A. Vagus nerve stimulation and inflammation: expanding the scope beyond
cytokines. *Bioelectron Med* **8**, 19, doi:10.1186/s42234-022-00100-3 (2022).
139 Tsaava, T. *et al.* Specific vagus nerve stimulation parameters alter serum cytokine levels
in the absence of inflammation. *Bioelectron Med* **6**, 8, doi:10.1186/s42234-020-00042-8
(2020).
140 Madisen, L. *et al.* A toolbox of Cre-dependent optogenetic transgenic mice for light-
induced activation and silencing. *Nat Neurosci* **15**, 793-802, doi:10.1038/nn.3078 (2012).
141 Rock, C., Zurita, H., Wilson, C. & Apicella, A. J. An inhibitory corticostriatal pathway. *Elife*
5, doi:10.7554/eLife.15890 (2016).
142 Rock, C., Zurita, H., Lebby, S., Wilson, C. J. & Apicella, A. J. Cortical Circuits of Callosal
GABAergic Neurons. *Cereb Cortex* **28**, 1154-1167, doi:10.1093/cercor/bhx025 (2018).
143 Rock, C. & Apicella, A. J. Callosal projections drive neuronal-specific responses in the
mouse auditory cortex. *J Neurosci* **35**, 6703-6713, doi:10.1523/jneurosci.5049-14.2015
(2015).
144 Zurita, H., Feyen, P. L. C. & Apicella, A. J. Layer 5 Callosal Parvalbumin-Expressing
Neurons: A Distinct Functional Group of GABAergic Neurons. *Front Cell Neurosci* **12**, 53,
doi:10.3389/fncel.2018.00053 (2018).
145 Bertero, A. & Apicella, A. J. Distinct electrophysiological properties of long-range
GABAergic and glutamatergic neurons from the lateral amygdala to the auditory cortex of
the mouse. *J Physiol* **602**, 1733-1757, doi:10.1113/jp286094 (2024).
146 Bertero, A., Garcia, C. & Apicella, A. J. Corticofugal VIP Gabaergic Projection Neurons in
the Mouse Auditory and Motor Cortex. *Front Neural Circuits* **15**, 714780,
doi:10.3389/fncir.2021.714780 (2021).
147 Bertero, A., Zurita, H., Normandin, M. & Apicella, A. J. Auditory Long-Range Parvalbumin
Cortico-Striatal Neurons. *Front Neural Circuits* **14**, 45, doi:10.3389/fncir.2020.00045
(2020).
1544