Conservation implications of dietary strategies in the critically endangered

Acacia gazelle (*Gazella arabica acacia*)

<u>Abstract</u>

Animals consume different foods in specific quantities and ratios to meet a multi-

dimensional nutrient target that maximizes their fitness. Attempts to reach this target in an ever-

changing nutritional landscape often influence animal foraging behavior, habitat choice, and

food-web interactions. Consequently, understanding the nutritional ecology of a species is

instrumental in developing efficient management plans for its conservation. We quantified the

dietary considerations of a critically endangered population of Acacia gazelles (*Gazella*

arabica acacia) in Southern Arava, Israel, and compared their diet and nutrition to that of the

sympatric Dorcas gazelle (*Gazella dorcas*), using behavioral observations coupled by

nutritional and secondary metabolite quantification, and morphological measurements of trees.

Acacia gazelles mainly consumed resources from two umbrella-thorn acacia (*Vachellia tortilis*)

subspecies that differed in morphology, nutritional composition, and phenology. Gazelles

maintained a rather narrow nutritional intake target year-round despite the dynamic dietary

landscape, tightly regulated their protein intake, and prioritized consumption of non-structural

carbohydrates. They did so by adjusting the consumption of leaves, pods, and flowers,

following the differential phenology of the two *Vachellia* trees. The sympatric Dorcas gazelles

generally avoided the thorny *Vachellia* subspecies and consumed more halophytic shrubs and

less tree foliage than Acacia gazelles. Despite these dietary differences, the two gazelle species

converged to a similar intake target. We conclude that suitable habitats for Acacia gazelle

reintroduction must include sympatric populations of the two *V. tortilis* subspecies, that

nutritional deficiency is unlikely to play a major role in explaining the slow rate of population

recovery, and that attempts to limit Dorcas gazelle's population size may be redundant due to

the seemingly low competition between the two gazelle species.

Introduction

Wildlife feed on various foods in specific ratios and quantities to meet a multifaceted nutrient intake target that maximizes their fitness (hereafter "intake target"; Simpson & Raubenheimer 2012; Solon-Biet et al. 2014; Raubenheimer & Simpson 2018). Attempts to meet this target may regulate animals' spatial activity and behavior (Charnov 1976; Seagle & McNaughton 1992; Johnson et al. 2017), and influence interactions with competitors, natural enemies, and anthropogenic risks (Kohler & McPeek 1989; Pettersson & Brönmark 1993; Weber 2001; Kay et al. 2010). A key step in understanding the ecology of endangered species is thus to uncover their dietary considerations (Birnie-Gauvin et al. 2017).

Human activities have dramatically changed the natural environment, altered the abundance, quality, and distribution of food resources, and limited their accessibility to wild animals (Acevedo-Whitehouse & Duffus 2009). These changes may prevent animals from meeting their intake target, adversely affecting their population dynamics (Armstrong & Perrott 2000; López-Bao et al. 2010). Managers tend to address these general concerns using supplementary feeding as a precautionary measure. This is despite growing evidence that supplementary feeding may negatively affect the population it is intended to help (Carrete et al. 2006; Blanco et al. 2011; Martínez-Abraín & Oro 2013), or the recipient ecosystem at large (Cortés-Avizanda et al. 2009a, 2009b). Managers can minimize these double-sided risks by revealing the dietary strategies of wildlife in nature, and by integrating this knowledge into conservation plans.

Attempts to quantify nutritional needs and other dietary considerations of wildlife in the field traditionally focused on a single currency such as biomass, energy, or protein consumed (Schoener 1971; Charnov 1976; White 1993). However, natural foods are complex mixtures of macronutrients and micronutrients in varying levels of availability and digestibility. Animals often need to ingest several food types, and occasionally digestive assistant agents (DAA) to meet their multidimensional nutritional target (Zaguri et al. 2024). Thus, focusing on a single food currency may provide only a limited understanding of the animals' true dietary considerations (Belovsky 1984; Simpson et al. 2004; Felton et al. 2009a, 2018). Despite this well-accepted realization, the lack of adequate methods to quantify wildlife intake targets in the field largely prevents adopting this approach for conservation. Progress in nutritional ecology, and especially the development of the right-angled mixture triangle (RMT) method that uses proportions rather than amounts of foods consumed to quantify wildlife intake targets marked a breakthrough in our ability to use nutritional ecology to inform conservation

(Raubenheimer 2011; Raubenheimer et al. 2015). Yet despite the clear potential, RMT was

used almost exclusively in primate research (Raubenheimer et al. 2015; Johnson et al. 2017;

Takahashi et al. 2019; DePasquale et al. 2023), and only rarely to quantify intake targets of

endangered wildlife including ungulates (Raubenheimer et al. 2014; Spitzer et al. 2023). In

these rare cases, the limited ability to directly observe the animal feeding behavior and to

quantify the nutritional composition of foods has led to the development of proxy methods like

fecal (Nie et al. 2015; Hecker et al. 2021; Spitzer et al. 2023) or rumen and gut content analyses

(Felton et al. 2021) to quantify the diet, and the use of crude nutritional estimates at the species

level to assess the food nutritional content (Costello et al. 2016; Shrestha et al. 2020; Spitzer et

al. 2023, 2024). These proxy methods are informative but produce less accurate and temporally

dubious intake target estimates that are less applicable for conservation.

Focusing solely on the nutritional composition without considering other food

characteristics may produce an incomplete understanding of animals' dietary considerations

and may lead to erroneous conservation decisions. Plants are often defended by various

chemical (e.g., tannins; alkaloids; salts) and physical defenses (e.g., thorns; opaline phytoliths;

trichomes), and include components of little nutritional value (e.g., lignin, Barboza et al. 2009).

These traits may alter herbivores' ability to ingest and digest foods, possibly influencing the

herbivore's dietary choices (Freeland & Janzen 1974; Cooper & Owen-Smith 1986).

Consequently, attempts to uncover the dietary strategies of endangered ungulates must include

quantification of plant traits that may influence food choice.

Our goal was to assist in the recovery of a critically endangered Acacia gazelle (*Gazella*

arabica acacia) population by exploring the gazelle dietary strategy using classic and modern

approaches. Acacia gazelles are a genetically unique and geographically isolated population of

Arabian gazelles (*Gazella arabica*, Hadas et al. 2015; IUCN SSC Antelope Specialist Group

2017). This remnant population of about 40 individuals is restricted to two small floodplains

in the Yotvata Nature Reserve, Southern Arava, Israel, and struggles to recover despite

substantial conservation efforts like water, food supplementation, and interventions to reduce

predation and competition (Shalmon 1991; Blank 1996, 2000; Polak et al. 2023). Historical

references suggest a much larger distribution before the 1950s when the entire population was

almost extirpated by hunting (Mendelssohn 1974), Yet, not enough is known about the historic

distribution to predict these gazelles' habitat requirements. In 2006, the Israel Nature and Parks

Authority (INPA) enclosed the remaining 13 gazelle individuals within a 3.5 km^2 predator-

proofed enclosure together with 82 Dorcas gazelles (*Gazella dorcas*) that presumably compete

with the Acacia gazelles for food (Shalmon et al. 2020; Breslau et al. 2020). Thus, another goal
of the project was to compare the diets and intake targets of the two gazelle species.

To reveal their dietary strategy, we observed the Acacia gazelles' natural foraging
behavior in monthly intervals for one year and collected samples from all plants consumed,
and from similar reference resources that were not consumed. We used a similar protocol to
assess the Dorcas gazelle dietary strategy during the summer months. We characterized the
chemical composition of all samples using wet chemistry and Near Infra-Red Spectroscopy
(NIRS). We then used right-angled mixture triangles (Raubenheimer 2011) to estimate the
gazelle nutritional intake target. We measured the physical traits and tannin composition of the
two *Vachellia* trees to identify the traits of individual trees that make them more palatable. We
compared the Acacia gazelle diet and nutrition to those of the sympatric Dorcas gazelle to
reveal the degree of resource competition between the two species.

Methods
Study site and gazelle populations

The study was conducted within the Acacia gazelle predator-proofed enclosure in
Yotvata Nature Reserve, Israel (29°51'-29°53'N, 35°01'-35°03'E; Fig. 1). This hyper-arid site
is located within the Jordan Rift Valley about 40 km north of the Gulf of Aqaba. The mean
annual temperature is 24°C, and the average annual precipitation is 40.5 mm with big
fluctuation between years. All precipitation occurs between October to May. The site includes
two alluvial fans dominated by scattered *Vachellia* (formerly *Acacia*) trees and their common
semi-parasite *Loranthus acaciae*. Several halophyte shrubs, mainly *Haloxylon salicornicum*,
Lycium shawii, *Salsola imbricata,* and *Nitraria retusa*, dominate the exposed stony gaps
between the *Vachellia* stands (Baharav 1982). Two *Vachellia* trees coexist in this site, both
considered subspecies of *V. tortilis (V. tortilis tortilis and V. tortilis raddiana). Locally, the two
subspecies exhibit consistent differences in their chemical and physical traits, and habitat
requirements (Halevy & Orshan 1972; Rohner & Ward 1997). Moreover, the two trees have
very distinct phenology, producing flowers and fruits at different times (Halevy & Orshan
1973). Thus, we treated the two *Vachellia* trees as two distinct resources hereafter *V. tortilis*
and *V. raddiana*.

The Acacia gazelle population of 35 individuals was first discovered at our study site
in 1964. The population was discovered only after all other Acacia gazelle populations in Israel
were extirpated. Since then, the INPA has attempted to restore the population using various

management interventions like supplementary feeding with natural or artificial foods, predator
culling, and removal of calves in an attempt to establish a breeding colony in captivity. Despite
those substantial efforts, the population failed to recover. In 2006, after a dramatic population
decline, the INPA enclosed the remaining 13 gazelles together with 82 Dorcas gazelles (*Gazella*
dorcas) in a 3.5 km^2 predator-proofed enclosure. By 2013, both Acacia and Dorcas gazelle
populations reached a maximum density of 42 and 173 individuals, respectively. However, in
the winter of this year, large parts of the fence were breached by flood water, enabling wolves
to penetrate and kill about 30 Acacia gazelles and 65 Dorcas gazelles. Between 2015 to 2019
the INPA actively removed 161 Dorcas gazelles from the enclosure as a precautionary step
against interspecific competition, leaving only 36 individuals in the enclosure. During 2022-
2023 only about 30 Acacia gazelle and about 40 Dorcas gazelle inhabited the enclosure (Polak
et al. 2023).

Fig. 1 Study site locality map showing the location of Yotvata NR (a), and the gazelle enclosure (b). the enclosure (bottom left) is presented with emphasis on the northern part (enlarged and highlighted) where observations were conducted. Marked in the northern part are the trees observed and sampled, colored by *Vachellia* subspecies, and the two hides used for observations. The Internal hide was used during morning sessions and the external hide was used during evening sessions.

 Gazelle feeding observations and forage sampling

To assess the gazelle diet, we conducted monthly group observations of the gazelles'
feeding behavior in the northern part of the gazelle enclosure during 2022 and 2023. We
observed the gazelles' behavior during their peak foraging activity in the early morning and
late afternoon for a minimum of 3 consecutive days each month. For each individual plant
resource, we recorded the number, age, and sex of the feeding gazelles, the feeding time, and
feeding behavior (either from the ground or from the canopy). We used QField mobile
application to collect this data with its location into a QGIS project (see Appendix 1 for full
protocol and details).

To compare the foraging behavior of Acacia gazelles to that of Dorcas gazelles, we used
observational data on the foraging behavior of the two gazelle species that we collected at the
same site during the summer of 2014, prior to the Dorcas gazelle removal. These foraging
observations were conducted using continuous time-budget observations on focal individuals.

After each observation, we collected samples from all plant resources that were
consumed by the gazelles and immediately transferred them to our field lab. There, we weighed
and oven-dried the samples (60°C at least 48 h), and re-weighed them for dry mass estimations.
Following each observation session, we milled all samples through a 1 mm sieve using a
laboratory grinding mill (Kinematica POLYMIX PX-MFC 90D). We stored the milled samples
in the freezer until further analyses.

Chemical analyses

We used classic chemical quantification methods to estimate the nutritional content of
209 out of 400 collected food samples (Table S1). Samples were chosen based on quantity and
Near Infrared Spectroscopy (NIRS) unique spectra. Each of the 209 samples was analyzed for
total non-structural carbohydrates, crude protein, lipids, fiber fractions, ash, and tannin content.
The non-structural carbohydrates were quantified using the phenol-sulfuric assay with acid
extraction (Dubois et al. 1956; Landhäusser et al. 2018). For crude protein estimation, we
estimated the Nitrogen (N) content by combustion, using a Leco FP-528 Nitrogen Analyzer
(Michigan, USA), and multiplied the N content by the widely used 6.25 conversion factor
(Jones 1931; Rothman et al. 2012). Lipids were measured using ether extract in an ANKOM
Fat Analyzer (New York, USA). Fiber fractions were determined using the sequential detergent
fibers assay (Van Soest et al. 1991) in an ANKOM A200 Fiber Analyzer: Neutral Detergent
Fibers (NDF), which contains non-soluble fibers (hemicellulose, cellulose, and lignin), Acid

Detergent Fibers (ADF, cellulose and lignin), and Acid Detergent Lignin (ADL). Samples were burned at 500–550°C for ash measurement (Rothman et al. 2012). Finally, the relative content of condensed tannins was estimated using the acid-butanol assay after extracting samples with 70% acetone (Porter et al. 1985; Hagerman & Butler 1994; Rothman et al. 2009). Additionally, we chose 56 samples of the food resources observed in 2014 (8-10 samples of each food resource) for mineral quantification to assess the ash composition. Minerals were assessed via atomic absorption spectroscopy after microwave digestion at Dairy One Commercial Forage Laboratory (Ithaca, NY, USA). Nutrient estimations are presented on a 105°C dry matter basis.

We then constructed a NIRS model for *Vachellia* leaves and pods of the two subspecies. *Vachellia* flowers, *Loranthus* leaves and flowers and perennial shrubs were analyzed only by wet chemistry and average nutrient content was applied, due to the low number of samples.

For NIRS analyses, we dried all samples for 48h. After homogenizing the sample, we randomly took a ~50 mg subsample and added it to a 1 ml clear glass shell vial. We then scanned all samples, using an Antaris II FT-NIR analyzer. We constructed two NIRS models using TQ analyst software, the first model included absolute concentrations of all measured macro-nutrients (Protein, Lipids, NSC, NDF, ADF, and ADL). The second model was for the relative concentration of tannins. NIRS model's protocol and performances are presented in Appendix 2.

Physical measurements of *Vachellia* trees

We randomly chose 20 *V. tortilis* and 20 *V. raddiana* trees in the northern part of the enclosure. Half of the trees were observed browsed by Acacia gazelles during the winter when mainly foliage is available. We randomly chose the 10 reference trees from all trees for which we did not detect feeding in the previous year. For each tree, we divided the canopy into 4 equal-sized sections and measured 2 randomly chosen branches in each section. All sampled branches were within the feeding range of an adult Acacia gazelle (up to 1.7 m, Fig. S1). Following the protocol described by Rohner & Ward (1997), on each branch, we counted all leaves and thorns >1mm within 50 cm from the branch distal end. We classified thorns as either straight (spines) or curved (prickles) and the lengths of all thorns >5mm were measured perpendicular to the branch. A sample of the 5 biggest leaves was measured for length and width as well. We also noted the existence of browsing marks on each branch.

 Data analysis

We estimated the Acacia gazelles' nutritional target and resources' composition using a
right-angled mixture triangle (Raubenheimer 2011), of the relative contents of the three main
macro-nutrients in the gazelle diet – Protein, non-structural carbohydrates (NSC), and
digestible fibers (calculated as NDF-ADL). We calculated the daily nutritional target by
multiplying each observation unit's (plant) nutritional values with the time and number of
gazelles spent feeding from it divided by the total daily feeding time. To unravel gazelle
regulation and prioritization of different macro-nutrients we conducted a permutation test for
the gazelles' intakes. We summarized each food resource mean nutritional concentrations and
its total feeding time in each observation session. Then, we permuted these feeding times. To
evaluate seasonal variation in gazelles' intake of NSCs and digestible fibers we performed a
Pearson's correlation test and fitted a linear model of the temperature effect. We obtained the
mean monthly minimal temperature from the daily ambient temperature measurements of the
nearest meteorological station in Yotvata, from August 2021 to April 2022 ("Meteorological
Database | Israel Meteorological Service" 2023).

We performed a Kruskal-Wallis test followed by a Bonferroni-corrected Conover-Iman
post-hoc test (Conover & Iman 1979) to compare the means of all macro-nutrients, minerals,
and tannins contents of the most prominent resources in the diets of both gazelle species as
observed in 2014 (i.e., *Vachellia* leaves and pods, *Haloxylon salicornicum*, and *Loranthus*
acacia leaves). We calculated fibers nutritional values (i.e. Hemicellulose, cellulose, and
lignin) by subtracting the chemical sequential fractions (NDF-ADF, ADF-ADL, ADL,
respectively). Since subtraction introduces a greater protentional error we report also the raw
values in the supplementary, where applicable.

For the physical traits measured, we averaged all traits for all branches of each tree. We
analyzed thorns data using counts within two length categories (with a 5 mm cutoff), resulting
in four thorns categories. Nutritional data were obtained for each tree from the NIRS model
predictive results, except for two *V. raddiana* trees which did not have nutritional data and were
therefore removed from this analysis. We used PCA to evaluate inter-specific variance between
Vachellia subspecies. To uncover differences between browsing categories we applied a
separate PLS Discriminant Analysis (PLS-DA, Barker & Rayens 2003) on each *Vachellia*
subspecies since the PCA revealed a large inter-specific variability. We standardized the data
before both ordinations. To estimate whether the separation between the two feeding categories

in the PLS analysis is significant, we performed two permutation tests on the feeding categories. First, we tested for cluster separation using the Mahalanobis distance between the two categories' centroids based on the pooled covariance matrix. We also tested for significance the proportion of explained variance between the feeding categories (R^2_y) and DR^2_y, correcting for false-negatives resulting from out-of-bounds estimation (Westerhuis et al. 2008). We used 1000 permutations and defined the p-value as the fraction of permutations greater than the chosen statistic. Last, we estimated each variable importance by randomly permuting it 1000 times and conducting the same analyses described. We compared the means of permutations R^2_y, DR^2_y, Mahalanobis distance between clusters, and the fraction of significant permutations for cluster separation.

All statistical analyses were conducted in R 4.3.0 software (R Core Team 2023). Linear models and correlation tests were performed using 'lm' and 'cor.test' functions from the 'stats' package. Kruskal-Wallis and Conover's tests were performed using 'kruskalTest' and 'kwAllPairsConoverTest' from 'PMCMRplus' package. PCA was fitted using 'prcomp' from 'stats' and PLS-DA was fitted using 'plsda' from the 'mixOmics' package (Rohart et al. 2017).

Results

Six different plant species were consumed by Acacia gazelles during the study. Their diet consisted mainly of different resources from two *Vachellia* subspecies, which together constituted about 88% of the annual gazelle diet, and no less than 70% of the monthly diet (Fig. 2). The consumed resources differed substantially in their nutritional and mineral content (Kruskal-Wallis $p < 0.001$ for all, see tables S3 and S4 for post-hoc comparisons).

These two *Vachellia* subspecies differ substantially in their chemical and physical characteristics. *V. raddiana* trees have bigger leaves, while *V. tortilis* trees have higher densities of prickles, and higher concentrations of proteins and lipids in the leaves (Fig. S2). PCA results revealed high intra-specific variability in spines counts, and the contents of NSC, fibers, and tannin in the leaves of both subspecies (Fig. S2).

Vachellia subspecies differed also in their phenology. *V. tortilis* trees flowered while shedding their leaves in early summer, whereas *V. raddiana* trees started flowering in late summer and throughout the fall without completely shedding their leaves. *V. tortilis* pods developed immediately after the blooming period and ripened in late summer, while *V. raddiana* green pods remained on the branches during winter and ripened only in the next spring (Fig. 3c). *V. tortilis* produced flowers and fruits in shorter and more synchronized time

windows than *V. raddiana* (Fig. 3c). The rate of gazelles' visits to the *Vachellia* trees, and their

relative feeding time were tightly associated with the above-listed phenological differences

(Fig. 3). For instance, gazelles seemed to prioritize feeding on flowers when those were

available. During June while green pods of *V. raddiana* were still available, the gazelles

switched to feeding almost exclusively on the *V. tortilis* flowers during the peak blooming

period (Fig. 3).

The variation in the gazelle's monthly nutritional intake was much smaller than the

variation in the nutritional landscape (Fig. 4, Table S3). Gazelles achieved this tighter balance

by eating differentially from the different available food resources. For example, in February

the gazelles ate *V. raddiana* leaves from the canopy and pods from the ground, as well as *L.*

Shawii leaves in order to reach their nutritional target (Fig. 2, Fig. 4, Fig. S3). This tight intake

target was reached despite substantial differences in resource availability (Fig. 3, Fig. 4). From

all macro-nutrients measured, protein intake varied the least (Table S3). The permutation test

shows no significant selection for protein-rich foods ($p=0.43$). Thus, the precise protein intake

may reflect the smaller variation in protein compared to other macro-nutrients in the consumed

resources (Fig. S5). Similarly, we observed no selective feeding for digestible fibers (i.e.

hemicellulose+cellulose, $p=0.29$, Fig. S5). However, the permutations test found that gazelles

strongly select for non-structural carbohydrates, as the observed intake was significantly higher

than the expected intake by random feeding ($p=0.11$, we consider this a significant result since

we permuted the intakes between consumed resources, depicting evolutionary resource

selection).

We observed a negative correlation between the monthly intake of NSC and digestible

fibers ($r=-0.54$, 95% CI [-0.85,0.043], $t_{10}=-2.0263$, $p=0.07$; Fig. 5a). Gazelles tended to

consume higher daily proportions of digestible fibers as monthly temperature decreased

(slope=-0.22 [-0.47,0.034], $R^2=0.1$, $F_{1,28}=3.143$, $p=0.087$ for both). In the coldest months

(January-February) gazelles increased both NSC and fiber intakes (Fig. 5a). Using NDF

measure showed similar patterns for temperature effect on NDF consumption (slope=-0.30 [-

0.53, -0.062], $R^2=0.2$, $F_{1,28}=6.725$, $p=0.015$ for both) and NDF to NSC trade-off ($r=-0.55$ [-

0.85,0.034], $t_{10}=-2.0866$, $p=0.064$). This seasonal fiber intake was not confounded by available

resources as different ratios could have been achieved in each season (Fig. 5b). In March,

gazelles consumed a high NSC to digestible fibers ratio that substantially deviated from this

general pattern. This was possibly because gazelles consumed large quantities of *V. tortilis*

leaves during this time (Fig. 2). Leaves and pods had a relatively low protein content during

March than in other months, but *V. tortilis* leaves had proportionally the highest protein content

(Fig. S4), suggesting that attempts to regulate protein intake may cause this dietary change.

During summer, gazelles increased their relative NSC intake by prioritizing *V. tortilis*

carbohydrates-rich flowers (Fig. 2, Fig. 4). Still, gazelles kept their tight protein intake by

consuming *V. raddiana* pods and *Vachellia* leaves from both subspecies.

We observed variation in the use of individual *Vachellia* trees, of both subspecies, where

some trees were heavily browsed while others were ignored (Fig. S6). The browsed and un-

browsed *V. tortilis* trees differed substantially (Fig. 6a, M_d=3.28, p=0.014). The variation in the

browsing category explained by the model was 75%-78%, depending on the type of fit-

measure, and was significantly higher than random for both fit-measures (R^2_y=0.75, DR^2_y=0.78,

p=0.021, 0.019, respectively). The separation was mainly along the first ordination axis (LV1)

driven predominantly by the high fiber fractions and bigger leaf size (expressed as width and

length measures) of the un-browsed trees, and the higher protein as well as higher number of

leaves of the browsed trees (Fig. 6a,c). Browsed trees were also characterized by having a lower

density of big prickles, and un-browsed trees had a relatively higher density of small prickles.

Hemicellulose, big prickles, and protein were the variables that their randomization had a major

effect on the model performance and cluster separation (Table S5). We did not find significant

differences between the browsed and un-browsed *V. raddiana* trees (M_d=2.74, R^2_y=0.67,

DR^2_y=0.72, p=0.14, 0.13, 0.11, respectively; Fig. 6b). Using raw analytical fiber fractions

instead of the nutritional calculated fractions slightly decreased the performances of both

models, but it remained significant for *V. tortilis* (Fig. S7).

We found major differences between the diet compositions of the two gazelle species.

Acacia gazelle foraged predominantly on two *Vachellia* subspecies while Dorcas gazelles' diet

included more shrubs, especially *Haloxylon salicornicum*. When feeding from the two

Vachellia trees, Acacia gazelles consumed a much higher proportion of *V. tortilis* in comparison

to Dorcas gazelles which fed almost exclusively from *V. raddiana* (Fig. 7a). Acacia gazelles

fed more on the foliage from both *Vachellias*, while Dorcas gazelles fed predominantly on *V.*

raddiana pods and flowers from the ground. Additionally, Acacia gazelle fed on *Loranthus*

acacia while Dorcas gazelle did not use this resource (Fig. 7a). Despite the large dietary

differences the two Gazelle species converged to a similar intake target. Both gazelle species

achieved very similar protein intake, but Acacia gazelles reached higher NSC and lower

digestible fibers than *G. dorcas* (Fig. 7b). Our two independent measurements of Acacia

Gazelle intake targets in the presence of high (2014) and low (2022) sympatric Dorcas gazelle

densities revealed very similar estimations (Fig. 7b), suggesting that intra-specific competition
had minimal influence on Acacia gazelle dietary considerations.

We observed consistent differences in ash and tannin levels between the diets of the
two gazelle species. The halophyte *H. salicornicum* which constituted a major part of Dorcas
gazelle's diet, had higher Na, Mg, and K content than all other plant resources (Table S4). The
differences in the concentration of these salts likely explain why ash content in *H. salicornicum*
was much higher than in other plant resources (*p*<0.03; Table S3). Leaves of the two *Vachellia*
trees had significantly higher tannin content compared to all other resources (*p*<0.001 for all;
Table S3), suggesting that Acacia gazelles potentially consumed higher quantities of tannins
and less salt than Dorcas gazelles due to higher ingestion of *Vachellia* leaves and lower
ingestion of *H. salicornicum*.

Fig. 2 Distribution of different food resources utilization by Acacia gazelles throughout the year. Proportions were calculated based on the time spent feeding from each resource.

Fig. 3 Acacia gazelles' response to *Vachellia* phenological changes. (a) the number of different trees utilized of each subspecies out of the total trees of that subspecies identified in the northern part of the enclosure. (b) proportional time spent each month feeding from the two *Vachellia* subspecies. (c) Phenological state throughout the year of both *Vachellia* subspecies.

Fig. 4 Right-angled Mixture Triangle (RMT) of the relative concentrations of the three major macro-nutrients in the Acacia gazelles' diet – Digestible fibers (hemicellulose+cellulose), non-structural carbohydrates (NSC), and Protein. Diagonal (Z) axis values are written in italics above the X-axis. Points are the mean monthly intake colored by month and the mean annual intake in black. In gray are the mean annual nutritional values of the different food resources in the gazelle diet. Ellipses represent 95% confidence interval of the mean intake in black, and the mean nutritional value of all resources in gray.

Fig. 5 Relationships between temperature and the intake of fiber and NSC. (a) NSC to fibers monthly intake colored by the monthly mean minimal temperature. Dashed lines are the annual nutritional target of each macro-nutrient (b) NSC to fiber mean intake and food resources available for the two coldest and two hottest months. Error bars are the standard error (n=4 days in summer and winter).

Fig. 6 PLS-DA ordination results. a;b. biplots for both *Vachellia* subspecies. Gray arrow sizes are equivalent to the loading scores for each ordination variable. Percentages on the axes are the variations explained by each. c;d. Loading scores of the first component are colored by the feeding category in which the median is maximum for each variable.

Fig. 7 Comparison of diets (a) and nutritional targets (b) of the two gazelle species. Colored intakes are calculated based on 2014 dietary observations and 2021-2022 mean resources nutritional content for August-October. The black intake and gray food resources are the same as in Fig. 4.

Discussion

Our goal was to reveal the dietary considerations of the critically endangered Acacia

gazelle in an attempt to assist population recovery and identify suitable habitats for

reintroduction. We found that Acacia gazelles feed almost exclusively on resources from two

Vachellia trees that differ in their phenology, nutritional composition, and defensive traits. The

Acacia gazelles maintained a tight nutritional intake year-round, despite an ever-changing

broader nutritional landscape. Specifically, Acacia gazelles regulated protein intake and

prioritized consumption of non-structural carbohydrates (NSC). We also found that Acacia

gazelles increased their fiber consumption in the cold season at the expense of NSC

consumption. Gazelles preferred browsing on *V. tortilis* individuals with higher protein content,

more leaves, and lower density of big prickles over trees with more fibers, bigger leaves, and

high density of small prickles. Despite a similar preference pattern, we failed to identify what

dictates the differential use of *V. raddiana* trees. The diet of Acacia gazelles differed

substantially from that of the sympatric Dorcas gazelles. While Acacia gazelles used the two

Vachellia subspecies and fed from their tree canopy, Dorcas gazelles' diet included more

halophytic shrubs and almost no *V. tortilis* resources. Dorcas gazelles ate less from the canopy

compared to Acacia gazelles, and used *V. raddiana* resources primarily by collecting them from

the ground. Consequently, Acacia gazelle consumed more tannin-rich foods and had to deal

with the high density of prickles and thrones, while Dorcas gazelles consumed more salt-rich

foods.

Acacia gazelle kept a precise multidimensional nutritional intake across seasons despite

a broader nutritional landscape and large seasonal fluctuations in the availability of food

resources. Protein intake was about 17% across seasons and showed the least variation in

comparison to other nutrients. To our best knowledge, such tight regulation of protein

consumption in wild herbivores was documented only in primates (Felton et al. 2009c, 2009b;

Hou et al. 2021; Takahashi et al. 2021). A common presumption in the herbivore-plant literature

is that herbivores must prioritize the consumption of protein-rich foods to fulfill their protein

demand (Milton 1979; Mattson 1980; White 1993). This presumption is based on the

acknowledged mismatch between the low protein content of many plant resources and the

herbivores' high physiological protein demand. We, however, found no evidence that gazelles

attempted to maximize protein ingestion to achieve the precise protein intake. We, therefore,

suggested that similarly high protein content in many available food resources allows these

gazelles to maintain tight protein intake while prioritizing the consumption of other important

nutrients (Codron et al. 2007; Felton et al. 2016; Ganzhorn et al. 2017).

Acacia gazelles prioritized consumption of NSC-rich foods, such as Vachellia flowers,

and aimed to increase NSC intake whenever possible. Prioritization of non-protein energy over

protein was observed in primates (Rothman et al. 2011) and in other ungulates (Asher et al.

2011; Felton et al. 2016), However, these studies found that attempts to ingest more NSC-rich

foods come at the expense of protein regulation. We found that Acacia gazelle can

simultaneously maintain a precise protein intake while prioritizing NSC intake. To increase

NSC intake, gazelles must feed on both *Vachellia* subspecies that provide flowers and ripe

fruits at different times. Consequently, a good habitat for Acacia gazelles must include

sufficiently large populations of both *Vachellia* subspecies. *V. raddiana* and *V. tortilis* co-occur

mainly along the Jordan Rift Valley, south of the Dead Sea (Halevy & Orshan 1972). This may

explain why the historic distribution of Acacia gazelles was limited to this area (Mendelssohn

1974), while the distribution of the sympatric Dorcas gazelle is much wider and includes

diverse desert habitats (Yom-Tov 2016).

The general pattern of maximizing NSC consumption was slightly disrupted by the

seasonal changes in fiber consumption. The relative intake of these two carbohydrate sources

varied across seasons in association with the monthly average minimal temperature and was affected by resource availability. Gazelles consumed proportionally more structural carbohydrates (digestible fibers; NDF) in the winter and increased intake of non-structural carbohydrates in the warmer seasons. These dietary changes seem intentional since the nutritional landscape allowed tighter regulation of fibers and NSC in both seasons. Our findings coincided with ungulate studies that found higher fiber consumption during cold periods. These studies hypothesized that the increase in fiber consumption is an undesirable outcome of resource scarcity or a costly byproduct of attempts to prioritize consumption of other nutrients (Stolter et al. 2005; Verheyden-Tixier et al. 2008, for example). This realization builds upon a common assumption of browsing theory that higher fiber content characterized by low digestible energy has lower physiological value for herbivores (Milton 1979; Van Soest 1994). Instead, we suggest that increased fiber consumption in cold periods may assist gazelles to thermoregulate. Fermentation of structural carbohydrates produces more heat than that produced by digestion of simple carbohydrates (Armstrong & Blaxter 1956; Cowsert & Montgomery 1969; Reynolds et al. 1991; Van Soest 1994). For example, the heat increment of feeding in white-tailed deer fawns can substitute up to 100% of their thermoregulation requirements. This effect was stronger for fawns that were fed on natural fiber-rich browse than those fed on NSC-rich pellets (Jensen et al., 1999). The domestic livestock literature supports this notion and recommends using lower fiber-to-NSC feed ratio as a way to improve livestock thermoregulation efficiency during warm seasons (John & Webster 1978; West 1999; Conte et al. 2018). Understanding the dietary considerations that are influenced by variations in animals' thermoregulatory needs is crucial for designing artificial diets and preventing adverse consequences of unsuitable supplementary feedings. Future studies should focus on the mechanistic details of this potentially important and thus far largely neglected dietary consideration by manipulating the feed under various climatic conditions and measuring the herbivores' dietary and physiological responses.

When examining diets and food preferences researchers often use resource species as a basic unit neglecting intraspecific variability that could be substantial (Lawler et al. 1998; Chapman et al. 2003; Rothman et al. 2012). Acacia gazelles chose to forage on specific trees and ignored others suggesting an intraspecific preference. The preferred *V. tortilis* trees were richer in protein and carbohydrate relative to fibers and had high densities of small leaves and lower densities of prickles relative to trees that were not used. We hypothesized that in their attempt to improve foraging efficiency gazelles consider both the nutritional intake and the

feeding costs. Understanding palatability variation across trees is necessary when estimating an area carrying capacity since not all potential resources are being used in similar ways (DeGabriel et al. 2014). Moreover, the nutritional content and chemical and physical defenses are plastic traits that can vary with age and level of browsing pressure that the tree experienced (Gowda 1997; Rohner & Ward 1997; Fornara & Du Toit 2007; Nzimande et al. 2022). High browsing pressure may reduce tree recruitment, altering the stand age structure and may force trees to invest more in anti-browsing defenses. These changes may generate a positive feedback loop in which gazelle may respond to these changes by browsing more on less defended trees. Thus, gazelle may experience nutritional limitations even if the number of trees and the foliage availability may erroneously suggest ample forage. We suggest that nutritional and defense traits of trees should be monitored to prevent such convoluted density-dependent reduction in resource quality and accessibility. This recommendation is especially relevant when the population is limited in space.

We found similar intraspecific variation in the use of *V. raddiana* trees but failed to identify what traits are associated with the gazelle preference. *V. raddiana* has higher phenolics and condensed tannins concentrations than *V. tortilis* (Rohner & Ward., 1997), suggesting those chemical compounds may play an important role in *V. raddiana*'s defense against herbivory and may affect gazelles' browsing preference in this subspecies. Our measurements of defense compounds did not allow testing this plausible hypothesis as a more elaborate secondary metabolite profile is required. *V. raddiana* has lower thorn densities and a much bigger leaf size than *V. tortilis*. Thus, these traits may play a much lesser role in predicting gazelle foraging preference in this subspecies. Blank (2005), who also noticed intraspecific variation in *Vachellia* use by Acacia gazelles, attributed this variation to differences in resource abundance (i.e., quantities of flowers and pods). We only used trees that were browsed for leaves during winter to minimize the possible effects of these confounding factors. Nevertheless, resource quantity may still affect our results regarding *V. raddiana* as dry pods can still be present below trees during winter. Future work should consider these potential sources of variation when exploring the intraspecific variation in *V. raddiana* use by Acacia gazelle.

The diet of the sympatric Acacia and Dorcas gazelles varied substantially, suggesting limited interspecific competition for food resources. Our continuous observation of Acacia gazelle individual-foraging from the summer of 2014 revealed a similar diet composition and nutritional intake to that observed in the summer of 2022, using a different observation approach. We found that Acacia gazelles consumed relatively high proportions of foliage from

the two *Vachellia* subspecies, ate the abundant parasitic plant *Loranthus acaciae,* and ingested pods of the two *Vachellia* subspecies in similar proportions. In comparison, the diet of the sympatric Dorcas gazelle was dominated by *V. raddiana* pods collected from the ground, a much lower proportion of *V. tortilis* pods, and just a small portion of *V. raddiana* leaves. Halophytic shrubs constituted 30% of the Dorcas gazelle's diet in comparison to only 7.2% percent in the Acacia gazelle's diet. Despite these large dietary differences, the protein intake of the two gazelle species converged to similar values and the intake of carbohydrate sources differed only slightly between the two gazelle species, suggesting similar physiological needs.

Similar physiological needs of sympatric species that are met by ingestion of different diets may suggest dietary niche separation that requires specific adaptations. Acacia gazelles forage high in the canopy by standing only on the hind legs or by leaning with the front legs on low branches (Blank 2005). These gazelles have to cope with high tannin levels and presumably other chemical defenses and avoid the thorns and prickles, especially those of *V. tortilis* that may limit foraging efficiency. On the other hand, Dorcas gazelles mainly consume halophytic plants and dry resources found under *Vachellia* trees and largely ignore resources from *V. tortilis* canopies. Consequently, needing to cope primarily with high salt levels and fluctuating and unpredictable food availability. Altogether our results suggest that the two sympatric gazelle species evolved to consume different diets. Our findings that Acacia gazelles' diet and nutritional intake did not differ before and after the sharp drop in the sympatric Dorcas gazelle population density coincided with this conclusion, suggesting limited inter-specific competition between the two gazelle species.

Our comparative observations were conducted during summer and fall. Thus, our inference of low interspecific competition between the two gazelle species may be limited to these seasons. We also found that Acacia gazelles increased consumption of *V. raddiana* in the winter, possibly intensifying competition with the Dorcas gazelles. However, previous work showed that Dorcas gazelles tend to substantially decrease the use of *V. raddiana* during the winter (Baharav 1980, 1982), supporting our conclusion of dietary niche separation between the two gazelle species. Acacia gazelles specialize in feeding from *Vachellia* trees and switch between subspecies and their resources to maintain a tight nutritional target. Tracking both *Vachellias'* phenology may allow this gazelle species to avoid seasonal nutritional shortages in this unpredictable hyper-arid environment. Moreover, *Vachellia* trees supply resources that are less prone to fluctuations in availability between years compared to annual plants or small perennial shrubs, which other herbivores rely upon. Stability in resource quality and quantity

may support a high reproductive rate allowing Acacia gazelles to calve in all seasons and even

twice a year (Shalmon 1991; Mendelssohn et al. 1997; Blank 2000), as we have also observed.

Other gazelle species, including the sympatric Dorcas gazelle, exhibit a synchronized seasonal

reproduction peak that is tightly timed to match the annual peak in resource availability

(Baharav 1983; Yom-Tov 2016). Thus, Acacia gazelles may have greater reproductive potential

than Dorcas gazelles, but, are limited to a more specific habitat where the two *Vachellia*

subspecies are present.

Conclusions and management implications

We used a nutritional geometry approach to uncover Acacia gazelles' dietary

considerations in attempts to guide this critically endangered population's recovery. Acacia

gazelles forage predominantly on two *Vachellia* subspecies that provide crucial resources at

different times. Acacia gazelle's macro-nutritional intake varied little between months in 2021-

2022 and was similar to that observed in the summer of 2014. This enforces confidence in the

estimated intake representing nutritional needs and suggests that since Acacia gazelles were

able to reach it year-round, currently, the population does not suffer from a nutritional

deficiency that requires supplement feeding. However, in case of future needs, identifying this

precise intake target may enable managers to provide feed that meets the species-specific needs.

Our findings that gazelle alternate fiber and NSC, possibly to improve thermoregulation

efficiency may further assist in this effort and should be considered whenever exploring the

nutritional ecology of wild ungulates. Acacia gazelles' specialization on *Vachellia* resources

suggests a higher reproductive potential than that of other desert ungulates. This specialization,

however, also poses a potential obstacle to the population's recovery, since the two *Vachellia*

subspecies coexist only in alluvial fans along the Jordan Rift Valley, south of the Dead Sea.

This restricted habitat is diminishing in an unprecedented way due to agricultural expansion.

Reintroductions of Acacia gazelles may thus necessitate a broader conservation plan to expand

the protection of their unique habitat. It was hypothesized that similar physiological needs may

induce intense interspecific competition for food between the two sympatric gazelles (Shalmon

et al., 2020). Instead, our findings suggest Acacia gazelles feed a different diet than Dorcas

gazelles and overall experience low interspecific competition that does not affect their diet or

nutrition. Thus, we recommend postponing the seemingly unjustified removal of Dorcas

gazelles from the enclosure that may counterintuitively harm the Acacia gazelles by denying

potential anti-predatory benefits (e.g., social vigilance, dilution effect) and by potentially

reducing selection against plant species that are less important for this species. We also found

that Acacia gazelles selectively feed on certain *Vachellia* individuals and identified the possible traits that affect this selection for *V. tortilis* but not for *V. raddiana*. Since trees subject to different browsing pressures may change the expression of these traits, continuous monitoring is important. We recommend monitoring the *Vachellia* recruitment, canopy height, and defensive traits, to prevent density-dependent limitations and identify otherwise obscured reductions in resources' accessibility and quality. Our case study highlights the benefit of adopting a nutritional ecology approach to guide management plans for endangered wildlife.

Acknowledgments:

We thank Tal Polak, Golan Reider, Eran Gissis, and Mor Ochnuna from INPA's Southern Arava district for their help throughout this project and especially in facilitating field work; Jenny Paltan, Shani Kandel, and Netta Shamir-Weller for their crucial help with nutritional analysis; and Liran Carmel and Michael Kalyuzhny for statistical advice.

Funding

This article was supported by an Israel Nature and Parks research grant to D.H. Hunter College provided funding to J.M.R. for a portion of the nutritional analysis.

Literature cited

Acevedo-Whitehouse K, Duffus ALJ. 2009. Effects of environmental change on wildlife health. Philosophical Transactions of the Royal Society B: Biological Sciences 364:3429–3438. Royal Society.

Armstrong DG, Blaxter KL. 1956. Heat Increments of Feeding in Ruminants. Nature 177:1183–1184. Nature Publishing Group.

Armstrong DP, Perrott JK. 2000. An Experiment Testing whether Condition and Survival are Limited by Food Supply in a Reintroduced Hihi Population. Conservation Biology 14:1171–1181.

Asher GW, Stevens DR, Archer JA, Barrell GK, Scott IC, Ward JF, Littlejohn RP. 2011. Energy and protein as nutritional drivers of lactation and calf growth of farmed red deer. Livestock Science 140:8–16.

Baharav D. 1980. Habitat utilization of the dorcas gazelle in a desert saline area. Journal of Arid Environments 3:161–167.

Baharav D. 1982. Desert habitat partitioning by the dorcas gazelle. Journal of Arid Environments 5:323–335.

Baharav D. 1983. Reproductive strategies in female Mountain and Dorcas gazelles (Gazella
gazella gazella and Gazella dorcas). Journal of Zoology 200:445–453.

Barboza PS, Parker KL, Hume ID, editors. 2009. Integrative Wildlife Nutrition. Springer,
Berlin, Heidelberg.

Barker M, Rayens W. 2003. Partial least squares for discrimination. Journal of Chemometrics
17:166–173.

Belovsky GE. 1984. Herbivore Optimal Foraging: A Comparative Test of Three Models. The
American Naturalist 124:97–115. The University of Chicago Press.

Birnie-Gauvin K, Peiman KS, Raubenheimer D, Cooke SJ. 2017. Nutritional physiology and
ecology of wildlife in a changing world. Conservation Physiology 5:cox030.

Blanco G, Lemus JA, García-Montijano M. 2011. When conservation management becomes
contraindicated: impact of food supplementation on health of endangered wildlife.
Ecological Applications 21:2469–2477.

Blank DA. 1996. The Acacia gazelle: extinction of a subspecies. Gnusletter 15:7–9.

Blank DA. 2000. Acacia gazelle increases with habitat improvement. Gnusletter 19:11.

Blank DA. 2005. Diet and Feeding Behavior of acacia gazelle (Gazella g. acaciae) in Israel.
Selevinia 13:147–155.

Breslau B, Polak T, Shalmon B, Groner E. 2020. Evidence of browsing pressure on the
critically endangered Acacia gazelle (Gazella acaciae). Journal of Arid Environments
173:104019.

Carrete M, Donázar JA, Margalida A. 2006. Density-Dependent Productivity Depression in
Pyrenean Bearded Vultures: Implications for Conservation. Ecological Applications
16:1674–1682.

Chapman CA, Chapman LJ, Rode KD, Hauck EM, McDowell LR. 2003. Variation in the
Nutritional Value of Primate Foods: Among Trees, Time Periods, and Areas.
International Journal of Primatology 24:317–333.

Charnov EL. 1976. Optimal foraging, the marginal value theorem. Theoretical Population
Biology 9:129–136.

Codron D, Lee-Thorp JA, Sponheimer M, Codron J. 2007. Nutritional content of savanna
plant foods: implications for browser/grazer models of ungulate diversification.
European Journal of Wildlife Research 53:100–111.

Conover WJ, Iman RL. 1979. Multiple-comparisons procedures. Informal report. LA-7677-
MS. Los Alamos National Lab. (LANL), Los Alamos, NM (United States). Available
from https://www.osti.gov/biblio/6057803 (accessed May 29, 2024).

Conte G, Ciampolini R, Cassandro M, Lasagna E, Calamari L, Bernabucci U, Abeni F. 2018.
586   Feeding and nutrition management of heat-stressed dairy ruminants. Italian Journal of
587   Animal Science 17:604–620. Taylor & Francis.

588 Cooper SM, Owen-Smith N. 1986. Effects of plant spinescence on large mammalian
589   herbivores. Oecologia 68:446–455.

590 Cortés-Avizanda A, Carrete M, Serrano D, Donázar JA. 2009a. Carcasses increase the
591   probability of predation of ground-nesting birds: a caveat regarding the conservation
592   value of vulture restaurants. Animal Conservation 12:85–88.

593 Cortés-Avizanda A, Selva N, Carrete M, Donázar JA. 2009b. Effects of carrion resources on
594   herbivore spatial distribution are mediated by facultative scavengers. Basic and
595   Applied Ecology 10:265–272.

596 Costello CM, Cain SL, Pils S, Frattaroli L, Haroldson MA, van Manen FT. 2016. Diet and
597   Macronutrient Optimization in Wild Ursids: A Comparison of Grizzly Bears with
598   Sympatric and Allopatric Black Bears. PLOS ONE 11:e0153702. Public Library of
599   Science.

600 Cowsert RL, Montgomery MJ. 1969. Effect of Varying Forage-to-Concentrate Ratio of
601   Isonitrogenous Rations on Feed Intake by Ruminants. Journal of Dairy Science
602   52:64–67.

603 DeGabriel JL, Moore BD, Felton AM, Ganzhorn JU, Stolter C, Wallis IR, Johnson CN, Foley
604   WJ. 2014. Translating nutritional ecology from the laboratory to the field: milestones
605   in linking plant chemistry to population regulation in mammalian browsers. Oikos
606   123:298–308.

607 DePasquale AN, Hogan JD, Villalobos Suarez C, Mah MA, Martin J-C, Fedigan LM,
608   Rothman JM, Melin AD. 2023. Does colour vision type drive dietary and nutritional
609   niche differentiation in wild capuchins (Cebus imitator)? Animal Behaviour 205:89–
610   106.

611 Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. 1956. Colorimetric Method for
612   Determination of Sugars and Related Substances. Analytical Chemistry 28:350–356.

613 Felton AM, Felton A, Lindenmayer DB, Foley WJ. 2009a. Nutritional goals of wild primates.
614   Functional Ecology 23:70–78.

615 Felton AM, Felton A, Raubenheimer D, Simpson SJ, Foley WJ, Wood JT, Wallis IR,
616   Lindenmayer DB. 2009b. Protein content of diets dictates the daily energy intake of a
617   free-ranging primate. Behavioral Ecology 20:685–690.

Felton AM, Felton A, Raubenheimer D, Simpson SJ, Krizsan SJ, Hedwall P-O, Stolter C.
2016. The Nutritional Balancing Act of a Large Herbivore: An Experiment with
Captive Moose (Alces alces L). PLOS ONE 11:e0150870. Public Library of Science.

Felton AM, Felton A, Wood JT, Foley WJ, Raubenheimer D, Wallis IR, Lindenmayer DB.
2009c. Nutritional Ecology of Ateles chamek in lowland Bolivia: How Macronutrient
Balancing Influences Food Choices. International Journal of Primatology 5:675–696.

Felton AM, Wam HK, Felton A, Simpson SJ, Stolter C, Hedwall P-O, Malmsten J, Eriksson
T, Tigabo M, Raubenheimer D. 2021. Macronutrient balancing in free-ranging
populations of moose. Ecology and Evolution 11:11223–11240.

Felton AM, Wam HK, Stolter C, Mathisen KM, Wallgren M. 2018. The complexity of
interacting nutritional drivers behind food selection, a review of northern cervids.
Ecosphere 9:e02230.

Fornara DA, Du Toit JT. 2007. Browsing Lawns? Responses of Acacia Nigrescens to
Ungulate Browsing in an African Savanna. Ecology 88:200–209.

Freeland WJ, Janzen DH. 1974. Strategies in Herbivory by Mammals: The Role of Plant
Secondary Compounds. The American Naturalist 108:269–289.

Ganzhorn JU et al. 2017. The importance of protein in leaf selection of folivorous primates.
American Journal of Primatology 79:e22550.

Gowda JH. 1997. Physical and chemical response of juvenile Acacia tortilis trees to
browsing. Experimental evidence. Functional Ecology 11:106–111.

Hadas L, Hermon D, Boldo A, Arieli G, Gafny R, King R, Bar-Gal GK. 2015. Wild Gazelles
of the Southern Levant: Genetic Profiling Defines New Conservation Priorities. PLOS
ONE 10:e0116401. Public Library of Science.

Hagerman AE, Butler LG. 1994. [42] Assay of condensed tannins or flavonoid oligomers and
related flavonoids in plants. Pages 429–437 Methods in Enzymology. Academic
Press.

Halevy G, Orshan G. 1972. Ecological studies on Acacia species in the Negev and Sinai. I.
Distribution of Acacia raddiana, A. tortilis and A. gerrardii ssp. negevensis as related
to environmental factors. Israel Journal of Botany 21:197–208.

Halevy G, Orshan G. 1973. Ecological studies on Acacia species in the Negev and Sinai. II.
Phenology of Acacia raddiana, A.tortilis and A.gerrardii ssp. negevensis. Israel
Journal of Botany 22:120–138.

Hecker LJ, Edwards MA, Nielsen SE. 2021. Assessing the nutritional consequences of
switching foraging behavior in wood bison. Ecology and Evolution 11:16165–16176.

Hou R, Chapman CA, Rothman JM, Zhang H, Huang K, Guo S, Li B, Raubenheimer D.

2021. The geometry of resource constraint: An empirical study of the golden snub-

nosed monkey. Journal of Animal Ecology 90:751–765.

IUCN SSC Antelope Specialist Group. 2017. Gazella arabica. The IUCN Red List of

Threatened Species 2017: e.T117582065A88018124DOI: https://dx.doi.org/10.2305.

Jensen PG, Pekins PJ, Holter JB. 1999. Compensatory effect of the heat increment of feeding

on thermoregulation costs of white-tailed deer fawns in winter. Canadian Journal of

Zoology 77:1474–1485. Canadian Science Publishing NRC Research Press, Ottawa,

Canada.

John A, Webster F. 1978. Prediction of the Energy Requirements for Growth in Beef Cattle.

Page in Bourne GH, editor. Human and Veterinary Nutrition, Biochemical Aspects of

Nutrients. S.Karger AG.

Johnson CA, Raubenheimer D, Chapman CA, Tombak KJ, Reid AJ, Rothman JM. 2017.

Macronutrient balancing affects patch departure by guerezas (Colobus guereza).

American Journal of Primatology 79:e22495.

Jones DB. 1931. Factors for converting percentages of nitrogen in foods and feeds into

percentages of proteins. US Department of Agriculture.

Kay AD, Zumbusch T, Heinen JL, Marsh TC, Holway DA. 2010. Nutrition and interference

competition have interactive effects on the behavior and performance of Argentine

ants. Ecology 91:57–64.

Kohler SL, McPeek MA. 1989. Predation Risk and The Foraging Behavior of Competing

Stream Insects. Ecology 70:1811–1825. Ecological Society of America.

Landhäusser SM et al. 2018. Standardized protocols and procedures can precisely and

accurately quantify non-structural carbohydrates. Tree Physiology 38:1764–1778.

Lawler IR, Foley WJ, Eschler BM, Pass DM, Handasyde K. 1998. Intraspecific variation in

Eucalyptus secondary metabolites determines food intake by folivorous marsupials.

Oecologia 116:160–169.

López-Bao JV, Palomares F, Rodríguez A, Delibes M. 2010. Effects of food supplementation

on home-range size, reproductive success, productivity and recruitment in a small

population of Iberian lynx. Animal Conservation 13:35–42.

Martínez-Abraín A, Oro D. 2013. Preventing the development of dogmatic approaches in

conservation biology: A review. Biological Conservation 159:539–547.

Mattson WJ. 1980. Herbivory in Relation to Plant Nitrogen Content. Annual Review of

Ecology and Systematics 11:119–161.

Mendelssohn H. 1974. The development of the population of gazelles in Israel and their
behavioural adaptions. Pages 723–743 in Geist V, Walther F, editors. The Behaviour
of Ungulates and Its Relation to Management. IUCN Publications.

Mendelssohn H, Groves CP, Shalmon B. 1997. A new subspecies of Gazella Gazella from the
Southern Negev. Israel Journal of Zoology 43:209–215. Taylor & Francis Group.

Meteorological Database | Israel Meteorological Service. 2023. Available from
https://ims.gov.il/en/data_gov (accessed November 10, 2023).

Milton K. 1979. Factors Influencing Leaf Choice by Howler Monkeys: A Test of Some
Hypotheses of Food Selection by Generalist Herbivores. The American Naturalist
114:362–378. The University of Chicago Press.

Nie Y, Zhang Z, Raubenheimer D, Elser JJ, Wei W, Wei F. 2015. Obligate herbivory in an
ancestrally carnivorous lineage: the giant panda and bamboo from the perspective of
nutritional geometry. Functional Ecology 29:26–34.

Nzimande A, Mgqatsa N, Kraai UM, Tsvuura Z, Kraai M. 2022. Goat browsing does not
induce physical defences but reduces leaf size in Vachellia trees. Acta Oecologica
114:103818.

Pettersson LB, Brönmark C. 1993. Trading off Safety against Food: State Dependent Habitat
Choice and Foraging in Crucian Carp. Oecologia 95:353–357. Springer.

Polak T, Rieder G, Gisis E, Ochnuna M. 2023. Acacia gazelle (Gazella arabica) annual count
(2023), summary and evaluation of population trends since fence collapse in 2013.
Nature and Parks Authority, Southern Arava, Israel. (In Hebrew).

Porter LJ, Hrstich LN, Chan BG. 1985. The conversion of procyanidins and prodelphinidins
to cyanidin and delphinidin. Phytochemistry 25:223–230.

R Core Team. 2023. R: a language and environment for statistical computing. manual. R
Foundation for Statistical Computing, Vienna, Austria. Available from https://www.R-
project.org/.

Raubenheimer D. 2011. Toward a quantitative nutritional ecology: the right-angled mixture
triangle. Ecological Monographs 81:407–427. John Wiley & Sons, Ltd.

Raubenheimer D, Machovsky-Capuska G, Felton A, Simpson S. 2014. Nutritional Geometry:
From Insects to Ruminants. Animal production 30:32–36.

Raubenheimer D, Machovsky-Capuska GE, Chapman CA, Rothman JM. 2015. Geometry of
nutrition in field studies: an illustration using wild primates. Oecologia 177:223–234.

Raubenheimer D, Simpson SJ. 2018. Nutritional ecology and foraging theory. Current
Opinion in Insect Science 27:38–45.

Reynolds CK, Tyrrell HF, Reynolds PJ. 1991. Effects of Diet Forage-to-Concentrate Ratio
and Intake on Energy Metabolism in Growing Beef Heifers: Whole Body Energy and
Nitrogen Balance and Visceral Heat Production. The Journal of Nutrition 121:994–
1003.

Rohart F, Gautier B, Singh A, Cao K-AL. 2017. mixOmics: An R package for 'omics feature
selection and multiple data integration. PLOS Computational Biology 13:e1005752.
Public Library of Science.

Rohner C, Ward D. 1997. Chemical and mechanical defense against herbivory in two
sympatric species of desert Acacia. Journal of Vegetation Science 8:717–726.

Rothman JM, Chapman CA, Van Soest PJ. 2012. Methods in Primate Nutritional Ecology: A
User's Guide. International Journal of Primatology 33:542–566.

Rothman JM, Dusinberre K, Pell AN. 2009. Condensed tannins in the diets of primates: a
matter of methods? American Journal of Primatology 71:70–76.

Rothman JM, Raubenheimer D, Chapman CA. 2011. Nutritional geometry: gorillas prioritize
non-protein energy while consuming surplus protein. Biology Letters 7:847–849.
Royal Society.

Schoener TW. 1971. Theory of Feeding Strategies. Annual Review of Ecology and
Systematics 2:369–404. Annual Reviews.

Seagle SW, McNaughton SJ. 1992. Spatial variation in forage nutrient concentrations and the
distribution of Serengeti grazing ungulates. Landscape Ecology 7:229–241.

Shalmon B. 1991. Gazella gazella acaciae conservation action plan. Gnusletter 11:9–11.

Shalmon B, Sun P, Wronski T. 2020. Factors driving Arabian gazelles (Gazella arabica) in
Israel to extinction: time series analysis of population size and juvenile survival in an
unexploited population. Biodiversity and Conservation 29:315–332.

Shrestha TK, Hecker LJ, Aryal A, Coogan SCP. 2020. Feeding preferences and nutritional
niche of wild water buffalo (Bubalus arnee) in Koshi Tappu Wildlife Reserve, Nepal.
Ecology and Evolution 10:6897–6905.

Simpson SJ, Raubenheimer D. 2012. The Nature of Nutrition: A Unifying Framework from
Animal Adaptation to Human Obesity. Princeton University Press, Princeton.

Simpson SJ, Sibly RM, Lee KP, Behmer ST, Raubenheimer D. 2004. Optimal foraging when
regulating intake of multiple nutrients. Animal Behaviour 68:1299–1311.

Solon-Biet SM et al. 2014. The Ratio of Macronutrients, Not Caloric Intake, Dictates
Cardiometabolic Health, Aging, and Longevity in Ad Libitum-Fed Mice. Cell
Metabolism 19:418–430.

Spitzer R et al. 2023. Macro-nutritional balancing in a circumpolar boreal ruminant under
winter conditions. Functional Ecology 37:1256–1268.

Spitzer R, Ericson M, Felton AM, Heim M, Raubenheimer D, Solberg EJ, Wam HK,
Rolandsen CM. 2024. Camera collars reveal macronutrient balancing in free-ranging
male moose during summer. Ecology and Evolution 14:e70192.

Stolter C, Ball JP, Julkunen-Tiitto R, Lieberei R, Ganzhorn JU. 2005. Winter browsing of
moose on two different willow species: food selection in relation to plant chemistry
and plant response. Canadian Journal of Zoology 83:807–819. NRC Research Press.

Takahashi MQ, Rothman JM, Raubenheimer D, Cords M. 2019. Dietary generalists and
nutritional specialists: Feeding strategies of adult female blue monkeys
(Cercopithecus mitis) in the Kakamega Forest, Kenya. American Journal of
Primatology 81:e23016.

Takahashi MQ, Rothman JM, Raubenheimer D, Cords M. 2021. Daily protein prioritization
and long-term nutrient balancing in a dietary generalist, the blue monkey. Behavioral
Ecology 32:223–235.

Van Soest PJ. 1994. Nutritional Ecology of the Ruminant, 2nd edition. Cornell University
Press.

Van Soest PJ, Robertson JB, Lewis BA. 1991. Methods for Dietary Fiber, Neutral Detergent
Fiber, and Nonstarch Polysaccharides in Relation to Animal Nutrition. Journal of
Dairy Science 74:3583–3597.

Verheyden-Tixier H, Renaud P-C, Morellet N, Jamot J, Besle J-M, Dumont B. 2008.
Selection for nutrients by red deer hinds feeding on a mixed forest edge. Oecologia
156:715–726.

Weber A. 2001. Interactions Between Predator Kairomone and Food Level Complicate the
Ecological Interpretation of Daphnia Laboratory Results. Journal of Plankton
Research 23:41–46.

West JW. 1999. Nutritional strategies for managing the heat-stressed dairy cow. Journal of
Animal Science 77:21–35.

Westerhuis JA, van Velzen EJJ, Hoefsloot HCJ, Smilde AK. 2008. Discriminant Q2 (DQ2)
for improved discrimination in PLSDA models. Metabolomics 4:293–296.

White TCR. 1993. The inadequate environment: Nitrogen and the abundance of animals, 1st
edition. Springer Berlin, Heidelberg.

Yom-Tov Y. 2016. Gazelles in Israel. Dan Pery Press, Jerusalem. (In Hebrew).

Zaguri M, Mogilevsky I, Raubenheimer D, Hawlena D. 2024. 'Dust you shall eat': The
788   complex nutritional and functional considerations underlying a simple diet. Ecology
789   Letters 27:e14414.