Large effect life-history genomic regions are associated with functional morphological

traits in Atlantic salmon

Tutku Aykanat[1], Paul V. Debes[2,6], Shadi Jansouz[1], Lison Gueguen[3], Jaakko Erkinaro[4], Victoria L. Pritchard[5], Craig

R. Primmer[1,6], Geir H. Bolstad[7]

1. Organismal and Evolutionary Biology Research Program, Faculty of Biological and Environmental Sciences, PO Box 56, FI-
00014 University of Helsinki, Finland. tutku.aykanat@helsinki.fi.

2. Department of Aquaculture and Fish Biology, Hólar University, 550 Sauðárkrókur, Iceland.

3. Oniris Graduate School of Veterinary Medicine, Nantes, France.

4. Natural Resources Institute Finland (Luke), Oulu, Finland

5. Institute for Biodiversity & Freshwater Conservation, UHI Inverness, Inverness, United Kingdom.

6. Institute of Biotechnology, Helsinki Institute of Life Science (HiLIFE), University of Helsinki, Helsinki, Finland.

7. Norwegian Institute for Nature Research (NINA), NO-7485 Trondheim, Norway.

Abstract

Understanding pleiotropic architectures of phenotypes is instrumental for identifying the functional basis of

adaptive genetic variation in the wild. Life-history variation may have a morphological basis that mediates

resource acquisition and allocation pathways, but identifying the underlying genetic basis of such traits is

challenging. In this study, using Atlantic salmon juveniles reared in common garden conditions, we test if two

life-history associated loci in Atlantic salmon, *six6* and *vgll3,* are also associated with functional morphological

traits. These loci had previously shown to exhibit strong signals of adaptation and are highly correlated with

age at maturity. Here, we show that genetic variation at the *vgll3* locus is linked to variation in morphological

traits that underlie swimming performance, along a trade-off axis between efficient cruising and maneuvering,

while the genetic variation at the *six6* locus was linked to variation in body-head proportions suggesting the

potential functional importance of these traits for resource acquisition efficiency. However, the direction of

changes in morphological traits associated with late- vs. early- maturing alleles was not always consistent with

the expected direction of an effect to maturation timing. Our results reveal a complex morphological landscape

associated with the genetic variation in these loci, possibly as a result of pleiotropy or linkage across these

genomic regions.

Keywords: Atlantic salmon, morphology, morphometrics, adaptive genetic variation, age at maturity,

pleiotropy, *six6*, *vgll3*

Introduction

Elucidating the genotype-phenotype relationship is instrumental to identifying the functional basis of adaptation, and the potential targets of selection (Barrett and Hoekstra, 2011; Kawecki and Ebert, 2004; Nielsen *et al*, 2009; Wagner and Altenberg, 1996). Methodological and statistical advances in genetics now allow the identification of genomic regions responding to local selection, but the phenotypic link between genotype and fitness is often missing. Furthermore, many genomic regions can affect a multitude of phenotypic traits (via pleiotropy or linkage disequilibrium), which makes identifying the phenotypic traits under selection even more challenging (Paaby and Rockman, 2013; Solovieff *et al*, 2013). Even large-genome-wide association studies might be underpowered to detect pleiotropy (Wagner and Zhang, 2011), a methodological bias by which the adaptive significance of pleiotropy can be overlooked especially in natural, non-model populations even if it may have substantial effects to limit, or promote adaptation (Orr, 2000; Otto, 2004). In natural systems, pleiotropy has been mostly studied at the phenotypic level to understand the evolutionary trade-offs by quantifying the physiological antagonisms between the senescence and reproduction age. Yet, some studies identify pleiotropic effects of genes associated with adaptive genetic variation. For example, adaptive divergence between lake vs. marine morphs of threespine stickleback (*Gasterosteus aculeatus*) was explained by both pleiotropy in the *Eda* gene, and via strong linkage with the surrounding loci (Archambeault *et al*, 2020), illustrating the significance of such genetic architectures for adaptation.

Genomic regions with large adaptive importance are more likely to be detected in genome scans for selection and are likely to have broad cellular and phenotypic effects (Chan *et al*, 2010; Colosimo *et al*, 2005). Major effect loci may have roles at critical nodes of the regulatory networks, which have high centrality and connectivity, and could thereby drive strong genetic interdependences among traits (Chan *et al*, 2010; Colosimo *et al*, 2005; Jimenez-Gomez *et al*, 2010; Weith *et al*, 2023) but see (Boyle *et al*, 2017; Jakobson and Jarosz, 2020). Therefore, loci exhibiting strong signals of adaptation may provide feasible empirical approaches to understand the significance of complex evolutionary processes, such as how pleiotropy and genetic interactions may hinder or promote adaptation in relation to the direction of natural selection between co-varying traits. Yet, such examples remain rare in non-model, wild organisms (Archambeault *et al*, 2020; Endler, 2000; Mackay *et al*, 2009; Olson-Manning *et al*, 2012).

Functional variation in morphological traits is integral in shaping resource use, and is correspondingly associated with life-history variation and ecotypic divergence in a large array of fish species, including salmonid fishes (Ferry-Graham *et al*, 2002; Fitzgerald *et al*, 2017; Keeley *et al*, 2005; Lundsgaard-Hansen *et al*, 2013; Mindel *et al*, 2016; Proulx and Magnan, 2004; Svanbäck and Eklöv, 2003; Taylor and McPhail, 1985; Varian and

Nichols, 2010; Webb, 1988). For example, a fusiform body and shorter fins appear adaptive to a long-distance migratory strategy, likely because this body shape reduces the drag (hence the energy consumed) during migration (Morinville and Rasmussen, 2007). Likewise, a robust body shape with larger head, caudal peduncle, and paired fins has been associated with piscivory in rainbow trout (*Oncorhynchus mykiss*, Keeley *et al*, 2005). Recent population genomics studies in Atlantic salmon have consistently identified two unlinked genomic regions, surrounding the genes *six6* in chromosome 9 and *vgll3* in chromosome 25 as strong targets for local selection within all European Atlantic salmon lineages (Barson *et al*, 2015; Miettinen *et al*, 2023; Pritchard *et al*, 2018; Zueva *et al*, 2020). These two loci have also been linked to a suite of life-history traits that strongly influence fitness. In particular, the *vgll3* and *six6* loci are strongly associated with age-at-first-maturity in seagoing salmon at the individual- and population- levels respectively (Barson *et al*, 2015) (see also, Ayllon *et al*, 2015; Sinclair-Waters *et al*, 2020), and the *vgll3* locus affects other alternative life-history strategies, such as precocious male maturation (Debes *et al*, 2021), and iteroparity (Aykanat *et al*, 2019). Similarly, these loci explain variation across a number of metabolic and behavior traits: *vgll3* has been shown to be associated with aggression, maximum metabolic rate, and aerobic scope at the juvenile stage (Bangura *et al*, 2022; Prokkola *et al*, 2022), while *six6* predicts diet acquisition at sea in maturing adults migrating towards fresh waters (Aykanat *et al*, 2020), and exhibits epistasis with *vgll*3 to effect aerobic scope variation (Prokkola *et al*, 2022). The strong genetic interdependence of these phenotypic traits across two loci points to the potential for a shared developmental origin (of phenotypes to the respective loci), which is supported by the role of *vgll3* and *six6* in the hormonal and cell-fate commitment gene-expression pathways (Ahi *et al*, 2022; Kurko *et al*, 2020; Verta *et al*, 2020; Verta *et al*, 2024) and broad expression of *six6* during early development (Moustakas-Verho *et al*, 2020). Finally, a recent study showed both loci are linked to early life survival in juvenile Atlantic salmon, albeit via parental genetic effects, further supporting broad functional roles across life stages (Aykanat *et al*, 2024). This collective evidence indicates that these loci are broad-scale modulators of salmon phenotype, yet their effect on morphological variation, if any, is not known.

In order to characterize the morphological variation associated with these locally adaptive genomic regions, in this study, we quantified morphometric variation in more than 700 juvenile Atlantic salmon reared in common garden conditions and related it to *six6* and *vgll3* genotype using a mixed modelling approach that also controls for confounding effects of length and condition factor. We identified a suite of morphological characters correlated to the variation in the *vgll3* and *six6* loci. Our study extends the broad pleiotropic effect of *vgll3* and *six6* loci to functional morphological traits, including phenotypes that are likely important targets of natural selection.

Materials and Methods

Fish breeding and rearing and common garden conditions

Details of breeding and rearing of individuals from the same cohort were provided in Debes *et al* (2021), using a similar experimental procedure and facilities, except that fish in the present study were reared in approximately 2° C cooler water temperature (See, Debes *et al* 2019, for the details of the rearing temperature). Briefly, the parents of the individuals used were 44 Baltic lineage Atlantic salmon from a broodstock maintained by the Natural Resources Institute Finland (Laukaa, Finland) that were initially originated from the Neva River in Russia (Gulf of Finland in the Eastern Baltic Sea) and has been used for juvenile salmon supplementation in the Kymi river in South-Eastern Finland (Asheim *et al*, 2023). Parents were crossed to have balanced representation of homozygote genotypes in the *vgll3* locus between early (E) and late (L) maturing alleles (i.e. *vgll3*^{EE} and *vgll3*^{LL}, Supplementary Table 1). *Six6* genotype distribution was not considered during the breeding design, but was nevertheless segregating in a number of families (Supplementary Table 1). Crosses were performed in November 2017 and offspring reared in the University of Helsinki fish facilities, first in replicated family-specific compartments in vertical incubators until the swim-up stage, and then subsequently offspring of all families were mixed in eight tanks with roughly balanced proportions of families per tank (Supplementary Table 2). The rearing temperature of fish in this study followed a seasonal cycle (range: 4.1– 16.0°C) and the natural light regime of the latitude (61.054° N) as described in Debes *et al* (2019). In August 2018, the fish were individually PIT-tagged and a fin clip sample was taken to identify the families via genetic pedigree reconstruction. Fish were fed ad libitum except for a short period of restricted feeding for half of the tanks, which included ad libitum feeding for two days and then no feeding for five days, each week for a five week period (Debes *et al*, 2021).

Landmarking and trait measurements

Morphological data were collected as part of a larger effort in December 2018. Fish of both sexes were euthanized using an overdose of tricaine methanesulphonate bath, and stored for a short time on ice until imaging was performed the same day. Each fish was sliced ventrally to collect data on gonad development for another study. The fish was placed on graphical paper and the body image was taken using a digital camera (Canon PowerShot S110) mounted on a tripod (Hama Star 61), with aperture-priority exposure at F:8, and auto focus option (Supplementary Figure 1). Cross sections were obtained by slicing the fish at the posterior end of the dorsal fin (hereafter termed as large cross section) and adipose fins (hereafter termed as small cross section), respectively. These were placed on a transparent sheet and digitized using a scanner (Epson XP-305) with a modified lid to accommodate the depth of the cross sections (Supplementary Figure 1).

Landmarks were placed on the whole body and the two cross-sections at a randomized order using imglab

software (Porto *et al*, 2020), by two of the co-authors (LG for landmarked body planes, and SC for landmarked

cross sections) who were blind to any information related to the genotypes or sex of the fish (Supplementary

Figure 2). Landmarks were not placed around the gut region due to the previous ventral slicing. Some

landmarks were solely used in the Procrustes analysis to quantify shape, while others used to quantify specific

trait values by calculating the distance between landmarks (Supplementary Figure 2, Table 1). Traits that were

measured twice on both the left and right side (Table 1), were averaged prior to analysis. All measurements

were reported in centimeters (cm).

Data Filtering

At the time of sampling, fish exhibited a bimodal length distribution which is often observed within juvenile

salmon from the same cohort (Hoar, 1988). At the time of sampling a majority of fish used in the analysis (i.e.

upper mode of size distribution, see below) had a complete smolt appearance (base on the disappearance of

parr marks in the skin and silvery coloration, e.g., Debes *et al*, 2020), or were at late stages of smoltification

(faint parr marks with close to silvery skin coloration). We excluded a small number of fish in the lower mode to

avoid potential interactions associated with this size bimodality (fish fork length < 9 cm, N=69, Supplementary

Figure 3). In addition, fish with missing *vgll3* and *six6* genotypes, and sex (N=2, N=12, and N=4, respectively),

and a small number of precociously matured males (N=19) were excluded from the analysis. Missing landmarks

(0.02 % of the total landmarks) were imputed using the *estimate.missing* function in the *geomorph* package

(version 4.0.6, Baken *et al*, 2021). A landmarked image was excluded from the analysis if more than one

landmark was identified as an outlier in the Procrustes analysis (i.e. > 4 sd away from the mean). These

landmarks were not corrected manually, to avoid introducing among technician variation. After data filtering,

821 individuals remained in the dataset, of which 731, 737, and 772 individuals were in the small cross section,

large cross section, and whole body datasets, respectively, (Supplementary Table 1). 16, 18, and 712 individuals

were landmarked more than once to estimate repeatability. Landmarkinng for the first 30 and 50 images per

researcher for body and cross sections, respectively, were discarded as training sets.

Trait measurements

We quantified variation in 21 traits across various body planes with potential functional significance using the

landmark positions (Supplementary Figure2) and as formulated in Table 1.

Rayed fins and caudal peduncle length: We measured the length of dorsal, anal, and tail fins, between interior

and exterior insertion points, as well as the caudal peduncle (Table 1, Supplementary Figure 2). The variation in

these traits may be associated with habitat use, feeding, and swimming performance in salmonids, and other

fishes (e.g., Jonsson and Jonsson, 2011a; Keeley *et al*, 2005). Increased dorsal and pectoral fin sizes in fish are

likely important for increase maneuverability and stability, and likely important adaptations for faster flowing

streams (Webb, 1984). Specifically, large paired fins (such as pectoral and pelvic fins) in Atlantic salmon are

associated with populations that occupy faster flowing and colder streams than slower flowing and warmer

rivers, an adaptation attributed to providing effective stability under fast flowing, and turbulent currents

(Riddell and Leggett, 1981; Webb, 1988). In Coho salmon, juveniles of interior populations with longer

migration distances appear to have smaller fins (Taylor and McPhail, 1985), which may help to reduce drag in

long migrations (Varian and Nichols, 2010).

Eye diameter and position, and snout length: The size of the eye was approximated from the diameter, and the

snout length was measured as the distance between the center of the eye to the anterior extreme of the head

(Table 1, Supplementary Figure 2). These traits are morphological features associated with trophic

specialization and habitat use across fish taxa (Elmer *et al*, 2010; Peris Tamayo *et al*, 2020; Snorrason, 1994).

Aerobic muscle: We approximate the area of aerobic muscle fibers (also called slow-twitch muscle, or red

muscle) by measuring the landmark distances in the cross sections from the lateral line towards the transverse

sections, and by measuring the extent of the red muscle along the lateral line (Table 1, Supplementary Figure

2). Aerobic muscle provides energy required during steady, cruise swimming, and helps with thermal

acclimation in fishes (Altringham and Ellerby, 1999; Shuman and Coughlin, 2018). Recently, it has been shown

that both *vgll3* and *six6* loci are associated with whole body aerobic activity (Prokkola *et al*, 2022), which is a

critical physiological process in salmonid life cycle (Eliason and Farrell, 2015), which may indicate that aerobic

muscle mass, as a proxy for aerobic activity, may be associated with these loci.

Cross sectional centroid size and diameter, and vertebral thickness: We measured cross sectional centroid size

and horizontal diameter from both large and small cross-section images (Table 1, Supplementary Figure 2), as

approximation for fusiform body shape. A slimmer and more torpedo-shape provides improved energy

efficiency during cruising and in high flow streams via reducing the water drag, and it is an important life

history adaptation among fishes (Crossin *et al*, 2004; Jonsson and Jonsson, 2011a; Taylor and McPhail, 1985;

Webb, 1988). In contrast, a robust cross-section facilitates burst swimming, which is advantages for prey

avoidance, and food acquisition (Taylor and McPhail, 1985).

We also quantified vertebral width as a proxy for cone diameter, to approximate vertebra's flexibility/stiffness

during the lateral bending movement in the undulated motion, which help maintaining steady swimming at the

expense of maneuvering capabilities (Baxter *et al*, 2022; Donatelli *et al*, 2021; Long and Nipper, 1996).

Head and body size: We quantified head length and width (Table 1, Supplementary Figure 2), which are

important determinants of ecotypic divergence via resource and habitat use in fishes (Peris Tamayo *et al*, 2020;

Simonsen *et al*, 2017). A larger head and larger snout length may be associated with feeding regime towards

piscivory (Keeley *et al*, 2005), or generally speaking, consuming larger prey items. We also measured head and

body centroid sizes as an approximation for the area of the morphometric planes, which is the square root of

the sum of squared distances of all the landmarks used in the Procrustes analysis from the point of center of

gravity (Klingenberg, 2016). The Atlantic salmon has a smaller head-body ratio, compared to the brown trout

(Jonsson and Jonsson 2011), which likely makes salmon more efficient swimmers both in strong freshwater

currents and during longer marine migrations.

Statistical analysis

Our analysis framework comprised of a series of analyses whereby Procrustes shape variable, and

morphological traits were used as response variables. These analyses were conducted independently for body,

small cross section, and large cross section. In all analyses length (centered to zero and scaled to have a unit

standard deviation), body condition (Fulton's condition factor, scaled), sex, and food treatment were used as

covariates. In all models, we implemented genetic variation as additive, where *vgll3* and *six3* genetic effects are

coded as 0, 1, and 2 for EE, EL, and LL genotypes respectively (E and L refer to the alleles associated with early

and late maturation, respectively), so that a unit change in genetic effect represent an additive allelic

substitution effect from E to L. Unless otherwise noted, all statistical analysis were performed in R software

version 4.4.1 (R Core Team, 2019).

We performed multivariate generalized Procrustes analysis to quantify shape variation (Gower, 1975), using

the default settings in the *gpagen* function in the geomorph package (version 4.0.8, Baken *et al*, 2021). This

function outputs Procrustes shape variables, centered on the mean value for each landmark and normalized for

centroid size. Centroid size is the square root of the sum of squared distances of landmarks from the center of

gravity (i.e., centroid) and it is highly correlated to size of the fish.

Procrustes shape data was then used to perform a Procrustes linear regression analysis using the *procD.lm*

function with default settings (i.e. residual randomization was used for significance testing and type III marginal

sums of squares for cross products computations). The model parameters were as follows:

$Y = \mu + L + CF + Feed + Tank + Sex + Vgll3 + Six6 + Fam + \varepsilon_{error}$

Where Y is the multivariate data frame consisting of x and y coordinates for Procrustes shape coordinates of

landmarks, μ is the intercept, L is the scaled fork length, CF is the scaled Fulton's condition factor, and Feed is

the factor associated with feed treatment (ad-libitum vs restricted feed). Other terms are the rearing tank ID

that the fish were reared after the swim-up stage (Tank), sex (Sex), family (Fam), and the additive genetic

effects for *vgll3* and *six6*, respectively. Since, the *procD.lm* function does not accommodate random terms, we

included tank and family effects as fixed terms. Trait values were averaged prior to the analyses when an

individual had replicate measurements. We reported multivariate analysis of variance Pillai's trace statistics,

estimated using the *manova.update* function in the RRPP package (version 2.0.3, Collyer *et al*, 2018), and plot

individuals landmark differences between loci using the *plotRefToTarget* in the geomorph package.

To decompose the variation associated with trait values and estimate parameter coefficients, we employed a

linear mixed effect (animal) model using the *relmatLmer* function in lme4qtl (version 0.2.2, Ziyatdinov *et al*,

2018), which is an implementation of *lmer* function in lme4 package (Bates, 2010), and can fit the animal model

(i.e., accommodate the additive relatedness matrix). When estimating fixed coefficients, the model was fitted

with maximum likelihood using the *REML = FALSE* option in the *relmatLmer* function. The model structure was

as follows:

$Y = \mu + L + CF + Feed + Sex + Vgll3 + Six6 + \epsilon_{ID} + \epsilon_{tank} + \epsilon_{animal} + \epsilon_{error}$

Fixed effect terms are as described above except that the fixed tank and family terms in the previous model

were replaced with random tank and animal terms, respectively. Random terms thus included, ϵ_{ID} which are

the individual effects, ϵ_{tank} are the effects associated with the tank effects, and ϵ_{error} is the measurement error.

ϵ_{animal} is the random term associated with the background genetic effects after controlling for *vgll3* and *six6*

fixed effects. The relatedness matrix was obtained using the pedigree data in Debes *et al* (2019) using *getA*

function in the pedigreem package (version 0.3.4, Vazquez *et al*, 2010). Since we used repeated measurements

in the analysis, ϵ_{ID} reflects the error term, while ϵ_{error} are residuals that quantifies deviations between repeated

samples (i.e. measurement error).

The significance of the fixed effects terms was estimated using the *as_lmerModLmerTest* function in the

lmerTest package (version 1.1-3, Kuznetsova *et al*, 2017) in R using type 3 F test with Satterthwaite's method to

estimate the effective degrees of freedom.

We then estimated the additive genetic variation, the contribution of *vgll3* and *six6* loci to the additive genetic

variation, evolvability, and repeatability. For these, we employed three models similar to above, but the first

model (Model$_{VA.full}$) did not contain *vgll3* or *six6* to estimate total additive genetic variation, and was fitted

using the *REML = TRUE* option in the *relmatLmer* function to obtain unbiased estimates of variance

components. As such, variance of ϵ_{animal} (V_{animal}) in the Model$_{VA.full}$ is the total additive genetic variation (V_A).

Percent contribution of *vgll3* and *six6* are calculated as $\%V_{A.vgll3} = 100 * ((2 \times L_{vgll3} \times E_{vgll3} \times (\beta_{vgll3}^2 - se(\beta)_{vgll3}^2) /$

V_{animal}), and %$V_{\text{A.six6}}$ = 100 * ((2 x L$_{six6}$ x E$_{six6}$ x (β_{six6}^2 – se(β)$_{six6}^2$))) / V_{animal}), as in (Luo *et al*, 2003), where L$_{vgll3}$,

E$_{vgll3}$, L$_{six6}$ and E$_{six6}$ denotes frequency of L and E alleles in *vgll3* and *six6* loci, respectively, β_{vgll3} and β_{six6} denotes

six6 and *vgll3* E to L substitution effects, respectively, and se denotes standard error of the estimate. Percent

repeatability for each model is calculated as R = V_{ID} / (V_{ID} + V_{error}) * 100, and heritability values were calculated

as h^2 = V_{animal} / (V_{animal} + V_{tank} + V_{ID}), where V_{tank}, V_{ID}, and V_{error} are the model variance estimates associated with

tank ($\varepsilon_{\text{tank}}$), individual ($\varepsilon_{\text{ID}}$), and error ($\varepsilon_{\text{error}}$) terms in Model$_{VA.full}$. Similarly, evolvability (I_{A}) was calculated as V_{A} /

$\bar{y}^2$ using Model$_{VA.full}$, where $\bar{y}$ is the mean trait value (Hansen *et al*, 2011; Houle, 1992). Evolvabilities are

interpreted as expected proportional responses in the trait mean under unit selection (i.e. the selection on

fitness itself) (Hansen *et al*, 2003; Hansen *et al*, 2011). For all variance components, 95% confidence intervals

were estimated by obtaining the posterior distributions of random effect terms in the model (Model$_{VA.full}$),

using the *sims* function in the arm package(version 1.1.4-4, Gelman and Su, 2020)) in R, using 10,000

permutations. The simulations generate 10000 estimates for random effect terms, which then used to

calculate the variance components as above, sorted, with 250[th] and 9750[th] estimates are recorded as 95% CIs.

We did not control for the effect of dominance in the analyses, which may have inflated the estimates of

additive genetic variance (Lynch and Walsh, 1998).

Results:

Shape analysis:

The multivariate analyses using Procrustes shape variation as the response showed that genetic variation in

six6, but not in *vgll3* is associated with shape change in the body plane (Table 2). The change in the body plane

appears to be related to changes in the head region associated with *six6*, whereby a substitution from *six6*[E] to

six6[L] elongates the head and slightly bends up the lateral body positioning at ends of the lateral line, while

vgll3[E] to *vgll3*[L] substitution reduces the head area relatively to the body area, and pushes the anterior anal fin

landmark toward the posterior end of the body plane (Figure 1a). On the other hand, the genetic variation in

vgll3 and *six6* are not associated with overall shape changes in small and large cross sections. Although not

statistically significant, a *vgll3*[E] to *vgll3*[L] substitution was linked with a narrower horizontal width, a smaller

aerobic muscle area in the small cross section (Figure 1b), and a narrower aerobic muscle extension

perpendicular to lateral line in the large cross section (Figure 1c). These results seem to be concordant with the

trait-based analysis (see below). Overall, family, tank, feed restriction, and length were important determinants

for shape variation in all planes, and condition factor was also significant in the analysis of the body plane

(Table 2).

Trait analysis: After correcting for body length, condition factor, feed restriction, and sex, out of 21

morphological traits, nine and four were statistically significantly associated with the *vgll3* and *six6* loci,

respectively (Table 1, Figure 2. See also Supplementary Table 3 for *t*-values and effective degrees of freedoms

of terms). In general, the $vgll3^E$ to $vgll3^L$ substitution effect decreased trait values (Table 1, Figure 2) with

changes in trait values between $vgll3^{EE}$ and $vgll3^{LL}$ genotypes being > 1 %, and up to 3 % of the mean trait

values (Figure 2). Notably, aerobic muscle depth in the small cross section was reduced by 3.45 % ($p < 0.001$)

and aerobic muscle vertical extension in the large cross section was reduced by 2.72 % ($p = 0.006$) by the $vgll3^E$

to $vgll3^L$ allelic substitution (Table 1, Figure 2). The *vgll3* allelic substitution effect associated with lateral red

muscle length was statistically significant when three aerobic muscles trait values were combined into one

composite value ($p < 0.001$, Supplementary Table 4). The $vgll3^L$ allele also decreased vertebral thickness in both

small (1.37 % decrease, $p = 0.009$) and large (1.21 % decrease, $p = 0.032$) cross sections (Table 1, Figure 2). The

$vgll3^E$ to $vgll3^L$ substitution effect significantly reduced the small cross section diameter by 1.43 % ($p = 0.012$),

whereas the small section centroid size was reduced by 0.86 % ($p = 0.012$, Table 1, Figure 2). Reduced eye

diameter, snout length, and the anal fin length was also associated with the $vgll3^L$ allele (Table 1, Figure 2). The

trait values for all ray fins (dorsal, anal, pectoral, and tail fins) was trending toward a smaller size in relation to

the late maturing genotype (Table 1, Figure 2), and their cumulative effect, i.e., effect size of all trait values

combined, was statistically significant ($p = 0.022$, Supplementary Table 4). We observed that in some traits the

variation explained by $vgll3^E$ to $vgll3^L$ substitution in *six6* and *vgll3* loci was up to 50% of the variation explained

by length (a factor with strong, positive allometric effect on other trait values, see Table 1), suggesting that this

variation is biologically important (Supplementary Figure 4).

Most differences explained by variation in the *six6* genomic region were related to changes in head and body

size. As such, the $six6^E$ to $six6^L$ substitution was associated with a 0.17 % decrease on body centroid ($t = -3.70$,

$p < 0.001$) and a 0.47 % increase on head centroid size ($t = 2.68$, $p = 0.008$, Table 1, Figure 2). Finally, the snout

length, measured as the distance between the center of the eye and maximum anterior point in the body, was

elongated by 0.73 % in fish with an additional $six6^L$ allele ($t = 3.05$, $p = 0.002$, Table 1, Figure 2).

Sex effects typically ranged from smaller and up to the same order of magnitude as the allele substitution

effects in the life-history loci (Table 1). Notably, the snout, pectoral fin, and anal fin were significantly larger in

males compared to the females ($p = 0.002$, $p = 0.007$, and $p = 0.021$, respectively. Table 1). In addition, the head

centroid was larger in males compared to females ($p = 0.040$, Table 1). As expected length was a strong

predictor of all traits and condition factor was also positively associated with most of the traits (Table 1). Feed restriction was notably associated with decrease in fin trait values, such as dorsal fin ($p < 0.001$), anal fin ($p = 0.040$), and tail fin ($p = 0.003$, Table 1). Interestingly, feed restriction substantially reduced the centroid size ($p = 0.011$), cross section diameter ($p = 0.003$), and the size of the aerobic muscle ($p = 0.005$) in the small cross section, but not in the large cross section (Table 1).

Trait heritabilities were low to moderate in most cases, with a mean heritability of 0.19 across all 21 traits (Table 3). Length of tail fin and body centroid size exhibited the highest heritability with 0.50 (95 % CI 0.47-0.54) and 0.47 (95 % CI 0.43-0.50), followed by pectoral fin with 0.41 (95 % CI 0.36-0.45). The repeatability of trait measurements was moderate (mean repeatability 73.0 %, Table 1).

Evolvabilities (mean-scaled additive genetic variances, denoted as I_A) differed greatly among traits (Table 3). Interestingly, body centroid size had the lowest evolvability (<0.001), meaning that under selection as strong as fitness itself (unit selection) the expected increase in this trait would be less than 0.1%. On the other hand, large cross section aerobic muscle had an evolvability as high as 0.72, meaning that the expected increase in this trait under strong selection could be as high as 71% of the mean trait value. In general, out of 21 traits measured, aerobic muscle and vertebrae thickness traits had higher I_A, while centroid sizes and cross-section diameters had lower I_A (Table 3).

Additive genetic variation explained by *vgll3* locus was non-zero for 12 traits (Table 3). Consistent with the fixed effect parameterization of the genotype, a substantial portion of the additive genetic variation in vertebra thickness was explained by *vgll3*, with 30.7 % and 13.0 % for small and large cross sections, respectively (Table 3). A significant portion of the additive genetic variation in small section aerobic muscle, cross section diameter, and body centroid was also explained by the *vgll3* locus, with 22.4 %, 12.7, and 6.3 % for each trait, respectively (Table 3). The contribution of *six6* locus to additive genetic variation associated included snout length (9.4 %), and head and body centroid size (5.2 % and 4.2 %, respectively, Table 3) consistent with the locus effect size.

As length is a strong a predictor of both trait values, and overall body shape, a spurious association between traits and the life-history loci may be expected if these loci affect trait values via length as a mediator. To test this, we constructed a model to include interaction between length and life-history loci. Across 21 traits, we did not observe a meaningful difference in main genetic effects, nor were interaction terms *t*-values substantial (Supplementary Figure 5), suggesting that length-by-genotype interaction did not drive the estimated relations.

Discussion

Intraspecific differences in functionally important morphological traits may provide important insights into the mechanistic basis of adaptation. We detected several morphological traits in juvenile Atlantic salmon that were linked with variation in genomic regions that were previously reported to be associated with life-history diversity and known to be under local selection. The genetic polymorphism in the *vgll3* locus appears to shape a suite of traits that are associated with swimming performance, while the genetic polymorphism in the *six6* locus shapes head and body size, and head features such as snout length, which potentially affects resource acquisition efficiencies for different foraging strategies. On the other hand, these genomic regions have only a small effect on the general shape of the morphological planes. The only statistically significant effect was detected in the body plane plan that was associated with the *six6* loci (Figure 1), for which the change in shape appears to be concordant with change in head/body proportions observed in the univariate trait analysis. Our results support these genomic regions having broad phenotypic effects, which may have emerged from either a form of pleiotropy or due to linkage disequilibrium. Our design does not allow to categorize the pleiotropic effects further, whether it is based on an effect of single causal variant (biological pleiotropy), or based on effects associated with different phenotypes along a biochemical pathway (mediated pleiotropy), or as a result of linkage disequilibrium between two independent causative loci (spurious pleiotropy, e.g., Solovieff *et al*, 2013). Yet, the strong genetic interdependence among traits associated with *six6* and *vgll3* may have a developmental origin, which is supported by the patterns of expression of genes in these genomic regions during development, and their molecular regulatory functions, (Ahi *et al*, 2022; Kurko *et al*, 2020; Mobley *et al*, 2021; Moustakas-Verho *et al*, 2020; Verta *et al*, 2020; Verta *et al*, 2024).

The suite of morphological traits associated with *vgll3* variation reflects changes along a functional axis that facilitates efficient cruising and steady swimming (*vgll3*[E]) versus maneuvering and acceleration (*vgll3*[L]). Efficient cruising is an important trait especially in the marine environment, while better maneuvering and acceleration help to better cope with biotic (i.e. competition and predation risk) and abiotic (flow regime) factors in the freshwater habitat. The larger aerobic muscle area associated with the *vgll3*[E] allele could indicate a higher potential for aerobic muscle activity. This result is in concordance with (Prokkola *et al*, 2022) who demonstrated that the juvenile Atlantic salmon with early maturing genotype has a higher maximum metabolic rate (MMR) and aerobic scope (AS). Aerobic locomotion is expected to be central for Atlantic salmon during the marine phase, where salmon sub-adults migrate and continuously forage for food. Our results suggests salmon with the early genotype are more likely to maintain sustained swimming and foraging activities, beneficial during marine exploration. On the other hand, it is not clear if the genetic association between life-history loci

and morphological traits persists between life stages. The parr-smolt transformation (smoltification) includes

morphological changes that would prepare juvenile Atlantic salmon for the marine environment (Jonsson and

Jonsson, 2011b), hence phenotype-genotype association might be different either as a result of smoltification

or change in the environment (Aguirre *et al*, 2014). However, since individuals used in this study have a

complete or close to complete smolt phenotype, conclusions may be more relevant to the marine compared to

the freshwater life stage.

Increased vertebral thickness associated with the *vgll3*^{E} allele could provide stiffness for energy efficient

undulation movement during steady swimming (Donatelli *et al*, 2021), though it may come at the expense of

body flexibility that may facilitate piscivorous prey capture. A larger aerobic muscle area and thicker vertebra

may eventually facilitate earlier maturation via more energy efficient resource acquisition at sea, suggesting

that local adaptation for optimal age structure could be partly mediated by conditions at sea. Hence, while our

results do not explain causation of early male maturation in fresh water, we cannot explicitly rule out that the

link between *vgll3* and age at maturity is mediated via its effect on body shape. Sex was also an important

factor shaping the trait variation. Intriguingly, significant change in shape associated with sex and *vgll3*

genotype was also in concordant with respect to maturation timing. As such, males, which on average mature

earlier than females, and *vgll3* allele associated with early maturation had similar direction of change in these

morphological traits.

Larger fin sizes in fish increase maneuverability and stability (Webb, 1984), and correspondingly in Atlantic

salmon larger fins occur in populations that occupy faster flowing streams (Riddell and Leggett, 1981). Overall,

we found that the ray-fins of fish were consistently larger for the *vgll3*^{E} allele than the *vgll3*^{L} allele

(Supplementary Table 4). However, this improvement for maneuverability and stability may come in the

expense of efficiency in longer marine migrations, as a result of increased drag. For example, in coho salmon,

juveniles of inland population with longer migration distances have smaller fins (Taylor and McPhail, 1985),

which may help to reduce drag in long migrations (Dynes *et al*, 2005; Varian and Nichols, 2010). A larger cross

section diameter and larger fin lengths associated with the *vgll3*^{E} allele likely facilitate maneuvering at the

expense of efficient cruising, which is in contrast to the *vgll3*^{E} effect that increases aerobic muscle area and

vertebral thickness. Similarly, the slender body (a smaller cross section diameter) associated with *vgll3*^{L} is

consistent with minimizing the drag for cruising body-form (e.g., Webb, 1988), advantageous for a multi-sea

winter strategy that migrates longer distances in the ocean (e.g., O'Sullivan *et al*, 2022). On the other hand, the

direction of effect on fin size associated with *vgll3*^{E} *to vgll3*^{L} substitution is concordant with the effect of

restricted-food regime, suggesting that the allelic substation effect may be correlated to food limitation

conditions. Thus overall, the direction of *vgll3* genetic effects are not functionally consistent in relation to

proposed ecological roles of these morphological traits.

The changes associated with the genetic variation in *six6* locus that are affecting body and head sizes are linked

to ecological processes related to resource acquisition mechanisms. A larger head (and a smaller body)

associated with the *six6*^{L} allele and larger snout length may be associated with feeding regime towards

piscivory (Keeley *et al*, 2005), or generally consuming larger prey items. This interpretation corroborates with

the study by Aykanat *et al* (2020), in which fish with *six6*^{LL} genotype were shown to be more efficient in

resource acquisition than fish with *six6*^{EE} genotype.

Evolvabilities (I_A) ranged from normal to very high compared to a recent meta-analysis (Hansen *et al*, 2011).

The median evolvability for length traits is 0.001 (or 0.1%, Hansen and Pélabon, 2021; Hansen *et al*, 2011).

Normal evolvability values were found for centroid sizes and cross-section diameters, while aerobic muscle and

vertebrae thickness had high evolvabilities, suggesting that these traits could evolve rapidly in response to

selection induced by environmental change. Interestingly, a substantial portion of the additive genetic variation

associated with some traits, such as aerobic muscle area and vertebrate thickness, was explained by the *vgll3*

locus, suggesting the locus' importance in shaping the additive genetic landscape.

Taken together, *vgll3* and *six6* loci have complex associations with morphology. These do not consistently align

with adaptive differences between early- and late-maturation genotypes associated with marine migration, but

could be important e.g., during spawning where higher maneuverability of smaller males may be beneficial for

reproductive success, or in early life stages. On the other hand, it is also plausible that these changes are

genetically correlated as part of broad changes during development, and that some of which have potentially

maladaptive consequences. In fact, when traits are genetically correlated, the cumulative adaptive value of

trait values cannot be optimal (Schluter, 1996), since breaking suboptimal trait combinations that are

genetically correlated may be challenging for evolution (Blows, 2007).

It is important to recognize that extrapolation to the marine stage of Atlantic salmon based on juvenile

morphology may have limitations. It is unknown if the genotype-associated morphological variation is

correlated between juvenile and adult stages. While there may be genetic correlation between freshwater and

marine environments (Fleming *et al*, 1994; Morinville and Rasmussen, 2006; Taylor and McPhail, 1985), the

morphological changes associated with adaption to the marine environment, the parr-smolt transformation

(smoltification) may break apart trait correlations, similar to how metamorphosis takes places many animal

taxa (e.g., Aguirre *et al*, 2014). Since the data in this study were composed of individuals that had completed

the parr-smolt transformation, or are close to completion, conclusions are likely more relevant to marine

stages than the freshwater stages. However, any adaptive inference associated with a specific life stage should

be taken with caution, until observed patterns are further tested and verified at different life stages. Finally,

while it is reasonable to hypothesize that the observed *vgll3* and *six6* associated morphological variation is

adaptively important, the fitness effects of this variation should be better assessed before any evolutionary

conclusions can be made (Kingsolver *et al*, 1995).

In conclusion, we have detected substantial morphological variation associated with two genomic regions

which are also major determinants of life-history variation and known to be under local selection. These loci

appear to influence multiple, strongly correlated traits, suggestive of a broad regulatory role with impacts

across the whole life cycle. How these genetic correlations are shaped – via single locus pleiotropy or close

physical linkage of multiple genes – remains to be determined.

Acknowledgment

We thank Artur Porto for discussion related to morphometric analyses, and Annukka Ruokolainen, Seija

Tillanen, Andrew H. House, Jacqueline E. Moustakas-Verho, Nikolai Piavchenko for helps with sampling. This

work was supported by the Research Council of Finland (grant nos. 328860, 353388, and 325964 to TA, and

314254, 314255, 327255 and 342851 to CRP), and the European Research Council under the European Articles

Union's Horizon 2020 and Horizon Europe research and innovation programs (grant no. 742312 and 101054307

to CRP). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect

those of the European Union or the European Research Council Executive Agency. Neither the European Union

nor the granting authority can be held responsible for them. LG was supported by Erasmus+ programme. The

authors declare no competing interests.

Ethical approval

452 The experiments were approved by the Project Authorisation Board (ELLA) on behalf of the Regional

453 Administrative Agency for Southern Finland (ESAVI) under experimental license ESAVI/2778/2018.

Author contributions

TA, VLP, and GHB conceptualized the paper. TA analyzed the data with significant input from GHB and PVD. SJ and LG performed landmarkings. JE, and CRP oversaw the resources and infrastructure. PVD and CRP devised the breeding design and reared the fish. TA and PVD sampled the fish and digitized the images. TA drafted the MS with significant inputs from PVD, VLP, and GHB, and all coauthors contributed to subsequent drafts.

References:

Aguirre JD, Blows MW, Marshall DJ (2014). The genetic covariance between life cycle stages separated by metamorphosis. *Proceedings Biological sciences / The Royal Society* **281**(1788)**:** 20141091.

Ahi EP, Sinclair-Waters M, Moustakas-Verho J, Jansouz S, Primmer CR (2022). Strong regulatory effects of *vgll3* genotype on reproductive axis gene expression in juvenile male Atlantic salmon. *Gen Comp Endocrinol* **325:** 114055.

Altringham JD, Ellerby DJ (1999). Fish swimming: patterns in muscle function. *J Exp Biol* **202**(Pt 23)**:** 3397-3403.

Archambeault SL, Bartschi LR, Merminod AD, Peichel CL (2020). Adaptation via pleiotropy and linkage: Association mapping reveals a complex genetic architecture within the stickleback *Eda* locus. *Evol Lett* **4**(4)**:** 282-301.

Asheim ER, Debes PV, House A, Liljestrom P, Niemela PT, Siren JP *et al* (2023). Atlantic salmon (*Salmo salar*) age at maturity is strongly affected by temperature, population and age-at-maturity genotype. *Conserv Physiol* **11**(1)**:** coac086.

Aykanat T, McLennan D, Metcalfe NB, Prokkola JM (2024). Early survival in Atlantic salmon is associated with parental genotypes at loci linked to timing of maturation. *Evolution* **78**(8)**:** 1441-1452.

Aykanat T, Ozerov M, Vaha JP, Orell P, Niemela E, Erkinaro J *et al* (2019). Co-inheritance of sea age at maturity and iteroparity in the Atlantic salmon *vgll3* genomic region. *J Evol Biol* **32**(4)**:** 343-355.

Aykanat T, Rasmussen M, Ozerov M, Niemela E, Paulin L, Vaha JP *et al* (2020). Life-history genomic regions explain differences in Atlantic salmon marine diet specialization. *J Anim Ecol* **89**(11)**:** 2677-2691.

Ayllon F, Kjaerner-Semb E, Furmanek T, Wennevik V, Solberg MF, Dahle G *et al* (2015). The *vgll3* locus controls age at maturity in wild and domesticated Atlantic salmon (*Salmo salar* L.) Males. *PLoS Genet* **11**(11)**:** e1005628.

Baken EK, Collyer ML, Kaliontzopoulou A, Adams DC (2021). geomorph v4.0 and gmShiny: Enhanced analytics
and a new graphical interface for a comprehensive morphometric experience. *Methods in Ecology and*
Evolution **12**(12)**:** 2355-2363.

Bangura PB, Tiira K, Niemela PT, Erkinaro J, Liljestrom P, Toikkanen A *et al* (2022). Linking *vgll3* genotype and
aggressive behaviour in juvenile Atlantic salmon (*Salmo salar*). *J Fish Biol* **100**(5)**:** 1264-1271.

Barrett RD, Hoekstra HE (2011). Molecular spandrels: tests of adaptation at the genetic level. *Nat Rev Genet*
12(11)**:** 767-780.

Barson NJ, Aykanat T, Hindar K, Baranski M, Bolstad GH, Fiske P *et al* (2015). Sex-dependent dominance at a
single locus maintains variation in age at maturity in salmon. *Nature* **528**(7582)**:** 405-408.

Bates DM (2010). lme4: Mixed-effects modeling with R.

Baxter D, Cohen KE, Donatelli CM, Tytell ED (2022). Internal vertebral morphology of bony fishes matches the
mechanical demands of different environments. *Ecology and evolution* **12**(11)**:** e9499.

Blows MW (2007). A tale of two matrices: multivariate approaches in evolutionary biology. *J Evol Biol* **20**(1)**:** 1-
8.

Boyle EA, Li YI, Pritchard JK (2017). An Expanded View of Complex Traits: From Polygenic to Omnigenic. *Cell*
169(7)**:** 1177-1186.

Chan YF, Marks ME, Jones FC, Villarreal G, Jr., Shapiro MD, Brady SD *et al* (2010). Adaptive evolution of pelvic
reduction in sticklebacks by recurrent deletion of a *Pitx1* enhancer. *Science* **327**(5963)**:** 302-305.

Collyer ML, Adams DC, Freckleton R (2018). RRPP: An R package for fitting linear models to high-dimensional
data using residual randomization. *Methods in Ecology and Evolution* **9**(7)**:** 1772-1779.

Colosimo PF, Hosemann KE, Balabhadra S, Villarreal G, Jr., Dickson M, Grimwood J *et al* (2005). Widespread
parallel evolution in sticklebacks by repeated fixation of Ectodysplasin alleles. *Science* **307**(5717)**:** 1928-1933.

Crossin GT, Hinch SG, Farrell AP, Higgs DA, Lotto AG, Oakes JD *et al* (2004). Energetics and morphology of
sockeye salmon: effects of upriver migratory distance and elevation. *J Fish Biol* **65**(3)**:** 788-810.

Debes PV, Piavchenko N, Erkinaro J, Primmer CR (2020). Genetic growth potential, rather than phenotypic size,
predicts migration phenotype in Atlantic salmon. *Proceedings Biological sciences / The Royal Society* **287**(1931)**:**
20200867.

Debes PV, Piavchenko N, Ruokolainen A, Ovaskainen O, Moustakas-Verho JE, Parre N *et al* (2019). Large single-
locus effects for maturation timing are mediated via condition variation in Atlantic salmon. *bioRxiv***:** 780437
(v780433).

Debes PV, Piavchenko N, Ruokolainen A, Ovaskainen O, Moustakas-Verho JE, Parre N *et al* (2021). Polygenic
and major-locus contributions to sexual maturation timing in Atlantic salmon. *Mol Ecol* **30**(18)**:** 4505-4519.

Donatelli CM, Roberts AS, Scott E, DeSmith K, Summers D, Abu-Bader L *et al* (2021). Foretelling the Flex-
Vertebral Shape Predicts Behavior and Ecology of Fishes. *Integr Comp Biol* **61**(2)**:** 414-426.

Dynes J, Magnan P, Bernatchez L, Rodríguez MA (2005). Genetic and morphological variation between two
forms of lacustrine brook charr. *J Fish Biol* **54**(5)**:** 955-972.

Eliason EJ, Farrell AP (2015). Oxygen uptake in Pacific salmon *Oncorhynchus* spp.: when ecology and physiology
meet. *J Fish Biol*.

Elmer KR, Kusche H, Lehtonen TK, Meyer A (2010). Local variation and parallel evolution: morphological and
genetic diversity across a species complex of neotropical crater lake cichlid fishes. *Philosophical transactions of*
the Royal Society of London Series B, Biological sciences **365**(1547)**:** 1763-1782.

Endler JA (2000). Adaptive Genetic Variation in the Wild. In: Mousseau TA, SinerVo B and Endler JA (eds)
Adaptive Genetic Variation in the Wild. Oxford University Press.

Ferry-Graham LA, Bolnick DI, Wainwright PC (2002). Using functional morphology to examine the ecology and
evolution of specialization. *Integr Comp Biol* **42**(2)**:** 265-277.

Fitzgerald DB, Winemiller KO, Sabaj Pérez MH, Sousa LM, Seymour C (2017). Using trophic structure to reveal
patterns of trait-based community assembly across niche dimensions. *Funct Ecol* **31**(5)**:** 1135-1144.

Fleming IA, Jonsson B, Gross MR (1994). Phenotypic Divergence of Sea-ranched, Farmed, and Wild Salmon. *Can*
J Fish Aquat Sci **51**(12)**:** 2808-2824.

Gelman A, Su YS (2020). arm: Data Analysis Using Regression and Multilevel/Hierarchical Models. R package
version 1.11-2.

Gower JC (1975). Generalized procrustes analysis. *Psychometrika* **40**(1)**:** 33-51.

Hansen TF, Pélabon C (2021). Evolvability: A Quantitative-Genetics Perspective. *Annual Review of Ecology,*
Evolution, and Systematics **52**(1)**:** 153-175.

Hansen TF, Pelabon C, Armbruster WS, Carlson ML (2003). Evolvability and genetic constraint in Dalechampia
blossoms: components of variance and measures of evolvability. *J Evol Biol* **16**(4)**:** 754-766.

574
575 Hansen TF, Pélabon C, Houle D (2011). Heritability is not Evolvability. *Evolutionary Biology* **38**(3)**:** 258-277.

576
577 Hoar WS (1988). 4 The Physiology of Smolting Salmonids. In: Hoar WS and Randall DJ (eds) *Fish Physiology*.
578 Academic Press. Vol. 11, pp 275-343.

579
580 Houle D (1992). Comparing evolvability and variability of quantitative traits. *Genetics* **130:** 195-204.

581
582 Jakobson CM, Jarosz DF (2020). What Has a Century of Quantitative Genetics Taught Us About Nature's Genetic
583 Tool Kit? *Annu Rev Genet* **54:** 439-464.

584
585 Jimenez-Gomez JM, Wallace AD, Maloof JN (2010). Network analysis identifies *ELF3* as a QTL for the shade
586 avoidance response in Arabidopsis. *PLoS Genet* **6**(9)**:** e1001100.

587
588 Jonsson B, Jonsson N (2011a). *Ecology of Atlantic salmon and brown trout : habitat as a template for life
589 histories*. Springer: Dordrecht.

590
591 Jonsson B, Jonsson N (2011b). Ecology of salmon and trout 2011.

592
593 Kawecki TJ, Ebert D (2004). Conceptual issues in local adaptation. *Ecol Lett* **7**(12)**:** 1225-1241.

594
595 Keeley ER, Parkinson EA, Taylor EB (2005). Ecotypic differentiation of native rainbow trout (*Oncorhynchus
596 mykiss*) populations from British Columbia. *Can J Fish Aquat Sci* **62**(7)**:** 1523-1539.

597
598 Kingsolver JG, Woods HA, Gilchrist G (1995). Traits related to fitness. *Science* **267**(5196)**:** 396.

599
600 Klingenberg CP (2016). Size, shape, and form: concepts of allometry in geometric morphometrics. *Dev Genes
601 Evol* **226**(3)**:** 113-137.

602
603 Kurko J, Debes PV, House AH, Aykanat T, Erkinaro J, Primmer CR (2020). Transcription Profiles of Age-at-
604 Maturity-Associated Genes Suggest Cell Fate Commitment Regulation as a Key Factor in the Atlantic Salmon
605 Maturation Process. *G3* **10**(1)**:** 235-246.

606
607 Kuznetsova A, Brockhoff PB, Christensen RHB (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *J
608 Stat Softw* **82**(13).

609
Long JH, Nipper KS (1996). The Importance of Body Stiffness in Undulatory Propulsion. *Am Zool* **36**(6)**:** 678-694.

Lundsgaard-Hansen B, Matthews B, Vonlanthen P, Taverna A, Seehausen O (2013). Adaptive plasticity and
genetic divergence in feeding efficiency during parallel adaptive radiation of whitefish (*Coregonus* spp.). *J Evol*
Biol **26**(3)**:** 483-498.

Luo L, Mao Y, Xu S (2003). Correcting the bias in estimation of genetic variances contributed by individual QTL.
Genetica **119**(2)**:** 107-113.

Lynch M, Walsh B (1998). *Genetics and analysis of quantitative traits*. Sinauer Associates: Sunderland, MA.

Mackay TF, Stone EA, Ayroles JF (2009). The genetics of quantitative traits: challenges and prospects. *Nat Rev*
Genet **10**(8)**:** 565-577.

Miettinen A, Dannewitz J, Palm S, Ahi EP, Romakkaniemi A, Vähä V *et al* (2023). Loci associated with
maturation, migration and appetite control are linked with signals of fine-scale local selection in a large Atlantic
salmon population. *bioRxiv***:** 2023.2008.2023.553800.

Mindel BL, Neat FC, Trueman CN, Webb TJ, Blanchard JL (2016). Functional, size and taxonomic diversity of fish
along a depth gradient in the deep sea. *PeerJ* **4:** e2387.

Mobley KB, Aykanat T, Czorlich Y, House A, Kurko J, Miettinen A *et al* (2021). Maturation in Atlantic salmon
(*Salmo salar*, Salmonidae): a synthesis of ecological, genetic, and molecular processes. *Rev Fish Biol Fish* **31**(3)**:**
523-571.

Morinville GR, Rasmussen JB (2006). Does life-history variability in salmonids affect habitat use by juveniles? A
comparison among streams open and closed to anadromy. *J Anim Ecol* **75**(3)**:** 693-704.

Morinville GR, Rasmussen JB (2007). Distinguishing between juvenile anadromous and resident brook trout
(Salvelinus fontinalis) using morphology. *Environ Biol Fishes* **81**(2)**:** 171-184.

Moustakas-Verho JE, Kurko J, House AH, Erkinaro J, Debes P, Primmer CR (2020). Developmental expression
patterns of *six6*: A gene linked with spawning ecotypes in Atlantic salmon. *Gene Expr Patterns* **38:** 119149.

Nielsen EE, Hemmer-Hansen J, Larsen PF, Bekkevold D (2009). Population genomics of marine fishes:
identifying adaptive variation in space and time. *Mol Ecol* **18**(15)**:** 3128-3150.

O'Sullivan RJ, Ozerov M, Bolstad GH, Gilbey J, Jacobsen JA, Erkinaro J *et al* (2022). Genetic stock identification
reveals greater use of an oceanic feeding ground around the Faroe Islands by multi-sea winter Atlantic salmon,
with variation in use across reporting groups. *ICES J Mar Sci* **79**(9)**:** 2442-2452.

Olson-Manning CF, Wagner MR, Mitchell-Olds T (2012). Adaptive evolution: evaluating empirical support for
theoretical predictions. *Nat Rev Genet* **13**(12)**:** 867-877.

653
654 Orr HA (2000). Adaptation and the cost of complexity. *Evolution* **54**(1)**:** 13-20.

655
656 Otto SP (2004). Two steps forward, one step back: the pleiotropic effects of favoured alleles. *Proceedings
657 Biological sciences / The Royal Society* **271**(1540)**:** 705-714.

658
659 Paaby AB, Rockman MV (2013). The many faces of pleiotropy. *Trends Genet* **29**(2)**:** 66-73.

660
661 Peris Tamayo AM, Devineau O, Praebel K, Kahilainen KK, Ostbye K (2020). A brain and a head for a different
662 habitat: Size variation in four morphs of Arctic charr (*Salvelinus alpinus* (L.)) in a deep oligotrophic lake. *Ecology
663 and evolution* **10**(20)**:** 11335-11351.

664
665 Porto A, Voje KL, Price S (2020). ML-morph: A fast, accurate and general approach for automated detection and
666 landmarking of biological structures in images. *Methods in Ecology and Evolution* **11**(4)**:** 500-512.

667
668 Pritchard VL, Makinen H, Vaha JP, Erkinaro J, Orell P, Primmer CR (2018). Genomic signatures of fine-scale local
669 selection in Atlantic salmon suggest involvement of sexual maturation, energy homeostasis and immune
670 defence-related genes. *Mol Ecol* **27**(11)**:** 2560-2575.

671
672 Prokkola JM, Asheim ER, Morozov S, Bangura P, Erkinaro J, Ruokolainen A *et al* (2022). Genetic coupling of life-
673 history and aerobic performance in Atlantic salmon. *Proceedings Biological sciences / The Royal Society*
674 **289**(1967)**:** 20212500.

675
676 Proulx R, Magnan P (2004). Contribution of phenotypic plasticity and heredity to the trophic polymorphism of
677 lacustrine brook charr (*Salvelinus fontinalis* M.). *Evol Ecol Res* **6**(4)**:** 503-522.

678
679 R Core Team. (2019). R Foundation for Statistical Computing, Vienna, Austria.

680
681 Riddell BE, Leggett WC (1981). Evidence of an adaptive basis for geographic variation in body morphology and
682 time of downstream migration of juvenile Atlantic salmon (*Salmo salar*). *Can J Fish Aquat Sci* **38**(3)**:** 308-320.

683
684 Schluter D (1996). Ecological causes of adaptive radiation. *Am Nat*.

685
686 Shuman JL, Coughlin DJ (2018). Red muscle function and thermal acclimation to cold in rainbow smelt, *Osmerus
687 mordax*, and rainbow trout, *Oncorhynchus mykiss*. *J Exp Zool A Ecol Integr Physiol* **329**(10)**:** 547-556.

689   Simonsen MK, Siwertsson A, Adams CE, Amundsen PA, Praebel K, Knudsen R (2017). Allometric trajectories of
690   body and head morphology in three sympatric Arctic charr (*Salvelinus alpinus* (L.)) morphs. *Ecology and
691   evolution* **7**(18)**:** 7277-7289.

692
693   Sinclair-Waters M, Odegard J, Korsvoll SA, Moen T, Lien S, Primmer CR *et al* (2020). Beyond large-effect loci:
694   large-scale GWAS reveals a mixed large-effect and polygenic architecture for age at maturity of Atlantic
695   salmon. *Genet Sel Evol* **52**(1)**:** 9.

696
697   Snorrason S (1994). Trophic specialization in Arctic charr *Salvelinus alpinus* (Pisces; Salmonidae): morphological
698   divergence and ontogenetic niche shifts. *Biol J Linn Soc* **52**(1)**:** 1-18.

699
700   Solovieff N, Cotsapas C, Lee PH, Purcell SM, Smoller JW (2013). Pleiotropy in complex traits: challenges and
701   strategies. *Nat Rev Genet* **14**(7)**:** 483-495.

702
703   Svanbäck R, Eklöv P (2003). Morphology dependent foraging efficiency in perch: a trade-off for ecological
704   specialization? *Oikos* **102**(2)**:** 273-284.

705
706   Taylor EB, McPhail JD (1985). Variation in body morphology among british columbia populations of coho
707   salmon, *Oncorhynchus kisutch*. *Can J Fish Aquat Sci* **42**(12)**:** 2020-2028.

708
709   Varian A, Nichols KM (2010). Heritability of morphology in brook trout with variable life histories. *PLoS One*
710   **5**(9)**:** e12950.

711
712   Vazquez AI, Bates DM, Rosa GJ, Gianola D, Weigel KA (2010). Technical note: an R package for fitting
713   generalized linear mixed models in animal breeding. *J Anim Sci* **88**(2)**:** 497-504.

714
715   Verta JP, Debes PV, Piavchenko N, Ruokolainen A, Ovaskainen O, Moustakas-Verho JE *et al* (2020). Cis-
716   regulatory differences in isoform expression associate with life history strategy variation in Atlantic salmon.
717   *PLoS Genet* **16**(9)**:** e1009055.

718
719   Verta JP, Moustakas-Verho JE, Donner I, Frapin M, Ruokolainen A, Debes PV *et al* (2024). A complex mechanism
720   translating variation of a simple genetic architecture into alternative life histories. *Proc Natl Acad Sci U S A*
721   **121**(48)**:** e2402386121.

722
723   Wagner GP, Altenberg L (1996). Perspective: Complex Adaptations and the Evolution of Evolvability. *Evolution*
724   **50**(3)**:** 967-976.

725
726   Wagner GP, Zhang J (2011). The pleiotropic structure of the genotype-phenotype map: the evolvability of
727   complex organisms. *Nat Rev Genet* **12**(3)**:** 204-213.

728
729   Webb PW (1984). Body form, locomotion and foraging in aquatic vertebrates. *Am Zool* **24**(1)**:** 107-120.

730
731 Webb PW (1988). Simple physical principles and vertebrate aquatic locomotion. *Am Zool* **28**(2)**:** 709-725.

732
733 Weith M, Grossbach J, Clement-Ziza M, Gillet L, Rodriguez-Lopez M, Marguerat S *et al* (2023). Genetic effects
734 on molecular network states explain complex traits. *Mol Syst Biol* **19**(8)**:** e11493.

735
736 Ziyatdinov A, Vazquez-Santiago M, Brunel H, Martinez-Perez A, Aschard H, Soria JM (2018). lme4qtl: linear
737 mixed models with flexible covariance structure for genetic studies of related individuals. *BMC Bioinformatics*
738 **19**(1)**:** 68.

739
740 Zueva KJ, Lumme J, Veselov AE, Primmer CR, Pritchard VL (2020). Population genomics reveals repeated signals
741 of adaptive divergence in the Atlantic salmon of north-eastern Europe. *J Evol Biol* **34**(6)**:** 866-878.

Tables (4) and Figures (2):

Table 1: Estimated fixed effect coefficients for 21 morphological traits obtained across three body planes. Trait lengths are given in centimeters (cm) for all traits. Coef. (SE) indicates coefficients and their standard error. B, S, and L in the body plane column indicates lateral view of the body, small and large cross sections, respectively. Significant *p* values are indicated with asterisk (* and ** denotes *p*<0.05 and *p*<0.01, respectively).

Trait	Body Plane[1]	Trait value formula[1]	Intercept coef. (SE)	Length (scaled) coef. (SE)	*p* value	Condition factor (scaled) coef. (SE)	*p* value	Feed restriction (*ad libitum* to restricted) coef. (SE)	*p* value	Sex (female to male) coef. (SE)	*p* value	vgll3 (E to L) coef. (SE)	*p* value	six6 (E to L) coef. (SE)	*p* value
Dorsal fin	B	B [12:13]	2.0439 (0.0305)	0.1560 (0.0064)	<0.001**	-0.0016 (0.0041)	0.696	-0.1222 (0.0176)	<0.001**	0.0228 (0.0124)	0.067	-0.0246 (0.0148)	0.106	-0.0142 (0.0126)	0.263
Anal fin	B	B [16:17]	1.35902 (0.0164)	0.06636 (0.0041)	<0.001**	0.00043 (0.0038)	0.909	-0.02549 (0.0094)	0.0402*	0.01759 (0.0076)	0.021*	-0.01979 (0.0077)	0.017*	0.00658 (0.0075)	0.378
Pectoral fin	B	B [10:11]	2.0973 (0.0202)	0.1148 (0.0029)	<0.001**	0.0082 (0.0022)	<0.001**	-0.0376 (0.0151)	0.059	0.0143 (0.0053)	0.007**	-0.0102 (0.0096)	0.29	0.0030 (0.0058)	0.601
Tail fin	B	B [19:20]	8.3e-01 (0.0098)	4.1e-02 (0.0015)	<0.001**	-8.2e-03 (0.0013)	<0.001**	-2.1e-02 (0.0037)	0.003**	2.8e-05 (0.0027)	0.992	-4.0e-03 (0.0051)	0.432	3.1e-03 (0.0030)	0.292
Caudal peduncle	B	B [15:18]	8.7e-01 (0.00428)	6.0e-02 (0.00078)	<0.001**	1.1e-02 (0.00071)	<0.001**	1.3e-03 (0.00198)	0.529	2.3e-03 (0.00144)	0.118	-2.3e-03 (0.00219)	0.294	-6.7e-05 (0.00153)	0.965
Vertebra thickness	Sml	S [3:4]	2.4e-01 (0.00265)	1.6e-02 (0.00081)	<0.001**	1.8e-03 (0.00082)	0.0241*	2.3e-03 (0.00165)	0.172	-2.2e-03 (0.00157)	0.168	-3.3e-03 (0.00117)	0.009**	5.3e-05 (0.00138)	0.969
Vertebra thickness	Lrg	L [6:7]	0.25170 (0.00282)	0.01394 (0.00077)	<0.001**	0.00467 (0.00078)	<0.001**	-0.00238 (0.00155)	0.125	-0.00244 (0.00146)	0.096	-0.00297 (0.00131)	0.032*	0.00023 (0.00137)	0.866
Cross section diameter	Sml	S [5:7]	1.1423 (0.0136)	0.0753 (0.0027)	<0.001**	0.0134 (0.0027)	<0.001**	-0.0562 (0.0114)	0.003**	0.0020 (0.0049)	0.683	-0.0157 (0.0058)	0.012*	-0.0054 (0.0050)	0.279
Cross section diameter	Lrg	L [12:8]	2.0269 (0.0117)	0.1477 (0.0021)	<0.001**	0.0312 (0.0021)	<0.001**	0.0050 (0.0108)	0.666	0.0019 (0.0039)	0.621	-0.0047 (0.0048)	0.335	-0.0046 (0.0039)	0.248
Aerobic muscle	Sml	S [(7:8+5:6)/2]	0.17549 (0.00353)	0.01024 (0.00082)	<0.001**	0.00137 (0.00083)	0.0981	-0.01237 (0.00290)	0.0051**	0.00056 (0.00153)	0.715	-0.00557 (0.00148)	<0.001**	-0.00210 (0.00149)	0.159
Aerobic muscle	Lrg	L [(8:9+12:13)/2]	0.2349 (0.00408)	0.0185 (0.00094)	<0.001**	0.0033 (0.00094)	<0.001**	0.0070 (0.00239)	0.025*	0.0017 (0.00175)	0.34	-0.0024 (0.00195)	0.222	-0.0022 (0.00174)	0.198
Aerobic muscle vertical extension	Lrg	L [(14:15+10:11)/2]	0.879 (0.0198)	0.087 (0.0039)	<0.001**	0.026 (0.0039)	<0.001**	0.038 (0.0176)	0.0916	-0.012 (0.0073)	0.089	-0.024 (0.0081)	0.006**	0.013 (0.0072)	0.067
Eye diameter[2]	B	B [(2:3+4:5)/2]	5.5e-01 (0.00358)	2.2e-02 (0.00094)	<0.001**	3.3e-04 (0.00092)	0.723	1.8e-03 (0.00285)	0.546	-1.7e-06 (0.00175)	0.999	-3.2e-03 (0.00152)	0.055	2.6e-03 (0.00163)	0.112
Snout length	B	B [1:0]	0.78061 (0.00458)	0.03839 (0.00097)	<0.001**	-0.00054 (0.00083)	0.509	-0.00277 (0.00196)	0.157	0.00557 (0.00183)	0.002**	-0.00607 (0.00232)	0.013*	0.00575 (0.00189)	0.002**
Eye vertical position	B	B [0:6]	0.40996 (0.0052)	0.02487 (0.0012)	<0.001**	0.00460 (0.0010)	<0.001**	0.00934 (0.0036)	0.0538	0.00024 (0.0022)	0.912	-0.00031 (0.0024)	0.899	0.00057 (0.0022)	0.794
Head length[3]	B	B [1:7]	2.1453 (0.0232)	0.1347 (0.0034)	<0.001**	0.0074 (0.0025)	0.0034*	-0.0370 (0.0224)	0.162	0.0106 (0.0064)	0.094	-0.0074 (0.0098)	0.455	0.0146 (0.0068)	0.032*
Head width[3]	B	B [8:9]	1.59855 (0.0052)	0.09654 (0.0015)	<0.001**	0.01633 (0.0013)	<0.001**	0.02432 (0.0030)	<0.001**	0.00420 (0.0028)	0.13	-0.00046 (0.0024)	0.85	0.00030 (0.0026)	0.907
Body centroid size	B	From Procrustes analysis	13.1556 (0.0201)	0.9084 (0.0031)	<0.001**	0.0090 (0.0026)	<0.001**	0.0505 (0.0124)	0.0118*	-0.0088 (0.0057)	0.124	0.0253 (0.0099)	0.014*	-0.0227 (0.0061)	<0.001**
Head centroid size	B	From Procrustes analysis	2.2095 (0.0131)	0.1312 (0.0020)	<0.001**	0.0143 (0.0016)	<0.001**	-0.0073 (0.0144)	0.632	0.0077 (0.0037)	0.040*	-0.0075 (0.0050)	0.148	0.0104 (0.0039)	0.008**
Centroid size	Sml	From Procrustes analysis	1.4166 (0.0129)	0.0968 (0.0023)	<0.001**	0.0199 (0.0023)	<0.001**	-0.0387 (0.0102)	0.0111*	0.0033 (0.0042)	0.432	-0.0119 (0.0058)	0.048*	-0.0048 (0.0044)	0.28
Centroid size	Lrg	From Procrustes analysis	3.5944 (0.0177)	0.2598 (0.0029)	<0.001**	0.0601 (0.0029)	<0.001**	0.0202 (0.0133)	0.209	-0.0015 (0.0053)	0.778	-0.0097 (0.0082)	0.244	-0.0054 (0.0056)	0.337

1: The trait values are based on landmarks as numbered and indicated in Supplementary Figure 2.
2: Eye origin is the center point of eye landmarks, and calculated using landmarks 2, 3, 4, and 5, which are the anterior, posterior, dorsal and ventral extremes of the bony orbit of the eye.
3: Landmark 7 is the posterior point of the bony operculum. Landmarks 8 and 9 are the points of maximum curvature of the bony operculum at the ventral and dorsal ends, respectively.

Table 2: Multivariate analysis using Procrustes shape variable across three body planes in juvenile Atlantic salmon. Pillai is the Pillai's trace test statistic to evaluate each term's significance in the multivariate analysis of variance.

Terms	Large cross section				Small cross section				Body			
	df	Pillai	z value	p value	df	Pillai	z value	p value	df	Pillai	z value	p value
Length (scaled)	1	0.207	8.231	0.001	1	0.042	2.574	0.008	1	0.530	15.369	0.001
Condition factor (scaled)	1	0.053	1.398	0.074	1	0.024	0.894	0.195	1	0.478	10.583	0.001
Feed restriction	1	0.096	3.924	0.001	1	0.102	6.035	0.001	1	0.284	12.050	0.001
Family	37	1.939	9.994	0.001	37	0.721	2.200	0.011	37	2.446	24.524	0.001
Tank	4	0.312	6.244	0.001	4	0.219	7.020	0.001	3	0.276	8.643	0.001
Sex	1	0.035	-0.118	0.548	1	0.013	-0.597	0.717	1	0.038	1.185	0.121
vgll3	1	0.042	0.495	0.318	1	0.018	0.108	0.461	1	0.032	0.562	0.285
six6	1	0.027	-0.992	0.831	1	0.013	-0.617	0.739	1	0.052	2.454	0.013
Full model	47	2.727	14.021	0.001	47	1.256	7.674	0.001	46	3.929	23.997	0.001
Residuals	689				681				724			

Table 3: Variance components for 21 morphological traits obtained from three body planes. V_A, h^2, and I_A are additive genetic variance, heritability, and evolvability, respectively, after controlling for length, condition factor, feed restriction, and sex. % $V_{A.vgll3}$ and %$V_{A.six6}$ are additive genetic variation explained by the *vgll3* and *six6* loci, respectively (values are truncated to zero if the estimate is less than zero). Values indicate medians, with 95% CI given in parenthesis.

Trait [1]	Body Plane [2]	V_A	% $V_{A.vgll3}$	%$V_{A.six6}$	h^2	I_A [3]	% rep
Dorsal fin	B	0.0032 (0.0027 - 0.0036)	2.6	0	0.083 (0.072 - 0.095)	0.045 (0.042 - 0.048)	95.2
Anal fin	B	7,7e-04 (6.8 e-04, 8.8e-04)	17.2	0	0.07 (0.061 - 0.08)	0.27 (0.26 - 0.3)	67.3
Pectoral fin	B	0.0027 (0.0024, 0.003)	0	0	0.41 (0.36 - 0.45)	0.016 (0.015 - 0.017)	83.7
Tail fin	B	8.2e-04 (7.3e-04, 9.1e-04)	0	0	0.5 (0.47 - 0.54)	0.051 (0.048 - 0.055)	69.7
Caudal peduncle	B	1e-04 (9.1e-05, 1.2e-04)	0	0	0.26 (0.24 - 0.29)	0.02 (0.019 - 0.021)	65.3
Vertebra thickness	Sml	1.3e-05 (1.1e-05, 1.5e-05)	30.7	0	0.039 (0.033 - 0.045)	0.31 (0.28 - 0.35)	63.7
Vertebra thickness	Lrg	2.0e-05 (1.7e-05, 2.3e-05)	13	0	0.06 (0.052 - 0.069)	0.21 (0.19 - 0.23)	71.4
Cross section diameter	Sml	6.8e-04 (5.9e-04, 7.9e-04)	12.7	0	0.12 (0.11 - 0.14)	0.0099 (0.009 - 0.011)	97.6
Cross section diameter	Lrg	3.3e-04 (2.9e-04, 3.8 e-04)	0	0	0.11 (0.093 - 0.12)	0.012 (0.01 - 0.013)	84.1
Aerobic muscle	Sml	5.8e-05 (5.1e-05, 6.7e-05)	22.4	0	0.19 (0.17 - 0.21)	0.69 (0.62 - 0.76)	56.5
Aerobic muscle	Lrg	7.5e-05 (6.6e-05, 8.7e-05)	0	0	0.37 (0.34 - 0.41)	0.72 (0.65 - 0.79)	24.0
Aerobic muscle vertical extension	Lrg	0.001 (8.6 e-04, 0.0012)	22.2	2.3	0.1 (0.087 - 0.12)	0.29 (0.26 - 0.32)	78.5
Eye diameter	B	3.1e-05 (2.7e-05, 3.5e-05)	8.6	1.6	0.11 (0.094 - 0.12)	0.23 (0.21 - 0.24)	25.9
Snout length	B	9.0e-05 (7.9e-05, 1e-04)	14.1	9.4	0.12 (0.11 - 0.14)	0.023 (0.021 - 0.024)	82.4
Eye vertical position	B	6.4e-05 (5.6e-05, 7.3e-05)	0	0	0.06 (0.052 - 0.069)	0.15 (0.14 - 0.16)	79.7
Head length	B	0.0022 (0.0019, 0.0025)	0	1.8	0.22 (0.19 - 0.26)	0.018 (0.017 - 0.02)	88.9
Head width	B	3.5e-05 (3e-05, 4e-05)	0	0	0.021 (0.018 - 0.025)	0.02 (0.019 - 0.022)	75.4
Body centroid size	B	0.0034 (0.0031, 0.0038)	6.3	4.2	0.47 (0.43 - 0.5)	7.6e-04 (7,1e-04, 8.2e-04)	73.8
Head centroid size	B	5e-04 (4.4e-04, 5.7e-04)	0.5	5.1	0.15 (0.12 - 0.17)	0.009 (0.0084 - 0.0096)	85.1
Centroid size	Sml	8.3e-04 (7.2e-04, 9.5e-04)	4.4	0	0.27 (0.24 - 0.3)	0.043 (0.039 - 0.048)	71.5
Centroid size	Lrg	0.0014 (0.0012, 0.0016)	0	0	0.22 (0.19 - 0.24)	0.0025 (0.0023 - 0.0028)	93.5

1: Trait length unit is cm for all traits.
2: The trait values are based on landmarks as numbered and indicated in Supplementary Figure 2. B, S, and L prefixes represent body, small cross section, and large cross sections, respectively.
3: I_A is measured as V_A divided by the squared trait mean, and it is unit free.

Figure 1: The difference between *vgll3*^{EE} and *vgll3*^{LL} (blue lines), and *six6*^{EE} and *six6*^{LL} (red lines) genotypes on body shape of juvenile Atlantic salmon using thin-plate spline in the body (a), and small (b) and large (b) cross sections. The gray dots are the predicted shape coordinate associated with *vgll3*^{EE} and *six6*^{EE} after the generalized Procrustes analysis. For all images, the effect sizes are magnified by 10x. Landmark numbers are based on Supplementary Figure 2. Note that the body and cross section images are to illustrate the location of landmarks on these body planes, and were slightly distorted to fit onto the average coordinates obtained by Procrustes analysis.

Figure 2: Percent trait value change associated with *vgll3*[E] to *vgll3*[L] (red lines), and *six6*[E] to *six6*[L] (blue lines) allelic substitutions relative to the average trait values. Asterisks denote significant genetic effect. Error bars and 1SE in the x-axis indicate one standard error.