1    Large effect life-history genomic regions are associated with functional morphological

2    traits in Atlantic salmon

4    Tutku Aykanat[1], Paul V. Debes[2,6], Shadi Jansouz[1], Lison Gueguen[3], Andrew H. House[1], Annukka Ruokolainen[1],

5    Jaakko Erkinaro[4], Victoria L. Pritchard[5], Craig R. Primmer[1,6], Geir H. Bolstad[7]

6        1. Organismal and Evolutionary Biology Research Program, Faculty of Biological and Environmental Sciences, PO Box 56, FI-
7           00014 University of Helsinki, Finland. tutku.aykanat@helsinki.fi.

2. Department of Aquaculture and Fish Biology, Hólar University, 550 Sauðárkrókur, Iceland.

3. Oniris Graduate School of Veterinary Medicine, Nantes, France.

4. Natural Resources Institute Finland (Luke), Oulu, Finland

5. Institute for Biodiversity & Freshwater Conservation, UHI Inverness, Inverness, United Kingdom., United Kingdom.

6. Institute of Biotechnology, Helsinki Institute of Life Science (HiLIFE), University of Helsinki, Helsinki, Finland.

7. Norwegian Institute for Nature Research (NINA), NO-7485 Trondheim, Norway.

**Short running title:** Functional morphology linked to major life history loci

**Keywords:** Atlantic salmon, morphology, morphometrics, adaptive genetic variation, age at maturity,

pleiotropy, *six6*, *vgll3*

## Abstract

Understanding pleiotropic architectures of phenotypes is instrumental for identifying the functional basis of adaptive genetic variation in the wild. Life-history variation may have a morphological basis that mediates resource acquisition allocation pathways, but identifying the underlying genetic basis of such traits is challenging. Using Atlantic salmon juveniles reared in common garden conditions, we test if two life-history associated loci in Atlantic salmon, s*ix6* and *vgll3,* are also associated with functional morphological traits. These loci had previously shown to exhibit strong signals of adaptation and are highly correlated with sea age at maturity. We show that genetic variation at the *vgll3* locus is linked to variation in morphological traits that underlie swimming performance, along a trade-off axis between efficient cruising and maneuvering, while the genetic variation at the *six6* locus was linked to variation in body-head proportions suggesting the potential functional importance of these traits for resource acquisition efficiency. However, the direction of changes in morphological traits associated with late- vs. early- maturing alleles was not always consistent with the expected direction of an effect to maturation timing. Our results reveal a complex morphological landscape associated with the genetic variation in these loci, possibly as a result of pleiotropy or linkage across these genomic regions.

**Keywords:** Atlantic salmon, morphology, morphometrics, adaptive genetic variation, age at maturity, pleiotropy, *six6*, *vgll3*

## Introduction

Elucidating the genotype-phenotype relationship is instrumental in identifying the functional basis of adaptation and the potential targets of selection (Barrett and Hoekstra, 2011; Kawecki and Ebert, 2004; Nielsen *et al*, 2009; Wagner and Altenberg, 1996). Methodological and statistical advances in genetics now allow the identification of genomic regions responding to local selection, but the phenotypic link between genotype and fitness is often missing. Furthermore, many genomic regions can affect a multitude of phenotypic traits (via pleiotropy or linkage disequilibrium), which makes identifying the phenotypic traits under selection even more challenging (Paaby and Rockman, 2013; Solovieff *et al*, 2013). Even large, genome-wide association studies might be underpowered to detect pleiotropy (Wagner and Zhang, 2011), a methodological bias by which the adaptive significance of pleiotropy can be overlooked especially in natural, non-model populations even if it may have substantial effects to limit or promote adaptation (Orr, 2000; Otto, 2004). In natural systems, pleiotropy has been mostly studied at the phenotypic level to understand the evolutionary trade-offs by quantifying the physiological antagonisms between senescence and reproduction age. Yet, some studies identify pleiotropic effects of genes associated with adaptive genetic variation. For example, adaptive divergence between lake vs. marine morphs of threespine stickleback (*Gasterosteus aculeatus*) was explained by both pleiotropy in the *Eda* gene, and via strong linkage with the surrounding loci (Archambeault *et al*, 2020), illustrating the significance of such genetic architectures for adaptation.

Genomic regions with large adaptive importance are more likely to be detected in genome scans  and are likely to have broad cellular and phenotypic effects (Shapiro *et al*, 2003, Jones *et al*, 2012, , Barson *et al*, 2015). Major effect loci may also have broad cellular and phenotypic effects, with key regulatory roles across development, and could thereby drive strong genetic interdependences among traits (Chan *et al*, 2010; Colosimo *et al*, 2005; Jimenez-Gomez *et al*, 2010; Weith *et al*, 2023) but see (Boyle *et al*, 2017; Jakobson and Jarosz, 2020). Therefore, loci exhibiting strong signals of adaptation may provide feasible empirical approaches to understand the significance of complex evolutionary processes, such as how pleiotropy and genetic interactions may hinder or promote adaptation in relation to the direction of natural selection between co-varying traits. Yet, such examples remain rare in non-model, wild organisms (Archambeault *et al*, 2020; Endler, 2000; Mackay *et al*, 2009; Olson-Manning *et al*, 2012).

Functional variation in morphological traits is integral in shaping resource use, and is correspondingly associated with life-history variation and ecotypic divergence in a large array of fish species, including salmonid fishes (Ferry-Graham *et al*, 2002; Fitzgerald *et al*, 2017; Keeley *et al*, 2005; Lundsgaard-Hansen *et al*, 2013; Mindel *et al*, 2016; Proulx and Magnan, 2004; Svanbäck and Eklöv, 2003; Taylor and McPhail, 1985; Varian and

Nichols, 2010; Webb, 1988). For example, a fusiform body and shorter fins appear adaptive to a long-distance

migratory strategy, likely because this body shape reduces the drag (hence the energy consumed) during

migration (Morinville and Rasmussen, 2007). Likewise, a robust body shape with a larger head, caudal

peduncle, and paired fins has been associated with piscivory in rainbow trout (*Oncorhynchus mykiss*, Keeley *et*

*al*, 2005). Recent population genomics studies in Atlantic salmon have consistently identified two unlinked

genomic regions, surrounding the genes *six6* in chromosome 9 and *vgll3* in chromosome 25 as strong targets

for local selection within all European Atlantic salmon lineages (Barson *et al*, 2015; Miettinen *et al*, 2023;

Pritchard *et al*, 2018; Zueva *et al*, 2020). These two loci have also been linked to a suite of life-history traits that

strongly influence fitness. In particular, the *vgll3* and *six6* loci are strongly associated with sea age at first

maturity (age at maturity from here on) in seagoing salmon at the individual- and population- levels

respectively (Barson *et al*, 2015, see also, Ayllon *et al*, 2015; Sinclair-Waters *et al*, 2020), and the *vgll3* locus

affects other alternative life-history strategies, such as precocious male maturation (Debes *et al*, 2021), and

iteroparity (Aykanat *et al*, 2019). For example, individuals with late maturation genotype in the *vgll3* gene

(*vgll3*[LL]) tended to stay at sea, on average, more than one year more than individuals with early maturation

(*vgll3*[EE]) genotype (Barson *et al*, 2015, Mobley *et al*, 2021). This pattern was also pertinent during the

freshwater phase within which juvenile males with *vgll3*[LL] genotype matured at a much higher probability (~

60%) than juvenile males with *vgll3*[LL] genotype (Debes *et al*, 2021).

Similarly, these loci explain variation across a number of metabolic and behavior traits. For example, *vgll3*[E]

allele has been shown to be associated with reduced aggression, increased maximum metabolic rate and

aerobic scope at the juvenile stage (Bangura *et al*, 2022; Prokkola *et al*, 2022), while *six6* locus predicts diet

acquisition patterns at sea in maturing adults with *six6*[E] allele linked to increased stomach weight despite

decreased foraging efforts (Aykanat *et al*, 2020). Likewise, *six6* locus exhibits epistasis with *vgll*3 to affect

aerobic scope variation (Prokkola *et al*, 2022). The strong genetic interdependence of these phenotypic traits

across two loci points to the potential for a shared developmental origin (of phenotypes to the respective loci),

which is supported by the role of *vgll3* and *six6* in the hormonal and cell-fate commitment gene-expression

pathways (Ahi *et al*, 2022; Kurko *et al*, 2020; Verta *et al*, 2020; Verta *et al*, 2024) and broad expression of *six6*

during early development (Moustakas-Verho *et al*, 2020). Finally, a recent study showed both loci are linked to

early life survival in juvenile Atlantic salmon, albeit via parental genetic effects, further supporting broad

functional roles across life stages (Aykanat *et al*, 2024). This collective evidence indicates that these loci are

broad-scale modulators of salmon phenotype. While many of these genetic effects have been demonstrated

for traits at the juvenile stage, including early male maturation, metabolism, and behavior, any concordance

with morphological variation, if any, is not known, but may be important to identify the architecture of

personality traits, underlying genetic basis, and evolutionary constrains (Bangura *et al*, 2022, Kern et al, 2016).

To characterize the morphological variation associated with these locally adaptive genomic regions, we collected

morphometric data form more than 700 juvenile immature Atlantic salmon at time when male parr maturation

may take place (Debes *et al*, 2021). The fish was reared in common garden conditions and the association

between morphology, and *six6* and *vgll3* genotype was estimated with a mixed-effect model that also controls for

confounding length and condition factor effects. We identified a suite of morphological characters correlated to

the genetic variation in both loci. Our study extends the broad pleiotropic effect of *vgll3* and *six6* loci to functional

morphological traits, including phenotypes that are likely important targets of natural selection.

## Materials and Methods

*Fish breeding and, rearing and common garden conditions*

Details of the breeding and rearing of individuals from the same cohort were provided in Debes *et al* (2021),

using a similar experimental procedure and facilities, except that the fish in the present study were reared in

approximately 2°C cooler water temperature (see, Debes *et al* (2019) for the details of the rearing

temperature). Briefly, the parents of the individuals used were from Baltic lineage Atlantic salmon from a

broodstock maintained by the Natural Resources Institute Finland (Laukaa, Finland) that were initially

originated from the Neva River in Russia (Gulf of Finland in the Eastern Baltic Sea) and has been used for

juvenile salmon supplementation in the Kymi river in South-Eastern Finland (Asheim *et al*, 2023). Parents were

crossed using a biallelic cross design where 44 parents were crossed in a 11 two by two factorials design, in

which, each factorial unit composed of a pair of males and females which are homozygote in early (E) and late

(L) maturing alleles in the *vgll3* locus (i.e. *vgll3*$^{EE}$ and *vgll3*$^{LL}$ genotypes, Supplementary Table 1), with one type

of genotype within each of the full-sib families with EE, EL, LE, and LL genotypes in the *vgll3* locus

(Supplementary Table 1). *Six6* genotype distribution was not considered during the breeding design, but was

nevertheless segregating in a number of families (Supplementary Table 1). Crosses were performed in

November 2017, and offspring were reared in the University of Helsinki fish facilities, first in replicated family-

specific compartments in vertical incubators until the swim-up stage. Subsequently offspring of all families

were mixed in eight tanks with roughly balanced proportions of families per tank (Supplementary Table 2). In

August 2018, fish were pit-tagged when and fin-clipped for later genotyping. Genotyping and genotypic sex of

the parent and offspring were performed by genotyping-by-sequencing using 177 nucleotide polymorphisms

(SNPs) which included *vgll3* and *six6* SNPs, which were also used for parentage assignment (see, Debes *et al,*

2021, for details). The rearing temperature of the fish in this study followed a seasonal cycle (range:  4.1–

16.0°C) and the natural light regime of the latitude (61.054° N) as described in Debes *et al* (2019). In August

2018, the fish were individually PIT-tagged and a fin clip sample was taken to identify the families via genetic

pedigree reconstruction. Fish were fed ad libitum except for a short period of restricted feeding for half of the

tanks, which included ad libitum feeding for two days and then no feeding for five days, each week for a five-

week period (Debes *et al*, 2021).

*Landmarking and trait measurements*

Morphological data were collected as part of a larger effort in December 2018. Fish of both sexes were

euthanized using an overdose of tricaine methanesulphonate bath, and stored briefly on ice until imaging was

performed the same day. Each fish was sliced ventrally to collect data on gonad development for another

study. The fish was placed on graphical paper, and the body image was taken using a digital camera (Canon

PowerShot S110) mounted on a tripod (Hama Star 61), with aperture-priority exposure at F:8, with autofocus

option (Supplementary Figure 1). Cross sections were obtained by slicing the fish at the posterior end of the

dorsal fin (hereafter termed as large cross section) and adipose fins (hereafter termed  as small cross section),

respectively. These were placed on a transparent sheet and digitized using a scanner (Epson XP-305) with a

modified lid to accommodate the depth of the cross sections (Supplementary Figure 1).

Landmarks were placed on the whole body and the two cross sections in a randomized order using imglab

software (King, 2009), by two of the co-authors (LG for landmarked body planes, and SC for landmarked cross

sections) who were blind to any information related to the genotypes or sex of the fish (Supplementary Figure

2). Landmarks were not placed around the gut region due to the previous ventral slicing. Some landmarks were

solely used in the Procrustes analysis to quantify shape, while others were used to quantify specific trait values

by calculating the distance between landmarks (Supplementary Figure 2, Table 1). Traits that were measured

twice along the transverse plane (both the left and right sides, Table 1) were averaged prior to analysis. All

measurements were reported in centimeters (cm).

*Data filtering*

At the time of sampling, fish exhibited a bimodal length distribution which is often observed within juvenile

salmon from the same cohort (Hoar, 1988). At the time of sampling a majority of fish used in the analysis (i.e.

upper mode of size distribution, see below) had a complete smolt appearance (base on the disappearance of

parr marks in the skin and silvery coloration, e.g., Debes *et al*, 2020), or were at late stages of smoltification

(faint parr marks with close to silvery skin coloration). We excluded a small number of fish in the lower mode to

avoid potential interactions associated with this size bimodality (fish fork length < 9 cm, N=69, Supplementary

Figure 3). In addition, fish with missing *vgll3* and *six6* genotypes, and sex (N=2, N=12, and N=4, respectively),

and a small number of precociously matured males (N=19) were excluded from the analysis. Missing landmarks

(0.02% of the total landmarks) were imputed using the *estimate.missing* function in the *geomorph* package

(version 4.0.6, Baken *et al*, 2021). A landmarked image was excluded from the analysis if more than one

landmark was identified as an outlier in the Procrustes analysis (i.e. > 4 sd away from the mean). These

landmarks were not corrected manually, to avoid introducing among technician variation.  After data filtering,

821 individuals remained in the dataset, of which 731, 737, and 772 individuals were in the small cross section,

large cross section, and whole body datasets, respectively (Supplementary Table 1). Within each dataset 16, 18,

and 712 individuals were landmarked more than once to estimate repeatability. Landmarking for the first 30

and 50 images per researcher for body and cross sections, respectively, were discarded as training sets.

*Trait measurements*

We quantified variation in 21 traits across various body planes with potential functional significance using the

landmark positions (Supplementary Figure 2) and as formulated in Table 1.

*Rayed fins and caudal peduncle length:* We measured the length of dorsal, anal, and tail fins, between interior

and exterior insertion points, as well as the caudal peduncle (Table 1, Supplementary Figure 2). The variation in

these traits may be associated with habitat use, feeding, and swimming performance in salmonids and other

fishes (e.g., Jonsson and Jonsson, 2011; Keeley *et al*, 2005). Increased dorsal and pectoral fin sizes in fish are

likely important for increased maneuverability and stability, and likely important adaptations for faster flowing

streams (Webb, 1984). Specifically, large paired fins (such as pectoral and pelvic fins) in Atlantic salmon are

associated with populations that occupy faster flowing and colder streams than slower flowing and warmer

rivers, an adaptation attributed to providing effective stability under fast flowing, and turbulent currents

(Riddell and Leggett, 1981; Webb, 1988). In Coho salmon, juveniles of interior populations with longer

migration distances appear to have smaller fins (Taylor and McPhail, 1985), which may help to reduce drag in

long migrations (Varian and Nichols, 2010).

*Eye diameter and position, and snout length*: The size of the eye was approximated from the diameter, and the

snout length was measured as the distance between the center of the eye and the anterior extreme of the

head (Table 1, Supplementary Figure 2). These traits are morphological features associated with trophic

specialization and habitat use across fish taxa (Elmer *et al*, 2010; Peris Tamayo *et al*, 2020; Snorrason, 1994).

*Aerobic muscle:* We approximate the area of aerobic muscle fibers (also called slow-twitch muscle, or red

muscle) by measuring the landmark distances in the cross sections from the lateral line towards the transverse

sections, and by measuring the extent of the red muscle along the lateral line (Table 1, Supplementary Figure

2). Aerobic muscle provides the energy required during steady, cruise swimming, and helps with thermal

acclimation in fishes (Altringham and Ellerby, 1999; Shuman and Coughlin, 2018). Recently, it has been shown

that both *vgll3* and *six6* loci are associated with whole body aerobic activity (Prokkola *et al*, 2022), which is a

critical physiological process in the salmonid life cycle (Eliason and Farrell, 2015), which may indicate that

aerobic muscle mass, as a proxy for aerobic activity, may be associated with these loci.

*Cross sectional centroid size and diameter, and vertebral thickness*: We measured cross sectional centroid size

and horizontal diameter from both large and small cross section images (Table 1, Supplementary Figure 2), as

an approximation for fusiform body shape. A slimmer and more torpedo-shaped body provides improved

energy efficiency during cruising and in high flow streams by reducing water drag, and it is an important life

history adaptation among fishes (Crossin *et al*, 2004; Jonsson and Jonsson, 2011a; Taylor and McPhail, 1985;

Webb, 1988). In contrast, a robust cross section facilitates burst swimming, which is advantageous for prey

avoidance and food acquisition (Taylor and McPhail, 1985).

We also quantified vertebral width as a proxy for cone diameter, to approximate vertebra's flexibility/stiffness

during the lateral bending movement in the undulated motion, which helps to maintain steady swimming at

the expense of maneuvering capabilities (Baxter *et al*, 2022; Donatelli *et al*, 2021; Long and Nipper, 1996).

*Head and body size:* We quantified head length and width (Table 1, Supplementary Figure 2), which are

important determinants of ecotypic divergence via resource and habitat use in fishes (Peris Tamayo *et al*, 2020;

Simonsen *et al*, 2017).  A larger head and larger snout length may be associated with a feeding regime towards

piscivory (Keeley *et al*, 2005), or generally speaking, consuming larger prey items. We also measured head and

body centroid sizes as an approximation for the area of the morphometric planes, which is the square root of

the sum of squared distances of all the landmarks used in the Procrustes analysis from the point of center of

gravity (Klingenberg, 2016).  The Atlantic salmon has a smaller head-body ratio compared to the brown trout

(Jonsson and Jonsson 2011), which likely makes salmon more efficient swimmers both in strong freshwater

currents and during longer marine migrations.

*Statistical analysis*

Our analysis framework comprised a series of analyses whereby Procrustes shape variables, and morphological

traits were used as response variables. These analyses were conducted independently for the body, small cross

section, and large cross section. In all analyses, length (centered to zero and scaled to have unit standard

deviation), body condition (Fulton's condition factor, scaled), sex, and food treatment were used as covariates.

In all models, we implemented genetic variation as additive, where *vgll3* and *six6* genetic effects are coded as

0, 1, and 2 for EE, EL, and LL genotypes respectively (E and L refer to the alleles associated with early and late

maturation, respectively), so that a unit change in genetic effect represents an additive allelic substitution

effect from E to L. Unless otherwise noted, all statistical analyses were performed in R software version 4.4.1 (R

Core Team, 2019).

We performed multivariate generalized Procrustes analysis to quantify shape variation (Gower, 1975), using

the default settings in the *gpagen* function in the geomorph package (version 4.0.8, Baken *et al*, 2021). This

function outputs Procrustes shape variables, centered on the mean value for each landmark and normalized for

centroid size. Centroid size is the square root of the sum of squared distances of landmarks from the center of

gravity (i.e., centroid) and it is highly correlated to the size of the fish.

Procrustes shape data was then used to perform a Procrustes linear regression analysis using the *procD.lm*

function with default settings (i.e. residual randomization was used for significance testing and type III marginal

sums of squares for cross products computations). The model parameters were as follows:

$Y = \mu + L + CF + Feed + Tank + Sex + Vgll3 + Six6 + Fam + \varepsilon_{error}$

Where Y is the multivariate data frame consisting of x and y coordinates for Procrustes shape coordinates of

landmarks, $\mu$ is the intercept, L is the scaled fork length, CF is the scaled Fulton's condition factor, and Feed is

the factor associated with feed treatment (ad-libitum vs restricted feed). Other terms are the rearing tank ID

that the fish were reared after the swim-up stage (Tank), sex (Sex), family (Fam), and the additive genetic

effects for *vgll3* and *six6*, respectively. Since the *procD.lm* function does not accommodate random terms, we

included tank and family effects as fixed terms. Trait values were averaged prior to the analyses when an

individual had replicate measurements. We reported multivariate analysis of variance Pillai's trace statistics,

estimated using the *manova.update* function in the RRPP package (version 2.0.3, Collyer *et al*, 2018), and plot

individual landmark differences between loci using the *plotRefToTarget* in the geomorph package.

To decompose the variation associated with trait values and estimate parameter coefficients, we employed a

linear mixed effect (animal) model using the *relmatLmer* function in lme4qtl (version 0.2.2, Ziyatdinov *et al*,

2018), which is an implementation of the *lmer* function in the lme4 package (Bates, 2010), and can fit the

animal model (i.e., accommodate the additive relatedness matrix). When estimating fixed coefficients, the

model was fitted with maximum likelihood using the *REML = FALSE* option in the *relmatLmer* function. The

model structure was as follows:

$Y = \mu + L + CF + Feed + Sex + Vgll3 + Six6 + \varepsilon_{ID} + \varepsilon_{tank} + \varepsilon_{animal} + \varepsilon_{error}$

Fixed effect terms are as described above except that the fixed tank and family terms in the previous model

were replaced with random tank and animal terms, respectively. Random terms thus included, $\varepsilon_{ID}$ which are

the individual effects, $\varepsilon_{tank}$ are the effects associated with the tank effects, and $\varepsilon_{error}$ is the measurement error.

$\varepsilon_{animal}$ is the random term associated with the background genetic effects after controlling for *vgll3* and *six6*

fixed effects. The relatedness matrix was obtained using the pedigree data in Debes *et al* (2019) using the *getA*

function in the pedigreem package (version 0.3.4, Vazquez *et al*, 2010). Since we used repeated measurements

in the analysis, $\varepsilon_{ID}$ reflects the error term, and $\varepsilon_{error}$ is the residuals that quantifies deviations between repeated

samples (i.e. measurement error).

The significance of the fixed effects terms was estimated using the *as_lmerModLmerTest* function in the

lmerTest package (version 1.1-3, Kuznetsova *et al*, 2017) in R using type 3 F test with Satterthwaite's method to

estimate the effective degrees of freedom.

We then estimated the additive genetic variation, the contribution of *vgll3* and *six6* loci to the additive genetic

variation, evolvability, and repeatability. For these, we employed three models similar to above, but the first

model (Model$_{VA.full}$) did not contain *vgll3* or *six6* to estimate total additive genetic variation, and was fitted

using the *REML = TRUE* option in the *relmatLmer* function to obtain unbiased estimates of variance

components. As such, the variance of $\varepsilon_{animal}$ ($V_{animal}$) in the Model$_{VA.full}$ is the total additive genetic variation ($V_A$).

Percent contribution of *vgll3* and *six6* are calculated as%$V_{A.vgll3}$ = 100 * ( ( 2 x L$_{vgll3}$ x E$_{vgll3}$ x ($\beta_{vgll3}^2$ − se($\beta$)$_{vgll3}^2$) /

$V_{animal}$ ), and %$V_{A.six6}$ = 100 * ( ( 2 x L$_{six6}$ x E$_{six6}$ x ($\beta_{six6}^2$ − se($\beta$)$_{six6}^2$) ) ) / $V_{animal}$ ), as in (Luo *et al*, 2003), where L$_{vgll3}$,

E$_{vgll3}$, L$_{six6}$ and E$_{six6}$ denotes frequency of L and E alleles in *vgll3* and *six6* loci, respectively, $\beta_{vgll3}$ and $\beta_{six6}$ denotes

*six6* and *vgll3* E to L substitution effects, respectively, and se denotes standard error of the estimate. Percent

repeatability for each model is calculated as R = $V_{ID}$ / ($V_{ID}$ + $V_{error}$) * 100, and heritability values were calculated

as $h^2$ = $V_{animal}$ / ($V_{animal}$ + $V_{tank}$ + $V_{ID}$), where $V_{tank}$, $V_{ID}$, and $V_{error}$ are the model variance estimates associated with

tank ($\varepsilon_{tank}$), individual ($\varepsilon_{ID}$), and error ($\varepsilon_{error}$) terms in Model$_{VA.full}$. Similarly, evolvability ($I_A$) was calculated as $V_A$ /

$\bar{y}^2$ using Model$_{VA.full}$, where $\bar{y}$ is the mean trait value (Hansen *et al*, 2011; Houle, 1992). Evolvabilities are

interpreted as expected proportional responses in the trait mean under unit selection (i.e. the selection on

fitness itself; Hansen *et al*, 2003; Hansen *et al*, 2011). For all variance components, 95% confidence intervals

were estimated by obtaining the posterior distributions of random effect terms in the model (Model$_{VA.full}$),

using the *sims* function in the arm package(version 1.1.4-4, Gelman and Su, 2020) in R, using 10000

permutations. The simulations generate 10000 estimates for random effect terms, which are then used to

calculate the variance components as above, with 250[th] and 9750[th] ranked estimates recorded as 95% CIs. We

did not control for the effect of dominance in the analyses, which may have inflated the estimates of additive

genetic variance (Lynch and Walsh, 1998).

## Results:

*Shape analysis:*

The multivariate analyses using Procrustes shape variation as the response showed that genetic variation in *six6*, but not in *vgll3*, is associated with shape change in the body plane (Table 2). The change in the body plane appears to be related to changes in the head region associated with *six6*, whereby a substitution from *six6*$^E$ to *six6*$^L$ elongates the head and slightly bends up the lateral body positioning at ends of the lateral line. In contrast, *vgll3*$^E$ to *vgll3*$^L$ substitution reduces the head area relative to the body area and pushes the anterior anal fin landmark toward the posterior end of the body plane, albeit the effect is statistically insignificant (Figure 1a). Genetic variation in *vgll3* and *six6* are not associated with overall shape changes in small and large cross sections. Although not statistically significant, a *vgll3*$^E$ to *vgll3*$^L$ substitution was linked with a narrower horizontal width, a smaller aerobic muscle area in the small cross section (Figure 1b), and a narrower aerobic muscle extension perpendicular to the lateral line in the large cross section (Figure 1c). These results are concordant with the trait-based analysis (see below). Overall, family, tank, feed restriction, and length were important determinants of shape variation in all planes, and condition factor was also significant in the analysis of the body plane (Table 2). Sex was an insignificant determinant of shape in all three body planes (Table 2).

*Trait analysis:* After correcting for body length, condition factor, feed restriction, and sex, nine out of 21 morphological traits were significantly associated with the *vgll3* locus, while four were associated with *six6* locus (Table 1, Figure 2. See also Supplementary Table 3 for *t*-values and effective degrees of freedoms of terms). In general, the *vgll3*$^E$ to *vgll3*$^L$ substitution effect decreased trait values (Table 1, Figure 2) with changes in trait values between *vgll3*$^{EE}$ and *vgll3*$^{LL}$ genotypes being > 1%, and up to 3% of the mean trait values (Figure 2). Notably, the aerobic muscle depth in the small cross section was reduced by 3.45% ($p< 0.001$) and the aerobic muscle vertical extension in the large cross section was reduced by 2.72% ($p = 0.006$) by the *vgll3*$^E$ to *vgll3*$^L$ allelic substitution (Table 1, Figure 2). The *vgll3* allelic substitution effect associated with lateral red muscle length was statistically significant when three aerobic muscles trait values were combined into one composite value ($p< 0.001$, Supplementary Table 4). The *vgll3*$^L$ allele also decreased vertebral thickness in both small (1.37% decrease, $p = 0.009$) and large (1.21% decrease, $p = 0.032$) cross sections (Table 1, Figure 2). The *vgll3*$^E$ to *vgll3*$^L$ substitution effect significantly reduced the small cross section diameter by 1.43% ($p = 0.012$), whereas the small section centroid size was reduced by 0.86% ($p = 0.012$, Table 1, Figure 2). Reduced eye diameter, snout length, and anal fin length were also associated with the *vgll3*$^L$ allele (Table 1, Figure 2). The trait values for all ray fins (dorsal, anal, pectoral, and tail fins) was trending toward a smaller size in relation to the late maturing genotype (Table 1, Figure 2), and their cumulative effect, i.e., effect size of all trait values

combined, was statistically significant ($p$ = 0.022, Supplementary Table 4). We observed that in some traits the

variation explained by $vgll3^E$ to $vgll3^L$ substitution was up to 50% of the variation explained by length (a factor

with strong, positive allometric effect on other trait values, see Table 1), suggesting that this variation is

biologically important (Supplementary Figure 4).

Most differences explained by variation in the $six6$ genomic region were related to changes in head and body

size. The $six6^E$ to $six6^L$ substitution was associated with a 0.17% decrease in body centroid ($t$ = -3.70, $p$< 0.001)

and a 0.47% increase in head centroid size ($t$ = 2.68, $p$ = 0.008, Table 1, Figure 2). The snout length, measured

as the distance between the center of the eye and the maximum anterior point in the body, was elongated by

0.73% in fish with an additional $six6^L$ allele ($t$ = 3.05, $p$ = 0.002, Table 1, Figure 2). Head length appears to be a

significant determinant of changes in head structure, the effect of an additional $six6^L$ allele explained a 0.33%

increase in average trait value ($t$ = 2.15, $p$ = 0.032, Table 1, Figure 2).

Sex effects typically ranged from smaller and up to the same order of magnitude as the allele substitution

effects in the life-history loci (Table 1). Notably, the snout, pectoral fin, and anal fin were significantly larger in

males compared to females ($p$ = 0.002, $p$ = 0.007, and $p$ = 0.021, respectively. Table 1). In addition, the head

centroid was larger in males compared to females ($p$ = 0.040, Table 1). As expected, length was a strong

predictor of all traits, and condition factor was also positively associated with most traits (Table 1). Feed

restriction was notably associated with a decrease in fin trait values, such as the dorsal fin ($p$ < 0.001), anal fin

($p$ = 0.040), and tail fin ($p$ = 0.003, Table 1). Interestingly, feed restriction substantially reduced the centroid size

($p$ = 0.011), cross section diameter ($p$ = 0.003), and the size of the aerobic muscle ($p$ = 0.005) in the small cross

section, but not in the large cross section (Table 1).

As length is a strong predictor of both trait values and overall body shape, a spurious association between traits

and the life-history loci may be expected if these loci affect trait values via length as a mediator. To test this,

we constructed a model to include the interaction between length and life-history loci. Across 21 traits, we did

not observe a meaningful difference in main genetic effects, nor were interaction term $t$-values substantial

(Supplementary Figure 5), suggesting that length-by-genotype interaction did not drive the estimated relations.

*Heritability, evolvability, and additive genetic* variation: Trait heritabilities were low to moderate in most cases,

with a mean heritability of 0.19 across all 21 traits (Table 3). Length of tail fin and body centroid size exhibited

the highest heritability with 0.50 (95% CI 0.47-0.54) and 0.47 (95% CI 0.43-0.50), followed by the pectoral fin

with 0.41 (95% CI 0.36-0.45). The repeatability of trait measurements was moderate (mean repeatability

73.0%, Table 3). The relatively low repeatability than expected (Takacs *et al*, 2016) is elevated likely due to low

intra-population variation obtained in the common garden setting relative to most morphometrics studies that

focuses on larger biological variation, e.g. between ecomorphs of conspecifics or across species, but complex

trait measurements combined with suboptimal fish imaging using gutted individuals, and measurer's expertise

may also contribute to such variation (Vrdoljak *et al,* 2020, Takacs *et al*, 2016).

Evolvabilities (mean-scaled additive genetic variances, denoted as $I_A$) differed greatly among traits (Table 3).

Interestingly, body centroid size had the lowest evolvability (<0.001), meaning that even under unit directional

selection ( the strength of selection on fitness itself, Hansen *et al,* 2003), the expected increase in this trait

value would be less than 0.1%. On the other hand, large cross section aerobic muscle had an evolvability as

high as 0.72, meaning that the expected increase in this trait under strong selection could be as high as 71% of

the mean trait value. In general, out of 21 traits measured, aerobic muscle and vertebrae thickness traits had

higher $I_A$, while centroid sizes and cross section diameters had lower $I_A$ (Table 3).

Additive genetic variation explained by the *vgll3* locus was non-zero for 12 traits (Table 3). Consistent with the

fixed effect parameterization of the genotype, a substantial portion of the additive genetic variation in vertebra

thickness was explained by *vgll3*, with 30.7% and 13.0% for small and large cross sections, respectively (Table

3). A significant portion of the additive genetic variation in small section aerobic muscle, cross section

diameter, and body centroid was also explained by the *vgll3* locus, with 22.4%, 12.7, and 6.3% for each trait,

respectively (Table 3). The contribution of *six6* locus to additive genetic variation associated included snout

length (9.4%), and head and body centroid size (5.2% and 4.2%, respectively, Table 3) consistent with the locus

effect size.

## Discussion

Intraspecific differences in functionally important morphological traits may provide important insights into the

mechanistic basis of adaptation. We detected several morphological traits in juvenile Atlantic salmon that were

linked with variation in genomic regions previously reported to be associated with life-history diversity and

known to be under local selection. The genetic polymorphism in the *vgll3* locus appears to shape a suite of

traits associated with swimming performance, while the genetic polymorphism in the *six6* locus shapes head

and body size, and head features such as snout length, which potentially affects resource acquisition

efficiencies for different foraging strategies. On the other hand, these genomic regions have only a small effect

on the general shape of the morphological planes. The only statistically significant effect was detected in the

body plane plan that was associated with the *six6* loci (Table 2), for which the most striking shape change at the

vector level elongates the head length (in relation to *six6*[E] to *six6*[L] substitution, Figure 1), which  is concordant

with change in the head/body proportions observed in the univariate trait analysis. Our results support these

genomic regions having broad phenotypic effects, which may have emerged from either a form of pleiotropy or

linkage disequilibrium. Our design does not allow us to categorize the pleiotropic effects further, whether it is

based on an effect of single causal variant (biological pleiotropy), or based on effects associated with different

phenotypes along a biochemical pathway (mediated pleiotropy), or as a result of linkage disequilibrium

between two independent causative loci (spurious pleiotropy, e.g., Solovieff *et al*, 2013). Yet, the strong

genetic interdependence among traits associated with *six6* and *vgll3* may have a developmental origin, which is

supported by the patterns of gene expression in these genomic regions during development (Moustakas-Verho

*et al*, 2020; Kurko *et al*, 2020), and their molecular regulatory functions, (Ahi *et al*, 2022; Kurko *et al*, 2020;

Mobley *et al*, 2021;; Verta *et al*, 2020; Verta *et al*, 2024). For example, being a co-transcription factor binding to

the TEAD protein, *vgll3* is likely an important regulator of the hippo pathway (Kurko *et al*, 2020) which is

involves in organ growth via controlling cell growth (Huang *et al,* 2005). Likewise, allele specific *vgll3* regulation

of pubertal development, interact with signaling pathways with diverse molecular functions, such as TGF- β

signaling, and wnt pathway which have developmental functions during body axes formation, as well as cell

fate commitment (Xue *et al,* 2025; Wu and Hill, 2009).

The suite of morphological traits associated with *vgll3* variation reflects changes along a functional axis that

facilitates energy-efficient cruising (*vgll3*$^L$) versus maneuvering and acceleration (*vgll3*$^E$). It appears that that

individuals with *vgll3*$^E$ genotype have a robust build to facilitate maneuvering and acceleration, and larger

aerobic activity (i.e. increased red muscle area, see also below), while individuals with *vgll3*$^L$ allele have a

slender body build for energy efficiency swimming (i.e. smaller cross section diameter, and smaller ray fins, see

also below). Energy efficient cruising is an important trait especially in the marine environment, while better

maneuvering and acceleration help coping with biotic (i.e. competition and predation risk) and abiotic (flow

regime) factors in the freshwater habitat. The larger aerobic muscle area associated with the *vgll3*$^E$ allele could

indicate a higher potential for aerobic muscle activity. This result is concordant with (Prokkola *et al*, 2022) who

demonstrated that juvenile Atlantic salmon with the early maturing genotype have a higher maximum

metabolic rate (MMR) and aerobic scope (AS). Aerobic locomotion is expected to be central for Atlantic salmon

during the marine phase, where salmon sub-adults migrate and continuously forage for food. Our results

suggest salmon with the early genotype are more likely to maintain sustained swimming and foraging activities,

beneficial during marine exploration, which may help to facilitate earlier maturation, with poor performers

having to delay reproduction until they can utilize sufficient energy for return migration. On the other hand,

having a slender body and energy-efficient cruising may be more advantageous in later years, since individuals

with more *vgll3*$^L$ alleles appears to perfrom better than indiviauls with *vgll3*$^E$ allale at older sea-age groups (i.e.

Barson *et al*, 2013). However, it is unclear if the genetic association between life-history loci and morphological

traits persists between life stages, or across different environments. The parr-smolt transformation

(smoltification) includes morphological changes that prepare juvenile Atlantic salmon for the marine

environment (Jonsson and Jonsson, 2011), hence phenotype-genotype association might differ either as a

result of smoltification or change in the environment (Aguirre *et al*, 2014). However, since individuals used in

this study have a complete or nearly complete smolt phenotype, conclusions may be more relevant to the

marine than to the freshwater life stage.

Increased vertebral thickness associated with the *vgll3*$^E$ allele could provide stiffness for energy efficient

undulation movement during steady swimming (Donatelli *et al*, 2021), though it may come at the expense of

body flexibility that may facilitate piscivorous prey capture. A larger aerobic muscle area and thicker vertebra

may eventually facilitate earlier maturation via more energy efficient resource acquisition at sea, suggesting

that local adaptation for optimal age structure could be partly mediated by conditions at sea. Hence, while our

results do not explain the causation of early male maturation in freshwater, we cannot explicitly rule out that

the link between *vgll3* and age at maturity is mediated via its effect on body shape. Sex was also an important

factor shaping the trait variation. Intriguingly, significant changes in shape associated with sex and *vgll3*

genotype was also in concordant with respect to maturation timing. As such, males, which on average mature

earlier than females, and the *vgll3* allele associated with early maturation had similar direction of change in

these morphological traits.

Larger fin sizes in fish increase maneuverability and stability (Webb, 1984), and correspondingly in Atlantic

salmon, larger fins occur in populations that occupy faster flowing streams (Riddell and Leggett, 1981). Overall,

we found that the ray-fins of fish were consistently larger for the *vgll3*$^E$ allele than the *vgll3*$^L$ allele

(Supplementary Table 4). However, this improvement for maneuverability and stability may come at the

expense of efficiency in longer marine migrations, as a result of increased drag. For example, in coho salmon,

juveniles of inland populations with longer migration distances have smaller fins (Taylor and McPhail, 1985),

which may help reduce drag in long migrations (Dynes *et al*, 2005; Varian and Nichols, 2010). A larger cross

section diameter and larger fin lengths associated with the *vgll3*$^E$ allele likely facilitate maneuvering at the

expense of efficient cruising, which is contrast with the *vgll3*$^E$ effect that increases aerobic muscle area and

vertebral thickness. Similarly, the slender body (a smaller cross section diameter) associated with *vgll3*$^L$ is

consistent with minimizing drag for a cruising body-form (Webb, 1988), advantageous for a multi-sea winter

strategy that migrates longer distances in the ocean (e.g., O'Sullivan *et al*, 2022). On the other hand, the

direction of effect on fin size associated with *vgll3*$^E$ *to vgll3*$^L$ substitution is concordant with the effect of the

restricted-food regime, suggesting that the allelic substitution effect may be correlated to food limitation

conditions. Thus, overall, the direction of *vgll3* genetic effects is not functionally consistent in relation to

proposed ecological roles of these morphological traits.

The changes associated with the genetic variation in the *six6* locus that are affecting body and head sizes are

linked to ecological processes related to resource acquisition mechanisms. A larger head (and a smaller body)

associated with the *six6*$^L$ allele and larger snout length may be associated with a feeding regime towards

piscivory (Keeley *et al*, 2005), or generally consuming larger prey items. This interpretation corroborates the

study by Aykanat *et al* (2020), in which fish with the *six6*$^{LL}$ genotype were shown to be more efficient in

resource acquisition than fish with the *six6*$^{EE}$ genotype.

Evolvabilities ($I_A$) ranged from normal to very high compared to a recent meta-analysis (Hansen *et al*, 2011).

The median evolvability for length traits is 0.001 (or 0.1%, Hansen and Pélabon, 2021; Hansen *et al*, 2011).

Normal evolvability values were found for centroid sizes and cross section diameters, while aerobic muscle and

vertebrae thickness had high evolvabilities, suggesting that these traits could evolve rapidly in response to

selection induced by environmental change. Interestingly, a substantial portion of the additive genetic variation

associated with some traits, such as aerobic muscle area and vertebrae thickness, was explained by the *vgll3*

locus, suggesting the locus' importance in shaping the additive genetic landscape.

Taken together, *vgll3* and *six6* loci have complex associations with morphology. These do not consistently align

with adaptive differences between early- and late-maturation genotypes associated with marine migration, but

could be important e.g., during spawning where higher maneuverability of smaller males may be beneficial for

reproductive success, or in early life stages. On the other hand, it is also plausible that these changes are

genetically correlated as part of broad changes during development, and some of which have potentially

maladaptive consequences. When traits are genetically correlated, the cumulative adaptive value of trait

values cannot be optimal (Schluter, 1996), since breaking suboptimal trait combinations that are genetically

correlated may be challenging for evolution (Blows, 2007). Finally, it is challenging to assess the biological

significance of the scale of trait variation between loci. However, effect size appears to be large for some trait,

i.e., as inferred by up to 50% effect size of allelic substitution effect relative to the effect of length

(Supplementary Figure 4), indicating that the genetic variation produces some static allometry between

individuals the same cohort, which may have functional and adaptive basis (Pelabon *et al*, 2014).

It is important to recognize that extrapolation to the marine stage of Atlantic salmon based on juvenile

morphology may have limitations. It is unknown if the genotype-associated morphological variation is

correlated between juvenile and adult stages. While there may be genetic correlation between freshwater and

marine environments (Fleming *et al*, 1994; Morinville and Rasmussen, 2006; Taylor and McPhail, 1985), the

469 morphological changes associated with adaptation to the marine environment, i.e., the parr-smolt

470 transformation (smoltification), may break apart trait correlations, similar to how metamorphosis takes place

471 in many animal taxa (e.g., Aguirre *et al*, 2014). Since the data in this study were composed of individuals that

472 had completed the parr-smolt transformation, or are close to completion, conclusions are likely more relevant

473 to marine stages than the freshwater stages. However, any adaptive inference associated with a specific life

474 stage should be taken with caution until observed patterns are further tested and verified at different life

475 stages. Finally, while it is reasonable to hypothesize that the observed *vgll3* and *six6* associated morphological

476 variation is adaptively important, the fitness effects of this variation should be better assessed before any

477 evolutionary conclusions can be made (Kingsolver *et al*, 1995).

478 While modelling non-additive architecture within a locus, such as dominance, overdominance, may be

479 relevant, since the *vgll3* locus exhibits sex dependent dominance (Barson *et al*, 2015), in this study, genetic

480 effects were modelled additively, to avoid conceptual and statistical complexities. However, when genetic

481 effects were modelled non-additively (separately for *vgll3* and *six6*), parsimony was meaningfully improved

482 (i.e., AIC for dominance gene effect model was 2 units smaller than the AIC for additive gene effect model) for

483 four out of 21 traits for the *vgll3* gene, while none of the non-additive models performed better than additive

484 models for the *six6* gene (Supplementary Figure 6a). In all three traits with significant non-additive genetic

485 effect, a dominance model with *vgll3*$^{EL}$ effect resembling *vgll3*$^{LL}$ was supported (Supplementary Figure 6 b-d),

486 suggesting a possible role of dominance shaping some of the trait values.

487 In conclusion, we have detected substantial morphological variation associated with two genomic regions,

488 which are also major determinants of life-history variation and known to be under local selection. These loci

489 appear to influence multiple, strongly correlated traits, suggestive of a broad regulatory role with impacts

490 across the whole life cycle. How these genetic correlations are shaped – via single locus pleiotropy or close

491 physical linkage of multiple genes – remains to be determined.

## 493 Acknowledgment

494 We thank Artur Porto for discussion related to morphometric analyses, and Seija Tillanen, Jacqueline E.

495 Moustakas-Verho, Nikolai Piavchenko for helps with sampling. This work was supported by the Research

496 Council of Finland (grant nos. 328860, 353388, and 325964 to TA, and 314254, 327255 and 342851 to CRP), and

497 the European Research Council under the European Articles Union's Horizon 2020 and Horizon Europe research

498 and innovation programs (grant no. 742312 and 101054307 to CRP). Views and opinions expressed are

499    however those of the author(s) only and do not necessarily reflect those of the European Union or the

500    European Research Council Executive Agency. Neither the European Union nor the granting authority can be

501    held responsible for them. LG was supported by Erasmus+ programme. The authors declare no competing

502    interests.

## 504    Ethical approval

505    The experiments were approved by the Project Authorisation Board (ELLA) on behalf of the Regional

506    Administrative Agency for Southern Finland (ESAVI) under experimental license ESAVI/2778/2018.

## 508    Author contributions

509    TA, VLP, and GHB conceptualized the paper. TA analyzed the data with significant input from GHB and PVD. SJ

510    and LG performed landmarking. AHH and AR prepared and scanned the cross sections. JE, and CRP oversaw the

511    resources and infrastructure. PVD and CRP devised the breeding design and reared the fish. TA and PVD

512    sampled the fish and digitized the images. TA drafted the MS with significant inputs from PVD, VLP, and GHB,

513    and all coauthors contributed to subsequent drafts.

## 515    Conflict of Interest

516    The authors declare no competing interests.

## 518    References:

519    Aguirre JD, Blows MW, Marshall DJ (2014). The genetic covariance between life cycle stages separated by
520    metamorphosis. *Proceedings Biological sciences / The Royal Society* **281**(1788)**:** 20141091.

521
522    Ahi EP, Sinclair-Waters M, Moustakas-Verho J, Jansouz S, Primmer CR (2022). Strong regulatory effects of *vgll3*
523    genotype on reproductive axis gene expression in juvenile male Atlantic salmon. *General and Comparative*
524    *Endocrinology* **325:** 114055.

525
Altringham JD, Ellerby DJ (1999). Fish swimming: Patterns in muscle function. *Journal of Experimental Biology*
**202**(Pt 23)**:** 3397-3403.

Archambeault SL, Bartschi LR, Merminod AD, Peichel CL (2020). Adaptation via pleiotropy and linkage:
Association mapping reveals a complex genetic architecture within the stickleback *Eda* locus. *Evolution Letters*
**4**(4)**:** 282-301.

Asheim ER, Debes PV, House A, Liljestrom P, Niemela PT, Siren JP *et al* (2023). Atlantic salmon (*Salmo salar*) age
at maturity is strongly affected by temperature, population and age-at-maturity genotype. *Conservation*
*Physiology* **11**(1)**:** coac086.

Aykanat T, McLennan D, Metcalfe NB, Prokkola JM (2024). Early survival in Atlantic salmon is associated with
parental genotypes at loci linked to timing of maturation. *Evolution* **78**(8)**:** 1441-1452.

Aykanat T, Ozerov M, Vaha JP, Orell P, Niemela E, Erkinaro J *et al* (2019). Co-inheritance of sea age at maturity
and iteroparity in the Atlantic salmon *vgll3* genomic region. *Journal of Evolutionary Biology* **32**(4)**:** 343-355.

Aykanat T, Rasmussen M, Ozerov M, Niemela E, Paulin L, Vaha JP *et al* (2020). Life-history genomic regions
explain differences in Atlantic salmon marine diet specialization. *Journal of Animal Ecology* **89**(11)**:** 2677-2691.

Ayllon F, Kjaerner-Semb E, Furmanek T, Wennevik V, Solberg MF, Dahle G *et al* (2015). The *vgll3* locus controls
age at maturity in wild and domesticated Atlantic salmon (*Salmo salar* L.) Males. *PLoS Genet* **11**(11)**:** e1005628.

Baken EK, Collyer ML, Kaliontzopoulou A, Adams DC (2021). geomorph v4.0 and gmShiny: Enhanced analytics
and a new graphical interface for a comprehensive morphometric experience. *Methods in Ecology and*
*Evolution* **12**(12)**:** 2355-2363.

Bangura PB, Tiira K, Niemela PT, Erkinaro J, Liljestrom P, Toikkanen A *et al* (2022). Linking *vgll3* genotype and
aggressive behaviour in juvenile Atlantic salmon (*Salmo salar*). *Journal Fish Biology* **100**(5)**:** 1264-1271.

Barrett RD, Hoekstra HE (2011). Molecular spandrels: tests of adaptation at the genetic level. *Nature Reviews*
*Genetics* **12**(11)**:** 767-780.

Barson NJ, Aykanat T, Hindar K, Baranski M, Bolstad GH, Fiske P *et al* (2015). Sex-dependent dominance at a
single locus maintains variation in age at maturity in salmon. *Nature* **528**(7582)**:** 405-408.

Bates DM (2010). lme4: Mixed-effects modeling with R.

Baxter D, Cohen KE, Donatelli CM, Tytell ED (2022). Internal vertebral morphology of bony fishes matches the
mechanical demands of different environments. *Ecology and evolution* **12**(11)**:** e9499.

Blows MW (2007). A tale of two matrices: multivariate approaches in evolutionary biology. *Journal of*
*Evolutionary Biology* **20**(1)**:** 1-8.

Boyle EA, Li YI, Pritchard JK (2017). An expanded view of complex traits: From polygenic to omnigenic. *Cell*
**169**(7)**:** 1177-1186.

Chan YF, Marks ME, Jones FC, Villarreal G, Jr., Shapiro MD, Brady SD *et al* (2010). Adaptive evolution of pelvic
reduction in sticklebacks by recurrent deletion of a *Pitx1* enhancer. *Science* **327**(5963)**:** 302-305.

Collyer ML, Adams DC, Freckleton R (2018). RRPP: An R package for fitting linear models to high-dimensional
data using residual randomization. *Methods in Ecology and Evolution* **9**(7)**:** 1772-1779.

Colosimo PF, Hosemann KE, Balabhadra S, Villarreal G, Jr., Dickson M, Grimwood J *et al* (2005). Widespread
parallel evolution in sticklebacks by repeated fixation of Ectodysplasin alleles. *Science* **307**(5717)**:** 1928-1933.

Crossin GT, Hinch SG, Farrell AP, Higgs DA, Lotto AG, Oakes JD *et al* (2004). Energetics and morphology of
sockeye salmon: effects of upriver migratory distance and elevation. *Journal Fish Biology* **65**(3)**:** 788-810.

Debes PV, Piavchenko N, Erkinaro J, Primmer CR (2020). Genetic growth potential, rather than phenotypic size,
predicts migration phenotype in Atlantic salmon. *Proceedings Biological sciences / The Royal Society* **287**(1931)**:**
20200867.

Debes PV, Piavchenko N, Ruokolainen A, Ovaskainen O, Moustakas-Verho JE, Parre N *et al* (2019). Large single-
locus effects for maturation timing are mediated via condition variation in Atlantic salmon. *bioRxiv***:** 780437
(v780433).

Debes PV, Piavchenko N, Ruokolainen A, Ovaskainen O, Moustakas-Verho JE, Parre N *et al* (2021). Polygenic
and major-locus contributions to sexual maturation timing in Atlantic salmon. *Molecular Ecology* **30**(18)**:** 4505-
4519.

Donatelli CM, Roberts AS, Scott E, DeSmith K, Summers D, Abu-Bader L *et al* (2021). Foretelling the flex-
vertebral shape predicts behavior and ecology of fishes. *Integrative and Comparative Biology* **61**(2)**:** 414-426.

Dynes J, Magnan P, Bernatchez L, Rodríguez MA (2005). Genetic and morphological variation between two
forms of lacustrine brook charr. *Journal Fish Biology* **54**(5)**:** 955-972.

Eliason EJ, Farrell AP (2015). Oxygen uptake in Pacific salmon *Oncorhynchus* spp.: when ecology and physiology
meet. *Journal Fish Biology*.

Elmer KR, Kusche H, Lehtonen TK, Meyer A (2010). Local variation and parallel evolution: morphological and
genetic diversity across a species complex of neotropical crater lake cichlid fishes. *Philosophical transactions of
the Royal Society of London Series B, Biological sciences* **365**(1547)**:** 1763-1782.

Endler JA (2000). Adaptive genetic variation in the wild. In: Mousseau TA, SinerVo B and Endler JA (eds)
*Adaptive Genetic Variation in the Wild*. Oxford University Press.

612
613  Ferry-Graham LA, Bolnick DI, Wainwright PC (2002). Using functional morphology to examine the ecology and
614  evolution of specialization. *Integrative and Comparative Biology* **42**(2)**:** 265-277.

615
616  Fitzgerald DB, Winemiller KO, Sabaj Pérez MH, Sousa LM, Seymour C (2017). Using trophic structure to reveal
617  patterns of trait-based community assembly across niche dimensions. *Functional Ecology* **31**(5)**:** 1135-1144.

618
619  Fleming IA, Jonsson B, Gross MR (1994). Phenotypic divergence of sea-ranched, farmed, and wild salmon.
620  *Canadian Journal of Fisheries and Aquatic Sciences* **51**(12)**:** 2808-2824.

621
622  Gelman A, Su YS (2020). arm: Data analysis using regression and multilevel/hierarchical models. R package
623  version 1.11-2.

624
625  Gower JC (1975). Generalized procrustes analysis. *Psychometrika* **40**(1)**:** 33-51.

626
627  Hansen TF, Pélabon C (2021). Evolvability: A quantitative-genetics perspective. *Annual Review of Ecology,*
628  *Evolution, and Systematics* **52**(1)**:** 153-175.

629
630  Hansen TF, Pelabon C, Armbruster WS, Carlson ML (2003). Evolvability and genetic constraint in *Dalechampia*
631  blossoms: components of variance and measures of evolvability. *Journal of Evolutionary Biology* **16**(4)**:** 754-
632  766.

633
634  Hansen TF, Pélabon C, Houle D (2011). Heritability is not evolvability. *Evolutionary Biology* **38**(3)**:** 258-277.

635
636  Hoar WS (1988). 4 The physiology of smolting salmonids. In: Hoar WS and Randall DJ (eds) *Fish Physiology*.
637  Academic Press. Vol. 11, pp 275-343.

638
639  Houle D (1992). Comparing evolvability and variability of quantitative traits. *Genetics* **130:** 195-204.

640
641  Huang, J, Wu S, Barrera J, Matthews K, Pan D (2005). The Hippo signaling pathway coordinately regulates cell
642  proliferation and apoptosis by inactivating Yorkie, the Drosophila Homolog of YAP. *Cell* **122**, 421-434.

643
644  Jakobson CM, Jarosz DF (2020). What has a century of quantitative genetics taught us about nature's genetic
645  tool kit? *Annual Review of Genetics* **54:** 439-464.

646
Jimenez-Gomez JM, Wallace AD, Maloof JN (2010). Network analysis identifies *ELF3* as a QTL for the shade
avoidance response in Arabidopsis. *PLoS Genetics* **6**(9)**:** e1001100.

Jones, F.C., Grabherr, M.G., Chan, Y.F., Russell, P., Mauceli, E., Johnson, J. et al. (2012). The genomic basis of
adaptive evolution in threespine sticklebacks. Nature 484, 55-61.

Jonsson B, Jonsson N (2011). *Ecology of Atlantic salmon and brown trout : habitat as a template for life*
*histories*. Springer: Dordrecht.

Kawecki TJ, Ebert D (2004). Conceptual issues in local adaptation. *Ecology Letters* **7**(12)**:** 1225-1241.

Keeley ER, Parkinson EA, Taylor EB (2005). Ecotypic differentiation of native rainbow trout (*Oncorhynchus*
*mykiss*) populations from British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences* **62**(7)**:** 1523-
1539.

Kern EMA, Robinson D, Gass E, Godwin J, Langerhans RB (2016). Correlated evolution of personality,
morphology and performance. *Animal Behaviour* **117**, 79-86.

King DE (2009). Dlib-Ml: A machine learning toolkit. *Journal of Machine Learning Research* **10** (Jul), 1755–1758.

Kingsolver JG, Woods HA, Gilchrist G (1995). Traits related to fitness. *Science* **267**(5196)**:** 396.

Klingenberg CP (2016). Size, shape, and form: concepts of allometry in geometric morphometrics. *Development*
*Genes and Evolution* **226**(3)**:** 113-137.

Kurko J, Debes PV, House AH, Aykanat T, Erkinaro J, Primmer CR (2020). Transcription profiles of age-at-
maturity-associated genes suggest cell fate commitment regulation as a key factor in the Atlantic salmon
Maturation Process. *G3* **10**(1)**:** 235-246.

Kuznetsova A, Brockhoff PB, Christensen RHB (2017). lmerTest package: Tests in linear mixed effects models.
*Journal of Statistical Software* **82**(13).

Long JH, Nipper KS (1996). The importance of body stiffness in undulatory propulsion. *American Zoologist*
**36**(6)**:** 678-694.

Lundsgaard-Hansen B, Matthews B, Vonlanthen P, Taverna A, Seehausen O (2013). Adaptive plasticity and
genetic divergence in feeding efficiency during parallel adaptive radiation of whitefish (*Coregonus* spp.). *Journal*
*of Evolutionary Biology* **26**(3)**:** 483-498.

Luo L, Mao Y, Xu S (2003). Correcting the bias in estimation of genetic variances contributed by individual QTL.
*Genetica* **119**(2)**:** 107-113.

Lynch M, Walsh B (1998). *Genetics and analysis of quantitative traits*. Sinauer Associates: Sunderland, MA.

Mackay TF, Stone EA, Ayroles JF (2009). The genetics of quantitative traits: challenges and prospects. *Nature*
*Reviews Genetics* **10**(8)**:** 565-577.

Miettinen A, Dannewitz J, Palm S, Ahi EP, Romakkaniemi A, Vähä V *et al* (2023). Loci associated with
maturation, migration and appetite control are linked with signals of fine-scale local selection in a large Atlantic
salmon population. *bioRxiv***:** 2023.2008.2023.553800.

Mindel BL, Neat FC, Trueman CN, Webb TJ, Blanchard JL (2016). Functional, size and taxonomic diversity of fish
along a depth gradient in the deep sea. *PeerJ* **4:** e2387.

Mobley KB, Aykanat T, Czorlich Y, House A, Kurko J, Miettinen A *et al* (2021). Maturation in Atlantic salmon
(*Salmo salar*, Salmonidae): a synthesis of ecological, genetic, and molecular processes. *Reviews in Fish Biology*
*and Fisheries* **31**(3)**:** 523-571.

Morinville GR, Rasmussen JB (2006). Does life-history variability in salmonids affect habitat use by juveniles? A
comparison among streams open and closed to anadromy. *Journal of Animal Ecology* **75**(3)**:** 693-704.

Morinville GR, Rasmussen JB (2007). Distinguishing between juvenile anadromous and resident brook trout
(*Salvelinus fontinalis*) using morphology. *Environmental Biology of Fishes* **81**(2)**:** 171-184.

Moustakas-Verho JE, Kurko J, House AH, Erkinaro J, Debes P, Primmer CR (2020). Developmental expression
patterns of six6: A gene linked with spawning ecotypes in Atlantic salmon. *Gene Expression Patterns* **38:**
119149.

Nielsen EE, Hemmer-Hansen J, Larsen PF, Bekkevold D (2009). Population genomics of marine fishes:
identifying adaptive variation in space and time. *Molecular Ecology* **18**(15)**:** 3128-3150.

O'Sullivan RJ, Ozerov M, Bolstad GH, Gilbey J, Jacobsen JA, Erkinaro J *et al* (2022). Genetic stock identification
reveals greater use of an oceanic feeding ground around the Faroe Islands by multi-sea winter Atlantic salmon,
with variation in use across reporting groups. *ICES Journal of Marine Science* **79**(9)**:** 2442-2452.

Olson-Manning CF, Wagner MR, Mitchell-Olds T (2012). Adaptive evolution: evaluating empirical support for
theoretical predictions. *Nature Reviews Genetics* **13**(12)**:** 867-877.

Orr HA (2000). Adaptation and the cost of complexity. *Evolution* **54**(1)**:** 13-20.

Otto SP (2004). Two steps forward, one step back: the pleiotropic effects of favoured alleles. *Proceedings*
*Biological sciences / The Royal Society* **271**(1540)**:** 705-714.

Paaby AB, Rockman MV (2013). The many faces of pleiotropy. *Trends in Genetics* **29**(2)**:** 66-73.

Pelabon C, Firmat C, Bolstad GH, Voje KL, Houle D, Cassara J et al (2014). Evolution of morphological allometry.
*Annals of the New York Academy of Sciences* **1320**, 58-75.

Peris Tamayo AM, Devineau O, Praebel K, Kahilainen KK, Ostbye K (2020). A brain and a head for a different
habitat: Size variation in four morphs of Arctic charr (*Salvelinus alpinus* (L.)) in a deep oligotrophic lake. *Ecology*
*and evolution* **10**(20)**:** 11335-11351.

Pritchard VL, Makinen H, Vaha JP, Erkinaro J, Orell P, Primmer CR (2018). Genomic signatures of fine-scale local
selection in Atlantic salmon suggest involvement of sexual maturation, energy homeostasis and immune
defence-related genes. *Molecular Ecology* **27**(11)**:** 2560-2575.

Prokkola JM, Asheim ER, Morozov S, Bangura P, Erkinaro J, Ruokolainen A *et al* (2022). Genetic coupling of life-
history and aerobic performance in Atlantic salmon. *Proceedings Biological sciences / The Royal Society*
**289**(1967)**:** 20212500.

Proulx R, Magnan P (2004). Contribution of phenotypic plasticity and heredity to the trophic polymorphism of
lacustrine brook charr (Salvelinus fontinalis M.). *Evolutionary Ecology Research* **6**(4)**:** 503-522.

R Core Team. (2019). R Foundation for Statistical Computing, Vienna, Austria.

Riddell BE, Leggett WC (1981). Evidence of an adaptive basis for geographic-variation in body morphology and
time of downstream migration of juvenile Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and*
*Aquatic Sciences* **38**(3)**:** 308-320.

Schluter D (1996). Ecological Causes of Adaptive Radiation. *American Naturalist*

Shapiro MD, Marks ME, Peichel CL, Blackman BK, Nereng KS, Jonsson B, *et al* (2004). Genetic and
developmental basis of evolutionary pelvic reduction in threespine sticklebacks. *Nature* **428**, 717-723.

Shuman JL, Coughlin DJ (2018). Red muscle function and thermal acclimation to cold in rainbow smelt, *Osmerus*
*mordax*, and rainbow trout, *Oncorhynchus mykiss*. *Journal of Experimental Zoology Part A: Ecological and*
*Integrative Physiology* **329**(10)**:** 547-556.
Simonsen MK, Siwertsson A, Adams CE, Amundsen PA, Praebel K, Knudsen R (2017). Allometric trajectories of
body and head morphology in three sympatric Arctic charr (*Salvelinus alpinus* (L.)) morphs. *Ecology and*
*evolution* **7**(18)**:** 7277-7289.

Sinclair-Waters M, Odegard J, Korsvoll SA, Moen T, Lien S, Primmer CR *et al* (2020). Beyond large-effect loci:
large-scale GWAS reveals a mixed large-effect and polygenic architecture for age at maturity of Atlantic
salmon. *Genetics Selection Evolution* **52**(1)**:** 9.

Snorrason S (1994). Trophic specialization in Arctic charr *Salvelinus alpinus* (Pisces; Salmonidae): morphological
divergence and ontogenetic niche shifts. *Biological Journal of the Linnean Society* **52**(1)**:** 1-18.

Solovieff N, Cotsapas C, Lee PH, Purcell SM, Smoller JW (2013). Pleiotropy in complex traits: challenges and
strategies. *Nature Reviews Genetics* **14**(7)**:** 483-495.

Svanbäck R, Eklöv P (2003). Morphology dependent foraging efficiency in perch: a trade-off for ecological
specialization? *Oikos* **102**(2)**:** 273-284.

Takacs P, Vital Z, Ferincz A, Staszny A (2016). Repeatability, reproducibility, separative power and subjectivity of
different fish morphometric analysis methods. *PLoS One* **11**, e0157890.

Taylor EB, McPhail JD (1985). Variation in body morphology among British Columbia populations of coho
salmon, *Oncorhynchus kisutch*. *Canadian Journal of Fisheries and Aquatic Sciences* **42**(12)**:** 2020-2028.

Varian A, Nichols KM (2010). Heritability of morphology in brook trout with variable life histories. *PLoS One*
**5**(9)**:** e12950.

Vazquez AI, Bates DM, Rosa GJ, Gianola D, Weigel KA (2010). Technical note: an R package for fitting
generalized linear mixed models in animal breeding. *Journal of Animal Science***88**(2)**:** 497-504.

Verta JP, Debes PV, Piavchenko N, Ruokolainen A, Ovaskainen O, Moustakas-Verho JE *et al* (2020). Cis-
regulatory differences in isoform expression associate with life history strategy variation in Atlantic salmon.
*PLoS Genetics* **16**(9)**:** e1009055.

Verta JP, Moustakas-Verho JE, Donner I, Frapin M, Ruokolainen A, Debes PV *et al* (2024). A complex mechanism
translating variation of a simple genetic architecture into alternative life histories. *Proceedings of the National*
*Academy of Sciences of the USA* **121**(48)**:** e2402386121.

Vrdoljak J, Sanchez KI, Arreola-Ramos R, Diaz Huesa EG, Villagra A, Avila LJ *et al* (2020). Testing repeatability,
measurement error and species differentiation when using geometric morphometrics on complex shapes: a
case study of Patagonian lizards of the genus *Liolaemus* (Squamata: Liolaemini). *Biological Journal of the*
*Linnean Society* **130**, 800-812.
Wagner GP, Altenberg L (1996). Perspective: Complex adaptations and the evolution of evolvability. *Evolution*
**50**(3)**:** 967-976.

Wagner GP, Zhang J (2011). The pleiotropic structure of the genotype-phenotype map: the evolvability of
complex organisms. *Nature Reviews Genetics* **12**(3)**:** 204-213.

Webb PW (1984). Body form, locomotion and foraging in aquatic vertebrates. *American Zoologist* **24**(1)**:** 107-
120.

Webb PW (1988). Simple physical principles and vertebrate aquatic locomotion. *American Zoologist* **28**(2)**:** 709-
725.

Weith M, Grossbach J, Clement-Ziza M, Gillet L, Rodriguez-Lopez M, Marguerat S *et al* (2023). Genetic effects
on molecular network states explain complex traits. *Molecular Systems Biology* **19**(8)**:** e11493.

Wu MY, Hill CS (2009). *Tgf-beta* superfamily signaling in embryonic development and homeostasis.
*Developmental Cell* **16**, 329-343.
Xue C, Chu Q, Shi Q, Zeng Y, Lu J, Li L (2025). Wnt signaling pathways in biology and disease: mechanisms and
therapeutic advances. *Signal Transduction and Targeted Therapy* **10**, 106.

Ziyatdinov A, Vazquez-Santiago M, Brunel H, Martinez-Perez A, Aschard H, Soria JM (2018). lme4qtl: linear
mixed models with flexible covariance structure for genetic studies of related individuals. *BMC Bioinformatics*
**19**(1)**:** 68.

Zueva KJ, Lumme J, Veselov AE, Primmer CR, Pritchard VL (2020). Population genomics reveals repeated signals
of adaptive divergence in the Atlantic salmon of north-eastern Europe. *Journal of Evolutionary Biology* **34**(6)**:**
866-878.

# Tables (3) and Figures (2):

**Table 1:** Estimated fixed effect coefficients for 21 morphological traits obtained across three body planes. Trait lengths are given in centimeters (cm) for all traits. Coef. (SE) indicates coefficients and their standard error. B, S, and L in the body plane column indicates lateral view of the body, small and large cross sections, respectively. Significant *p* values are indicated with asterisk (* and ** denotes *p*<0.05 and *p*<0.01, respectively).

| Trait | Body Plane[1] | Trait value formula[1] | Intercept coef. (SE) | Length (scaled) coef. (SE) | Length (scaled) p value | Condition factor (scaled) coef. (SE) | Condition factor (scaled) p value | Feed restriction (*ad libitum* to restricted) coef. (SE) | Feed restriction p value | Sex (female to male) coef. (SE) | Sex p value | vgll3 (E to L) coef. (SE) | vgll3 p value | six6 (E to L) coef. (SE) | six6 p value |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Dorsal fin | B | B [12:13] | 2.0439 (0.0305) | 0.1560 (0.0064) | <0.001** | -0.0016 (0.0041) | 0.696 | -0.1222 (0.0176) | <0.001** | 0.0228 (0.0124) | 0.067 | -0.0246 (0.0148) | 0.106 | -0.0142 (0.0126) | 0.263 |
| Anal fin | B | B [16:17] | 1.35902 (0.0164) | 0.06636 (0.0041) | <0.001** | 0.00043 (0.0038) | 0.909 | -0.02549 (0.0094) | 0.0402* | 0.01759 (0.0076) | 0.021* | -0.01979 (0.0077) | 0.017* | 0.00658 (0.0075) | 0.378 |
| Pectoral fin | B | B [10:11] | 2.0973 (0.0202) | 0.1148 (0.0029) | <0.001** | 0.0082 (0.0022) | <0.001** | -0.0376 (0.0151) | 0.059 | 0.0143 (0.0053) | 0.007** | -0.0102 (0.0096) | 0.29 | 0.0030 (0.0058) | 0.601 |
| Tail fin | B | B [19:20] | 8.3e-01 (0.0098) | 4.1e-02 (0.0015) | <0.001** | -8.2e-03 (0.0013) | <0.001** | -2.1e-02 (0.0037) | 0.003** | 2.8e-05 (0.0027) | 0.992 | -4.0e-03 (0.0051) | 0.432 | 3.1e-03 (0.0030) | 0.292 |
| Caudal peduncle | B | B [15:18] | 8.7e-01 (0.00428) | 6.0e-02 (0.00078) | <0.001** | 1.1e-02 (0.00071) | <0.001** | 1.3e-03 (0.00198) | 0.529 | 2.3e-03 (0.00144) | 0.118 | -2.3e-03 (0.00219) | 0.294 | -6.7e-05 (0.00153) | 0.965 |
| Vertebra thickness | Sml | S [3:4] | 2.4e-01 (0.00265) | 1.6e-02 (0.00081) | <0.001** | 1.8e-03 (0.00082) | 0.0241* | 2.3e-03 (0.00165) | 0.172 | -2.2e-03 (0.00157) | 0.168 | -3.3e-03 (0.00117) | 0.009** | 5.3e-05 (0.00138) | 0.969 |
| Vertebra thickness | Lrg | L [6:7] | 0.25170 (0.00282) | 0.01394 (0.00077) | <0.001** | 0.00467 (0.00078) | <0.001** | -0.00238 (0.00155) | 0.125 | -0.00244 (0.00146) | 0.096 | -0.00297 (0.00131) | 0.032* | 0.00023 (0.00137) | 0.866 |
| Cross section diameter | Sml | S [5:7] | 1.1423 (0.0136) | 0.0753 (0.0027) | <0.001** | 0.0134 (0.0027) | <0.001** | -0.0562 (0.0114) | 0.003** | 0.0020 (0.0049) | 0.683 | -0.0157 (0.0058) | 0.012* | -0.0054 (0.0050) | 0.279 |
| Cross section diameter | Lrg | L [12:8] | 2.0269 (0.0117) | 0.1477 (0.0021) | <0.001** | 0.0312 (0.0021) | <0.001** | 0.0050 (0.0108) | 0.666 | 0.0019 (0.0039) | 0.621 | -0.0047 (0.0039) | 0.335 | -0.0046 (0.0039) | 0.248 |
| Aerobic muscle | Sml | S [(7:8+5:6)/2] | 0.17549 (0.00353) | 0.01024 (0.00082) | <0.001** | 0.00137 (0.00083) | 0.0981 | -0.01237 (0.00290) | 0.0051** | 0.00056 (0.00153) | 0.715 | -0.00557 (0.00148) | <0.001** | -0.00210 (0.00149) | 0.159 |
| Aerobic muscle | Lrg | L [(8:9+12:13)/2] | 0.2349 (0.00408) | 0.0185 (0.00094) | <0.001** | 0.0033 (0.00094) | <0.001** | 0.0070 (0.00239) | 0.025* | 0.0017 (0.00175) | 0.34 | -0.0024 (0.00195) | 0.222 | -0.0022 (0.00174) | 0.198 |
| Aerobic muscle vertical extension | Lrg | L [(14:15+10:11)/2] | 0.879 (0.0198) | 0.087 (0.0039) | <0.001** | 0.026 (0.0039) | <0.001** | 0.038 (0.0176) | 0.0916 | -0.012 (0.0073) | 0.089 | -0.024 (0.0081) | 0.006** | 0.013 (0.0072) | 0.067 |
| Eye diameter[2] | B | B [(2:3+4:5)/2] | 5.5e-01 (0.00358) | 2.2e-02 (0.00094) | <0.001** | 3.3e-04 (0.00092) | 0.723 | 1.8e-03 (0.00285) | 0.546 | -1.7e-06 (0.00175) | 0.999 | -3.2e-03 (0.00152) | 0.055 | 2.6e-03 (0.00163) | 0.112 |
| Snout length | B | B [1:0] | 0.78061 (0.00458) | 0.03839 (0.00097) | <0.001** | -0.00054 (0.00083) | 0.509 | -0.00277 (0.00196) | 0.157 | 0.00557 (0.00183) | 0.002** | -0.00607 (0.00232) | 0.013* | 0.00575 (0.00189) | 0.002** |
| Eye vertical position | B | B [0:6] | 0.40996 (0.0052) | 0.02487 (0.0012) | <0.001** | 0.00460 (0.0010) | <0.001** | 0.00934 (0.0036) | 0.0538 | 0.00024 (0.0022) | 0.912 | -0.00031 (0.0024) | 0.899 | 0.00057 (0.0022) | 0.794 |
| Head length[3] | B | B [1:7] | 2.1453 (0.0232) | 0.1347 (0.0034) | <0.001** | 0.0074 (0.0025) | 0.0034* | -0.0370 (0.0224) | 0.162 | 0.0106 (0.0064) | 0.094 | -0.0074 (0.0098) | 0.455 | 0.0146 (0.0068) | 0.032* |
| Head width[3] | B | B [8:9] | 1.59855 (0.0052) | 0.09654 (0.0015) | <0.001** | 0.01633 (0.0013) | <0.001** | 0.02432 (0.0030) | <0.001** | 0.00420 (0.0028) | 0.13 | -0.00046 (0.0024) | 0.85 | 0.00030 (0.0026) | 0.907 |
| Body centroid size | B | From Procrustes analysis | 13.1556 (0.0201) | 0.9084 (0.0031) | <0.001** | 0.0090 (0.0026) | <0.001** | 0.0505 (0.0124) | 0.0118* | -0.0088 (0.0057) | 0.124 | 0.0253 (0.0099) | 0.014* | -0.0227 (0.0061) | <0.001** |
| Head centroid size | B | From Procrustes analysis | 2.2095 (0.0131) | 0.1312 (0.0020) | <0.001** | 0.0143 (0.0016) | <0.001** | -0.0073 (0.0144) | 0.632 | 0.0077 (0.0037) | 0.040* | -0.0075 (0.0050) | 0.148 | 0.0104 (0.0039) | 0.008** |
| Centroid size | Sml | From Procrustes analysis | 1.4166 (0.0129) | 0.0968 (0.0023) | <0.001** | 0.0199 (0.0023) | <0.001** | -0.0387 (0.0102) | 0.0111* | 0.0033 (0.0042) | 0.432 | -0.0119 (0.0058) | 0.048* | -0.0048 (0.0044) | 0.28 |
| Centroid size | Lrg | From Procrustes analysis | 3.5944 (0.0177) | 0.2598 (0.0029) | <0.001** | 0.0601 (0.0029) | <0.001** | 0.0202 (0.0133) | 0.209 | -0.0015 (0.0053) | 0.778 | -0.0097 (0.0082) | 0.244 | -0.0054 (0.0056) | 0.337 |

1: The trait values are based on landmarks as numbered and indicated in Supplementary Figure 2.
2: Eye origin is the center point of eye landmarks, and calculated using landmarks 2, 3, 4, and 5, which are the anterior, posterior, dorsal and ventral extremes of the bony orbit of the eye.
3: Landmark 7 is the posterior point of the bony operculum. Landmarks 8 and 9 are the points of maximum curvature of the bony operculum at the ventral and dorsal ends, respectively.

**Table 2:** Multivariate analysis using Procrustes shape variable across three body planes in juvenile Atlantic salmon. Pillai is the Pillai's trace test statistic to evaluate each term's significance in the multivariate analysis of variance.

| Terms | Large cross section | | | | Small cross section | | | | Body | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | df | Pillai | z value | p value | df | Pillai | z value | p value | df | Pillai | z value | p value |
| Length (scaled) | 1 | 0.207 | 8.231 | 0.001 | 1 | 0.042 | 2.574 | 0.008 | 1 | 0.530 | 15.369 | 0.001 |
| Condition factor (scaled) | 1 | 0.053 | 1.398 | 0.074 | 1 | 0.024 | 0.894 | 0.195 | 1 | 0.478 | 10.583 | 0.001 |
| Feed restriction | 1 | 0.096 | 3.924 | 0.001 | 1 | 0.102 | 6.035 | 0.001 | 1 | 0.284 | 12.050 | 0.001 |
| Family | 37 | 1.939 | 9.994 | 0.001 | 37 | 0.721 | 2.200 | 0.011 | 37 | 2.446 | 24.524 | 0.001 |
| Tank | 4 | 0.312 | 6.244 | 0.001 | 4 | 0.219 | 7.020 | 0.001 | 3 | 0.276 | 8.643 | 0.001 |
| Sex | 1 | 0.035 | -0.118 | 0.548 | 1 | 0.013 | -0.597 | 0.717 | 1 | 0.038 | 1.185 | 0.121 |
| *vgll3* | 1 | 0.042 | 0.495 | 0.318 | 1 | 0.018 | 0.108 | 0.461 | 1 | 0.032 | 0.562 | 0.285 |
| *six6* | 1 | 0.027 | -0.992 | 0.831 | 1 | 0.013 | -0.617 | 0.739 | 1 | 0.052 | 2.454 | 0.013 |
| Full model | 47 | 2.727 | 14.021 | 0.001 | 47 | 1.256 | 7.674 | 0.001 | 46 | 3.929 | 23.997 | 0.001 |
| Residuals | 689 | | | | 681 | | | | 724 | | | |

**Table 3**: Variance components for 21 morphological traits obtained from three body planes. $V_A$, $h^2$, and $I_A$ are additive genetic variance, heritability, and evolvability, respectively, after controlling for length, condition factor, feed restriction, and sex.% $V_{A.vgll3}$ and%$V_{A.six6}$ are additive genetic variation explained by the *vgll3* and *six6* loci, respectively (values are truncated to zero if the estimate is less than zero). Values indicate medians, with 95% CI given in parenthesis. % rep. indicate the percent repeatability estimated from repeated measurements of individuals, and to assess the contribution of measurement error to trait measurements.

| Trait [1] | Body Plane[2] | $V_A$ | % $V_{A.vgll3}$ | %$V_{A.six6}$ | $h^2$ | $I_A$[3] | % rep. |
|---|---|---|---|---|---|---|---|
| Dorsal fin | B | 0.0032 (0.0027 - 0.0036) | 2.6 | 0 | 0.083 (0.072 - 0.095) | 0.045 (0.042 - 0.048) | 95.2 |
| Anal fin | B | 7,7e-04 (6.8 e-04, 8.8e-04) | 17.2 | 0 | 0.07 (0.061 - 0.08) | 0.27 (0.26 - 0.3) | 67.3 |
| Pectoral fin | B | 0.0027 (0.0024, 0.003) | 0 | 0 | 0.41 (0.36 - 0.45) | 0.016 (0.015 - 0.017) | 83.7 |
| Tail fin | B | 8.2e-04 (7.3e-04, 9.1e-04) | 0 | 0 | 0.5 (0.47 - 0.54) | 0.051 (0.048 - 0.055) | 69.7 |
| Caudal peduncle | B | 1e-04 (9.1e-05, 1.2e-04) | 0 | 0 | 0.26 (0.24 - 0.29) | 0.02 (0.019 - 0.021) | 65.3 |
| Vertebra thickness | Sml | 1.3e-05 (1.1e-05, 1.5e-05) | 30.7 | 0 | 0.039 (0.033 - 0.045) | 0.31 (0.28 - 0.35) | 63.7 |
| Vertebra thickness | Lrg | 2.0e-05 (1.7e-05, 2.3e-05) | 13 | 0 | 0.06 (0.052 - 0.069) | 0.21 (0.19 - 0.23) | 71.4 |
| Cross section diameter | Sml | 6.8e-04 (5.9e-04, 7.9e-04) | 12.7 | 0 | 0.12 (0.11 - 0.14) | 0.0099 (0.009 - 0.011) | 97.6 |
| Cross section diameter | Lrg | 3.3e-04 (2.9e-04, 3.8 e-04) | 0 | 0 | 0.11 (0.093 - 0.12) | 0.012 (0.01 - 0.013) | 84.1 |
| Aerobic muscle | Sml | 5.8e-05 (5.1e-05, 6.7e-05) | 22.4 | 0 | 0.19 (0.17 - 0.21) | 0.69 (0.62 - 0.76) | 56.5 |
| Aerobic muscle | Lrg | 7.5e-05 (6.6e-05, 8.7e-05) | 0 | 0 | 0.37 (0.34 - 0.41) | 0.72 (0.65 - 0.79) | 24.0 |
| Aerobic muscle vertical extension | Lrg | 0.001 (8.6 e-04, 0.0012) | 22.2 | 2.3 | 0.1 (0.087 - 0.12) | 0.29 (0.26 - 0.32) | 78.5 |
| Eye diameter | B | 3.1e-05 (2.7e-05, 3.5e-05) | 8.6 | 1.6 | 0.11 (0.094 - 0.12) | 0.23 (0.21 - 0.24) | 25.9 |
| Snout length | B | 9.0e-05 (7.9e-05, 1e-04) | 14.1 | 9.4 | 0.12 (0.11 - 0.14) | 0.023 (0.021 - 0.024) | 82.4 |
| Eye vertical position | B | 6.4e-05 (5.6e-05, 7.3e-05) | 0 | 0 | 0.06 (0.052 - 0.069) | 0.15 (0.14 - 0.16) | 79.7 |
| Head length | B | 0.0022 (0.0019, 0.0025) | 0 | 1.8 | 0.22 (0.19 - 0.26) | 0.018 (0.017 - 0.02) | 88.9 |
| Head width | B | 3.5e-05 (3e-05, 4e-05) | 0 | 0 | 0.021 (0.018 - 0.025) | 0.02 (0.019 - 0.022) | 75.4 |
| Body centroid size | B | 0.0034 (0.0031, 0.0038) | 6.3 | 4.2 | 0.47 (0.43 - 0.5) | 7.6e-04 (7,1e-04, 8.2e-04) | 73.8 |
| Head centroid size | B | 5e-04 (4.4e-04, 5.7e-04) | 0.5 | 5.1 | 0.15 (0.12 - 0.17) | 0.009 (0.0084 - 0.0096) | 85.1 |
| Centroid size | Sml | 8.3e-04 (7.2e-04, 9.5e-04) | 4.4 | 0 | 0.27 (0.24 - 0.3) | 0.043 (0.039 - 0.048) | 71.5 |
| Centroid size | Lrg | 0.0014 (0.0012, 0.0016) | 0 | 0 | 0.22 (0.19 - 0.24) | 0.0025 (0.0023 - 0.0028) | 93.5 |

1: Trait length unit is cm for all traits.
2: The trait values are based on landmarks as numbered and indicated in Supplementary Figure 2. B, S, and L prefixes represent body, small cross section, and large cross sections, respectively.
3: $I_A$ is measured as $V_A$ divided by the squared trait mean, and it is unit free.

**Figure 1**: The difference between *vgll3*[EE] and *vgll3*[LL] (blue lines), and *six6*[EE] and *six6*[LL] (red lines) genotypes on body shape of juvenile Atlantic salmon using thin-plate spline in the body (a), and large (b) and small (b) cross sections. The black dots in a, and green dots in b and c are the predicted shape coordinate associated with *vgll3*[EE] and *six6*[EE] after the generalized Procrustes analysis. For all images, the effect sizes are magnified by 10x. Landmark numbers are based on Supplementary Figure 2. Note that the body and cross section images are to illustrate the location of landmarks on these body planes, and were slightly distorted to fit onto the average coordinates obtained by Procrustes analysis.

**Figure 2:** Percent trait value change associated with *vgll3*E to *vgll3*L (red lines), and *six6*E to *six6*L (blue lines) allelic substitutions relative to the average trait values. Asterisks denote significant genetic effect. Error bars and 1SE in the x-axis indicate one standard error.