Structural mechanisms of PIP$_2$ activation and SEA0400 inhibition in human cardiac

sodium-calcium exchanger NCX1

Authors: Jing Xue[1,2,#], Weizhong Zeng[1,2], Scott John[3], Nicole Attiq[4], Michela Ottolia[4,*] &

Youxing Jiang[1,2,*]

Affiliations:

[1]Howard Hughes Medical Institute and Department of Physiology, the University of Texas

Southwestern Medical Center, Dallas, Texas, USA

[2]Department of Biophysics, the University of Texas Southwestern Medical Center, Dallas,

Texas, USA

[3]Department of Medicine (Cardiology), UCLA, Los Angeles, California, USA

[4]Department of Anesthesiology and Perioperative Medicine, Division of Molecular Medicine,

David Geffen School of Medicine, University of California Los Angeles, Los Angeles,

California, USA

[#]Current address: Renji Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai,

China

[*]**Address correspondence to:**

Youxing Jiang, Ph.D., Department of Physiology, UT Southwestern Medical Center, 5323 Harry

Hines Blvd., Dallas, Texas 75390-9040, Tel. 214 645-6027; Fax. 214 645-6042; E-Mail:

youxing.jiang@utsouthwestern.edu

Michela Ottolia, Ph.D., Department of Anesthesiology and Perioperative Medicine, Division of

Molecular Medicine, David Geffen School of Medicine, University of California Los Angeles,

Los Angeles, California; E-Mail: mottolia@g.ucla.edu

Abstract

Na^+/Ca^{2+} exchangers (NCXs) transport Ca^{2+} across the plasma membrane in exchange for Na^+ and play a vital role in maintaining cellular Ca^{2+} homeostasis. Our previous structural study of human cardiac NCX1 (HsNCX1) reveals the overall architecture of the eukaryotic exchanger and the formation of the inactivation assembly by the intracellular regulatory domain that underlies the cytosolic Na^+-dependent inactivation and Ca^{2+} activation of NCX1. Here we present the cryo-EM structures of HsNCX1 in complex with a physiological activator phosphatidylinositol 4,5-bisphosphate (PIP_2), or pharmacological inhibitor SEA0400 that enhances the inactivation of the exchanger. We demonstrate that PIP_2 binding stimulates NCX1 activity by inducing a conformational change at the interface between the TM and cytosolic domains that destabilizes the inactivation assembly. In contrast, SEA0400 binding in the TM domain of NCX1 stabilizes the exchanger in an inward-facing conformation that facilitates the formation of the inactivation assembly, thereby promoting the Na^+-dependent inactivation of NCX1. Thus, this study reveals the structural basis of PIP_2 activation and SEA0400 inhibition of NCX1 and provides some mechanistic understandings of cellular regulation and pharmacology of NCX family proteins.

Introduction

Sodium-calcium exchangers (NCX) are transporters that control the flux of Ca^{2+} across the plasma membrane and play a vital role in maintaining cellular calcium homeostasis for cell signaling[1-6]. NCXs facilitate the exchange of 3 Na^+ for 1 Ca^{2+} in an electrogenic manner, primarily responsible for extruding Ca^{2+} from the cytoplasm. However, this process can be reversed to permit Ca^{2+} entry, depending on the chemical gradients of Na^+ and Ca^{2+}, as well as the membrane potential[3,7-13]. Three NCX isoforms (NCX1-3) are present in mammals, with each isoform bearing multiple splice variants expressed in distinct tissues, thereby modulating numerous fundamental physiological events[4,14-19]. Dysfunctions of NCXs are associated with a plethora of human pathologies including cardiac hypertrophy, arrhythmia, and postischemic brain damage[3,20-22]. The cardiac variant NCX1.1 has been extensively studied, with its function playing a central role in cardiac excitation and contractile activity[23-27].

The eukaryotic NCX consists of a transmembrane (TM) domain with 10 TM helices and a large intracellular regulatory domain between TMs 5 and 6[4,28-33]. The TM domain is responsible for the ion exchange function in NCX. It consists of two homologous halves (TMs 1-5 and TMs 6-10) with TMs 2-3 and TMs 7-8 forming the core of the TM domain and hosting the residues that coordinate the transported Na^+ and Ca^{2+}[34,35]. The large regulatory domain contains two calcium-binding domains (CBD1 and CBD2). Ca^{2+} binding at CBDs enhances NCX activity and rescues it from the Na^+-dependent inactivation, a process that manifests as slow decay of the exchange current due to elevated levels of cytosolic Na^+ ions[12,30,31,36-41]. A stretch of residues known as XIP region (eXchanger Inhibitory Peptide) at the N terminus of the regulatory domain plays a pivotal role in the Na^+ inactivation process[42,43].

Several other cellular cues can also modulate NCX1 activity, including phosphatidylinositol 4,5-bisphosphate (PIP_2) in the membrane. PIP_2 has been shown to stimulate the NCX1 activity by reducing the Na^+-dependent inactivation and the XIP region was suggested to participate in the PIP_2 activation[39,44-46]. In addition, several small molecule NCX inhibitors have been developed to provide valuable tools for studying the physiological and pharmacological properties of NCXs, among which compound SEA0400 is a highly potent and selective NCX1 inhibitor that promotes the Na^+-dependent inactivation of the exchanger[20,47-52].

We previously determined the human cardiac NCX1 structure in an inward-facing inactivated state in which XIP and the β-hairpin between TMs 1 and 2 form a TM-associated

four-stranded β-hub and mediate a tight packing between the TM and cytosolic domains,
resulting in the formation of a stable inactivation assembly that blocks the TM movement
required for ion exchange function[32]. The study also provides a mechanistic insight into how
cytosolic Ca^{2+} binding at CBD2 destabilizes the inactivation assembly and activates the
exchanger[32]. To expand our understanding of NCX modulation by PIP_2 lipid and small molecule
inhibitors, we present the cryo-EM structures of human cardiac NCX1 in complex with PIP_2 or
SEA0400, revealing the structural basis underlying PIP_2 activation and SEA0400 inhibition of
NCX1.

107
108

Results

PIP$_2$ activation of NCX1

Phosphatidylinositol 4,5-bisphosphate (PIP$_2$) has been shown to modulate NCX1 activity

by reducing the Na$^+$-dependent inactivation of the exchanger[39,44-46]. To characterize the effect of

PIP$_2$ on HsNCX1, we expressed the exchanger in Xenopus laevis oocytes and recorded the

outward exchanger currents using the giant patch technique in the inside-out configuration with

or without applying porcine brain PIP$_2$ (Fig. 1, and Methods). The recording was performed with

12 μM free cytosolic $[Ca^{2+}]_i$ (bath) and the outward NCX1 current was elicited by the rapid

replacement of 100 mM Cs$^+$ with 100 mM Na$^+$ in the bath solution. In the patches without

applying PIP$_2$, the outward exchanger currents quickly decay and reach a steady state due to Na$^+$-

dependent inactivation (Fig. 1A). Introducing 10 μM PIP$_2$ at the steady state progressively

increases the current that plateaus at about two-fold of the peak current measured before PIP$_2$

addition (Fig. 1A). After PIP$_2$ washout, the outward current remains at the pre-removal level

without obvious inactivation, indicating high-affinity PIP$_2$ binding and its positive modulation of

NCX1 by both potentiating the peak current and reducing the Na$^+$-dependent inactivation.

Intriguingly, the commonly used shorter chain PIP$_2$ substitute (PIP$_2$ diC8) does not have the

equivalent activation effect on NCX1 and the exchanger remains susceptible to Na$^+$-dependent

inactivation when recorded in the presence of 10 μM PIP$_2$ diC8 (Fig. 1B). However, PIP$_2$ diC8

still binds and stimulates both the peak and steady currents of the exchanger. This stimulation

effect is abolished after PIP$_2$ diC8 washout, indicating a lower affinity of short-chain PIP$_2$ than

that of the long-chain native PIP$_2$.

To verify that the short-chain PIP$_2$ diC8 and the long-chain brain PIP$_2$ share the same

binding site, we performed a competition assay. As shown in Fig. 1C, introducing high-affinity

brain PIP$_2$ at the steady state yields a long-lasting potentiation of NCX1 current that is

irreversible even after a 5-minute washout. Applying PIP$_2$ diC8 can steadily decrease the brain

PIP$_2$-potentiated NCX1 current, suggesting that both lipids compete for the same binding site.

Structural insight into PIP$_2$ binding in NCX1

To reveal the structural mechanism of PIP$_2$ activation, we tried to obtain the EM structure

of HsNCX1 in the presence of the long-chain porcine brain PIP$_2$. However, the exchanger

becomes highly dynamic, yielding a low-resolution EM map with an overall shape similar to a

cytosolic Ca^{2+}-activated NCX1 whose β-hub-mediated inactivation assembly is destabilized and

cytosolic domain (CBD1 and CBD2) is detached from the TM domain[32] (Fig. S1). We suspect

the long-chain PIP_2 exerts the same activation effect on NCX1 as high cytosolic Ca^{2+} by

destabilizing the inactivation assembly. As NCX1 retains its Na^+-dependent inactivation property

in the presence of the short-chain PIP_2, we reasoned that the PIP_2 diC8-bound NCX1 likely

remain in an inward-facing inactivated state in high Na^+ low Ca^{2+} condition and its structure

would still reveal how PIP_2 binds in NCX1. We therefore determined the EM structure of NCX1

in complex with PIP_2 diC8 at 3.5Å (Fig. 2, Fig. S2, Table S1, and Methods), which indeed

adopts an inward-facing conformation with intact inactivation assembly. Due to the relative

dynamic movement between the TM and cytosolic domains, we also performed local refinement

to improve the map quality for each domain (Fig. S2). The density of the IP_3 head group from the

bound PIP_2 diC8 is well-defined in the local-refined EM map focused on the TM domain (Fig.

2A and Fig. S2B). This density is not present in the apo NCX1 structure (Fig. S3). The acyl

chains, however, are flexible and could not be resolved in the structure (Fig. S2). The lipid is

attached to the cytosolic sides of TMs 4 and 5 with its head group positioned at the C-terminal

end of TM5 (Fig. 2A). Four positively charged residues including K164 and R167 from the N-

terminus of TM4 and R220 and K225 from the C-terminus of TM5 are positioned in the vicinity

of the PIP_2 head group and likely participate in the electrostatic interactions with the head group

(Fig. 2A).

PIP_2 diC8-induced conformational changes in NCX1

Two major conformational changes occur in NCX1 upon PIP_2 diC8 binding (Fig. 2B-D).

The first change occurs at the C-terminus of TM5 which ends at R220 and is connected to the 2-

stranded XIP β-sheet (β3 and β4) via a 6-residue loop in the apo structure. When PIP_2 binds, the

TM5 helix extends by one helical turn. This loop-to-helix transition significantly changes the

locations of these connecting loop residues and their side-chain orientations to accommodate

PIP_2, enabling K225 to reorient and interact with the IP_3 head group (Fig. 2D, top panel). The

second conformational change is the partial detachment of the CBD2 domain from XIP upon

PIP_2 binding, resulting in a downward swing of cytosolic CBD domains (CBD1 and CBD2) as a

rigid body (Fig. 2B&C). This CBD2 detachment is a result of the first PIP_2-induced

conformational change at TM5 that disrupts part of the interactions between CBD2 and XIP as
follows: In the apo inactivated state, Y226 and R247, the two termini residues of the 2-stranded
XIP β-sheet, form H-bonds with several CBD2 residues including E554 sidechain and backbone
carbonyls of I518 and I520 (Fig. 2D, bottom panel); The loop-to-helix transition of TM5 upon
PIP$_2$ binding leads to a dramatic rotation of Y226 that allows it to move closer to and directly
interact with R247, resulting in the loss of H-bonding interactions between XIP and CBD2 (Fig.
2D, bottom panel); In addition, the rotation of Y226 also causes a direct collision with CBD2 if it
remains closely attached to XIP. Thus, the PIP$_2$-induced TM5 movement, particularly the
reorientation of Y226, abolishes some local interactions between CBD2 and XIP and also pushes
CBD2 away from XIP, causing the partial detachment of CBD2. However, the short-chain PIP$_2$
only partially destabilizes rather than completely disassembles the inactivation assembly as the
CH2 helix of CBD2 still engages in extensive interactions with the C-shaped β-hub.

To test if the PIP$_2$-interacting residues play a critical role in the native long-chain PIP$_2$
activation, we performed mutagenesis to those positively charged residues, including K164A,
R167A, R220A, and K225A single mutants. We also generated an R220A/K225A double mutant
as these two residues undergo PIP$_2$-induced conformational change at TM5. While all mutants
remain susceptible to current potentiation upon brain PIP$_2$ application as seen in the wild-type
NCX1, the PIP$_2$-potentiation effects on peak and steady-state currents are weakened in some
mutants, most notably in R220A and R220A/K225A (Fig. 3A-C). Interestingly, these two
mutants also have reduced Na$^+$-dependent inactivation as demonstrated by their higher fractional
activity (ratio between stead-state and peak currents) before applying PIP$_2$ (Fig. 3D). The
retained PIP$_2$ activation on these mutants implies that the longer acyl chain from native PIP$_2$
plays an important role in its interaction and activation of NCX1. Furthermore, as the
interactions between PIP$_2$ and NCX1 are both electrostatic involving multiple charged residues
and hydrophobic involving the long lipid acyl chain, those single or double amino acid
substitutions may only decrease the affinity of PIP$_2$ rather than abolish its binding. To test that,
we also mutated all four positively charged residues to alanine. The
K164A/R167A/R220A/K225A mutant is no longer sensitive to PIP$_2$ activation. The currents
from this quadruple mutant are small in most recordings and show no Na$^+$-dependent
inactivation. The unresponsiveness to PIP$_2$ and lack of Na$^+$-dependent inactivation in this mutant
is consistent with previous studies demonstrating that PIP$_2$ activates NCX by tuning the amount

of Na$^+$-dependent inactivation and any mutation that decreases NCX sensitivity to PIP$_2$ will also

affect the extent of Na$^+$-dependent inactivation[44]. This quadruple NCX1 mutant likely abolishes

the potentiation effect from both endogenous and externally applied PIP$_2$ and thereby functions

in a Na$^+$-inactivated steady state.

Structure of NCX1 in complex with SEA0400 inhibitor

SEA0400 is known to potently inhibit cardiac NCX1 by facilitating the inactivation of

the exchanger[50-52]. To reveal the structural mechanism of its inhibition, we determined the

structure of HsNCX1 in complex with SEA0400 at 2.9 Å (Fig. 4, Fig. S4, Table S1, and

Methods). The SEA0400-bound NCX1 structure adopts an inward-facing, inactivated state

identical to the apo NCX1 structure obtained at high Na$^+$, nominal Ca^{2+}-free condition [32] (Fig.

4A and Fig. S5). SEA0400 binds at the TM domain in a pocket enclosed by the internal halves of

TMs 8 (8a segment), 2 (2ab segments), 4, and 5 (Fig. 4B). The pocket has a lateral fenestration

in the middle of the membrane (Fig. 4B), which provides a portal for the SEA0400 entrance. The

pocket is sealed off at the cytosolic side by E244 from the XIP β-sheet. As demonstrated in our

previous study[32], XIP and the linker β-hairpin (β1 and β2) between TMs 1 and 2ab form a β-hub

that stabilizes the exchanger in an inactivated state. This β-hub has to be dissembled in an

activated NCX1 and XIP is expected to be detached from the TM domain, which would lead to

the opening of the SEA400-binding pocket to the cytosol and provide a cytosolic portal for the

release of the inhibitor release as further discussed below. Fig. 4C summarizes the interactions

between SEA0400 and NCX1 and demonstrates that SEA0400 fits perfectly in the pocket,

making extensive contact with surrounding residues. Mutations of some key interacting residues,

such as F213 and G833, have been shown to compromise the inhibitor binding[52]. The same

SEA0400 binding was also demonstrated in the recent study of human NCX1.3 and mutations at

some pocket-forming residues mitigate the SEA400 inhibition of the exchanger[33].

Inhibition mechanism of SEA0400

The TM domain of NCX1 shares a similar overall architecture to the archaea exchanger

NCX_Mj[34]. The structural comparison between the TM domains of the inward-facing NCX1 and

the outward-facing NCX_Mj reveals the conformational changes that occur during ion exchange,

providing structural insight into the SEA0400 inhibition mechanism (Fig. 5A). The inward-
outward transition mainly involves the sliding motion of TMs 1 and 6 and the bending
movement of TMs 2ab and 7ab[32,34,35,53]. As TM2ab directly interacts with SEA0400 in the
inward-facing state, its bending movement towards the outward conformation would cause a
direct collision with the inhibitor (Fig. 5A). Thus, the binding of SEA0400 stabilizes the
exchanger in the inward-facing state and blocks the conformational change from the inward to
the outward state. The Na^+-dependent NCX1 inactivation occurs when the exchanger is in a Na^+-
loaded, inward-facing state with low cytosolic $[Ca^{2+}]$[40,41,54]. As demonstrated in our previous
study, only in this state, the β-hub can form and readily interact with the cytosolic CBD domains,
generating the inactivation assembly that locks TMs 1 and 6 and prevents the TM module from
transporting ions[32]. Thus, SEA0400 promotes NCX1 inactivation by stabilizing NCX1 in the
inward-facing conformation which facilitates the formation of the inactivation assembly. The
formation of the inactivation assembly also reciprocally stabilizes SEA0400 binding as the XIP
of the assembly interacts with the TM domain and seals off the inhibitor binding pocket from the
cytosolic side (Fig. 5B). Under conditions in which the inactivation assembly cannot form in
NCX1, such as chymotrypsin treatment or forward exchange mode (Na^+ influx/Ca^{2+} efflux), the
removal or detachment of XIP from the TM domain would generate a cytosolic open portal for
SEA0400 release and effectively reduce its binding affinity (Fig. 5C)[50,51]. Indeed, cysteine
scanning mutagenesis studies have shown that the pocket-forming G833 residue is accessible to
intracellular sulfhydryl reagents in the chymotrypsin-treated inward-facing NCX1, confirming
the cytosolic exposure of the SEA400-binding pocket upon XIP removal[55]. This cytosolic
opening of the SEA0400 pocket explains the low efficacy of SEA0400 inhibition in NCX1 when
the Na^+-dependent inactivation is absent.

Discussion

In this study, we provide mechanistic underpinnings of cellular regulation and
pharmacology of NCX family proteins by revealing the structural basis of PIP_2 activation and
SEA0400 inhibition in HsNCX1. Both compounds modulate NCX1 activity by reducing or
enhancing the physiologically relevant Na^+-dependent inactivation process that occurs when the
exchanger is in an inward-facing, Na^+-loaded state with high $[Na^+]$ and low $[Ca^{2+}]$ on the

cytosolic side[27,40]. In this inactivation state, XIP and β-hairpin can assemble into a TM-
associated four-stranded β-hub that mediates a tight packing between the TM and cytosolic
domains, resulting in the formation of an inactivation assembly that blocks the TM
conformational changes required for ion exchange function[32,33]. PIP_2 or SEA0400 binding
changes the stability of the inactivation assembly in NCX1, resulting in a reduction or
potentiation of inactivation.

Although short-chain PIP_2 cannot fully recapitulate the NCX1 activation effect from
long-chain native PIP_2, the structure of PIP_2 diC8 NCX1 allows us to define the lipid-binding site
and the lipid-induced conformational changes at the interface between CBD2 and XIP that
destabilize the inactivation assembly. The SEA0400-bound NCX1 structure presented here,
along with the recent study by Dong et al.[33], suggests that the drug can directly inhibit NCX1 by
blocking the inward-outward conformational change at TM2ab. Our structural analysis also
explains the strong connection between SEA0400 binding and the Na^+-dependent inactivation -
SEA0400 is ineffective in an exchanger lacking Na^+-dependent inactivation whereas enhancing
the extent of Na^+-dependent inactivation increases the affinity for SEA0400. As SEA0400
binding traps NCX1 in an inward-facing state that facilitates the formation of the inactivation
assembly, the interaction between XIP and the TM domain in the assembly can in turn stabilize
SEA0400 binding by sealing the drug-binding pocket from the cytosol. When XIP is removed as
in the chymotrypsin-treated NCX1 or when the inactivation assembly cannot form as in the
NCX1 at forward exchange mode, the SEA0400-binding pocket becomes exposed to the cytosol,
mitigating the inhibition efficacy of SEA0400[50,51].

While we expect the long-chain PIP_2 to bind in the same location as PIP_2 diC8, it is
unclear how it exacerbates the destabilization effect on the inactivation assembly. The long acyl
chain of native PIP_2 may engage in some interactions with the TM module of NCX1 that are not
present in the short-chain lipid, rendering a higher affinity binding and more profound
destabilization effect on the inactivation assembly than PIP_2 diC8. The acyl-chain length-
dependent PIP_2 activation is consistent with some previous studies. Before PIP_2 was
demonstrated to regulate NCX, some earlier studies showed that negatively charged long-chain
lipids such as phosphatidylserine (PS) or phosphatidic acid (PA) could have the same
potentiation effects on NCX1 as PIP_2[56,57]. Furthermore, long-chain acyl-CoAs could also have
the same potentiation effects on NCX as PIP_2[58]. All these studies demonstrated that activation of

NCX by the anionic lipids depends on their chain length with the short chain being ineffective or
less effective. These findings have two implications. First, it is the negative surface charge rather
than the specific IP_3 head group of the lipid that is important for stimulating NCX1 activity,
suggesting non-specific electrostatic interactions between the negatively charged lipids and those
positively charged residues at the binding site. Second, a longer acyl chain is required for the
high-affinity binding of PIP_2 or negatively charged lipids. In the PIP_2 diC8-bound structure, the
tail of the acyl chain is positioned right at the fenestration of NCX1 that serves as the portal for
the SEA0400 binding. Two pieces of evidence lead us to suggest that the tail of the long acyl
chain from a native lipid can enter the same binding pocket for SEA0400 and thereby rendering
its higher affinity binding than a shorter chain lipid. Firstly, a docking analysis[59,60] of both long-
chain and short-chain PIP_2 at the binding site showed that the tail portion of the acyl chain from
the native brain PIP_2 inserts into the SEA0400-binding pocket through the fenestration of NCX1
in all docking models with highest calculated affinity (Fig. S6A). This pocket is not reachable for
PIP_2 diC8 due to its shorter chain length (Fig. S6B). Second, a stretch of density likely from a
native lipid acyl chain is observed in the apo NCX1 structure inside the SEA0400-binding pocket
(Fig. S6C), indicating its accessibility to a long-chain lipid. Thus, the accommodation of the
long acyl chain in the open pocket of NCX1 through the fenestration in the middle of the
membrane likely contributes to the high affinity binding of native lipids.
312
313
314

Methods

Protein expression and purification

The expression and purification of the HsNCX1 (cardiac isoform NCX1.1, indicated as

HsNCX1 or NCX1 throughout the manuscript) were carried out as described previously[32].

Truncated HsNCX1 (Δ341-365aa) containing a C-terminal Strep-tag was cloned into a pEZT-

BM vector and baculoviruses were produced in *Sf9* cells[61]. For protein expression, cultured

Expi293F GnTI- cells were infected with the baculoviruses at a ratio of 1:20 (virus: GnTI-, v/v)

for 10 hours. 10 mM sodium butyrate was then introduced to boost protein expression level and

cells were cultured in suspension at 30°C for another 60 hours and harvested by centrifugation at

4,000 g for 15 mins. All purification procedures were carried out at 4°C. The cell pellet was

resuspended in lysis buffer (25 mM HEPES pH 7.4, 300 mM NaCl, 2 μg/ml DNase I, 0.5 μg/ml

pepstatin, 2 μg/ml leupeptin, 1 μg/ml aprotinin, and 0.1 mM PMSF) and homogenized by

sonication. NCX1 was extracted with 2% (w/v) N-dodecyl-β-D-maltopyranoside (DDM,

Anatrace) supplemented with 0.2% (w/v) cholesteryl hemisuccinate (CHS, Sigma Aldrich) by

gentle agitation for 2 hours and supernatant collected by centrifugation at 40,000 g for 30 mins

was incubated with Strep-Tactin affinity resin (IBA) for 1 hour. The resin was then collected on

a disposable gravity column (Bio-Rad) and washed with 30 column volumes of buffer A (25 mM

HEPES pH 7.4, 200 mM NaCl, 0.06% digitonin). NCX1 was eluted in buffer A supplemented

with 50 mM biotin and further purified by size-exclusion chromatography on a Superdex200

10/300 GL column (GE Healthcare). For the generation of NCX1-Fab 2E4 complex, NCX1 was

incubated with purified Fab in a molar ratio of 1:1.2 (NCX1: Fab 2E4) for 2 hours and further

purified by size-exclusion chromatography in buffer A. The peak fractions were collected and

concentrated to ~5-6 mg/ml for cryo-EM analysis. To prepare the protein samples in complex

with various compounds, 0.9 mM SEA0400, 0.47 mM PI(4,5)P_2 diC8 or 0.42 mM brain

PI(4,5)P_2 were added to the protein samples 2 hours before grid preparation.

Expi293F GnTI- cells were purchased from and authenticated by Thermo Fisher

Scientific. The cell lines were tested negative for mycoplasma contamination.

Cryo-EM sample preparation and data acquisition

HsNCX1-Fab 2E4 samples (~5-6 mg/ml) in various conditions were applied to a glow-

discharged Quantifoil R1.2/1.3 300-mesh gold holey carbon grid (Quantifoil, Micro Tools

GmbH, Germany), blotted for 4.0 s under 100% humidity at 4 °C and plunged into liquid ethane

using a Mark IV Vitrobot (FEI). For the SEA0400-bound NCX1-Fab 2E4, raw movies were

acquired on a Titan Krios microscope (FEI) operated at 300 kV with a K3 camera (Gatan) at 0.83

Å per pixel and a nominal defocus range of -0.9 to -2.2 μm. Each movie was recorded for about

5 s in 60 subframes with a total dose of 60 $e^-/Å^2$. For other samples, raw movies were acquired

on a Titan Krios microscope operated at 300 kV with a Falcon 4i (Thermo Fisher Scientific) at

0.738 Å per pixel and a nominal defocus range of -0.8 to -1.8 μm. Each movie was recorded for

4 s with a total dose of 60 $e^-/Å^2$.

Image processing

Cryo-EM data were processed following the general scheme described below with some

modifications to different datasets (Figs. S1, S2, and S4). First, movie frames were motion-

corrected and dose-weighted using MotionCor2[62]. The CTF parameters of the micrographs were

estimated using the GCTF program[63]. After CTF estimation, micrographs were manually

inspected to remove images with bad defocus values and ice contamination. Particles were

picked using program Gautomatch (Kai Zhang, https://www.mrc-lmb.cam.ac.uk/kzhang/) or

crYOLO[64] and extracted with a binning factor of 3 in RELION[65,66]. Extracted particles were

subjected to 2D classification, ab initio modeling, and 3D classification. The particles from the

best-resolving 3D class were then re-extracted with the original pixel size and subjected to

heterogenous 3D refinement, non-uniform refinement, CTF refinement, and local refinement in

cryoSPARC[67]. The quality of the EM density maps for the transmembrane (TM) and cytosolic

domains was further improved through focused refinement, allowing for accurate model building

for a major part of the protein. For the dataset of NCX1 in complex with brain PI(4,5)P$_2$, the

maps of apo inactive NCX1 (PDB 8SGJ) and Ca^{2+}-bound active NCX1 (PDB 8SGT) are used as

references for heterogeneous refinement. Due to the highly dynamic nature of the protein

samples, the particles sorted into the active state produce a map with very low resolution. All

resolution was reported according to the gold-standard Fourier shell correlation (FSC) using the

0.143 criterion[68]. Local resolution was estimated using cryoSPARC.

Model building, refinement, and validation

The EM maps of HsNCX1 in the SEA0400-bound and PI(4,5)P$_2$ diC8-bound states show

high-quality density and model building is facilitated by previous apo NCX1 structure (PDB

8SGJ)[32]. Models were manually adjusted in Coot [69] and refined against maps using the

phenix.real_space_refine with secondary structure restraints applied[70]. The final NCX1structural

model contains residues 17-248, 370-467, 482-644, 652-698, 707-718, and 738-935. The EM

map of HsNCX1 in complex with brain PI(4,5)P$_2$ is relatively poor. The Ca^{2+}-bound activated

NCX1 structure (PDB 8SGT) is directly docked into the EM map without adjustment.

The statistics of the geometries of the models were generated using MolProbity[71]. All the

figures were prepared in PyMol (Schrödinger, LLC.), Chimera[72], and ChimeraX[73].

Electrophysiological experiments

The wild-type HsNCX1 and its mutants were cloned into a pGEMHE vector and

expressed in oocytes for electrophysiological recordings. RNA was synthesized using mMessage

mMachine (Ambion) and injected into Xenopus laevis oocytes as described in[74]. Oocytes were

isolated from at least 3 different frogs and kept at 18°C for 4-7 days. Outward HsNCX1 currents

were recorded using the giant patch technique in the inside-out configuration. Each data point

shown in this study represents a recording obtained from a single oocyte. The external solution

(pipette solution) contained the following (mM): 100 CsOH (cesium hydroxide), 10 HEPES (4-

(2-hydroxyethyl)-1-piperazineethanesulfonic acid), 20 TEAOH (tetraethyl-ammonium

hydroxide), 0.2 niflumic acid, 0.2 ouabain, 8 Ca(OH)$_2$ (calcium hydroxide), pH = 7 (using MES,

(2-(N-morpholino) ethanesulfonic acid)); bath solution (mM): 100 CsOH or 100 NaOH (sodium

hydroxide), 20 TEAOH, 10 HEPES, 10 EGTA(ethylene glycol-bis(β-aminoethyl ether)-

N,N,N',N'-tetra acetic acid) or HEDTA (N-(2-Hydroxyethyl) ethylenediamine-N,N',N'-triacetic

acid) and different Ca(OH)$_2$ concentrations to obtain the desired final free Ca^{2+} concentrations,

pH = 7 (using MES). Free Ca^{2+} concentrations were calculated according to WEBMAXc

program and confirmed with a Ca^{2+} electrode.

HsNCX1 currents were evoked by the rapid replacement of 100 mM Cs$^+$ with 100 mM

Na$^+$, using a computer-controlled 20-channel solution switcher. As HsNCX1 does not transport

Cs^+, there is no current and only upon application of Na^+ does the exchange cycle initiate. Data

were acquired at 4 ms/point and filtered at 50 Hz using an 8-pole Bessel filter. Experiments were

performed at 35°C and at a holding potential of 0 mV. The effects of the Na^+-dependent

inactivation were analyzed by measuring fractional currents calculated as the ratio of the steady-

state current to the peak current (fractional activity). All P values were calculated using an

unpaired, two-sided Welch's t-test.

PI(4,5)P_2 diC8 (Phosphatidylinositol 4,5-bisphosphate diC8, Echelon Bioscience) and

brain PI(4,5)P_2 (L-α-phosphatidylinositol-4,5-bisphosphate, Brain, Porcine, Avanti Polar Lipids)

were dissolved in water and kept as stock at -20 ℃. Immediately prior to recordings, PI(4,5)P_2

was diluted in the bath solution to 10 μM final concentration and perfused cytosolically for the

indicated time.

Data availability

The cryo-EM density maps of the human NCX1 have been deposited in the Electron

Microscopy Data Bank under accession numbers EMD-40456 for the SEA0400-bound state and

EMD-60921 for the PI(4,5)P_2 diC8-bound state, respectively. Atomic coordinates have been

deposited in the Protein Data Bank under accession numbers 8SGI for SEA0400-bound structure

and 9IV8 for the PI(4,5)P_2 diC8-bound structure.

Acknowledgments

Single particle cryo-EM data were collected at the University of Texas Southwestern Medical

Center Cryo-EM Facility which is funded by the CPRIT Core Facility Support Award

RP170644. Cryo-EM sample grids were prepared at the Structural Biology Laboratory at UT

Southwestern Medical Center which is partially supported by grant RP170644 from CPRIT. This

work was supported in part by the Howard Hughes Medical Institute (to Y.J.) and by grants from

the National Institute of Health (R35GM140892 to Y.J. and R01HL152296 to M.O.), and the

Welch Foundation (Grant I-1578 to Y.J.).

Author contributions

J.X. prepared the samples and performed data acquisition, image processing, and structure

determination; M.O., S.J., N.A., and W.Z. performed electrophysiology recording; Y.J.

supervised the work; all authors participated in research design, data analysis, discussion, and

manuscript preparation.

Competing interests

The authors declare no competing interests.

References

1.   Ottolia, M., John, S., Hazan, A., and Goldhaber, J.I. (2021). The Cardiac Na(+) -Ca(2+)
Exchanger: From Structure to Function. Compr Physiol *12*, 2681-2717.
10.1002/cphy.c200031.
2.   Clapham, D.E. (2007). Calcium signaling. Cell *131*, 1047-1058.
10.1016/j.cell.2007.11.028.
3.   Blaustein, M.P., and Lederer, W.J. (1999). Sodium/calcium exchange: its physiological
implications. Physiol Rev *79*, 763-854. 10.1152/physrev.1999.79.3.763.
4.   Philipson, K.D., and Nicoll, D.A. (2000). Sodium-calcium exchange: a molecular
perspective. Annu Rev Physiol *62*, 111-133. 10.1146/annurev.physiol.62.1.111.
5.   Berridge, M.J., Bootman, M.D., and Roderick, H.L. (2003). Calcium signalling: dynamics,
homeostasis and remodelling. Nat Rev Mol Cell Biol *4*, 517-529. 10.1038/nrm1155.
6.   DiPolo, R., and Beauge, L. (2006). Sodium/calcium exchanger: influence of metabolic
regulation on ion carrier interactions. Physiol Rev *86*, 155-203.
10.1152/physrev.00018.2005.
7.   Hilgemann, D.W., Nicoll, D.A., and Philipson, K.D. (1991). Charge movement during Na+
translocation by native and cloned cardiac Na+/Ca2+ exchanger. Nature *352*, 715-718.
10.1038/352715a0.
8.   Reeves, J.P., and Hale, C.C. (1984). The stoichiometry of the cardiac sodium-calcium
exchange system. J Biol Chem *259*, 7733-7739.
9.   Blaustein, M.P., and Russell, J.M. (1975). Sodium-calcium exchange and calcium-calcium
exchange in internally dialyzed squid giant axons. J Membr Biol *22*, 285-312.
10.1007/BF01868176.
10. Rasgadoflores, H., and Blaustein, M.P. (1987). Na/Ca Exchange in Barnacle Muscle-Cells
Has a Stoichiometry of 3 Na+/1 Ca2+. Am J Physiol *252*, C499-C504. DOI
10.1152/ajpcell.1987.252.5.C499.
11. Kimura, J., Noma, A., and Irisawa, H. (1986). Na-Ca Exchange Current in Mammalian
Heart-Cells. Nature *319*, 596-597. DOI 10.1038/319596a0.
12. Matsuoka, S., and Hilgemann, D.W. (1992). Steady-State and Dynamic Properties of
Cardiac Sodium-Calcium Exchange - Ion and Voltage Dependencies of the Transport
Cycle. J Gen Physiol *100*, 963-1001. DOI 10.1085/jgp.100.6.963.
13. Kang, T.M., and Hilgemann, D.W. (2004). Multiple transport modes of the cardiac
Na+/Ca2+ exchanger. Nature *427*, 544-548. 10.1038/nature02271.

14.   Quednau, B.D., Nicoll, D.A., and Philipson, K.D. (2004). The sodium/calcium exchanger
474   family - SLC8. Pflug Arch Eur J Phy *447*, 543-548. 10.1007/s00424-003-1065-4.

475   15.   Lee, S.L., Yu, A.S.L., and Lytton, J. (1994). Tissue-Specific Expression of Na+-Ca2+
476   Exchanger Isoforms. Journal of Biological Chemistry *269*, 14849-14852.

477   16.   Kofuji, P., Lederer, W.J., and Schulze, D.H. (1994). Mutually Exclusive and Cassette Exons
478   Underlie Alternatively Spliced Isoforms of the Na/Ca Exchanger. Journal of Biological
479   Chemistry *269*, 5145-5149.

480   17.   Linck, B., Qiu, Z., He, Z., Tong, Q., Hilgemann, D.W., and Philipson, K.D. (1998).
481   Functional comparison of the three isoforms of the Na+/Ca2+ exchanger (NCX1, NCX2,
482   NCX3). Am J Physiol *274*, C415-423. 10.1152/ajpcell.1998.274.2.C415.

483   18.   Lytton, J. (2007). Na+/Ca2+ exchangers: three mammalian gene families control Ca2+
484   transport. Biochem J *406*, 365-382. 10.1042/BJ20070619.

485   19.   Dyck, C., Omelchenko, A., Elias, C.L., Quednau, B.D., Philipson, K.D., Hnatowich, M., and
486   Hryshko, L.V. (1999). Ionic regulatory properties of brain and kidney splice variants of
487   the NCX1 Na(+)-Ca(2+) exchanger. J Gen Physiol *114*, 701-711. 10.1085/jgp.114.5.701.

488   20.   Watanabe, Y., Koide, Y., and Kimura, J. (2006). Topics on the Na+/Ca2+ exchanger:
489   pharmacological characterization of Na+/Ca2+ exchanger inhibitors. J Pharmacol Sci
490   *102*, 7-16. 10.1254/jphs.fmj06002x2.

491   21.   Pott, C., Eckardt, L., and Goldhaber, J.I. (2011). Triple threat: the Na+/Ca2+ exchanger in
492   the pathophysiology of cardiac arrhythmia, ischemia and heart failure. Curr Drug Targets
493   *12*, 737-747. 10.2174/138945011795378559.

494   22.   Matsuda, T., Takuma, K., and Baba, A. (1997). Na(+)-Ca2+ exchanger: physiology and
495   pharmacology. Jpn J Pharmacol *74*, 1-20. 10.1254/jjp.74.1.

496   23.   Shigekawa, M., and Iwamoto, T. (2001). Cardiac Na(+)-Ca(2+) exchange: molecular and
497   pharmacological aspects. Circ Res *88*, 864-876. 10.1161/hh0901.090298.

498   24.   Kimura, J., Miyamae, S., and Noma, A. (1987). Identification of sodium-calcium exchange
499   current in single ventricular cells of guinea-pig. J Physiol *384*, 199-222.
500   10.1113/jphysiol.1987.sp016450.

501   25.   Bridge, J.H.B., Smolley, J.R., and Spitzer, K.W. (1990). The Relationship between Charge
502   Movements Associated with Ica and Ina-Ca in Cardiac Myocytes. Science *248*, 376-378.
503   DOI 10.1126/science.2158147.

504   26.   Ottolia, M., Torres, N., Bridge, J.H., Philipson, K.D., and Goldhaber, J.I. (2013). Na/Ca
505   exchange and contraction of the heart. J Mol Cell Cardiol *61*, 28-33.
506   10.1016/j.yjmcc.2013.06.001.

507   27.   Scranton, K., John, S., Angelini, M., Steccanella, F., Umar, S., Zhang, R., Goldhaber, J.I.,
508   Olcese, R., and Ottolia, M. (2024). Cardiac function is regulated by the sodium-
509   dependent inhibition of the sodium-calcium exchanger NCX1. Nat Commun *15*, 3831.
510   10.1038/s41467-024-47850-z.

511   28.   Ren, X., and Philipson, K.D. (2013). The topology of the cardiac Na(+)/Ca(2)(+)
512   exchanger, NCX1. J Mol Cell Cardiol *57*, 68-71. 10.1016/j.yjmcc.2013.01.010.

513   29.   Sharma, V., and O'Halloran, D.M. (2014). Recent Structural and Functional Insights into
514   the Family of Sodium Calcium Exchangers. Genesis *52*, 93-109. 10.1002/dvg.22735.

30. Matsuoka, S., Nicoll, D.A., Reilly, R.F., Hilgemann, D.W., and Philipson, K.D. (1993). Initial
localization of regulatory regions of the cardiac sarcolemmal Na(+)-Ca2+ exchanger.
Proc Natl Acad Sci U S A *90*, 3870-3874. 10.1073/pnas.90.9.3870.

31. Hilgemann, D.W. (1990). Regulation and Deregulation of Cardiac Na+-Ca2+ Exchange in
Giant Excised Sarcolemmal Membrane Patches. Nature *344*, 242-245. DOI
10.1038/344242a0.

32. Xue, J., Zeng, W., Han, Y., John, S., Ottolia, M., and Jiang, Y. (2023). Structural
mechanisms of the human cardiac sodium-calcium exchanger NCX1. Nat Commun *14*,
6181. 10.1038/s41467-023-41885-4.

33. Dong, Y., Yu, Z., Li, Y., Huang, B., Bai, Q., Gao, Y., Chen, Q., Li, N., He, L., and Zhao, Y.
(2024). Structural insight into the allosteric inhibition of human sodium-calcium
exchanger NCX1 by XIP and SEA0400. EMBO J *43*, 14-31. 10.1038/s44318-023-00013-0.

34. Liao, J., Li, H., Zeng, W.Z., Sauer, D.B., Belmares, R., and Jiang, Y.X. (2012). Structural
Insight into the Ion-Exchange Mechanism of the Sodium/Calcium Exchanger. Science
335, 686-690. 10.1126/science.1215759.

35. Liao, J., Marinelli, F., Lee, C., Huang, Y., Faraldo-Gomez, J.D., and Jiang, Y. (2016).
Mechanism of extracellular ion exchange and binding-site occlusion in a sodium/calcium
exchanger. Nat Struct Mol Biol *23*, 590-599. 10.1038/nsmb.3230.

36. Matsuoka, S., Nicoll, D.A., Hryshko, L.V., Levitsky, D.O., Weiss, J.N., and Philipson, K.D.
(1995). Regulation of the cardiac Na(+)-Ca2+ exchanger by Ca2+. Mutational analysis of
the Ca(2+)-binding domain. J Gen Physiol *105*, 403-420. 10.1085/jgp.105.3.403.

37. Hilge, M., Aelen, J., and Vuister, G.W. (2006). Ca2+ regulation in the Na+/Ca2+
exchanger involves two markedly different Ca2+ sensors. Mol Cell *22*, 15-25.
10.1016/j.molcel.2006.03.008.

38. Ottolia, M., Nicoll, D.A., and Philipson, K.D. (2009). Roles of two Ca2+-binding domains
in regulation of the cardiac Na+-Ca2+ exchanger. J Biol Chem *284*, 32735-32741.
10.1074/jbc.M109.055434.

39. Hilgemann, D.W., Collins, A., and Matsuoka, S. (1992). Steady-state and dynamic
properties of cardiac sodium-calcium exchange. Secondary modulation by cytoplasmic
calcium and ATP. J Gen Physiol *100*, 933-961. 10.1085/jgp.100.6.933.

40. Hilgemann, D.W., Matsuoka, S., Nagel, G.A., and Collins, A. (1992). Steady-state and
dynamic properties of cardiac sodium-calcium exchange. Sodium-dependent
inactivation. J Gen Physiol *100*, 905-932. 10.1085/jgp.100.6.905.

41. Matsuoka, S., and Hilgemann, D.W. (1994). Inactivation of outward Na(+)-Ca2+
exchange current in guinea-pig ventricular myocytes. J Physiol *476*, 443-458.
10.1113/jphysiol.1994.sp020146.

42. Matsuoka, S., Nicoll, D.A., He, Z., and Philipson, K.D. (1997). Regulation of cardiac Na(+)-
Ca2+ exchanger by the endogenous XIP region. J Gen Physiol *109*, 273-286.
10.1085/jgp.109.2.273.

43. Li, Z., Nicoll, D.A., Collins, A., Hilgemann, D.W., Filoteo, A.G., Penniston, J.T., Weiss, J.N.,
Tomich, J.M., and Philipson, K.D. (1991). Identification of a peptide inhibitor of the
cardiac sarcolemmal Na(+)-Ca2+ exchanger. J Biol Chem *266*, 1014-1020.

44.   He, Z., Feng, S., Tong, Q., Hilgemann, D.W., and Philipson, K.D. (2000). Interaction of
558   PIP(2) with the XIP region of the cardiac Na/Ca exchanger. Am J Physiol Cell Physiol *278*,
559   C661-666. 10.1152/ajpcell.2000.278.4.C661.

560   45.   Hilgemann, D.W., and Ball, R. (1996). Regulation of cardiac Na+,Ca2+ exchange and
561   KATP potassium channels by PIP2. Science *273*, 956-959. 10.1126/science.273.5277.956.

562   46.   Yaradanakul, A., Feng, S., Shen, C., Lariccia, V., Lin, M.J., Yang, J., Kang, T.M., Dong, P.,
563   Yin, H.L., Albanesi, J.P., and Hilgemann, D.W. (2007). Dual control of cardiac Na+ Ca2+
564   exchange by PIP(2): electrophysiological analysis of direct and indirect mechanisms. J
565   Physiol *582*, 991-1010. 10.1113/jphysiol.2007.132712.

566   47.   Tanaka, H., Nishimaru, K., Aikawa, T., Hirayama, W., Tanaka, Y., and Shigenobu, K.
567   (2002). Effect of SEA0400, a novel inhibitor of sodium-calcium exchanger, on myocardial
568   ionic currents. Br J Pharmacol *135*, 1096-1100. 10.1038/sj.bjp.0704574.

569   48.   Matsuda, T., Arakawa, N., Takuma, K., Kishida, Y., Kawasaki, Y., Sakaue, M., Takahashi,
570   K., Takahashi, T., Suzuki, T., Ota, T., et al. (2001). SEA0400, a novel and selective inhibitor
571   of the Na+-Ca2+ exchanger, attenuates reperfusion injury in the in vitro and in vivo
572   cerebral ischemic models. J Pharmacol Exp Ther *298*, 249-256.

573   49.   Watano, T., Kimura, J., Morita, T., and Nakanishi, H. (1996). A novel antagonist, No.
574   7943, of the Na+/Ca2+ exchange current in guinea-pig cardiac ventricular cells. Br J
575   Pharmacol *119*, 555-563. 10.1111/j.1476-5381.1996.tb15708.x.

576   50.   Lee, C., Visen, N.S., Dhalla, N.S., Le, H.D., Isaac, M., Choptiany, P., Gross, G.,
577   Omelchenko, A., Matsuda, T., Baba, A., et al. (2004). Inhibitory profile of SEA0400 [2-[4-
578   [(2,5-difluorophenyl)methoxy]phenoxy]-5-ethoxyaniline] assessed on the cardiac Na+-
579   Ca2+ exchanger, NCX1.1. J Pharmacol Exp Ther *311*, 748-757. 10.1124/jpet.104.070805.

580   51.   Bouchard, R., Omelchenko, A., Le, H.D., Choptiany, P., Matsuda, T., Baba, A., Takahashi,
581   K., Nicoll, D.A., Philipson, K.D., Hnatowich, M., and Hryshko, L.V. (2004). Effects of
582   SEA0400 on mutant NCX1.1 Na+-Ca2+ exchangers with altered ionic regulation. Mol
583   Pharmacol *65*, 802-810. 10.1124/mol.65.3.802.

584   52.   Iwamoto, T., Kita, S., Uehara, A., Imanaga, I., Matsuda, T., Baba, A., and Katsuragi, T.
585   (2004). Molecular determinants of Na+/Ca2+ exchange (NCX1) inhibition by SEA0400. J
586   Biol Chem *279*, 7544-7553. 10.1074/jbc.M310491200.

587   53.   Marinelli, F., and Faraldo-Gomez, J.D. (2023). Conformational free-energy landscapes of
588   a Na(+)/Ca(2+) exchanger explain its alternating-access mechanism and functional
589   specificity. bioRxiv. 10.1101/2023.01.20.524959.

590   54.   Hilgemann, D.W. (2020). Regulation of ion transport from within ion transit pathways. J
591   Gen Physiol *152*. 10.1085/jgp.201912455.

592   55.   John, S.A., Liao, J., Jiang, Y., and Ottolia, M. (2013). The cardiac Na+-Ca2+ exchanger has
593   two cytoplasmic ion permeation pathways. Proc Natl Acad Sci U S A *110*, 7500-7505.
594   10.1073/pnas.1218751110.

595   56.   Hilgemann, D.W., and Collins, A. (1992). Mechanism of cardiac Na(+)-Ca2+ exchange
596   current stimulation by MgATP: possible involvement of aminophospholipid translocase.
597   J Physiol *454*, 59-82. 10.1113/jphysiol.1992.sp019254.

598   57.   Vemuri, R., and Philipson, K.D. (1988). Phospholipid composition modulates the Na+-
599   Ca2+ exchange activity of cardiac sarcolemma in reconstituted vesicles. Biochim Biophys
600   Acta *937*, 258-268. 10.1016/0005-2736(88)90248-9.

601   58.   Riedel, M.J., Baczko, I., Searle, G.J., Webster, N., Fercho, M., Jones, L., Lang, J., Lytton, J.,
602   Dyck, J.R., and Light, P.E. (2006). Metabolic regulation of sodium-calcium exchange by
603   intracellular acyl CoAs. EMBO J *25*, 4605-4614. 10.1038/sj.emboj.7601321.
604   59.   Eberhardt, J., Santos-Martins, D., Tillack, A.F., and Forli, S. (2021). AutoDock Vina 1.2.0:
605   New Docking Methods, Expanded Force Field, and Python Bindings. J Chem Inf Model
606   *61*, 3891-3898. 10.1021/acs.jcim.1c00203.
607   60.   Bugnon, M., Röhrig, U.F., Goullieux, M., Perez, M.A.S., Daina, A., Michielin, O., and
608   Zoete, V. (2024). SwissDock 2024: major enhancements for small-molecule docking with
609   Attracting Cavities and AutoDock Vina. Nucleic Acids Res *52*, W324-w332.
610   10.1093/nar/gkae300.
611   61.   Morales-Perez, C.L., Noviello, C.M., and Hibbs, R.E. (2016). Manipulation of Subunit
612   Stoichiometry in Heteromeric Membrane Proteins. Structure *24*, 797-805.
613   10.1016/j.str.2016.03.004.
614   62.   Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A. (2017).
615   MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron
616   microscopy. Nat Methods *14*, 331-332. 10.1038/nmeth.4193.
617   63.   Zhang, K. (2016). Gctf: Real-time CTF determination and correction. J Struct Biol *193*, 1-
618   12. 10.1016/j.jsb.2015.11.003.
619   64.   Wagner, T., Merino, F., Stabrin, M., Moriya, T., Antoni, C., Apelbaum, A., Hagel, P., Sitsel,
620   O., Raisch, T., Prumbaum, D., et al. (2019). SPHIRE-crYOLO is a fast and accurate fully
621   automated particle picker for cryo-EM. Commun Biol *2*, 218. 10.1038/s42003-019-0437-
622   z.
623   65.   Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., and
624   Scheres, S.H. (2018). New tools for automated high-resolution cryo-EM structure
625   determination in RELION-3. Elife *7*. 10.7554/eLife.42166.
626   66.   Scheres, S.H. (2012). RELION: implementation of a Bayesian approach to cryo-EM
627   structure determination. J Struct Biol *180*, 519-530. 10.1016/j.jsb.2012.09.006.
628   67.   Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). cryoSPARC:
629   algorithms for rapid unsupervised cryo-EM structure determination. Nat Methods *14*,
630   290-296. 10.1038/nmeth.4169.
631   68.   Henderson, R., Sali, A., Baker, M.L., Carragher, B., Devkota, B., Downing, K.H., Egelman,
632   E.H., Feng, Z., Frank, J., Grigorieff, N., et al. (2012). Outcome of the first electron
633   microscopy validation task force meeting. Structure *20*, 205-214.
634   10.1016/j.str.2011.12.014.
635   69.   Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and development
636   of Coot. Acta Crystallogr D Biol Crystallogr *66*, 486-501. 10.1107/S0907444910007493.
637   70.   Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J.,
638   Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010). PHENIX: a comprehensive
639   Python-based system for macromolecular structure solution. Acta Crystallogr D Biol
640   Crystallogr *66*, 213-221. S0907444909052925 [pii]
641   10.1107/S0907444909052925.
642   71.   Chen, V.B., Arendall, W.B., 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J.,
643   Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: all-atom

structure validation for macromolecular crystallography. Acta Crystallogr D Biol
Crystallogr *66*, 12-21. 10.1107/S0907444909042073.
72. Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C.,
and Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory research
and analysis. J Comput Chem *25*, 1605-1612. 10.1002/jcc.20084.
73. Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H., and
Ferrin, T.E. (2018). UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. Protein Sci *27*, 14-25. 10.1002/pro.3235.
74. John, S., Kim, B., Olcese, R., Goldhaber, J.I., and Ottolia, M. (2018). Molecular
determinants of pH regulation in the cardiac Na(+)-Ca(2+) exchanger. J Gen Physiol *150*,
245-257. 10.1085/jgp.201611693.
655

Fig. 1. PIP$_2$ enhances HsNCX1 activity.

(A) Representative outward currents recorded from oocytes expressing the human NCX1 before and after application of long-chain brain PIP$_2$. Currents were activated by replacing cytosolic Cs$^+$ with Na$^+$. Application of 10 μM brain PIP$_2$ enhanced HsNCX1 current and abolished the Na$^+$-dependent inactivation irreversibly. Perfusion time of PIP$_2$ is indicated above traces while lines below traces indicate solution exchange. Arrows mark the peak and steady currents used to measure the fold of increase upon PIP$_2$ application. The fold of current increase was calculated by comparing the peak or steady-state current before and after PIP$_2$ application.

(B) Representative outward currents recorded before and after application of short-chain PIP$_2$ diC8 (10 μM). PIP$_2$ diC8 was perfused from the cytosolic side before HsNCX1 activation (in the presence of Cs$^+$ for 30 s) and during transport (in the presence of Na$^+$). Both peak and steady-state currents of HsNCX1 are enhanced by PIP$_2$ diC8 and the effect is reversible. The Na$^+$-dependent inactivation remains in the presence of PIP$_2$ diC8.

(C) Representative outward currents recorded with the application of brain PIP$_2$ and PIP$_2$ diC8. The NCX1 current was first potentiated by applying 10 μM brain PIP$_2$ at the steady state. The PIP$_2$ effect was not reversible over the 5-minute washout with a solution containing 100 mM Na$^+$ and 12 μM Ca^{2+}. The same patch was then perfused with the same solution in the presence of 10 μM PIP$_2$ diC8. Application of the short-chain PIP$_2$ diC8 facilitates the decrease of brain PIP$_2$-potentiated current, suggesting that both lipids compete for the same binding site.

Fig. 2. PIP₂ binding in NCX1.

(**A**) Structure of the TM and β-hub regions of PIP₂ diC8-bound NCX1 with a zoomed-in view of the lipid binding site. The density of PIP₂ diC8 is shown as a grey mesh contoured at 5.5σ.

(**B**) Structural comparison between apo (grey) and PIP₂ diC8-bound (color) NCX1. Upon PIP2 diC8 binding, there is a rigid-body downward swing movement (marked by an arrow) at CBDs caused by the partial detachment of the CBD2 domain from XIP. The conformational change at the TM domain is subtle and mainly occurs at the C-terminus of TM5 as illustrated in (**C**) and (**D**).

(**C**) Zoomed-in view of the structural comparison (boxed area in (B)). The two major conformational changes occur in the boxed regions. (**D**) Zoomed-in views of the two conformational changes between apo (left in grey) and PIP₂ diC8-bound (right in color) state. Top: conformational change 1 at the C-terminus of TM5. Bottom: conformational change 2 at the interface between XIP and CBD2.

Fig. 3. Mutagenesis at the PIP$_2$ binding.
(A) Representative NCX1 currents of PIP$_2$ site mutants before and after perfusion of 10 µM brain PIP$_2$ to the cytosolic side of the patch.
(B&C) Summary graphs demonstrating the effects of PIP$_2$ on the enhancement of peak (B) and steady-state currents (C). Potentiation (fold of increase) was measured by comparing the current magnitude before and after PIP$_2$ application. Mutants R167A, R220A, and K225A showed some decreased response to PIP$_2$ whereas R220A/K225A mutant shows more profound decrease in PIP$_2$ response. Compared to WT, the PIP$_2$ potentiation of R220A/K225A at the steady state is decreased by ~70-90% (Fold of increase WT=8.9±1.6, n=10 vs R220A/K225A=1.9±0.1, n=6). PIP$_2$ has no potentiation effect on the quadruple K164A/R167A/R220A/K225A mutant. Data points are mean ± s.e.m. (* p <0.1).
(D) The extent of Na$^+$-dependent inactivation was measured as the ratio between steady state and peak currents (fractional activity) and values for WT and the indicated mutants are shown. Mutants R220A and R220A/K225A displayed significantly higher fractional activity values when compared to WT, indicating that the Na$^+$-dependent inactivation was less pronounced in these mutant exchangers. K164A/R167A/R220A/K225A shown no Na$^+$-dependent inactivation.

Fig. 4. SEA0400 binding in NCX1.

(**A**) Overall structure of human NCX1 in complex with SEA0400 obtained in high Na^+ and low Ca^{2+} conditions. Yellow spheres represent the bound Ca^{2+} in CBD1 and XIP.

(**B**) Cartoon representation of the TM domain and β-hub of the complex with surface-rendered view of the fenestration in the middle of the membrane. The β-hub is assembled by β-hairpin (β1 and β2) and XIP (β3 and β4).

(**C**) Zoomed-in view of SEA0400 binding site and the schematic diagram detailing the interactions between NCX1 residues and SEA0400.

Fig. 5. Structural mechanism of SEA0400 inhibition.
(A) Structural comparison at the core part of the TM domain between the SEA0400-bound, inward-facing HsNCX1 and the outward-facing NCX_Mj (PDB 3V5U). Red arrows mark the sliding movement of TMs 1 and 6 and the bending of TM2ab from inward to outward conformation.
(B) Surface-rendered views of the SEA0400-binding pocket sealed off from the cytosolic side by E244 from XIP in the inactivated state.
(C) Removal of XIP would generate a cytosolic portal that facilitates the release of SEA0400.

MotionCor2
Gctf
crYOLO picking

8604 micrographs

1,742,058 particles

Heterogeneous Refinement

365,702 particles

Further
heterogeneous Refinement

FSC Resolution: 11.45Å

No Mask (12Å)
Spherical (12Å)
Loose (11Å)

Fab

282,455 particles

Ca^{2+}-activated model fitted into the map

Fig. S1. Cryo-EM data processing of HsNCX1 in the presence of long-chain porcine brain PIP₂. The structural model of Ca^{2+}-activated HsNCX1 from a previous study (PDB 8SGT) was directly fitted into the low-resolution EM-map (~11.5 Å). The Fab fragment from a monoclonal antibody against NCX1 was used as a fiducial marker to facilitate the single-particle alignment.

Fig. S2. Structure determination of HsNCX1-PIP$_2$ diC8 complex.
(A) Cryo-EM data processing of HsNCX1 in complex with short-chain PIP$_2$ diC8. The Fab fragment from a monoclonal antibody against NCX1 was used as a fiducial marker to facilitate the single-particle alignment.
(B) Zoomed-in view of density map of the bound PIP$_2$ diC8 contoured at the threshold level of 0.35 using ChimeraX.
(C) The Fourier shell correlation (FSC) curves for cross-validation between the maps and the models.

PIP₂-bound   apo

Fig. S3. Side-by-side comparison of the densities at the PIP₂-binding site between the PIP₂-bound structure (EMD-60921) and the apo structure (EMD-40457). The local-refined maps focused on the TM domain are used in the comparison. The density map of the bound PIP₂ is contoured at a threshold level of 0.35 using ChimeraX.

Fig. S4. Structure determination of HsNCX1-SEA0400 complex.
(A) Cryo-EM data processing scheme of HsNCX1 in complex with SEA0400 inhibitor. The Fab fragment from a monoclonal antibody against NCX1 was used as a fiducial marker to facilitate the single-particle alignment.
(B) Zoomed-in view of density map of the bound SEA0400 inhibitor contoured at the threshold level of 0.52 using ChimeraX.
(C) The Fourier shell correlation (FSC) curves for cross-validation between the maps and the models.

Fig. S5. Structural comparison between the apo (grey) and SEA0400-bound (color) HsNCX1.

Fig. S6. Proposed structural basis underlying the different binding affinity between long and short-chain PIP$_2$.

(A) A docking model for native brain PIP$_2$ binding in NCX1 showing the insertion of its long acyl chain into the SEA0400-binding pocket.

(B) A docking model for short-chain PIP$_2$ diC8 binding in NCX1 showing that the SEA0400-binding pocket is not accessible to shorter acyl chain.

(C) The density (red mesh, contoured at 6σ) of an acyl chain likely from a native lipid is observed in the SEA0400-binding pocket of the apo NCX1 structure (EMD-40457, local-refined map at the TM domain).

Table S1. Cryo-EM data collection and model statistics.

Sample preparation conditions	25 mM Hepes pH 7.4, 200 mM NaCl 0.9 mM SEA0400	25 mM Hepes pH 7.4, 200 mM NaCl, 0.47 mM PI(4,5)P₂ diC8
	SEA0400-bound state (EMD-40456, PDB 8SGI)	**PI(4,5)P₂ diC8-bound state** (EMD-60921, PDB 9IV8)
Data collection and processing		
Magnification	105k	105k
Voltage (kV)	300	300
Electron exposure ($e^-/Å^2$)	60	60
Defocus range (μm)	-0.9 - -2.2	-0.9 - -2.2
Pixel size (Å)	0.83	0.84
Symmetry imposed	C1	C1
Initial particle images (no.)	1,249,151	2,233,044
Final particle images (no.)	368,227	117,748
Map resolution (Å)	2.93	3.47
FSC threshold	0.143	0.143
Refinement		
Initial model used (PDB code)	8SGJ	8SGJ
Model resolution (Å)	3.36	3.81
FSC threshold	0.5	0.5
Model composition		
Non-hydrogen atoms	7667	5947
Protein residues	982	750
Ligands	3: Na 6: Ca 1: H_2O 1: SEA0400	5: Ca 1: PI(4,5)P₂ diC8
B factors ($Å^2$)		
Protein	66.21	50.47
Ligands	58.55	106.20
R.m.s. deviations		
Bond lengths (Å)	0.005	0.004
Bond angles (°)	0.696	0.679
Validation		
MolProbity score	1.36	1.28
Clashscore	5.24	5.22
Poor rotamers (%)	0	0
Ramachandran plot		
Favored (%)	97.62	98.24
Allowed (%)	2.38	1.76
Disallowed (%)	0	0