Vocal species are more central in Eastern Himalayan Mixed-Species

bird flocks

Satyam Gupta[1*], Akshay Bharadwaj[2,3], Akshiti Bhat[3], Aiti Thapa[4], Aman Biswakarma[4],

Bharat Tamang[4], Binod Munda[4], Biren Biswakarma[4], Dema Tamang[4], Dambar

Pradhan[4], Mangal K Rai[4], Rohit Rai[4], Shambhu Rai[4] and Umesh Srinivasan[4]

[1]Department of Ecology and Environmental Sciences, Pondicherry University, India

[2]Conservation Biology Lab, University of Neuchâtel, Switzerland

[3]Organismic and Evolutionary Biology Lab, University of Massachusetts Amherst, USA

[4]Centre for Ecological Sciences, Indian Institute of Science, Bangalore, India

*Corresponding author: Satyam Gupta, Department of Ecology and Environmental Sciences,

Pondicherry University, India, Satmaysatyam@gmail.com, +919654560597

ABSTRACT

Mixed-species flocks (MSF) represent an important form of social organisation in bird

communities worldwide. Despite its likely importance in flock formation and cohesion, the

role of vocal communication in the formation and maintenance of MSF in birds is hitherto

understudied. In this study, we examine if a species' centrality within a mixed-species flock

is influenced by its vocal behaviour during the dawn chorus, i.e., the time of MSF formation.

Using acoustic sampling and field observations, we studied the bird species found in MSF in

the Eastern Himalayas. Our results show differential vocal activity patterns among MSF-

forming bird species and suggest a positive correlation between calling rates and closeness

centrality (species' importance in a flock) in understory MSFs. We also found a more

synchronised vocalisation pattern in the understory MSFs, with a consistent peak in vocal

activity in the early morning hours, whereas no consistent vocal pattern was found for canopy

flocks. Overall, our results suggest a potential mechanism that drives MSF formation wherein

the vocal activity of central species precedes and likely attracts participation from other

attendant species.

KEYWORDS: Animal Communication, Bioacoustics, Eastern Himalaya, Mixed-species

Flock, Network Analysis, Social Organization

INTRODUCTION

Multi-species associations are common in nature. Interspecific interactions consisting of

individuals of two or more species can aid in predator avoidance and enhance foraging

success (Morse 1977; Sridhar et al. 2009; Goodale et al. 2017). Mixed-species flocks

(hereafter, MSF or "flocks") are interspecific associations of two or more species,

representing an important feature of bird communities worldwide (Greenberg 2000; Harrison

and Whitehouse 2011).

MSF consist of species exhibiting a continuum of social cohesion within flocks, ranging from

loose aggregations such as those of herons and seabirds (Sealy 1973, Baltz 1977, Kushlan

1977) to highly structured winter foraging groups of chickadees (*Parus sp.*) (Hogstad 1978)

and multi-species territoriality in the Amazon (Jullien and Thiollay 1998). The formation and

maintenance of these flocks are influenced by a complex interplay of factors, including

species-specific ecological traits, foraging strategies, and social behaviours. While traditional

models categorized species into rigid roles such as 'nuclear' and 'satellite' (Winterbottom

1943; Powell 1985; Hutto 1994; Farley et al. 2008), more recent research has highlighted the

dynamic nature of these roles and the potential for significant behavioural flexibility in

response to changing environmental conditions and social interactions (Dolby & Grubb,

2000; Srinivasan et al. 2010, Borah et al. 2018).

While several studies have explored the composition of MSF, the vocal behaviour of its

constituent species remains understudied. Acoustic communication plays a pivotal role in

forming and maintaining MSF (Suzuki 2012). Bird vocalisations can serve multiple

functions, ranging from species recognition to maintenance of interactions and alarm calling

(Marler, 2004). Indeed, some bird species have been shown to produce specific calls to attract

others to join MSF (Diamond 1987, Goodale & Kotagama, 2006). In the context of MSF,

these distinct and recognisable calls can allow individuals to locate and join the flock. By

investigating the patterns in species-specific vocalizations, we can elucidate how these

signals contribute to flock formation, cohesion and information exchange (Goodale &

Kotagama, 2005).

In this study, we examined the role of vocalisation patterns in forming and maintaining MSF.

By examining temporal patterns in species-specific vocalizations, we aim to understand how

vocalisation could influence the role of each species in the MSF formation process. Our *a*

priori hypothesis was that species exhibiting higher calling rates would be more central in the

MSF networks than species with lower calling rates.

MATERIALS AND METHODS

STUDY AREA

We conducted this study in the Eaglenest Wildlife Sanctuary, located in the West Kameng

district of Arunachal Pradesh, India (27.07°N; 92.40°E), which is situated within the Eastern

Himalaya global biodiversity hotspot (Figure 1). We sampled MSF in the montane wet

temperate forest, ~2000 m above sea level. The forest is characterised by dominant tree

species such as *Quercus lamellosa*, *Alnus nipalensis*, and *Michelia doltsopa* in the canopy,

along with an understorey dominated by bamboo species (*Chimonobambusa sp.*; Srinivasan,

2019).

Figure 1: A map of the study area showing the location of Eaglenest Wildlife

Sanctuary, India.

ETHICAL NOTE

All procedures complied with the ASAB/ABS principles. Our study was confined to

noninvasive and non-disruptive behavioral observation techniques. Given the non-

invasive nature of our research, no animals were harmed, and it did not require review

from an institutional or governmental regulatory body. The Arunachal Forest

Department granted permission for the study under permit number

CWLW/GEN/998/2023/Pt-VIII(B)/3493-96.

SAMPLING METHODS

We used two sampling methods for data collection:

1. Mixed-Species Flock Observation

We observed MSF between 15[th] January and 11[th] March 2024 at Eaglenest Wildlife

Sanctuary near Bompu camp. An MSF is a single observation of a group of different species

of birds moving together over space and time (usually at least 5 minutes). We focused on two

stretches of a fairweather road, where observers monitored MSF daily (except during rain)

from 6:00 AM to 10:00 AM and 2:00 PM to 4:00 PM.

Mixed-species flocks were defined as groups of birds from different species foraging and

moving together within a short distance (<=5m) of each other. For each flock encountered,

we recorded the number of species, the number of individuals of each species, the flock type

(based on their habitat stratum), time, and location (recorded using a GARMIN Etrex 30x

GPS device) for each observed flock. We used a Sony ICD-PX240 MP3 recorder for data

collection, which proved more efficient than manual recording due to the dynamic nature of

MSF. Our observations were confined to road stretches for improved species identification.

To enhance the network analysis, we incorporated MSF data collected in 2023 at the same

site and during the same season, with the 2024 data.

2. Acoustic Sampling

2.1. Active Acoustic Sampling

We collected "active" acoustic data from mixed-species flocks passing through specific

locations along the stretch of Fairweather Road (detailed in Figure 2). Acoustic sampling was

done between January 15th and March 11th, 2024. Active sampling involved us being present

and adjusting the microphone's orientation to optimize recording as the flocks moved

(Britzke 2002). Our sampling took place from 6:00 AM to 10:00 AM each day. We had two

teams stationed on different roads near Bompu camp, using Audiomoth recorders (Hill et al.,

2019) for active acoustic sampling. At the same time as recording, observers documented

flock composition, time, and GPS location using a custom Epicollect5 form (Aanensen et al.,

2009) (Eaglenest Acoustic Sampling SAS).

Figure 2 Active Acoustic Flock Sampling Locations

2.2 Opportunistic Acoustic Sampling

We collected additional acoustic recordings opportunistically from MSF-participating bird

species by recording individual calling bouts from January 15th to March 15th, 2024. We

conducted this sampling daily from 6:00 AM to 10:00 AM Indian Standard Time (UTC

+5:30), alongside the mixed-species flock observation detailed above.

3.   Preliminary Results

We observed a total of 188 MSF during the study period, comprising 55 unique species. This

observational data was combined with the 2023 dataset of 280 MSF collected in the same

study site and methodology. In total, we had species-level flock composition data for 347

MSFs (280 in 2023 and 188 in 2024) comprising 68 unique species. In the active acoustic

sampling, we observed a total of 204 MSFs, comprising 40 unique species. Additionally, we

opportunistically collected 1804 unique individual vocalization bouts of birds participating in

MSFs.

ANALYSIS

1. Social Network Analysis

The MSF data were filtered to remove species encountered in less than ten observed flocks,

following recommendations from Farine & Whitehead (2015). We used a walk-run clustering

algorithm, implemented using the 'cluster' package (v1.3.8; Maechler et al., 2023) in R

statistical software (v4.3.2; R Core Team, 2023), which facilitates the classification of flocks

into distinct clusters based on species co-occurrences. The 'igraph' package (v2.1.3; Csardi &

Nepusz, 2006) in R statistical software (v4.3.2; R Core Team, 2023) was used to construct

species-level networks, where species were represented as nodes and co-occurrences as

edges. Additionally, the 'igraph' package (v2.1.3; Csardi & Nepusz, 2006) was also employed

to calculate the closeness centrality of each species in the MSF data.

We have used closeness centrality to evaluate the relative position of species in flock

networks, due to its suitability in traditional information-sharing property (Wey & Blumstein,

2010). The closeness centrality of a node (here, species) in a graph is the inverse of the

average shortest path distance from the node to other nodes in the graph (Wasserman &

Faust, 1994). It can be viewed as the efficiency of each node (species) in spreading

information to all other nodes (species) (Wasserman & Faust, 1994). The higher the closeness

centrality of a node, the shorter the average distance from the node to any other node, and

thus the better positioned the node is in potentially spreading information to other nodes

(Wasserman & Faust, 1994). All analysis was done in the R statistical software (v4.3.2; R

Core Team 2023).

2. Acoustic Analysis

Each MSF recording was segmented into one-minute files, and MSF species vocalizing

within these segments were manually annotated through visual inspection of spectrograms

and audio analysis, with corresponding marked timestamps. This process was systematically

repeated for all the observed MSF. Acoustic analysis was done using Raven Pro (v1.6.1;

Charif et al., 2010).

By combining opportunistic acoustic data with active sampling data, we were able to gather a

more comprehensive dataset for calculating the various vocal parameters of a species. This

integrated dataset included the species name, date, and time of vocalization for each

recording. We organised the vocalization data for each species into 30-minute intervals,

creating eight-time windows between 6:00 AM and 10:00 AM (6:00 - 6:30 A.M. as

Window#1). Subsequently, we calculated the number of calls in all windows and identified

the time window with the highest number of call counts as the "peak vocalization period" or

"peak vocalization category". We used the 'dplyr' package in R Statistical Software (v4.3.2; R

Core Team 2023) for data manipulation. We used the 'ggplot2' package (Wickham et al.,

2016) in R Statistical Software (v4.3.2; R Core Team 2023) to create heatmaps for visualising

vocal activity patterns.

3. Correlation Analysis

We analyzed the correlation between acoustic metrics and closeness centrality in two flock

types out of three: the canopy flock (12 species analyzed) and the understory flock (9

species). We didn't include the mid-story flock in the analysis since it consisted of only two

species overall. The flock types were defined based on the social network analysis and

aligned with prior natural history observations at the site.

We performed Pearson correlation tests in R Statistical Software (v4.3.2; R Core Team 2023)

to examine relationships between social network properties (namely, closeness centrality of a

species) and the vocal activity patterns (peak vocalisation period and mean call count). We

visualised the results using the 'ggplot2' package (Wickham et al., 2016).

RESULTS

1. Social Network Analysis

We analysed mixed-species flock data across 35 species observed during our data collection.

We generated a matrix showing which species tend to co-occur and used this information to

build the network at the species level.

Subsequently, using the network clustering algorithm, we identified three distinct types of

clusters (Figure 3), which corresponded to the MSFs as follows -

1. Understory Flock (12 Species):

Understory flocks consisted of flocks led by the Yellow-throated Fulvetta

(*Schoeniparus cinereus*) observed during field observation. Species belonging to this

flocktype are the Yellow-throated Fulvetta (*Schoeniparus cinereus*), Rufous-capped

Babbler (*Cyanoderma ruficeps*), Golden Babbler (*Cyanoderma chrysaeum*), Grey-

cheeked Warbler (*Phylloscopus poliogenys*) and other species. The full list of species

can be found in Table 1 (Supplementary Data).

2. Midstory Flock – 3 Species

Mid-story flocks were led by the Rusty-fronted Barwing (*Actinodura egertoni*)

observed during field observation. Species belonging to this flock type are Rusty-

fronted Barwing (*Actinodura egertoni*), Coral-billed Scimitar Babbler (*Pomatorhinus*

ferruginosus) and White-breasted Parrotbill (*Psittiparus ruficeps*).

3. Canopy Flock – 20 Species

Canopy flocks were the most species-rich flock types found in the study area, with

several flocks sometimes exceeding 20 species in number. They consisted of flocks

led by Yellow-cheeked Tit (*Parus spilonotus*) and Yellow-browed Tit (*Sylviparus*

modestus), as observed during field observation. Species belonging to this flock type

include the Lesser Racket-tailed Drongo (*Dicrurus remifer*), Red-tailed Minla (*Minla*

ignotincta), Ashy-throated Warbler (*Phylloscopus maculipennis*), Himalayan Cutia

(*Cutia nipalensis*) and others species. The full list of species can be found in Table 1

(Supplementary Data).

Figure 3: Species-level network visualising the co-occurrence of various species seen within mixed-species flocks. Each node represents a distinct bird species, and the edges (connected lines) between them represent relationships or interactions. The thickness of each edge represents the frequency of co-occurrence – a thicker edge signifies two species which are more commonly observed together. The network contains visible clusters of species (each "flock type"), and also shows strong connections between different types of flocks, indicating that while species may favour specific flock types (clusters), there are interactions across flock types.

2. *Vocal Activity Pattern*

The acoustic analysis revealed distinct vocal activity patterns among species within the

canopy and understory flocks (Figure 4). In the canopy flocks, species such as the Chestnut-

crowned Warbler and Green-tailed Sunbird showed peak calling activity between 6:30 A.M.

and 7:00 A.M. Other species, including the Ashy-throated Warbler and Himalayan Cutia,

displayed moderate call activity across time, indicating a broader distribution of vocalization.

In contrast, species such as the Lesser Racket-tailed Drongo (*Dicrurus remifer*) and Black-

throated Tit (*Aegithalos concinnus*) vocalised less. These observations indicate that the

canopy flock contains species with more diverse vocalization patterns compared with the

understory flock, with some species being more active at specific times while others remain

relatively inactive.

In contrast, understory MSF birds exhibited earlier peak vocal activity as compared to the

canopy MSF birds, with all the members vocalising highest between 6:30 A.M. and 7:00

A.M. The Yellow-throated Fulvetta and Grey-cheeked Warbler had particularly high call

counts during this period, while species like the Grey-headed Canary-Flycatcher (*Culicicapa*

ceylonensis) and Golden Babbler maintained moderate and consistent vocal activity

throughout the day. Species such as the Golden-breasted Fulvetta and White-throated Fantail

vocalised less. The understory flock generally followed a pattern of heightened morning

activity, with different species showing peak calling activity as compared to the canopy flock.

Figure 4: Vocal Activity Pattern: The heatmap displays the vocal activity patterns of different bird species in the canopy and understory flock types. The Y-axis represents species ordered in increasing closeness centrality values. The X-axis represents time in

263 **30-minute intervals from 6:00 A.M. to 10:00 A.M. Each row corresponds to a specific**

264 **bird species, and each column represents a 30-minute time interval. The colour intensity**

265 **within each cell corresponds to the number of calls observed for that species during a**

266 **specific vocal period**

267 **.**

268

269 *3. Correlation Analysis*

270

271 Our analysis revealed different results for each type of MSF.

272

273 *3.1 Canopy Flock*

274 Our analyses revealed a weak association between the peak vocalisation period, mean call

275 count and closeness centrality. Correlation results indicated that the peak vocalization period

276 explained a negligible amount of variance in closeness centrality ($R^2 = 0.07$, $p = 0.62$, F-

277 values $= 0.14$, df $= 10$). Similarly, mean call count exhibited a minimal relationship with

278 closeness centrality ($R^2 = 0.08$, $p = 0.79$, F-values $= 0.06$, df $= 7$).

279

280 *3.2 Understory Flock*

281 In contrast, our analysis revealed a positive correlation between the mean call count and the

282 closeness centrality of species within the understory flocks. ($R^2 = 0.73$; F-statistic $= 8.10$; $p =$

283 0.02) (Figure 5). However, the peak vocalization period was not tractable to statistical

284 analysis due to its homogeneity within the understory flock. This lack of variability precluded

285 its inclusion in the linear regression model. Furthermore, visual inspection (Suppl. Figure 4)

286 revealed no discernible relationship between peak vocalization period and closeness

287 centrality, suggesting that changes in the latter did not influence the former.

Figure 5: Correlation of Mean Call Count and Closeness Centrality. The Y-axis

illustrates the values of closeness centrality, whereas the X-axis displays the mean call

count over time. The plots indicate a positive correlation between closeness centrality

and mean call count for understory flocks, suggesting that species with higher call

counts are typically more central in understory flocks. This correlation is more

pronounced in the Understory Flock, while it does not achieve statistical significance in

the Canopy Flock.

DISCUSSION

The phenomenon of MSF in birds has long fascinated researchers (Bates 1859); however, the

mechanisms underpinning their formation and dynamics remain to be fully understood.

Communication between different species plays a pivotal role in orchestrating the assembly

and maintenance of these diverse avian congregations (Goodale et al., 2010).

Early research conducted by Munn (1985) and McDonald and Henderson (1977) provided

insights into the significance of vocalization in the formation of MSF. Their findings

indicated that certain bird species, such as titmice and antshrikes, produce loud calls in the

morning, serving as a signal to attract other species to gather and form a flock. Once

assembled, the vocalisation of the central species, often referred to as "leader species," seems

to reinforce flock cohesion and momentum (Bell, 1983; Diamond, 1987). These leaders

usually initiate vocalizations that act as cues for others to follow, thereby influencing the

structure and dynamics of the flock.

In addition, McDonald (1989) showcased the effectiveness of playback experiments and

vocal muting techniques in gaining insight into the role of vocalization in flock formation.

Monkkonen et al. (1996) undertook a thorough investigation on the impact of the territorial

song of the willow tit (*Parus montanus*) on the creation of loose mixed feeding groups during

the breeding season in Finland. Their observations of foraging aggregations among various

species of boreal forest birds revealed varying responses to the willow tit's song (Greenberg,

2000). They discovered that other boreal forest insectivores notably joined the willow tit in

these mixed feeding groups (Greenberg, 2000). However, the level of distinction in the effect

playback method was somewhat obscured due to experimental comparisons made with the

song of a North American tyrant flycatcher and a piece of classical music (Greenberg, 2000).

As a result, this study did not definitively uphold the idea that specific vocalizations are

essential for promoting the formation of tit flocks (Greenberg, 2000).

Our research aimed to explore the connection between a species' vocal behaviour during the

crucial phase of flock formation and its importance within a flock.Overall, we identified three

distinct types of flocks which are understory, canopy and midstory flocks in our study area,

consistent with previous findings (Bharadwaj et al., 2023). Using network analyses, we

identified several species, including the Yellow-bellied Fairy-fantail (*Chelidorhynx*

hypoxanthus), Ashy-throated Warbler (*Phylloscopus maculipennis*), Grey-cheeked Warbler

(*Phylloscopus poliogenys*), and White-throated Fantail (*Rhipidura albicollis*), that actively

participated in multiple flock types, indicating high interspecies interaction and potential

information sharing across different flock types. This dynamic interplay highlights the

complexity of avian social networks and emphasizes the importance of considering individual

species' roles within larger ecological communities.

We postulated that species having higher vocalization rates, particularly during the early

hours of the day when flock formation occurs are more important, or "central species". Our

hypothesis was based on the concept that central species play a crucial role in initiating and

maintaining group cohesion through vocal communication. To investigate this hypothesis, we

conducted field observations and acoustic recordings, meticulously documenting the vocal

behaviour of different bird species within mixed-species flocks and their social interactions.

Our findings revealed intriguing patterns in vocal activity among different flock types. Birds

in understory flocks, led by species such as the Yellow-throated Fulvetta (*Schoeniparus*

cinereus), exhibited synchronized vocalization patterns, with a consistent peak in vocal

activity during the early hours. However, we did not observe a clear pattern in canopy flock

types, suggesting potential variation in the role of vocalizations in influencing MSF

formation and cohesion.

Our study found that within the understory flock type, species with higher closeness

centrality (referred to as central species) exhibited higher mean call counts, signalling that

central species are more vocal than other flock members. These findings align with previous

research, such as Smith (1991), which noted that key species such as tits (*Parus*) consistently

vocalized within the flock. Similarly, in a tropical study, Wiley (1971) observed the dot-

winged antwren, a central species in Central American antwren flocks, continuously

vocalizing while moving between trees. This behaviour was used by other species to locate

flocks as they moved slowly through dense vegetation, maintaining flock cohesion.

Additionally, Munn (1985) in the Neotropical forests of Peru, discovered that the leader

species, the White-winged Shrike-tanager (*Lanio versicolor*), made chattering vocalizations,

prompting other flock members to join in. Therefore, the higher calling rate of central species

observed in our study may serve as a mechanism for maintaining the cohesion of other

species and forming the flock. Our results highlight the crucial role of key species in

initiating and maintaining flock cohesion through vocal communication.

Understanding the extent to which species exchange information with each other is crucial

for comprehending the structure of animal groups. Measuring the level of information flow

between species can offer valuable insights into the dynamics of species interactions that

underpin group dynamics and may also indicate the extent to which certain species play a

disproportionate role in maintaining group coherence.

In our research, while our primary focus was on the vocalization patterns and their correlation

with the significance of species in mixed-species foraging (MSF) flocks, integrating metrics

of information exchange between species could add further depth to our analysis. By

quantifying the information exchange both within and between species, we can gain a better

understanding of the communication dynamics within MSF, which in turn can provide

valuable insights into the processes driving the formation and stability of these flocks.

Additionally, in terms of conservation, evaluating the symmetry of information flow can aid

in identifying species that have key roles in maintaining group cohesion (Martinez, 2013).

Previous studies have noted the absence of flock formation in the absence of certain key

species (Cody & Diamond, 1975; Maldonado-Coelho and Marini, 2000). Species that engage

in high levels of information exchange with others can act as keystone species within multi-

species flocks, influencing the behaviour and spatial dynamics of other members (Martinez,

2013). Therefore, prioritizing the conservation of these species is crucial for preserving the

composition and functionality of avian communities (Martinez, 2013).

Although our study provides valuable insights into the formation of MSF and avian

behaviour, it is important to recognise its limitations. Our observations were based on two

years of observing flocks and two months of collecting acoustic samples, emphasizing the

necessity for long-term data to strengthen the reliability of our findings. Future research

should involve longer sampling periods and investigate the functions of specific vocalizations

or the vocalizations of particular species, potentially through playback experiments or vocal

muting experiments (McDonald 1989).

By incorporating measures of information flow into future studies on MSF formation and

dynamics, we can gain valuable insights into the underlying mechanisms of avian social

behaviour and contribute to more effective conservation strategies aimed at preserving

biodiversity and ecosystem stability. In summary, our study adds to the expanding knowledge

of MSF formation and underscores the importance of key species in driving flock dynamics.

Understanding the complex interactions among individuals and species within different

habitats is critical for the effective conservation and maintenance of community structure.

Further exploration of the vocalizations of MSF birds and their influence on flock behaviour

holds the promise of unravelling the intricacies of this fascinating ecological phenomenon,

paving the way for more comprehensive conservation strategies.

CONFLICT OF INTEREST
410
411   The authors have declared that no competing interests exist.
412

REFERENCES

1. Aanensen DM, Huntley D, Feil EJ, al-Own F, & Spratt BG. (2009). EpiCollect:

linking smartphones to web applications for epidemiology, ecology and community

data collection. *PLoS ONE*, *4*(9).

2. Baltz, D. M., & Morejohn, G. V. (1977). Food Habits and Niche Overlap of Seabirds

Wintering on Monterey Bay, California. *The Auk*, *94*(3), 526–543.

https://doi.org/10.1093/AUK/94.3.526

3. Bell, H. L. (1983). A Bird Community of Lowland Rainforest in New Guinea. 5.

Mixed-Species Feeding Flocks. *Emu - Austral Ornithology*, *82*(5), 256–275.

https://doi.org/10.1071/MU9820256S

4. Bharadwaj, A., Thapa, A., Bhat, A., Biswakarma, A., Tamang, B., Munda, B.,

Biswakarma, B., Pradhan, D. K., Tamang, D., & Pradhan, K. (2023). Understanding

inter-individual social networks in mixed-species bird flocks. *BioRxiv*, 2023–2024.

5. Billerman, S. M., Keeney, B. K., Rodewald, P. G., & Schulenberg, T. S. (2022). Birds

of the World. *Cornell Laboratory of Ornithology*.

6. Britzke, E. R. (2004). Designing monitoring programs using frequency-division bat

detectors: active versus passive sampling. *Bat Echolocation Research: Tools,*

Techniques and Analysis, 79.

7. Charif, R., AM Waack, & LM Strickman. (2010). *Raven Pro 1.4 User's Manual.*

8. Cody, M. L., & Diamond, J. M. (1975). *Ecology and evolution of communities.*

Harvard University Press.

9. Csardi, G., & Nepusz, T. (2006). The Igraph Software Package for Complex Network

Research. *InterJournal, Complex Systems*, *1695*(5), 1–9.

https://www.researchgate.net/publication/221995787

10. Diamond, J. (1987). Flocks of Brown and Black New Guinean Bird: A Bicolored

Mixed-species Foraging Association. *Emu*, *87*(4), 201–211.

https://doi.org/10.1071/MU9870201

11. Dolby, A. S., & Grubb, T. C. (2000). Social context affects risk taking by a satellite

species in a mixed-species foraging group. *Behavioral Ecology*, *11*(1), 110–114.

https://doi.org/10.1093/BEHECO/11.1.110

12. Farine, D. R., & Whitehead, H. (2015). Constructing, conducting and interpreting

animal social network analysis. *Journal of Animal Ecology*, *84*(5), 1144–1163.

https://doi.org/10.1111/1365-2656.12418

13. Farley, E. A., Sieving, K. E., & Contreras, T. A. (2008). Characterizing complex

mixed-species bird flocks using an objective method for determining species

participation. *Journal of Ornithology*, *149*(3), 451–468.

https://doi.org/10.1007/S10336-008-0284-Z/TABLES/3

14. Goodale, E., & Beauchamp, G. (2010). The relationship between leadership and

gregariousness in mixed-species bird flocks. *Journal of Avian Biology*, *41*(1), 99–103.

https://doi.org/10.1111/j.1600-048X.2009.04828.x

15. Goodale, E., Beauchamp, G., Magrath, R. D., Nieh, J. C., & Ruxton, G. D. (2010).

Interspecific information transfer influences animal community structure. *Trends in*

Ecology & Evolution, *25*(6), 354–361. https://doi.org/10.1016/J.TREE.2010.01.002

16. Goodale, E., Beauchamp, G., & Ruxton, G. D. (2017). Adaptive Implications of

Mixed-Species Grouping: Foraging, Physical, and Reproductive Factors. *Mixed-*

Species Groups of Animals, 59–80. https://doi.org/https://doi.org/10.1016/B978-0-12-

805355-3.00004-X

17. Goodale, E., & Kotagama, S. W. (2006). Vocal mimicry by a passerine bird attracts

other species involved in mixed-species flocks. *Animal Behaviour*, *72*(2), 471–477.

https://doi.org/10.1016/j.anbehav.2006.02.004

18. Goodale, E., & Kotagama W., S. (2005). Alarm Calling in Sri Lankan Mixed-Species

Bird Flocks. *The Auk*, *122*(1), 108–120.

https://academic.oup.com/auk/article/122/1/108/5562418

19. Greenberg, R. (2000). Birds of Many Feathers: The Formation and Structure of

Mixed-Species Flocks of Forest Birds. In S. Boinski & P. Garber (Eds.), *On the move:*

how and why animals travel in groups (pp. 521–558). University of Chicago Press.

20. Harrison, N. M., & Whitehouse, M. J. (2011). Mixed-species flocks: An example of

niche construction? *Animal Behaviour*, *81*(4), 675–682.

https://doi.org/10.1016/j.anbehav.2011.01.013

21. Hill, A. P., Prince, P., Snaddon, J. L., Doncaster, C. P., & Rogers, A. (2019).

AudioMoth: A low-cost acoustic device for monitoring biodiversity and the

environment. *HardwareX*, *6*, e00073.

https://doi.org/https://doi.org/10.1016/j.ohx.2019.e00073

22. Hogstad, O. (1978). Sexual Dimorphism in Relation to winter foraging and Territorial

Behaviour of three-toed woodpecker (Picoides tridactylus) and three Dendroccops

species. *Ibis*, *120*(2), 198–203. https://doi.org/10.1111/J.1474-919X.1978.TB06775.X

23. Hutto, R. L. (1994). The Composition and Social Organization of Mixed-Species

Flocks in a Tropical Deciduous Forest in Western Mexico. *The Condor*, *96*(1), 105–

118. https://doi.org/10.2307/1369068

24. Jullien, M., & Thiollay, J.-M. (1998). Multi-Species Territoriality and Dynamic of

Neotropical Forest Understorey Bird Flocks. *Journal of Animal Ecology*, *67*(2), 227–

252.

25. Kushlan, J. A. (1977). The Significance of Plumage Colour in the Formation of

Feeding Aggregations of Ciconiiformes. *Ibis*, *119*(3), 361–364.

https://doi.org/10.1111/J.1474-919X.1977.TB08256.X

26. Macdonald, D. W., & Henderson, D. G. (1977). Aspects of the Behaviour And

Ecology of Mixed-Species Bird Flocks in Kashmir. *Ibis*, *119*(4), 481–493.

https://doi.org/10.1111/j.1474-919X.1977.tb02055.x

27. Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M., & Hornik, K. (2023). *cluster:*

Cluster Analysis Basics and Extensions.

28. Maldonado-Coelho, M., & Marini, M. A. ˆ. (2000). Effects Of Forest Fragment Size

And Successional Stage On Mixed-Species Bird Flocks In Southeastern Brazil. *The*

Condor, *120*, 585–594. https://academic.oup.com/condor/article/102/3/585/5562985

29. Marler, P. (2004). Bird Calls: Their Potential for Behavioral Neurobiology. *Annals of*

the New York Academy of Sciences, *1016*(1), 31–44.

https://doi.org/10.1196/annals.1298.034

30. Martinez, A. E. (2013). *Interspecific Communication And its Consequences for the*

Organization of Permanent Mixed-Species Bird Flocks.

31. McDonald, M. V. (1989). Function of song in Scott's seaside sparrow, Ammodramus

maritimus peninsulae. *Animal Behaviour*, *38*(3), 468–485.

https://doi.org/https://doi.org/10.1016/S0003-3472(89)80040-5

32. Mönkkönen, M., Forsman, J., Helle, P., & Monkkonen, M. (1996). Mixed-Species

Foraging Aggregations and Heterospecific Attraction in Boreal Bird Communities.

Oikos, *77*, 127. https://doi.org/10.2307/3545592

33. Morse, D. H. (1970). Ecological Aspects of Some Mixed-Species Foraging Flocks of

Birds. *Ecological Monographs*, *40*(1), 119–168. https://doi.org/10.2307/1942443

34. Morse, D. H. (1977). Feeding Behavior and Predator Avoidance in Heterospecific

Groups. *BioScience*, *27*(5), 332–339. https://doi.org/10.2307/1297632

35. Munn, C. A. (1985). Permanent Canopy and Understory Flocks in Amazonia: Species

Composition and Population Density. *Ornithological Monographs*, *36*, 683–712.

https://doi.org/10.2307/40168312

36. Powell, G. V. N. (1985). Sociobiology and Adaptive Significance of Interspecific

Foraging Flocks in the Neotropics. *Ornithological Monographs*, *36*, 713–732.

https://doi.org/10.2307/40168313

37. R Core Team. (2023). R: A Language and Environment for Statistical Computing. *R*

Foundation for Statistical Computing, Vienna, Austria.

38. Sealy, S. (1973). Interspecific Feeding Assemblages of Marine Birds off British

Columbia. *The Auk*, *90*, 796–802. https://doi.org/10.2307/4084359

39. Smith, S. M. (1991). The Black-Capped Chickadee. In *Behavioral Ecology and*

Natural History. Cornell University Press.

40. Sridhar, H., Beauchamp, G., & Shanker, K. (2009). Why do birds participate in

mixed-species foraging flocks? A large-scale synthesis. *Animal Behaviour*, *78*(2),

337–347. https://doi.org/10.1016/j.anbehav.2009.05.008

41. Srinivasan, U. (2019). Morphological and behavioral correlates of long-term bird

survival in selectively logged forest. *Frontiers in Ecology and Evolution*, *7*(FEB).

https://doi.org/10.3389/fevo.2019.00017

42. Srinivasan, U., Raza, R. H., & Quader, S. (2010). The nuclear question: rethinking

species importance in multi-species animal groups. *Journal of Animal Ecology*, *79*(5),

948–954. https://doi.org/10.1111/J.1365-2656.2010.01707.X

43. Suzuki, T. N. (2012). Long-distance calling by the willow tit, Poecile montanus,

facilitates formation of mixed-species foraging flocks. *Ethology*, *118*(1), 10–16.

https://doi.org/10.1111/j.1439-0310.2011.01982.x

44. Wasserman, S., & Faust, K. (1994). *Social Network Analysis*. Cambridge University

Press. https://doi.org/10.1017/CBO9780511815478

45. Wey, T.-W., & Blumstein, D. T. (2010). Social network analysis of animal behaviour:

a promising tool for the study of sociality. Animal Behaviour, 79(2), 277-286.

46. Wickham, H., Chang, W., & Wickham, M. H. (2016). Package 'ggplot2.' *Create*

Elegant Data Visualisations Using the Grammar of Graphics. Version, *2*(1), 1–189.

47. Wickham, H., Francois, R., Muller, K., & Vaughan, D. (2023). dplyr: A Grammar of

Data Manipulation. In *R package version 1.1.4, https://github.com/tidyverse/dplyr,*

https://dplyr.tidyverse.org.

48. Wiley, R. H. (1971). Cooperative Roles in Mixed Flocks of Antwrens

(Formicariidae). *The Auk*, *88*(4), 881–892. https://doi.org/10.2307/4083845

49. Winterbottom., J. M. (1943). On Woodland Bird Parties in Northern Rhodesia. *Ibis*,

85(4), 437–442. https://doi.org/10.1111/J.1474-919X.1943.TB03857.

557 **Supplementary Material**

Sr. No.	Species	FlockType
1	Ashy-throated Warbler	Canopy Flock
2	Black-headed Shrike Babbler	Canopy Flock
3	Black-throated Tit	Canopy Flock
4	Buff-barred Warbler	Canopy Flock
5	Chestnut-crowned Warbler	Canopy Flock
6	Chestnut-tailed Minla	Canopy Flock
7	Green-tailed Sunbird	Canopy Flock
8	Grey-chinned Minivet	Canopy Flock
9	Grey-hooded Warbler	Canopy Flock
10	Himalayan Cutia	Canopy Flock
11	Lesser Racket-tailed Drongo	Canopy Flock
12	Red-tailed Minla	Canopy Flock
13	Sikkim Treecreeper	Canopy Flock
14	Yellow-browed Tit	Canopy Flock
15	Yellow-cheeked Tit	Canopy Flock
16	Coral-billed Scimitar Babbler	Mid-story Flock
17	Rusty-fronted Barwing	Mid-story Flock
18	White-breasted Parrotbill	Mid-story Flock
19	Black-faced Warbler	Understory Flock
20	Black-throated Parrotbill	Understory Flock
21	Broad-billed Warbler	Understory Flock
22	Golden-breasted Fulvetta	Understory Flock
23	Golden Babbler	Understory Flock
24	Grey-cheeked Warbler	Understory Flock
25	Grey-headed Canary-flycatcher	Understory Flock

26	Rufous-capped Babbler	Understory Flock
27	Rufous-winged Fulvetta	Understory Flock
28	White-throated Fantail	Understory Flock
29	Yellow-bellied Fairy-fantail	Understory Flock
30	Yellow-throated Fulvetta	Understory Flock

Table 1 MSF participating birds in different flock types

Figure 1 MSF Observation Sampling Locations

568   **Figure 2 Opportunistic Acoustic Sampling Locations**

Figure 3 Distribution of Closeness Centrality in the MSF

Figure 4 Correlation of Peak Vocalisation Period and Closeness Centrality

The Y-axis illustrates the values of closeness centrality, whereas the X-axis displays the peak

vocalisation period. The plots indicate a positive correlation between closeness centrality and

576   peak vocalization period for canopy flocks, although this relationship was insignificant. A

577   correlation could not be established for the understory flock, as all variables on the X-axis

578   were identical, indicating a synchronised peak vocalisation between the different species.

579