Vocal species are more central in Eastern Himalayan Mixed-Species bird flocks

Satyam Gupta[1*], Akshay Bharadwaj[2], Akshiti Bhat[3], Aiti Thapa[4], Aman Biswakarma[4], Bharat Tamang[4], Binod Munda[4], Biren Biswakarma[4], Dema Tamang[4], Dambar Pradhan[4], Mangal K Rai[4], Rohit Rai[4], Shambh Rai[4]and Umesh Srinivasan[4]

[1]Department of Ecology and Environmental Sciences, Pondicherry University, India

[2]Conservation Biology Lab, University of Neuchâtel, Switzerland

[3]Organismic and Evolutionary Biology Lab, University of Massachusetts Amherst, USA

[4]Centre for Ecological Sciences, Indian Institute of Sciences, Bangalore, India

*Corresponding author: Satmaysatyam@gmail.com

ABSTRACT

Mixed-species flocks (MSF) represent an important form of social organisation in bird communities worldwide. Despite its likely importance in flock formation and cohesion, the role of vocal communication in the formation and maintenance of MSF in birds is hitherto understudied. In this study, we examine if a species' centrality within a mixed-species flock is influenced by its vocal behaviour during the dawn chorus, i.e., the time of MSF formation. Using acoustic sampling and field observation, we studied bird species found in MSF in the Eastern Himalayas. Our results show differential vocal activity patterns among MSF-forming bird species and suggest a positive correlation between calling rates and closeness centrality in understory MSFs. We also found a more synchronised vocalisation pattern in the understory MSFs, with a consistent peak in vocal activity in the early morning hours, whereas no consistent vocal pattern was found for canopy flocks. Overall, our results suggest a potential mechanism that drives MSF formation wherein the vocal activity of central species precedes and likely attracts participation from other attendant species.

KEYWORDS: Animal Communication, Bioacoustics, Eastern Himalaya, Mixed-species Flock, Network Analysis, Social Organization

INTRODUCTION

Multi-species associations are common in nature. Interspecific interactions consisting of individuals of two or more species can aid in predator avoidance and enhance foraging success (Morse 1977; Sridhar et al. 2009; Goodale et al. 2017). Mixed-species flocks (hereafter, MSF or "flocks") are non-trophic interspecific associations of two or more species, representing an important feature of bird communities worldwide (Greenberg 2000; Harrison and Whitehouse 2011).

MSF consist of species exhibiting a continuum of social cohesion within flocks, ranging from loose aggregations such as those of herons and seabirds (Sealy 1973, Baltz 1977, Kushlan 1977) to highly structured winter foraging groups of chickadees (*Parus sp.*) (Hogstad 1978) and multi-species territoriality in the Amazon (Jullien and Thiollay 1998). The formation and maintenance of these flocks are influenced by a complex interplay of factors, including species-specific ecological traits, foraging strategies, and social behaviours. While traditional models categorized species into rigid roles such as 'nuclear' and 'satellite' (Winterbottom 1943; Powell 1985; Hutto 1994; Farley et al. 2008), more recent research has highlighted the dynamic nature of these roles and the potential for significant behavioural flexibility in response to changing environmental conditions and social interactions (Dolby & Grubb, 2000; Srinivasan et al. 2010, Borah et al. 2018).

While several studies have explored the composition of MSF, the vocal behaviour of its constituent species remains understudied. Nevertheless, this is crucial for a mechanism that facilitates the formation and coordination within MSF. Acoustic communication plays a pivotal role in forming and maintaining MSF (Suzuki 2012). Bird vocalisations can serve multiple functions, ranging from species recognition to maintenance of interactions and alarm calling (Marler, 2004). Indeed, some bird species have been shown to produce specific calls to attract others to join MSF (Diamond 1987, Goodale & Kotagama, 2006). In the context of MSF, these distinct and recognisable calls can allow individuals to locate and join the flock. By investigating the patterns in species-specific vocalizations, we can elucidate how these signals contribute to flock formation, cohesion and information exchange (Goodale & Kotagama, 2005).

In this study, we examined the role of vocalisation patterns in forming and maintaining MSF. By examining temporal patterns in species-specific vocalizations, we aim to understand how

vocalisation could influence the role of each species in the MSF formation process. Our *a*

priori hypothesis was that species exhibiting higher calling rates would be more central in the

MSF networks than species with lower calling rates.

Material and Methods

STUDY AREA

We conducted this study in the Eaglenest Wildlife Sanctuary, West Kameng district,

Arunachal Pradesh, India (27.07°N; 92.40°E), which lies in the Eastern Himalaya global

biodiversity hotspot (Fig. 1). We sampled MSF in the montane wet temperate forest, ~2000 m

above sea level. The forest is characterised by dominant tree species such as *Quercus*

lamellosa, *Alnus nipalensis*, and *Michelia doltsopa* in the canopy, alongside bamboo species

(*Chimonobambusa sp.*) in the understorey (Srinivasan, 2019).

Figure 1: A map of the study area showing the location of Eaglenest Wildlife

Sanctuary, India.

SAMPLING METHODS

We used two sampling methods for data collection:

1. Mixed-Species Flock Observation

We observed MSF between 15[th] January and 11[th] March 2024 at Eaglenest Wildlife Sanctuary near Bompu camp. An MSF is a single observation of a group of different species of birds moving together over space and time (usually at least 5 minutes). We focused on two stretches of a fairweather road, where observers monitored MSF daily (except during rain) from 6:00 AM to 10:00 AM and 2:00 PM to 4:00 PM. Mixed-species flocks were defined as groups of birds from different species foraging and moving together within a short distance of each other. For each flock encountered, we recorded the number of species, number of individuals of each species, flock type (based on their habitat stratum), time, and location (recorded using a GARMIN Etrex 30x GPS device) for each observed flock. We used a Sony ICD-PX240 MP3 recorder for data collection, which proved more efficient than manual recording due to the dynamic nature of MSF. Our observations were confined to road stretches for improved species identification. We transcribed the collected data for subsequent analysis. To enhance the network analysis, we incorporated MSF data collected in 2023 at the same site and during the same season, with the 2024 data.

2. Acoustic Sampling

2.1. Active Acoustic Sampling

We collected "active" acoustic data from mixed-species flocks passing through a specific location between January 15th and March 11th, 2024. Active sampling involved us being present and adjusting the microphone's orientation to optimize recording as the flocks moved (Britzke 2002). Our sampling took place from 6:00 AM to 10:00 AM each day. We had two teams stationed on different roads near Bompu camp, using Audiomoth recorders (Hill et al., 2019) for active acoustic sampling. At the same time as recording, observers documented flock composition, time, and GPS location using a custom Epicollect5 form (Aanensen et al., 2009) (Eaglenest Acoustic Sampling SAS).

2.2 Opportunistic Acoustic Sampling

We collected acoustic calls opportunistically from MSF-participating bird species by recording individual call records from January 15th to March 15th, 2024. We conducted this sampling daily from 6:00 AM to 10:00 AM alongside active acoustic sampling and mixed-species flock observation. The purpose of this sampling method was to gather acoustic data without a dedicated, planned effort by recording the vocalizations of MSF birds when they were observed outside the MSF. We used a customized Epicollect5 form (Aanensen et al., 2009) (Opportunistic Acoustic Sampling OAS) to record species name, observation time, date, and location coordinates for each vocalization.

3. Preliminary Results

We observed a total of 188 MSF during the study period, comprising 55 unique species. This observational data was combined with the 2023 dataset of 280 MSF collected in the same location. After combining, we had species-level information for 347 MSF (188, 2024 & 280, 2023) comprising 68 unique species. Upon filtering out rare species in MSF, we then considered 35 MSF-participating species.

In the active acoustic sampling, we observed and analyzed a total of 204 MSF. This resulted in data for 40 unique species. During opportunistic acoustic sampling, we recorded 1804 unique individual vocalization records of birds participating in MSF.

ANALYSIS

1. Social Network Analysis

The MSF data were filtered to remove species encountered in less than ten observed flocks, following recommendations from Farine & Whitehead (2015). We used a walk-run algorithm clustering, implemented using the 'cluster' package (Maechler et al., 2023), which facilitated the classification of flocks into distinct clusters based on species co-occurrences. The 'igraph' package (Csardi & Nepusz, 2006) was used to construct species-level networks, where species were represented as nodes and co-occurrences as edges. Additionally, *igraph* (Csardi & Nepusz, 2006) was also employed to calculate the closeness centrality of each species in the MSF data.

The closeness centrality of a node (here, species) in a graph is the inverse of the average shortest path distance from the node to other nodes in the graph (Wasserman & Faust, 1994).

It can be viewed as the efficiency of each node (species) in spreading information to all other nodes (species) (Wasserman & Faust, 1994). The higher the closeness centrality of a node, the shorter the average distance from the node to any other node, and thus the better positioned the node is in potentially spreading information to other nodes (Wasserman & Faust, 1994). Therefore, because of the information-sharing property, we used closeness centrality as the metric to evaluate the relative position of species in flock networks. All analysis was done in the R statistical software (v4.3.2; R Core Team 2023).

2. Acoustic Analysis

Each MSF recording was segmented into one-minute files and MSF species vocalizing within these segments were manually annotated through visual inspection of spectrograms and audio analysis, with corresponding marked timestamps. This process was systematically repeated for all the observed MSF. Acoustic analysis was done using Raven Pro (v1.6.1; Charif et al., 2010).

By combining opportunistic acoustic data with active sampling data, we were able to gather a more comprehensive dataset for calculating the various vocal parameters of a species. This integrated dataset included the species name, date, and time of vocalization for each recording. We organized the vocalization data for each species into 30-minute intervals, creating eight-time windows between 6:00 AM and 10:00 AM (6:00 - 6:30 A.M. as Window#1). Subsequently, we calculated the number of calls in all windows and identified the time window with the highest number of call counts as the "peak vocalization period". We used the 'dplyr' package in R Statistical Software (v4.3.2; R Core Team 2023) to analyze the data. We used the 'ggplot2' package (Wickham et al., 2016) in R Statistical Software (v4.3.2; R Core Team 2023) to create heatmaps for visualising vocal activity patterns.

3. Correlation Analysis

We conducted Pearson correlation tests in R Statistical Software (v4.3.2; R Core Team 2023). We used the correlations to examine relationships between the dependent variable closeness centrality and the independent variable's peak vocalisation category and Mean call count. We constructed correlation plots using the 'ggplot2' package (Wickham et al., 2016).

We analyzed the correlation between acoustic metrics and closeness centrality in two flock types out of three: the canopy flock, (12 species analyzed), and the understory flock, (9

species). We didn't include the mid-story flock in the analysis since it had only two species, which was insufficient for correlation analysis.

RESULTS

1. Social Network Analysis

We used data from 35 species observed in the flock and created a social network based on their interactions. We generated a matrix showing which species tend to co-occur and used this information to build the network at the species level.

Subsequently, using the network clustering algorithm we identified three distinct types of clusters, which corresponded to the MSFsas follows -

1. Understory Flock – 12 Species
2. Midstory Flock – 3 Species
3. Canopy Flock – 20 Species

1.1 **Understory flocks** consisted of flocks led by the Yellow-throated Fulvetta (*Schoeniparus cinereus*). Species belonging to this flocktype are the Yellow-throated Fulvetta (*Schoeniparus cinereus*), Rufous-capped Babbler (*Cyanoderma ruficeps*), Golden Babbler (*Cyanoderma chrysaeum*) and Grey-cheeked Warbler (*Phylloscopus poliogenys*) and other species (Fig. 2).

Figure 2: Species found in understory flock type (clockwise): Grey-cheeked Warbler, White-throated Fantail, Yellow-bellied Fairy-fantail, Yellow-throated Fulvetta, Golden Babbler, Rufous-capped Babbler, Black-faced Warbler, Golden-breasted Fulvetta. Images: Satyam Gupta

1.2. **Mid-story flocks** were led by the Rusty-fronted Barwing (*Actinodura egertoni*). Species belonging to this flock type are Rusty-fronted Barwing (*Actinodura egertoni*), Coral-billed Scimitar Babbler (*Pomatorhinus ferruginosus*) and White-breasted Parrotbill (*Psittiparus ruficeps*) (Fig. 3).

Figure 3: Species found in Midstory flock type (left to right): Rusty-fronted Barwing, White-breasted Parrotbill, Coral-billed Scimitar Babbler. Images: Satyam Gupta

1.3. **Canopy flocks** were the most-species-rich flock types found in the study area, with several flocks sometimes exceeding 20 species in number. They consisted of flocks led by Yellow-cheeked Tit (*Parus spilonotus*) and Yellow-browed Tit (*Sylviparus modestus*). Species belonging to this flock type are Lesser Racket-tailed Drongo (*Dicrurus remifer*), Red-tailed Minla (*Minla ignotincta*), Ashy-throated Warbler (*Phylloscopus maculipennis*), Himalayan Cutia (*Cutia nipalensis*) and many others species (Fig. 4).

Figure 4 Species found in canopy flock type (clockwise): Himalayan Cutia,
Grey-chinned Minivet, Chestnut-tailed Minla, Red-tailed Minla, Chestnut-crowned
Warbler, Black-headed Shrike-Babbler, Sikkim Treecreeper, Rufous-bellied
Woodpecker, Black-throated Sunbird (centre). Images: Satyam Gupta

Figure 5: Species-level network representation of co-occurrence of various species seen within mixed-species flocks. Each node represents a distinct bird species, and the edges between them denote relationships or interactions. The thickness of each edge represents the frequency of co-occurrence – a thicker edge signifies two species which are more commonly observed together. The network reveals visible clusters of species within each flock type and also shows strong connections between different types of flocks, particularly between the canopy and understory flocks. This shows that while species may favour specific flock types (clusters), there are interactions across flock types.

2. Vocal Activity Pattern

The acoustic analysis revealed distinct vocal activity patterns among species within the canopy and understory flocks (Fig 6). In the canopy flocks, species such as the Chestnut-crowned Warbler and Green-tailed Sunbird showed peak calling activity between 6:30 A.M. and 7:00 A.M. Other species, including the Ashy-throated Warbler and Himalayan Cutia, displayed moderate call activity across time, indicating a broader distribution of vocalization. In contrast, species such as the Lesser Racket-tailed Drongo (*Dicrurus remifer*) and Black-throated Tit (*Aegithalos concinnus*) vocalised less. These observations indicate that the canopy flock contains species with more diverse vocalization patterns compared with the understory flock, with some species being more active at specific times while others remain relatively inactive.

In contrast, understory MSF birds exhibited earlier peak vocal activity as compared to the canopy MSF birds, with all the members vocalising highest between 6:30 A.M. and 7:00 A.M. The Yellow-throated Fulvetta and Grey-cheeked Warbler had particularly high call counts during this period, while species like the Grey-headed Canary-Flycatcher (*Culicicapa ceylonensis*) and Golden Babbler maintained moderate and consistent vocal activity throughout the day. Species such as the Golden-breasted Fulvetta and White-throated Fantail vocalised less. The understory flock generally followed a pattern of heightened morning activity, with different species showing peak calling activity as compared to the canopy flock.

Figure 6: Vocal Activity Pattern The heatmap displays the vocal activity patterns of different bird species in the canopy and understory flock types. The Y-axis represents species ordered in increasing closeness centrality values. In contrast, the X-axis represents time in 30-minute intervals from 6:00 A.M. to 10:00 A.M. Each row

corresponds to a specific bird species, and each column represents a 30-minute time interval. The colour intensity within each cell corresponds to the number of calls observed for that species during a specific vocal category. Lighter colours indicate a higher call count, while darker colours represent fewer calls.

3. Correlation Analysis

Our analysis revealed different results for each type of MSF.

3.1 Canopy Flock

Our analyses revealed a weak association between the peak vocalisation category, mean call count and closeness centrality. Linear regression models indicated that the peak vocalization category explained a negligible amount of variance in closeness centrality ($R^2 = 0.076$, p = 0.620, F-values =0.144, df = 10). Similarly, mean call count exhibited a minimal relationship with closeness centrality ($R^2 = 0.083$, p = 0.798, F-values =0.069, df = 7).

3.2 Understory Flock

In contrast, we detected a positive association between the mean call count and closeness centrality of species within the understory flocks ($R^2 = 0.73$; F-statistic = 8.10; p = 0.025) (Fig 7). However, the peak vocalization category was not tractable to statistical analysis due to its homogeneity within the understory flock. This lack of variability precluded its inclusion in the linear regression model. Furthermore, visual inspection (Suppl. Fig 5) revealed no discernible relationship between peak vocalization category and closeness centrality, suggesting that changes in the latter did not influence the former.

Figure 7: Correlation of Mean Call Count and Closeness Centrality. The Y-axis illustrates the values of closeness centrality, whereas the X-axis displays the mean call count over time. The plots indicate a positive correlation between closeness centrality and mean call count for both types of flocks, suggesting that species with higher call

**counts are typically more central. This correlation is more pronounced in the
Understory Flock, while it does not achieve statistical significance in the Canopy Flock.**

DISCUSSION

The phenomenon of MSF in birds has long fascinated researchers (Bates 1859); however, the mechanisms underpinning their formation and dynamics remain to be fully understood. Despite significant advances in our understanding, particularly regarding flocking benefits and participation, there is still much to uncover about how birds come together in mixed-species groups daily and how they are maintained throughout the day. Communication between different species plays a pivotal role in orchestrating the assembly and maintenance of these diverse avian congregations (Goodale et al., 2010).

Early research conducted by Munn (1985) and McDonald and Henderson (1977) provided insights into the significance of vocalization in the formation of MSF. Their findings indicated that certain bird species, such as titmice and antshrikes, produce loud calls in the morning, serving as a signal to attract other species to gather and form a flock. Once assembled, the vocalisation of the central species, often referred to as "leader species," seems to reinforce flock cohesion and momentum (Bell, 1983; Diamond, 1987). These leaders usually initiate vocalizations that act as cues for others to follow, thereby influencing the structure and dynamics of the flock.

In addition, McDonald (1989) showcased the effectiveness of playback experiments and vocal muting techniques in gaining insight into the role of vocalization in flock formation. Monkkonen et al. (1996) undertook a thorough investigation on the impact of the territorial song of the willow tit (*Parus montanus*) on the creation of loose mixed feeding groups during the breeding season in Finland. Their observations of foraging aggregations among various species of boreal forest birds revealed varying responses to the willow tit's song (Greenberg, 2000). They discovered that other boreal forest insectivores notably joined the willow tit in these mixed feeding groups (Greenberg, 2000). However, the level of distinction in the effect playback method was somewhat obscured due to experimental comparisons made with the song of a North American tyrant flycatcher and a piece of classical music (Greenberg, 2000).

As a result, this study did not definitively uphold the idea that specific vocalizations are essential for promoting the formation of tit flocks (Greenberg, 2000).

Our research aimed to explore the connection between a species' vocal behaviour during the crucial phase of flock formation and its importance within a flock. Using network analysis clustering, we identified three distinct types of flocks which are understory, canopy and Midstory flocks in our study area, consistent with previous findings (Bharadwaj et al., 2023). Additionally, our study uncovered complex interactions between bird species within and across flock types. Through detailed behavioural observations and species interaction analyses, we identified several species, including the Yellow-bellied Fairy-fantail (*Chelidorhynx hypoxanthus*), Ashy-throated Warbler (*Phylloscopus maculipennis*), Grey-cheeked Warbler (*Phylloscopus poliogenys*), and White-throated Fantail (*Rhipidura albicollis*), that actively participated in multiple flock types, indicating high interspecies interaction and potential information sharing across different flock types. This dynamic interplay highlights the complexity of avian social networks and emphasizes the importance of considering individual species' roles within larger ecological communities.

We postulated that species having higher vocalization rates, particularly during the early hours of the day when flock formation occurs are more important, or "central species". Our hypothesis was based on the concept that central species play a crucial role in initiating and maintaining group cohesion through vocal communication. To investigate this hypothesis, we conducted field observations and acoustic recordings, meticulously documenting the vocal behaviour of different bird species within mixed-species flocks and their social interactions.

Our findings revealed intriguing patterns in vocal activity among different flock types. Birds in understory flocks, led by species such as the Yellow-throated Fulvetta (*Schoeniparus cinereus*), exhibited synchronized vocalization patterns, with a consistent peak in vocal activity during the early hours. However, we did not observe a clear pattern in canopy flock types, suggesting potential variation in the role of vocalizations in influencing MSF formation and cohesion.

Our study found that within the understory flock type, species with higher closeness centrality (referred to as central species) exhibited higher mean call counts, signalling that central species are more vocal than other flock members. These findings align with previous

research, such as Smith (1991), which noted that key species such as tits (*Parus*) consistently vocalized within the flock. Similarly, in a tropical study, Wiley (1971) observed the dot-winged antwren, a central species in Central American antwren flocks, continuously vocalizing while moving between trees. This behaviour was used by other species to locate flocks as they moved slowly through dense vegetation, maintaining flock cohesion. Additionally, Munn (1985) in the Neotropical forests of Peru, discovered that the leader species, the White-winged Shrike-tanager (*Lanio versicolor*), made chattering vocalizations, prompting other flock members to join in. Therefore, the higher calling rate of central species observed in our study may serve as a mechanism for maintaining the cohesion of other species and forming the flock. Our results highlight the crucial role of key species in initiating and maintaining flock cohesion through vocal communication.

Understanding the extent to which species exchange information with each other is crucial for comprehending the structure of animal groups. Measuring the level of information flow between species can offer valuable insights into the dynamics of species interactions that underpin group dynamics, and may also indicate the extent to which certain species play a disproportionate role in maintaining group coherence.

In our research, while our primary focus was on the vocalization patterns and their correlation with the significance of species in mixed-species foraging (MSF) flocks, integrating metrics of information exchange between species could add further depth to our analysis. By quantifying the information exchange both within and between species, we can gain a better understanding of the communication dynamics within MSF, which in turn can provide valuable insights into the processes driving the formation and stability of these flocks.

Additionally, in terms of conservation, evaluating the symmetry of information flow can aid in identifying species that have key roles in maintaining group cohesion (Martinez, 2013). Previous studies have noted the absence of flock formation in the absence of certain key species (Cody & Diamond, 1975; Maldonado-Coelho and Marini, 2000). Species that engage in high levels of information exchange with others can act as keystone species within multi-species flocks, influencing the behaviour and spatial dynamics of other members (Martinez, 2013). Therefore, prioritizing the conservation of these species is crucial for preserving the composition and functionality of avian communities (Martinez, 2013).

Although our study provides valuable insights into the formation of MSF and avian behaviour, it is important to recognise its limitations. Our observations were based on two years of observing flocks and two months of collecting acoustic samples, emphasizing the necessity for long-term data to strengthen the reliability of our findings. Future research should involve longer sampling periods and investigate the functions of specific vocalizations or the vocalizations of particular species, potentially through playback experiments or vocal muting experiments (McDonald 1989).

By incorporating measures of information flow into future studies on MSF formation and dynamics, we can gain valuable insights into the underlying mechanisms of avian social behaviour and contribute to more effective conservation strategies aimed at preserving biodiversity and ecosystem stability. In summary, our study adds to the expanding knowledge of MSF formation and underscores the importance of key species in driving flock dynamics.

Understanding the complex interactions among individuals and species within different habitats is critical for the effective conservation and maintenance of community structure. Further exploration of the vocalizations of MSF birds and their influence on flock behaviour holds the promise of unravelling the intricacies of this fascinating ecological phenomenon, paving the way for more comprehensive conservation strategies.

Acknowledgement

We extend our heartfelt thanks to Pranav Balasubramanian, Anisha Mandal, and Tarun Menon for their unwavering support during the fieldwork period. Our gratitude also goes to the Arunachal Pradesh Forest Department for granting permission for this study. In addition, we sincerely appreciate the support from our peers, family, and the larger research community.

References

1. Aanensen DM, Huntley D, Feil EJ, al-Own F, & Spratt BG. (2009). EpiCollect: linking smartphones to web applications for epidemiology, ecology and community data collection. *PLoS ONE*, *4*(9).

2. Baltz, D. M., & Morejohn, G. V. (1977). Food Habits and Niche Overlap of Seabirds Wintering on Monterey Bay, California. *The Auk*, *94*(3), 526–543. https://doi.org/10.1093/AUK/94.3.526

3. Bell, H. L. (1983). A Bird Community of Lowland Rainforest in New Guinea. 5. Mixed-Species Feeding Flocks. *Emu - Austral Ornithology*, *82*(5), 256–275. https://doi.org/10.1071/MU9820256S

4. Bharadwaj, A., Thapa, A., Bhat, A., Biswakarma, A., Tamang, B., Munda, B., Biswakarma, B., Pradhan, D. K., Tamang, D., & Pradhan, K. (2023). Understanding inter-individual social networks in mixed-species bird flocks. *BioRxiv*, 2023–2024.

5. Billerman, S. M., Keeney, B. K., Rodewald, P. G., & Schulenberg, T. S. (2022). Birds of the World. *Cornell Laboratory of Ornithology*.

6. Britzke, E. R. (2004). Designing monitoring programs using frequency-division bat detectors: active versus passive sampling. *Bat Echolocation Research: Tools, Techniques and Analysis*, 79.

7. Charif, R., AM Waack, & LM Strickman. (2010). *Raven Pro 1.4 User's Manual.*

8. Cody, M. L., & Diamond, J. M. (1975). *Ecology and evolution of communities*. Harvard University Press.

9. Csardi, G., & Nepusz, T. (2006). The Igraph Software Package for Complex Network Research. *InterJournal, Complex Systems*, *1695*(5), 1–9. https://www.researchgate.net/publication/221995787

10. Diamond, J. (1987). Flocks of Brown and Black New Guinean Bird: A Bicolored Mixed-species Foraging Association. *Emu*, *87*(4), 201–211. https://doi.org/10.1071/MU9870201

11. Dolby, A. S., & Grubb, T. C. (2000). Social context affects risk taking by a satellite species in a mixed-species foraging group. *Behavioral Ecology*, *11*(1), 110–114. https://doi.org/10.1093/BEHECO/11.1.110

12. Farine, D. R., & Whitehead, H. (2015). Constructing, conducting and interpreting animal social network analysis. *Journal of Animal Ecology*, *84*(5), 1144–1163. https://doi.org/10.1111/1365-2656.12418

13. Farley, E. A., Sieving, K. E., & Contreras, T. A. (2008). Characterizing complex mixed-species bird flocks using an objective method for determining species participation. *Journal of Ornithology*, *149*(3), 451–468. https://doi.org/10.1007/S10336-008-0284-Z/TABLES/3

14. Goodale, E., & Beauchamp, G. (2010). The relationship between leadership and gregariousness in mixed-species bird flocks. *Journal of Avian Biology*, *41*(1), 99–103. https://doi.org/10.1111/j.1600-048X.2009.04828.x

15. Goodale, E., Beauchamp, G., Magrath, R. D., Nieh, J. C., & Ruxton, G. D. (2010). Interspecific information transfer influences animal community structure. *Trends in Ecology & Evolution*, *25*(6), 354–361. https://doi.org/10.1016/J.TREE.2010.01.002

16. Goodale, E., Beauchamp, G., & Ruxton, G. D. (2017). Adaptive Implications of Mixed-Species Grouping: Foraging, Physical, and Reproductive Factors. *Mixed-Species Groups of Animals*, 59–80. https://doi.org/https://doi.org/10.1016/B978-0-12-805355-3.00004-X

17. Goodale, E., & Kotagama, S. W. (2006). Vocal mimicry by a passerine bird attracts other species involved in mixed-species flocks. *Animal Behaviour*, *72*(2), 471–477. https://doi.org/10.1016/j.anbehav.2006.02.004

18. Goodale, E., & Kotagama W., S. (2005). Alarm Calling in Sri Lankan Mixed-Species Bird Flocks. *The Auk*, *122*(1), 108–120. https://academic.oup.com/auk/article/122/1/108/5562418

19. Greenberg, R. (2000). Birds of Many Feathers: The Formation and Structure of Mixed-Species Flocks of Forest Birds. In S. Boinski & P. Garber (Eds.), *On the move: how and why animals travel in groups* (pp. 521–558). University of Chicago Press.

20. Harrison, N. M., & Whitehouse, M. J. (2011). Mixed-species flocks: An example of niche construction? *Animal Behaviour*, *81*(4), 675–682. https://doi.org/10.1016/j.anbehav.2011.01.013

21. Hill, A. P., Prince, P., Snaddon, J. L., Doncaster, C. P., & Rogers, A. (2019). AudioMoth: A low-cost acoustic device for monitoring biodiversity and the environment. *HardwareX*, *6*, e00073. https://doi.org/https://doi.org/10.1016/j.ohx.2019.e00073

22. Hogstad, O. (1978). Sexual Dimorphism in Relation to winter foraging and Territorial Behaviour of three-toed woodpecker (Picoides tridactylus) and three Dendroccops species. *Ibis*, *120*(2), 198–203. https://doi.org/10.1111/J.1474-919X.1978.TB06775.X

23. Hutto, R. L. (1994). The Composition and Social Organization of Mixed-Species Flocks in a Tropical Deciduous Forest in Western Mexico. *The Condor*, *96*(1), 105–118. https://doi.org/10.2307/1369068

24. Jullien, M., & Thiollay, J.-M. (1998). Multi-Species Territoriality and Dynamic of Neotropical Forest Understorey Bird Flocks. *Journal of Animal Ecology*, *67*(2), 227–252.

25. Kushlan, J. A. (1977). The Significance of Plumage Colour in the Formation of Feeding Aggregations of Ciconiiformes. *Ibis*, *119*(3), 361–364. https://doi.org/10.1111/J.1474-919X.1977.TB08256.X

26. Macdonald, D. W., & Henderson, D. G. (1977). Aspects of the Behaviour And Ecology of Mixed-Species Bird Flocks in Kashmir. *Ibis*, *119*(4), 481–493. https://doi.org/10.1111/j.1474-919X.1977.tb02055.x

27. Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M., & Hornik, K. (2023). *cluster: Cluster Analysis Basics and Extensions*.

28. Maldonado-Coelho, M., & Marini, M. A. ˆ. (2000). Effects Of Forest Fragment Size And Successional Stage On Mixed-Species Bird Flocks In Southeastern Brazil. *The Condor*, *120*, 585–594. https://academic.oup.com/condor/article/102/3/585/5562985

29. Marler, P. (2004). Bird Calls: Their Potential for Behavioral Neurobiology. *Annals of the New York Academy of Sciences*, *1016*(1), 31–44. https://doi.org/10.1196/annals.1298.034

30. Martinez, A. E. (2013). *Interspecific Communication And its Consequences for the Organization of Permanent Mixed-Species Bird Flocks*.

31. McDonald, M. V. (1989). Function of song in Scott's seaside sparrow, Ammodramus maritimus peninsulae. *Animal Behaviour*, *38*(3), 468–485. https://doi.org/https://doi.org/10.1016/S0003-3472(89)80040-5

32. Mönkkönen, M., Forsman, J., Helle, P., & Monkkonen, M. (1996). Mixed-Species Foraging Aggregations and Heterospecific Attraction in Boreal Bird Communities. *Oikos*, *77*, 127. https://doi.org/10.2307/3545592

33. Morse, D. H. (1970). Ecological Aspects of Some Mixed-Species Foraging Flocks of Birds. *Ecological Monographs*, *40*(1), 119–168. https://doi.org/10.2307/1942443

34. Morse, D. H. (1977). Feeding Behavior and Predator Avoidance in Heterospecific Groups. *BioScience*, *27*(5), 332–339. https://doi.org/10.2307/1297632

35. Munn, C. A. (1985). Permanent Canopy and Understory Flocks in Amazonia: Species Composition and Population Density. *Ornithological Monographs*, *36*, 683–712. https://doi.org/10.2307/40168312

36. Powell, G. V. N. (1985). Sociobiology and Adaptive Significance of Interspecific Foraging Flocks in the Neotropics. *Ornithological Monographs*, *36*, 713–732. https://doi.org/10.2307/40168313

37. R Core Team. (2023). R: A Language and Environment for Statistical Computing. *R Foundation for Statistical Computing, Vienna, Austria*.

38. Sealy, S. (1973). Interspecific Feeding Assemblages of Marine Birds off British Columbia. *The Auk*, *90*, 796–802. https://doi.org/10.2307/4084359

39. Smith, S. M. (1991). The Black-Capped Chickadee. In *Behavioral Ecology and Natural History*. Cornell University Press.

40. Sridhar, H., Beauchamp, G., & Shanker, K. (2009). Why do birds participate in mixed-species foraging flocks? A large-scale synthesis. *Animal Behaviour*, *78*(2), 337–347. https://doi.org/10.1016/j.anbehav.2009.05.008

41. Srinivasan, U. (2019). Morphological and behavioral correlates of long-term bird survival in selectively logged forest. *Frontiers in Ecology and Evolution*, *7*(FEB). https://doi.org/10.3389/fevo.2019.00017

42. Srinivasan, U., Raza, R. H., & Quader, S. (2010). The nuclear question: rethinking species importance in multi-species animal groups. *Journal of Animal Ecology*, *79*(5), 948–954. https://doi.org/10.1111/J.1365-2656.2010.01707.X

43. Suzuki, T. N. (2012). Long-distance calling by the willow tit, Poecile montanus, facilitates formation of mixed-species foraging flocks. *Ethology*, *118*(1), 10–16. https://doi.org/10.1111/j.1439-0310.2011.01982.x

44. Wasserman, S., & Faust, K. (1994). *Social Network Analysis*. Cambridge University Press. https://doi.org/10.1017/CBO9780511815478

45. Wickham, H., Chang, W., & Wickham, M. H. (2016). Package 'ggplot2.' *Create Elegant Data Visualisations Using the Grammar of Graphics. Version*, *2*(1), 1–189.

46. Wickham, H., Francois, R., Muller, K., & Vaughan, D. (2023). dplyr: A Grammar of Data Manipulation. In *R package version 1.1.4, https://github.com/tidyverse/dplyr, https://dplyr.tidyverse.org*.

47. Wiley, R. H. (1971). Cooperative Roles in Mixed Flocks of Antwrens (Formicariidae). *The Auk*, *88*(4), 881–892. https://doi.org/10.2307/4083845

48. Winterbottom., J. M. (1943). On Woodland Bird Parties in Northern Rhodesia. *Ibis*, *85*(4), 437–442. https://doi.org/10.1111/J.1474-919X.1943.TB03857.X

Supplementary Material

Figure 1 MSF Observation Sampling Locations

 Figure 2 Opportunistic Acoustic Sampling Locations

Active Acoustic Sampling Points - Eaglenest WS

Figure 3 Active Acoustic Flock Sampling Locations

 Figure 4 Distribution of Closeness Centrality in the MSF

Figure 5 Correlation of Peak Vocalisation Category and Closeness Centrality

The Y-axis illustrates the values of closeness centrality, whereas the X-axis displays the peak vocalisation category. The plots indicate a positive correlation between closeness centrality and peak vocalization category for canopy flocks, although this relationship was insignificant. A correlation could not be established for the understory flock, as all variables

565 on the X-axis were identical, indicating a synchronised peak vocalisation between the

566 different species.