In situ structural analysis of mammalian cells using a 200 kV electron cryomicroscope – implications for research infrastructure.

Piotr Szwedziak[†*]

Center for Microscopy and Image Analysis
University of Zurich

[†]corresponding author:

Piotr Szwedziak

[†]Center for Microscopy and Image   *ReMedy-International Research
Analysis   Agenda Unit
University of Zurich   University of Warsaw
Winterthurerstrasse 190   Stefana Banacha 2c
8057 Zurich   02-097 Warsaw
Switzerland   Poland

email: piotr.szwedziak@zmb.uzh.ch

keywords: electron cryomicroscopy, electron cryotomography, *in situ*, subtomogram averaging, core facility, research infrastructure

Abstract

Background

Electron cryotomography is a powerful imaging technique allowing for studying
functional cellular modules in their native environment with macromolecular
resolution. However, it requires access to complex and expensive instrumentation,
typically a 300 kV electron cryomicroscope equipped with an energy filter.
Simpler and cheaper 200 and 100 kV instruments have been successfully used for
single particle cryoEM analyses, which has helped to democratize the technique
and broaden access. It has not been systematically studied if 200 kV electron
cryomicroscopes can deliver meaningful and interpretable data with respect to
electron cryotomography applications.

Methods

Here, we set out to investigate if a 200 kV electron cryomicroscope without an
energy filter can be utilized for *in situ* structural studies of mammalian cells by
electron cryotomography of thin cell edges followed by extensive image analysis
including segmentations, subtomogram averaging and molecular sociology
studies of lipid droplets.

Results

We demonstrate that the resulting tomograms of thin edges of U2OS cells are of
sufficient quality to annotate the contents of the cell and observe spatial inter-
relationships among macromolecules. In particular, we undertook a molecular
sociology analysis of lipid droplets and addressed their subcellular distribution
and interactions with other organelles. Additionally, we performed subtomogram
averaging of purified 70S ribosomes that resulted in ~15 Å resolution 3D
reconstruction. Finally, we examined geographical distribution and scientific
output of the two most common electron cryomicroscopy platforms and deduced
that 200 kV instruments are heavily underutilized with respect to electron
cryotomography applications.

Discussion

This study demonstrates that 200 kV electron cryomicroscopes can be utilized for

structural cell biology studies by electron cryotomography. Given the favorable

ratio of their versatility versus costs we foresee that 200 kV electron

cryomicroscopes will become workhorses of local electron cryomicroscopy

facilities.

Introduction

It has been a decade since the resolution revolution in the electron

cryomicroscopy (cryoEM) field occurred (Kühlbrandt, 2014). Single particle

analysis (SPA) cryoEM has been a tremendously successful structural biology

technique (as evidenced by the exponentially growing number of relevant entries

in the Electron Microscopy Data Bank) enabling routine determination of high-

resolution 3D structures of purified macromolecules such as heterogenous or

flexible protein complexes (Punjani & Fleet, 2023; Schwab et al., 2024).

However, biological functions are not executed by individual molecules in

isolation but rather complex networks of interactions between all the cellular

components, including proteins, nucleic acids and lipids, are responsible for

maintaining cellular homeostasis. Therefore, the need for structural studies by

cryoEM in unperturbed cellular environments is steadily growing (Nogales &

Mahamid, 2024). Electron cryotomography (cryoET) is a cryoEM modality that

provides 3D molecular resolution images of isolated/*in vitro* reconstituted

cellular components as well as intact cellular environments, which is known as *in*

situ cryoET. CryoET preserves the full spectrum of conformations and interactions

of each molecule, enabling detailed ultrastructural analyses of cells and tissues in

health and disease that has been described as 'visual proteomics' (Beck &

Baumeister, 2016; Förster et al., 2010). If structures of interest are present in

multiple copies within a tomogram, they can be extracted, aligned and averaged

in a process called subtomogram averaging, which is analogous to SPA but makes

use of 3D volumes rather than 2D projections. In favorable conditions this can

yield a near-atomic resolution reconstructions (Tegunov et al., 2021). Additionally,

the position and context of molecules and how they relate to the cellular context

are available.

Given that the mean free path for inelastic scattering for 300 kV electrons is ~300-

400 nm (Martynowycz et al., 2021; Rice et al., 2018), a critical requirement and

limitation for cellular cryoET is sample thickness. The resulting tilt series images

must have enough signal to noise ratio in order to be reliably aligned when

reconstructing the 3D volume (tomogram). Therefore, initially the analyses were

limited to *in vitro* specimens, bacteria or thin edges of eukaryotic cells (Medalia et

al., 2002; Szwedziak et al., 2014; Szwedziak & Pilhofer, 2019). In the last ten years

techniques based on dual-beam FIB-SEM instruments have become available to

prepare thin lamellae of cells and tissues for subsequent cryoET analyses (Rigort

& Plitzko, 2015).

Complex sample preparation techniques, fragmented data processing pipelines

and lower throughput, as compared to SPA cryoEM, are primary reasons why

cryoET still lags SPA in terms of maturity and output. Cellular cryoET is a costly

technique as it frequently requires access to high-end 300 kV TEMs equipped with

energy filters that remove inelastically scattered electrons in order to increase the

signal-to-noise ratio. In the future the costs will become even more prohibitive

given the implementation of laser phase plates and Cc correctors that will yield an

ultimate, cryoET purpose-built instrument (Russo et al., 2022).

Given that the mean free path at 200 kV is ~260 nm and inspired by the study

(Martynowycz et al., 2021) suggesting that useful cryoEM data can be collected up

to twice the inelastic mean free path without the need for energy filters, we set out

to investigate if a 200 kV TEM equipped with a direct electron detector is sufficient

to provide meaningful cryoET data.

Here, we imaged by cryoET thin edges of mammalian U2OS cells using a 200 kV

Glacios G1 cryoTEM fitted with a Falcon 3EC camera without an energy filter. The

tomograms enabled us to observe a plethora of cellular structures ranging from

mitochondria through cytoskeletal filaments to macromolecular complexes, e.g.

ribosomes. By combining segmentation with subtomogram averaging we

generated detailed ultrastructural cellular snapshots that might be useful in

addressing various cell biology problems, e.g. lipid droplets (LDs) metabolism.

LDs are central to lipid and energy homeostasis and are important storage

organelles (Olzmann & Carvalho, 2019). LDs are known to originate in the

endoplasmic reticulum (ER) and associate with other cellular structures and a

myriad of proteins (Olzmann & Carvalho, 2019). To better understand the role of

LDs in cellular metabolism we undertook a molecular sociology analysis of LDs

and addressed their subcellular distribution and interactions with other

organelles. We also provide visual evidence for direct interactions between LDs

and membrane trafficking proteins. Finally, we obtained a ~15 Å subtomogram

average of purified and vitrified *Escherichia coli* 70S ribosomes.

The current trend towards democratizing cryoEM in the SPA field is gaining

momentum and it has been demonstrated that 200 kV (Herzik et al., 2017) and

100 kV (McMullan et al., 2023) cryoTEMs are sufficient to obtain high-quality SPA

cryoEM data. Here, we prove that 200 kV TEMs without an energy filter can be

utilized for cellular cryoET investigations. We suggest that 200 kV cryoTEMs can

be cost-effective workhorse cryoEM instruments capable of providing

complementary information from both major cryoEM modalities and with the

recent development of electron detectors optimized for lower electron energies

(Chan et al., 2024; McMullan et al., 2023) their performance and significance will

steadily grow.

Results

CryoET imaging of mammalian cells.

CryoET has been frequently employed to address cell biology problems that

otherwise were not accessible to other imaging techniques, e.g. coronavirus

replication (Wolff et al., 2020) or detailed membrane architecture (Wozny et al.,

2023). To ensure optimal performance and data quality these studies made use of

complex 300 kV TEMs equipped with energy filters. Due to prohibitive costs of

purchase and maintenance, such instruments are predominantly available in

major research centers with limited access. Given that the mean free path for

inelastic scattering at 200 kV is ∼260 nm, which is comparable to the average

cryoET specimen thickness, we set out to investigate if a 200 kV TEM without an

energy filter can deliver interpretable cryoET results. Specifically, we were

interested in understanding the interplay and spatial arrangement of LDs in the

context of intact cells.

U2OS cells were seeded on electron microscopy grids with a hole carbon support

film and vitrified by plunge-freezing. These cells are known to adopt a flat,

pancake-like shape that offers a significant cryoET imaging surface over thin cells

areas without the need for micromachining by cryoFIB. The grids were imaged by

a Glacios G1 cryoTEM equipped with a Falcon 3EC camera without an energy filter.

The cells could be easily recognized at low magnifications (**Fig. 1a, left**) with the

cell nucleus surrounded by the electron transparent cytoplasmic regions. This

allowed for a relatively straightforward setup of automated tilt series acquisition

and great care was taken to tune the 2-condenser lens system to parallel

illumination conditions. It was possible to acquire multiple tilt series from a single

cell, and the final thickness of the 3D volume was dictated by the local cell

thickness (**Fig. 1a, small panels**). The imaged areas were structurally well

preserved and showed no sign of crystalline ice contamination. In total, 64 cells

were imaged over which we recorded almost 230 tomograms with an average

thickness of 240 ± 75 nm (**Fig. 1b**). Most tomograms were between 200 and 250

nm thick, with the thinnest being 50 nm and thickest 400 nm (**Fig. 1c**). Attempts

to record tilt series on thicker areas failed due to the low contrast of resulting

images and poor tracking. To assess the quality of the data quantitively we fitted

the contrast transfer function (CTF) on the 2D zero tilt images and using

CTFFIND4 we estimated the resolution up to which the CTF is fitted reliably. We

found that the precision of the fit is linearly dependent on the local cell thickness

(**Fig. 1d**). Encouraged by these results we moved on to a more detailed inspection

of the tomograms.

Molecular atlases of native cellular environments.

Since the aim of this study was to generate low nanometer resolution 3D volumes

of intact mammalian cells in order to qualitatively and quantitatively describe LDs

molecular neighborhoods, we adopted an appropriate data collection strategy

(**Table 1**): we utilized relatively high defocus (range from -5 to -7 μm) and high

total dose (140 eÅ^{-2}) to boost image contrast and large pixel size (5.1 Å) to benefit

from the broad cellular context provided by the large field of view (~2 μm). The

figures presented were prepared using tomograms generated by SIRT

(Simultaneous Iterative Reconstruction Technique) as it offers high contrast that

is useful for downstream cellular tomography applications such as segmentations

and annotations. The quality of the tomograms can be judged from the clearly

distinguishable cellular features, e.g.: microtubules, intermediate filaments, actin

filaments, lipid tubules, ribosomes (**Fig. 2a, left panels**). A representative

tomogram of thickness of 180 nm is shown in **Fig. 2a (left large panel)** and **Movie**

S1. In the total volume of ~0.72 μm^3 we could find the three classes of intracellular

filaments (microtubules, intermediate filaments, actin) intertwined with each

other as well as with membranous structures of the endoplasmic reticulum (ER)

and ribosome particles. Subsequently we trained a neural network within the

EMAN2 software package (Chen et al., 2017) to recognize lipid bilayers,

intermediate filaments, double membranes of mitochondria, microtubules and

ribosomes and applied it to the tomograms (**Fig. 2a, middle panels**). This allowed

for reproducible identification of features of interest and their separation from the

background. By applying the neural network, we identified and automatically

extracted ∼1500 ribosomal particles and subjected them to subtomogram

averaging. This resulted in a moderate resolution 3D reconstruction at 38 Å with

the conservative threshold of 0.5 for FSC (**Fig. 2a, small panel**). Including more

particles in the average didn't improve the resolution suggesting that the dataset

is limited by data collection strategy used (high defocus and large total electron

dose) and sample thickness. Importantly, resolution is not the best criterion for

data quality in cryoET as the contextual information provided by cellular

tomograms can be of higher significance. Accordingly, by mapping the averaged

structure to the calculated positions and orientations of each particle in the

tomograms and combining it with segmentations of other cellular components we

generated 3D molecular maps of thin edges of intact mammalian cells (**Fig. 2a,**

right panel, Movie S2). Such maps are visually appealing to human perception

and easier in interpretation than raw tomograms. We find the prospect of

producing detailed annotations of 3D cryoET volumes obtained by imaging with a

200 kV cryoTEM without an energy filter particularly exciting since such

segmentations aid interpretation and are essential for many downstream cryoET

analysis steps, e.g. observing spatial inter-relationships among macromolecules.

We went on to evaluate if the 200 kV cryoET data allow for reliable identification

of other known subcellular structures (**Fig. 2b**). In 13 tomograms we recognized

mitochondria that adopted either spherical or rod-shaped morphologies (**Fig. 2b**

(I), Movie S3 and Fig. S1). The measured cristae width was 12.8 nm, which stays

in accordance with the previously published 14.5±5.8 nm (Fry et al., 2024). Within

the mitochondrial matrix we could observe abundant granules characterized by

high contrast – we attributed those densities to calcium phosphate storage, as first

described in (Wolf et al., 2017). Calcium phosphate granules are used to regulate

calcium homeostasis and to maintain mitochondrial function. Similar densities

were observed by cryoET in neuronal mitochondria (G. H. Wu et al., 2023). CryoET

at 200 kV could be an excellent diagnostic platform for *in situ* organelle

phenotyping as it has been shown that mitochondrial ultrastructure, including its

membrane architecture, has profound implications for human disease

(Navaratnarajah et al., 2021).

CryoET at 200 kV was sufficient to resolve coated vesicles/buds that play a role in

transport between various cellular compartments (**Fig. 2b (II), Movie S4**). The

overall morphology and the localization in the cell peripheries indicate that the

presented membrane-bound compartment could be an early endosome. A closer

look at the dense protein coat and comparison to the previously published cryoET

data (Bykov et al., 2017) suggests it is COPI. COPI coat is known to play an

important role in mediating transport between the Golgi and ER within the

secretory pathway but it has also been demonstrated to act in early endosome

function (Daro et al., 1997). The observed buds/vesicles had an average diameter

of 55±7 nm (n=7), which stays in accordance with other reports (Bykov et al.,

2017; Faini et al., 2012). To fully appreciate the complex membrane architecture,

we recommend seeing **Movie S4**.

In the imaged cells we found extensive areas of ER. The observed ER is

characterized by the lumen deprived of other intracellular

organelles/macromolecular complexes and is tightly coated on the cytoplasmic

face by ribosomes where individual particles could be discerned (**Fig. 2b (III) and**

Movie S5). The ER network consisted of cisterna interconnected by an array of

branching tubules. In 10 tomograms we also identified a recently reported form

of ER – ribosome associated vesicles (RAVs). RAVs were originally described as

dynamic ER subcompartments found primarily in the periphery of secretory cells

(Carter et al., 2020). Here, we found numerous RAVs interacting tightly with

microtubules that in the most severe cases triggered membrane flattening (**Fig.**

2b (IV) and Movie S6 and Fig. S2). RAV-bound ribosomes were excluded from

the RAV-microtubule interface. The observed RAVs were interconnected via

membrane tubules that were 8 nm in diameter (**Fig. 2b (V) and Movie S7**).

Membrane-bound polyribosome, which is a cluster of ribosomes bound to an

mRNA, was found on the RAV surface (**Fig. S3 and Movie S8**). Given that actively

translating mRNAs are frequently associated with multiple ribosomes we suspect

that RAVs might be sites of membrane/extracellular protein synthesis, similarly

to the canonical ER.

We also observed multilamellar compartments containing vesicles as well as

membrane sheets and other cellular debris/dense deposits. These are most likely

degradative compartments, for example autophagosomes (**Fig. 2b (VI) and**

Movie S9) or endolysosomes (**Fig. 2b (VII) and Movie S10**), as described in

(Foster et al., 2022). The luminal density within these organelles was highly

variable, suggesting that their content is different and not well defined.

Detailed analysis of LDs.

In 84 tomograms we noticed the presence of LDs. They are characterized by an

amorphous, electron-dense core made of neutral lipids with a single phospholipid

layer on the outside, which is manifested as the electron-dense line delineating

the LDs circumference in the tomograms (**Fig. 3a, top panel and Movie S11**). In

total we recognized 231 LDs that were 205±70 nm in diameter. LDs were often

found in clusters (**Fig. 3a, bottom panels and Movie S12**) and 29% of LDs stayed

in direct contact with each other, which frequently resulted in shape changes and

flattening (**Fig. 3a, bottom right panel and Fig. S4**), as reported previously

(Ganeva et al., 2023). ER double leaflet is the site of LDs synthesis (Fujimoto &

Parton, 2011) and they remain tightly associated with the ER throughout their life

cycle (**Fig. 3b, top right panel and Movie S13**): indeed, 69% of the cytoplasmic

LDs were in contact with ER whose membrane curvature followed LD's shape (**Fig.

3b, top left panel, Movie S14**). Interestingly, 5.6% of the observed LDs were

positioned in the ER lumen (**Fig. 3b, top right panel and Fig. S5**). The luminal

LDs are the least characterized members of the mammalian LD family, and more

investigations are needed to establish what triggers LDs movement towards

either cytoplasmic or luminal side. LDs association with ER was described as very

promiscuous (Valm et al., 2017) and we could observe several distinct instances

of this interaction. For example, luminal and cytoplasmic LDs in direct contact

presumably during material exchange (**Fig. 3b, bottom middle panel**) or

cytoplasmic LDs impacting the ER membrane and remaining at a distance of 10-

12 nm that is sufficient to accommodate proteins mediating the interaction (**Fig.

3b, bottom right panel**). The lipid monolayer is a busy hub for interaction with

proteins that frequently are LD-specific (Olarte et al., 2022) and mediate LDs

interactions with various organelles (Olzmann & Carvalho, 2019): for example,

3% of all LDs interacted with mitochondria (**Fig. 3c, left panel and Movie S15**)

and this association has been reported before as being crucial for metabolism (Fan

& Tan, 2024). 40% of the identified LDs interacted with microtubules (**Fig. 3c,**

middle bottom panel, Movie S16 and Fig. S6). Direct physical contact between

LDs and microtubules can aid with transport of LDs and facilitate finding their

cognate organelles. Rapid LD movement on microtubules has been observed by

time-lapse fluorescent light microscopy imaging (Gross et al., 2000; Kilwein &

Welte, 2019). A small fraction (1.7% of total observed LDs) of LDs interacted with

putative lysosomes and MVBs (**Fig. 3c, middle top and right panels**). These LDs

are most likely destined for degradation by lysosomal acid lipase and the resulting

fatty acids can enter metabolic pathways in mitochondria (Zhang et al., 2022). We

could discern direct association of membrane trafficking proteins on the LDs

surface. COPI has been demonstrated to act directly on lipid droplets *in vitro*

(Thiam et al., 2013) and *in vivo* to enable their connection to the ER for protein

targeting (Wilfling et al., 2014). Here, we observed the dense protein coating on

the surface of an LD that promoted budding of lipid monolayer resulting in an

empty void between the LD core and the monolayer (**Fig. 3d left panels and**

Movie S17). Based on the protein coat morphological similarity to previously

published coated vesicles (Bykov et al., 2017) we assume that the observed

protein density might be attributed to COPI. It is likely that COPI-induced

membrane deformation could help establish continuous membrane bridges

between the LD and outer ER membrane leaflet (Wilfling et al., 2013), which has

been observed in our tomograms (**Fig. 3b, bottom left panel**). Here, the LD is

tightly embedded within the lipid bilayer. Another kind of vesicle protein coat,

clathrin, has been detected in LDs by mass spectrometry (Bartz et al., 2007;

Nonoyama et al., 2019) and our study provides visual confirmation of this

interaction (**Fig. 3d, middle and right panels**). The triskelion building block

assembling into pentagonal and hexagonal lattice was readily visible and the

overall arrangement was similar to clathrin cage architecture reported by SPA

cryoEM (Fotin et al., 2006). The role of clathrin in association with LDs is unclear

but it has been suggested that it might be important for lipid hydrolysis as it

facilitates lysosome formation that is essential for lipid droplet digestion

(Nonoyama et al., 2019).

LDs are ubiquitous in the eukaryotic world and essential for lipid and energy

metabolism. With the emergence of cryoET imaging we are now beginning to

appreciate how LDs interact with other cellular compartments (summarized in

Table 2), and how these interactions play a role in cellular homeostasis. We

believe that structural insights into LDs *in situ* biology obtained by cryoET imaging

at 200 kV might further advance the field.

1 **Discussion**
2
CryoET is a versatile and unique structural biology technique as it can bridge
information from several orders of magnitude (Å to µm) and is an ideal tool for
integrative studies (Ziemianowicz & Kosinski, 2022). In this investigation we
sought to evaluate whether a 200 kV cryoTEM equipped with a direct electron
detector without an energy filter can deliver interpretable cryoET data that could
be used to address structural cell biology problems. Our primary objective was to
utilize the mammalian U2OS cell line as the model specimen, compare obtained
findings into cellular architecture with the previously published data acquired at
300 kV cryoTEMs fitted with direct electron detectors and energy filters and
possibly provide novel insights. It has been demonstrated that radiation damage
to biological material is more pronounced at 200 kV than at 300 kV (Peet et al.,
2019). Here, we experimentally assessed and carefully tuned the maximal
acceptable electron dose to obtain high contrast micrographs suitable for cellular
studies by cryoET. This work resulted in several significant outcomes: firstly, the
utilized setup allowed for cryoET tilt series acquisition over thin edges of
mammalian cells with thicknesses ranging from 50 to 300 nm that could
subsequently be reliably reconstructed into 3D volumes; secondly, the resulting
tomograms could be reproducibly segmented and all the major cellular
components readily recognized; thirdly, the images of RAVs, LDs and other
cellular compartments were qualitatively comparable to previously published
tomograms acquired by 300 kV cryoTEMs with energy filters (Dudka et al., 2024;
Ganeva et al., 2023; Mahamid et al., 2019); fourthly, quantitative analysis of LDs
in their native cellular environment revealed the molecular sociology of these
organelles. Additionally, in this work we provide molecular resolution direct
visual evidence of LDs interaction with membrane trafficking components. We
expect that a 200 kV electron cryomicroscope without an energy filter will
produce similar results when imaging lamella of cellular/tissue specimens
obtained by cryo-focused ion beam milling using either gallium or plasma sources
which typically are micromachined to the final thickness of 150-200 nm. It
remains to be seen how a 200 kV cryoTEM without an energy filter can perform
regarding high-resolution subtomogram averaging, including both *in vitro* and

cellular specimens. With this respect we have performed an analysis of purified

70S *Escherichia coli* ribosomes. Subtomogram averaging of ~4000 ribosomal

particles yielded a 15 Å resolution 3D reconstruction with the threshold of 0.5 for

FSC (**Fig. S7**). This demonstrates that a 200 kV cryoEM instrument without an

energy filter can generate medium resolution cryoEM maps useful for

understanding the overall architecture of macromolecular complexes.

The field of *in situ* structural studies by cryoET is lagging behind SPA cryoEM – as

of August 2024 83.4% EMDB entries were derived from SPA analyses vs. 11.4%

obtained by cryoET approaches. However, the dynamics of the growth for the

cryoET modality is greater – in 2023 year-over-year growth was 50% as

compared to 24% for SPA. This rapid growth has been partially possible due to

higher throughput data collection strategies (Eisenstein et al., 2023, 2024) and

attempts to streamline data processing (Burt et al., 2024; Chen et al., 2019a).

Current hardware developments including C_C correction, laser phase plates and

reduced radiation damage at liquid helium temperatures will further boost the

field of cryoET but with a hefty price tag. Such cryoET-purpose built instruments

will be prohibitively expensive to purchase and maintain and will be available in

a handful of institutions only.

To better understand how current cryoEM resources are utilized we reviewed the

output (based on the EMDB database) and distribution (based on the map of high-

end cryoEM worldwide (Noble, 2024) – please note that map data might be

incomplete since it is based on voluntary contributions from the community) of

the two most popular 200 and 300 kV cryoEM platforms: Thermo Scientific

Glacios and Thermo Scientific Krios, respectively (**Fig. 4a**). In 2023 there were

5379 EMDB entries attributed to SPA and 837 entries attributed to cryoET

obtained from the Krios platform. Whereas Glacios contribution to SPA cryoEM

was only 0.95% of Krios in 2020, it steadily rose to 5.9% in 2023. In contrast, the

Glacios platform delivered only 4 cryoET EMDB entries in the whole of 2023.

As of August 2024, there were ~364 Krioses and ~230 Glacioses around the world

suggesting that 200 kV cryoTEMs are predominantly used as screening devices

and heavily underutilized for primary data collection, as evidenced by the number

of relevant EMDB entries. In further analysis we focused on Europe that is home

to 102 Krioses and 74 Glacioses. As shown on the map, the distribution of these

instruments across the continent is not equal. 17 European countries are in

possession of either Krios or Glacios or both, 2 countries have access to other

cryoEM platforms whereas the 25 remaining countries don't have access to

cryoEM instrumentation at all (**Fig. 4b, c**). The maturity and complexity of cryoEM

infrastructure reflects to a certain degree economic development - the top 3

Glacios/Krios owner countries are also 3 top countries by nominal GDP (according

to the International Monetary Fund (IMF), August 2024). Next, we surveyed the

distribution of the Glacios and Krios cryoEM platforms across academic hubs

defined here as cities or agglomerations characterized by the presence of research

institutes, either academic or commercial. The Krioses were spread across 39

academic hubs whereas Glacioses across 49, which accounts for 2.6 Krios per hub

and 1.5 Glacios per hub. Such clustering of state of the art 300 kV cryoTEMs and

better spatial distribution of 200 kV microscopes is related to the fact that the

former ones are usually maintained by national/international facilities (eBIC UK,

EMBL Germany, NeCEN Netherlands, SciLifeLab Sweden). This trend is visible

worldwide, e.g. in the United States there are 3 major SPA cryoEM centers and 4

cryoET centers under the umbrella of NIH Common Fund Transformative High

Resolution Cryo-Electron Microscopy Program Centers, each containing multiple

high-end instruments. Such centers provide not only access to state-of-the-art

microscopes but also to know-how and sample preparation services that,

particularly for cellular cryoET, are complex and expensive. The clustering of

cryoEM research infrastructure will become more prominent with the arrival of

another generation of cryoET purpose-built cryoTEMs. Such microscopes can cost

>5 M CHF with added costs for space, service contracts and qualified staff. With

this respect it is worth noting that mid-range 200 kV systems have a significantly

lower price tag and long-term running costs, e.g. service contracts, which is

important in planning facility operations and ensuring optimal instruments

performance. The data presented were acquired by operating the Falcon 3EC

camera in the linear mode without dose fractionation (due to the impracticalities

of long exposure times while operating the Falcon 3EC camera in the counting

mode). We suggest that such qualitative studies could also be performed using less

costly CMOS-based cameras (e.g. CETA), further bringing down the total

investment costs.

200 kV cryoTEMs have been demonstrated to deliver high-quality SPA data, e.g.

(Brouwer et al., 2024; Träger et al., 2024) as well as cryoET data from *in vitro*

samples, e.g. (Tian et al., 2020). Here, we show that these instruments are capable

of delivering excellent cryoET images of cellular specimens. We believe that the

200 kV platform can be a viable option for smaller universities and research

institutes that will help democratize cryoEM access and can serve as workhorse

for both major cryoEM modalities: SPA and cryoET. We expect that these 1st

degree cryoEM access points will further boost the field by attracting top-tier

experts, training independent users/operators, facilitating methods development

and fostering interdisciplinary research in local academic centers and will greatly

complement the 2nd degree cryoEM access points offering the latest and ultimate

technological solutions (**Fig. 4d**).

1 Acknowledgements

We thank Dr Ishier Raote (CRG Barcelona) for providing the U2OS cell line, Dr
Tomasz Góral (University of Warsaw) for the microscope maintenance, Sonia
Ngati (University of Warsaw) for the cell culture technical assistance and Dr
Fabian Eisenstein (ETH Zurich) for help with setting up and troubleshooting
SPACEtomo installation. 70S *E. coli* ribosomes were the generous gift from Dr
Alain Scaiola (ETH Zurich). Members of the Center for Microscopy and Image
Analysis at the University of Zurich are acknowledged for insightful discussions.
This work was supported by the "Regenerative Mechanisms for Health-ReMedy"
grant MAB/20172 from the Foundation for Polish Science.

12 Author contributions

13 PS: conception and design, acquisition of data, analysis and interpretation of data,
14 drafting and revising the article

Figure Legends

Figure 1. CryoET analysis of intact mammalian cells using a 200 kV cryoTEM

without an energy filter.

a) A low magnification overview of a U2OS cell seeded on a Quantifoil grid

and vitrified in liquid ethane. The electron opaque area encircled in blue

corresponds to the nucleus/perinuclear area. The cell boundaries are

encircled in red. Six tilt series were acquired in the electron-transparent

area corresponding to the thin cell peripheries. Central slices of

reconstructed tomograms from each area are shown in the panels on the

right and color-coded to mark the acquisition areas. Measured tomogram

thicknesses are displayed. Scale bars: 10 µm and 100 nm.

b) Table summarizing the dataset acquired in this study.

c) Histogram of the tomogram thicknesses. The thickness varied from 50 to

400 nm with the most populated class in the range of 200-250 nm.

d) Plot of resolution up to which CTF was reliably fitted for a central tilt in

every tilt series against the tomogram/local cell thickness. The red line

shows a linear fit through all data points.

Figure 2. CryoET at 200 kV provides molecular resolution atlases of intact

mammalian cells.

a) Central slice (14 nm thick) through a tomogram of an intact U2OS cell with

a local thickness of 180 nm (left large panel). Small panels on the left

display common mammalian cell features that were identified in the

tomogram, from top to bottom: microtubules, intermediate filaments,

actin, membrane tubules and ribosomes (red arrows). By employing

machine learning-based approaches we could identify, extract and average

subvolumes containing ribosome particles and map the resulting low

nanometer resolution average back in the tomogram (middle panels). The

quality of the tomogram was sufficient to segment other components of the

cell: membranes, mitochondria, intermediate filaments and microtubules

(middle panels). By combining all the segmentations, we generated a

simplified model of the cell that can aid interpretation (right panel). Scale

bars: small panels – 25 nm, main panel – 100 nm.

b) Examples of some of the functional cellular modules that could be readily

identified in the tomograms: I) Mitochondrion with calcium phosphate

inclusions (red arrow). Measured cristae diameter is highlighted in white.

II) Putative COPI-coated vesicles budding off an early endosome. Color-

coded are enlarged views of highlighted areas at a different tomographic

slice. COPI coat is characterized by a very dense and compact appearance.

III) endoplasmic reticulum (ER) with its empty lumen and the cytoplasmic

surface covered with ribosomes (red arrow). IV) ribosome-associated

vesicle (RAV) covered densely with ribosome particles except where it

interacts with a microtubule filament that ultimately leads to a local

ribosome deprivation and membrane flattening of otherwise spherical

structure (red arrow). V) RAVs and ER subcompartments are

interconnected by a dense network of thin membrane tubules that are 8

nm in diameter, as highlighted in the enlarged view that is a snapshot at a

different tomographic slice of the area encircled in blue. VI) A membrane-

embedded compartment that could be an autophagosome. VII) A bell-like

shaped membrane compartment with internal debris that could be an

endolysosome that is a result of the fusion of a lysosome with a

multivesicular body. Scale bars: major panels – 100 nm, enlarged views –

25 nm.

Figure 3. Molecular sociology of lipid droplets (LDs) in U2OS cell periphery as

revealed by cryoET at 200 kV.

a) LDs were found as spherical, amorphous entities characterized by distinct

contrast (top left panel). One of the unique structural features of LDs is

their lipid monolayer surface that manifests itself as an electron dense line

around perimeter (enlarged view, red arrow). LDs were frequently found

near each other (bottom left panel), including direct contact that resulted

in flattened and deformed morphology (bottom right panel, red arrow).

Scale bars: major panels – 100 nm, enlarged view – 25 nm.

b) LDs lifecycle is inherently connected to ER, which is the site of their synthesis. Cytoplasmic LDs were in close association with the ER whose membrane curvature mirrored the LD's shape (top left panel, blue arrowheads). Three distinct modes of LDs interactions with ER are shown on the top right panel and in enlarged views in the bottom panels, color-coded accordingly. The LD highlighted in orange is embedded within the ER lipid bilayer, which splits to accommodate the bulky volume of the LD (red arrow); in blue are highlighted luminal and cytoplasmic LDs interacting across the ER lipid bilayer (red arrow) and in red a cytoplasmic LD impacting the ER membrane and remaining at a distance of 10-12 nm that is sufficient to accommodate proteins mediating the interaction (Hugenroth & Bohnert, 2020). Scale bars: major panels – 100 nm, enlarged view – 25 nm.

c) LDs remain in close association (marked with red arrows) with various cellular modules: mitochondria (left panel), microtubules (middle bottom panel), putative MVBs (middle top panel), membrane embedded compartment that could be a lysosome (right panel). Scale bars: 100 nm.

d) LDs interact with membrane trafficking machinery. Top left panel depicts an LD with apparent lipid monolayer deformation (blue square). More detailed analysis (bottom left panel, which is a view of the region marked with the blue square at a different tomographic slice) suggests that the lipid monolayer might be coated with the dense COPI protein layer (red arrow, see also Fig. 2b (II) for comparison). More data and analyses are required to unequivocally identify the protein forming the coat.

Top middle and right panels show tomographic slices through central sections of two LDs and below their corresponding views at different tomographic slices. The surface of the two LDs is covered with clathrin that adopts its characteristic architecture with the triskelion forming both hexagonal and pentagonal units (red arrows). Scale bars: top left panel – 100 nm, other panels – 25 nm.

Figure 4. Current utilization of high-end cryoTEMs.

a) Table summarizing scientific output generated by two most common 200
kV and 300 kV cryoTEMs: Glacios and Krios, respectively. Whereas the
Glacios platform contributed less than 1% of the Krios SPA output in 2020,
this fraction rose to 5.9% in 2023. With respect to cryoET, the Glacios
platform is heavily underutilized, with only a marginal contribution. Based
on the EMDB database as of August 2024.

b) Geographical distribution of the Thermo Scientific Glacios and Krios
platforms in Europe. As of August 2024, there were ~102 Krioses and ~74
Glacioses scattered across 17 European countries. Eastern and South-
Eastern Europe regions are underequipped. Please note that in Spain,
United Kingdom, Switzerland, Belgium and Germany there are also JEOL
cryoARM platform cryoTEMs. Based on the map of high end cryoEM
worldwide as of August 2024 (Noble, 2024).

c) Number of Glacios and Krios cryoTEMs in European countries. Countries
are sorted from left to right according to their nominal gross domestic
product (according to IMF, August 2024).

d) Implications of this study for core facilities: it is expected that 200 kV
instruments will become versatile workhorses in the majority of local
cryoEM facilities. These 1st degree access facilities are essential for
democratizing access to cryoEM. The 2nd degree access facilities that
operate at the national/international level will offer access to cutting-edge
imaging technology and advanced sample preparation techniques for
cryoET. The technologies for laser phase plates (Schwartz et al., 2019), Cc
correctors (J. Wu et al., 2025) and liquid helium cryostages (He et al., 2024)
exist but these solutions are not commonly used or not yet available from
commercial suppliers.

Table 1. CryoET data collection.

Microscope	Thermo Fisher Glacios G1
Camera	Falcon 3EC
Energy filter	N/A
Magnification	28000
Voltage (kV)	200
Electron exposure (eÅ^{-2})	140
Defocus range (μm)	5.0-7.0
Pixel size (Å)	5.1
Tilt range	$\pm54^0$
Increment	3^0

3   **Table 2. Lipid droplets interactions summary.**

#total LDs	231
#cytoplasmic LDs	218
#ER luminal LDs	13
#LDs in contact with each other	68
#LDs in contact with microtubules	94
#LDs in contact with ER	151
#LDs in contact with mitochondria	7
#LDs in contact with lysosomes/MVBs	4

Supplementary Figures and Movies Legends

Figures:

Figure S1. Representative cryoET slices showing mitochondria with calcium
phosphate deposits. Corresponding to **Fig. 2b (I)**. Scale bars: 100 nm.

Figure S2. Representative cryoET slice showing a RAV closely associated with a
microtubule (red arrow). Corresponding to **Fig. 2b (IV)**. Scale bar: 100 nm.

Figure S3. Representative cryoET images showing a RAV at two corresponding
views at different tomographic slices. Red arrow highlights a polysome associated
with the surface of the RAV. Corresponding to **Fig. 2b (V)** and **Movie S8**. Scale
bars: 100 nm.

Figure S4. Representative cryoET slice showing several LDs, two of them closely
associated with each other (red arrow). Corresponding to **Fig. 3a**. Scale bar: 100
nm.

Figure S5. Representative cryoET slices showing LDs residing in the ER lumen.
Corresponding to **Fig. 3b**. Scale bars: 100 nm.

Figure S6. Representative cryoET slice showing LD closely associated with a
microtubule (red arrow). Corresponding to **Fig. 3c**. Scale bar: 100 nm.

Figure S7. Subtomogram averaging of purified 70S *E. coli* ribosomes. Shown are
the FSC curve (top) and overviews of the obtained cryoEM density map.

Movies:

Movie S1. Tomogram of a thin edge of an U2OS cell. Corresponding to **Fig. 2a**.
Scale bar: 100 nm.

Movie S2. Segmentation of a thin edge of an U2OS cell. Corresponding to **Fig. 2a**
and **Movie S1**.

Movie S3. Tomogram of a mitochondrion found in the thin edge of an U2OS cell.
Corresponding to **Fig. 2b (I)**. Scale bar: 100 nm.

Movie S4. Tomogram of protein-coated membranes found in the thin edge of an

U2OS cell. Corresponding to **Fig. 2b (II)**. Scale bar: 100 nm.

Movie S5. Tomogram of ribosome-coated endoplasmic reticulum found in the

thin edge of an U2OS cell. Corresponding to **Fig. 2b (III)**. Scale bar: 100 nm.

Movie S6. Tomogram of a ribosome-associated vesicle attached to a microtubule

found in the thin edge of an U2OS cell. Corresponding to **Fig. 2b (IV)**. Scale bar:

100 nm.

Movie S7. Tomogram of two ribosome-associated vesicles connected by a thin

membrane tubule found in the thin edge of an U2OS cell. Corresponding to **Fig. 2b**

(V). Scale bar: 100 nm.

Movie S8. Tomogram of a large ribosome-associated vesicle with an internal LD

and with polysomes positioned on its top surface found in the thin edge of an U2OS

cell. Corresponding to **Fig. S3**. Scale bar: 100 nm.

Movie S9. Tomogram of a membrane-embedded compartment found in the thin

edge of an U2OS cell. Corresponding to **Fig. 2b (VI)**. Scale bar: 100 nm.

Movie S10. Tomogram of a membrane-embedded compartment found in the thin

edge of an U2OS cell. Corresponding to **Fig. 2b (VII)**. Scale bar: 100 nm.

Movie S11. Tomogram of a single lipid droplet found in the thin edge of an U2OS

cell. Corresponding to **Fig. 3a**. Scale bar: 100 nm.

Movie S12. Tomogram of a bunch of lipid droplets found in the thin edge of an

U2OS cell. Corresponding to **Fig. 3a**. Scale bar: 100 nm.

Movie S13. Tomogram of lipid droplets closely interacting with an endoplasmic

reticulum found in the thin edge of an U2OS cell. Corresponding to **Fig. 3b**. Scale

bar: 100 nm.

Movie S14. Tomogram of a lipid droplet impacting the lipid bilayer of an

endoplasmic reticulum found in the thin edge of an U2OS cell. Corresponding to

Fig. 3b. Scale bar: 100 nm.

Movie S15. Tomogram of a lipid droplet closely interacting with a mitochondrion

found in the thin edge of an U2OS cell. Corresponding to **Fig. 3c**. Scale bar: 100 nm.

Movie S16. Tomogram of a lipid droplet closely interacting with microtubules

found in the thin edge of an U2OS cell. Corresponding to **Fig. 3c**. Scale bar: 100 nm.

Movie S17. Tomogram of a lipid droplet putatively coated with COPI found in the

thin edge of an U2OS cell. Corresponding to **Fig. 3d**. Scale bar: 25 nm.

Movie S18 and Movie S19. Tomograms of lipid droplets coated with clathrin

found in the thin edge of an U2OS cell. Corresponding to **Fig. 3d**. Scale bars: 25 nm.

Methods

Sample preparation

Samples for cellular electron cryotomography were prepared as follows:
Quantifoil Au200 R2/2 grids were glow discharged for 90 s at 20 mA, coated with
fibronectin (25 μg/ml) and transferred to a 12 well plate in 3D printed cell culture
grid holders to facilitate handling (Fäßler et al., 2020). After sterilization for 30
min under UV light, the grids were seeded with U2OS cells (~6000 per well) and
incubated at 37^0C overnight. Before plunge freezing the grids were briefly washed
in PBS and 2.5 μl of 10 nm colloidal gold fiducial markers was applied.
Subsequently, the grids were back-blotted and vitrified in liquid ethane using a
Thermo Scientific Vitrobot.

Samples for electron cryotomography of purified macromolecular complexes
were prepared as follows: Quantifoil Au200 R1.2/1.3 grids were glow discharged
for 90 s at 20 mA. Then 4 μl of *Escherichia coli* 70S ribosomes at 250 nM was
applied and vitrified in liquid ethane-propane using a Leica EM GP2.

Data collection

Cellular electron cryotomography was performed using a Thermo Scientific
Glacios G1 TEM operating at 200 kV, equipped with a 4k x 4k Falcon 3EC direct
electron detection camera set in the linear mode without dose fractionation at a
magnification of 28k with parallel illumination conditions, corresponding to a
pixel size of 5.1 Å at the specimen level. Samples were tilted between
approximately -54^0 and $+54^0$ with a 3^0 increment using the dose symmetric
scheme in the TOMO software. The defocus was set between -5 and -7 μm and the
total dose for each tilt series was 140 e$Å^{-2}$.

Electron cryotomography of the 70S ribosomes was performed using a Thermo
Scientific Glacios G2 TEM operating at 200 kV, equipped with a 4k x 4k Falcon 4i
direct electron detection camera set in the counting mode at a magnification of
92k with parallel illumination conditions, corresponding to a pixel size of 1.6 Å at
the specimen level. Samples were tilted between approximately -54^0 and $+54^0$
with a 3^0 increment using the dose symmetric scheme and beam image shift data
collection strategy implemented in the (S)PACEtomo/SerialEM software was used
to record tilt series (Eisenstein et al., 2023, 2024; Mastronarde, 2005). The
defocus was set between -1.5 and -3 μm and 1.25 s exposures were divided into 8
frames each. The total dose for each tilt series was 130 e$Å^{-2}$.

Data analysis

Tomographic reconstruction from tilt series were calculated using RAPTOR (Amat
et al., 2008) and the IMOD tomography reconstruction package (Kremer et al.,

1996) followed by Weighted Back Projection or Simultaneous Iterative
Reconstruction Technique (SIRT) with the tomo3d package (Agulleiro &
Fernandez, 2011). Measurements of distances, including tomogram thicknesses,
were carried out within IMOD. Here, two model points were placed (to mark the
top and bottom surface of the biological density in the tomogram) and the distance
in 3D between them calculated in 3dmod. Single threshold surface representation
and movies were prepared using Chimera, with a similar threshold level to
exclude most background noise for each data set. EMAN2 (Chen et al., 2017,
2019b) was used for tomogram segmentations/annotations with the use of the
neural network and subtomogram averaging of *in situ* and purified ribosomes. The
neural network was trained to recognize different cellular features by selecting 5
tomograms with slightly various defocus values, then identifying in every
tomogram ~20 tiles containing the feature of interest and ~100 tiles deprived of
the feature, after which the content of the tiles was manually annotated and fed
into the neural network. CTFFIND4 (Rohou & Grigorieff, 2015) was used to obtain
information about CTF fit.

Data deposition

Representative 3D volumes have been deposited in the Electron Microscopy Data
Bank (EMDB) under accession numbers: EMD-51957, EMD-51959, EMD-51960,
EMD-51961, EMD-51962 and EMD-54370 and representative tilt series in the
EMPIAR database under accession numbers: EMPIAR-12763, EMPIAR-12764,
EMPIAR-12765, EMPIAR-12766 and EMPIAR-12767.

Declarations

Ethics approval and consent to participate

Not applicable, as no patient data was used in this research. Cell line used in this
study is not relevant material under the Human Research Act, so no ethical
approval was required.

6
7 *Consent for publication*

8 Not applicable

9
Availability of data and material

Data is provided within the manuscript or supplementary information files.
Additionally, representative 3D volumes have been deposited in the Electron
Microscopy Data Bank (EMDB) under accession numbers: EMD-51957, EMD-
51959, EMD-51960, EMD-51961, EMD-51962 and EMD-54370 and
representative tilt series in the EMPIAR database under accession numbers:
EMPIAR-12763, EMPIAR-12764, EMPIAR-12765, EMPIAR-12766 and EMPIAR-
12767.

Funding

This work was supported by the "Regenerative Mechanisms for Health - ReMedy"
project (MAB/2017/2) carried out within the International Research Agendas
programme of the Foundation for Polish Science co-financed by the European
Union under the European Regional Development Fund.

Competing interests

The author declares no competing interests.

References

Agulleiro, J. I., & Fernandez, J. J. (2011). Fast tomographic reconstruction on
multicore computers. *Bioinformatics*, *27*(4), 582–583.
https://doi.org/10.1093/bioinformatics/btq692

Amat, F., Moussavi, F., Comolli, L. R., Elidan, G., Downing, K. H., & Horowitz, M.
(2008). Markov random field based automatic image alignment for electron
tomography. *Journal of Structural Biology*, *161*(3), 260–275.
https://doi.org/10.1016/j.jsb.2007.07.007

Bartz, R., Zehmer, J. K., Zhu, M., Chen, Y., Serrero, G., Zhao, Y., & Liu, P. (2007).
Dynamic activity of lipid droplets: Protein phosphorylation and GTP-
mediated protein translocation. *Journal of Proteome Research*, *6*(8), 3256–
3265. https://doi.org/10.1021/pr070158j

Beck, M., & Baumeister, W. (2016). Cryo-Electron Tomography: Can it Reveal the
Molecular Sociology of Cells in Atomic Detail? In *Trends in Cell Biology* (Vol.
26, Issue 11, pp. 825–837). Elsevier Ltd.
https://doi.org/10.1016/j.tcb.2016.08.006

Brouwer, P. J. M., Perrett, H. R., Beaumont, T., Nijhuis, H., Kruijer, S., Burger, J. A.,
Bontjer, I., Lee, W.-H., Ferguson, J. A., Schauflinger, M., Müller-Kräuter, H.,
Sanders, R. W., Strecker, T., van Gils, M. J., & Ward, A. B. (2024). Defining
bottlenecks and opportunities for Lassa virus neutralization by structural
profiling of vaccine-induced polyclonal antibody responses. *Cell Reports*,
43(9), 114708. https://doi.org/10.1016/j.celrep.2024.114708

Burt, A., Toader, B., Warshamanage, R., von Kügelgen, A., Pyle, E., Zivanov, J.,
Kimanius, D., Bharat, T. A. M., & Scheres, S. H. W. (2024). An image
processing pipeline for electron cryo-tomography in RELION-5. *FEBS Open*
Bio. https://doi.org/10.1002/2211-5463.13873

Bykov, Y. S., Schaffer, M., Dodonova, S. O., Albert, S., rgen Plitzko, J. M.,
Baumeister, W., Engel, B. D., & Briggs, J. A. (2017). *The structure of the COPI*
coat determined within the cell. https://doi.org/10.7554/eLife.32493.001

Carter, S. D., Hampton, C. M., Langlois, R., Melero, R., Farino, Z. J., Calderon, M. J.,
Li, W., Wallace, C. T., Han Tran, N., Grassucci, R. A., Siegmund, S. E.,
Pemberton, J., Morgenstern, T. J., Bartolini, F., Volchuk, A., Murray, S. A.,
Aridor, M., Fish, K. N., Walter, P., … Freyberg, Z. (2020). Ribosome-associated
vesicles: A dynamic subcompartment of the endoplasmic reticulum in
secretory cells. In *Despoina Aslanoglou* (Vol. 10). https://www.science.org

Chan, L. M., Courteau, B. J., Maker, A., Wu, M., Basanta, B., Mehmood, H., Bulkley,
D., Joyce, D., Lee, B. C., Mick, S., Czarnik, C., Gulati, S., Lander, G. C., & Verba, K.
A. (2024). High-resolution single-particle imaging at 100–200 keV with the
Gatan Alpine direct electron detector. *Journal of Structural Biology*, *216*(3).
https://doi.org/10.1016/j.jsb.2024.108108

Chen, M., Bell, J. M., Shi, X., Sun, S. Y., Wang, Z., & Ludtke, S. J. (2019a). A complete
data processing workflow for cryo-ET and subtomogram averaging. *Nature*
Methods, *16*(11), 1161–1168. https://doi.org/10.1038/s41592-019-0591-8

Chen, M., Bell, J. M., Shi, X., Sun, S. Y., Wang, Z., & Ludtke, S. J. (2019b). A complete
data processing workflow for cryo-ET and subtomogram averaging. *Nature*
Methods, *16*(11), 1161–1168. https://doi.org/10.1038/s41592-019-0591-8

Chen, M., Dai, W., Sun, S. Y., Jonasch, D., He, C. Y., Schmid, M. F., Chiu, W., & Ludtke,
S. J. (2017). Convolutional neural networks for automated annotation of
cellular cryo-electron tomograms. *Nature Methods*, *14*(10), 983–985.
https://doi.org/10.1038/nmeth.4405

Daro, E., Sheff, D., Gomez, M., Kreis, T., & Mellman, I. (1997). Inhibition of
Endosome Function in CHO Cells Bearing a Temperature-sensitive Defect in
the Coatomer (COPI) Component-COP. In *The Journal of Cell Biology* (Vol.
139, Issue 7). http://www.jcb.org

Dudka, W., Salo, V. T., & Mahamid, J. (2024). Zooming into lipid droplet biology
through the lens of electron microscopy. In *FEBS Letters* (Vol. 598, Issue 10,
pp. 1127–1142). John Wiley and Sons Inc. https://doi.org/10.1002/1873-
3468.14899

Eisenstein, F., Fukuda, Y., & Danev, R. (2024). Smart parallel automated cryo-
electron tomography. *Nature Methods*, *21*(9), 1612–1615.
https://doi.org/10.1038/s41592-024-02373-9

Eisenstein, F., Yanagisawa, H., Kashihara, H., Kikkawa, M., Tsukita, S., & Danev, R.
(2023). Parallel cryo electron tomography on in situ lamellae. *Nature*
Methods, *20*(1), 131–138. https://doi.org/10.1038/s41592-022-01690-1

Faini, M., Prinz, S., Beck, R., Schorb, M., Riches, J. D., Bacia, K., Brügger, B., Wieland,
F. T., & Briggs, J. A. G. (2012). The structures of COPI-coated vesicles reveal
alternate coatomer conformations and interactions. *Science*, *336*(6087),
1451–1454. https://doi.org/10.1126/science.1221443

Fan, H., & Tan, Y. (2024). Lipid Droplet–Mitochondria Contacts in Health and
Disease. In *International Journal of Molecular Sciences* (Vol. 25, Issue 13).
Multidisciplinary Digital Publishing Institute (MDPI).
https://doi.org/10.3390/ijms25136878

Fäßler, F., Zens, B., Hauschild, R., & Schur, F. K. M. (2020). 3D printed cell culture
grid holders for improved cellular specimen preparation in cryo-electron
microscopy. *Journal of Structural Biology*, *212*(3).
https://doi.org/10.1016/j.jsb.2020.107633

Förster, F., Han, B. G., & Beck, M. (2010). Visual Proteomics. In *Methods in*
Enzymology (Vol. 483, Issue C, pp. 215–243). Academic Press Inc.
https://doi.org/10.1016/S0076-6879(10)83011-3

Foster, H. E., Santos, C. V., & Carter, A. P. (2022). A cryo-ET survey of microtubules and intracellular compartments in mammalian axons. *Journal of Cell Biology*, *221*(2). https://doi.org/10.1083/jcb.202103154

Fotin, A., Kirchhausen, T., Grigorieff, N., Harrison, S. C., Walz, T., & Cheng, Y. (2006). Structure determination of clathrin coats to subnanometer resolution by single particle cryo-electron microscopy. *Journal of Structural Biology*, *156*(3), 453–460. https://doi.org/10.1016/j.jsb.2006.07.001

Fry, M. Y., Navarro, P. P., Hakim, P., Ananda, V. Y., Qin, X., Landoni, J. C., Rath, S., Inde, Z., Lugo, C. M., Luce, B. E., Ge, Y., Mcdonald, J. L., Ali, I., Ha, L. L., Kleinstiver, B. P., Chan, D. C., Sarosiek, K. A., & Chao, L. H. (2024). *In situ architecture of Opa1-dependent mitochondrial cristae remodeling.* https://doi.org/10.1101/2023.01.16.524176

Fujimoto, T., & Parton, R. G. (2011). Not just fat: The structure and function of the lipid droplet. *Cold Spring Harbor Perspectives in Biology*, *3*(3), 1–17. https://doi.org/10.1101/cshperspect.a004838

Ganeva, I., Lim, K., Boulanger, J., Hoffmann, P. C., Muriel, O., Borgeaud, A. C., Hagen, W. J. H., Savage, D. B., & Kukulski, W. (2023). The architecture of Cidec-mediated interfaces between lipid droplets. *Cell Reports*, *42*(2). https://doi.org/10.1016/j.celrep.2023.112107

Gross, S. P., Welte, M. A., Block, S. M., & Wieschaus, E. F. (2000). Dynein-mediated Cargo Transport In Vivo: A Switch Controls Travel Distance. In *The Journal of Cell Biology* (Vol. 148, Issue 5). http://www.jcb.org

He, X., Kostin, R., Knight, E., Han, M. G., Mun, J., Bozovic, I., Jing, C., & Zhu, Y. (2024). Development of a liquid-helium free cryogenic sample holder with mK temperature control for autonomous electron microscopy. *Ultramicroscopy*, *267*. https://doi.org/10.1016/j.ultramic.2024.114037

Herzik, M. A., Wu, M., & Lander, G. C. (2017). Achieving better-than-3-Å resolution by single-particle cryo-EM at 200 keV. *Nature Methods*, *14*(11), 1075–1078. https://doi.org/10.1038/nmeth.4461

Hugenroth, M., & Bohnert, M. (2020). Come a little bit closer! Lipid droplet-ER contact sites are getting crowded. In *Biochimica et Biophysica Acta - Molecular Cell Research* (Vol. 1867, Issue 2). Elsevier B.V. https://doi.org/10.1016/j.bbamcr.2019.118603

Kilwein, M. D., & Welte, M. A. (2019). Lipid droplet motility and organelle contacts. *Contact (Thousand Oaks)*, *585*, 276–296.

Kremer, J. R., Mastronarde, D. N., & Mcintosh, J. R. (1996). *Computer Visualization of Three-Dimensional Image Data Using IMOD*.

Kühlbrandt, W. (2014). The resolution revolution. In *Science* (Vol. 343, Issue 6178, pp. 1443–1444). American Association for the Advancement of Science. https://doi.org/10.1126/science.1251652

Mahamid, J., Tegunov, D., Maiser, A., Arnold, J., Leonhardt, H., Plitzko, J. M., & Baumeister, W. (2019). Liquid-crystalline phase transitions in lipid droplets are related to cellular states and specific organelle association. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(34), 16866–16871. https://doi.org/10.1073/pnas.1903642116

Martynowycz, M. W., Clabbers, M. T. B., Unge, J., Hattne, J., & Gonen, T. (2021). *Benchmarking the ideal sample thickness in cryo-EM. 118*, 2108884118. https://doi.org/10.1073/pnas.2108884118/-/DCSupplemental

Mastronarde, D. N. (2005). Automated electron microscope tomography using robust prediction of specimen movements. *Journal of Structural Biology*, *152*(1), 36–51. https://doi.org/10.1016/j.jsb.2005.07.007

McMullan, G. I., Naydenova, K. I., Mihaylov, D., Yamashita, K. I., Peet ID, M. J., Wilson, H. I., Dickerson ID, J. L., Chen, S. I., Cannone, G. I., Lee, Y. I., Hutchings ID, K. A., Gittins, O. I., Sobhy ID, M. A., Wells, T. I., El-Gomati ID, M. M., Dalby, J., Meffert, M. I., Schulze-Briese, C. I., Henderson, R. I., & Russo, C. J. (2023). *Structure determination by cryoEM at 100 keV*. https://doi.org/10.1073/pnas

Medalia, O., Weber, I., Frangakis, A. S., Nicastro, D., Gerisch, G., & Baumeister, W. (2002). Macromolecular Architecture in Eukaryotic Cells Visualized by Cryoelectron Tomography. *Science*, *5596*, 1209–1213. https://doi.org/DOI:10.1126/science.1076184

Navaratnarajah, T., Anand, R., Reichert, A. S., & Distelmaier, F. (2021). The relevance of mitochondrial morphology for human disease. In *International Journal of Biochemistry and Cell Biology* (Vol. 134). Elsevier Ltd. https://doi.org/10.1016/j.biocel.2021.105951

Noble, A. J. (2024). *HighEnd CryoEM Worldwide*.

Nogales, E., & Mahamid, J. (2024). Bridging structural and cell biology with cryo-electron microscopy. In *Nature* (Vol. 628, Issue 8006, pp. 47–56). Nature Research. https://doi.org/10.1038/s41586-024-07198-2

Nonoyama, T., Nojima, D., Maeda, Y., Noda, M., Yoshino, T., Matsumoto, M., Bowler, C., & Tanaka, T. (2019). Proteomics analysis of lipid droplets indicates involvement of membrane trafficking proteins in lipid droplet breakdown in the oleaginous diatom Fistulifera solaris. *Algal Research*, *44*. https://doi.org/10.1016/j.algal.2019.101660

Olarte, M. J., Swanson, J. M. J., Walther, T. C., & Farese, R. V. (2022). The CYTOLD and ERTOLD pathways for lipid droplet–protein targeting. In *Trends in Biochemical Sciences* (Vol. 47, Issue 1, pp. 39–51). Elsevier Ltd. https://doi.org/10.1016/j.tibs.2021.08.007

Olzmann, J. A., & Carvalho, P. (2019). Dynamics and functions of lipid droplets. In *Nature Reviews Molecular Cell Biology* (Vol. 20, Issue 3, pp. 137–155). Nature Publishing Group. https://doi.org/10.1038/s41580-018-0085-z

Peet, M. J., Henderson, R., & Russo, C. J. (2019). The energy dependence of
contrast and damage in electron cryomicroscopy of biological molecules.
Ultramicroscopy, *203*, 125–131.
https://doi.org/10.1016/j.ultramic.2019.02.007

Punjani, A., & Fleet, D. J. (2023). 3DFlex: determining structure and motion of
flexible proteins from cryo-EM. *Nature Methods*, *20*(6), 860–870.
https://doi.org/10.1038/s41592-023-01853-8

Rice, W. J., Cheng, A., Noble, A. J., Eng, E. T., Kim, L. Y., Carragher, B., & Potter, C. S.
(2018). Routine determination of ice thickness for cryo-EM grids. *Journal of*
Structural Biology, *204*(1), 38–44.
https://doi.org/10.1016/j.jsb.2018.06.007

Rigort, A., & Plitzko, J. M. (2015). Cryo-focused-ion-beam applications in
structural biology. In *Archives of Biochemistry and Biophysics* (Vol. 581, pp.
122–130). Academic Press Inc. https://doi.org/10.1016/j.abb.2015.02.009

Rohou, A., & Grigorieff, N. (2015). CTFFIND4: Fast and accurate defocus
estimation from electron micrographs. *Journal of Structural Biology*, *192*(2),
216–221. https://doi.org/10.1016/j.jsb.2015.08.008

Russo, C. J., Dickerson, J. L., & Naydenova, K. (2022). Cryomicroscopy in situ:
what is the smallest molecule that can be directly identified without labels
in a cell? *Faraday Discussions*, *240*, 277–302.
https://doi.org/10.1039/d2fd00076h

Schwab, J., Kimanius, D., Burt, A., Dendooven, T., & Scheres, S. H. W. (2024).
DynaMight: estimating molecular motions with improved reconstruction
from cryo-EM images. *Nature Methods*. https://doi.org/10.1038/s41592-
024-02377-5

Schwartz, O., Axelrod, J. J., Campbell, S. L., Turnbaugh, C., Glaeser, R. M., & Müller,
H. (2019). Laser phase plate for transmission electron microscopy. *Nature*
Methods, *16*(10), 1016–1020. https://doi.org/10.1038/s41592-019-0552-2

Szwedziak, P., & Pilhofer, M. (2019). Bidirectional contraction of a type six
secretion system. *Nature Communications*, *10*(1).
https://doi.org/10.1038/s41467-019-09603-1

Szwedziak, P., Wang, Q., Bharat, T. A., Tsim, M., & Löwe, J. (2014). Architecture of
the ring formed by the tubulin homologue FtsZ in bacterial cell division.
ELife, *3*. https://doi.org/10.7554/eLife.04601

Tegunov, D., Xue, L., Dienemann, C., Cramer, P., & Mahamid, J. (2021). Multi-
particle cryo-EM refinement with M visualizes ribosome-antibiotic complex
at 3.5 Å in cells. *Nature Methods*, *18*(2), 186–193.
https://doi.org/10.1038/s41592-020-01054-7

Thiam, A. R., Antonny, B., Wang, J., Delacotte, J., Wilfling, F., Walther, T. C., Beck,
R., Rothman, J. E., & Pincet, F. (2013). COPI buds 60-nm lipid droplets from

reconstituted water-phospholipid- triacylglyceride interfaces, suggesting a tension clamp function. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(33), 13244–13249. https://doi.org/10.1073/pnas.1307685110

Tian, Y., Liang, R., Kumar, A., Szwedziak, P., & Viles, J. H. (2020). 3D-visualization of amyloid-β oligomer and fibril interactions with lipid membranes by cryo-electron tomography. In *bioRxiv*. https://doi.org/10.1101/2020.07.21.214072

Träger, T. K., Kyrilis, F. L., Hamdi, F., Tüting, C., Alfes, M., Hofmann, T., Schmidt, C., & Kastritis, P. L. (2024). Disorder-to-order active site capping regulates the rate-limiting step of the inositol pathway. *Proceedings of the National Academy of Sciences of the United States of America*, *121*(34). https://doi.org/10.1073/pnas.2400912121

Valm, A. M., Cohen, S., Legant, W. R., Melunis, J., Hershberg, U., Wait, E., Cohen, A. R., Davidson, M. W., Betzig, E., & Lippincott-Schwartz, J. (2017). Applying systems-level spectral imaging and analysis to reveal the organelle interactome. In *Nature* (Vol. 546, Issue 7656, pp. 162–167). Nature Publishing Group. https://doi.org/10.1038/nature22369

Wilfling, F., Thiam, A. R., Olarte, M. J., Wang, J., Beck, R., Gould, T. J., Allgeyer, E. S., Pincet, F., Bewersdorf, J., Farese, R. V., & Walther, T. C. (2014). Arf1/COPI machinery acts directly on lipid droplets and enables their connection to the ER for protein targeting. *ELife*, *2014*(3). https://doi.org/10.7554/eLife.01607

Wilfling, F., Wang, H., Haas, J. T., Krahmer, N., Gould, T. J., Uchida, A., Cheng, J. X., Graham, M., Christiano, R., Fröhlich, F., Liu, X., Buhman, K. K., Coleman, R. A., Bewersdorf, J., Farese, R. V., & Walther, T. C. (2013). Triacylglycerol synthesis enzymes mediate lipid droplet growth by relocalizing from the ER to lipid droplets. *Developmental Cell*, *24*(4), 384–399. https://doi.org/10.1016/j.devcel.2013.01.013

Wolf, S., Mutsafi, yael, Dadosh, T., Ilani, T., Lansky, Z., Horowitz, B., Rubin, S., Elbaum, M., & Fass, D. (2017). 3D visualization of mitochondrial solid-phase calcium stores in whole cells. *ELife*. https://doi.org/10.7554/eLife.29929.001

Wolff, G., Limpens, R. W. A. L., Zevenhoven-Dobbe, J. C., Laugks, U., Zheng, S., De Jong, A. W. M., Koning, R. I., Agard, D. A., Grünewald, K., Koster, A. J., Snijder, E. J., & Bárcena, M. (2020). A molecular pore spans the double membrane of the coronavirus replication organelle. *Science*, 1395–1398. https://doi.org/10.1126/science.abd3629

Wozny, M. R., Di Luca, A., Morado, D. R., Picco, A., Khaddaj, R., Campomanes, P., Ivanović, L., Hoffmann, P. C., Miller, E. A., Vanni, S., & Kukulski, W. (2023). In situ architecture of the ER–mitochondria encounter structure. *Nature*, *618*(7963), 188–192. https://doi.org/10.1038/s41586-023-06050-3

Wu, G. H., Smith-Geater, C., Galaz-Montoya, J. G., Gu, Y., Gupte, S. R., Aviner, R.,
Mitchell, P. G., Hsu, J., Miramontes, R., Wang, K. Q., Geller, N. R., Hou, C.,
Danita, C., Joubert, L. M., Schmid, M. F., Yeung, S., Frydman, J., Mobley, W.,
Wu, C., … Chiu, W. (2023). CryoET reveals organelle phenotypes in
huntington disease patient iPSC-derived and mouse primary neurons.
Nature Communications, *14*(1). https://doi.org/10.1038/s41467-023-
36096-w

Wu, J., Liu, C., Wang, A. J., Gao, Y. Z., Fu, L. T., Liu, Z., Dickerson, J. L., Russo, C. J., &
Wang, P. (2025). Chromatic aberration (Cc) corrected cryo-EM: The
structure of pseudorabies virus (PRV) using both zero-loss and energy loss
electrons. *Ultramicroscopy*, *276*.
https://doi.org/10.1016/j.ultramic.2025.114182

Zhang, S., Peng, X., Yang, S., Li, X., Huang, M., Wei, S., Liu, J., He, G., Zheng, H., Yang,
L., Li, H., & Fan, Q. (2022). The regulation, function, and role of lipophagy, a
form of selective autophagy, in metabolic disorders. In *Cell Death and
Disease* (Vol. 13, Issue 2). Springer Nature.
https://doi.org/10.1038/s41419-022-04593-3

Ziemianowicz, D. S., & Kosinski, J. (2022). New opportunities in integrative
structural modeling. In *Current Opinion in Structural Biology* (Vol. 77).
Elsevier Ltd. https://doi.org/10.1016/j.sbi.2022.102488

Figure 1

b)

Data collection summary	
#imaged grids	9
#imaged cells	64
#tomograms	228
average thickness	240±75 nm

Figure 2

a)

b)

Figure 3

Figure 4

a)

EMDB entries by cryoTEM instrument and modality						
	SPA cryoEM		cryoET		Others	
Year	Glacios	Krios	Glacios	Krios	Glacios	Krios
2020	24	2519	0	243	0	112
2021	50	3239	0	260	1	165
2022	149	4233	0	449	3	248
2023	318	5379	4	837	18	333

b)

c)

d)

1° **Local cryoEM facilities**
200 kV instruments suitable for SPA cryoEM and cryoET data collection and screening

2° **National/international cryoEM hubs**
300 kV cryoET purpose-built instruments fitted with laser phase plate/Cc corrector/liquid helium cryostage.
Complex cryoET sample prep (HPF, cryo-pFIB-SEM with lift-out)

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

Figure S7

resolution at FSC$_{0.5}$ ~ 15 Å

Fourier Shell Correlation

Spatial frequency [1/Å]

90º