1 **A genomic resource for exploring bacterial-viral dynamics in seagrass ecosystems**
2

3 Cassandra L. Ettinger[1,#], Jason E. Stajich[1,2]
4
5 [1]Department of Microbiology and Plant Pathology, University of California, Riverside, CA, United
6 States
7
8 [2]Institute for Integrative Genome Biology, University of California, Riverside, Riverside, CA,
9 United States
10
11 [#]**Corresponding Author:** Cassandra L. Ettinger, clettinger@ucdavis.edu

12
13 **Keywords:** *Zostera marina*, eelgrass, metagenomics, viral discovery, viral diversity, phage,

14 bacteria, carbon cycling, metagenome-assembled genomes

Abstract:

Background: Seagrasses are globally distributed marine flowering plants that play foundational

roles in coastal environments as ecosystem engineers. While research efforts have explored

various aspects of seagrass-associated microbial communities, including describing the

diversity of bacteria, fungi and microbial eukaryotes, little is known about viral diversity in these

communities.

Results: To begin to address this, we leveraged metagenomic sequencing data to generate a

catalog of bacterial metagenome-assembled genomes (MAGs) and phage genomes from the

leaves of the seagrass, *Zostera marina*. We expanded the robustness of this viral catalog by

incorporating publicly available metagenomic data from seagrass ecosystems. The final MAG

set represents 85 high-quality draft and 62 medium-quality draft bacterial genomes. While the

viral catalog represents 354 medium-quality, high-quality, and complete viral genomes.

Predicted auxiliary metabolic genes in the final viral catalog had putative annotations largely

related to carbon utilization, suggesting a possible role for phage in carbon cycling in seagrass

ecosystems.

Conclusions: These genomic resources provide initial insight into bacterial-viral interactions in

seagrass meadows and are a foundation on which to further explore these critical interkingdom

interactions. These catalogs highlight a possible role for viruses in carbon cycling in seagrass

beds which may have important implications for blue carbon management and climate change

mitigation.

Background:

Seagrasses are submerged marine flowering plants that have critical roles as foundation

species in coastal ecosystems worldwide. They provide essential ecosystem services such as

stabilizing the seafloor, filtering pollutants, supporting fisheries, and driving biogeochemical

cycles [1–3]. One of their most significant contributions is their ability to sequester carbon in

both their tissues and surrounding sediments (i.e., blue carbon) [2,4]. Despite their ecological

importance, seagrass ecosystems are under increasing threat from pollution, climate change,

and coastal development. Preserving these ecosystems is critical not only for their carbon

sequestration potential but also for maintaining the biodiversity and ecological services they

provide to coastal communities.

In recent years, there has been growing recognition of the importance of microorganisms in

maintaining the health of plants [5–7]. Studies have begun to describe the composition and

structure of the bacterial community associated with seagrasses, particularly *Zostera marina*,

the dominant seagrass in the Northern hemisphere [8–14]. These bacterial communities are

thought to have important roles in facilitating nitrogen and sulfur cycling to benefit seagrass

growth and survival [8,9,11,12,15–17].

While research has largely focused on the bacterial component of these microbial communities,

advances in sequencing and bioinformatics now allow us to explore the role of viruses in these

systems. Phages, in particular, have been shown to influence global ocean and soil

biogeochemical cycles by modulating host population dynamics through auxiliary metabolic

genes (AMGs), which can directly alter bacterial metabolism to increase overall host fitness [18–

21]. However, the diversity and ecological roles of phage in seagrass ecosystems remains

largely unexplored.

In this study, we generated a catalog of viruses from seagrass-associated metagenomic

samples, and then investigated host-viral dynamics of viral operational taxonomic units (vOTUs)

and bacterial metagenome-assembled genomes (MAGs) generated from *Z. marina* leaf

samples from Bodega Bay, CA. Specifically our objectives were to: (i) create a catalog of

vOTUs from *Z. marina* and other seagrass species using publicly available metagenomic

sequencing data, (ii) assemble a catalog of bacterial MAGs from *Z. marina* leaf tissue, and (iii)

explore bacterial-phage dynamics with a focus on exploring AMGs involved in nitrogen and

sulfur metabolisms.

107
108   **Methods:**
109
110   *Sequence generation*
111
We extracted DNA from epiphytic washes from *Z. marina* leaves as part of previous work

focused on characterizing the mycobiome using high-throughput sequencing of the ITS2 region

[22]. We chose three DNA extracts from that work here for deep metagenomic sequencing with

the goal of obtaining high quality metagenome-assembled genomes. We provided DNA to the

UC Davis Genome Center DNA Technologies Core for sequencing and library preparation. DNA

libraries were sequenced on an Illumina HiSeq4000 to generate 150 bp paired-end reads.

118
119   *Metagenomic processing*
120
We trimmed sequence reads using bbDuk v. 37.68 [23] with the following parameters: qtrim=rl

trimq=10 maq=10. We then mapped against and removed any reads from the metagenomes

matching the available genome for *Z. marina* v. 3.1 [24] using bowtie2 v. 2.4.5 [25] and

samtools v. 1.11 [26]. We co-assembled the remaining reads from all three metagenomic

samples using MEGAHIT v. 1.2.9 [27].

We identified and assessed metagenome-assembled genomes (MAGs) using the anvi'o v. 7.2

workflow [28]. First, we used bowtie2 v. 2.4.5 [25] and samtools v. 1.11 [26] to obtain read

coverage for each metagenomic sample against the assembly. Then we used "anvi-gen-

contigs-database" to generate a database from the co-assembly and to predict open-reading

frames using Prodigal v. 2.6.3 [29]. This command also identifies single-copy bacterial [30],

archaeal [31], and protista [32] genes using HMMER v. 3.2.1 [33] and ribosomal RNA genes

using barrnap [34]. We predicted taxonomic assignments for each gene call using Kaiju v. 1.8.2

[35] with the NCBI BLAST non redundant protein database nr including fungi and microbial

eukaryotes v. v. 2020-05-25. For each individual metagenomic sample, we then used "anvi-

profile" to construct an anvi'o profile for contigs >1 kbp with the "–cluster-contigs" option. Next

we ran several automatic binning algorithms including MetaBAT2 v. 2.15, MaxBin v. 2.2.1,

BinSanity v.0.5.4 and CONCOCT v. 1.1.0 [36–39] to generate preliminary sets of bacterial

MAGs. We provided the resulting set from each algorithm to DAStool v. 1.1.2 [40] to generate a

single optimal MAG set. MAGs from this set were then manually assessed for contamination

and refined using "anvi-refine". After manual refinement, MAGs were further de-contaminated

using MAGpurify v. 2.1.2 [41].

We assessed MAG completeness and contamination using CheckM v. 1.2.1 [42] and CheckM2

v. 1.0.1 [43]. In this work we report all identified MAGs with >80% completion and <10%

contamination. We refer to MAGs as high quality if they were >90% complete with <5%

contamination, and medium quality if they were >50% complete with <10% contamination per

established guidelines [44]. To obtain a putative taxonomy for each MAG, we used GTDB-Tk v.

2.2.6 [45], which uses a combination of average nucleotide identity and phylogenetic placement

in the context of the Genome Taxonomy Database to taxonomically identify MAGs.

Viral identification

In order to have a robust dataset to compare our recovered virus catalog in the context of other

studies, we downloaded publicly available data from NCBI GenBank from nine studies

[11,12,17,46–51] representing 65 metagenomic and 18 metatranscriptomic samples collected

from seagrass environments (Table S1).

We filter and trimmed public metagenomic data using bbDuk v. 37.68 [23] with the following

parameters: ktrim=r k=23 mink=11 hdist=1 tpe tbo qtrim=rl trimq=10 maq=10. Then for each

study, samples were co-assembled using MEGAHIT v. 1.2.9. We used the co-assemblies for

each study, as well as the co-assembly from this study, when identifying viral sequences using

a workflow similar to Guo et al. [52]

Briefly, we identified viral sequences from metagenomic co-assemblies using VirSorter2 v. 2.2.3

[53], a tool that uses multiple random forest classifiers to predict whether a sequence contains a

DNA or RNA virus, with the following parameters: --min-length 5000, --min-score 0.5, --include-

groups dsDNAphage,RNA,ssDNA,lavidaviridae. We then ran CheckV v0.8.1 [54] on the

VirSorter2 predicted viral sequences using the "end_to_end" workflow. To be conservative in

our analyses, we removed viral sequences with a CheckV quality score of "not-determined" and

"low-quality" prior to downstream analysis. We then ran VirSorter2 again on the viral sequences

from the CheckV workflow with the --prep-for-dramv option. We used DRAM-v v. 1.2.2 [55] to

"annotate" viral sequences and then "distill" annotations into predicted auxiliary metabolic genes

(AMGs) for phage.

175
Virus clustering and analysis

We clustered viral sequences ≥ 10 kbp in length into 95% similarity viral operational taxonomic

units (vOTUs) using dRep v. 3.2.2 [56]. We used Prodigal v. 2.6.3 [29] to predict open reading

frames in vOTUs using the -p meta option. We then provided the predicted proteins from the

phage vOTUs to VContact2 v. 0.9.19, as well as predicted proteins from the INPHARED August

2023 viral reference database, to generate viral clusters (VCs) based on viral gene-sharing

networks [57,58]. We further used geNomad v. 1.7.4 to assign taxonomy to phage vOTUs [59].

We used iPHoP v. 1.3.2 [60], which integrates across multiple methods in a machine learning

framework to assign host taxonomy at the genus level, to predict host-virus linkages using a

combination of the final bacterial MAG collection from this study and iPHoP's reference host

database.

We mapped reads from each metagenome to vOTUs using bowtie2 --sensitive with a

minid=0.90 to quantify vOTU relative abundance [61]. We then used SAMtools and BEDTools

genomecov to obtain coverage estimates for each vOTU across each individual metagenomic

sample [26,62]. We used coverM in contig mode to parse bam files and calculate the trimmed

pileup coverage (tpmean) of vOTUs which displayed ≥ 75% coverage over the length of the viral

sequence. Thresholds for analysis of vOTUs were based on community guidelines for length

(i.e. ≥ 10 kbp), similarity (i.e. ≥ 95% similarity), and detection (i.e. ≥ 75% of the viral genome

length covered ≥ 1x by reads at ≥ 90% average nucleotide identity) [63,64]. The viral relative

abundance (tpmean), CheckV quality, geNomad taxonomy, iPHoP host-prediction, MAG

diversity, and DRAM-v annotation results were analyzed and visualized in R v. 4.3.0 [65] using

the tidyverse v. 2.0.0 and phyloseq v. 1.44.0 [66,67].

To compare viral diversity between metagenomic samples (i.e. beta diversity), we calculated the

Hellinger distance, the Euclidean distance of Hellinger transformed relative abundance (tpmean)

data. We performed Hellinger transformations using the transform function in the microbiome v.

1.22.0 package [68], calculated the Hellinger distance using the ordinate function in phyloseq,

and then visualized these distances using principal-coordinate analysis (PCoA).

206
Results and Discussion

208
Refined viral catalog from seagrass ecosystems

210
In total, we recovered 17,145 predicted viral genomic fragments which we clustered at 95%

average nucleotide identity into 3,633 vOTUs. To ensure a high quality viral catalog, we filtered

this dataset further using community thresholds for length and quality [63,64]. The refined viral

collection represents 354 viral sequences comprising 351 double-stranded DNA phage, two

lavidaviridae, and one RNA phage based on VirSorter2 random forest classification, with each

sequence representing a unique vOTU (Table S2). Of these sequences, nine are integrated

prophage (Figure 1A). The final catalog includes 44 complete, 28 high-quality and 282 medium-

quality draft viral genomic fragments (Figure 1B).

To explore the taxonomy of the viral sequences reported here, we used two complementary

approaches: VContact2 [57], a cluster based method, and geNomad [59], an alignment based

tool. We ran VContact2 [57] with INPHARED [58] reference genomes to cluster phage vOTUs

into 136 VCs, which represent genus-level groupings based on gene-sharing networks. Of

these, 35 VCs represented clusters that contained reference genomes, while the other 101 VCs

were unique, potentially representing novel phage genera (Figure S1A). However, 148 vOTUs

could not be assigned confidently to any VC.

Taxonomic classification of viruses was further refined using geNomad [59]. Most vOTUs

(98.58% of vOTUs) were assigned to the *Caudoviricetes* class of tailed double-stranded DNA

bacteriophage (Figure S1B). However, viral taxonomy is currently in revision [69], and the

majority of sequences could not be confidently placed into finer taxonomic ranks: 331 vOTUs

were unclassified at the order level, and 352 at the family level. While geNomad provided

higher-level taxonomic predictions for 351 of the 354 vOTUs, more comprehensive classification

awaits future updates to formal viral taxonomy definitions.

The recovery of phage genomes from metagenomes varied across experimental studies (Figure

1C). Of the 354 viral sequences in the final catalog, 50 were recovered from the new

metagenomic data in this study, with the remainder derived from publicly available

metagenomes. Studies that contributed more viral sequences to the final catalog generally

employed deeper sequencing, with average depths ranging from 9.7 to 28.2 Gb per sample,

compared to studies with few recovered viral sequences (1.2 to 9.5 Gb per sample).

When we examined viral community composition across metagenomes, clustering by scientific

study was evident (Figure 1D). Notably, the viral communities in the metagenomes generated in

this study formed a distinct cluster. However, technical differences (e.g., sequencing depth)

across datasets prevented us from drawing broader conclusions regarding viral diversity based

on seagrass species or tissue type.

Despite the use of deep metagenomic sequencing and the integration of public metagenomes,

we recovered a relatively small, refined catalog of viral sequences. This highlights the limitations

of relying solely on short-read metagenomic sequencing to capture viral diversity, at least in

seagrass ecosystems. Viromics approaches (e.g., [70]) may provide more comprehensive

insights into these viral communities, as even with improved bioinformatics pipelines, true viral

diversity and abundance remains elusive.

255
256 *MAG collection reflects abundant bacterial groups on Z. marina leaves*
257
We assembled 147 total bacterial MAGs including 85 high-quality draft MAGs (>90%

completion, <5% contamination), and 62 medium-quality MAGs (>80% completion, <10%

contamination) (Table S3). These MAGs largely belong to the following taxonomic classes

(Figure 2A): Alphaproteobacteria (43), Gammaproteobacteria (35), Bacteroidia (29), and

Plactomycetia (16). Within these classes, the most frequently recovered orders were:

Rhodobacterales (28), Pirellulales (14), Flavobacteriales (12), Chitinophagales (11) and

Pseudomonadales (11). Notably, using GTDB-Tk 30 MAGs could not be assigned to any known

genus, six were unclassified at the family level, and one lacked an assignment at the order

level, highlighting potential evolutionary novelty compared to the current GTDB database. These

findings suggest that seagrass ecosystems may harbor novel bacterial species that remain

uncharacterized.

The taxonomic distribution of the recovered MAGs is broadly consistent with previous DNA-

based surveys of *Zostera*-associated bacteria. These studies also identified

Alphaproteobacteria, Gammaproteobacteria, and Bacteroidia as dominant classes, along with

orders such as Rhodobacterales and Flavobacteriales [9–13]. Further a global study of *Z.*

marina reported that Planctomycetia was enriched on leaves relative to surrounding water [8].

Together, these results suggest that this MAG collection complements previous DNA-based

surveys, providing an opportunity for exploring deeper insight into the genomics and functional

potential of abundant bacterial taxa within *Z. marina* ecosystems.

278
279   *Recovered phage largely predicted to infect most frequent MAG phyla*
280
We used iPHoP [60] to predict bacteria-virus linkages but were only able to predict bacterial

hosts for 18 vOTUs (Figure 2B). Of these, hosts generally were distributed across the most

frequently recovered MAG classes (i.e., Alphaproteobacteria, Gammaproteobacteria, and

Bacteroidia). While the MAG collection was added to the iPHoP database to enable prediction

of direct links between MAGs and viruses, only one MAG, SGMAG-05 (a Saprospiraceae

bacterium in the class Bacteroidia) was predicted to be host to a virus from the high-quality

catalog. This phage, vOTU566, was identified as a prophage belonging to the Caudoviricetes.

The remaining host-virus predictions were derived from iPHoP's default genome library rather

than the MAG collection. The limited number of host-virus links identified between the MAG and

viral collections suggests that much more work remains to uncover these interactions within

seagrass ecosystems. Future studies should endeavor to sequence the bacterial community

alongside deeper virome sequencing to better capture viral diversity. Additionally, employing

physical linking techniques such as Hi-C may help characterize novel host-viral interactions.

294
295   *Auxiliary metabolic genes point to phage role in carbon cycling*
296
We used DRAM-v [55] to explore predicted AMG functions in the viral collection (Figure 2C).

Over half of putative AMGs were unannotated (57.47%), similar to other recent studies of phage

from the environment [71,72]. Given the importance of nitrogen and sulfur cycling in seagrass

ecosystems, we searched the annotated AMGs for putative functions related to these

processes. However no annotated AMGs had predictions related to nitrogen fixation or sulfur

cycling. This absence could be biological (e.g., such genes may be more likely to be present in

root or rhizosphere samples vs. leaves) or could reflect the limitations of recovering viruses from

metagenomic datasets. Virome sequencing will likely be necessary to confirm whether viruses

play a role in these nutrient cycles.

In contrast, we identified a variety of predicted AMGs with putative annotations of functions

related to carbon utilization, particularly carbohydrate-active enzymes (CAZymes) involved in

organic carbon cycling, sugar processing and plant degradation (Figure 2D). Seagrass beds are

known as hot spots for carbon sequestration and key contributors to blue carbon storage [2,4].

Additionally previous work in the Mediterranean seagrass *Posidonia oceanica* has shown that

rhizosphere sediments are enriched in sugars [46]. These findings may suggest a possible role

for viruses in carbon cycling in seagrass beds, which may have important implications for blue

carbon management and climate change mitigation.

315
316   **Conclusion**

317
This study provides a valuable community resource and catalog of refined seagrass-associated

bacterial and viral genomes, serving as a genomic foundation for future research. These

comprehensive collections likely capture the most abundant groups associated with the leaves

of *Z. marina*. While we found no evidence of AMGs related to nitrogen or sulfur metabolism, we

instead report the presence of AMGs putatively linked to carbon utilization, particularly a diverse

array of CAZymes that may be involved in organic carbon cycling. These results suggest that

viruses may play an important role in carbon sequestration and blue carbon storage within

seagrass ecosystems, which may have significant implications for climate change mitigation.

Moving forward, research should prioritize investigating the role of phages through viromic

techniques, which would offer deeper insights into viral ecology and phage contributions to

carbon cycling in seagrass habitats.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

The raw metagenomic sequencing data was deposited at GenBank under accession no.

PRJNA1140276. MAGs with > 90% completion estimates were deposited on NCBI under this

BioProject at GenBank under accession no. PRJNA1140276, while MAGs with completion

estimates between 80 and 90% were deposited on Zenodo (DOI: 10.5281/zenodo.14225973).

Viral genomes were also deposited on Zenodo (DOI: 10.5281/zenodo.14226038). All code used

in this work has been deposited on Github (casett/Seagrass_MAG_and_Virus_Catalog) and

archived in Zenodo (DOI: 10.5281/zenodo.14226514).

Competing Interests

The authors declare no competing interests.

Funding

The sequencing data in this work was generated by grants from the UC Davis H. A. Lewin

Family Fellowship and the UC Davis Center for Population Biology to CLE. CLE was supported

by the National Science Foundation (NSF) under a NSF Ocean Sciences Postdoctoral

Fellowship (Award No. 2205744). JES is a CIFAR Fellow in the program Fungal Kingdom:

Threats and Opportunities and was partially supported by NSF awards EF-2125066 and IOS-

2134912. Computations were performed using the computer clusters and data storage

366   resources of the UC Riverside HPCC, which were funded by grants from NSF (MRI-2215705,

367   MRI-1429826) and NIH (1S10OD016290-01A1). The funders had no role in study design, data

368   collection and analysis, decision to publish, or preparation of the manuscript.

369   *Authors' Contributions*

370
371   CLE conceived and designed the experiments, performed sampling, analyzed the data,

372   prepared figures and/or tables, wrote and reviewed drafts of the paper. JES reviewed drafts of

373   the paper.

374   *Acknowledgements*

375
376   We would like to thank Jonathan A. Eisen for feedback on this manuscript. We would like to

377   thank Katherine Dynarski (ORCID: 0000-0001-5101-9666) and Sonia Ghose (ORCID: 0000-

378   0001-5667-6876) for their help with sample collection. We would also like to thank John J.

379   Stachowicz for use of his scientific sampling permit, California Department of Fish and Wildlife

380   Scientific Collecting Permit # SC 4874.

References

1. Hemminga MA, Duarte CM. Seagrass Ecology. Cambridge University Press; 2000. doi:10.1017/CBO9780511525551

2. Fourqurean JW, Duarte CM, Kennedy H, Marbà N, Holmer M, Mateo MA, et al. Seagrass ecosystems as a globally significant carbon stock. Nat Geosci. 2012;5: 505–509. doi:10.1038/ngeo1477

3. Orth RJ, Carruthers TJB, Dennison WC, Duarte CM, Fourqurean JW, Heck KL, et al. A Global Crisis for Seagrass Ecosystems. BioScience. 2006;56: 987. doi:10.1641/0006-3568(2006)56[987:agcfse]2.0.co;2

4. Mcleod E, Chmura GL, Bouillon S, Salm R, Björk M, Duarte CM, et al. A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO2. Front Ecol Environ. 2011;9: 552–560. doi:10.1890/110004

5. Trivedi P, Batista BD, Bazany KE, Singh BK. Plant–microbiome interactions under a changing world: responses, consequences and perspectives. New Phytol. 2022;234: 1951–1959. doi:10.1111/nph.18016

6. Vandenkoornhuyse P, Quaiser A, Duhamel M, Le Van A, Dufresne A. The importance of the microbiome of the plant holobiont. New Phytol. 2015;206: 1196–1206. doi:10.1111/nph.13312

7. Compant S, Samad A, Faist H, Sessitsch A. A review on the plant microbiome: Ecology, functions, and emerging trends in microbial application. J Advert Res. 2019;19: 29–37. doi:10.1016/j.jare.2019.03.004

8. Fahimipour AK, Kardish MR, Lang JM, Green JL, Eisen JA, Stachowicz JJ. Global-Scale Structure of the Eelgrass Microbiome. Appl Environ Microbiol. 2017;83. doi:10.1128/AEM.03391-16

9. Ettinger CL, Voerman SE, Lang JM, Stachowicz JJ, Eisen JA. Microbial communities in sediment from *Zostera marina* patches, but not the leaf or root microbiomes, vary in relation to distance from patch edge. PeerJ. 2017;5: e3246. doi:10.7717/peerj.3246

10. Cúcio C, Engelen AH, Costa R, Muyzer G. Rhizosphere Microbiomes of European Seagrasses Are Selected by the Plant, But Are Not Species Specific. Front Microbiol. 2016;7: 440. doi:10.3389/fmicb.2016.00440

11. Cúcio C, Overmars L, Engelen AH, Muyzer G. Metagenomic Analysis Shows the Presence of Bacteria Related to Free-Living Forms of Sulfur-Oxidizing Chemolithoautotrophic Symbionts in the Rhizosphere of the Seagrass *Zostera marina*. Front Mar Sci. 2018;5: 343793. doi:10.3389/fmars.2018.00171

12. Crump BC, Wojahn JM, Tomas F, Mueller RS. Metatranscriptomics and Amplicon Sequencing Reveal Mutualisms in Seagrass Microbiomes. Front Microbiol. 2018;9: 332425. doi:10.3389/fmicb.2018.00388

13. Hurtado-McCormick V, Kahlke T, Petrou K, Jeffries T, Ralph PJ, Seymour JR. Regional and Microenvironmental Scale Characterization of the Seagrass Microbiome. Front

Microbiol. 2019;10: 1011. doi:10.3389/fmicb.2019.01011

14. Ettinger CL, Williams SL, Abbott JM, Stachowicz JJ, Eisen JA. Microbiome succession
during ammonification in eelgrass bed sediments. PeerJ. 2017;5: e3674.
doi:10.7717/peerj.3674

15. Sun F, Zhang X, Zhang Q, Liu F, Zhang J, Gong J. Seagrass (*Zostera marina*) Colonization
Promotes the Accumulation of Diazotrophic Bacteria and Alters the Relative Abundances of
Specific Bacterial Lineages Involved in Benthic Carbon and Sulfur Cycling. Appl Environ
Microbiol. 2015;81. doi:10.1128/AEM.01382-15

16. Wang L, Tomas F, Mueller RS. Nutrient enrichment increases size of *Zostera marina*
shoots and enriches for sulfur and nitrogen cycling bacteria in root-associated microbiomes.
FEMS Microbiol Ecol. 2020;96. doi:10.1093/femsec/fiaa129

17. Mohr W, Lehnen N, Ahmerkamp S, Marchant HK, Graf JS, Tschitschko B, et al. Terrestrial-
type nitrogen-fixing symbiosis between seagrass and a marine bacterium. Nature.
2021;600: 105–109. doi:10.1038/s41586-021-04063-4

18. Kristensen DM, Mushegian AR, Dolja VV, Koonin EV. New dimensions of the virus world
discovered through metagenomics. Trends Microbiol. 2010;18.
doi:10.1016/j.tim.2009.11.003

19. Jin M, Guo X, Zhang R, Qu W, Gao B, Zeng R. Diversities and potential biogeochemical
impacts of mangrove soil viruses. Microbiome. 2019;7. doi:10.1186/s40168-019-0675-9

20. Hurwitz BL, Westveld AH, Brum JR, Sullivan MB. Modeling ecological drivers in marine
viral communities using comparative metagenomics and network analyses. Proc Natl Acad
Sci U S A. 2014;111. doi:10.1073/pnas.1319778111

21. Breitbart MYA, Thompson LR, Suttle CA, Sullivan MB. Exploring the vast diversity of
marine viruses. Oceanography. 2007;20: 135–139. Available:
https://www.jstor.org/stable/24860053

22. Ettinger CL, Eisen JA. Characterization of the Mycobiome of the Seagrass, *Zostera marina*,
Reveals Putative Associations With Marine Chytrids. Front Microbiol. 2019;10: 2476.
doi:10.3389/fmicb.2019.02476

23. Bushnell B. BBTools. 2022. Available: https://sourceforge.net/projects/bbmap/

24. Ma X, Olsen JL, Reusch TBH, Procaccini G, Kudrna D, Williams M, et al. Improved
chromosome-level genome assembly and annotation of the seagrass, *Zostera marina*
(eelgrass). F1000Res. 2021;10. doi:10.12688/f1000research.38156.1

25. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. Genome Biology. 2009;10: R25.
doi:10.1186/gb-2009-10-3-r25

26. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.
doi:10.1093/bioinformatics/btp352

27. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution

for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics.
2015;31: 1674–1676. doi:10.1093/bioinformatics/btv033

28. Eren AM, Murat Eren A, Esen ÖC, Quince C, Vineis JH, Morrison HG, et al. Anvi'o: an
advanced analysis and visualization platform for 'omics data. PeerJ. 2015;3: e1319.
doi:10.7717/peerj.1319

29. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic
gene recognition and translation initiation site identification. BMC Bioinformatics. 2010;11:
119. doi:10.1186/1471-2105-11-119

30. Lee MD. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics. 2019;35:
4162–4164. doi:10.1093/bioinformatics/btz188

31. Lee MD. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics. 2019;35:
4162–4164. doi:10.1093/bioinformatics/btz188

32. Delmont T. Assessing the completion of eukaryotic bins with anvi'o. 2018. Available:
http://merenlab.org/delmont-euk-scgs

33. Eddy SR. Accelerated Profile HMM Searches. PLoS Comput Biol. 2011;7: e1002195.
doi:10.1371/journal.pcbi.1002195

34. Seemann T. barrnap: BAsic Rapid Ribosomal RNA Predictor. Available:
https://github.com/tseemann/barrnap

35. Menzel P, Ng KL, Krogh A. Fast and sensitive taxonomic classification for metagenomics
with Kaiju. Nat Commun. 2016;7: 11257. doi:10.1038/ncomms11257

36. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, et al. Binning
metagenomic contigs by coverage and composition. Nat Methods. 2014;11: 1144–1146.
doi:10.1038/nmeth.3103

37. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive binning
algorithm for robust and efficient genome reconstruction from metagenome assemblies.
PeerJ. 2019;7: e7359. doi:10.7717/peerj.7359

38. Graham ED, Heidelberg JF, Tully BJ. BinSanity: unsupervised clustering of environmental
microbial assemblies using coverage and affinity propagation. PeerJ. 2017;5: e3035.
doi:10.7717/peerj.3035

39. Wu Y-W, Tang Y-H, Tringe SG, Simmons BA, Singer SW. MaxBin: an automated binning
method to recover individual genomes from metagenomes using an expectation-
maximization algorithm. Microbiome. 2014;2: 1–18. doi:10.1186/2049-2618-2-26

40. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, et al. Recovery of
genomes from metagenomes via a dereplication, aggregation, and scoring strategy. Nature
Microbiology. 2018;3: 836–843. doi:10.1101/107789

41. Nayfach S, Shi ZJ, Seshadri R, Pollard KS, Kyrpides NC. New insights from uncultivated
genomes of the global human gut microbiome. Nature. 2019;568: 505–510.
doi:10.1038/s41586-019-1058-x

42.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the
quality of microbial genomes recovered from isolates, single cells, and metagenomes.
Genome Res. 2015;25: 1043–1055. doi:10.1101/gr.186072.114

43.   Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. CheckM2: a rapid, scalable and
accurate tool for assessing microbial genome quality using machine learning. Nat Methods.
2023;20: 1203–1212. doi:10.1038/s41592-023-01940-w

44.   Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK, et al.
Minimum information about a single amplified genome (MISAG) and a metagenome-
assembled genome (MIMAG) of bacteria and archaea. Nat Biotechnol. 2017;35: 725–731.
doi:10.1038/nbt.3893

45.   Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. Bioinformatics. 2019.
doi:10.1093/bioinformatics/btz848

46.   Sogin EM, Michellod D, Gruber-Vodicka HR, Bourceau P, Geier B, Meier DV, et al. Sugars
dominate the seagrass rhizosphere. Nature Ecology & Evolution. 2022;6: 866–877.
doi:10.1038/s41559-022-01740-z

47.   Rubio-Portillo E, Martin-Cuadrado A-B, Ramos-Esplá AÁ, Antón J. Metagenomics Unveils
Posidonia oceanica "Banquettes" as a Potential Source of Novel Bioactive Compounds and
Carbohydrate Active Enzymes (CAZymes). mSystems. 2021.
doi:10.1128/msystems.00866-21

48.   Miranda K, Weigel BL, Fogarty EC, Veseli IA, Giblin AE, Murat Eren A, et al. The Diversity
and Functional Capacity of Microbes Associated with Coastal Macrophytes. mSystems.
2022. doi:10.1128/msystems.00592-22

49.   Schorn S, Ahmerkamp S, Bullock E, Weber M, Lott C, Liebeke M, et al. Diverse
methylotrophic methanogenic archaea cause high methane emissions from seagrass
meadows. Proceedings of the National Academy of Sciences. 2022;119: e2106628119.
doi:10.1073/pnas.2106628119

50.   Fraser MW, Martin BC, Wong HL, Burns BP, Kendrick GA. Sulfide intrusion in a habitat
forming seagrass can be predicted from relative abundance of sulfur cycling genes in
sediments. Sci Total Environ. 2023;864: 161144. doi:10.1016/j.scitotenv.2022.161144

51.   Fraser MW, Gleeson DB, Grierson PF, Laverock B, Kendrick GA. Metagenomic Evidence
of Microbial Community Responsiveness to Phosphorus and Salinity Gradients in Seagrass
Sediments. Front Microbiol. 2018;9: 378381. doi:10.3389/fmicb.2018.01703

52.   Guo J, Vik D, Pratama AA, Roux S, Sullivan M. Viral sequence identification SOP with
VirSorter2. protocols.io. 2021. Available: https://www.protocols.io/view/viral-sequence-
identification-sop-with-virsorter2-btv8nn9w

53.   Guo J, Bolduc B, Zayed AA, Varsani A, Dominguez-Huerta G, Delmont TO, et al.
VirSorter2: a multi-classifier, expert-guided approach to detect diverse DNA and RNA
viruses. Microbiome. 2021;9: 37. doi:10.1186/s40168-020-00990-y

54.   Nayfach S, Camargo AP, Schulz F, Eloe-Fadrosh E, Roux S, Kyrpides NC. CheckV

assesses the quality and completeness of metagenome-assembled viral genomes. Nat
Biotechnol. 2021;39: 578–585. doi:10.1038/s41587-020-00774-7

55. Shaffer M, Borton MA, McGivern BB, Zayed AA, La Rosa SL, Solden LM, et al. DRAM for
distilling microbial metabolism to automate the curation of microbiome function. Nucleic
Acids Res. 2020;48: 8883–8900. doi:10.1093/nar/gkaa621

56. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate genomic
comparisons that enables improved genome recovery from metagenomes through de-
replication. ISME J. 2017;11: 2864–2868. doi:10.1038/ismej.2017.126

57. Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al. Taxonomic
assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing
networks. Nat Biotechnol. 2019;37: 632–639. doi:10.1038/s41587-019-0100-8

58. Cook R, Brown N, Redgwell T, Rihtman B, Barnes M, Clokie M, et al. INfrastructure for a
PHAge REference Database: Identification of Large-Scale Biases in the Current Collection
of Cultured Phage Genomes. PHAGE. 2021;2: 214–223. doi:10.1089/phage.2021.0007

59. Camargo AP, Roux S, Schulz F, Babinski M, Xu Y, Hu B, et al. Identification of mobile
genetic elements with geNomad. Nat Biotechnol. 2024;42: 1303–1312.
doi:10.1038/s41587-023-01953-y

60. Roux S, Camargo AP, Coutinho FH, Dabdoub SM, Dutilh BE, Nayfach S, et al. iPHoP: An
integrated machine learning framework to maximize host prediction for metagenome-
derived viruses of archaea and bacteria. PLoS Biol. 2023;21: e3002083.
doi:10.1371/journal.pbio.3002083

61. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature methods.
2012;9. doi:10.1038/nmeth.1923

62. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 2010;26: 841–842. doi:10.1093/bioinformatics/btq033

63. Roux S, Adriaenssens EM, Dutilh BE, Koonin EV, Kropinski AM, Krupovic M, et al.
Minimum Information about an Uncultivated Virus Genome (MIUViG). Nat Biotechnol.
2018;37: 29–37. doi:10.1038/nbt.4306

64. Roux S, Emerson JB, Eloe-Fadrosh EA, Sullivan MB. Benchmarking viromics: an
evaluation of metagenome-enabled estimates of viral community composition and diversity.
PeerJ. 2017;5: e3817. doi:10.7717/peerj.3817

65. R Core Team. R: A Language and Environment for Statistical Computing. 2021. Available:
https://www.r-project.org/

66. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis
and Graphics of Microbiome Census Data. PLoS One. 2013;8: e61217.
doi:10.1371/journal.pone.0061217

67. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome to
the Tidyverse. Journal of Open Source Software. 2019;4: 1686. doi:10.21105/joss.01686

68. Lahti L, Shetty S. Tools for microbiome analysis in R - Microbiome package. 2017.

Available: http://microbiome.github.com/microbiome

69. Turner D, Kropinski AM, Adriaenssens EM. A Roadmap for Genome-Based Phage
Taxonomy. Viruses. 2021;13. doi:10.3390/v13030506

70. Santos-Medellin C, Zinke LA, Ter Horst AM, Gelardi DL, Parikh SJ, Emerson JB. Viromes
outperform total metagenomes in revealing the spatiotemporal patterns of agricultural soil
viral communities. ISME J. 2021;15: 1956–1970. doi:10.1038/s41396-021-00897-y

71. Ettinger CL, Saunders M, Selbmann L, Delgado-Baquerizo M, Donati C, Albanese D, et al.
Highly diverse and unknown viruses may enhance Antarctic endoliths' adaptability.
Microbiome. 2023;11: 103. doi:10.1186/s40168-023-01554-6

72. An L, Liu X, Wang J, Xu J, Chen X, Liu X, et al. Global diversity and ecological functions of
viruses inhabiting oil reservoirs. Nat Commun. 2024;15: 6789. doi:10.1038/s41467-024-
51101-6

Figure 1. Putative viral sequences identified from seagrass ecosystems. (A) The proviral status

of predicted viral genomes is shown as a bar graph, with the number above the bar representing

the total number of sequences. (B) CheckV quality metrics for viral sequences are shown as a

bar graph with bars colored by quality and the number above the bar representing the total

number of sequences in each bar. (C) Bar chart depicting the number of viral sequences

identified in the metagenomic co-assembly from each study, with the number to the right of the

bar representing the total number of sequences in each bar. (D) Principal-coordinate analysis

(PCoA) visualization of Hellinger distances of relative abundance of viral communities across

metagenomic samples. Samples are colored by study, and have shapes based on sample type.

Figure 2. Viral-Bacterial interactions likely related to carbon metabolism. (A) Bar chart depicting

the number of metagenome-assembled genomes (MAGs) identified in this study, colored by

taxonomic class, and with the number to the right of the bar representing the total number of

MAGs in each bar. (B) Bar chart depicting the viral host predictions, colored by host taxonomic

class, and with the number to the right of the bar representing the total number of viral

sequences in each bar. (C) Bar chart showing the number of predicted phage auxiliary

metabolic genes (AMGs) summarized by DRAM-v distilled metabolic categories, with the

number to the right of the bar representing the total number of AMGs in each bar. (D) Bar chart

showing the number of predicted phage AMGs in different CAZyme families with-in the carbon

utilization category, with the number to the right of the bar representing the total number of

AMGs in each bar.