Establishing reliable blood biomarkers for trimethylamine N-oxide status in rodents: effects of oral choline challenge, dietary choline and fasting conditions

Ahmad Ud Din[a], Michael G. Sweet[a], Ashley M. McAmis[a], Juanita G. Ratliff[1a], Anandh Babu Pon Velayutham[b], Andrew P. Neilson[a,c]*

[a]Plants for Human Health Institute, North Carolina State University, 600 Laureate Way, Kannapolis, NC, 28081, USA

[b]Department of Nutrition and Integrative Physiology, College of Health, University of Utah, 250 South 1850 East, Room 214, Salt Lake City, Utah 84112, USA

[c]Department of Food, Bioprocessing, and Nutrition Sciences, North Carolina State University, Campus Box 7624, Raleigh, NC, 27695-7624, USA

Ahmad Ud Din: adin@ncsu.edu
Michael G. Sweet1: mgsweet@ncsu.edu
Ashley M. McAmis1: ammcamis@ncsu.edu
Juanita Ratliff: jgratlif@ncsu.edu
Anandh Babu Pon Velayutham: anandh.velayutham@utah.edu
Andrew P. Neilson: aneilso@ncsu.edu

Corresponding author: 600 Laureate Way, Kannapolis, NC, 28081, USA. Phone: +1(704)250-5495; email: aneilso@ncsu.edu

Abstract

Circulating concentrations of the gut microbial-mammalian metabolite trimethylamine N-oxide (TMAO) is linked to atherosclerosis risk. TMAO biosynthesis begins when dietary choline is converted to trimethylamine (TMA) by gut microbial TMA lyase. TMA is transported to the liver, where flavin-containing monooxygenases convert it to TMAO. While dietary modifications regulate TMAO production, the impact of different intake methods, including oral gavage, dietary supplementation, and conditions such as fasting versus non-fasting, has not been fully explored. Twelve female Sprague-Dawley rats were divided into three diet groups (n = 4 per group): no-choline (0% choline), low-choline (0.08% choline), and high-choline (1% choline). Choline and TMAO fasting and non-fasting blood concentrations, and their kinetics following an acute choline challenge, were assessed before and after a 2-week dietary intervention with the distinct choline dietary levels. Fasting choline was under tight control, with little effect of dietary choline. Non-fasting choline was more variable, with high dietary choline reflected in higher blood choline. Greater levels of dietary choline were reflected in significantly greater levels of TMAO, particularly for non-fasting levels. Kinetic profiling demonstrated additional information regarding the appearance and clearance of these compounds from blood. These results suggest that acute oral choline gavage is likely most suitable for studies targeting acute (direct) inhibitors, whereas a choline-rich diet with assessment of fasting and non-fasting blood levels is more suitable for studying alterations to TMAO production capacity. Future research should examine the impact on atherosclerosis biomarkers and microbiome diversity to deepen the understanding of TMAO regulation and its cardiovascular implications.

Keywords: Cardiovascular disease, atherosclerosis, choline, TMAO, fasting, dietary intervention, oral gavage

1. Introduction

Cardiovascular disease (CVD) is the leading cause of death worldwide and in the US [1,2]. According to a recent estimate, global deaths from CVD are 19.8 million annually [3]. Multiple risk factors are associated with CVD, such as poor diet, smoking, obesity, lack of physical activity, etc. The composition and function of the gut microbiome are also associated with various cardiovascular events [4] through the production of diet derived microbial metabolites [5]. Gut microbes can produce both beneficial and harmful metabolites, such as short-chain fatty acids and trimethylamine N-oxide (TMAO) [5,6], respectively.

Trimethylamine *N*-oxide (TMAO) was identified as a novel risk factor for CVD, specifically atherosclerosis, in 2011 by the Hazen group at the Cleveland Clinic [7,8]. Since then, elevated blood TMAO has been reported as a CVD and mortality risk factor [9,10], although controversy remains regarding its utility as a predictor of CVD risk [11,12]. TMAO is produced by sequential action of gut microbiota and the liver [8]. First, gut bacteria possessing choline trimethylamine (TMA) lyase enzymes (*cutC/D* gene cluster) metabolize choline into TMA [8]. TMA is then absorbed into the circulation and oxidized to TMAO by hepatic flavin-containing monooxygenase 3 (FMO3) [13]. Other quaternary amines, such as carnitine, can also be metabolized to TMA by similar microbial enzymes.

Currently, there is no FDA-approved drug to manage elevated TMAO [14,15]. One strategy that has shown promise is direct inhibition of TMA lyases. For example, 3,3-dimethyl-1-butanol (DMB), a choline analog, inhibits TMA lyase and has shown a non-toxic potential to inhibit TMAO formation and prevent atherosclerosis *in vivo* [16,17]. However, due to the lack of approved pharmaceuticals to date, complementary non-drug strategies are needed. Dietary interventions are a logical strategy to reduce TMAO formation with the goal of preventing or slowing atherosclerosis/CVD [18,19]. Previous studies have demonstrated the cardioprotective roles of plant-based dietary interventions, including berries [18,20]. For example, blueberry supplementation mediates vascular inflammation, improves vascular dysfunction, and reduces circulating TMAO concentrations [21,22]. Polyphenols such as chlorogenic acid (CGA) are abundant in blueberries and have the potential to reduce choline microbial metabolism into TMA [23,24]. Chronic CGA supplementation improves the gut microbiome and protects against CVD in animal models [23,25]. It is currently unknown how CGA-rich foods may reduce TMA and TMAO, but it may act by acute mechanisms (such as direct TMA lyase inhibition) requiring the inhibitor and substrate to be present simultaneously, or mechanisms requiring chronic exposure, including shifts in microbiome taxonomic distribution, altered levels of *cutC/D*, and/or changes to FMO3

expression. The key step in lowering TMAO is reducing the conversion of choline to
TMA by gut bacteria [16]. We recently developed an *ex vivo-in vitro* methodology to
screen compounds for the ability to reduce microbial biotransformation of choline into
TMA [26,27]. Using our fecal fermentation method, we tested the ability of CGA to
inhibit TMA production [28]. We found CGA reduces both choline use and TMA
production [28].

In vivo studies are needed to validate the efficacy of compounds identified *in vitro* as
having potential TMA- and TMAO-lowering activities. There are various methods in
place to use choline administration models either in a chronic diet or acute oral
challenge (by gavage) [29,30]. However, there is a lack of consensus on the utility of
various potential TMAO measures such as blood vs. urinary levels, or single time point
vs. kinetics over time. There is no clearly accepted method to evaluate the efficacy of
interventions that may reduce choline conversion to TMA and TMAO. Given that such
interventions might work via acute or chronic intervention, experimental conditions
such as blood collection in a fasting or non-fasting state, or use of chronic dietary
choline vs. acute exposure , could yield vastly different results [31]. In the present
study, we assessed the plasma TMAO responses with different experimental conditions
such as acute oral choline gavage, dietary choline intervention, fasting, and non-fasting
conditions. We found a clear difference in circulating TMAO between treatments of
fasting, oral gavage, and dietary interventions in the preclinical model. This outcome
could be considered for future in vivo studies to evaluate the efficacy of therapeutic
compounds on circulating TMAO.

2. Materials and Methods

2.1 Reagents and chemicals. Choline chloride (C7017-5G), 3,3-dimethyl-1-butanol (183105-
10G), ethyl 2-bromoacetate (133973-100G), ammonia (AM180-500MLGL), TMA (243205-
100G), TMA-$^{13}C_3$-^{15}N (730092), TMAO (317594-1G), and TMAO-d_9 (DLM47791), were
purchased from Sigma-Aldrich/Millipore (St. Louis, MO, USA). Ammonium formate
(70221-25G-F), LC-MS grade acetonitrile (JT9829-3), and water were purchased from
VWR International (Suwanee, GA, USA).

2.2 Animals. All animal experiments and procedures were approved by The North
Carolina Research Campus (Kannapolis, NC) Institutional Animal Care and Use
Committee (IACUC) under approval #20-017. The design of the study is shown in **Fig.**
1. A total of 12 female SAS Sprague-Dawley rats (age 7-8 weeks, 150-200g) were
purchased from Envigo RMS (Indianapolis, IN, USA). On arrival, rats were kept in the
Center for Laboratory Animal Science animal facility at the North Carolina Research
Campus. Female rats are the best model of human FMO3 as they have higher FMO3

than males, allowing efficient conversion of TMA to TMAO **[32–35]**. Rats were caged in
pairs and were allowed to assimilate for a week, provided with a standard chow
(0.253% Choline; Teklad 8604, Envigo) diet, and access to tap water *ad libitum.*
138

139
140   *Figure 1. A summary of the experimental timeline and treatments.*
141

2.3 Initial choline challenge. Prior to the start of the study, all rats were assimilated for 1
week on the standard chow diet. Non-fasting blood was collected from the saphenous
vein in the fed state in the afternoon (5-7 hours prior to fasting) and then rats were
fasted overnight for 9-10 h (water provided). The next morning, rats were weighed, and
fasting blood was collected from the saphenous vein. Then each rat was orally gavage
with choline (200 mg/kg choline, provided as 268 mg/kg choline chloride. Gavage time
was noted, and blood was collected at 3, 6, 9, 12, 15 and 24 h after gavage (the fasting
blood was time 0 h). Post-gavage, the rats were immediately provided with the chow
diet. Immediately following gavage, each rat was observed for dyspnea and signs of
fluid in the chest cavity (lethargy, hunched posture, unresponsiveness, changes in
normal respiration, etc.) within 0-10 min post-gavage. Rats were observed for signs of
distress, such as rough coat appearance, etc, at the end of the day and in the morning of
the next day. No animal showed any such adverse signs and symptoms. Blood was
directly collected into 0.8 mL lithium Heparin Sep tubes (MiniCollect® Capillary Blood
Collection System USA) from saphenous vein using 4 mm lancets. Blood was
centrifuged at 3000×g for 10 min, plasma was briefly stored on ice, and then stored at
−80°C for further processing.
159

2.4 Diets. After the first choline challenge was completed (24 h), animals were
immediately randomized to three diets (*n*=4/diet): no choline (NC, AIN-93M mature
rodent diet, without added choline, 0% choline bitartrate, D13120601), low-choline (LC,
AIN-93M mature rodent diet with 0.08 % choline bitartrate, D10012M), or high-choline
(HC, AIN-93M mature rodent diet with 1% choline bitartrate, D230113903) (**Table 1**).
The HC diet was designed to stimulate chronic TMA/TMAO overproduction [16]
compared to the LC diet. All diets (NC, LC, HC) were purchased from Research Diets
(RDI, New Brunswick, NJ, USA). Food and water were provided *ab libitum* and
monitored regularly.

Table 1. Treatment groups and respective 2-week diet and acute choline challenge parameters			
Group ID	**No choline (NC)**	**Low choline (LC)**	**High choline (HC)**
2-week diet	0% choline bitartrate (RDI D131206010)	0.08% choline bitartrate (RDI D13120601)	1% choline bitartrate (RDI D230113903)
Acute oral gavage	200 mg/kg choline		

2.5 Final choline challenge. The same procedures for the first choline challenge (non-fasting blood collection in the afternoon, fasting, gavage, and blood kinetics) were reproduced after the 2-week dietary intervention, except that rats were returned to their respective diets after gavage.

2.6 Euthanasia. Animals were euthanized by CO_2 following current American Veterinary Medical Association guidelines in a closed vessel with CO_2 from a compressed gas cylinder. CO_2 flow was controlled to displace 30-70% of the cage volume/min. Animals were exposed to CO_2 until 2~3 minutes following apparent clinical death (cessation of respiration). Following euthanasia, the abdominal cavity was opened ventrally from the rectum to the neck, and bilateral pneumothorax was performed to ensure death.

2.7 Choline and TMAO extraction and analysis. Choline, and TMAO in plasma samples were quantified by stable-isotope dilution UHPLC-MS/MS as described previously for animal samples [21,36–38]. Plasma samples (25 μl) were plated manually into 96-well plates, followed by automated sample preparation using an OT-2 liquid handler (Opentrons, NY, USA). Acetonitrile (100 μl) and zinc sulfate (10 μL) (5% w/v in water) were added to each well and 20 μL of a mixture of choline-d_9 and TMAO-d_9 as internal standards (IS, 2.5 μM each). The 96-well plate was sealed with an Adhesive Plate Seal (Waters, Milford, MA, USA) and sonicated for five minutes at room temperature in a water bath. After sonication, samples were transferred to AcroprepAdv 0.2 μm WWPTFE 96-well filtering plates (Pall Life Sciences, Port Washington, NY) and spin filtered (4 ^{0}C, 3400 x g for 10 min) onto Waters Acquity UPLC 700 μL Round 96-well Sample Plates and stored at −80°C until analysis. Choline and TMAO were analyzed by UHPLC-ESI-MS/MS as described previously [28]. Separation was performed on a Waters Acquity UPLC system equipped with an ACQUITY BEH HILIC column (1.7 μm, 2.1 × 100 mm) and pre-column (1.7 μm, 2.1 × 5 mm). The mobile phases consisted of 15 mM ammonium formate (pH 3.5) in water (A) and acetonitrile (B), with an isocratic gradient (80% B) at 0.65 mL/min. Column and autosampler temperatures were 30 °C and 10 °C, respectively. Quantification utilized a Waters Acquity triple quadrupole (TQD) mass spectrometer, operating under positive-mode ESI, with source and capillary temperatures at 150°C and 400°C. The capillary voltage was set at +0.60 kV, and nitrogen flows for desolvation and cone gas were 800 and 20 L/h, respectively. Data

204 acquisition was performed using multi-reaction monitoring (MRM) in MS/MS mode.
205 MRM fragmentation conditions for analytes and IS compounds are reported in **Table 2**.
206

Table 2. Optimized multi-reaction monitoring conditions for detection of choline and TMAO, and their internal standards

Compound	MW	RT (min)	MS/MS transition	Cone Voltage (V)	Collision Energy (eV)
Choline	104.2	1.05	104.2→60.0	38	16
Choline-d$_9$	113.2	1.05	113.3→69.1	40	16
TMAO	75.11	1.25	76.16→58.91	40	10
TMAO-d$_9$	84.17	1.20	85.21→68.10	40	12
MW, molecular weight; RT, retention time, CV, cone voltage; CE, collision energy					

207
For sample quantification, LCMS-grade water was spiked with 24 different
concentrations of choline and TMAO standards to obtain external calibration curves.
Standard samples (25 μL) were then prepared identically to blood samples. Samples
were quantified by calculating the analyte/IS peak abundance ratios and then
interpolating the concentrations from the standard curves. Data acquisition was carried
out using Masslynx software (V4.1 version, Waters).
214
2.8 Data analysis and statistics. Kinetics (concentration vs. time) were plotted and area-
under-the-curve (AUC) values were calculated for choline and TMAO from acute
challenges (pre- and post-dietary intervention). Values for each analyte were compared
by 2-way repeated measures (RM) ANOVA (to test effects of time, treatment, and their
interactions) and appropriate *post hoc* tests between treatments. Fasting and non-fasting
blood levels and kinetic AUCs were compared pre- and post-intervention (within
treatments) to assess adaptation over time, and between treatments (post-intervention).
To assess the effects of non-fasting and fasting between different dietary conditions, i.e.,
before diet and after comparison on choline utilization and TMAO production, a two-
way ANOVA followed by an appropriate test for multiple comparison was used.
225
226 **3. Results**
227
3.1 Choline blood concentrations in fasting and non-fasting conditions. Blood
concentrations of TMAO are assessed in humans and rodents as biomarkers of CVD
risk. However, there is no consensus on whether fasting- or fed-state blood levels or
kinetics in response to an acute challenge should be measured, or which is most useful.
Therefore, we first assessed fed and fasted blood levels before and after a 2-week
dietary intervention with various levels of choline in the diet.
234
Plasma choline levels in a non-fasting state before the dietary intervention had no
significant difference between groups as expected due to all groups being assimilated

on the same standard lab chow rodent diet (**Fig 2A**, blue color bars). After the dietary
intervention, non-fasting choline levels were higher in the HC group, followed by LC
and NC (**Fig 2A**, red color bars). When comparing non-fasting choline within each diet
group before and after the intervention, no difference was observed for either NC or LC
groups before and after the diet, while a significant difference was observed in the HC
group before vs. after the diet (**Fig 2A**). It is interesting to note that the fasting choline
blood levels of the NC and LC groups were not significantly different after going from
the chow diet (0.253% choline) both the NC diet (0% choline) or LC diet (0.08% choline).
Taken together, these data suggest that large differences in dietary choline levels are
reflected in non-fasting blood samples for very high dietary choline. However, at lower
dietary levels, some differences are not reflected in non-fasting levels, suggesting
somewhat tight control (as the dietary choline in the LC and HC groups differed by
12.5-fold, but non-fasting blood choline was only 2-fold different).
250

Fasting plasma choline levels are shown in **Fig 2B**. No differences were observed
between the 3 diets before or after the intervention. There were also no significant
differences within the NC and LC diets between time points. For the HC diet, the
plasma choline levels significantly increased after the diet (however, the pre-diet fasting
levels were lowest for HC, albeit not significantly lower than NC or HC). This suggests
again that circulating choline levels are under tight control, and that fasting levels may
not be useful reflections of dietary choline.
258

3.2 TMAO blood concentrations in fasting and non-fasting conditions. Non-fasting
levels of TMAO before and after the dietary intervention are shown in **Fig. 2C**. Before
the dietary intervention, there were no differences between groups (blue bars), again as
expected due to all groups being assimilated on the same standard lab chow rodent
diet. However, after the diet, TMAO levels for the HC diet were significantly greater
than the LC or NC diets (red bars). Within each diet, only HC was significantly different
between pre- and post-diet time points (increasing by ~50x from pre-diet levels).
Interestingly, LC and NC levels decreased after the diet, albeit non-significantly
compared to baseline. This is because the lab chow diets fed during assimilation was
0.253% choline, which is higher than both NC (0%) and LC (0.08%) diets. These data
suggest that high (but not normal or moderate) levels of dietary choline massively
increase non-fasting blood levels. Furthermore, increases in TMAO were greatly
disproportionate to dietary choline increases: in the HC diet, non-fasting TMAO
increased~50x from pre-diet levels, despite the dietary choline levels only rising ~4x,
from 0.253% in the chow diet to 1% in the HC diet).
274

275   Fasting plasma TMAO levels are shown in **Fig. 2D**. Note the distinct scales between
276   **Figs. 2C** and **2D**. Fasting levels showed similar trends as non-fasting levels. Before the

diet intervention, there were no differences in TMAO between the 3 diets as anticipated (blue bars). After the diet intervention, the blood TMAO for the HC diet was significantly greater than the LC or NC diets (red bars). Within each diet, only HC was significantly different between pre- and post-diet time points (increasing by ~10x from pre-diet levels, despite only an ~ 4X increase in dietary choline, from 0.025% to 1%). Comparing non-fasting (**Fig. 2C**) with fasting (**Fig. 2D**) levels, the TMAO values were roughly the same, except for the HC levels post-diet, for which on-fasting levels were ~350 µM and fasting levels were ~60 µM. This suggests that high dietary choline significantly increases both non-fasting and fasting TMAO, but that non-fasting levels drop significantly when fasted. This suggests that non-fasting blood collection may be advantageous for observing the most significant differences between treatment groups, with the caveat that some increased variability might be observed due to differences in time since the last meal (although that was not observed in this case, when rodents eat more frequently).

*Figure 2. Non-fasting and fasting plasma concentrations of choline and TMAO. A) Non-fasting plasma choline before and after dietary intervention. B). Fasting choline before and after dietary intervention. C) Non-fasting TMAO before and after dietary intervention. D) Fasting TMAO before and after dietary intervention. All graphs are (n=4/group). Values are mean ± SEM. Significant differences within diet groups before and after dietary intervention are shown by * (ns= no significant difference (p>0.05), * p≤ 0.05, ** p ≤ 0.01, *** p ≤ 0.001, **** p≤ 0.0001); while the difference in various dietary groups before diet and after diet is shown by letters above the graph (lowercase letters for before diet and uppercase letters for after diet); bars sharing letters are not significantly different.*

3.3 Acute challenge choline kinetics. Beyond the information provided by fed or fasting levels, an acute choline challenge can be useful to assess choline and TMAO responses in blood to a single, defined choline dose (where metabolic capacity is the only variable, as substrate availability is held constant). The controlled dose and timing of a gavage, and the subsequent plasma response, may provide additional information beyond fed state levels (where the time since the last ingestion of TMAO precursors and their amounts is often unknown) or fasting state levels (which depend heavily on elimination efficiency). Thus, a choline challenge can provide information that may more accurately reflect primarily TMAO production capacity.

For the pre-dietary intervention challenge (**Fig. 3A**), choline levels in blood demonstrated that the acute challenge worked as hypothesized to moderately elevate circulating choline. All groups had roughly the same fasting choline (0 h), and increased at time point 3 h, indicating oral acute challenge choline appearing in circulation. All the groups peaked at the highest concentration at time point 3 h, after which choline concentrations declined slightly and steadily, with decline back to baseline levels by 24 h.

319
Figure 3. Choline plasma kinetics following acute choline challenge. A) Choline kinetics
from an acute choline challenge before dietary intervention. B) Choline kinetics from an acute
choline challenge after dietary intervention. Data represent the mean ± SEM from n=4
rats/group. The asterisk above at each time point represents significant difference compared to
NC (* p≤0.05, ** p ≤0.01, *** p≤0.001). C). Areas under the curve (AUCs) for choline, results
presented is a mean of time (h) × Concentration (μM) +SEM (n=4). Lowercase letters indicate
statistical differences between groups before the dietary intervention. Uppercase letters represent
statistical differences between the groups after dietary intervention. Bars sharing the same letters
are not significantly different.

329
In the acute choline challenge after the 2-week dietary intervention (**Fig. 3B**), baseline (0
h) levels were similar between the 3 groups, but the separation was very clear at time
point 3 h after receiving the oral gavage of choline. Concentration between the groups
and afterwards continues to separate based on various dietary interventions of either
NC, LC or HC. This indicates that even after fasting, the HC group was efficient in
maintaining the choline concentration to a considerable level. Also, it's notable that the
choline in NC and LC was very low, and there were slight differences even after the
oral gavage of choline both before diet and after diet (Fig **3A**, **3B**). A high choline diet
(HC; 1%) effectively alters choline uptake so that serum choline levels rise acutely when

given an oral gavage, whereas an NC or LC diet is less effective at increasing choline levels in response to acute administration (**Fig. 3B**).

The areas under the curve (AUCs) for plasma choline kinetics before and after the diet intervention were calculated (**Fig. 3C**). As expected, no differences were found in choline AUCs between the 3 groups before the dietary intervention. There was a clear difference observed between all 3 diet groups following the 2-week intervention. Within diets, no difference was detected for choline AUCs between the pre- and post-diet intervention challenges for the NC group. Both LC and HC groups showed statistically different AUCs between the pre- and post-diet intervention challenges. These data suggest that high dietary choline significantly increases choline plasma concentrations.

3.4 TMAO kinetics after acute choline challenge. Plasma TMAO levels were monitored following the acute choline challenges to track TMAO formation and presence in the blood. In the pre-diet intervention challenge, similar to the choline kinetics, the TMAO levels dramatically dropped to a base level in all the groups when the rats were fasted (**Fig. 2D, Fig. 4A**). After the choline gavage, TMAO levels started to increase slightly at the same rate after the 3 h time point in all the groups (**Fig. 4A**). Maximum levels (~200 µM) were achieved around 12 h and gradually declined back to near baseline levels around 24 h. Notably, the peak TMAO levels (12 h, **Fig. 4A**) significantly trailed peak choline levels (3 h, **Fig. 3A**), as TMAO formation requires choline to proceed to the lower gut for microbial conversion to TMA and then conversion to TMAO by FMO3 in the liver before appearing in circulation. Also seen for choline (**Fig. 3A**), there were no kinetic differences between groups in TMAO prior to the 2-week dietary intervention. For the post-dietary intervention challenge, there was a clear difference in TMAO kinetics between groups, particularly HC (**Fig. 4B**). Also, it was clear that in the after diet regime, the TMAO concentration was higher in the HC group even after the fasting (**Fig. 4B, 2D**), which indicated that 1% dietary choline dietary intervention could be a potential method for future studies, providing very high TMAO acute production which could then be subject to potential inhibitory treatments. Interestingly, the NC and LC groups did not have a major difference in TMAO kinetics after the diet intervention. For these 2 groups, TMAO increased gradually up to 12-15 h (~200 µM for NC at 15 h and ~150 µM for LC at 12 h). However, the HC group started higher and went up to around 350-400 µM at 6 h and remained high through 24 h. These data suggest that high levels of dietary choline significantly elevate fasting as well as postprandial TMAO levels, whereas lower levels (LC, 0.08%) do not.

Looking at AUCs of TMAO kinetics following the acute choline challenge (**Fig. 4C**) before the dietary intervention, there were no differences between groups (blue bars), as expected. After the dietary treatment intervention, NC and LC treatments had the same

379 AUCs (which did not differ from pre-intervention levels). The HC diet significantly
380 increased TMAO after the dietary phase, compared to the HC group pre-intervention
381 and compared to the other two groups post-intervention. This suggests that HC dietary
382 treatment can be used in future chronic and acute studies to assess the efficacy of
383 potential TMAO-reducing agents (**Fig. 4C**).

385
Figure 4. TMAO plasma kinetics following acute choline challenge. A) TMAO kinetics
from an acute choline challenge before dietary intervention. B) TMAO from an acute choline
challenge after dietary intervention. Data represent the mean ± SEM from n=4 rats. The asterisk
above at each time point represents significant differences compared to NC (* p≤ 0.05, ** p ≤
0.01, *** p ≤ 0.001). C). Area under the curve for TMAO, results presented as mean of time (h) ×
concentration (μM) +SEM (n=4). Lowercase letters indicate statistical differences between
groups before the dietary intervention. Uppercase letters represent statistical differences between
groups after dietary intervention. Bars sharing the same letters are not significantly different.

4. Discussion
In the past few decades, the prevalence of CVD and associated mortality have increased
significantly [39]. The ecology and associated metabolome of the gut microbiome has
been found to be associated with CVD progression and as a strong contributing factor.
Especially microbial metabolites such as TMAO and tryptophan-derived indoxyl sulfate

are known prognostic microbial metabolites that lead to various CVD events [40,41]. To
date, there is no approved treatment to reduce levels of TMAO to counteract its
vascular consequences, such as atherosclerosis, heart failure, and coronary heart events.
Researchers are looking for dietary-based approaches with lower or fewer side effects
than conventional therapeutic interventions. While reducing TMAO is the goal of many
interventions, there is no consensus on whether interventions targeting choline
conversion to TMAO can be more rigorously evaluated by acute oral challenges or
fasting/fed levels during chronic dietary studies. Further, to the best of our knowledge,
there are no data assessing the comparative utility of fasting and non-fasting conditions
in dietary and oral gavage in any in *vivo* interventions. Fasting and non-fasting TMAO
levels are commonly used both in human and animal studies. It is worth mentioning
that non-fasting represents a state where choline and TMAO are being consumed and
produced, respectively, and eliminated. Fasting, on the other hand, represents a state
that persists after choline intake, TMAO production, and TMAO elimination. These are
distinct biomarkers, and it is unclear which is more important as a risk factor for CVD.
In the present study, we evaluated (i) the plasma TMAO responses to acute oral choline
gavage compared to a chronic choline diet, (ii) compared plasma choline and TMAO
levels in fasting and non-fasting conditions, and (iii) investigated the impact of diets
with or without choline on circulating choline and TMAO. We found a clear difference
between treatments of fasting, oral gavage, and dietary interventions in the preclinical
model. This outcome could be considered for future therapeutic studies using either
oral gavage or dietary choline. It also provides a basis for using choline as a precursor
and helps establish a rationale for testing therapeutic interventions targeting TMAO *in*
vivo.
424
In our study, all groups had similar choline levels in non-fasting conditions before the
intervention with different levels of choline in the diets. However, after the dietary
intervention, a noticeable difference in choline levels was observed in the non-fasting
condition. As shown, there is a range based on the type of diet, with the NC group
having the lowest choline levels, followed by the LC group, and the HC group showing
the highest choline levels (**Fig. 2A**). Non-fasting choline values, as expected, thus reflect
differences in dietary choline levels. Previous studies also demonstrated that fasting
could be useful in excluding the inter individual variability and to accurately get a basal
level for any intervention in place [42]. In our study, we also measured choline in the
fasting condition (**Fig. 2B**). Our results show no difference in fasted choline among
dietary groups pre- or post-diet intervention, although fasted choline did increase
significantly for the HC diet after the 2-week diet (Fig. 2B). A previous study
demonstrated that various blood biomarkers can vary and require optimized fasting
conditions [43]. While we did not focus specifically on evaluating fasting times in this
study, we selected the minimal fasting period commonly used in rats to minimize stress

on the animals **[44–46]**. Our results show that fasted choline is somewhat under tight
control, and dietary differences are poorly reflected in fasting choline. Thus, non-fasting
choline appears to be the better biomarker, particularly where differences in choline are
suspected or intentionally administered.
444
Dietary choline levels affected plasma TMAO fasting and non-fasting levels much more
than plasma choline levels. The 2-week dietary intervention resulted in a massive
increase in non-fasting TMAO for HC, while NC and LC were not different (**Fig. 2C**).
This suggests that at some level, TMAO production is kept in check by TMAO
elimination, but this process is significantly overwhelmed at higher TMAO production
levels (a 4-fold increase in dietary choline resulted in a 50-fold increase in non-fasting
TMAO levels for the HC group). A similar trend was observed in fasting TMAO levels
after the dietary intervention (**Fig. 2D**), but fasting TMAO levels in the HC group were
reduced ~5x (from 450 to 70 µM) from non-fasting levels. These data suggest that both
non-fasting and fasting TMAO levels are useful biomarkers of dietary choline and
conversion to TMAO. However, fasting levels will show lower separation among
treatments. Non-fasting TMAO may be the most sensitive and discriminating treatment
differentiator.
458
While fasting and non-fasting markers are useful, the drawback of these methods is that
the choline intake amount and time since the last choline intake are not as finely
controlled. Additionally, these markers cannot be utilized to determine if a potential
TMAO-lowering treatment acts via acute vs. chronic mechanisms. By controlling the
timing and amount of choline administered and comparing post-administration
kinetics, a more refined biomarker is obtained. Additionally, acute choline challenges
administered simultaneously with a potential TMAO-lowering can identify acute
inhibitors (which require the inhibitor and substrate to be present at the same time)
without prior administration of the treatment of interest. Compounds that act via
chronic mechanisms (including shifts in microbiome taxonomic distribution, altered
levels of *cutC/D gene copy or CutC/D protein*, and/or changes to FMO3 expression)
require an adaptation period and also do not require the TMAO-lowering agent to be
present. Thus, with no prior treatment, chronic acting agents would theoretically not
inhibit acute TMAO production because they are not CutC/D inhibitors. For this reason,
we explored the choline and TMAO kinetics in blood following an acute choline
challenge.
475
In the acute choline challenge, our initial results from before the diet showed that
gavage-administered choline entered the bloodstream, reaching a certain level that was
maintained for a few hours. The concentration peaked at the 3-hour time point (**Fig.
3A**), aligning with findings from a related previous study conducted in mice, although

with different objectives than ours [47,48]. Dietary choline intervention clearly altered
the acute kinetics for choline (**Fig. 3B**). Specifically, the HC group sustained a
significantly higher choline concentration, as the diet contained added choline. In
contrast, choline levels in the LC and NC groups began to decline after the 3-hour point.
This demonstrated that a 1% (HC) dietary choline level is effective in maintaining
substantial choline levels in circulation.
486
TMAO kinetics proved highly informative. In the pre-diet challenge, a slight delay was
observed, with TMAO beginning to appear and gradually increasing between 3-6 h
(**Fig. 4A**). Interestingly, pre-diet TMAO levels in all groups gradually increased. After 9-
12 hours, TMAO leveled off and began to decline. This suggests that oral gavage
interventions offer a narrow but reliably steady time window. Specifically, the 6-12 h
period seems optimal for focused analysis, providing valuable insights into the
metabolic responses to oral choline intake. However, this limited timeframe presents
challenges for conducting therapeutic intervention studies due to its practical
constraints. In the after-diet kinetics of TMAO, a clear variation was observed across
treatment groups, particularly in the HC group. Unlike the LC and NC groups, which
did not appear to exhibit changes in TMAO kinetics after the 2-week intervention
compared to the pre-diet intervention, the HC group demonstrated a rapid and
significant increase in TMAO, with a steady TMAO level after 12 hours. This suggests
that dietary choline has a significant impact on TMAO levels, making it particularly
useful in studies focused on this metabolite. The difference in TMAO levels in the after-
diet condition can likely be attributed to changes in microbiome *cutC/D* copy number,
CutC/D protein expression, or gerater activity of *cutC/D*-bearing genera caused by the
dietary choline intake [49], and/or increased FMO3 expression.
505
The type of inhibition studied—whether lowering of fed/fasted levels versus blunting of
acute kinetics—depends on the mechanism or proposed action of the inhibitor or
TMAO-lowering agent. Interventions that target direct enzyme inhibition of CutC/D or
FMO3, for instance, require both the substrate (choline or TMA, respectively) and the
prospective inhibitor to be present simultaneously in the gut or liver. These
mechanisms are best studied using acute challenges, where the inhibitor is co-
administered with choline via gavage. In contrast, treatments that reduce TMAO
through changes in microbial composition (e.g., decreasing *cutC/D*-bearing bacteria),
microbial function (e.g., reduced CutC/D expression), or hepatic function (e.g., reduced
FMO3 expression) are better suited to chronic treatments. These may involve
monitoring both fed and fasted states and conducting acute challenges to fully assess
the intervention's effects. The present data suggest that fasting and fed levels are useful
for assessing overall TMAO exposure, whereas acute challenges may be useful for
parsing out mechanisms of action.

This study is not without limitations. First, we focused on choline and did not include an analysis of carnitine, betaine, or other related precursor molecules of trimethylamine N-oxide (TMAO). The findings and conclusions drawn from this research, therefore, are only applicable to choline. Second, TMA acts as an intermediate metabolite that undergoes conversion into TMAO in the liver before it enters general circulation. Consequently, our study did not include direct measurement of TMA levels due to the inherent challenges involved in such an assessment. The difficulty arises from the rapid and extensive conversion of TMA to TMAO in the liver, making it impractical to accurately capture TMA concentrations. Third, we allowed rats to re-feed on their respective experimental diets immediately following the second challenge instead of reverting to a standard chow diet (which was done for the first acute challenge). This methodological choice may have contributed to differences in the observed kinetics and area under the curve (AUC) measurements. These variations could potentially be attributed to the consumption of the diets with differing nutrient levels post-gavage, rather than to differences in the absorption of TMA or the production of TMAO. Finally, while it would be preferable to use both male and female subjects to ensure a comprehensive analysis, this is complicated by the fact that males are not ideal models for TMAO research due to their low expression of flavin-containing monooxygenase 3 (FMO3), as supported by prior studies[32]. The inclusion of male subjects would require the use of a larger number of animals to compensate for the variability and potentially unreliable data, which may not be a practical or efficient approach **[32,33,50–52]**. Using twice as many animals for questionable data is not practical. Finally, use of unlabeled choline reduced the specificity of the blood levels of choline and TMAO, as choline from other sources (phospholipids, etc.) could be converted into free choline and TMAO. Use of labeled choline (such as choline-d_9) in the diet and measurement of the labeld substrate and products (such as choline-d_9 and TMAO-d_9) would increase the specificity and sensitivity of these blood biomarkers in animals.

It would be ideal to include specific atherosclerosis-related parameters and microbiome diversity analysis in future studies. This approach could provide valuable insights into how the microbiome shifts in response to varying dietary concentrations of specific metabolites. Including both fasting and non-fasting periods in dietary and oral gavage regimens is crucial. This dual approach could help clarify the reasons behind differing TMAO concentrations observed under various conditions. Understanding these variations could reveal how metabolic responses are influenced by different dietary strategies. Such analyses may shed light on the relationship between choline intake, microbiome changes, and TMAO levels. Overall, this could enhance our comprehension of dietary impacts on atherosclerosis risk and metabolic health.

5. Conclusions

The outcomes of this study highlighted the importance of differentiating between fasting and non-fasting levels when studying TMAO. The inclusion of a fasting period establishes a baseline that enables a more accurate comparison of subsequent results. However, the non-fasting samples may indicate greater differences between treatments and be more representative of typical blood profiles. Moreover, our findings indicated that administering choline via oral gavage can be used to assess TMAO kinetics. Using dietary choline—particularly at high concentration such as 1% choline—may be the most suitable approach for studies aiming to explore therapeutic strategies involving TMAO. The HC diet appears to provide a consistent and reliable means of raising TMAO levels, which can be beneficial for long-term research into metabolic and cardiovascular outcomes associated with TMAO. However, if the primary research focus involves the study of acute inhibitors of TMAO production or metabolism, the oral gavage method presents a distinct advantage. This approach facilitates the immediate ingestion of choline, followed by rapid absorption and metabolism into TMA and subsequently into TMAO. Such a setup allows researchers to monitor the immediate effects of choline and any interventions on TMAO synthesis, making it an ideal procedure for acute-phase studies where time-sensitive responses are critical.

Abbreviations: 3,3-dimethyl-1-butanol (DMB); area-under-the-curve (AUC); cardiovascular disease (CVD); chlorogenic acid (CGA); flavin-containing monooxygenase 3 (FMO3); high-choline (HC); low-choline (LC); no choline (NC); repeated measures (RM); trimethylamine (TMA); trimethylamine N-oxide (TMAO).

Author Contributions: Conceptualization: APN, AD; methodology: APN, AD; formal analysis, AD; investigation, AD, AMM, MGS, JGR; resources: APN; data curation, AD; writing—original draft preparation, AD; writing—review and editing, APN, AMM, JGR, MGS, ABPV; visualization, AD; supervision, APN; project administration, APN; funding acquisition, APN, ABPV. All authors have read and agreed to the published version of the manuscript.

Funding: Funding for this work was provided via USDA NIFA AFRI Foundational program award 2024-67017-42462. The sponsor had no role in study design, collection, analysis and interpretation of data, writing of the report, decision to submit the article for publication, or selection of the journal.

Institutional Review Board Statement: All animal experiments and procedures were approved by The North Carolina Research Campus Institutional Animal Care and Use Committee (IACUC) under approval #20-017.

Acknowledgments: The authors wish to acknowledge the contributions of Dr. Lisard Iglesias-Carres, who contributed to the original ideation of the study.

Conflicts of Interest: The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results".

References

[1] Ahmad FB, Anderson RN. The Leading Causes of Death in the US for 2020. JAMA 2021;325:1829–30. https://doi.org/10.1001/jama.2021.5469.

[2] Ms S, At H, A B de G, Nd F. Leading Causes of Death in the US During the COVID-19 Pandemic, March 2020 to October 2021. JAMA Internal Medicine 2022;182. https://doi.org/10.1001/jamainternmed.2022.2476.

[3] Mensah GA, Fuster V, Roth GA. A Heart-Healthy and Stroke-Free World. Journal of the American College of Cardiology 2023;82:2343–9. https://doi.org/10.1016/j.jacc.2023.11.003.

[4] Vijay A, Valdes AM. Role of the gut microbiome in chronic diseases: a narrative review. Eur J Clin Nutr 2022;76:489–501. https://doi.org/10.1038/s41430-021-00991-6.

[5] Witkowski M, Weeks TL, Hazen SL. Gut Microbiota and Cardiovascular Disease. Circulation Research 2020;127:553–70. https://doi.org/10.1161/CIRCRESAHA.120.316242.

[6] Spivak I, Fluhr L, Elinav E. Local and systemic effects of microbiome-derived metabolites. EMBO Reports 2022;23:e55664. https://doi.org/10.15252/embr.202255664.

[7] Tang WW, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, et al. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. New England Journal of Medicine 2013;368:1575–84.

[8] Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, DuGar B, et al. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. Nature 2011;472:57–63.

[9] Roncal C, Martínez-Aguilar E, Orbe J, Ravassa S, Fernandez-Montero A, Saenz-Pipaon G, et al. Trimethylamine-N-Oxide (TMAO) Predicts Cardiovascular Mortality in Peripheral Artery Disease. Sci Rep 2019;9:15580. https://doi.org/10.1038/s41598-019-52082-z.

[10] Tang WHW, Li XS, Wu Y, Wang Z, Khaw K-T, Wareham NJ, et al. Plasma trimethylamine N-oxide (TMAO) levels predict future risk of coronary artery disease in apparently healthy individuals in the EPIC-Norfolk prospective population study. American Heart Journal 2021;236:80–6. https://doi.org/10.1016/j.ahj.2021.01.020.

[11] Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. Circulation 2005;112:2735–52. https://doi.org/10.1161/CIRCULATIONAHA.105.169404.

[12] Nutrients | Free Full-Text | Trimethylamine-N-oxide (TMAO) as Novel Potential Biomarker of Early Predictors of Metabolic Syndrome n.d. https://www.mdpi.com/2072-6643/10/12/1971 (accessed August 20, 2024).

[13] Fennema D, Phillips IR, Shephard EA. Trimethylamine and Trimethylamine N-Oxide, a Flavin-Containing Monooxygenase 3 (FMO3)-Mediated Host-Microbiome Metabolic Axis Implicated in Health and Disease. Drug Metab Dispos 2016;44:1839–50. https://doi.org/10.1124/dmd.116.070615.

[14] Steinke I, Ghanei N, Govindarajulu M, Yoo S, Zhong J, Amin RH. Drug Discovery and Development of Novel Therapeutics for Inhibiting TMAO in Models of Atherosclerosis and Diabetes. Frontiers in Physiology 2020;11.

[15] Dannenberg L, Zikeli D, Benkhoff M, Ahlbrecht S, Kelm M, Levkau B, et al. Targeting the human microbiome and its metabolite TMAO in cardiovascular prevention and therapy. Pharmacology & Therapeutics 2020;213:107584. https://doi.org/10.1016/j.pharmthera.2020.107584.

[16] Wang Z, Roberts AB, Buffa JA, Levison BS, Zhu W, Org E, et al. Non-lethal Inhibition of Gut Microbial Trimethylamine Production for the Treatment of Atherosclerosis. Cell 2015;163:1585–95. https://doi.org/10.1016/j.cell.2015.11.055.

[17] Chen K, Zheng X, Feng M, Li D, Zhang H. Gut microbiota-dependent metabolite Trimethylamine N-oxide contributes to cardiac dysfunction in western diet-induced obese mice. Frontiers in Physiology 2017;8:1–9. https://doi.org/10.3389/fphys.2017.00139.

[18] Iglesias-Carres L, Hughes MD, Steele CN, Ponder MA, Davy KP, Neilson AP. Use of dietary phytochemicals for inhibition of trimethylamine N-oxide formation. Journal of Nutritional Biochemistry 2021;91:108600. https://doi.org/10.1016/j.jnutbio.2021.108600.

[19] Sikand G, Severson T. Top 10 dietary strategies for atherosclerotic cardiovascular risk reduction. American Journal of Preventive Cardiology 2020;4:100106. https://doi.org/10.1016/j.ajpc.2020.100106.

[20] Cosme P, Rodríguez AB, Espino J, Garrido M. Plant Phenolics: Bioavailability as a Key Determinant of Their Potential Health-Promoting Applications. Antioxidants 2020;9:1263. https://doi.org/10.3390/antiox9121263.

[21] Satheesh Babu AK, Petersen C, Iglesias-Carres L, Paz HA, Wankhade UD, Neilson AP, et al. Blueberry intervention mitigates detrimental microbial metabolite trimethylamine N-oxide by modulating gut microbes. BioFactors n.d.;n/a. https://doi.org/10.1002/biof.2014.

[22] Petersen C, Bharat D, Cutler BR, Gholami S, Denetso C, Mueller JE, et al. Circulating metabolites of strawberry mediate reductions in vascular inflammation and endothelial dysfunction in db/db mice. International Journal of Cardiology 2018;263:111–7. https://doi.org/10.1016/j.ijcard.2018.04.040.

[23] Chen J, Yu B, Chen D, Zheng P, Luo Y, Huang Z, et al. Changes of porcine gut microbiota in response to dietary chlorogenic acid supplementation. Applied Microbiology and Biotechnology 2019;103:8157–68. https://doi.org/10.1007/s00253-019-10025-8.

[24] Nguyen V, Taine EG, Meng D, Cui T, Tan W. Chlorogenic Acid: A Systematic Review on the Biological Functions, Mechanistic Actions, and Therapeutic Potentials. Nutrients 2024;16:924. https://doi.org/10.3390/nu16070924.

[25] Wang Z, Lam KL, Hu J, Ge S, Zhou A, Zheng B, et al. Chlorogenic acid alleviates obesity and modulates gut microbiota in high-fat-fed mice. Food Science and Nutrition 2019;7:579–88. https://doi.org/10.1002/fsn3.868.

[26] Fu H, Kong B, Zhu J, Huang H, Shuai W. Phenylacetylglutamine increases the susceptibility of ventricular arrhythmias in heart failure mice by exacerbated activation of the TLR4/AKT/mTOR signaling pathway. International Immunopharmacology 2023;116:109795. https://doi.org/10.1016/j.intimp.2023.109795.

[27] Iglesias-Carres L, Essenmacher LA, Racine KC, Neilson AP. Development of a High-Throughput Method to Study the Inhibitory Effect of Phytochemicals on Trimethylamine Formation. Nutrients 2021;13:1466. https://doi.org/10.3390/nu13051466.

[28] Iglesias-Carres L, Essenmacher LA, Racine KC, Neilson AP. Development of a high-throughput method to study the inhibitory effect of phytochemicals on trimethylamine formation. Nutrients 2021;13:1466.

[29] Mehta AK, Gaur SN, Arora N, Singh BP. Effect of choline chloride in allergen-induced mouse model of airway inflammation. European Respiratory Journal 2007;30:662–71. https://doi.org/10.1183/09031936.00019307.

[30] Xu F, Thomas JD, Goldowitz D, Hamre KM. The ameliorative effects of choline on ethanol-induced cell death in the neural tube of susceptible BXD strains of mice. Front Neurosci 2023;17. https://doi.org/10.3389/fnins.2023.1203597.

[31] Ge P-X, Tai T, Jiang L-P, Ji J-Z, Mi Q-Y, Zhu T, et al. Choline and trimethylamine N-oxide impair metabolic activation of and platelet response to clopidogrel through activation of the NOX/ROS/Nrf2/CES1 pathway. Journal of Thrombosis and Haemostasis 2023;21:117–32. https://doi.org/10.1016/j.jtha.2022.10.010.

[32] Ripp SL, Itagaki K, Philpot RM, Elfarra AA. Species and sex differences in expression of flavin-containing monooxygenase form 3 in liver and kidney microsomes. Drug Metabolism and Disposition 1999;27:46–52.

[33] Janmohamed A, Hernandez D, Phillips IR, Shephard EA. Cell-, tissue-, sex- and developmental stage-specific expression of mouse flavin-containing monooxygenases (Fmos). Biochemical Pharmacology 2004;68:73–83. https://doi.org/10.1016/j.bcp.2004.02.036.

[34] Falls JG, Ryu DY, Cao Y, Levi PE, Hodgson E. Regulation of mouse liver flavin-containing monooxygenases 1 and 3 by sex steroids. Archives of Biochemistry and Biophysics 1997;342:212–23. https://doi.org/10.1006/abbi.1997.9965.

[35] Falls JG, Blake BL, Cao Y, Levi PE, Hodgson E. Gender differences in hepatic expression of flavin-containing monooxygenase isoforms (FMO1, FMO3, and FMO5) in Mice. Journal of Biochemical Toxicology 1995;10:171–7. https://doi.org/10.1002/jbt.2570100308.

[36] Satheesh Babu AK, Petersen C, Paz HA, Iglesias-Carres L, Li Y, Zhong Y, et al. Gut Microbiota Depletion Using Antibiotics to Investigate Diet-Derived Microbial Metabolites: An Efficient Strategy. Molecular Nutrition & Food Research n.d.;n/a:2300386. https://doi.org/10.1002/mnfr.202300386.

[37] Brunt VE, Gioscia-Ryan RA, Casso AG, VanDongen NS, Ziemba BP, Sapinsley ZJ, et al. Trimethylamine-N-oxide promotes age-related vascular oxidative stress and endothelial dysfunction in mice and healthy humans. Hypertension 2020;76:101–12.

[38] Casso AG, VanDongen NS, Gioscia-Ryan RA, Clayton ZS, Greenberg NT, Ziemba BP, et al. Initiation of 3,3-dimethyl-1-butanol at midlife prevents endothelial dysfunction and attenuates in vivo aortic stiffening with ageing in mice. The Journal of Physiology 2022;600:4633–51. https://doi.org/10.1113/JP283581.

[39] Hughes BB, Kuhn R, Peterson CM, Rothman DS, Solórzano JR, Mathers CD, et al. Projections of global health outcomes from 2005 to 2060 using the International Futures integrated forecasting model. Bull World Health Organ 2011;89:478–86. https://doi.org/10.2471/BLT.10.083766.

[40] Hou K, Wu Z-X, Chen X-Y, Wang J-Q, Zhang D, Xiao C, et al. Microbiota in health and diseases. Sig Transduct Target Ther 2022;7:1–28. https://doi.org/10.1038/s41392-022-00974-4.

[41] Luqman A, Hassan A, Ullah M, Naseem S, Ullah M, Zhang L, et al. Role of the intestinal microbiome and its therapeutic intervention in cardiovascular disorder. Front Immunol 2024;15:1321395. https://doi.org/10.3389/fimmu.2024.1321395.

[42] Rubio WB, Cortopassi MD, Ramachandran D, Walker SJ, Balough EM, Wang J, et al. Not so fast: Paradoxically increased variability in the glucose tolerance test due to food withdrawal in continuous glucose-monitored mice. Molecular Metabolism 2023;77:101795. https://doi.org/10.1016/j.molmet.2023.101795.

[43] Carper D, Coué M, Laurens C, Langin D, Moro C. Reappraisal of the optimal fasting time for insulin tolerance tests in mice. Mol Metab 2020;42:101058. https://doi.org/10.1016/j.molmet.2020.101058.

[44] National Research Council (US) Committee for the Update of the Guide for theCare and Use of Laboratory Animals. Guide for the Care and Use of Laboratory Animals. 8th ed. Washington (DC): National Academies Press (US); 2011.

[45] Rowland NE. Food or fluid restriction in common laboratory animals: balancing welfare considerations with scientific inquiry. Comp Med 2007;57:149–60.

[46] Nowland MH, Hugunin KM, Rogers KL. Effects of Short-Term Fasting in Male Sprague–Dawley Rats. Comparative Medicine 2011;61:138.

[47] Gao X, Jiang C, Xu J, Yanagita T, Xue C, Wang Y. Serum pharmacokinetics of choline, trimethylamine, and trimethylamine-N-oxide after oral gavage of phosphatidylcholines with different fatty acid compositions in mice. Bioscience, Biotechnology, and Biochemistry 2016;80:2217–23. https://doi.org/10.1080/09168451.2016.1206813.

[48] Yu J, Zhang T, Gao X, Xue C, Xu J, Wang Y. Fish oil affects the metabolic process of trimethylamine N-oxide precursor through trimethylamine production and flavin-containing monooxygenase activity in male C57BL/6 mice. RSC Adv 2017;7:56655–61. https://doi.org/10.1039/C7RA10248H.

[49] Koeth RA, Wang Z, Levison BS, Buffa JA, Org E, Sheehy BT, et al. Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. Nat Med 2013;19:576–85. https://doi.org/10.1038/nm.3145.

[50] Lattard V, Lachuer J, Buronfosse T, Garnier F, Benoit E. Physiological factors affecting the expression of FMO1 and FMO3 in the rat liver and kidney. Biochemical Pharmacology 2002;63:1453–64.

[51] Cherrington NJ, Cao Y, Cherrington JW, Rose RL, HODGSON* E. Physiological factors affecting protein expression of flavin-containing monooxygenases 1, 3 and 5. Xenobiotica 1998;28:673–82.

[52] Novick RM, Mitzey AM, Brownfield MS, Elfarra AA. Differential localization of flavin-containing monooxygenase (FMO) isoforms 1, 3, and 4 in rat liver and kidney and evidence for expression of FMO4 in mouse, rat, and human liver and kidney microsomes. Journal of Pharmacology and Experimental Therapeutics 2009;329:1148–55.