Title: Small Extracellular Vesicles Isolated from Cardiac Tissue Matrix or Plasma Display

Distinct Aging-Related Changes in Cargo Contributing to Chronic Cardiovascular Disease

Authors: George Ronan[1,2], Jun Yang[1,3], Pinar Zorlutuna[1,2,3,4]*

[1]Bioengineering Graduate Program, University of Notre Dame, Notre Dame, IN, 46556, USA

[2]Department of Aerospace and Mechanical Engineering, University of Notre Dame, Notre Dame,

IN, 46556, USA

[3]Department of Chemical and Biomolecular Engineering, University of Notre Dame, Notre Dame,

IN, 46556, USA

[4]Harper Cancer Research Institute, University of Notre Dame, Notre Dame, 46556, USA

*Corresponding author. Email: pzorlutu@nd.edu

Abstract: Aging is a major risk factor for cardiovascular disease, the leading cause of death

worldwide, and numerous other diseases, but the mechanisms of these aging-related effects remain

elusive. Chronic changes in the microenvironment and paracrine signaling behaviors have been

implicated, but remain understudied. Here, for the first time, we directly compare extracellular

vesicles obtained from young and aged patients to identify therapeutic or disease-associated

agents, and directly compare vesicles isolated from heart tissue matrix (TEVs) or plasma (PEVs).

While young EVs showed notable overlap of miRNA cargo, aged EVs differed substantially,

indicating differential age-related changes between TEVs and PEVs. TEVs overall were uniquely

enriched in miRNAs which directly or indirectly demonstrate cardioprotective effects, with 45

potential therapeutic agents implicated in our analysis. Both populations also showed increased

predisposition to disease with aging, though through different mechanisms. PEVs were largely

correlated with chronic systemic inflammation, while TEVs were more related to cardiac

homeostasis and local inflammation. From this, 17 protein targets unique to TEVs were implicated

as aging-related changes which likely contribute to the development of cardiovascular disease.

Main Text:

INTRODUCTION

Advanced age is quickly becoming the most prevalent risk factor worldwide for numerous

diseases, including cardiovascular disease (CVD), leading cause of death worldwide[1], obesity and

obesity-related diseases[2], and many cancers[3], and by 2050 the global population aged 65 or older

is expected to double to over 2 billion or nearly 20% of predicted global population[4]. In some

rapidly aging countries, such as Japan and Italy, 65+ aged individuals already account for over

20% of the population[5], with Japan quickly approaching 30%. Recent literature has implicated

long-term changes in the cardiac microenvironment in aging-related CVD pathogenesis[6–9], and

suggests that these changes can be manipulated to improve patient outcomes. The use of

extracellular matrix (ECM) or ECM-derived materials has seen some success in pre-clinical trials

by leveraging previously established effects of ECM treatment, including

immunomodulation[7,10,11], stem cell recruitment[7,12,13], and functional revitalization[6,12–14]. These

properties enhance the reparative effects of novel biomaterials, commonly being integrated with

hydrogels or other bioscaffolds in treatment of CVDs[9,10,13,15,16]. However, the mechanisms by

which ECM promotes these effects is not well understood, and the drivers of aging-related changes

in the microenvironment are critically understudied.

Extracellular vesicles (EVs) have been recently identified as potential agents of aging-related

changes in the microenvironment[6,7,14] and key mediators of many therapeutic effects observed in

ECM treatment[11,13,17]. However, many of these studies have been largely limited to EVs isolated

from plasma and have been identified as exosomes[18,19]. Exosomes are a specific subcategory of

small EV (sEV), or EVs with typical sizes of under 200 nm[20,21], that express characteristic surface

protein markers and serve as common vehicles for endocrine, paracrine, and even autocrine

transport of RNAs and proteins[21–23]. While more recent efforts have focused on the broader

category of sEVs, due to difficulties in separating exosomes from other sEV subgroups such as

supermeres and exomeres[7,24], exosomes have been investigated as "tissue-free" theragnostic

nanoparticles. Stem cell-derived exosomes are being investigated in a clinical trial as vital

components for an "off-the-shelf" cardiac patch that emulates the benefits of direct ECM

incorporation[13,25], and plasma exosomes have provided a wealth of real-time diagnostic

information for patients in both CVDs[15,26] and cancer diagnosis[27,28].

Tissue matrix-derived sEVs provide a unique avenue to evaluate the effects of aging-related

changes in the microenvironment on the progression of CVDs, a topic that has remained elusive

and difficult to address clinically[8,10,13,14,17,29]. While CVD-related mortality is often actuated by the

onset of myocardial infarction (MI)[1,30], the risk of mortality is predominantly regulated by a

combination of risk factors including previous cardiac event[30–32], obesity and metabolic health[30,33],

and age[34,35], and recent studies have clearly demonstrated correlative effects between age and the

risk of MI or chronic CVD[4,8,35–37].

Previously, we discovered and characterized cardiac tissue ECM-resident exosome-like sEVs from

donor hearts from young or aged donors, and found that the size, morphology, and miRNA and

protein contents of these sEVs significantly changed with donor age[7]. Despite a growing body of

literature supporting the use of sEVs as dynamic theragnostic tools in CVD and other diseases[9,17],

currently in literature there is no single resource that directly investigates and establishes aging-

related changes in sEV content for tissue-resident sEVs, and any head-to-head comparison of

miRNA and whole protein profiles of tissue-resident or plasma-derived sEVs is similarly absent.

This may be due to the relatively recent discovery of ECM-resident sEVs[38], but remains a notable

gap in knowledge.

Herein we, for the first time in literature, characterize differences in size profile, surface protein distribution, and uptake mechanics of sEVs derived from cardiac tissue ECM, or tissue sEVs (TEVs), and sEVs derived from plasma, or plasma sEVs (PEVs). We also establish previously unreported differential aging-related changes in these metrics between TEVs and PEVs by assessing both populations of EVs from young (<40 y.o., n = 6 for tissue & plasma) and aged (>50 y.o., n = 6 for tissue & plasma) donors. After this, we provide a whole miRNA profile for TEVs and PEVs from young and aged donors, a dataset currently absent in literature, and identify targets and pathways of interest for CVD within our dataset. Following this, we do the same for a whole protein profile, again from TEVs and PEVs from both young and aged donors for the first time in literature. From these analyses, we can successfully bifurcate the cargo profiles of TEVs and PEVs, with PEVs tending to be largely correlated with systemic immune response and cardiac TEVs tending to be related to cardiac homeostasis and CVD, as expected. TEVs were also uniquely enriched in cardioprotective miRNAs involved in the regulation and prevention of fibrosis and anoxia-related cell death. Most interestingly, aged-source EVs in both populations tended to exhibit higher enrichment of cargo involved in disease-associated pathways than young-source EVs not through nucleic acid cargo, but through chauffeured proteins. While the specific pathways involved were different, with aged PEVs mostly trafficking proteins related to increases in chronic systemic inflammation and neurological diseases and aged TEVs transporting proteins relating to increased risk of heart failure, increased local inflammation, and overall metabolic shift, both populations suggested that EVs can propagate aging-related damage-inducing proteins. From these cargo analyses, we identified 45 miRNA targets which provide cardioprotective effects and 17 proteins which are implicated in the development of CVDs.

With these data we achieve several goals. First, we provide a resource for future studies

investigating aging-related changes in tissue-resident sEVs, a recently discovered source of EVs

that are currently poorly understood but provide exciting avenues for understanding and

manipulating disease-related intra-tissue signaling. Second, we contrast both these tissue-resident

EVs and classical plasma EVs as well as directly assess aged and young cargo from both

populations to identify aging-related changed in EV content and signaling, a sorely understudied

topic in EV research. Lastly, from the EV cargo we have identified both therapeutic miRNAs and

damage-associated protein cargo to direct future target-based studies and drug development for

notoriously difficult to target diseases, such as cardiac fibrosis. The dissemination of these data

will help promote more in-depth study of sEVs as both diagnostic and therapeutic vehicles to

investigate the effects of chronic changes to tissues, such as aging, and provide a useful tool for

the future investigation of comprehensive comparison of sEV populations.

RESULTS

Characteristics of sEV size profile change depending on biological source: Transmission

electron microscopy (TEM) of sEVs isolated from tissue (TEVs) or plasma (PEVs) of young (<40

y.o., n = 6 for tissue & plasma, Supplemental Table S1) and aged (>50 y.o., n = 6 for tissue &

plasma, Supplemental Table S1) donors showed that all particles were similar in overall

morphology and staining characteristics (Figure 1A). sEV size profile was validated via

Nanoparticle Tracking Analysis (NTA), and the resultant statistics showed that size was mostly

consistent between young and aged samples in PEVs (young mode: 129.5 nm; aged mode: 159.5

nm) but not TEVs (young mode: 128.5 nm; aged mode: 49.5 nm) (Figure 1B). Furthermore, TEVs

were predominantly smaller than PEVs overall, although all sEVs fell within the expected size

range for exosome-like sEVs. All populations were found to be monodisperse, though young

tissue-source TEVs had a notably larger PDI than any other cohort (Supplemental Table S2).

Different tissue source sEVs demonstrate differential responses to aging effects: Western blot

was performed to identify characteristic exosome markers CD9, CD63, and TSG101 on the sEVs

isolated, with each protein showing up clearly (Figure 1C). The measured quantity of each protein

was also assessed from the blot using ImageJ for standardized blot quantification, and the results

were Z-scored for comparison (Figure 1D). These show that changes in expression of these

particular exosomal surface proteins behave inversely with respect to age between TEVs and

PEVs, with PEVs showing a 0.5-fold to 1-fold decrease in expression of CD9 and CD63 between

the young and aged cohorts and TEVs showing a greater than 1-fold increase in expression across

that same comparison (Figure 1E). CD63 expression remained mostly consistent with CD9

expression in both populations with respect to age, though the change in expression was opposite

between TEVs and PEVs, but TSG101 availability was increased with age only in TEVs while

remaining consistent in PEVs, although it should be noted that these differences were not

significant with the metrics used. Taken with the above results, this suggests a shift in the

tetraspanin web and targeting abilities[39–41] that substantially differs between the TEVs and PEVs

for the proteins assessed.

Figure 1. Tissue resident sEVs display substantive differences from those isolated from plasma in size profile, surface protein expression, and behavior. Measurements were taken from n = 6 biological replicates per cohort, equally spread between male and female donors. (A) Transmission electron microscopy representative images for each cohort. (B) Nanoparticle tracking analysis differentiating young (gold) or aged (blue) source tissue sEVs (left) or plasma sEVs (right). The overall expression of exosomal surface proteins was measured by western blot (C) and compared (D) and the change in expression between groups was compared with respect to tetraspanin (E). (F) Uptake rate of TEVs or PEVs from young donors (n >= 3 for each) on human or mouse fibroblasts in 2D culture (n >= 3 replicates for each) over 24 hours, presented as change in uptake rate per hour between timepoints taken ± standard error. (G) Endpoint uptake rate of TEVs or PEVs on human or mouse fibroblasts, bars represent mean of biological replicates ± standard deviation and points represent mean of replicates for each biological replicate. * $p < 0.05$, ** $p < 0.01$, assessed by T-test with Welsch's correction for (E) and (G).

Uptake kinetics of sEVs varies between tissue source and destination tissue: Uptake kinetics of

sEVs with typical or atypical cell culture populations was also assessed (Figure 1F). When applied

to murine cells, both PEVs and TEVs maintained steady uptake by cells over 24 h, resulting in a

nearly flat line (right). When applied to human cardiac cells, however, PEVs showed a slight

increase in uptake over time whereas TEVs showed exponential increase in uptake rate over 24 h,

reaching several-fold higher uptake rates and uptake overall than PEVs (Figure 1G). Additional

assays with young vs aged cells (artificially aged iCFs or aged mice) showed that the age of the

cell had no significant effect on either the uptake rate of EVs or the size of the EVs produced

(Supplemental Figure S1), an unexpected result.

Profiling of miRNA content: miRNA profiling via Nanostring analysis revealed highly distinct

miRNA profiles between PEVs and TEVs (Figure 2A), with sources tending to cluster together

both when compiled and in individual replicates (Supplemental Figure S2). While sEVs from

young donors of both sources showed some notable overlap, the miRNA profiles of sEVs from

aged donors were almost completely distinct, with less than 100 miRNAs in common. Relative

expression values for all 798 detected miRNA expressed in these samples is provided in the

supplement (Supplemental Table S3). Prevalence of miRNAs in all sEVs was mostly normally

distributed (Figure 2B), though both PEV populations had a notable number of very highly

expressed (detected count >30 in chip) miRNAs. In general, sEVs from young donors tended to

express a larger number of miRNAs overall but at lower counts, while sEVs from aged donors

tended to express fewer miRNAs but at higher concentrations, consistent with previous findings[7].

Unsupervised analysis of the data to identify distinguishing miRNAs between TEVs and PEVs

revealed 125 potential targets (Figure 2C). Of these 125 miRNAs, 59 were expressed in TEVs but

not PEVs, and 66 were highly expressed in PEVs but not TEVs (Supplemental Table S4).

Figure 2. Summary of total miRNA profile from tissue or plasma-source sEVs. Measurements

were taken from n = 6 biological replicates per cohort, equally spread between male and female

donors. (A) Heatmap showing the full miRNA profile for each cohort. Supplementary figures are

available for individual replicates (Supplemental Figure S2) and for separation by both age and

sex (Supplemental Figure S3) of total miRNA profile. (B) Histogram detailing the overall

prevalence and distribution of miRNAs measured for each cohort. (C) Heatmap of the miRNA

targets identified via unsupervised analysis that distinguish between TEVs and PEVs. A total of

125 miRNAs were identified and were nearly evenly split between TEVs and PEVs, with TEVs

overexpressing 59 miRNAs and PEVs claiming 66. A full list of these miRNAs can be found in

the supplement (Supplemental Table S4). * $p < 0.05$.

Mapping of miRNA targets implicates TEV signaling in multi-point regulation of

cardiovascular health-related pathways: Of the 59 miRNA targets overexpressed in TEVs, 45

have been identified as "potentially involved" in CVD according to PubMed and other online

databases[26]. These 45 miRNAs were pulled out and visualized (Figure 3A), evalutated for potential

gene targets and overlap (Figure 3B, C), assessed via gene ontology (Figure 3D), and mapped

using MetaCore Pathway Analysis (Figure 3E-H). Likely gene targets were assessed for

overlapping targets of miRNAs and contributions of miRNAs to individual gene targets. Analysis

revealed 11,366 potential gene targets (Supplemental Table S5), though more than two-thirds were

targeted by only a single target miRNA. Of the target miRNAs, miR-519, miR-181, and miR-548

contributed to the largest number of gene targets (Supplemental Figure S4). Genes which were

targeted by 10 or more target miRNAs were selected for gene ontology analysis, resulting in

selection of 167 genes (Supplemental Figure S4, Supplemental Table S6). Within this cohort of

genes, miR-548 and miR-519 remained the top contributors with miR-129 supplanting miR-181,

though miR-181 remained a major contributor (Figure 3B). The top 10 targeted genes were also

identified (Supplemental Figure S5) along with contributing miRNAs (Figure 3C, Supplemental

Figure S5). Of the top 10, the disruption of regular signaling for ZBTB20, CLOCK, TNRC6B,

RORA, CREB1, MECP2, MINDY2, TAOK1, and AAK1 was linked to cardiovascular disease[42–

51], typically through analogous pathways to chronic CVD or similar chronic fibrosis diseases

although many of these genes were also identified targets in neurological disorders and ischemic

stroke. Gene ontology analysis of the 167 common gene targets was largely unsurprising, though

there was noted involvement in cellular metabolism, macromolecule biosynthesis, and cAMP, co-

SMAD, and JNK-kinase activity (Supplemental Figure S6). MetaCore analysis of the 45 miRNA

targets revealed four major pathways of interest which were regulated at several points by different

Figure 3. Cardiovascular-disease related target miRNAs and identified pathways identified from age and sex-related differences in sEV miRNA profile. Measurements were taken from n = 6 biological replicates per cohort, equally spread between male and female donors. (A) Heatmap showing identified target miRNAs enriched in TEVs relative to PEVs. (B) Analysis of miRNA contributors to predicted gene targets, showing by number the frequency of miRNAs contributing to different identified target genes (full gene list available in Supplemental Table S5). (C) Chord diagram showing overlap of miRNAs and the top 10 enriched predicted gene targets. (D) Gene ontology analysis of the predicted gene targets of the identified miRNAs showing the top 10 significant pathways identified (full graphs available in Supplemental Figure S6). (E-H) Results of MetaCore pathway analysis for the identified target miRNA data. * p < 0.05. Identified pathways were considered for analysis only for p-value < 0.05.

identified miRNAs along the pathway (Figure 3E-H), suggesting deliberate targeting of certain

pathways in a highly specific manner. The pathways of interest were the HNF4-a pathway (Figure

3E), with downstream regulation that promoted exosome trafficking and inhibited inflammation

and apoptotic signaling[52–56], the Smad3/STAT5 pathway (Figure 3F), with overall inhibition of

signaling early in the Ang2 cardiac fibrosis cascade[57–62], PPARg pathway/MAPK cascade (Figure

3G), with more multi-point regulation of Ang2-mediated fibrosis signaling and downstream targets

parallel to the HNF4-a pathway[63–67], and an SDF-1a/systemic stem cell recruitment pathway

implicated in pro-regenerative healing from MI[68–73] (Figure 4H).

Profiling of protein content: Protein profiling via mass spectrometry and subsequent analysis

revealed that TEVs and PEVs tended to separate into two distinct clusters depending on tissue

source (Figure 4A), although this was less consistent when comparing individual biological

replicates (Supplemental Figure S4). This may be due to some proteins being entirely absent from

detection in many biological replicates. To present a more readily comparable visual, the 93

proteins common across all biological replicates for all cohorts were also mapped (Figure 4B,

Supplemental Figure S7, Supplemental Table S7). While in the overall population TEVs showed

a slightly higher overall protein enrichment, with aged donor TEVs still having higher overall

protein enrichment than young donor TEVs, comparison of only the common proteins across all

replicates revealed a substantial increase in enrichment in aged donor TEVs compared to all groups

while young donor TEVs displayed notably low levels of relative protein enrichment. Further

subdivision of cohorts by sex in addition to age additionally reveals that this is not a sex-dependent

shift, though the overall protein enrichment may be sex-dependent (Supplemental Figure S8). This

is consistent with our previous findings on relative protein content between aged and young donor

TEVs[7]. Visualization of the overall protein population in terms of prevalence or absence of

proteins by number also revealed that many proteins were enriched inconsistently between

biological replicates (Figure 4C), with less than 500 (< 20% of all identified proteins via mass

spec) were universally present in any given cohort, and a similar number was universally absent

in every cohort.

Aging-correlated shifts in protein profile is similar between TEVs and PEVs: Breakdown of

protein enrichment by cohort was recorded via Venn Diagram (Figure 4D), and reveals that overall

both TEVs and PEVs have a similar number of overlapping and distinctive protein cargo. Both

populations, however, show similar changes in cargo with respect to age. In both TEVs and PEVs,

comparison of young donor sEV unique protein contents with that of aged donor sEV content

reveals a nearly 3-fold increase in unique proteins consistently expressed in all biological

replicates. Aged donor TEVs have a higher number of unique proteins compared to TEVs.

Gene ontology reveals functional differences in protein content between TEVs and PEVs: The

proteins enriched in TEVs or PEVs were also assessed via gene ontology analysis (Figure 4E, F).

Both TEV and PEV proteins were closely associated with extracellular vesicles and the

extracellular space, as expected, however the associated biological processes predicted were

notably different. TEVs were identified with muscular development and function (Figure 4E) as

well as a number of pathways corresponding to specifically cardiac development and homeostasis

and redox pathways (Supplemental Figure S9). PEV proteins, alternatively, corresponded more to

acute inflammatory response and related pathways (Figure 4F), and was associated with IgG and

other immunoglobulins and components (Supplemental Figure S9). Interestingly, the gene

ontology analysis returned no significant results for molecular functions pertaining to PEV

proteins.

263
Figure 4. Summary of total internal protein profile of differing age and sex cohorts from tissue or plasma-source sEVs. Measurements were taken from n = 6 biological replicates per cohort, equally spread between male and female donors. (A) Heatmap showing all 2547 proteins detected via mass spec that were measured for each cohort. (B) Heatmap showing the relative expression of the 93 proteins that were expressed in every biological replicate across all cohorts (Supplemental Table S7). Supplementary figures are available for individual replicates (Supplemental Figure S4) and for separation by both age and sex (Supplemental Figure S5) of both protein profiles. (C) Stacked bar plot showing the expression profile of all 2547 detected proteins for individual cohorts, separating between proteins that are absent in all biological replicates (red, top), present in all biological replicates (green, bottom), or neither (grey, middle). (D) Venn diagrams showing overlap of "present" proteins across cohorts, comparing between TEVs and PEVs overall (top left), only aged donor TEVs and PEVs (top right), young and aged donor TEVs (bottom left), and young and aged PEVs (bottom right). In comparisons where one side shows >33% greater quantity of proteins, the number of proteins was underlined. Gene ontology analysis was also performed for proteins isolated from TEVs (E) or PEVs (F) with the top 10 significant pathways identified (full graphs available in Supplemental Figure S9). * p < 0.05.

Tissue-resident sEVs showcase greater involvement in metabolism and homeostasis than

plasma sEVs: Proteomap profiles (Figure 5A) were generated from the unique proteins identified

for both TEVs (106 proteins) and PEVs (132 proteins) (Supplemental Table S8), as well as for

aged donor TEVs (274 proteins) and PEVs (202 proteins) (Supplemental Table S9). Interestingly,

overall TEV pathways were balanced between metabolism, cardiac muscle activity, and

environmental signal transduction, whereas overall PEV pathways were dominated by

immunoregulation and exosome handling. Overall TEVs also uniquely called a pathway for

degenerative diseases (black, far left, top left proteomap). Aged donor sEV protein pathways,

however, were fairly different. Aged donor TEVs showed less overall involvement in cardiac

muscle activity and greater involvement in environmental signaling and metabolism. Aged donor

TEVs, compared to overall, also became newly involved in immunoregulatory signaling as well

as developing pathways associated specifically with cancer development, hypertrophic

cardiomyopathy, and arrythmogenic right ventricular cardiomyopathy, as opposed to the

previously observed degenerative diseases. Aged donor PEVs on the other hand became solely

dominated by immunoregulatory signaling, excluding exosome handling almost entirely, and

developing involvement in pathways associated with Alzheimer's Disease. Potential individual

protein targets were also identified from the unique proteins from TEVs and PEVs via supervised

analysis (Figure 5B). These 17 proteins, while not necessarily directly implicated in major

pathways, are common biomarkers for identification of disease state or other pathology and may

allow for more rapid detection of critical marker quantification in clinical scenarios.

Tissue-resident sEVs from older hearts are directly implicated in CVDs, but not plasma sEVs

from older patients: The unique proteins from TEVs and PEVs were mapped via MetaCore

pathway analysis, and the resulting pathways provided further insight into the downstream

activities of TEVs (Figure 5C, D) and PEVs (Figure 5E). TEV proteins mediated NFkB/Interleukin

signaling downstream of c-Jun and TRAF6 (Figure 5C), activities highly involved in cardiac

immunoregulation and matrix homeostasis[74–78], as well as directly interfacing with PI3k-1a,

mTOR, and Laminin response (Figure 5D), which together can be implicated in response to cardiac

fibrosis or other cardiac matrix remodeling[79–85]. Additionally, dysregulation of PI3k-1a and mTOR

via the complex "PI3K/mTOR signaling pathway" network has been directly implicated in aging-

related CVD onset and other aging-related pathologies, and the targeted inhibition of this pathway

has reversed aging-related risk of post-MI cardiac fibrosis and pathological signaling post-cardiac

insult[86,87]. PEVs also interacted with the surrounding matrix through integrin mediation (Figure

5E) but displayed more focus on modulating interactions with white blood cells via Igg

manipulation, which may indicate a role in chronic inflammation in chronic CVD and heart

failure[88–90].

317
Figure 5. Cardiovascular-disease related target proteins and identified pathways identified
from age-related differences in sEV protein contents. Measurements were taken from n = 6
biological replicates per cohort, equally spread between male and female donors. Red: cellular
processes; Yellow: metabolism; Pink: organismal systems; Blue: genetic information processing;
Teal: environmental information processing; Purple: immune system; Black: human diseases. (A)
Proteomaps showing the KEGG pathways affected by the unique protein profiles of TEVs and
PEVs overall, as well as for only aged donor TEVs and PEVs. (B) Heatmap showing the 17
proteins of interest identified by supervised analysis of unique proteins for TEVs and PEVs. (C-
E) Results of MetaCore pathway analysis for the unique protein profiles of TEVs (C, D) or PEVs
(E). * p < 0.05. Identified pathways were considered for analysis only for p-value < 0.05.

DISCUSSION

In this study, we isolated and characterized small extracellular vesicles (sEVs) isolated from patient blood plasma samples and from donor heart tissue samples. All sEV samples were taken from an equal number of men (n = 6 for tissue & plasma) and women (n = 6 for tissue and plasma) to minimize the impact of sex variance, and were bifurcated between aged (>50 y.o., n = 6 for tissue & plasma) and young (<40 y.o., n = 6 for tissue & plasma) for analysis (Supplemental Table S1). Both the plasma sEVs (PEVs) and tissue sEVs (TEVs) were observed under transmission electron microscopy (TEM) and population size distribution was assayed via nanoparticle tracking analysis (NTA), with mean and mode population sizes being consistent between both assays. Interestingly, the mode size of the sEV population differed with age in TEVs, but not PEVs, suggesting at least some degree of differential function with respect to aging between each category of EV. Surface protein composition was also assessed between aged and young PEVs and TEVs using western blot for classical exosome markers CD9, CD63, and TSG101[21]. While overall measured expression of these proteins increased with age in TEVs and decreased with age in PEVs, the relative expression of these proteins did not significantly change in either group, suggesting that the quantity, but not overall composition, of tetraspanin web targeting proteins changes with age, and that these changes are different in PEVs and TEVs. Finally, the miRNA and protein content of young and aged PEVs and TEVs were assayed. The miRNA content was measured using Nanostring technology, providing a full profile of human miRNAs, and unsupervised analysis was performed to identify distinguishing miRNAs between PEVs and TEVs which were independent of age. The TEV-specific age-independent miRNAs were further analyzed in MetaCore pathway analysis to provide a preliminary overview of processes uniquely modulated by TEVs. The protein content of all cohorts was similarly assessed using mass

spectrometry, revealing 2547 proteins with varying expression levels across all biological

replicates. Analysis of the dataset revealed 93 proteins consistently expressed in all biological

replicates of all cohorts, with aged sEVs tending to have more commonly expressed proteins than

young sEVs and TEVs tending to have more commonly expressed proteins than PEVs. Unique

cohort proteins were also identified, and assessed for downstream function using KEGG

proteomapping software and MetaCore pathway analysis.

The size discrepancy in age-related size changes between TEVs and PEVs (Figure 1A, B) is an

interesting result due to the known consistency of size distribution in EV populations[21] and our

previous study showing that these size differences are also not sex-dependent[7]. We first evaluated

whether or not this discrepancy was due to different cell ages in the heart, as the difference due to

cell age could have been mitigated in PEVs by the large number of organs which contribute to the

PEV population. We evaluated this for both artificially aged iCFs (high passage cells at high

confluency) and mCFs from aged mice (72 weeks), but found that neither population showed

significantly different EV sizes produced compared to the young cells (Supplemental Figure S1).

In previous studies, we have shown that the stiffness of the surrounding matrix can significantly

alter the behavior of cells[10,13], so an alternative cause may be differences in the matrix stiffness

between young and aged hearts. This discrepancy will be further evaluated in future works to

identify the precise relationship, if any, between matrix stiffness and produced EV size.

The uptake rate of EVs was also assessed for both populations and revealed novel interactions of

TEVs and PEVs with tissue specificity, though only young PEVs and TEVs were used for this

assay due to the substantially smaller size of aged TEVs compared to all other cohorts, which we

have previously shown increases the uptake rate by cells[7]. To assess the specificity of TEVs and

PEVs, uptake was assays for both populations for human and non-human (mouse) CFs.

Interestingly, the uptake rate for both PEV groups and the non-human uptake of the TEVs was

essentially the same at all timepoints measured (Figure 1F) but the uptake of TEVs on human CFs

was significantly higher (Figure 1G) and in fact was increasing exponentially by the endpoint

(Figure 1F). This suggests some level of specificity of the TEVs that is not present in the PEVs

which may distinguish how EVs are selected to be exported to the plasma or to the surrounding

ECM. The consistency of tetraspanin expression in TEV populations (Figure 1E) may be a partial

contributor to this, as the tetraspanin web has been implicated in EV organ targeting

systemically[39–41], with several key, unidentified tetraspanins that are unique to TEVs contributing

to the uniqueness of this specificity. Additionally, the exponential increase in rate of uptake of

TEVs by human cells suggests that human cells taking up TEVs undergo some change which

facilitates further TEV uptake. This has not yet been reported in literature and may be a key aspect

of how TEVs provide therapeutic effects in young hearts, as an exponential increase in the uptake

of therapeutic compounds over a short period of time would actively counteract increased

inflammation and damage-associated effects common in the aftereffects of cardiac insult. These

results demand further study, and will be more fully evaluated in future works specifically

assessing the dynamics between uptake rate of natural or engineered EVs for the time-sensitive

delivery of therapeutic agents.

The differential miRNA cargo of TEVs and PEVs provides an interesting avenue for the

identification of therapeutic agents, as we have previously established that TEVs, but not PEVs,

can promote anti-fibrotic effects in heart cells post-insult[7]. In light of this, the overlap in miRNA

expression in the young PEVs and TEVs presents an interesting conundrum, although this is

resolved by the nearly bifurcated miRNA expression in the aged EV populations (Figure 2A),

suggesting that miRNA cargo substantially diverges between TEVs and PEVs with age. Despite

this, it was possible to completely distinguish between TEVs and PEVs as a whole via 125

identified miRNAs selected by untargeted analysis of the entire miRNA profile for all cohorts

(Figure 2B), providing further evidence that TEVs and PEVs are specialized for different

functions. Of the 125 differentiating miRNAs, 59 were enriched in TEVs and 66 were enriched in

PEVs, providing ample targets for further analysis of the specific function of each population. To

better understand the unique cardiac-related function of the TEVs, we manually assessed the 59

TEV-enriched miRNAs for potential involvement in any CVD or CVD-related pathways via

public databases and identified 45 targets for downstream analysis. MetaCore pathway analysis of

the 45 revealed several pathways which were consistently regulated by several of the target

miRNAs (Figure 3). These combined effects served to inhibit targets in the HNF4-α pathway,

Smad3/STAT5 pathway, and PPARγ pathway, and promote a therapeutic SDF-1α pathway. These

results imply several exciting possibilities. First, we have previously hypothesized the cooperative

effects of TEV miRNAs in promoting cardioprotective effects via synergistic inhibition of CVD-

associated pathways[7], but were only able to demonstrate this with 4 identified target miRNAs. The

combined regulation of 3 damage associated pathways at multiple points by 45 target miRNAs

provides a plethora of additional targets to evaluate in order to generate a TEV-mimetic cocktail

of beneficial miRNAs for therapeutic applications. The use of a large number of miRNAs which

endogenously cooperate to act on a small number of pathways would be able to provide much

greater effect specificity than existing single-target drugs and previously proposed single-miRNA

therapeutics. Additionally, these miRNAs together not only inhibit damage-associated effects but

also promote reparative signaling. The SDF-1α is a known cardioprotective pathway which is

difficult to activate without off-target effects. While the utility of these specific miRNAs require

additional study to determine efficacy and viability, this is an exciting first step in the development

of highly specific therapeutics using endogenous agents.

The protein cargo of both EV populations also provided exciting results. While comparison of all

identified proteins separated PEVs and TEVs, this result was not consistent across all biological

replicates (Figure 4A, Supplemental Figure S4). This was likely due to the inconsistency of protein

expression across biological replicates (Figure 3C), with >60% of proteins being expressed in

some, but not all, biological replicates of a given cohort. To be consistent in our analysis, we

focused on proteins which were either expressed consistently in all cohorts (Figure 3B), or were

uniquely expressed in a single cohort (Figure 4A), although cohort-unique targets were still

expressed in all biological replicates of a given cohort. Of the proteins consistently expressed in

all cohorts, aged EVs tended to express more proteins overall for both TEVs and PEVs (Figure

4B, C), which is consistent with our previous hypothesis[7]. This suggests that overall proteins

included in both systemic and tissue-resident increases with increasing age, which is consistent

with the expected effects of inflammaging. Inflammaging has been identified as a potential cause

of many aging-related pathologies, but is poorly defined and not well understood. The evaluation

of EVs as part of aging-related increases in inflammation may provide a novel avenue for increased

understanding of inflammaging and the development of intervention strategies.

Analysis of the proteins uniquely expressed in TEVs or PEVs via gene ontology and KEGG

proteomapping revealed highly different functions of each population. PEVs were largely involved

in immune system regulation (Figure 4F, Figure 5A), while TEVs were involved in a host of

metabolic and regulatory functions in the heart (Figure 4E, Figure 5A, Supplemental Figure S9).

Interestingly, only TEVs were involved in disease progression (Figure 5A). To evaluate the aging-

related changes in protein expression, the proteins unique to aged TEVs and PEVs were also

mapped (Figure 5A). Again, PEVs were largely involved with the immune system, though immune

system interactions made up a much larger proportion of involved pathways than PEVs overall,

and now showed involvement with the onset of Alzheimer's Disease pathways, which was an

interesting pathway to remain consistent across all biological replicates. TEVs, however, showed

much greater involvement in metabolism-related activities, less regulation of cardiac homeostasis,

and new involvement in local inflammation and the direct onset of CVDs, differentiating aged

TEVs from overall TEVs by direct implication in increased inflammation and disease progression.

From the proteins uniquely expressed in TEVs or PEVs overall, 17 proteins of interest were

identified which may be involved in the onset and progression of CVD. MetaCore analysis of these

proteins revealed mulit-point involvement in three pathways which, while not directly implicated

in CVD, are closely linked to the onset of several CVDs. These pathways included

NFkB/Interleukin signaling downstream of c-Jun and TRAF6, PI3k-1a, mTOR, and Laminin

response, which are each critical to regulating cardiac homeostasis and normal function. In

particular, PI3k-1a and mTOR together are a regulatory pathway that is increasingly dysregulated

with age, leading to increased risk of CVD, fibrosis, and heart failure[86,87], while the dysregulation

of Interleukin signaling and Laminin response pathways are closely related to the onset of chronic

fibrosis and other pathological cardiac matrix remodeling [79–85]. This data suggests that these

pathways are, in some part, regulated by TEV signaling and that aging-related changes in TEV

signaling, such as those we have shown, may disrupt these vital pathways and lead to increased

risk of CVD. While more direct evaluation of the roles of TEVs on the regulation of these pathways

is essential for understanding the complex relationship of aging-related EV changes and aging-

related pathologies, this data provides a first step into understanding these processes.

Overall, this study serves as an in-depth, comprehensive resource for identifying the cargo of sEVs

isolated from plasma or cardiac tissue, as well as distinguishing between samples from young or

aged individuals. These specific delineations are currently not well documented in literature, and

increased assessment and validation of age-related and organ-specific paracrine signaling

mechanisms will be essential in the development of next-generation therapeutics.

MATERIALS & METHODS

Human Heart Tissue Preparation: Donor human heart tissue was prepared according to our

previous protocol[7]. Briefly, the left ventricle from frozen heart tissue was excised of fat and

connective tissue and section into sub-300 µm slices, which were agitated in an aqueous acetic

acid (0.1%, Sigma Aldrich, USA) and ethanol (4%, Sigma Aldrich, USA) solution at 200 rpm at

room temperature (RT) for 2 hours. Tissues were then washed in sterile PBS for at least 30 minutes

to remove free debris, then re-agitated in solution overnight at RT. Matrices were then vigorously

washed in sterile PBS and sterile DI water, lyophilized, pulverized with pre-chilled mortar and

pestle, and separated into 200 mg (dry weight) powder aliquots. ECM powder was resuspended in

1 mL of aqueous Collagenase Type II (0.2 mg/mL, Corning, USA), Tris buffer (50 mM, Sigma

Aldrich, USA), and $CaCl_2$ (5 mM, Amresco, USA) and mixed vigorously. Suspension was reacted

overnight at RT with re-mixing every 6 h to prevent settling.

Plasma Sample Collection: Whole blood was collected from patients via direct veinous puncture

into ethylenediaminetetraacetic acid (EDTA)-treated tubes to prevent coagulation. Following

collection, whole blood was centrifuged at 1000g and 4°C for 5 min to separate plasma. Patient

plasma was then aliquoted to sterile RNAse-free tubes and shipped from the University of Florida

College of Medicine to the University of Notre Dame at -80°C. After arrival, samples were stored

at -80°C until use.

Vesicle Extraction and Isolation: For both digested heart tissue ECM and collected plasma

samples, the solution was centrifuged three times at 500g for 10 minutes, 2500g for 20 minutes,

and 10,000g for 30 minutes. The pellet was discarded after each centrifugation to remove any

insoluble contaminants. After the final centrifugation, the collected supernatant was

ultracentrifuged at 100,000g for 70 minutes at 4°C using an ultracentrifuge (Optima MAX-XP

Tabletop Ultracentrifuge, Beckman Coulter). The resulting pellet from ultracentrifugation was

either used immediately or stored dry at -80°C.

Transmission Electron Microscopy: Single pellets were fixed via resuspension in 2.5%

glutaraldehyde solution for 2 h at RT in the dark. Fixed samples were then loaded onto plasma-

cleaned Formvar/carbon-coated copper 200 mesh grids (Polysciences) and negative-stained with

Vanadium staining solution (Abcam, ab172780). The resulting grids were imaged at 80 kV with a

TEM (JEOL 2011, Japan).

Nanoparticle Tracking Analysis: Single pellets were resuspended in 500 µL of sterile, particle-

free PBS and measured using a NanoSight NS300 machine (Malvern Panalytical) with NTA

software version 3.2.16. Resuspended samples were kept at 4 °C until measurement, and

measurements were taken at RT.

Western Blot: Single pellets were lysed in RIPA buffer containing 1% proteinase inhibitor cocktail

(Brand, Country) at 4°C for 30 minutes and resultant protein concentration was measured via

bicinchoninic acid (BCA) assay (Pierce Chemical). Equal amounts of protein for each sample were

separated by 12% SDS-PAGE and transferred to blotting membranes. Blotting membranes were

incubated on a rocker overnight at 4°C with rabbit polyclonal primary antibodies against CD9

(Abcam, ab223052), CD63 (Abcam, ab216130), and TSG101 (Abcam, ab30871) at (1:1000), then

for 1 h at RT with HRP-conjugated goat anti-rabbit secondary antibody (Abcam, ab205718).

Stained membranes were then treated with a chemiluminescent substrate (Clarity ECL, Bio-Rad)

and imaged using a ChemiDoc-It2 imager (UVP, Analytik Jena) equipped with VisionWorks

software. Images were processed using ImageJ (NIH).

Cell Culture: In-house differentiated induced pluripotent stem cell (iPSC)-derived cardiac

fibroblasts (iCFs) were generated and cultured according to our previous protocol[13], and were

seeded on fibronectin (50 µL/mL) -coated 24-well cell culture plates. Mouse primary cardiac

fibroblasts were isolated from collected mouse hearts according to our previous protocol[7], and

were seeded on fibronectin-coated 24-well cell culture plates as with iCFs. Both iCFs and mCFs

were cultured in Dulbecco's Modified Eagle's Medium (DMEM, Thermo Fisher, USA),

supplemented with fetal bovine serum (FBS, 10%, Gibco, USA), penicillin/streptomycin

antibiotics (P/S, 1%, Life Technologies, USA), and SD208 (3 µM, Sigma Aldrich, USA).

Cellular Uptake of sEVs: Cellular uptake of sEVs was performed based on our previous protocol[7].

Briefly, sEV pellets were resuspending in PBS and concentration was assessed via BCA assay and

subsequently stained using the ExoGlow-Membrane staining kit (System Biosciences, USA) for

30 min at RT. Stained sEVs or a blank control were then purified in a salt gradient column for a

final sEV concentration of 25 µg/mL in DMEM, supplemented with exosome-free FBS (10%) and

1% P/S. Cells were incubated with the sEV-conditioned media at 37°C and imaged at 0, 3, 8, 16,

and 24 h incubation. During imaging the conditioned media was exchanged with sterile, particle-

free PBS, and the same media was re-placed in the same wells immediately preceding return to

incubation.

miRNA Isolation: Internal RNAs were isolated from sEVs using the Total Exosome RNA &

Protein Isolation Kit (Thermo Fisher Scientific, USA) according to the manufacturer's protocol.

Briefly, single pellets were resuspended in sterile, particle-free PBS and incubated with an equal

volume of provided denaturation buffer 4 °C for 5 min. The solution was then mixed with an equal

volume of Acid-Phenol:Chloroform by vortexing for 30 seconds and centrifuged for 5 min at

15,000g. The resulting aqueous phase was extracted and combined with 1.25x volume of 100%

ethanol, then transferred to the provided spin column. The spin column was centrifuged at 10,000g

for 15 seconds to bind and wash the RNA, then the RNA was eluted in 100 µL of the provided

elution solution. Eluted RNA was then concentrated using 3 kDa microcentrifuge spin filters

(Amicon, USA). 100 µL miRNA solution was worked up to 420 µL with RNAse-free water and

placed into a filter, then centrifuged at 14,000g for 90 minutes. Next, the filter was inverted into a

fresh collection tube, and centrifuged at 8,000g for 2 minutes. The resulting 20-25 µL of RNA

concentrate was immediately quantified by microvolume spectrophotometery (Nanodrop 2000,

Thermo Fisher Scientific, USA).

Profiling of Total miRNA Population: Concentrated RNA isolate was prepared for Nanostring

miRNA profiling according to the manufacturer's protocol. Briefly, the provided miRNA codeset

was mixed with the provided hybridization buffer to produce a master mix, and spike-in miRNA

controls were prepared at 200 pm. In order, the master mix, concentrated sample miRNA, spike-

in miRNA, and provided probes were mixed in a PCR plate and incubated at 65 °C for 16 h. The

hybridized solution was then mixed with 15 µL of provided hybridization buffer, for a total volume

of 30-35 µL, and added to the provided microfluidic cartridge. The assay was run with the provided

protocol for total miRNA analysis, and data was processed and analyzed using the provided

software using the recommended settings. The dataset was then exported for further analysis and

target selection.

miRNA Gene Target Identification and Gene Ontology: Likely gene targets of miRNAs were

predicted by miRDB[91], miRTarBase[92], and TargetScan[93], and commonly predicted genes were

compiled for further analysis (Supplemental Table S5). In-house software was used to identify

unique genes and match miRNAs to the corresponding gene targets to determine overlapping gene

targets between miRNAs of interest. Gene targets were selected as genes which were predicted

targets for greater than or equal to 10 miRNA targets (Supplemental Table S6).

Gene Ontology Analysis (GOA) was performed on the identified gene targets using an online

database via PANTHER Gene Ontology classification for enrichment analysis[94–96]. Data was

extracted from the output and graphed using R.

miRNA Pathway Analysis: Exported Nanostring data was normalized via Log10 normalization

and uploaded to the Clarivate MetaCore system for pathway analysis, then cleaned to remove

erroneous miRNA detection, if any, by the system. Automated network analysis was conducted

with 50 nodes per network. Results were presented as pathways obtained from the software.

Protein Isolation: Single pellets were lysed in 1% Triton-X solution at 4°C overnight, and

resultant protein concentration was measured via bicinchoninic acid (BCA) assay (Pierce

Chemical, USA). The resulting protein solution was solubilized in an SDS buffer solution

consisting of 5% SDS, 50 nM TEAB (pH 7.55) at room temperature until fully combined.

Total Protein Profiling: Total protein profiling was performed via mass spectrometery (Q

Exactive HF Mass Spectrometer) according to the manufacturer's protocol. Briefly, solubilized

protein samples were denatured in 20 mM DTT, then alkylated with 40 mM iodoacetamide.

Alkylated samples were digested in a provided column with trypsin (1:25 wt:wt) in 50 mM TEAB

(pH 8) at 37°C overnight. Digested samples were then desalted via ziptip C18 and dried, then

reconstituted in acetonitrile. Mass spectrometry was run with 1 µL of acetonitrile-reconstituted

solution for bottom-up proteomics.

Protein pathway mapping: GOA was performed on enriched proteins using the same protocol as

described above. Data was extracted from the output dataset and graphed using R. Proteomic

interactions of uniquely enriched proteins were classified through KEGG-based proteomapping

software[97] and are presented as obtained.

Statistical Analysis: Results were analyzed by one-way analysis of variance (ANOVA) with post-

hoc Tukey's HSD, two-way ANOVA with post-hoc Tukey's multiple comparison test, or a two-

tailed Student's t-test with Welch's correction for unequal standard deviation. Values are presented

as the mean ± standard deviation (SD) unless otherwise indicated, and differences were considered

significant when $p \leq 0.05$.

List of Supplementary Materials:

Fig S1 to S9

Table S1 to S9

REFERENCES

1.   Rahimi, K., Duncan, M., Pitcher, A., Emdin, C. A. & Goldacre, M. J. Mortality from heart failure,
acute myocardial infarction and other ischaemic heart disease in England and Oxford: a trend
study of multiple-cause-coded death certification. *J Epidemiol Community Health* **69**, 1000–1005
(2015).

2.   Reyes-Farias, M., Fos-Domenech, J., Serra, D., Herrero, L. & Sánchez-Infantes, D. White adipose
tissue dysfunction in obesity and aging. *Biochem Pharmacol* **192**, 114723 (2021).

3.   Laconi, E., Marongiu, F. & DeGregori, J. Cancer as a disease of old age: changing mutational and
microenvironmental landscapes. *British Journal of Cancer 2020 122:7* **122**, 943–952 (2020).

4.   Xi, J. Y., Lin, X. & Hao, Y. T. Measurement and projection of the burden of disease attributable to
population aging in 188 countries, 1990-2050: A population-based study. *J Glob Health* **12**,
(2022).

5.   Population Reference Bureau (PRB). Countries With the Oldest Populations in the World.
https://www.prb.org/resources/countries-with-the-oldest-populations-in-the-world/ (2020).

6.   Ozcebe, S. G., Bahcecioglu, G., Yue, X. S. & Zorlutuna, P. Effect of cellular and ECM aging on
human iPSC-derived cardiomyocyte performance, maturity and senescence. *Biomaterials* **268**,
120554 (2021).

7.   Ronan, G., Bahcecioglu, G., Yang, J. & Zorlutuna, P. Age and Sex-Dependent Differences in Human
Cardiac Matrix-Bound Exosomes Modulate Fibrosis through Synergistic miRNA Effects. *bioRxiv*
2022.11.14.516464 (2022) doi:10.1101/2022.11.14.516464.

8.   Ozcebe, S. G. & Zorlutuna, P. In need of age-appropriate cardiac models: Impact of cell age on
extracellular matrix therapy outcomes. *Aging Cell* **22**, e13966 (2023).

9.   Basara, G. *et al.* Myocardial infarction from a tissue engineering and regenerative medicine point
of view: A comprehensive review on models and treatments. *Biophys Rev* **3**, 031305 (2022).

10.   Basara, G., Gulberk Ozcebe, S., Ellis, B. W. & Zorlutuna, P. Tunable Human Myocardium Derived
Decellularized Extracellular Matrix for 3D Bioprinting and Cardiac Tissue Engineering. *Gels 2021,*
Vol. 7, Page 70 **7**, 70 (2021).

11.   Yang, J., Bahcecioglu, G. & Zorlutuna, P. The Extracellular Matrix and Vesicles Modulate the
Breast Tumor Microenvironment. *Bioengineering (Basel)* **7**, (2020).

12.   Ellis, B. W. *et al.* Adipose Stem Cell Secretome Markedly Improves Rodent Heart and hiPSC-
derived Cardiomyocyte Recovery from Cardioplegic Transport Solution Exposure. *Stem Cells* **39**,
170 (2021).

13.   Basara, G., Celebi, L. E., Ronan, G., Discua Santos, V. & Zorlutuna, P. 3D bioprinted aged human
post-infarct myocardium tissue model. *Health Sci Rep* **7**, e1945 (2024).

14.   Ozcebe, S. G., Tristan, M. & Zorlutuna, P. Adult Human Heart ECM Improves Human iPSC-CM
Function via Mitochondrial and Metabolic Maturation. *bioRxiv* 2023.10.31.565062 (2023)
doi:10.1101/2023.10.31.565062.

15.   Ellis, B. W. *et al.* Human Heart Anoxia and Reperfusion Tissue (HEART) Model for the Rapid Study
of Exosome Bound miRNA Expression As Biomarkers for Myocardial Infarction. *Small* **18**, 2201330
(2022).

16.   Ellis, B. W., Acun, A., Isik Can, U. & Zorlutuna, P. Human iPSC-derived myocardium-on-chip with
capillary-like flow for   personalized medicine. *Biomicrofluidics* **11**, 024105 (2017).

17.   Ronan, G., Bahcecioglu, G., Aliyev, N. & Zorlutuna, P. Engineering the cardiac tissue
microenvironment. *Progress in Biomedical Engineering* **6**, 012002 (2023).

18.   D'Anca, M. *et al.* Exosome Determinants of Physiological Aging and Age-Related
Neurodegenerative Diseases. *Front Aging Neurosci* **11**, 232 (2019).

19.   Eitan, E. *et al.* Age-Related Changes in Plasma Extracellular Vesicle Characteristics and
Internalization by Leukocytes. *Sci Rep* **7**, 1342 (2017).

20.   Sheta, M., Taha, E. A., Lu, Y. & Eguchi, T. Extracellular Vesicles: New Classification and Tumor
Immunosuppression. *Biology (Basel)* **12**, (2023).

21. Welsh, J. A. *et al.* Minimal information for studies of extracellular vesicles (MISEV2023): From
basic to advanced approaches. *J Extracell Vesicles* **13**, e12404 (2024).

22. Xu, M. Y., Ye, Z. S., Song, X. T. & Huang, R. C. Differences in the cargos and functions of exosomes
derived from six cardiac cell types: a systematic review. *Stem Cell Res Ther* **10**, 194 (2019).

23. Yáñez-Mó, M. *et al.* Biological properties of extracellular vesicles and their physiological
functions. *J Extracell Vesicles* **4**, 27066 (2015).

24. Jia, Y. *et al.* Small extracellular vesicles isolation and separation: Current techniques, pending
questions and clinical applications. *Theranostics* **12**, 6548 (2022).

25. Huang, K. *et al.* An off-the-shelf artificial cardiac patch improves cardiac repair after myocardial
infarction in rats and pigs. *Sci Transl Med* **12**, (2020).

26. Ren, X. *et al.* A multiplexed ion-exchange membrane-based miRNA (MIX·miR) detection platform
for rapid diagnosis of myocardial infarction. *Lab Chip* **21**, 3876–3887 (2021).

27. Bahcecioglu, G. *et al.* Aged Breast Extracellular Matrix Drives Mammary Epithelial Cells to an
Invasive and Cancer-Like Phenotype. *bioRxiv* 2020.09.30.320960 (2020)
doi:10.1101/2020.09.30.320960.

28. Yang, J., Bahcecioglu, G., Ronan, G. & Zorlutuna, P. Aged breast matrix bound vesicles promote
breast cancer invasiveness. *Biomaterials* **306**, 122493 (2024).

29. Basara, G. *et al.* Electrically conductive 3D printed Ti3C2Tx MXene-PEG composite constructs for
cardiac tissue engineering. *Acta Biomater* **139**, 179 (2022).

30. Law, M. R., Watt, H. C. & Wald, N. J. The underlying risk of death after myocardial infarction in
the absence of treatment. *Arch Intern Med* **162**, 2405–2410 (2002).

31. Cupples, L. A., Gagnon, D. R., Wong, N. D., Ostfeld, A. M. & Kannel, W. B. Preexisting
cardiovascular conditions and long-term prognosis after initial myocardial infarction: The
Framingham Study. *Am Heart J* **125**, 863–872 (1993).

32. Narins, C. R. *et al.* Relationship Between Intermittent Claudication, Inflammation, Thrombosis,
and Recurrent Cardiac Events Among Survivors of Myocardial Infarction. *Arch Intern Med* **164**,
440–446 (2004).

33. Zhu, J. *et al.* The incidence of acute myocardial infarction in relation to overweight and obesity: a
meta-analysis. *Arch Med Sci* **10**, 855 (2014).

34. Yusuf, S. *et al.* Effect of potentially modifiable risk factors associated with myocardial infarction in
52 countries (the INTERHEART study): case-control study. *Lancet* **364**, 937–952 (2004).

35. Florio, M. C., Magenta, A., Beji, S., Lakatta, E. G. & Capogrossi, M. C. Aging, MicroRNAs, and Heart
Failure. *Curr Probl Cardiol* **45**, 100406 (2020).

36. Anand, S. S. *et al.* Risk factors for myocardial infarction in women and men: insights from the
INTERHEART study. *Eur Heart J* **29**, 932–940 (2008).

37. Piccinini, A. M. & Midwood, K. S. Illustrating the interplay between the extracellular matrix and
microRNAs. *Int J Exp Pathol* **95**, 158–180 (2014).

38. Huleihel, L. *et al.* Matrix-bound nanovesicles within ECM bioscaffolds. *Sci Adv* **2**, (2016).

39. Smith, Z. J. *et al.* Single exosome study reveals subpopulations distributed among cell lines with
variability related to membrane content. *J Extracell Vesicles* **4**, 28533 (2015).

40. Rana, S., Yue, S., Stadel, D. & Zöller, M. Toward tailored exosomes: The exosomal tetraspanin
web contributes to target cell selection. *Int J Biochem Cell Biol* **44**, 1574–1584 (2012).

41. Barranco, I. *et al.* Extracellular vesicles isolated from porcine seminal plasma exhibit different
tetraspanin expression profiles. *Scientific Reports 2019 9:1* **9**, 1–9 (2019).

42. Li, F. *et al.* Zinc Finger Protein ZBTB20 protects against cardiac remodelling post-myocardial
infarction via ROS-TNFα/ASK1/JNK pathway regulation. *J Cell Mol Med* **24**, 13383–13396 (2020).

43. Corella, D. *et al.* CLOCK gene variation is associated with incidence of type-2 diabetes and
cardiovascular diseases in type-2 diabetic subjects: Dietary modulation in the PREDIMED
randomized trial. *Cardiovasc Diabetol* **15**, 1–12 (2016).

44. Edwards, T. L., Michels, K. A., Hartmann, K. E. & Velez Edwards, D. R. BET1L and TNRC6B associate
with uterine fibroid risk among European Americans. *Hum Genet* **132**, 943–953 (2013).

45. Bai, Z., Luo, Y. & Tian, L. ERCC5, HES6 and RORA are potential diagnostic markers of coronary
artery disease. *FEBS Open Bio* **12**, 1814–1827 (2022).

46. Zhang, T. & Ge, J. Mechanism of CREB1 in cardiac function of rats with heart failure via regulating
the microRNA-376a-3p/TRAF6 axis. *Mammalian Genome* **33**, 490–501 (2022).

47. Wang, C. *et al.* The Effect of Mecp2 on Heart Failure. *Cellular Physiology and Biochemistry* **47**,
2380–2387 (2018).

48. Bone, W. P. *et al.* Multi-trait association studies discover pleiotropic loci between Alzheimer's
disease and cardiometabolic traits. *Alzheimers Res Ther* **13**, 1–14 (2021).

49. Fang, C. Y., Lai, T. C., Hsiao, M. & Chang, Y. C. The Diverse Roles of TAO Kinases in Health and
Diseases. *International Journal of Molecular Sciences 2020, Vol. 21, Page 7463* **21**, 7463 (2020).

50. Song, Y. *et al.* M2 Microglia Extracellular Vesicle miR-124 Regulates Neural Stem Cell
Differentiation in Ischemic Stroke via AAK1/NOTCH. *Stroke* **54**, 2629–2639 (2023).

51. Xin, X. *et al.* Development and therapeutic potential of adaptor-associated kinase 1 inhibitors in
human multifaceted diseases. *Eur J Med Chem* **248**, 115102 (2023).

52. Rafiq, K. *et al.* C-Cbl inhibition improves cardiac function and survival in response to myocardial
ischemia. *Circulation* **129**, 2031–2043 (2014).

53. Tang, Z. *et al.* Suppression of c-Cbl tyrosine phosphorylation inhibits neointimal formation in
balloon-injured rat arteries. *Circulation* **118**, 764–772 (2008).

54.   De Melker, A. A., Van Der Horst, G., Calafat, J., Jansen, H. & Borst, J. c-Cbl ubiquitinates the EGF
710   receptor at the plasma membrane and remains receptor associated throughout the endocytic
711   route. *J Cell Sci* **114**, 2167–2178 (2001).

712   55.   Cormont, M. *et al.* CD2AP/CMS Regulates Endosome Morphology and Traffic to the Degradative
713   Pathway Through its Interaction with Rab4 and c-Cbl. *Traffic* **4**, 97–112 (2003).

714   56.   Ravandi, A. *et al.* Relationship of IgG and IgM autoantibodies and immune complexes to oxidized
715   LDL with markers of oxidation and inflammation and cardiovascular events: Results from the
716   EPIC-norfolk study. *J Lipid Res* **52**, 1829–1836 (2011).

717   57.   Zheng, Y. *et al.* Modulation of STAT3 and STAT5 activity rectifies the imbalance of Th17 and Treg
718   cells in patients with acute coronary syndrome. *Clinical Immunology* **157**, 65–77 (2015).

719   58.   Kurdi, M., Zgheib, C. & Booz, G. W. Recent Developments on the Crosstalk Between STAT3 and
720   Inflammation in Heart Function and Disease. *Front Immunol* **9**, 421289 (2018).

721   59.   Pang, Q. *et al.* Regulation of the JAK/STAT signaling pathway: The promising targets for
722   cardiovascular disease. *Biochem Pharmacol* **213**, 115587 (2023).

723   60.   Liu, J. ; *et al.* The Role of JAK/STAT Pathway in Fibrotic Diseases: Molecular and Cellular
724   Mechanisms. *Biomolecules 2023, Vol. 13, Page 119* **13**, 119 (2023).

725   61.   Hanna, A., Humeres, C. & Frangogiannis, N. G. The role of Smad signaling cascades in cardiac
726   fibrosis. *Cell Signal* **77**, 109826 (2021).

727   62.   Zhang, M. *et al.* Notch3 Ameliorates Cardiac Fibrosis After Myocardial Infarction by Inhibiting the
728   TGF-β1/Smad3 Pathway. *Cardiovasc Toxicol* **16**, 316–324 (2016).

729   63.   Lu, Q. *et al.* Targeting peroxisome proliferator-activated receptors: A new strategy for the
730   treatment of cardiac fibrosis. *Pharmacol Ther* **219**, 107702 (2021).

731   64.   Liu, H. J., Liao, H. H., Yang, Z. & Tang, Q. Z. Peroxisome proliferator-activated receptor-γ is critical
732   to cardiac fibrosis. *PPAR Res* **2016**, (2016).

733   65.   Li, X. *et al.* Aspirin Reduces Cardiac Interstitial Fibrosis by Inhibiting Erk1/2-Serpine2 and P-Akt
734   Signalling Pathways. *Cellular Physiology and Biochemistry* **45**, 1955–1965 (2018).

735   66.   Zhang, Z., Yang, Z., Wang, S., Wang, X. & Mao, J. Targeting MAPK-ERK/JNK pathway: A potential
736   intervention mechanism of myocardial fibrosis in heart failure. *Biomedicine & Pharmacotherapy*
737   **173**, 116413 (2024).

738   67.   Mohamad, H. E., Askar, M. E. & Hafez, M. M. Management of cardiac fibrosis in diabetic rats; The
739   role of peroxisome proliferator activated receptor gamma (PPAR-gamma) and calcium channel
740   blockers (CCBs). *Diabetol Metab Syndr* **3**, 1–12 (2011).

741   68.   Pillarisetti, K. & Gupta, S. K. Cloning and relative expression analysis of rat stromal cell derived
742   factor-1 (SDF-1)1: SDF-1 α mRNA is selectively induced in rat model of myocardial infarction.
743   *Inflammation* **25**, 293–300 (2001).

69. Boyle, A. J. *et al.* Myocardial production and release of MCP-1 and SDF-1 following myocardial
infarction: Differences between mice and man. *J Transl Med* **9**, 1–8 (2011).

70. Chen, Y. *et al.* Testosterone replacement therapy promotes angiogenesis after acute myocardial
infarction by enhancing expression of cytokines HIF-1a, SDF-1a and VEGF. *Eur J Pharmacol* **684**,
116–124 (2012).

71. Zhong, J. & Rajagopalan, S. Dipeptidyl peptidase-4 regulation of SDF-1/CXCR4 axis: Implications
for cardiovascular disease. *Front Immunol* **6**, 159972 (2015).

72. Döring, Y., Pawig, L., Weber, C. & Noels, H. The CXCL12/CXCR4 chemokine ligand/receptor axis in
cardiovascular disease. *Front Physiol* **5 JUN**, 88349 (2014).

73. Wen, J., Zhang, J.-Q., Huang, W. & Wang, Y. SDF-1α and CXCR4 as therapeutic targets in
cardiovascular disease. *Am J Cardiovasc Dis* **2**, 20 (2012).

74. Fiordelisi, A., Iaccarino, G., Morisco, C., Coscioni, E. & Sorriento, D. NFkappaB is a Key Player in
the Crosstalk between Inflammation and Cardiovascular Diseases. *International Journal of
Molecular Sciences 2019, Vol. 20, Page 1599* **20**, 1599 (2019).

75. Cheng, W. *et al.* NF-κB, A Potential Therapeutic Target in Cardiovascular Diseases. *Cardiovascular
Drugs and Therapy 2022 37:3* **37**, 571–584 (2022).

76. Przybyszewski, E. M., Targher, G., Roden, M. & Corey, K. E. Nonalcoholic Fatty Liver Disease and
Cardiovascular Disease. *Clin Liver Dis (Hoboken)* **17**, 19–22 (2021).

77. Abdullah, M., Berthiaume, J. M. & Willis, M. S. Tumor necrosis factor receptor-associated factor 6
as a nuclear factor kappa B-modulating therapeutic target in cardiovascular diseases: at the heart
of it all. *Translational Research* **195**, 48–61 (2018).

78. Huang, W. *et al.* MicroRNA-3614 regulates inflammatory response via targeting TRAF6-mediated
MAPKs and NF-κB signaling in the epicardial adipose tissue with coronary artery disease. *Int J
Cardiol* **324**, 152–164 (2021).

79. Ghigo, A., Laffargue, M., Li, M. & Hirsch, E. PI3K and Calcium Signaling in Cardiovascular Disease.
Circ Res **121**, 282–292 (2017).

80. Ghigo, A., Morello, F., Perino, A. & Hirsch, E. Therapeutic Applications of PI3K Inhibitors in
Cardiovascular Diseases. *Future Med Chem* **5**, 479–492 (2013).

81. Qin, W., Cao, L. & Massey, I. Y. Role of PI3K/Akt signaling pathway in cardiac fibrosis. *Molecular
and Cellular Biochemistry 2021 476:11* **476**, 4045–4059 (2021).

82. Chong, Z. Z., Shang, Y. C. & Maiese, K. Cardiovascular Disease and mTOR Signaling. *Trends
Cardiovasc Med* **21**, 151–155 (2011).

83. Yang, Z. & Ming, X. F. mTOR signalling: the molecular interface connecting metabolic stress, aging
and cardiovascular diseases. *Obesity Reviews* **13**, 58–68 (2012).

84. Sciarretta, S., Forte, M., Frati, G. & Sadoshima, J. New Insights Into the Role of mTOR Signaling in
the Cardiovascular System. *Circ Res* **122**, 489–505 (2018).

85. Samidurai, A., Kukreja, R. C. & Das, A. Emerging role of mTOR signaling-related miRNAs in
cardiovascular diseases. *Oxid Med Cell Longev* **2018**, (2018).

86. Das, A. & Reis, F. mTOR Signaling: New Insights into Cancer, Cardiovascular Diseases, Diabetes
and Aging. *International Journal of Molecular Sciences 2023, Vol. 24, Page 13628* **24**, 13628
(2023).

87. Ghafouri-Fard, S. *et al.* Interplay between PI3K/AKT pathway and heart disorders. *Mol Biol Rep*
49, 9767–9781 (2022).

88. Shen, A. R. *et al.* Integrin, Exosome and Kidney Disease. *Front Physiol* **11**, 627800 (2021).

89. van den Hoogen, P. *et al.* Increased circulating IgG levels, myocardial immune cells and IgG
deposits support a role for an immune response in pre- and end-stage heart failure. *J Cell Mol*
Med **23**, 7505–7516 (2019).

90. Tsimikas, S. *et al.* Relationship of IgG and IgM autoantibodies to oxidized low density lipoprotein
with coronary artery disease and cardiovascular events. *J Lipid Res* **48**, 425–433 (2007).

91. Chen, Y. & Wang, X. miRDB: an online database for prediction of functional microRNA targets.
Nucleic Acids Res **48**, D127 (2019).

92. Huang, H. Y. *et al.* miRTarBase update 2022: an informative resource for experimentally validated
miRNA–target interactions. *Nucleic Acids Res* **50**, D222 (2021).

93. Agarwal, V., Bell, G. W., Nam, J. W. & Bartel, D. P. Predicting effective microRNA target sites in
mammalian mRNAs. *Elife* **4**, e05005 (2015).

94. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium. *Nat Genet* **25**, 25–29 (2000).

95. Consortium, G. O. The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids Res* **49**,
D325–D334 (2021).

96. Mi, H., Muruganujan, A., Ebert, D., Huang, X. & Thomas, P. D. PANTHER version 14: more
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. *Nucleic Acids*
Res **47**, D419–D426 (2019).

97. Liebermeister, W. *et al.* Visual account of protein investment in cellular functions. *Proc Natl Acad*
Sci U S A **111**, 8488–8493 (2014).

ACKNOWLEDGEMENTS

We thank Dr. Kieth March for providing the plasma samples utilized in this study.

The lyophilization of decellularized ECM was conducted at the Center for Environmental Science

and Technology (CEST) at the University of Notre Dame.

We thank the Biophysics Instrumentation (BIC) Core Facility for the use of Optima MAX-XP

Tabletop Ultracentrifuge.

The authors acknowledge the use of the Electron Microscopy Core of the Notre Dame Integrated

Imaging Facility, a designated core of the NIH-funded Indiana Clinical and Translational Sciences

Institute.

The Nanoparticle Tracking Analysis was conducted using the NanoSight NS300 at the Harper

Cancer Research Institute (HCRI) Tissue Core Facility.

Funding: Research reported in this publication was supported by NSF-CAREER Award #

1651385, NSF CBET Award # 1805157 and NIH Award # 1 R01 HL141909-01A1

Competing Interests Statement: The authors have no competing interest to disclose.

Data Availability Statement: All data required for production of the manuscript is included in

this submission. Additional raw data can be provided upon request.

Ethics Statement: Human heart tissue was collected from donors whose hearts were deemed

unsuitable for transplantation through the Indiana Donor Network (IDN) or National Disease

Research Institute (NDRI). IRB approval was waived, as no identifying information was provided

by the Indiana Donor Network or NDRI. De-identified human whole blood was collected from

live donors at the University of Florida College of Medicine through an IRB approved standard

collection protocol (IRB#201901232). All tissue collection was performed in accordance with the

declaration of Helsinki.