PARKINSON'S DISEASE-ASSOCIATED *PINK1* LOSS DISRUPTS ENSHEATHING GLIA AND CAUSES DOPAMINERGIC NEURON SYNAPSE LOSS

Lorenzo Ghezzi[1,2], Sabine Kuenen[1,2], Ulrike Pech[1,2], Nils Schoovaerts[1,2], Ayse Kilic[1,2], Suresh Poovathingal[1,2,3], Kristofer Davie[1,2,], Jochen Lamote[1,3], Roman Praschberger[1,2,4, #], Patrik Verstreken[1,2,#]

[1]VIB-KU Leuven Center for Neuroscience; Leuven, Belgium.

[2]KU Leuven, Department of Neurosciences, Leuven Brain Institute, Leuven, Belgium.

[3]VIB Flow Core Leuven, VIB Technologies, Leuven, Belgium

[4]Medical University of Innsbruck, Institute of Human Genetics; Innsbruck, Austria.

[#]Correspondence: patrik.verstreken@kuleuven.be, roman.praschberger@i-med.ac.at

ABSTRACT

Parkinson's disease (PD) is commonly associated with the loss of dopaminergic neurons in the *substantia nigra*, but many other cell types are affected even before neuron loss occurs. Recent studies have linked oligodendrocytes to early stages of PD, though their precise role is still unclear. *PINK1* is mutated in familial PD, and through unbiased single-cell sequencing of the entire brain of *Drosophila Pink1* models, we observed significant gene deregulation in ensheathing glia (EG); cells that share functional similarities with oligodendrocytes. We found that the loss of *Pink1* leads to abnormalities in EG, similar to the reactive response of EG seen upon nerve injury. Using cell-type-specific transcriptomics, we identified deregulated genes in EG as potential functional modifiers. Specifically downregulating two trafficking factors in EG, Vps35 and Vps13, also mutated in PD, was sufficient to rescue neuronal function and protect against dopaminergic synapse loss. Our findings demonstrate that *Pink1* loss in neurons triggers an injury-like response in EG, and that *Pink1* loss in EG in turn disrupts neuronal function. Vesicle trafficking components, which may regulate membrane interactions between organelles in EG, seem to play a role in maintaining neuronal health and ultimately preventing dopaminergic synapse loss. Our work highlights the essential role of glial support cells in the pathogenesis of PD and identifies vesicle trafficking within these cells in disease progression.

INTRODUCTION

Parkinson's disease (PD) is characterized by motor symptoms such as bradykinesia, rigidity and tremor that are caused by the progressive loss of dopaminergic neurons in the substantia nigra (Lang & Lozano, 1998). However, most of the patients report non-motor symptoms, such as constipation, hyposmia and sleep defects, even before the onset of motor symptoms (Munhoz et al., 2015). This suggests that PD is a progressive neurodegenerative disease that initiates many years before the diagnosis and involves different neuronal systems, multiple anatomical areas and different cell types. While therapies symptomatically improve motor functioning temporarily by restoring dopaminergic tone, they do not effectively prevent neurodegeneration and disease progression. This highlights the necessity to identify cell-types and biological mechanisms that act at the earliest stages of disease.

With the advent of single-cell sequencing applied to PD and control brains in combination with evidence from genome-wide association studies (GWAS), it is now possible to identify which cells and mechanisms are at play in PD models that recapitulate early stages of PD (Kaempf et al., 2026; Pech et al., 2025). Among these, one study reported that genes related to GWAS loci are enriched for oligodendrocyte-specific gene expression (Bryois et al., 2020). Interestingly, differential gene expression in post-mortem brains with Braak scores of 1-2, 3-4, and 5-6 and controls, suggested a role for oligodendrocytes at an early stage of PD prior to the overt loss of dopaminergic neurons (Bryois et al., 2020). Additionally, two independent single-cell sequencing studies, one of the human substantia nigra and one of the midbrain, both associated PD genetic risk with oligodendrocyte-specific expression patterns (Agarwal et al., 2020; Smajic et al., 2022). While these studies started to link dysfunction of oligodendrocytes to the early stages of PD, they did not address their role in the pathophysiology of the disease.

Here, we describe a *Drosophila* model to start assessing how neuron-glia cross-talk contributes to PD. *Drosophila* has been successfully used to investigate cellular and molecular dysfunction preceding the onset of age-related symptoms, including dopaminergic neuron-dependent motor symptoms (Kaempf et al., 2026; Pech et al., 2025; Valadas et al., 2018). Furthermore, *Drosophila* glial cells display several anatomical and functional features that show remarkable similarity to their mammalian counterparts (Freeman & Doherty, 2006; Kremer et al., 2017). While the small size of the *Drosophila* brain does not necessitate elaborate myelination, ensheathing glia (EG) in this species share not only anatomical features with oligodendrocytes, such as the ability to wrap around nerve tracts in the Central Nervous System (CNS) (Kremer et al., 2017; Yildirim et al., 2019), but also

important molecular and functional features, such as providing metabolic support and regulating neuronal activity (Delgado et al., 2018; Otto et al., 2018).

In a brain-wide single-cell sequencing experiment, we identified EG as the most deregulated cell type in a young (pre-motor) *Pink1* loss-of-function *Drosophila* model. Using cell-specific labeling, immunohistochemistry, and electrophysiological recordings, we correlated the transcriptional deregulation in EG with defects that appear similar to those seen upon nerve injury. Additionally, we find that healthy EG are fundamental to support dopaminergic synapse integrity in *Pink1* mutants. Finally, using cell-type-specific transcriptomic, high-throughput screening and immunohistochemistry, we identified vesicle trafficking as a genetic modifier and showed that *Vps13* and *Vps35* are EG-expressed regulators that maintain dopaminergic synapse integrity. Our study shows that neuron-ensheathing glia crosstalk is already disrupted at a young age by the loss of *Pink1* in neurons, but also in EG, and that EG are fundamental to support dopaminergic synapse-integrity, suggesting a role for oligodendrocytes in the progression of PD.

RESULTS

Ensheathing glia in Pink1 loss-of-function flies show cell non-autonomous defects

We previously created a whole-brain single-cell RNAseq dataset from different *Drosophila* PD knock-in models. This dataset was generated from young 5-day-old flies to capture 'early' changes (Pech et al., 2025). We re-analyzed the data from *Pink1*[P399L] knock-in loss-of-function mutant flies and isogenic controls and performed differential gene expression (DEG) analysis for each cell type using DESeq2 (Figure 1A). Glial cell subtypes show the highest degree of deregulation; particularly, ensheathing glia are strongly affected (Figure 1A). While flies do not myelinate neurons, ensheathing glia (EG) serve similar supporting functions as oligodendrocytes (Otto et al., 2018; Yildirim et al., 2019). These results suggest that ensheathing glia are deregulated in young *Pink1*[P399L] mutants (and also *Pink1*[KO-WS], see below) at an age prior to dopaminergic neuron-dependent motor defects (Kaempf et al., 2026; Pech et al., 2025).

When neurons are damaged, activated EG invade the neuropil (Doherty et al., 2009; MacDonald et al., 2006). For example, when fly antennal lobes are removed, severing the Olfactory Receptor Neurons (ORNs), EG invade the antennal lobe neuropil (MacDonald et al., 2006), potentially as a protective response (Figure 1B'-C). To test if EG invaded the neuropil in *Pink1* mutant flies, we expressed a transmembrane fluorescent protein (*UAS-CD8GFP*) under the control of a promoter specific to EG (*MZ709-Gal4*) (Doherty et al., 2009, Supplementary figure 1) in control and *Pink1*[KO]

flies. Samples were labeled with anti-GFP and the pre-synaptic protein Bruchpilot (NC82). Similar to ORN-severed controls, GFP signal, lining the membrane of EG, is visible inside the antennal neuropil in *Pink1^{KO}*, and severing the ORN in *Pink1^{KO}* does not further exacerbate this phenotype (Figure 1B"-C). Even though we cannot exclude that in *Pink1* mutants the GFP signal is merely upregulated, *Pink1* loss and neuron-severing that triggers EG activation seem to have a similar phenotype.

In order to determine whether this phenotype is cell-autonomous, we generated cell-type-specific *Pink1* perturbations. We either downregulated *Pink1* using a previously well-validated RNAi line specifically in EG (EG-specific *Pink1* loss-of-function – Figure 1D'), or we re-expressed wild-type *Pink1* specifically in EG in *Pink1^{KO-WS}* flies (all cells, but the EG, loss-of-function - Figure 1D"). When *Pink1* is knocked-down in EG (but present in other cells, including neurons), no GFP signal is detected in the neuropil (Figure 1D''-1E). However, when *Pink1* is not present in neurons and expressed in EG, the GFP is present inside the antennal neuropil (Figure 1D"-E). This suggests that *Pink1*-defective neurons activate the EG in a cell-nonautonomous fashion to invade, and potentially protect, the neuropil.

Ensheathing glia function supports synaptic integrity

Our finding that *Pink1^{KO-WS}* in neurons elicits a cell non-autonomous response in EG suggested that EG might, in turn, also functionally modulate neuronal integrity in mutant flies. To test this, we made use of a readily accessible model circuit in the fly visual system with an easy electrophysiological read-out, electroretinograms (ERG). Notably, whereas our functional readout here interrogates neuronal activity in the visual system, our analysis of EG morphology was performed in the antennal lobe, a distinct brain region. We therefore make the explicit assumption that EG perform comparable functions across fly brain regions. This assumption is supported by our single-cell sequencing dataset, in which we did not detect region-specific segregation of EG into distinct clusters that could be attributed to different brain areas. The ERG waveforms are susceptible to changes in intracellular signaling, neuronal function, and synaptic transmission, and have been well characterized and amply used to assess neuronal and cellular function (Hardie & Raghu, 2001; Praschberger et al., 2023; Soukup et al., 2016; C. F. Wu & Wong, 1977). We exploited this method to understand the role of *Pink1* in EG on neuronal function and synaptic transmission in young flies.

Comparing control with young *Pink1^{KO-WS}* flies does not show a difference in the "depolarization-response", indicating that -as expected- there is no neurodegeneration occurring in the photoreceptors at this stage (Figure 2A). However, when comparing the ON peak – which represents synaptic transmission from the photoreceptors and within the underlying neuronal circuit, and is a more robust readout of this than the OFF peak (Vilinsky & Johnson, 2012) – we detect a significant

reduction in mutant compared to control animals, suggesting that in $Pink1^{KO-WS}$ flies, synaptic communication in this circuit is partially impaired, already at 5-day-old (Figure 2A-B).

In order to discern the role of EG in this defective ERG response, we (1) downregulated $Pink1$ specifically in EG and (2) we expressed $Pink1$ specifically in the EG of $Pink1^{KO-WS}$. When $Pink1$ is knocked down specifically in EG, we observe the same reduction in ON peak response as in the $Pink1^{KO-WS}$ flies. This indicates that EG integrity is necessary for efficient neurotransmission in this circuit. Conversely, the ON peak defect of $Pink1^{KO-WS}$ mutants is rescued when $Pink1$ is reintroduced in EG only. This suggests that activated EG are able to, at least partially, buffer the detrimental effects of $Pink1$ deficiency in neurons (Figure 2A-B).

To also test EG function in relation to PD-relevant dopaminergic (DA) synapses, we evaluated the loss of dopaminergic neuron afferents. In a previous paper from our laboratory, we have shown that the loss of $Pink1$ function causes a progressive loss of protocerebral anterior medial dopaminergic neuron (PAM DAN) afferents in the Mushroom Body (MB) neuropil in 25-day-old animals, but not when they are 5-day-old (Kaempf et al., 2026). Indeed, we confirm that 25-day-old animals show a decrease in the dopaminergic synaptic area in MB lobes (Figure 2 C-D). Hence, while ensheathing glia are already activated at a young age in $Pink1^{KO-WS}$, DA synapses are structurally still intact (at 5 days of age) and only deteriorate in older flies. We then knocked down $Pink1$ specifically in EG and found a similar strong decrease in DA synaptic area (Figure 2C'-D). Conversely, when $Pink1$ is reintroduced specifically in EG of $Pink1^{KO-WS}$ animals, DA synapse-loss is significantly rescued (Figure 2C"-D and Supplementary figure 2). Together, these results indicate that $Pink1$ in EG is necessary to protect dopaminergic neurons from synapse loss.

EG cell type-specific transcriptomics reveals modifiers of neuronal dysfunction

To identify pathways in EG that are deregulated by $Pink1$ loss-of-function, we optimized a method to isolate EGs from the rest of the brain, enabling us to perform much higher-sensitivity transcriptomics than what we achieved with our 10x droplet-based single-cell sequencing (Pech et al., 2025). We expressed a fluorescently-tagged histone (*UAS-HisTag-eGFP*) in EG (*GMR-56-Gal4*) and used Fluorescence-activated Cell Sorting (FACS) to isolate EG. We sorted 300 cells per technical replicate and performed Bulk-RNA sequencing using a SMART-seq2 protocol (Figure 3A). The sequencing results of EG-sorted cells and of neuron-sorted cells (*nSyb-Gal4* instead of *GMR-56-Gal4*) show strong enrichment of EG cell markers in the former and strong enrichment of neuronal markers in the latter (Figure 3B). We then compared the transcriptomic profile of EG in $Pink1^{KO-WS}$ to that of control using differential expression analysis and found 617 deregulated genes in EG (Appendix 1, padj<0.05) (Figure 3C). To understand whether the deregulated genes were enriched for a specific pathway or

associated with a cellular component compared to non-significantly deregulated genes, we performed Gene Ontology analysis of the deregulated genes in Pink1^{KO-WS} flies. Our analysis resulted in generic terms and very diverse pathways that did not allow us to draw further conclusions.

To test if the genes deregulated in EG are modifiers of the neuronal pink1^{KO-WS} phenotypes, we downregulated and upregulated (when possible) the 50 most deregulated genes (Figure 3C), for which we could find RNAi fly lines, specifically in the EG of Pink1^{KO-WS} flies and recorded ERGs in 5-day-old animals. Knock down of one of the genes (CG17660) resulted in a rescue of the ON transient defects in Pink1^{KO-WS} (suppressors at p<0.05), and a few exacerbated the defect (Figure 3D). Interestingly, the human orthologs of CG17660, TMEM87A/B, have an established role in endosomal sorting (Gaudet et al., 2011; Hirata et al., 2015). Hence, our genetic screening suggests that Pink1^{KO-WS} phenotypes might be suppressed by downregulating genes involved in vesicle trafficking in EG.

PD causative vesicle trafficking gene downregulation in EG rescues synaptic dopaminergic neurons afferent loss

Because our ERG-based modifier screen for the Pink1^{KO-WS} phenotype identified a gene with a known role in vesicle trafficking, we next asked whether Pink1 in EG also genetically interacts with other PD-causative genes that function in vesicle trafficking, specifically vesicle trafficking components affecting membrane interactions between mitochondria and the endoplasmic reticulum (ER), like Vps35 and Vps13 (Brickner & Fuller, 1997; Lesage et al., 2016a; Vilariño-Güell et al., 2011; Zimprich et al., 2011). We find that the knock down of each of these genes in EG also rescues the ON transient defect of Pink1^{KO-WS} mutants (Figure 4A-B).

To determine if manipulation of this pathway in EG can also rescue the synaptic innervation defect of PAM DAN onto the mushroom bodies in old (20-25-day-old) Pink1^{KO-WS} mutants, we expressed RNAi to Vps13 in EG using MZ709-Gal4. Anti-TH labeling (marking DAN synapses) in the MB area (marked by anti-DLG) is significantly reduced in aged Pink1^{KO-WS}, but is rescued to control levels when Vps13 is knocked down in EG (Figure 4C-D, Supplementary figure 3). These results suggest that manipulation of vesicle trafficking components affecting membrane interactions between mitochondria and the ER in EG regulate glia-neuron crosstalk to maintain synaptic integrity of dopaminergic neuron synapses in the fly brain of Pink1 mutants.

DISCUSSION

In this work, we provide evidence for early, non-cell-autonomous activation of EG in a PD-relevant Drosophila Pink1 model. This occurs already in young Pink1-mutant flies when neuronal defects in

dopaminergic neurons are not yet measurable. The EG activation and invasion phenotype appears as a protective response, as *Pink1*-deficient neurons seem to show defects in EG that mimic the response to nerve injury (Doherty et al., 2009; MacDonald et al., 2006). In addition to this, we show that there is also a non-autonomous role of EG to support *Pink1*-deficient neurons. When we manipulate the expression of membrane-lipid trafficking genes, the homologs of the established PD genes VPS13C and VPS35 (Lesage et al., 2016; Vilariño-Güell et al., 2011; Zimprich et al., 2011), specifically in EG, we rescue the dysfunction of *Pink1* mutant neuron defects (ERG and PAM DAN synapse loss). Our work shows the importance of neuron-glia cross-talk in the context of *Pink1*-deficiency. We suggest (1) there is early EG activation secondary to *Pink1*-loss-induced neuronal impairments; (2) there are specific lipid- and membrane trafficking problems caused by *Pink1* loss in EG that center on mitochondria/ER contacts and (3) there is a convergence of Parkinson-relevant genetic factors that act in EG to maintain neuronal function.

The loss of Pink1 function in the context of Parkinson's disease has been amply linked to the regulation of mitochondrial health. Pink1 maintains the integrity of the electron transport chain by phosphorylating NDUFA10 (Morais et al., 2009, 2014; Pogson et al., 2014) and when mitochondria are damaged, it phosphorylates Parkin and Ubiquitin to facilitate mitophagy (Kane et al., 2014; Narendra et al., 2010; Narendra & Youle, 2024). The regulation of mitophagy is complex and requires mitochondrial rearrangements controlled by mitochondria-organelle contacts (ER and lysosomes) (Wong et al., 2018, 2019; Yamano et al., 2018). Such contacts mediate inter-organellar lipid exchange as well as facilitate organellar fusion (Kumar et al., 2018; Valadas et al., 2018a). In *Pink1* mutants, these contacts appear to be increased in number, and this causes cellular defects (Grossmann et al., 2023; Valadas et al., 2018).

We found that in EG, *Vps35* functionally interacts with Pink1-induced neuronal dysfunction: its genetic manipulation in EG rescues cell non-autonomous neuronal dysfunction. Loss of *Vps35* itself only in EG is sufficient to rescue neuronal *Pink1* phenotypes. *Pink1* loss-of-function models show increased numbers of mitochondria-ER contact sites (Valadas et al., 2018), affecting mitochondrial calcium levels (Barazzuol et al., 2020; De Brito & Scorrano, 2008; Ham et al., 2023) and dysregulating lipidic ER composition (Valadas et al., 2018) (Figure 5A). Indeed, Pink1 is necessary for ubiquitination and degradation of Mitofusin (Poole et al., 2010), which is fundamental for the mitochondria-ER tethering (De Brito & Scorrano, 2008). Interestingly, *Vps35* deficiency promotes untethering of mitochondria-ER contact sites by increasing the mitochondrial levels of MUL1, which is necessary for the ubiquitination and degradation of Mitofusin (Puri et al., 2019; Tang et al., 2015; Yun et al., 2014). Hence, conditions that affect mitochondria-ER contact site regulation in EG, by modifying an endosomal trafficking regulator such as *Vps35,* are expected to rescue Pink1 neuronal dysfunction,

possibly by restoring mitochondrial calcium levels and the lipid composition of the ER (Figure 5B); but further work is needed to sort these mechanistic elements in a cell-specific manner.

Our observation that downregulation of *Vps13* in EG also rescues *Pink1* is in further support of a role for ER-mitochondria contact site regulation. *Vps13* has two mammalian homologs, VPS13A and VPS13C (Velayos-Baeza et al., 2004; Vonk et al., 2017; Vrijsen et al., 2022), which are also mutated in familial Parkinsonism/neurodegenerative disease (Lesage et al., 2016). VPS13A tethers ER to mitochondria and lipid droplets, and VPS13C tethers ER to late endosomes, lysosomes and lipid droplets (Kumar et al., 2018; Muñoz-Braceras et al., 2019; Vrijsen et al., 2022). The function of these proteins is to facilitate lipid transfer between organelles, a function we previously showed to be increased in *Pink1* mutants (Valadas et al., 2018). Hence, the downregulation of *Vps13* would counteract the effects of excessive organelle membrane contact formation and dysregulated lipid homeostasis in *Pink1* mutants (Figure 5B).

Lipid homeostasis in glial cells is crucial for supporting neuronal function by providing energy, protecting against oxidative stress, and regulating synaptic health. In oligodendrocytes, maintaining myelin sheaths is critical (Ettle et al., 2016; Lappe-Siefke et al., 2003). While fly EG do not generate such structures, they do wrap processes around neuropil regions, also requiring lipid membrane production (Kremer et al., 2017; Pogodalla et al., 2021). Through lipid metabolism, glia also supply neurons with essential metabolites, cholesterol, and signaling molecules that aid in membrane integrity, synaptic remodeling, and neuroprotection (Delgado et al., 2018; Otto et al., 2018; Saab et al., 2016; Suárez-Pozos et al., 2020). Further work is now required to define which functions, supported by organelle contact sites in glial cells, drive cell-nonautonomous protective mechanisms in neurons. Our work in flies indicates these glial defects precede dopaminergic synapse loss, and, importantly, that rescuing the glial defects helps to prevent dopaminergic problems later in life.

METHODS

Resource availability

Lead Contact

Further information and requests for resources and reagents should be directed to the lead

contact, Patrik Verstreken (patrik.verstreken@kuleuven.be).

Materials Availability

Data, code, Drosophila models, and reagents are available upon request.

Fly stocks

The fruit flies were maintained in an incubator at 25°C under a 12h:12h light-dark cycle and provided with a standard diet consisting of corn meal and molasses. For experiments, only male flies were used. Flies were raised in parallel on the same batch of food, which was exchanged every three to four days and kept in only male population of similar density. Flies were aged to 5 ± 1-day-old, and 22 ± 2-day-old, as indicated for immunohistochemistry and ERGs, respectively.

To create a set of *Drosophila* models for Parkinson's disease, CRISPR/Cas9-based gene editing was utilized, as outlined in (Kaempf et al., 2026; Pech et al., 2025). In brief, each knock-out line contained an attP-flanked w+ reporter cassette that replaced the first shared exon across different isoforms of the targeted gene. For this study, we used knockout flies for *Pink1* from this collection (Kaempf et al., 2026). The "control" strain denotes a semi-isogenized w^{1118} strain that underwent backcrossing to Canton-S for 10 successive generations, resulting in a strain termed Canton-S-w1118. To generate the fly line *yw; UAS-His2Av::eGFP.VK27/TM3Sb*, we created the pUASTattB_His2AV-GFP plasmid, linearizing the pUAST.attB (Bischof et al., 2007) with EcoRI-BamHI. A gBlock, containing the His2AV, GS linker and eGFP sequence, was cloned into this linearized plasmid with Gibson Assembly. The plasmid was inserted into the VK27 landing site by BestGene. The sequence His2AV-GS-eGFP gblock is:

TGAATAGGGAATTGGGcAAacATGGCTGGCGGTAAAGCAGGCAAGGATTCGGGCAAGGCCAAGGCGAAGGC
GGTATCGCGTTCCGCGCGCGCGGGTCTTCAGTTCCCCGTGGGTCGCATCCATCGTCATCTCAAGAGCCGCACTA
CGTCACATGGACGCGTCGGAGCCACTGCAGCCGTGTACTCCGCTGCCATATTGGAATACCTGACCGCCGAGGT
CCTGGAGTTGGCAGGCAACGCATCGAAGGACTTGAAAGTGAAACGTATCACTCCTCGCCACTTACAGCTCGCC
ATTCGCGGAGACGAGGAGCTGGACAGCCTGATCAAGGCAACCATCGCTGGTGGCGGTGTCATTCCGCACATA
CACAAGTCGCTGATCGGCAAAAAGGAGGAAACGGTGCAGGAcCCGCAGCGGAAGGGCAACGTCATTCTGTCG
CAGGCCTACGGTTCAGGCGGAGGTGGCAGCGGCGGTGGCGGATCCATGGTGAGCAAGGGCGAGGAGCTGTT
CACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGA
GGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCGTGCC
CTGGCCCACCCTCGTGACCACCCTGACCTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAG
CACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCA
ACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCG
ACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACAACTACAACAGCCACAACGTCTATATCAT
GGCCGACAAGCAGAAGAACGGCATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCA
GCTCGCCGACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCGACAACCACTACCTG
AGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGACCG
CCGCCGGGATCACTCTCGGCATGGACGAGCTGTACAAATAAGGGTACCTCTAGAGGATCTT.

Flies were backcrossed for five generations into an inbred Canton-S strain harboring the w^{1118} mutation. The genotypes used in this study are listed in the key resource table and Supplementary table 1.

Olfactory Receptor Neurons (ORNs) axotomy

ORN bilateral axotomy was performed on 4±1 day old flies by surgical ablation of the third antennal segment. Briefly, flies were anesthetized with CO_2, and then both antennae were removed with Dumont #5 forceps and returned to the vial (M. Purice, 2020; M. D. Purice et al., 2017). Flies recovered for 24 h on food at 25°C before dissection.

Immunohistochemistry and confocal imaging – invasion phenotype

Immunohistochemistry was performed on adult fly brains of 5±1-day-old, with at least two independent experiments. The brains were dissected in ice-cold PBS and fixed for 20 min in freshly prepared 3.7% paraformaldehyde (in 1x PBS, 0.3% Triton X-100 [PBX] [Sigma]) at room temperature (RT), followed by three 15-min washes in PBX at RT on a shaker. Then, the brains were incubated for 1 h in blocking solution (PBX, 10% normal goat serum [NGS]) at RT. Following blocking, the brains were incubated with primary antibodies (rabbit anti-GFP [Thermo Fisher Scientific], 1:1000, mouse anti-brp [DSHB, nc82], 1:100) in blocking solution at 4°C overnight, followed by three 15-min washes in PBX at RT. Secondary antibodies (goat anti-rabbit Alexa488, goat anti-mouse Alexa555, both at 1:1000 [Thermo Fisher Scientific]) in PBX with 10% NGS were applied overnight at 4°C. Afterwards, the brains were washed three times for 15 min in PBT at RT on a shaker and mounted with the anterior facing up in RapiClear 1.47 (SUNJin Lab). Imaging was performed using a Nikon A1R confocal microscope with a 40x (NA 1.15) water immersion lens; Z-stacks of the entire antennal lobes were acquired. The acquisition was carried out using a Galvano scanner, with a zoom factor of 1, scan speed of 0.5, and line averaging set to 2. All images were captured with a pinhole of 0.9 Airy units and a resolution of 1024 × 1024. Z-stacks (with 0,5 μm step intervals) were obtained for data acquisition, and the same imaging settings were applied across all genotypes and sessions. Image analysis was performed using Fiji (Schindelin et al., 2009). Image analysis was performed using Fiji (Schindelin et al., 2009). Each antennal lobe is composed of multiple glomeruli, and EG separate them and invade them when active (Doherty et al., 2009; Pogodalla et al., 2021). To quantify EG invasion in the antennal lobe, we calculated the invasion in every single glomerulus. To do so and be consistent with the section of the antennal lobe analyzed, anti-brp was used to identify the 3 Z-planes where DM6 and DM1 glomeruli were present (B. Wu et al., 2017). Then, to automatically detect and segment each glomerulus of the antennal lobe section, regions of interest (ROIs) were defined in the SUM projection of the 3 Z-planes by using the Fiji plugin Stardist (Weigert et al., 2019).

To quantify the EG invasion in the ROIs, anti-GFP SUM projection of the same 3 Z-planes was used. The GFP intensity was detected in each ROI, and it was normalized to the area of each ROI (GFP intensity/glomeruli area) to calculate the EG invasion in each glomerulus. Then, the normalized GFP intensity of each glomerulus of an antennal lobe was summed and normalized for the number of glomeruli detected in the antennal lobe. For each experiment, the EG antennal lobe invasion of each fly was normalized to the mean of the control. Representative images show the SUM projections of the 3 Z-planes.

Immunohistochemistry and confocal imaging – TH staining

Immunohistochemistry was performed on adult fly brains of 22 ± 2-day-old, with at least two independent experiments. The brains were dissected in ice-cold PBS and fixed for 20 min in freshly prepared 3.7% paraformaldehyde (in 1x PBS, 0.2% Triton X-100 [PBX]) at room temperature (RT), followed by three 15-min washes in PBX at RT on a shaker. Then, the brains were incubated for 1 h in blocking solution (PBX, 10% normal goat serum [NGS]) at RT. Following blocking, the brains were incubated with primary antibodies (rabbit anti-TH [Sigma], 1:200, mouse anti-DLG [DSHB], 1:100) in blocking solution at 4°C for 1.5 to 2 days, followed by three 15-min washes in PBX at RT. Secondary antibodies (goat anti-rabbit Alexa488, goat anti-mouse Alexa555, both at 1:500 [Thermo Fisher Scientific]) in PBX with 10% NGS were applied overnight at 4°C. Afterwards, the brains were washed three times for 15 min in PBT at RT on a shaker and mounted with the anterior facing up in RapiClear 1.47 (SUNJin Lab). Imaging was performed using a Nikon A1R confocal microscope with a 20x (NA0.95) water immersion lens; Z-stacks of the entire brain were acquired. The acquisition was carried out using a Galvano scanner, with a zoom factor of 1, scan speed of 0.5, and line averaging set to 2. All images were captured with a pinhole of 2.3 Airy units and a resolution of 1024×1024. Z-stacks (with 3 μm step intervals) were obtained for data acquisition, and the same imaging settings were applied across all genotypes and sessions. Image analysis was performed using Fiji (Schindelin et al., 2009). To quantify dopaminergic neuron innervation of the mushroom body (MB), anti-DLG was used to identify the five Z-planes containing the synaptic region of the MB lobes. The ROI for the MB was defined in the sum projection of the five z-planes. To exclude background signal comparable to control, the area of anti-TH fluorescence was thresholded (using the default threshold for all z-planes) within the selected z-stacks. Quantification of the thresholded area within the ROI was performed in each z-plane, then summed and normalized to the MB ROI area for each brain individually (TH+ area/MB area). For each experiment, the individual TH+ area/MB area values were normalized to the mean of the control. Representative images show the maximum projection of five z-planes and the thresholded middle z-plane.

Electroretinograms (ERG)

ERGs were recorded from flies 5±1-day-old as previously described (Heisenberg, 1971; Slabbaert et al., 2016). Flies were immobilized on glass microscope slides using double-sided tape. For recordings, glass electrodes (borosilicate, 1.5 mm outer diameter) filled with 3 M NaCl were placed in the thorax as a reference and on the fly eye for recordings. Each fly was exposed to 5 cycles of 3 s of darkness, followed by a 1-s of light stimuli with LED illumination. Response to the stimuli was recorded using Axoscope 10.7 and analyzed using Clampfit 10.7 software (Molecular Devices). ERG traces were analyzed with Igor Pro 6.37 (Wave Metrics) using a custom-made macro.

Single-cell dissociation for cell-type specific transcriptomics

To test that EG could be isolated from the rest of the brain, we used two cohorts *GMR56F03> UAS-His2Av::eGFP.VK27* and *nSyb-Gal4> UAS-His2Av::eGFP.VK27*. To identify differentially expressed genes in EG, two cohorts were used: the control w^{1118} expressing *GMR56F03> UAS-His2Av::eGFP.VK27* and *Pink1^{KO-WS}* expressing *GMR56F03> UAS-His2Av::eGFP.VK27*. For each genotype, ten brains of flies 5±1-day-old were collected, with most of the laminae removed, from each experimental repeat. The dissections were performed in ice-cold PBS that contained 5 µM Actinomycin D (Sigma) to inhibit changes in gene expression during tissue processing(Y. E. Wu et al., 2017). To prevent any bias caused by different experimental batches, all the genotypes were processed in parallel. To minimize variability across batches, the same reagents and procedures were used throughout the experiment. Dissections were completed within one hour. The dissection order and the assignment of genotypes to dissectors were alternated to reduce any experimenter-related variation.

For the dissociation of brain tissue, a mix of Dispase I (0.6 mg/ml) (Sigma), Collagenase I (30 mg/ml) (Thermo Fisher), Trypsin (0.5x) (Thermo Fisher), and 5 µM Actinomycin D was used. The brains were incubated for 20 min at 25°C with 1000 rpm shaking, performing four triturations at 5-min intervals. After dissociation, the cell suspension was washed with PBS containing 5 µM Actinomycin D and then filtered through a 10 µm strainer (Pluriselect), using 300 µl of PBS EDTA (Sigma) 1µM. DAPI (Sigma) (1:300.000) was added after filtration.

Fluorescence Activated Cell Sorting (FACS) for cell-type specific transcriptomics

Immediately after the dissociation, cells were sorted using a FACSAria Fusion (BD Biosciences) flow cytometer equipped with 4 lasers (405nm, 488nm, 561nm and 640nm) and a 100 µm nozzle at 20 psi. Live (DAPI-negative), GFP-positive cells were sorted into 96-well PCR plates at 300 cells per well (3 wells per genotype) using a 4-way purity mask with the FACSDiva software v9.0.1 (BD Biosciences).

Bulk transcriptomics of EG cells

Library preparations and sequencing for the Bulk RNA seq of the EG cells were performed using a modified Smart-seq2 protocol based on a previously published protocol (Picelli et al., 2013). Briefly, 3 µl of cell lysis buffer (0.1% Triton X-100, Sigma Aldrich; 1U/ul of RNase Inhibitor, RNase OUT, Thermo Fisher; 2.5 µM Oligo dT(25), IDT; and 2.5 mM dNTP, Promega) was aliquoted in triplicate to 96-well plates (4titude, Cat. No. 4TI-0960/C). Roughly 300 cells were sorted into the wells with lysis buffer and the plate was spun down at 2000xg for 1 min prior to storage at -80 °C. For the first strand synthesis, the Smart-Seq 2 plates were thawed to RT for a minute and subsequently spun down at 2000 xg for a minute. The RNA denaturation is carried out at 72°C for 10 min and then flash cooled on ice for 5 min. First strand synthesis mix (1X SuperScript II first strand synthesis buffer, Thermo Fisher; 5 mM DTT, Thermo Fisher; 100 U SuperScript II reverse transcriptase enzyme, Thermo Fisher; 10 U RNAse OUT, Thermo Fisher; 1M Betaine, Sigma Aldrich; 6 mM MgCl2, Thermo Fisher; and 1 uM TSO, IDT Technologies) is added to the cell lysis mix to a total volume of 10 µl. The First strand synthesis reaction was carried out with following program: 42°C for 90 min; 10 cycles of {50°C for 2 min; 42°C for 2 min}; 72°C for 15 min; 4°C indefinite hold. PCR amplification of the first strand product was performed by adding 15 µl of the PCR mix to the first strand product (1X KAPA HiFi HotStart Ready Mix, Roche; and 0.1 µM of IS PCR primer, IDT Technologies). 22 cycles of the following PCR program were used to amplify the first strand product: 98°C for 3 min; 22 cycles of {98°C for 20 sec; 67°C for 15 sec; 72°C for 6 min}; 72°C for 5 min; 4C indefinite hold. 20 µl of Ampure XP was added to each well and mixed. Standard Ampure XP purification was carried out as per manufacturer's recommendation, and the cDNA library was eluted in 17.5 µl of Elution buffer. 17 µl of the eluted library was ,transferred to a fresh 96-well plate (4titude, Cat. No. 4TI-0960/C). 5 µl of Tn5 tagmentation mix (1X Tagment DNA buffer, Illumina; ATM mix, Illumina, and cDNA library 1 ng) was prepared for cDNA fragmentation. Tn5 tagmentation was performed with the following program: 55°C for 10 min; 4°C indefinite hold. Tagmentation reaction was stopped by quenching the reaction with 1.25 µl of NT buffer for 5 min (Illumina). 1.25 µl of the i5 and i7 Illumina indexes were added to the plates. Finally 3.75 µl of NPM master mix was added to the plate and mixed well and put for the index PCR amplification: 72°C for 3 min; 95°C for 30 sec; 12 cycles of {95°C for 10 sec; 55°C for 30 sec; 72°C for 30 sec}; 72°C for 5 min, and 4°C indefinite hold.

The indexed PCR products were pooled in a single tube, and 0.8X Ampure XP purification (Beckman Coulter) was carried out as per the manufacturer's recommendation, and finally, the sequencing library was eluted in 35.5 µl of Elution buffer (Qiagen).

SMART seq 2 libraries were sequenced on the NextSeq 500 (Illumina) sequencing platform. Sequencing was done as per the protocol recommendations: paired-end read of 76 bps (read 1), 76 bps (read 2), 8 bps (index 1) and 8 bps (index 2). For a targeted sequencing depth of ~30 million reads per sample.

Analysis of RNAseq data - Neuron vs EG

Raw reads from neuron- vs. glia-specific FACsorting experiments were processed with the nf-core/rnaseq pipeline (Patel et al., 2020) with default parameters and the BDGP6 reference genome. The STAR-aligned and salmon-quantified expression matrix was then subset to genes known to be enriched in neurons or glia and displayed as a per gene scaled heatmap.

Analysis of RNAseq data - EG-specific differentially expressed genes

After FACsorting of $Pink^{KO-ws}$ vs. control ensheathing glia, library preparation and sequencing resulting FASTQ files were cleaned with fastp (Chen et al., 2018) with default settings. Aligning and counting were carried out with STAR (Dobin et al., 2013) and the 4th 2020 FlyBase *Drosophila melanogaster* release (r6.35) with the –quantMode flag set as GeneCounts. For differential gene expression testing we used DESeq2 (Love et al., 2014), where we fit a negative binomial model and carried out the Wald-test. We removed the experimental batch as a covariate according to the design formula ~date+genotype. All genes below a Benjamini-Hochberg corrected p-value of 0.05 were considered deregulated.

ScRNAseq data loading, filtering, clustering, and differential expression analysis

Raw counts data from (Pech et al., 2025), alongside cell annotations provided by the authors, was loaded into Scanpy (v1.9.1) (Wolf et al., 2018) and subset to only include cells in the young control (<= 6-day-old) or $Pink1^{KO-WS}$ samples. After subsetting the data, filtering was performed to remove any cell with less than 250 genes expressed or a percentage of counts coming from mitochondrial genes greater than 15%, the data was then normalized to a total of 10.000 counts and log transformed. Highly variable genes were detected, total_counts and percent of mitochondrial reads were regressed out, and finally the data was scaled to unit variance with a zero mean, clipped to a max of 10.

After data pre-processing, a PCA was performed, Harmony (Korsunsky et al., 2019) was used to correct batches using "sample_id" as a batch key, and 122 components were used for dimensionality reduction and clustering analyses. Leiden clustering was performed with a resolution of 8.0 and using the annotations from (Pech et al., 2025), all cells within each cluster were labeled by taking the most common, original annotation per cluster.

Per cluster, a differential analysis was performed. First, pseudobulk counts were generated by summing the counts per cell from all cells per sample. These counts were then used in DESeq2 (v1.44.0) (Love et al., 2014) comparing Control vs $Pink1^{KO-WS}$ samples. The total number of genes significantly up- or downregulated per cluster (padj <= 0.05 and an abs (log2foldchange) >= 1.5) was counted and plotted.

Statistical analysis

GraphPad Prism was used for visualization and to determine statistical significance. Datasets were tested for normal distribution using the D'Agostino-Person Omnibus and the Shapiro-Wilk normality test. For a normally distributed dataset ordinary one-way ANOVA was used, followed to correct for multiple comparisons by a post hoc Tukey's test when comparing all the datasets with each other or Dunnett's test when comparing all the datasets to a general control. For non-normally distributed datasets Kruskal-Wallis test is used, followed by a post hoc Dunn's test to correct for multiple comparisons. Significance levels are defined as * is $p<0.05$, ** is $p<0.01$, *** is $p<0.001$, and ns, not significant. Effect size is calculated with eta-squared when one-way ANOVA is used, indicated in the figure legend as η^2. 'N' in the legends is used to indicate how many animals were analyzed. Data are plotted as mean ± SD. Specifics on the statistical test used for each analysis are reported in the figure legends.

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifier	Additional information
Antibody	Rabbit polyclonal anti-GFP	Thermo Fisher Scientific	Cat#A-11122; RRID: AB_221569	
Antibody	Mouse monoclonal anti-Brp	DSHB	Cat#nc82; RRID: AB_2314866	
Antibody	Mouse monoclonal anti-DLG	DSHB	Cat#4F3; RRID: AB 528203	
Antibody	Rabbit polyclonal anti-TH	Sigma Aldrich	Cat#AB 152	
Antibody	Alex Flour 488 goat anti-rabbit	Invitrogen	Cat#A11034	
Antibody	Alexa Fluor 555 goat anti-mouse IgG2a	Invitrogen	Cat#A21137	
Commercial assay, kit	Agencourt AMPure XP	Beckman Coulter	Cat#A63880	
Commercial assay, kit	KAPA HiFi HotStart ReadyMix	Roche	Cat# 07958927001	

Commercial assay, kit	Nextera XT DNA Library Preparation Kit	Illumina	Cat#FC-131-1096	
Peptide, recombinant protein	Dispase I	Sigma Aldrich	Cat#D4818	
Peptide, recombinant protein	Collagenese I	Thermo Fisher Scientific	Cat#17100017	
Peptide, recombinant protein	SuperScript™ II Reverse Transcriptase	Thermo Fisher Scientific	Cat#18064022	
Chemical compound, drug	Trypsin-EDTA (0,5%)	Thermo Fisher Scientific	Cat#15400054	
Chemical compound, drug	Triton X-100 Solution	Sigma Aldrich	Cat#93443-100MI	
Chemical compound, drug	Paraformaldehyde	Sigma Aldrich	Cat#252549	
Chemical compound, drug	Actinomycin D	Sigma Aldrich	Cat#A1410	
Chemical compound, drug	EDTA	Sigma Aldrich	Cat#E6511	
Chemical compound, drug	DAPI	Sigma Aldrich	Cat#D9542	
Chemical compound, drug	$MgCl_2$ (1M)	Thermo Fisher Scientific	Cat#AM9530G	
Chemical compound, drug	RNaseOUT	Thermo Fisher Scientific	Cat#10777019	
Chemical compound, drug	Betaine	Sigma Aldrich	Cat#B0300	
Chemical compound, drug	DTT (100mM Solution)	Thermo Fisher Scientific	Cat#707265ML	
Chemical compound, drug	Buffer EB	Qiagen	Cat#19086	
Chemical compound, drug	RapiClear 1.47	Sunjin Lab	Cat#RC147001	
Sequence-based reagent	dNTP Mix	Promega	Cat#U1511	
Sequnce-based reagent	Primers, gRNAs, oligos, gBlocks	Integrated DNA technologies (IDT)		
Strain, strain backgrou	$w[1118]$ (w^{1118})	(Kaempf et al., 2026)	NA	

nd (D. melanoga ster)			
Strain, strain backgrou nd (D. melanoga ster)	w[1118] M{w+} (w[1118] w+)	(Kaempf et al., 2026)	NA
Gene (D. melanoga ster)	w[1118] TI{w[+]=white-STAR}Pink1[KO-WS]/FM7a (Pink1[KO-WS])	(Kaempf et al., 2026)	NA
Genetic reagent (D. melanoga ster)	Yw; ; UAS-His2Av::eGFP.VK27/TM3S b (UAS-HisTag-eGFP)	This study	NA
Genetic reagent (D. melanoga ster)	;;MZ0709-Gal4	(Ito et al., 1995)	NA
Genetic reagent (D. melanoga ster)	y[1] w[*]; P{w[+mC]=UAS-mCD8::GFP.L}LL5, P{UAS-mCD8::GFP.L}2 (UAS-mCD8-GFP)	(Lee & Luo, 1999)	BDSC_5137
Genetic reagent (D. melanoga ster)	w[*]; P{y[+t7.7] w[+mC]=GMR56F03-GAL4}attP24/CyO (Gmr56-gal4)	(Jenett et al., 2012)	BDSC_77469
Genetic reagent (D. melanoga ster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF01672}att P2 (Pink1[RNAi])	(Perkins et al., 2015)	BDSC_31170
Genetic reagent (D. melanoga ster)	w[*]; P{w[+mC]=UAS-Pink1.C}A (UAS-Pink1)	Blooming ton Drosophi la Stock Center	BDSC_51648
Genetic reagent (D. melanoga ster)	w[1118]; P{y[+t7.7] w[+mC]=GMR57C10-GAL4}attP2 (nSyb-gal4)	(Jenett et al., 2012)	BDSC_39171
Genetic reagent (D. melanoga ster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS02460} attP40 (Vps13[RNAi])	(Perkins et al., 2015)	BDSC_38270
Genetic reagent (D. melanoga ster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01858} attP40 (Vps35[RNAi])	(Perkins et al., 2015)	BDSC 38944
Genetic reagent (D. melanoga ster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS05488} attP40 (CG17660[RNAi])	(Perkins et al., 2015)	BDSC_67022

Genetic reagent (D. melanogaster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF03249}attP2 (Proc[RNAi])	(Perkins et al., 2015)	BDSC_29570	
Genetic reagent (D. melanogaster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC05229} attP40 (fiz[RNAi])	(Perkins et al., 2015)	BDSC_62222	
Genetic reagent (D. melanogaster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF01165}attP2 (CG15011[RNAi])	(Perkins et al., 2015)	BDSC_31589	
Genetic reagent (D. melanogaster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC04857} attP40 (CG17612[RNAi])	(Perkins et al., 2015)	BDSC_57540	
Genetic reagent (D. melanogaster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC05717} attP40 (Prpk[RNAi])	(Perkins et al., 2015)	BDSC 64844	
Genetic reagent (D. melanogaster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMJ22434}attP40 (Lnpk[RNAi])	(Perkins et al., 2015)	BDSC_64036	
Genetic reagent (D. melanogaster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01827} attP2 (PIG-B[RNAi])	(Perkins et al., 2015)	BDSC_38359	
Genetic reagent (D. melanogaster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01840} attP2/TM3, Sb[1] (spag[RNAi])	(Perkins et al., 2015)	BDSC_38371	
Genetic reagent (D. melanogaster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS00070} attP2 (Rel[RNAi])	(Perkins et al., 2015)	BDSC_33661	
Genetic reagent (D. melanogaster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF03249}attP2 (Proc[RNAi])	Bloomington Drosophila Stock Center	BDSC 29570	
Genetic reagent (D. melanogaster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC04640} attP40 (Atac1[RNAi])	(Perkins et al., 2015)	BDSC_57250	
Genetic reagent (D. melanogaster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.GLV21056}attP2 (l(3)07882[RNAi])	(Perkins et al., 2015)	BDSC_35691	
Genetic reagent (D.	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS00529}	(Perkins et al., 2015)	BDSC_32839	

melanoga ster)	attP2 (Pld[RNAi])		
Genetic reagent (D. melanoga ster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS02243} attP2 (Rpp25[RNAi])	(Perkins et al., 2015)	BDSC_41679
Genetic reagent (D. melanoga ster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03184} attP40 (GMF[RNAi])	(Perkins et al., 2015)	BDSC_51452
Genetic reagent (D. melanoga ster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01358} attP2/TM3, Sb[1] (Atg7[RNAi])	(Perkins et al., 2015)	BDSC_34369
Genetic reagent (D. melanoga ster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMJ03126}a ttP40 (Ada2d[RNAi])	(Perkins et al., 2015)	BDSC_50905
Genetic reagent (D. melanoga ster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC06224} attP2 (CG12773[RNAi])	(Perkins et al., 2015)	BDSC_65949
Genetic reagent (D. melanoga ster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS00328} attP2 (CG7627[RNAi])	(Perkins et al., 2015)	BDSC_32337
Genetic reagent (D. melanoga ster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMS00870} attP2 (Dark[RNAi])	(Perkins et al., 2015)	BDSC_33924
Genetic reagent (D. melanoga ster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF03275}att P2 (CG3703[RNAi])	(Perkins et al., 2015)	BDSC_29596
Genetic reagent (D. melanoga ster)	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01175} attP2/TM3, Sb[1] (NiPp1[RNAi])	(Perkins et al., 2015)	BDSC_34696
Genetic reagent (D. melanoga ster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMJ22813}a ttP40 (CG31370[RNAi])	(Perkins et al., 2015)	BDSC_60456
Genetic reagent (D. melanoga ster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMJ23526}a ttP40 (Irbp[RNAi])	(Perkins et al., 2015)	BDSC_61942
Genetic reagent (D. melanoga ster)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF02313}att P2 (CG32532[RNAi])	Click or tap here to enter text.	BDSC_26750
Genetic	y[1] v[1]; P{y[+t7.7]	(Perkins	BDSC 62936

reagent (*D. melanoga ster*)	*v[+t1.8]=TRiP.HMJ30013}a ttP40/CyO (Rhau[RNAi])*	et al., 2015)	
Genetic reagent (*D. melanoga ster*)	*y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HM05047}at tP2 (hfw[RNAi])*	(Perkins et al., 2015)	BDSC_28561
Genetic reagent (*D. melanoga ster*)	*y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF02192}att P2 (achi[RNAi])*	(Perkins et al., 2015)	BDSC_31903
Genetic reagent (*D. melanoga ster*)	*y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMJ23972}a ttP40/CyO (TTLL6A[RNAi])*	(Perkins et al., 2015)	BDSC_62488
Genetic reagent (*D. melanoga ster*)	*y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03076} attP2 (Nos[RNAi])*	(Perkins et al., 2015)	BDSC_50675
Genetic reagent (*D. melanoga ster*)	*w[1118]; P{w[+mC]=UAS-Nos.L}2 (UAS-Nos)*	Blooming ton Drosophi la Stock Center	BDSC_56823
Plasmids	pUASTattB	(Bischof, 2007)	https://www.fly c31.org
Software, algorithm	Fiji	(Schindel in et al., 2009)	RRID:SCR_00228 5	https://imagej.net/Fiji
Software, algorithm	GraphPad Prism	GraphPa d software, CA, USA	RRID:SCR_00279 8	https://www.graphpad.com/scientific-software/prism/
Software, algorithm	Python (v 3.7.3)	Python	RRID:SCR_00839 4	http://www.python.org/
Software, algorithm	scanpy	(Wolf et al., 2018)	RRID:SCR_01813 9	https://github.com/theislab/scanpy
Software, algorithm	DESeq2	(Love et al., 2014)	RRID:SCR_01568 7	https://bioconductor.org/packages/release/bioc /html/DESeq2.html
Software, algorithm	Clampfit	Molecula r Devices		https://www.moleculardevices.com
Software, algorithm	Axoscope	Molecula r Devices		https://www.moleculardevices.com
Software, algorithm	Igor Pro	WaveMe trics	RRID: CR_000325	https://www.wavemetrics.com/products/ igorpro/igorpro.htm
Software, algorithm	FACSDiva software v9.0.1	BD Bioscienc es	RRID:SCR_00145 6	https://www.bdbiosciences.com/en-be/products/software/instrument-software/bd-facsdiva-software
Software, algorithm	nf-core/rnaseq	(Patel et al., 2020)		https://nf-co.re/rnaseq/3.14.0/
Software, algorithm	fastp	(Chen et al., 2018)		https://github.com/OpenGene/fastp
Software, algorithm	R Studio	RStudio, PBC / Posit		
Software, algorithm	BioRender	BioRend er		https://BioRender.com

Software, algorithm	STAR	(Dobin et al., 2013)	RRID:SCR_00446 3	https://github.com/alexdobin/STAR
Software, algorithm	Harmony	(Korsunsky et al., 2019)		https://github.com/immunogenomics/harmony

Acknowledgments

We thank the Vlaams Supercomputer Centrum, the VIB Nucleomic and Bioimaging Cores, and the Bloomington Drosophila Stock Center (NIH P40OD018537). We are grateful to the members of the Verstreken lab for valuable discussions. Research support was provided by Leuven University Fund and Opening the Future, ERC, the Chan Zuckerberg Initiative, a Methusalem grant from the Flemish government, KU Leuven BOF, Fund Jacqueline Cigrang administered by the KBS, the KBS, fund Generet and FWO Vlaanderen to PV an EMBO long-term fellowship to RP, a DFG fellowship to UP and an FWO PhD mandate to AK. Cartoons/models/illustrations were created in https://BioRender.com. P.V. is an alumnus of the FENS-Kavli Network of Excellence.

Author contributions

Conceptualization, L.G., R.P., P.V.;

Methodology, L.G., U.P., N.S., J.L., K.D., R.P., P.V., S.K., A.K.;

Software, L.G., K.D., R.P.;

Validation, L.G., K.D., R.P.;

Formal analysis, L.G., K.D., R.P.;

Investigation, L.G., U.P., N.S., J.L., K.D., R.P., S.K., A.K.;

Writing -Original Draft, L.G., R.P., P.V.;

Writing – review and editing, all co-authors read and edited the manuscript.

Visualization, L.G., K.D., R.P., S.K.;

Supervision, R.P., P.V.;

Project administration, L.G.;

Funding acquisition, R.P., P.V.;

Bibliography

Agarwal, D., Sandor, C., Volpato, V., Caffrey, T. M., Monzón-Sandoval, J., Bowden, R., Alegre-Abarrategui, J., Wade-Martins, R., & Webber, C. (2020). A single-cell atlas of the human substantia nigra reveals cell-specific pathways associated with neurological disorders. *Nature Communications*, *11*(1), 1–11. https://doi.org/10.1038/s41467-020-17876-0

Barazzuol, L., Giamogante, F., Brini, M., & Calì, T. (2020). PINK1/Parkin Mediated Mitophagy, Ca2+ Signalling, and ER-Mitochondria Contacts in Parkinson's Disease. *International Journal of Molecular Sciences*, *21*(5). https://doi.org/10.3390/ijms21051772

Bischof, J., Maeda, R. K., Hediger, M., Karch, F., & Basler, K. (2007). An optimized transgenesis system for Drosophila using germ-line-specific φC31 integrases. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(9), 3312–3317. https://doi.org/10.1073/pnas.0611511104

Brickner, J. H., & Fuller, R. S. (1997). SOI1 encodes a novel, conserved protein that promotes TGN-endosomal cycling of Kex2p and other membrane proteins by modulating the function of two TGN localization signals. *Journal of Cell Biology*, *139*(1), 23–36. https://doi.org/10.1083/jcb.139.1.23

Bryois, J., Skene, N. G., Hansen, T. F., Kogelman, L. J. A., Watson, H. J., Liu, Z., Adan, R., Alfredsson, L., Ando, T., Andreassen, O., Baker, J., Bergen, A., Berrettini, W., Birgegård, A., Boden, J., Boehm, I., Boni, C., Boraska Perica, V., Brandt, H., … Sullivan, P. F. (2020). Genetic identification of cell types underlying brain complex traits yields insights into the etiology of Parkinson's disease. *Nature Genetics*, *52*(5), 482–493. https://doi.org/10.1038/s41588-020-0610-9

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). Fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, *34*(17), i884–i890. https://doi.org/10.1093/bioinformatics/bty560

De Brito, O. M., & Scorrano, L. (2008). Mitofusin 2 tethers endoplasmic reticulum to mitochondria. *Nature*, *456*(4), 605–610. https://doi.org/10.1038/nature07534

Delgado, M. G., Oliva, C., López, E., Ibacache, A., Galaz, A., Delgado, R., Barros, L. F., & Sierralta, J. (2018). Chaski, a novel Drosophila lactate/pyruvate transporter required in glia cells for survival under nutritional stress. *Scientific Reports*, *8*(1), 1–13. https://doi.org/10.1038/s41598-018-19595-5

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21. https://doi.org/10.1093/bioinformatics/bts635

Doherty, J., Logan, M. A., Taşdemir, Ö. E., & Freeman, M. R. (2009a). Ensheathing Glia Function as Phagocytes in the Adult Drosophila Brain. *The Journal of Neuroscience*, *29*(15), 4768–4781. https://doi.org/10.1523/JNEUROSCI.5951-08.2009

Doherty, J., Logan, M. A., Taşdemir, Ö. E., & Freeman, M. R. (2009b). Ensheathing Glia Function as Phagocytes in the Adult Drosophila Brain. *The Journal of Neuroscience*, *29*(15), 4768–4781. https://doi.org/10.1523/JNEUROSCI.5951-08.2009

Ettle, B., Schlachetzki, J. C. M., & Winkler, J. (2016). Oligodendroglia and Myelin in Neurodegenerative Diseases: More Than Just Bystanders? *Molecular Neurobiology*, *53*(5), 3046–3062. https://doi.org/10.1007/s12035-015-9205-3

Freeman, M. R., & Doherty, J. (2006). Glial cell biology in Drosophila and vertebrates. In *Trends in Neurosciences* (Vol. 29, Issue 2, pp. 82–90). Elsevier Current Trends. https://doi.org/10.1016/j.tins.2005.12.002

Gaudet, P., Livstone, M. S., Lewis, S. E., & Thomas, P. D. (2011). Phylogenetic-based propagation of functional annotations within the Gene Ontology consortium. *Briefings in Bioinformatics*, *12*(5), 449–462. https://doi.org/10.1093/bib/bbr042

Grossmann, D., Malburg, N., Glaß, H., Weeren, V., Sondermann, V., Pfeiffer, J. F., Petters, J., Lukas, J., Seibler, P., Klein, C., Grünewald, A., & Hermann, A. (2023). Mitochondria–Endoplasmic Reticulum Contact Sites Dynamics and Calcium Homeostasis Are Differentially Disrupted in PINK1-PD or PRKN-PD Neurons. *Movement Disorders*, *38*(10), 1822–1836. https://doi.org/10.1002/mds.29525

Ham, S. J., Yoo, H., Woo, D., Lee, D. H., Park, K. S., & Chung, J. (2023). PINK1 and Parkin regulate IP3R-mediated ER calcium release. *Nature Communications*, *14*(1). https://doi.org/10.1038/s41467-023-40929-z

Hardie, R. C., & Raghu, P. (2001). Visual transduction in Drosophila. *Nature*, *413*(6852), 186–193.

Heisenberg, M. (1971). Separation of receptor and lamina potentials in the electroretinogram of normal and mutant Drosophila. *Journal of Experimental Biology*, *55*(1), 85–100. https://doi.org/10.1242/jeb.55.1.85

Hirata, T., Fujita, M., Nakamura, S., Gotoh, K., Motooka, D., Murakami, Y., Maeda, Y., & Kinoshita, T. (2015). Post-Golgi anterograde transport requires GARP-dependent endosome-to-TGN retrograde transport. *Molecular Biology of the Cell*, *26*(17), 3071–3084. https://doi.org/10.1091/mbc.E14-11-1568

Ito, K., Urban, J., & Technau, G. M. (1995). Distribution, classification, and development of Drosophila glial cells in the late embryonic and early larval ventral nerve cord. *Roux's Archives of Developmental Biology*, *204*(5), 284–307. https://doi.org/10.1007/BF02179499

Jenett, A., Rubin, G. M., Ngo, T.-T. B., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B. D., Cavallaro, A., Hall, D., Jeter, J., Iyer, N., Fetter, D., Hausenfluck, J. H., Peng, H., Trautman, E. T., Svirskas, R., Myers, E. W., Iwinski, Z. R., Aso, Y., … Zugates, C. T. (2012). A GAL4-Driver Line Resource for Drosophila Neurobiology. *Cell Reports*, *2*(4), 991–1001. https://doi.org/10.1016/j.celrep.2012.09.011

Kaempf, N., Valadas, J. S., Robberechts, P., Schoovaerts, N., Ortega, A., Nachman, E., Ghezzi, L., Kilic, A., Chabot, D., Pech, U., Kuenen, S., Vilain, S., Baz, E.-S., Davis, J., Liu, S., & Verstreken, P. (2026). Behavioral screening defines the molecular Parkinsonism subgroups in Drosophila. *Nature Communications, in print*.

Kane, L. A., Lazarou, M., Fogel, A. I., Li, Y., Yamano, K., Sarraf, S. A., Banerjee, S., & Youle, R. J. (2014). PINK1 phosphorylates ubiquitin to activate Parkin E3 ubiquitin ligase activity. *Journal of Cell Biology*, *205*(2), 143–153. https://doi.org/10.1083/jcb.201402104

Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner, M., Loh, P. ru, & Raychaudhuri, S. (2019). Fast, sensitive and accurate integration of single-cell data with Harmony. *Nature Methods*, *16*(12), 1289–1296. https://doi.org/10.1038/s41592-019-0619-0

Kremer, M. C., Jung, C., Batelli, S., Rubin, G. M., & Gaul, U. (2017). The glia of the adult Drosophila nervous system. *Glia*, *65*(4), 606–638. https://doi.org/10.1002/glia.23115

Kumar, N., Leonzino, M., Hancock-Cerutti, W., Horenkamp, F. A., Li, P., Lees, J. A., Wheeler, H., Reinisch, K. M., & De Camilli, P. (2018). VPS13A and VPS13C are lipid transport proteins differentially localized at ER contact sites. *Journal of Cell Biology*, *217*(10), 3625–3639. https://doi.org/10.1083/jcb.201807019

Lang, A. E., & Lozano, A. M. (1998). Parkinson's Disease. *New England Journal of Medicine*, *339*(15), 1044–1053. https://doi.org/10.1056/NEJM199810083391506

Lappe-Siefke, C., Goebbels, S., Gravel, M., Nicksch, E., Lee, J., Braun, P. E., Griffiths, I. R., & Nave, K.-A. (2003). Disruption of Cnp1 uncouples oligodendroglial functions in axonal support and myelination. *Nature Genetics*, *33*(3), 366–374. https://doi.org/10.1038/ng1095

Lee, T., & Luo, L. (1999). Mosaic Analysis with a Repressible Neurotechnique Cell Marker for Studies of Gene Function in Neuronal Morphogenesis. *Neuron*, *22*, 451–461. https://doi.org/10.1016/s0896-6273(00)80701-1

Lesage, S., Drouet, V., Majounie, E., Deramecourt, V., Jacoupy, M., Nicolas, A., Cormier-Dequaire, F., Hassoun, S. M., Pujol, C., Ciura, S., Erpapazoglou, Z., Usenko, T., Maurage, C. A., Sahbatou, M., Liebau, S., Ding, J., Bilgic, B., Emre, M., Erginel-Unaltuna, N., … Brice, A. (2016a). Loss of VPS13C Function in Autosomal-Recessive Parkinsonism Causes Mitochondrial Dysfunction and Increases PINK1/Parkin-Dependent Mitophagy. *American Journal of Human Genetics*, *98*(3), 500–513. https://doi.org/10.1016/j.ajhg.2016.01.014

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 1–21. https://doi.org/10.1186/s13059-014-0550-8

MacDonald, J. M., Beach, M. G., Porpiglia, E., Sheehan, A. E., Watts, R. J., & Freeman, M. R. (2006). The Drosophila Cell Corpse Engulfment Receptor Draper Mediates Glial Clearance of Severed Axons. *Neuron*, *50*(6), 869–881. https://doi.org/10.1016/j.neuron.2006.04.028

Morais, V. A., Haddad, D., Craessaerts, K., De Bock, P.-J., Swerts, J., Vilain, S., Aerts, L., Overbergh, L., Grünewald, A., Seibler, P., Klein, C., Gevaert, K., Verstreken, P., & De Strooper, B. (2014). PINK1 Loss-of-Function Mutations Affect Mitochondrial Complex I Activity via NdufA10 Ubiquinone Uncoupling. *Science*, *344*(6180), 203–207. https://doi.org/10.1126/science.1249161

Morais, V. A., Verstreken, P., Roethig, A., Smet, J., Snellinx, A., Vanbrabant, M., Haddad, D., Frezza, C., Mandemakers, W., Vogt-Weisenhorn, D., Van Coster, R., Wurst, W., Scorrano, L., & De Strooper, B. (2009). Parkinson's disease mutations in PINK1 result in decreased Complex I activity and deficient synaptic function. *EMBO Molecular Medicine*, *1*(2), 99–111. https://doi.org/10.1002/emmm.200900006

Munhoz, R. P., Moro, A., Silveira-moriyama, L., & Teive, H. A. (2015). Non-motor signs in Parkinson's disease: a review. *Arq. Neuropsiquiatr.*, *January*, 454–462. https://doi.org/10.1590/0004-282X20150029

Muñoz-Braceras, S., Tornero-Écija, A. R., Vincent, O., & Escalante, R. (2019). VPS13A, a closely associated mitochondrial protein, is required for efficient lysosomal degradation. *Disease Models & Mechanisms*, *12*(2). https://doi.org/10.1242/dmm.036681

Narendra, D. P., Jin, S. M., Tanaka, A., Suen, D.-F., Gautier, C. A., Shen, J., Cookson, M. R., & Youle, R. J. (2010). PINK1 Is Selectively Stabilized on Impaired Mitochondria to Activate Parkin. *PLoS Biology*, *8*(1), e1000298. https://doi.org/10.1371/journal.pbio.1000298

Narendra, D. P., & Youle, R. J. (2024). The role of PINK1–Parkin in mitochondrial quality control. *Nature Cell Biology*, *26*(10), 1639–1651. https://doi.org/10.1038/s41556-024-01513-9

Otto, N., Marelja, Z., Schoofs, A., Kranenburg, H., Bittern, J., Yildirim, K., Berh, D., Bethke, M., Thomas, S., Rode, S., Risse, B., Jiang, X., Pankratz, M., Leimkühler, S., & Klämbt, C. (2018). The sulfite oxidase Shopper controls neuronal activity by regulating glutamate homeostasis in Drosophila ensheathing glia. *Nature Communications*, *9*(1), 3514. https://doi.org/10.1038/s41467-018-05645-z

Patel, H., Ewels, P., Manning, J., Garcia, M. U., Peltzer, A., Hammarén, R., Botvinnik, O., Talbot, A., Sturm, G., Bot, N., Zepper, M., Moreno, D., Vemuri, P., Binzer-Panchal, M., Greenberg, E., Silviamorins, Pantano, L., Syme, R., Kelly, G., … Gabernet, G. (2020). nf-core/rnaseq: nf-core/rnaseq. In *Zenodo*. https://doi.org/10.5281/zenodo.1400710.

Pech, U., Janssens, J., Schoovaerts, N., Kuenen, S., Aristoy, C. C., Gallego, S. F., Makhzami, S., Hulselmans, G., Poovathingal, S., Davie, K., Bademosi, A. T., Swerts, J., Vilain, S., Aerts, S., & Verstreken, P. (2025). Synaptic deregulation of cholinergic projection neurons causes olfactory dysfunction across 5 fly Parkinsonism models. *ELife*. https://doi.org/10.7554/eLife.98348.3

Perkins, L. A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., Yang-Zhou, D., Flockhart, I., Binari, R., Shim, H. S., Miller, A., Housden, A., Foos, M., Randkelv, S., Kelley, C., Namgyal, P., Villalta, C., Liu, L. P., Jiang, X., … Perrimon, N. (2015). The transgenic RNAi project at Harvard medical school: Resources and validation. *Genetics*, *201*(3), 843–852. https://doi.org/10.1534/genetics.115.180208

Picelli, S., Björklund, Å. K., Faridani, O. R., Sagasser, S., Winberg, G., & Sandberg, R. (2013). Smart-seq2 for sensitive full-length transcriptome profiling in single cells. *Nature Methods*, *10*(11), 1096–1100. https://doi.org/10.1038/nmeth.2639

Pogodalla, N., Kranenburg, H., Rey, S., Rodrigues, S., Cardona, A., & Klämbt, C. (2021). Drosophila ßHeavy-Spectrin is required in polarized ensheathing glia that form a diffusion-barrier around the neuropil. *Nature Communications*, *12*(1), 6357. https://doi.org/10.1038/s41467-021-26462-x

Pogson, J. H., Ivatt, R. M., Sanchez-Martinez, A., Tufi, R., Wilson, E., Mortiboys, H., & Whitworth, A. J. (2014). The Complex I Subunit NDUFA10 Selectively Rescues Drosophila pink1 Mutants through a Mechanism Independent of Mitophagy. *PLoS Genetics*, *10*(11), e1004815. https://doi.org/10.1371/journal.pgen.1004815

Poole, A. C., Thomas, R. E., Yu, S., Vincow, E. S., & Pallanck, L. (2010). The mitochondrial fusion-promoting factor mitofusin is a substrate of the PINK1/parkin pathway. *PLoS ONE*, *5*(4). https://doi.org/10.1371/journal.pone.0010054

Praschberger, R., Kuenen, S., Schoovaerts, N., Kaempf, N., Singh, J., Janssens, J., Swerts, J., Nachman, E., Calatayud, C., Aerts, S., Poovathingal, S., & Verstreken, P. (2023). Neuronal identity defines α-synuclein and tau toxicity. *Neuron*, *111*(10), 1577-1590.e11. https://doi.org/10.1016/j.neuron.2023.02.033

Puri, R., Cheng, X. T., Lin, M. Y., Huang, N., & Sheng, Z. H. (2019). Mul1 restrains Parkin-mediated mitophagy in mature neurons by maintaining ER-mitochondrial contacts. *Nature Communications*, *10*(1). https://doi.org/10.1038/s41467-019-11636-5

Purice, M. (2020). Olfactory receptor neuron (ORN) axotomy assay. *Bio-Protocol Preprint*. https://doi.org/bio-protocol.org/prep625

Purice, M. D., Ray, A., Münzel, E. J., Pope, B. J., Park, D. J., Speese, S. D., & Logan, M. A. (2017). A novel Drosophila injury model reveals severed axons are cleared through a draper/MMP-1 signaling cascade. *ELife*, *6*, 1–30. https://doi.org/10.7554/eLife.23611

Saab, A. S., Tzvetavona, I. D., Trevisiol, A., Baltan, S., Dibaj, P., Kusch, K., Möbius, W., Goetze, B., Jahn, H. M., Huang, W., Steffens, H., Schomburg, E. D., Pérez-Samartín, A., Pérez-Cerdá, F., Bakhtiari, D., Matute, C., Löwel, S., Griesinger, C., Hirrlinger, J., … Nave, K. A. (2016). Oligodendroglial NMDA Receptors Regulate Glucose Import and Axonal Energy Metabolism. *Neuron*, *91*(1), 119–132. https://doi.org/10.1016/j.neuron.2016.05.016

Schindelin, J., Arganda-Carrera, I., Frise, E., Verena, K., Mark, L., Tobias, P., Stephan, P., Curtis, R., Stephan, S., Benjamin, S., Jean-Yves, T., Daniel, J. W., Volker, H., Kevin, E., Pavel, T., & Albert, C. (2009). Fiji - an Open platform for biological image analysis. *Nature Methods*, *9*(7). https://doi.org/10.1038/nmeth.2019.Fiji

Slabbaert, J. R., Kuenen, S., Swerts, J., Maes, I., Uytterhoeven, V., Kasprowicz, J., Fernandes, A. C., Blust, R., & Verstreken, P. (2016). Shawn, the Drosophila homolog of SLC25A39/40, is a mitochondrial carrier that promotes neuronal survival. *Journal of Neuroscience*, *36*(6), 1914–1929. https://doi.org/10.1523/JNEUROSCI.3432-15.2016

Smajic, S., Prada-Medina, C. A., Landoulsi, Z., Ghelfi, J., Delcambre, S., Dietrich, C., Jarazo, J., Henck, J., Balachandran, S., Pachchek, S., Morris, C. M., Antony, P., Timmermann, B., Sauer, S., Pereira, S. L., Schwamborn, J. C., May, P., Grunewald, A., & Spielmann, M. (2022). Single-cell sequencing of human midbrain reveals glial activation and a Parkinson-specific neuronal state. *Brain*, *145*(3), 964–978. https://doi.org/10.1093/brain/awab446

Soukup, S., Kuenen, S., Vanhauwaert, R., Geens, A., Strooper, B. De, & Verstreken, P. (2016). *A LRRK2-Dependent EndophilinA Phosphoswitch Is Critical for Macroautophagy at Presynaptic Article A LRRK2-Dependent EndophilinA Phosphoswitch Is Critical for Macroautophagy at Presynaptic Terminals*. 829–844. https://doi.org/10.1016/j.neuron.2016.09.037

Suárez-Pozos, E., Thomason, E. J., & Fuss, B. (2020). Glutamate Transporters: Expression and Function in Oligodendrocytes. *Neurochemical Research*, *45*(3), 551–560. https://doi.org/10.1007/s11064-018-02708-x

Tang, F. L., Liu, W., Hu, J. X., Erion, J. R., Ye, J., Mei, L., & Xiong, W. C. (2015). VPS35 Deficiency or Mutation Causes Dopaminergic Neuronal Loss by Impairing Mitochondrial Fusion and Function. *Cell Reports*, *12*(10), 1631–1643. https://doi.org/10.1016/j.celrep.2015.08.001

Valadas, J. S., Esposito, G., Vandekerkhove, D., Miskiewicz, K., Deaulmerie, L., Raitano, S., Seibler, P., Klein, C., & Verstreken, P. (2018a). ER Lipid Defects in Neuropeptidergic Neurons Impair Sleep Patterns in Parkinson's Disease. *Neuron*, *98*(6), 1155-1169.e6. https://doi.org/10.1016/j.neuron.2018.05.022

Velayos-Baeza, A., Vettori, A., Copley, R. R., Dobson-Stone, C., & Monaco, A. P. (2004). Analysis of the human VPS13 gene family. *Genomics*, *84*(3), 536–549. https://doi.org/10.1016/j.ygeno.2004.04.012

Vilariño-Güell, C., Wider, C., Ross, O. A., Dachsel, J. C., Kachergus, J. M., Lincoln, S. J., Soto-Ortolaza, A. I., Cobb, S. A., Wilhoite, G. J., Bacon, J. A., Bahareh Behrouz, Melrose, H. L., Hentati, E., Puschmann, A., Evans, D. M., Conibear, E., Wasserman, W. W., Aasly, J. O., Burkhard, P. R., … Farrer, M. J. (2011). VPS35 mutations in parkinson disease. *American Journal of Human Genetics*, *89*(1), 162–167. https://doi.org/10.1016/j.ajhg.2011.06.001

Vilinsky, I., & Johnson, K. G. (2012). Electroretinograms in Drosophila: A Robust and Genetically Accessible Electrophysiological System for the Undergraduate Laboratory. *The Journal of Undergraduate Neuroscience Education*, *11*(1), 149–157. http://fly.bio.indiana.edu/

Vonk, J. J., Yeshaw, W. M., Pinto, F., Faber, A. I. E., Lahaye, L. L., Kanon, B., van der Zwaag, M., Velayos-Baeza, A., Freire, R., van IJzendoorn, S. C., Grzeschik, N. A., & Sibon, O. C. M. (2017). Drosophila Vps13 Is Required for Protein Homeostasis in the Brain. *PLOS ONE*, *12*(1), e0170106. https://doi.org/10.1371/journal.pone.0170106

Vrijsen, S., Vrancx, C., Del Vecchio, M., Swinnen, J. V., Agostinis, P., Winderickx, J., Vangheluwe, P., & Annaert, W. (2022). Inter-organellar Communication in Parkinson's and Alzheimer's Disease: Looking Beyond Endoplasmic Reticulum-Mitochondria Contact Sites. *Frontiers in Neuroscience*, *16*. https://doi.org/10.3389/fnins.2022.900338

Weigert, M., Schmidt, U., Haase, R., Sugawara, K., & Myers, G. (2019). *Star-convex Polyhedra for 3D Object Detection and Segmentation in Microscopy*. http://arxiv.org/abs/1908.03636

Wolf, F. A., Angerer, P., & Theis, F. J. (2018). SCANPY: large-scale single-cell gene expression data analysis. *Genome Biology*, *19*(1), 15. https://doi.org/10.1186/s13059-017-1382-0

Wong, Y. C., Peng, W., & Krainc, D. (2019). Lysosomal Regulation of Inter-mitochondrial Contact Fate and Motility in Charcot-Marie-Tooth Type 2. *Developmental Cell*, *50*(3), 339-354.e4. https://doi.org/10.1016/j.devcel.2019.05.033

Wong, Y. C., Ysselstein, D., & Krainc, D. (2018). Mitochondria–lysosome contacts regulate mitochondrial fission via RAB7 GTP hydrolysis. *Nature*, *554*(7692), 382–386. https://doi.org/10.1038/nature25486

Wu, B., Li, J., Chou, Y.-H., Luginbuhl, D., & Luo, L. (2017). Fibroblast growth factor signaling instructs ensheathing glia wrapping of Drosophila olfactory glomeruli. *Proceedings of the National Academy of Sciences*, *114*(29), 7505–7512. https://doi.org/10.1073/PNAS.1706533114

Wu, C. F., & Wong, F. (1977). Frequency characteristics in the visual system of Drosophila: Genetic dissection of electroretinogram components. *Journal of General Physiology*, *69*(6), 705–724. https://doi.org/10.1085/jgp.69.6.705

Wu, Y. E., Pan, L., Zuo, Y., Li, X., & Hong, W. (2017). Detecting Activated Cell Populations Using Single-Cell RNA-Seq. *Neuron*, *96*(2), 313-329.e6. https://doi.org/10.1016/j.neuron.2017.09.026

Yamano, K., Wang, C., Sarraf, S. A., Münch, C., Kikuchi, R., Noda, N. N., Hizukuri, Y., Kanemaki, M. T., Harper, W., Tanaka, K., Matsuda, N., & Youle, R. J. (2018). Endosomal Rab cycles regulate Parkin-mediated mitophagy. *ELife*, *7*, 1–32. https://doi.org/10.7554/eLife.31326

Yildirim, K., Petri, J., Kottmeier, R., & Klämbt, C. (2019). Drosophila glia: Few cell types and many conserved functions. *Glia*, *67*(1), 5–26. https://doi.org/10.1002/glia.23459

Yun, J., Puri, R., Yang, H., Lizzio, M. A., Wu, C., Sheng, Z.-H., & Guo, M. (2014). MUL1 acts in parallel to the PINK1/parkin pathway in regulating mitofusin and compensates for loss of PINK1/parkin. *ELife*, *3*. https://doi.org/10.7554/elife.01958

Zimprich, A., Benet-Pagès, A., Struhal, W., Graf, E., Eck, S. H., Offman, M. N., Haubenberger, D., Spielberger, S., Schulte, E. C., Lichtner, P., Rossle, S. C., Klopp, N., Wolf, E., Seppi, K., Pirker, W., Presslauer, S., Mollenhauer, B., Katzenschlager, R., Foki, T., … Strom, T. M. (2011). A mutation in VPS35, encoding a subunit of the retromer complex, causes late-onset parkinson disease. *American Journal of Human Genetics*, *89*(1), 168–175. https://doi.org/10.1016/j.ajhg.2011.06.008

Figure 1. EG are affected non-cell autonomously by *Pink1* loss-of-function (with supplementary fig 1)

(**A**) tSNE of the cells of *Pink1^{P399L}* knock-in mutants (5-day-old). Cell types are labeled with colors indicating the number of deregulated genes compared to control. EG are encircled and labeled.

(**B-B"**) Maximum intensity projections of confocal images of fly brains (5 ± 1-day-old) stained with anti-GFP (Green) and anti-Brp (Magenta), where anti-GFP marks EG and anti-Brp marks presynaptic sites of the antennal lobes in flies where CD8GFP is expressed via the EG driver MZ709-Gal4. Scale bar: 20 µm (**B'**) Maximum intensity projection of confocal images of controls vs. controls 24 hours after ORN-severing (injury). (**B"**) Maximum intensity projection of confocal images of *Pink1^{KO-WS}* vs *Pink1^{KO-WS}* 24 hours after ORNs severing (injury).

(**C**) Quantification of GFP intensity within the glomeruli of the antennal lobe area in 5 ± 1-day-old flies (as in B) relative to controls. ANOVA with Dunnet's multiple comparison test, * is p<0.05, ** is p<0.01, *** is p<0.001. Effect size: η²=0.23. Bars: mean ± SD; points are individual animals N≥13 per genotype, 4 replicates.

(**D-D"**) Maximum intensity projection of confocal images of fly brains (5 ± 1-day-old) stained with anti-GFP (Green) and anti-Brp (Magenta), where anti-GFP marks EG and anti-Brp marks presynaptic sites of the antennal lobes in flies where CD8GFP is expressed via the EG driver MZ709-Gal4. Scale bar: 20 µm. (**D'**) Maximum intensity projection of confocal images of control (*w^{1118}*) and *Pink1^{KO-WS}* animals (**D"**) Maximum intensity projection of confocal images of animals with *Pink1* downregulation in EG and *Pink1^{KO-WS}* with *Pink1* rescued in EG.

(**E**) Quantification of GFP intensity within the glomeruli of the antennal lobe area in 5-day-old flies (as in D) relative to the control. ANOVA with Tukey's multiple comparison test, * is p<0.05, ** is p<0.01,

*** is p<0.001. Effect size: $\eta^2=0.36$ Bars: mean ± SD; points are individual animals N≥10 per genotype, 4 replicates.

Supplementary Figure 1

Representative confocal image of a (5 ± 1-day-old) MZ709-Gal4/+;; UAS-HisTageGFP/+ brain stained with anti-GFP (Cyan) and anti-ELAV (Magenta). Scale bar: 50µm; N= 3, 1 replicate.

Figure 2. Pink1 in EG is necessary to support synaptic integrity (with Supplementary figure 2)

(**A**) Representative ERG traces of indicated genotypes: ON peak is highlighted by the arrow.

(**B**) Normalized ON peak amplitude of flies (5 ± 1-day-old). ANOVA with Tukey's multiple comparison test, * is p<0.05, ** is p<0.01, *** is p<0.001. Effect size: $\eta^2=0.55$ Bars: mean ± SD; points are individual animals N≥11 per genotype, 3 replicates.

(**C-C"**) (**C**) Maximum intensity projection of confocal images of control and $Pink1^{KO-WS}$ in Mushroom Bodies (MB) of aged flies (22 ± 2-day-old), stained with anti-TH (cyan) and anti-DLG (magenta) antibodies - DLG is used to mark post-synaptic sites of MB. The black and white image is the middle Z-plane within the region of interest of the MB (ROI, yellow), which is used to represent the thresholded TH area (white). Scale bar: 20 µm. (**C'**) Maximum intensity projection of confocal images of w^{1118} with *Pink1* downregulation in EG. (**C"**) Maximum intensity projection of confocal images of $Pink1^{KO-WS}$ with *Pink1* rescued in EG.

(**D**) Quantification of the dopaminergic synaptic area at MB neuropil in aged flies (22 ± 2-day-old). ANOVA with Tukey's multiple comparison test, * is p<0.05, ** is p<0.01, *** is p<0.001. Effect size: $\eta^2= 0.38$. Bars: mean ± SD; points are individual animals N≥12 per genotype, 3 replicates.

Supplementary Figure 2

Examples of confocal Z-stacks of anti-DLG labeled fly brains (22 ± 2-day-old) of the indicated genotypes. Such images were used to delineate ROIs to quantify DAN innervation onto the MBs. (A) control (w1118/y;; MZ709-Gal4/+); (B) Pink1KO-WS/Y;; MZ709-Gal4/+; (C) flies where Pink1 is downregulated in EG (w1118/y;; MZ709-Gal4/ Pink1RNAi); (D) Pink1KO-WS flies with expression of wild type Pink1 in EG (Pink1KO-WS/Y;; MZ709-Gal4/UAS-Pink1).

Figure 3. Cell type-specific transcriptomics reveals modifiers of neuronal dysfunction

(**A**) Scheme of cell-type specific transcriptomics (Created in BioRender. Verstreken, P. (2025) https://biorender.com/p56u250).

(**B**) Scaled gene expression of representative genes for EG and neurons after sorting EG or neurons using the protocol described in (A) (R Core Team, 2022). N=2, 2 replicates.

(**C**) Differentially expressed genes in EG in $Pink1^{KO-WS}$ compared to control flies, plotted according to their Log2 Fold change and the -Log10 of the adjusted p-value. Intercept in red (-Log10 adjusted p-value = 4.31); light green dots are all detected genes, dark green are the 50 most deregulated genes, and the pink dot is from a gene positive in the genetic screen (D). *Two data points are outside the boundaries of the plot. To determine the transcriptomic profile of each genotype, a N=3 was used, and 3 replicates were performed.

(**D**) ERG ON peak value differences to $Pink1^{KO-WS}$ flies (5 ± 1-day-old) of control (red) and of $Pink1^{KO-WS}$ flies with DEGs downregulated, or upregulated, specifically in EG (5 ± 1 days old). ON peak values are expressed as the difference to $Pink1^{KO-WS}$. ANOVA with Dunnett's test, * is p<0.05, ** is p<0.01, *** is p<0.001. Effect size: η^2= 0.44. Bars: mean ± SD; points are individual animals N≥3 per genotype. *One data point is outside the boundaries of the plot.

Figure 4. Vps35 and Vps13 downregulation in EG rescues synaptic deficits in $Pink1^{KO-WS}$ flies (with supplementary figure 3)

(**A**) Representative ERG traces of control,$Pink1^{KO-WS}$ flies and $Pink1^{KO-WS}$ flies with $Vps35$ or $Vps13$ downregulated in EG.

(**B**) Quantification of the normalized ON peak response of flies with the genotypes in (A) (5 ±1-day-old). ANOVA with Dunnett's multiple comparison test, * is p<0.05, ** is p<0.01, *** is p<0.001. Effect size: η^2= 0.48. Bars: mean ± SD; points are individual animals N≥10 per genotype, 3 replicates.

(**C**) Maximum intensity projection of confocal images of Mushroom Bodies (MB) of aged flies (22 ± 2-day-old) of control, $Pink1^{KO-WS}$ flies and $Pink1^{KO-WS}$ flies with $Vps13$ downregulated in EG , labeled with anti-TH (cyan) and anti-DLG (magenta), DLG is used to mark the MB neuropil. The black and white image is the middle Z-plane within the region of interest of the MB (ROI, yellow), which is used to represent the thresholded TH area (white). Scale bar: 20 µm.

(**D**) Quantification of the dopaminergic synaptic area within MB of aged flies (22 ± 2-day-old). ANOVA with Dunnett's multiple comparison test, * is p<0.05, ** is p<0.01, *** is p<0.001. Effect size: η^2=0.23. Bars: mean ± SD; points are individual animals n≥22 per genotype, 5 replicates.

Supplementary Figure 3

Examples of confocal Z-stacks of anti-DLG labeled fly brains (22 ± 2-day-old) of the indicated genotypes. Such images were used to delineate ROIs to quantify DAN innervation onto the MBs. (A) control (w1118/y;; MZ709-Gal4/+); (B) Pink1KO-WS/Y;; MZ709-Gal4/+; (C) Pink1KO-WS flies with Vps13 downregulation in EG (Pink1KO-WS/Y;; Vps13RNAi/MZ709-Gal4).

Figure 5. Modulation of ER–mitochondria contact sites and lipid transfer in ensheathing glia (EG) rescues Pink1-dependent neuronal dysfunction.

Schematic representation of the suggested model (Created in BioRender. Verstreken, P. (2026) https://BioRender.com/sawj1pw).

(A) Loss of Pink1 leads to an abnormal increase in endoplasmic reticulum (ER)–mitochondria contact sites (represented by blue thick lines), resulting in enhanced ER-to-mitochondria lipid transfer and dysregulation of ER lipid composition (represented by yellow lipids). Increased organelle membrane contacts and lipid flux in EG contribute to neuronal dysfunction in a non–cell autonomous manner.

(B) Genetic downregulation of ER–mitochondria contact and lipid transfer regulators in EG rescues Pink1-induced neuronal phenotypes through two convergent mechanisms. Reduction of Vps35 decreases the number of ER–mitochondria contact sites, likely via MUL1-mediated Mitofusin (Mfn) turnover, leading to normalization of calcium and lipid homeostasis. In parallel, downregulation of Vps13, a lipid transfer facilitator at organelle contact sites, limits ER-to-mitochondria lipid transfer capacity, counteracting the excessive lipid flux induced by Pink1 loss. Both interventions restore organelle homeostasis in EG and result in rescue of neuronal dysfunction through a non–cell autonomous mechanism.

Declaration of interests

P.V. is the scientific founder of Jay Therapeutics. All other authors declare no competing interests.

Inclusion and diversity

We support inclusive, diverse, and equitable conduct of research.

Supplementary figure 1

Supplementary figure 2

A Control

B Pink1^{KO-WS}/y

C Pink1^{RNAi}

D Pink1^{KO-WS/y;} >UAS-Pink1

Supplementary figure 3