Structure and Cl⁻ Conductance Properties of the Open State of Human CFTR

Zhi-Wei Zeng[1,2], Christopher E. Ing[1,2], Régis Pomès[1,2]

[1] Molecular Medicine, Hospital for Sick Children, Toronto ON, M5G 0A4, Canada

[2] Department of Biochemistry, University of Toronto, Toronto ON, M5S 1A8, Canada

Abstract

The cystic fibrosis transmembrane conductance regulator (CFTR) is an anion channel that plays a vital role in water and ion secretion on epithelial surfaces. Loss of function in CFTR causes the life-threatening disease cystic fibrosis (CF). The functionally open state of CFTR has so far eluded detailed structural characterization. Although multiple near-atomic resolution structures of CFTR have been solved under conditions that promote channel opening, they all lack a continuous ion conduction pathway. In recent molecular dynamics (MD) simulations, structural fluctuations of human CFTR in a hydrated lipid bilayer led to the observation of transient Cl⁻ conducting states, but the stability and conduction properties of these putative open states were not established. Here, we conduct massively repeated simulations initiated from these Cl⁻ permeable conformations. Reproducible structural relaxation of the pore leads to a stable open conformation featuring five symmetrically arranged pore-lining helices. Unlike previously reported structures, this novel penta-helical arrangement reproduces experimentally determined properties of the open pore, including a Cl⁻ conductance close to that measured at physiological voltages. Together, our results support the validity of this newly identified pore conformation as a model of the fully open channel. Detailed analysis highlights the role of cationic pore-lining residues in the Cl⁻ permeation mechanism and suggests that the kinks observed in several transmembrane helices play a role in channel gating.

Introduction

CFTR is an anion channel that facilitates water and ion secretion from the cells of epithelial tissues (Berger et al., 1991; Riordan et al., 1989; Saint-Criq and Gray, 2017). It is vital for the healthy function of exocrine organs in respiratory (Smith et al., 1996), digestive (Ishiguro et al., 2009), and reproductive systems (Liu et al., 2012). Loss of function in CFTR causes cystic fibrosis, a devastating genetic disease that shortens life expectancy primarily due to its deleterious impact on lung health (Elborn, 2016; Smith et al., 1996). Over the past three decades, CFTR has been extensively studied with the aim of understanding the molecular pathology of cystic fibrosis – from its folding and processing (Bose et al., 2020; Lukacs and Verkman, 2012) to gating and ion conduction mechanisms (Hwang et al., 2018; Linsdell, 2017).

Even though CFTR functions as an ion channel, it belongs to the superfamily of ATP-binding cassette transporters – specifically, the ABCC exporter subfamily with type IV transmembrane fold (Dassa and Bouige, 2001; Ikuma and Welsh, 2000; Thomas et al., 2020). Like other ABCC proteins, CFTR consists of two transmembrane domains (TMDs), which form a gated pore in the membrane, and two nucleotide-binding domains (NBDs) which bind and hydrolyze ATP during gating. In addition, CFTR possesses a disordered R-domain that becomes phosphorylated by protein kinase A (PKA) before the channel undergoes ATP-dependent gating (Cheng et al., 1991). During gating, ATP binding promotes dimerization of the NBDs (Ikuma and Welsh, 2000; Levring et al., 2023); as a result, two ATP-binding sites are formed at the interface between the NBDs: a catalytic site and a degenerate site which are differentiated by their ability to hydrolyze ATP. Once it has been stabilized by bound ATP, the dimerization of NBDs promotes

conformational changes in the TMDs and opening of the gate. Subsequent hydrolysis of

ATP at the catalytic site ultimately results in the separation of the NBDs and closure of

the gate.

Over the past decade, invaluable structural insights into the functional states of the

gating cycle have been gained. Near-atomic resolution cryo-EM structures have been

determined for the unphosphorylated, ATP-free state, as well as for the phosphorylated,

ATP-bound state (Fiedorczuk and Chen, 2022a; Liu et al., 2019, 2017; Zhang et al.,

2018, 2017; Zhang and Chen, 2016). The folded domains of CFTR can be regarded as

consisting of two half-channels, each of which consists of one NBD, four TM helices

from a TMD, and two crossover TM helices from the other TMD (Figure 1A) – namely,

half-channel 1 (TM helices 1, 2, 3, 6, 10, 11 and NBD1) and half-channel 2 (TM helices

7, 8, 9, 12, 4, 5 and NBD2). Chloride ions enter the channel cavity through a lateral

portal between the two half-channels. A bottleneck region, sometimes referred to in the

literature as the channel gate (Gao and Hwang, 2015) or selectivity filter (Gao and

Hwang, 2015; Levring and Chen, 2024; Linsdell, 2016), separates the lumen of the

channel (inner vestibule) from the extracellular space (Figure 1B). The structures

provide insights into the mechanism of channel activation via phosphorylation.

Comparison of cryo-EM maps for unphosphorylated and phosphorylated CFTR

revealed that, upon phosphorylation, the R-domain is disengaged from the two half-

channels, allowing ATP-induced NBD dimerization to occur (Zhang et al., 2017; Zhang

and Chen, 2016).

Although more than ten near-atomic resolution (< 4 Å) cryo-EM structures of NBD-

dimerized CFTR have been determined to this date (Fiedorczuk et al., 2024; Fiedorczuk

and Chen, 2022a, 2022b; Liu et al., 2019; Zhang et al., 2018, 2017), none of them

appear to provide an open pathway for Cl⁻ permeation – a defining feature of the open

state. In these structures, the channel appears closed at the bottleneck despite its

activation by phosphorylation, ATP and, in some cases, activity enhancing drugs

(Fiedorczuk and Chen, 2022a; Liu et al., 2019; Zhang et al., 2018, 2017). Whether

these phosphorylated, ATP-bound structures represent the functional open state has

been elusive and somewhat contentious. On the one hand, some of these structures

were proposed to represent the *bona fide* open state of the channel as they could

narrowly accommodate fully dehydrated Cl⁻ ions in the gate region (Levring and Chen,

2024). However, this hypothesis disregards the high dehydration penalty of Cl⁻ and the

fact that CFTR is also permeable to anions larger than Cl⁻, both of which considerations

suggest that the pore bottleneck is significantly wider in the open state than in existing

experimentally derived structures (Linsdell et al., 1997; Linsdell and Hanrahan, 1998).

Accordingly, these ATP-bound structures have also been called "near-open" (Linsdell,

2018) or "quasi-open" (Simon and Csanády, 2021), implying that they closely resemble,

but do not represent, the truly functional open state of CFTR. Uncertainties regarding

the physiological relevance of structural models have been raised in studies of other ion

channels (Dämgen and Biggin, 2020; Deng et al., 2020; Gonzalez-Gutierrez et al.,

2017). In particular, there is evidence that the transmembrane solvent environment

(lipids vs detergents) during structural determination can affect the structure of

membrane proteins (Dalal et al., 2024; Deng et al., 2020). The use of detergents has

been suggested to explain the observed closed-pore structures of CFTR despite the

activating conditions (Cottrill et al., 2020; Zhang et al., 2017).

The above considerations raise the possibility that the near-open state may

spontaneously relax to the open state in a lipid bilayer—a rationale that has been

exploited successfully in MD simulation studies of other ion channels (Amaral et al.,

2012; Cerdan et al., 2018; Chakrabarti et al., 2010; Shrivastava and Sansom, 2000) and

motivated two recent computational studies of CFTR (Farkas et al., 2020; Zeng et al.,

2023). In simulations of the near-open state of zebrafish CFTR, a structure highly

similar to that of human CFTR, embedded in a lipid bilayer, the radius of the pore

bottleneck was only wide enough to allow Cl^- passage transiently, with no spontaneous

Cl^- permeation event observed on a sub-microsecond time scale (Farkas et al., 2020).

Using an enhanced sampling technique to help Cl^- ions overcome the energy barrier in

the bottleneck, two divergent Cl^- permeation pathways were identified (Farkas et al.,

2020). In the other study, repeated µs-long simulations of the putative near-open state

of human CFTR (Zeng et al., 2023) led to the observation of rare and transient Cl^-

conducting bursts: in two out of ten simulations, a total of 17 Cl^- ions traversed the pore

bottleneck in the presence of a strong TM voltage of -500 mV. Three distinct exit

pathways were observed. Consistent with hydrophobic gating from a closed (dry or de-

wetted) to a putative open (hydrated or wetted) state, these permeation bursts were

accompanied by structural fluctuations of pore-lining TM helices leading to widening and

wetting of the pore bottleneck.

These two simulation studies afford insight into the mechanism of Cl^- permeation that

goes beyond the ion pathway inferred from static structures of CFTR (Farkas et al.,

2020; Zeng et al., 2023). Consistent with the assumption that the latter are nearly open,

the second study demonstrates that conformational fluctuations in the sub-microsecond

time scale can lead to gate opening and ion permeation (Zeng et al., 2023). However,

this result was obtained under an artificially high TM voltage, raising the question of

whether the true open state of CFTR was attained. Indeed, the transient nature of the

conduction bursts observed suggests that further structural relaxation may be required

to reach a stable, fully open state that can support Cl^- currents at physiological voltages.

Here, we aim to elucidate the functional open state structure of human CFTR by

building upon that previous study. To overcome sampling limitations, we performed

hundreds of µs-long MD simulations extended from Cl^- permeable conformations of

ATP-bound, NBD-dimerized human CFTR obtained previously. Over a total sampling

time of 719 µs and across a broad range of voltage, further structural relaxation of the

channel involving the reproducible rearrangement of helices in the outer region of the

TMD led to a stable conformation featuring five symmetrically arranged pore-lining

helices. This novel arrangement satisfies experimentally determined properties of the

open pore, including Cl^- conductance, and likely represents a valid model of fully open

human CFTR under physiological conditions. The Cl^- permeation mechanism and

implications for channel gating are analyzed and discussed.

Results & Discussion

A novel, penta-helical pore conformation

We performed 719 replicates of microsecond-long unrestrained MD simulations of ATP-

bound, NBD-dimerized human CFTR, most of which were initiated from Cl^- permeable

conformations obtained in our previous study (Zeng et al., 2023). The overall structure

of the protein remained stable, with an average RMSD < 2 Å for Cα atoms in the TM

region (Figure 2 – figure supplement 1). The NBDs remained dimerized and the

fluctuation of TM helices in the plane of the inner leaflet was limited (Figure 2B). In

contrast, the amplitude of structural fluctuations in the plane of the outer leaflet, which

hosts the extracellular end of the pore, was noticeably larger (Figure 2A). Remarkably,

while the planar distribution of most TM helices relative to the rest of the helical bundle

was essentially unimodal, both TM1 and TM11 sampled bimodal distributions, with one

state closer to the TM helical bundle ("in") and the other ~4 Å farther away ("out")

(Figure 2A and C). The two states were sampled in nearly equal proportions by both

helices (in:out ratio of 54:46 for TM1; 60:40 for TM11). Transitions between the two

states of TM1 and TM11 were also well sampled, occurring in 240 and 328 out of 719

trajectories, respectively. Importantly, the positions of TM1 and 11 were anti-correlated,

in that as TM1 moved away from the centre of the pore, TM11 moved towards it,

resulting in a symmetrical pentagonal pore lined by TM helices 1, 6, 8, 11, and 12

(Figure 2C and D). The anti-correlation of TM1 and 11 positions is confirmed by

principal component analysis (PCA) of the spatial coordinates of TM helices, which

reveals that four conformational states, labelled α to δ, were sampled in our simulations

(Figure 3A). The first principal component (PC1), which accounts for 25% of structural

fluctuations in the conformational ensemble, captures the collective movements of TM

helices 1, 2 and 11 leading to the symmetrical pentagonal pore or α state (Figure 3B

and C, Figure 3 – figure supplement 2). In contrast, PC2, which accounts for 8% of

structural fluctuations, captures comparatively more subtle movements of TM helices,

as both β and γ states are characterized by a "squeezed" arrangement of pore-lining

helices (Figure 3B and D). This squeezed arrangement also dominates in the δ state,

which includes the near-open cryo-EM structure (PDB: 6MSM) and occupies a location

intermediate between the other three conformational states (Figure 3A and B). To gauge

the stability of the four conformational states, we analyzed the time evolution of their

fractional population by considering all 719 MD trajectories in parallel. While the

populations of α, β, and γ either increased or remained stable, the population of state δ

decreased over the 1-μs simulation time, confirming that the cryo-EM structure is a

metastable intermediate state in a lipid bilayer environment (Figure 3 – figure

supplement 1).

Among all states sampled, the structure of state α is the most compatible with the open

state of human CFTR. As a result of the symmetric arrangement of the five pore-lining

helices, the radius of the pore bottleneck was larger by 0.5~1 Å compared to the other

states (Figure 4, Figure 4 – figure supplement 1). Notably, α is the only state featuring

TM11 as a pore-lining helix due to the movement of this helix towards the centre of the

pore. Such movement brings TM11 closer to TM6 and away from TM12, consistent with

cross-linking experiments suggesting the close proximity of bottleneck-lining residues of

TM6 and TM11 in the open state (Fig. 4 – figure supplement 2) (Wang and Linsdell,

2012). Furthermore, the solvent-accessible surface areas (SASA) of bottleneck-lining

residues T1115 and S1118 from TM11 are both greater compared to states β-δ,

consistent with experimental findings that these residues are accessible to aqueous

reagents in the open state (Fig. 4 – figure supplement 2) (Fatehi and Linsdell, 2009).

Overall, state α exhibits experimentally demonstrated features of the open pore,

indicating that it most likely represents the open state of CFTR compared to the existing

cryo-EM structures as well as all the other TM conformational states sampled in our

simulations.

Cl⁻ permeation pathway

In our previous study of human CFTR, Cl⁻ permeation events occurred in 2 out of 10

microsecond-long MD simulations carried out at -500 mV TM voltage (Zeng et al.,

2023). We identified three distinct Cl⁻ permeation pathways through the pore bottleneck:

the "1-6" and "1-12" pathways, in which Cl⁻ ions exited through lateral openings between

the extracellular ends of TM helices 1 & 6 and 1 & 12, respectively; and the

"intermediate" pathway, in which Cl⁻ ions travelled along a straight path in between the

other two pathways (Figure 5A and B). It was also found that the pathway taken by

permeating Cl⁻ ions depended on minor differences in the structure and arrangement of

pore-lining helices (Zeng et al., 2023). Remarkably, only one of these two simulations

briefly sampled putative open state α. Moreover, in addition to α, Cl⁻ permeation events

also occurred in states γ and δ, from which some Cl⁻ permeable snapshots served as

starting configurations for the extended simulations in the present study (Figure 5 –

figure supplement 1). Due to the small number of permeation events, the relative

importance of the three permeation pathways as well as the corresponding pore

structures could not be determined from the previous study. Our current simulation set

contains more than 2000 Cl⁻ permeation events, amongst which all three permeation

pathways were sampled. However, the pathway taken by Cl⁻ ions is strongly influenced

by the conformation of the channel pore (Figure 5C and D). Whereas virtually all Cl⁻

permeation followed the 1-6 pathway in states β and γ, all three permeation pathways

were sampled in both α and δ. Note that the enlarged pore bottleneck in state α allowed

permeant Cl⁻ ions to occupy the region that previously separated the intermediate and

1-12 pathways, blurring the boundary between them (Figure 5A and B). As such, the

intermediate and 1-12 pathways shall henceforth be collectively referred to as the

"central" pathway.

Multiple Cl⁻ ion binding sites are present inside the pore. Many protein residues,

primarily ones with polar and positively charged sidechains, engaged in direct contact

with Cl⁻, displacing water molecules in its first solvation shell (Figure 6B and C).

Consistent with our previous findings, there are two main Cl⁻ binding sites inside the

channel cavity (Zeng et al., 2023): one near the intracellular entrance involving K190,

R248, R303, R352, W356, and the other deeper inside the pore with R1097, K95, and

R134 (Figure 6A). In addition, a third Cl⁻ binding site consisting of S341 and N1138 is

found below the bottleneck.

Despite the abundance of hydrophobic residues in the bottleneck region (-5 < z < 5 Å),

permeant Cl⁻ ions are partially solvated by polar and charged residues, including S341,

T338, R334, Y914, Y917 and S1118 (Figure 6C, Figure 6 -- figure supplement 1). Direct

binding to pore residues is responsible for the partial dehydration of Cl⁻ ions, which lose

1 to 3 water molecules out of an average of 7.3 ± 0.8 from their first solvation shell in

bulk water (Figure 6D and E). Remarkably, in ~15% of permeation events, the

sidechains of non-polar residues lining the bottleneck, such as F337, also formed direct

contacts with Cl⁻ ions and disrupted their solvation shell. Experimental studies have

shown that mutations at residues R334, F337, T338, and Y914 affect the relative

permeability among anions according to their hydration free energy – the greater the

dehydration cost, the less permeable the ion, a mechanism known as lyotropic

selectivity (Levring and Chen, 2024; Linsdell et al., 2000, 1998; Negoda et al., 2019).

Taken together, these findings support the role of these residues in partial dehydration

of Cl$^-$ ions in the rate limiting step of permeation. Because these residues appear to line

the 1-6 pathway in the cryo-EM structure, this pathway was recently suggested to be

the only Cl$^-$ permeation route (Levring and Chen, 2024). As noted above, it was

proposed that the existing "near-open" cryo-EM structures in fact represent the

functional open state of the channel, whereby fully dehydrated Cl$^-$ ions pass through the

1-6 pathway without requiring further pore widening (Levring and Chen, 2024). Our

findings challenge this proposed picture of Cl$^-$ permeation in two respects. First,

following TM movements, the bottleneck region of the pore expands, which allows Cl$^-$

ions to permeate through two diverging pathways instead of a single pathway. Second,

permeant Cl$^-$ ions only ever undergo partial dehydration, regardless of which pathway

(1-6 or central) is taken (Figure 6D and E, Figure 6 – figure supplement 1).

Cl$^-$ conductance

From numerous Cl$^-$ permeation events at different TM voltages, we next evaluate the Cl$^-$

conductance of the channel. To quantify the Cl$^-$ conductance of states α–γ, we

partitioned the simulation trajectories by state and computed the Cl$^-$ current in terms of

the number of translocation events over time spent in each state at different voltages.

Across all negative TM voltages, the putative open state α is most permeable to Cl$^-$

ions, exhibiting a Cl$^-$ conductance of 3±1 pS, a value approaching the range of 6~10 pS

measured experimentally at physiological voltages (Sheppard and Welsh, 1999) (Figure

7). In contrast, states β-δ showed almost no Cl$^-$ conductance at low negative TM

voltages (-100 < U < 0 mV) and only produced Cl$^-$ currents lower than state α at high

negative voltages, suggesting that they are unlikely to be functional open states. At

positive TM voltages, however, all four states virtually lost the ability to conduct Cl⁻ ions

(e.g. 0.5±1 pS for state α at $0 < U < +100$ mV). This result is at odds with the linear IV

relationship across all voltages, from negative to positive (Sheppard and Welsh, 1999).

Although some studies have reported minor inward rectification at positive voltages for

the wild-type channel (Cai et al., 2003), the rectification of IV-curves computed from our

MD simulations is much stronger.

To explain the observed low conductance and strong inward rectification compared to

experiment, we examined factors influencing Cl⁻ permeation. Chloride ions traversing

the bottleneck region made frequent direct contact with positively charged R334.

Notably, the sidechain of R334 can adopt two distinct conformations differing by 6 Å in

the axial position of the center of charge: one pointing up and away from the pore

bottleneck ("up") as seen in the PDB structure; the other dunking down and extending

into the bottleneck region ("down") to form an ion pair with permeant Cl⁻ ions (Figure 8A

and B). When R334 is down, the average Cl⁻ ion density in the bottleneck as well as Cl⁻

conductance are increased (Figure 8C-E). As a control, the Cl⁻ conductance of the

channel was unaffected by the conformational state of the sidechain of R134, another

pore-lining arginine from the inner vestibule whose sidechain sampled two slightly

different conformations (Figure 8 – figure supplement 1), indicating that the observed

conductance enhancement is unique to R334.

Furthermore, the sidechain of R334, whether in the up or down conformation, can form

a salt-bridge with either E1124 or E1126, two residues in the middle of flexible

extracellular loop 6 (ECL-6) linking TM11 and 12. This interaction occurs when the loop

is closer to the centre of the pore, resulting in the dunking of either glutamate sidechains

(Figure 9A-B), blockage of the central ion permeation pathway (Figure 9E-F), and

neutralization of the positive charge of R334 (Figure 9C-D). Glutamate dunking results

in a 5-fold decrease in conductance in the range -100 < U < 100 mV in the penta-helical

state (Figure 9G).

Our analysis shows that the configuration of charged residues at the extracellular mouth

of the pore affect the conductance of the penta-helical pore. The most favourable

conditions for Cl⁻ conduction are: 1) the dunked conformation of R334, which assist Cl⁻

ions to cross the narrowest region of the pore; and 2) the disengagement of two ECL-6

Glu residues from the conduction pathway. The apparent conductance of the penta-

helical pore at each voltage is strongly correlated with the fraction of time when these

two conditions are met (Figure 9 – figure supplement 1). The fact that these conditions

were seldom satisfied at positive voltages explains the observed rectification. This bias

is likely due at least in part to the fact that the positive electric field tends to push

cationic R334 upwards and anionic ECL-6 Glu sidechains downwards towards the pore,

resulting in a greater likelihood of charge pairing between these residues. In addition, it

is possible that initial conformations of the ECL-6 loop do not reflect equilibrium and

more generally, that the conformational equilibria of the extracellular loops are not

adequately sampled in the simulations. In particular, interactions with other extracellular

loops that could help keep ECL-6 away and disengaged from the central pore may not

be modelled accurately. In that perspective, the omission of 11 residues in glycosylated

loop ECL-4 linking TM7 and 8, which was not modeled in structural studies to date or in

the present study but is close to ECL-6, could alter the conformational properties ECL-6.

Experimental evidence supports the role of R334 in enhancing Cl^- conductance in

CFTR. For instance, loss-of-charge, CF-causing mutation R334W shows reduced Cl^-

conductance compared to wild-type (cftr2.org, n.d.; Sheppard et al., 1993). Our results

suggest that the ability of R334 sidechain to extend into the pore bottleneck is an

important feature of the functional open state of CFTR. In support of this finding, it was

shown that R334 becomes less accessible from the extracellular space in the open

state than in the closed state (Zhang et al., 2005). The down conformation of the R334

sidechain could be stabilized through cation-π and π-π interactions with the aromatic

ring of F337 (Figure 8B, Figure 8 – figure supplement 2). Consistent with this

observation, both R334K and F337A single-point mutants displayed reduced

conductance (Ge et al., 2004; Gong and Linsdell, 2004; Linsdell et al., 2000). It should

be noted that classical forcefields such as the one that we are using, which approximate

electronic distributions as atomic point charges, do not explicitly capture the electric

quadrupole contributions of aromatic rings in π-interactions (Lemkul, 2020; Mackerell,

2004; MacKerell et al., 1998). As such, it is possible that these interactions stabilize the

down conformation of R334 sidechain to a greater extent than captured in our

simulations, which could result in greater Cl^- conductance and a better agreement with

experimental single channel recordings. On the other hand, there is no evidence

supporting the roles of E1124 or E1126 in CFTR function. Instead, recent study

suggests that E1124A and E1126P mutations retain or slightly improve the function of

the channel (Simon and Csanády, 2021), suggesting that the presence of negative

charges at these two positions is not required for function and may even be inhibitory,

consistent with our conclusion.

Taken together, the above results validate state α as the functional open state of human

CFTR. Although limitations were identified in the accuracy of the conformational

ensemble of loop residues at the extracellular mouth of the pore, the pentameric

conformation of pore helices derived in the current study is stable and consistent with

CFTR function. Our findings provide a realistic, physiological depiction of Cl$^-$ permeation

through the open state of CFTR, which features a wide central pathway and a side

pathway between TM helices 1 and 6, with a preference for the central pathway (Figure

5D). We shall henceforth refer to state α as the "MD-open" state. In contrast, ion

translocation only occurs at large, non-physiological voltages for states β-δ, confirming

their status as "near-open" conformations. States β and γ shall be referred to as stable

"MD-closed" states to distinguish them from the metastable near-open state δ.

Insights into the gating mechanism

Having characterized the structure of the functional open state of the channel, we next

seek to gain structural insights into CFTR gating. In addition to extensive simulations of

the NBD-dimerized, ATP-bound CFTR, we performed 20 replicates of 1-microsecond-

long simulations of the ATP-unbound, NBD-separated cryo-EM structure of closed state

human CFTR (PDB: 5UAK) in a hydrated lipid bilayer. Consistent with past MD

simulation studies (Corradi et al., 2018; Tordai et al., 2017), the NBDs either partially or

fully dimerized in 15 out of 20 trajectories (Figure 10 – figure supplement 1).

Dimerization occurred very rapidly due to the fact that the structural model is missing

the R-domain that keeps two half-channels apart in the closed, unphosphorylated state

of the channel. This omission makes the initial NBD-separated conformation highly

unstable. In the presence of the R-domain, one expects that the dimerization of NBDs

would occur much more slowly as the R-domain gradually disengages from the two half-

channels upon phosphorylation.

Despite the rapid dimerization of the NBDs, the arrangement of the extracellular TM

helical segments remained unchanged over the 1-µs simulation time (Figure 10 – figure

supplement 2), indicating that further conformational changes in the TM region need to

occur for the gate to open following NBD dimerization. Accordingly, comparison of the

MD-open state with the structural ensemble obtained after fast NBD dimerization of the

closed state highlights persistent structural differences in the extracellular segments of

TM helices, especially in half-channel 2. In particular, pore opening requires movement

of the extracellular end of TM8 towards the center of the pore by ~8 Å and TM12 away

from it by ~7 Å (Figure 10A-C, Figure 10 – video 1). To accommodate the outward

movement of TM12, the extracellular ends of TM helices 10 and 11 must move apart

from each other (Figure 10D). The same conformational changes of TM helices 8 and

12 from the ATP-unbound closed state to the near-open state were previously identified

based on the comparison between closed and near-open cryo-EM structures (Zhang et

al., 2017). Contrary to the previous hypothesis, however, our results suggest that

reaching the open state from near-open state does not involve further conformational

changes in these two helices (Figure 2D).

Whereas reaching the open state from the ATP-unbound, NBD-separated closed state

primarily involves the closure of two half-channels at the NBDs and rearrangements of

TM helices 8 and 12 within half-channel 2, reaching the MD-open state from near-open

states primarily involves the movements of TM helices within half-channel 1, with the

analysis above showing that collective motions of TM helices 1, 2, and 11 were the

dominant changes involved in channel opening (Figures 2 and 3). What is the structural

mechanism underlying such a movement? A notable aspect of the CFTR structure is the

abundance of kinks in the TM helices, of which all except TM7 contain at least one kink.

Helical kinks often result from structural distortions due to proline or glycine residues,

which impart structural flexibility within the helix and have been suggested to facilitate

conformational changes in other transmembrane proteins (Chamberlain et al., 2003;

S.P. Sansom and Weinstein, 2000). In human CFTR, we identified five sites where

helical kinks undergo changes between MD-closed and MD-open states: two on TM1, at

G85 and P99; two on TM2, at G126 and P140; and one on TM11 at W1098 (Figure

11A). At P99, the average kink angle increases slightly from ~20° in the MD-closed

states to ~25° in the MD-open state (Figure 11 – figure supplement 1A and B). In

contrast, at G85, G126, and P140, reaching the MD-open state involves changes in the

orientation of the kinks rather than in the magnitude of kink angles. Specifically, upon

reaching the MD-open state from the MD-closed states: (1) the middle segment of TM1

(residues 85-96) swivels with respect to the intracellular segment (residues 78-85) by

~20° (Figure 11F and G); (2) the extracellular segment of TM2 (residues 117-128)

swivels with respect to the middle segment (residues 129-138) by ~40° (Figure 11B and

C); and (3) the middle segment of TM2 swivels with respect to the intracellular segment

(residues 140-149) by ~30° (Figure 11 – figure supplement 1C and D). Finally, at the

W1098 hinge point, the kink angle is larger in the MD-open state by ~8° compared to

the MD-closed states, while the extracellular segment (residues 1098-1118) also swivels

around the middle segment (residues 1077-1097) by ~20° (Figure 11D). Among all

observed patterns implicated for channel opening, the changes in kink at W1098 and in

swivel at G85 and G126 are the most strongly correlated with the transition between

MD-closed and MD-open states (Figure 11C, E and G). Altogether, changes in both the

magnitude and the orientation of these helical kinks, even if not necessarily all occurring

simultaneously, underlie the structural transition between MD-open and MD-closed

states (Figure 11 – video 1).

Studies of CFTR gating have understandably been focused on the coupling between

conformational changes in the TMDs and ATP-binding-induced NBD dimerization. In this

process, the gate opens in the TMDs in response to NBD dimerization, resulting in ion

conduction bursts as observed in single channel recordings (Cai et al., 2011).

Nevertheless, even during conduction bursts with dimerized NBDs, brief interruptions

occur frequently (Cai et al., 2011; Levring et al., 2023). The transitions between

conducting state and such "flicker closures" have been interpreted as ATP-independent

gating events (Cai et al., 2003; Chen et al., 2017). Correspondingly, it is possible that

conformational dynamic transitions such as those observed in our simulations of MD-

open and MD-closed states correspond to such ATP-independent gating. In support of

the functional relevance of helical kink flexibility at these sites, single-point mutations of

G85, P99, and G126 are known to cause CF (cftr2.org, n.d.; Decaestecker et al., 2004;

Sheppard et al., 1996; Wagner et al., 1994). Although functional and clinical data

associated with P140 are not currently available, this residue is noteworthy for being in

a π-bulge, a type of helical defect linked to functional roles in other proteins (Cooley et

al., 2010). Finally, our analysis shows that changes in the magnitude of the kink angle

near W1098 are strongly correlated to transitions between MD-closed and MD-open

states (Figure 11E). This finding may seem surprising considering that unlike proline

and glycine, tryptophan is not usually associated with helix kinks. However, consistent

with the functional importance of this locus, W1098R and W1098C are CF-causing

mutations (cftr2.org, n.d.).

Our results show that changes in helical kinks can affect the relative arrangement of

pore-forming helices and are involved in near-open to open transitions in CFTR. Such

changes are also likely to play a role in gating, as suggested by the fact that multiple

residues located at helical kinks in TM helices are loci of CF mutations. The role of small

changes in helical conformations in gate opening and closing has been noted in other

ion channels. In the CorA magnesium channel, gating is coupled to a 10-degree change

in the tilt of kinked pore-lining helices that modulates the diameter of the pore

(Chakrabarti et al., 2010). In Kv and MthK channels, gating involves a ~10-degree

change in the internal kink of pore-lining helices (Fowler and Sansom, 2013; Gu and de

Groot, 2020). In NavM channels, the open state is reached through a local α-to-π

transition of a pore-lining helix (Choudhury et al., 2023; Choudhury and Delemotte,

2023). These examples, along with our findings on CFTR, show that the gating of ion

channels can involve changes in helical structure as well as global conformational

changes.

Conclusions

In this study, we have derived and characterized a structural model of the open state of

human CFTR through massively repeated MD simulations of a near-open structure of

the ATP-bound, NBD-dimerized channel. Reproducible structural relaxation of the pore

in a lipid bilayer leads to a stable open state structure featuring an approximately

symmetric arrangement of five transmembrane helices lining an expanded pore

bottleneck region. The functional relevance of this penta-helical open state is

demonstrated by its ability to sustain Cl⁻ currents at physiological voltages with a

conductance comparable to experimentally measured values. The detailed analysis of

ion solvation and ion-channel interactions highlights the role of cationic residues in the

ion conduction mechanism throughout the permeation pathway and reveals how the

R334 sidechain facilitates the translocation of partly dehydrated Cl⁻ ions through the

hydrophobic pore bottleneck.

While the present study does not address the allosteric mechanism linking NBD

dimerization to channel opening, it provides insights into the gating mechanism in the

pore domain (TMDs). The comparison of closed and open state structures suggests that

ATP-dependent gating at long timescale primarily involves rearrangements of TM

helices within half-channel 2, whereas the changes governing near-open to open

transitions, which may be relevant to ATP-independent gating on a faster timescale,

occur within half-channel 1. In this process, opening and closing of the hydrophobic

pore bottleneck involves changes in the orientation and magnitude of flexible kinks in

TM helices.

Understanding the structure and functional dynamics of CFTR is an important step

towards unraveling the molecular pathology of CF. As a case in point, we note that

several residues shown in this study to be implicated in pore opening and Cl⁻

permeation are loci of CF-causing mutations. Future studies of the atomistic open-state

structure of human CFTR derived in this work should help gain a better understanding

of gating and conduction as well as their defects in CF-causing mutants.

Methods

Molecular Systems

The preparation of the system containing ATP-bound, NBD-dimerized CFTR was

described in detail in our previous work (Zeng et al., 2023). Briefly, the structural model

of phosphorylated, ATP-bound human E1371Q CFTR (PDB: 6MSM) was used. The

PDB structure is missing certain loop regions (residues 410-434, 890-899, 1174-1201,

and 1452-1489) and the disordered R-domain (residues 638-844), which are not

modelled in our study. All chains were acetylated at the N-termini and amidated at the

C-termini. Two Mg-ATP moieties bound to the protein at the NBD interface in the PDB

structure were kept, while all other putative lipids and an unknown helix at the TMD-

NBD interface were removed. The Mg-ATP bound human CFTR was embedded in a

hydrated POPC bilayer with 150 mM NaCl using CHARMM-GUI server (Jo et al., 2008).

For the ATP-free, NBD-separated CFTR, the simulation box was prepared in a similar

way. The structural model of unphosphorylated, ATP-free human wild-type CFTR (PDB:

5UAK) was used (Liu et al., 2017). Residues that were missing in the PDB structure

were not modeled, including a short N-terminal segment (1-4), the loops connecting

each TMD-NBD pair (403-438, 1173-1206), the R-region (646-843), the extracellular

loop between TM7 and TM8 (884-908), and the segment at the C-terminal end of NBD2

(1437-1489). A putative helix from the R-domain in the PDB structure with unknown

amino acid composition was removed. Before embedding the CFTR protein into lipids,

the PPM server of Orientations of Proteins in Membranes (OPM) database was used to

determine the starting position of the lipid bilayer (Lomize et al., 2012). The model for

CFTR with Mg-ATP bound was then embedded in a POPC bilayer and solvated in water

with 150 mM NaCl using the CHARMM-GUI server without water or ions inside the

channel pore (Jo et al., 2008). A total of 255 POPC molecules added around the protein

per periodic box.

Simulation setup and protocol

All MD simulations were conducted using GROMACS versions 2016 and later

(Abraham et al., 2015). The CHARMM36 forcefield was used for protein, lipids, ions,

ATP, together with the TIP3P water model (Best et al., 2012; Huang and MacKerell,

2013; Jorgensen et al., 1983; Klauda et al., 2010). Simulations were run at constant

temperature and pressure (T = 300 K, p = 1 atm) at 2 fs integration timesteps. Constant

temperature was maintained using the Nosé–Hoover thermostat (τ_T = 0.5) (Hoover,

1985; Nosé, 1984); constant pressure was maintained using the Parrinello-Rahman

barostat (τ_p = 2.0) (Nosé and Klein, 1983; Parrinello and Rahman, 1980). Semi-isotropic

pressure coupling was used, with isothermal compressibility set to 4.5×10^{-5} bar^{-1} both in

the *xy*-plane and along the *z*-axis. Nonbonded interactions were calculated using Verlet

neighbor lists (Páll and Hess, 2013; Verlet, 1967). Lennard-Jones interactions were cut

off at 1.2 nm and a force-based switching function with a range of 1.0 nm was used. The

particle-mesh Ewald (PME) method was used to compute electrostatic interactions with

a real-space cut-off of 1.2 nm (Darden et al., 1993; Essmann et al., 1995). The LINCS

algorithm was used to constrain covalent bonds involving H atoms (Hess, 2008).

For initial simulations started from the PDB structures of CFTR, the system was

subjected to energy minimization followed by a multi-step *NpT* equilibration protocol.

The system was first subjected to steepest descent energy minimization until maximum

force dropped below 1000 kJ/mol/nm. Random velocities were generated at the

beginning of the *NpT* equilibration phase, which was conducted in three 10-ns stages,

successively with protein heavy atoms, protein backbone atoms, and protein Cα atoms

restrained (force constant k = 1000 kJ/mol/nm^2 in *xyz* directions). *NpT* equilibration was

followed by production run without restraints. All production run was 1-µs-long. For

simulations with TM voltage, a constant electric field of strength between -32 mV/nm

and +32 mV/nm was applied along the *z*-axis of the simulation box which is normal to

the lipid bilayer. During the production runs, the *z*-dimension of the simulation box

stabilized to and fluctuated around 16.5 nm. In all simulations, new random velocities

were generated at the beginning of production runs. For extended simulations,

snapshots from different timepoints of other MD trajectories were used as the initial

configurations. No extra energy minimization or equilibration phase precedes extended

production runs.

Among all simulations incorporated in the analysis, 649 of which started from both Cl$^-$

permeable and impermeable configurations obtained in our previous study, and 50 of

which were further extended from these simulation trajectories. For completeness, the

dataset analyzed herein also includes 20 simulation trajectories obtained from our

previous study, which started from the experimental "near-open" structure (PDB:

6MSM). Information about the initial configurations and how the extended trajectories

are spawned are presented in Table 1 and Supplementary file 1.

General analysis methodology of simulation data

Prior to analysis and visualization, CFTR structure from all snapshots of production run

were aligned through minimization of RMSD of Cα atoms of TMDs. After alignment, the

overall permeation pathway is parallel to the *z*-axis, which was used as the channel

pore axis. The origin of the z-axis was approximately at the level of T338, and the POPC

bilayer spanned the range from -30 Å to 10 Å on the z-axis. All visualizations and

renderings of molecules were done with VMD (Humphrey et al., 1996). Scripts written in

Tcl and python were used to perform analyses.

Analysis of conformational ensemble of TM helices

The *xy*-positions of TM-helices were represented by the Cα of residues that lie

approximately at the same *z*-level. At the inner leaflet level, Cα of residues 82, 142, 197,

241, 303, 356, 863, 938, 990, 1033, 1097, and 1152 were used to represent TM helices

1-12. At the outer leaflet level, Cα of residues 106, 121, 219, 324, 334, 882, 914, 1011,

1118, and 1131 were used. Due to that TM3, 4, 9, and 10 end at this level

extracellularly, residues 219 and 1011 were used to represent the positions of helix

pairs TM3-4 and TM9-10, respectively.

Principal component analysis

Principal component analysis (PCA) was performed on the 3D Cartesian coordinates of

TM helical backbone atoms (i.e. carbonyl carbon and oxygen, α-carbon, and amide

nitrogen atoms) from all production run snapshots. K-means algorithm was used to

cluster TM conformational states of CFTR. Both PCA and clustering were aided by

Scikit-learn python library (Pedregosa et al., 2011).

Pore dimension analysis

Pore dimensions were analyzed by HOLE2 program implemented through MDAnalysis

(Gowers et al., 2016; Michaud-Agrawal et al., 2011; Smart et al., 1996). Residues 1-380

and 846-1173, including all hydrogens, were used to define the channel for HOLE2

analysis. The radius of the bottleneck in a given snapshot is represented as the

minimum of the pore radius in the region $-5 < z < 5$ Å computed by HOLE2. Other

parameters for running HOLE2 are included in the associated repository.

Solvent accessibility analysis

Calculation of solvent accessible surface areas (SASAs) for residues T1115 and S1118

using VMD *measure* functionality (Humphrey et al., 1996). A probe radius of 1.4 Å was

used, which corresponds to a coarse-grained spherical model for water. As T1115 is

located at the protein-lipid interface, both protein and POPC lipids contribute to the

solvent-excluded region during calculation.

Analysis of Cl⁻ translocation and coordination

A Cl⁻ permeation/translocation event occurred if it: 1) crossed through $z = 0$ in the

middle of the pore bottleneck; and 2) left the pore region subsequently without re-

crossing $z = 0$. The time of translocation is when crossing $z = 0$ occurs. Details about

the pore region definition are provided in the associated repository.

Time series for the 3D positions of the 17 Cl⁻ ions that underwent translocation from our

previous study (Zeng et al., 2023) were grouped into three clusters based on similarity

measured by dynamic time warping (DTW) distance (Sakoe and Chiba, 1978). The

three clusters were visually verified to correctly correspond to the 1-6, intermediate, and

1-12 pathways. To assign translocation paths to all Cl⁻ permeation events in a high

throughput manner, all trajectories of permeating Cl⁻ ions were collected and clustered

into three categories based on their DTW distance to the cluster centroids. DTW

analysis and clustering were aided by Tslearn python library (Tavenard et al., 2020).

Analysis of binding and solvation of Cl⁻ ions were aided by MDAnalysis. A molecule or

residue is considered to form a direct contact with a Cl⁻ ion if it contains atom within 3 Å

of the Cl⁻ ion, based on radial pair distribution functions computed in our previous study

(Zeng et al., 2023). Low voltage simulations (i.e. -120 mV ~ +120 mV) were used to

analyze Cl⁻ binding and solvation.

Analysis of current-voltage relationships

Transmembrane voltages *U* were computed by $U = E_z L_z$, where *Ez* is the electric field

strength in the *z*-direction, and *Lz* is the length of the periodic simulation box in the *z*-

direction (Gumbart et al., 2012; Roux, 2008). At each given TM voltage, the simulation

dataset was partitioned by the states of CFTR (i.e. TM helix conformation state α-δ,

arginine sidechain conformational state, or ECL-6 conformational state). Chloride

current at a certain voltage in a certain state is computed by $I = eN/t$, where *e* is the

elementary charge, *N* is the number of translocation events, and *t* is the total time spent

in the state. Simple linear regression without intercept was used to construct the

current-voltage curve and determine the channel conductance (i.e. slope of the fit to

current vs voltage plot).

Helix kink analysis

The descriptors used to characterize helix kink, specifically kink and wobble/swivel

angles, are inspired by the ProKink approach (Visiers et al., 2000). The kink angle

between two helical segments were calculated as the angle between their principal

axes. To compute the swivel angle, the protein structure from all snapshots were

reoriented such that one of the helical segments is aligned with the *z*-axis. The swivel

angle is represented as the azimuth angle of the other helical segment around the z-

axis.

Acknowledgements

602 We thank Paul Linsdell and Christine E. Bear for discussion of early results. This work

603 was supported by Canadian Institutes of Health Research Grant MOP130461. MD

604 simulations and analyses were enabled by supercomputing resources and support

605 provided by SciNet (www.scinet.ca) and the Digital Research Alliance of Canada

606 (www.alliancecan.ca). Z-W.Z. is supported by SickKids Restracomp.

Data availability

608 Data generated from the study is provided on Zenodo at 10.5281/zenodo.14323352.
609 Code created to analyze the data is provided on Github at
610 https://github.com/wilzzw/cftr2.

Author contributions

612 Z-W.Z. and R.P. conceptualized the study; Z.W.Z. and C.E.I. performed the MD
613 simulations; Z.W.Z. analyzed the data; Z.W.Z. and R.P. wrote and edited the manuscript.

Conflict of interest

The authors declare no conflict of interest.

Figure 1. Overview of CFTR channel structure. (A) Cartoon representations of the

3D structure of CFTR (PDB: 6MSM). Different parts of the channel are coloured

according to the domain (left) or the half channel (HC; right) to which they belong. The

dashed lines indicate the membrane region. **(B)** Side view of the channel lumen. Cl⁻

ions (cyan spheres) from multiple simulation snapshots are overlaid to TM helices (gray

cartoon), outlining the space accessible to ions inside the pore. Intracellular Cl⁻ ions

cross a lateral intracellular (IC) entrance (*z*~-40 Å) into the inner vestibule which leads

to the pore bottleneck (*z*~0 Å) separating the inner vestibule from its outer or

extracellular (EC) counterpart. Note the absence of Cl⁻ from the bottleneck region in this

"near-open" state of the channel.

Figure 2. Conformational ensemble of TM helices. (A) Spatial distribution of TM helices in the plane (*x*,*y*) of the outer lipid leaflet. Conformational transitions result in well resolved bimodal distributions of TM helices 1 and 11. The extracellular ends of TM helices 3-4 and 9-10 overlap in this plane. **(B)** In contrast to the outer leaflet, the spatial distribution of TM helices in the plane of the inner membrane leaflet (*x*,*y*) is conserved throughout the simulations. **(C)** 2D histograms of outer leaflet positions of pore-lining helices (TM1, 6, 8, 11, and 12) and TM2 in the squished conformation of the pore (blue) versus expanded pentagonal conformation (red). Helices in the blue-shaded region belong to half-channel 1, whereas those in the unshaded region belong to half-channel 2. **(D)** Top (extracellular) view of representative structures of the squished (blue) and pentagonal (red) conformations superimposed. Cl⁻ ions (green spheres) are overlaid from multiple simulation snapshots to indicate the location of the channel pore.

Figure 2 – figure supplement 1. Conservation of overall structure of ATP-bound

CFTR in MD simulations. Left: Distributions of RMSDs from the average of the MD

structural ensemble. Average values are indicated in the legend. Right: Distribution of

the distance between the centres of mass of the two NBDs (blue). The distances in

experimental structures of ATP-bound (PDB: 6MSM; red dashed line) and ATP-free

(PDB: 5UAK; green dashed line) states are shown for comparison.

Figure 3. Principal component analysis of the CFTR conformational ensemble in the TM region. (A) Distribution of TM conformations projected onto the plane of the first two principal components reveals 4 conformational states labelled α-δ. The cross indicates the location of the initial cryo-EM structure (PDB: 6MSM). Contour levels are evenly spaced values of probability density. **(B)** Spatial distribution of pore-lining helices (TM1, 6, 8, 11, 12) in the plane of the outer leaflet (x,y) for conformational states α-δ. Contour levels are evenly spaced values of probability density. Concentric circles are centered on the pore axis and spaced by 4 Å. **(C-D)** Porcupine representation (red) of eigenvectors **(C)** PC1 and **(D)** PC2 overlaid on the CFTR structure (cyan) to illustrate the collective movements in each dimension. The largest collective motion, PC1, captures the concerted movement of TM1, 2, and 11 from near-open states β and γ to the pentagonal pore conformation, α (see also Fig. 2C).

Figure 3 – figure supplement 1. Time evolution of the fractional population of conformational states α-δ indicating their relative stability on the timescale of the MD simulations.

Figure 3 – figure supplement 2. Supplementary information for PCA analysis. Left: fractional variance explained by each component. Right: cumulative fractional variance of principal components.

Figure 4. Structural features of the pore bottleneck. (A) Cartoon representation of

the bottleneck region in putative open state α. Pore-lining helices are shown in cyan

except for TM6 (tan) and TM11 (magenta). The blue surface encapsulating the interior

of the open pore was computed with HOLE2 (Smart et al., 1996). **(B)** Distributions of the

minimum radius (R_{min}) in the pore bottleneck. Pore radii are calculated with HOLE2.

Average values are indicated in the legend. **(C)** Distribution of the distances of pore-

lining helices (TM1, 6, 8, 11, and 12) to the centre of the pore for conformational states

α-δ. State α is unique for showing a unimodal distribution, consistent with the

symmetrical arrangement of pore-lining helices.

Figure 4 – figure supplement 1. Symmetry of TM helix conformations in the pore bottleneck. Distributions of the distances of the five pore-lining helices to the centre of the pore in conformational states α-δ. In contrast to states β-δ, the strong overlap between narrow TM helix distributions in state α reflects their nearly symmetrical arrangement.

Figure 4 – figure supplement 2. Detailed structural features of the pore bottleneck. (A-B) Distributions of distance between the Cα of residues R334 and T338 onTM6, and that of T1122 and S1118 on TM11 in the bottleneck region for conformational states α-δ. **(C-D)** Distributions of solvent accessible surface area (SASA) of bottleneck-lining residues T1115 and S1118 from TM11.

Figure 5. Cl⁻ permeation pathways through the pore bottleneck. (A) Side and **(B)**
top views of the bottleneck region in the putative open state α with overlaid Cl⁻ ions
(coloured spheres) showing three different Cl⁻ permeation routes. Blue: 1-6 pathway
between TM1 and 6; red: 1-12 pathway between TM1 and 12; green: intermediate
pathway. **(C)** Pore conformations in which Cl⁻ permeation occurs are indicated as scatter
points on the (PC1,PC2) plane, coloured by permeation pathway. **(D)** Breakdown of Cl⁻
permeation pathways followed in each conformational state (N: number of permeation
events) for (top) the full voltage range (-500 < U < 500 mV) and (bottom) the
physiological voltage range (-120 < U < 120 mV). The number of times each permeation
pathway was followed is indicated on top of the bar graphs.

Figure 5 – figure supplement 1. Projection of the two earlier trajectories in which Cl⁻

permeation events occurred in our previous study (Zeng et al., 2023) onto the

(PC1,PC2) plane. **(Left)** traces of the two trajectories coloured by timestep. **(Right)** The

conformations sampled in the two simulations indicated as red scatter points.

Conformations in which Cl⁻ translocations occurred are marked in black "x". The

snapshots from these two trajectories used to initiate the extended trajectories reported

in this work are marked in cyan "x".

Figure 6. Solvation of Cl⁻ by protein and water inside the pore of CFTR. (A) Cl⁻
binding sites inside the inner vestibule. Regions of high Cl⁻ density (blue mesh) indicate
the binding sites where Cl⁻ ions preferentially reside. Protein sidechains coming into
direct contact with Cl⁻ ions in its first solvation shell are shown; blue labels highlight
cationic residues among them. **(B-C)** Solvation environment of Cl⁻ ions that underwent
translocation. The bottleneck region is indicated as the gray shaded region ($-5 < z < 5$
Å). **(B)** The average numbers of polar groups from protein sidechains (black) and of
water molecules (red) in the first solvation shell of Cl⁻ ions (n_{solv}) along the channel axis
are shown together with their sum (green). The average number of non-polar protein
residues in the first solvation shell of Cl⁻ ions is indicated in purple. **(C)** Axial distribution
of protein residues making significant sidechain contributions to the solvation of Cl⁻ ions.
F337 is also shown. **(D-E)** Distribution of the number of water molecules (N_{H2O}) in the
first solvation shell of Cl⁻ ions in the bottleneck region of **(D)** the 1-6 pathway (blue) and
of **(E)** in the central pathway (green) compared to bulk water (gray).

Figure 6 – figure supplement 1. Solvation of Cl⁻ ions by permeation pathway in the

pore bottleneck. Statistics of solvation by water and protein of Cl⁻ ions that followed **(A-**

B) the 1-6 pathway and **(C-D)** the central pathway (combined 1-12 and intermediate

pathways) in the pore bottleneck. The grey shaded region indicates the pore bottleneck

$(-5 < z < 5$ Å). For details, see the caption of Figure 6.

Figure 7. Current-voltage (I-V) relationship for each of the four conformational states. (A) Entire voltage range sampled and **(B)** physiological and experimentally accessible range. Zero-intercept linear regression fits of data points in the positive (0 < U < +120 mV) and in the negative (-120 < U < 0 mV) TM voltages are shown for state α as red straight lines with conductance (slope) indicated. The gray shaded regions are bound by the experimental single channel conductance range of 6-10 pS (Sheppard and Welsh, 1999). Negative and positive currents indicate Cl⁻ efflux to and influx from the extracellular space, respectively.

Figure 8. Effect of R334 conformation on Cl⁻ permeation. (A-B) Snapshots of the

extracellular mouth of the pore with R334 in the **(A)** up and **(B)** down states. TM helices

6 (tan), 11 (magenta) and 12 (yellow) are shown together with loop ECL-6. In the down

conformation, R334 can form cation-π and π-π interactions with F337 and an ion pair

with Cl⁻ in the pore bottleneck. **(C-D)** Iso-surfaces reveal an increase in the average Cl⁻

ion density in the bottleneck region in state α from **(C)** the up state to **(D)** the down state

of R334. **(E)** The current-voltage relationship shows that, in putative open state α, the

down conformation of R334 (red) increases Cl⁻ conductance 5-fold in the physiological

voltage range (inset) compared to the up conformation (blue).

Figure 8 – figure supplement 1. The sidechain conformation of R134 does not

influence Cl⁻ permeation in CFTR. All analyses are restricted to the α state. **(A)** The

axial position of the centre of charge (Cζ atom) of the R134 sidechain samples two main

states "up" and "down". **(B-C)** Estimates of Cl⁻ conductance computed from these two

conformational states do not differ significantly.

Figure 8 – figure supplement 2. Distribution of distance (*d*) between centre of charge

(Cζ atom) of R334 and centre of the phenyl ring of F337 in conformational states R334-

up (blue) and R334-down (red).

Figure 9. Effect of salt-bridge interactions between R334 and ECL-6 loop residues

on Cl⁻ permeation. (A-B) Snapshots of the extracellular mouth of the pore showing the

dunked sidechain of R334 forming salt bridges with **(A)** E1124 and **(B)** E1126. **(C-D)**

Iso-surfaces reveal a greater Cl⁻ ion density in the extracellular mouth of the pore in

open state α when **(C)** the glutamate sidechains are disengaged from the central pore,

compared to **(D)** when they form a salt bridge with R334. **(E-F)** Top view of

representative conformations of ECL-6 (cyan) in the **(E)** loop-outward and **(F)** loop-

inward states. The helices lining the pentagonal pore are shown in gray. **(G)** The

current-voltage relationship shows that the salt-bridge interactions decrease Cl⁻

conductance in the R334-down, open state of CFTR, especially in the physiological

voltage range (inset).

Figure 9 – figure supplement 1. Analysis of charged sidechain arrangements in the extracellular mouth of the penta-helical pore (state α). (A) Axial distribution of the centres of charge of R334 (Cζ atom), E1124 and E1126 (Cδ atoms). Ion pairing between R334 and either glutamate residue is precluded when glutamate sidechains are oriented outwards. In the R334 vs E1126 contour plot (bottom left panel), the boxed region highlights the "down-out" state with R334 down and E1126 out. In the E1124 vs E1124 contour plot (bottom middle panel), the boxed region highlights the "out-out" state with both glutamate residues out. **(B)** Scatter plot of apparent Cl⁻ conductance (*g*) versus fractional population of R334 down without salt-bridge with ECL-6 glutamate residues [$f_{down-out-out}$, i.e. intersection of the two red boxes in **(A)**] at different TM voltages (indicated in mV in black). The circles are colour coded according to the sign of the voltage (yellow: positive; black: negative). The positive correlation shows that at negative voltages, the sidechain conformations of the ECL-6 loop glutamate residues and R334 modulate Cl⁻ conductance. At positive voltages, the systematic under-sampling of the R334-down and glutamates-out conformation provides a plausible explanation for the apparent inward rectification of CFTR in the simulations.

Figure 10. Structural differences between the ATP-unbound, closed state and the open state of human CFTR in the gate region. (A) Top and **(B)** side views of representative structures taken from MD simulations of ATP-free CFTR (left; black) and of the MD-open state of ATP-bound CFTR (right; red). To reach the open state, the outer segments of TM10 (cyan) and 11 (magenta) must separate to accommodate the intercalation of TM12 (yellow) between them. **(C)** 2D histograms of outer leaflet positions of pore-lining TM helices (1, 6, 8, 11, and 12) together with TM 2 and 10 from MD simulations of (black) ATP-free CFTR versus (red) MD-open state of CFTR. The largest movements of TM helices needed to reach the open state are indicated by arrows. Helices in the blue-shaded region belong to half-channel 1, whereas those in the unshaded region belong to half-channel 2. **(D)** Distribution of distances separating the centres of mass of the outer segments of TM10 and 11 in the ATP-free closed state (black) and in the four ATP-bound conformational states α-δ.

Figure 10 – figure supplement 1. NBD1-NBD2 separation in the 20 simulations of ATP-free CFTR. Time evolution of the distances separating the two halves of the catalytic site (orange) and of the degenerate site (blue). Dashed lines indicate the distances in ATP-bound, NBD-dimerized PDB structure 6MSM (red: catalytic site; blue: degenerate site). Background colors highlight the simulations in which (green) full dimerization; (caramel) partial dimerization; or (white) no dimerization occurred.

Figure 10 – figure supplement 2. Conservation of TM helix arrangement at the

extracellular end of the pore despite fast NBD dimerization in simulations of ATP-

free, closed state CFTR. (Left) Distribution of centre-of-mass distance between NBDs

showing an NBD-separated population (d > 40 Å) and NBD-dimerized population (d <

40 Å). Distributions of (x,y) positions of TM helices at the extracellular end **(middle)**

before and **(right)** after NBD-dimerization show that their relative arrangement is

conserved. Blue crosses in the 2D contour plots indicate the position of TM helices in

the ATP-free, unphosphorylated, and NBD-separated human CFTR (PDB: 5UAK) at the

start of the simulations.

Figure 10 – video 1. Conformational changes from ATP-free closed state to open

state illustrated by linear morph between the two states. Top view of the pore.

Opening of the channel involves reorganization of pore helices TM 12 (yellow) and 8

(green) relative to TM 1 (blue), 2 (red), 6 (tan), and 11 (magenta). TM11 separates from

TM10 (cyan) to accommodate TM12.

Figure 11. Analysis of changes in helical kinks underlying the transition between near-open and open states. (A) Cartoon representation of CFTR highlighting residues located at helical kinks on TM 1 (blue), 2 (red), and 11 (magenta) where structural changes occur upon channel opening. The sidechains of these residues (Hα atoms for glycine) are shown. The three helices completing the pore bottleneck are shown in dark gray. All other parts of the protein are translucent. **(B-G)** Analysis of helical kink magnitude and orientation. **(Left)** Distribution of kink angle θ (kink magnitude) and wobble or swivel angle ▢ (kink orientation) are shown for each of the four conformational states at **(B)** G126 on TM2; **(D)** W1098 on TM11; and **(F)** G85 on TM1. **(Right)** Contour plots showing the correlation between kink or swivel angles and principal component 1 (PC1), which captures conformational transitions between MD-closed and MD-open states (see Figure 3). Note that the uncertainty in kink orientation ▢ is high for small kink angles θ.

Figure 11 – figure supplement 1. Analysis of helical kink and orientation angles at **(A-B)** P99 on TM1 and **(C-D)** P140 on TM2. See the caption of Fig. 11 for details.

Figure 11 – video 1. Conformational changes from MD-closed state β to MD-open state α visualized using 50 randomly selected snapshots from each state (first 50 frames: state β, last 50 frames: state α). Note the changes in the conformation and orientation of TM helices 1 (blue), 2 (red), and 11 (magenta) relative to TM12 (yellow) and the other pore lining helices, TM 6 and 8 (grey).

References

References

Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, Lindahl E. 2015. GROMACS: High performance
molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX*
1–2:19–25. doi:10.1016/j.softx.2015.06.001

Amaral C, Carnevale V, Klein ML, Treptow W. 2012. Exploring conformational states of the bacterial
voltage-gated sodium channel NavAb via molecular dynamics simulations. *Proceedings of the*
National Academy of Sciences **109**:21336–21341. doi:10.1073/pnas.1218087109

Berger HA, Anderson MP, Gregory RJ, Thompson S, Howard PW, Maurer RA, Mulligan R, Smith AE, Welsh
MJ. 1991. Identification and regulation of the cystic fibrosis transmembrane conductance regulator-
generated chloride channel. *Journal of Clinical Investigation* **88**:1422–1431. doi:10.1172/JCI115450

Best RB, Zhu X, Shim J, Lopes PEM, Mittal J, Feig M, MacKerell AD. 2012. Optimization of the Additive
CHARMM All-Atom Protein Force Field Targeting Improved Sampling of the Backbone ϕ, ψ and Side-
Chain χ_1 and χ_2 Dihedral Angles. *J Chem Theory Comput* **8**:3257–3273. doi:10.1021/ct300400x

Bose SJ, Krainer G, Ng DRS, Schenkel M, Shishido H, Yoon JS, Haggie PM, Schlierf M, Sheppard DN, Skach
WR. 2020. Towards next generation therapies for cystic fibrosis: Folding, function and
pharmacology of CFTR. *Journal of Cystic Fibrosis* **19**:S25–S32. doi:10.1016/j.jcf.2019.12.009

Cai Z, Scott-Ward TS, Sheppard DN. 2003. Voltage-dependent gating of the cystic fibrosis transmembrane
conductance regulator Cl- channel. *J Gen Physiol* **122**:605–20. doi:10.1085/jgp.200308921

Cai Z, Sohma Y, Bompadre SG, Sheppard DN, Hwang T-C. 2011. Application of High-Resolution Single-
Channel Recording to Functional Studies of Cystic Fibrosis Mutants. pp. 419–441. doi:10.1007/978-
1-61779-117-8_27

Cerdan AH, Martin NÉ, Cecchini M. 2018. An Ion-Permeable State of the Glycine Receptor Captured by
Molecular Dynamics. *Structure* **26**:1555-1562.e4. doi:10.1016/j.str.2018.07.019

cftr2.org. n.d. The Clinical and Functional TRanslation of CFTR (CFTR2). *https://cftr2.org/*.

Chakrabarti N, Neale C, Payandeh J, Pai EF, Pomès R. 2010. An Iris-Like Mechanism of Pore Dilation in the
CorA Magnesium Transport System. *Biophys J* **98**:784–792. doi:10.1016/j.bpj.2009.11.009

Chamberlain AK, Faham S, Yohannan S, Bowie JU. 2003. Construction of helix-bundle membrane
proteins. pp. 19–46. doi:10.1016/S0065-3233(03)63002-0

Chen J, Xu W, Sheppard DN. 2017. Altering intracellular pH reveals the kinetic basis of intraburst gating in
the CFTR Cl⁻ channel. *J Physiol* **595**:1059–1076. doi:10.1113/JP273205

Cheng SH, Rich DP, Marshall J, Gregory RJ, Welsh MJ, Smith AE. 1991. Phosphorylation of the R domain
by cAMP-dependent protein kinase regulates the CFTR chloride channel. *Cell* **66**:1027–1036.
doi:10.1016/0092-8674(91)90446-6

Choudhury K, Delemotte L. 2023. Modulation of Pore Opening of Eukaryotic Sodium Channels by π-
Helices in S6. *J Phys Chem Lett* **14**:5876–5881. doi:10.1021/acs.jpclett.3c00803

Choudhury K, Howard RJ, Delemotte L. 2023. An α–π transition in S6 shapes the conformational cycle of
the bacterial sodium channel NavAb. *Journal of General Physiology* **155**.
doi:10.1085/jgp.202213214

Cooley RB, Arp DJ, Karplus PA. 2010. Evolutionary Origin of a Secondary Structure: π-Helices as Cryptic
but Widespread Insertional Variations of α-Helices That Enhance Protein Functionality. *J Mol Biol*
404:232–246. doi:10.1016/j.jmb.2010.09.034

Corradi V, Gu R-X, Vergani P, Tieleman DP. 2018. Structure of Transmembrane Helix 8 and Possible
Membrane Defects in CFTR. *Biophys J* **114**:1751–1754. doi:10.1016/j.bpj.2018.03.003

Cottrill KA, Farinha CM, McCarty NA. 2020. The bidirectional relationship between CFTR and lipids.
Commun Biol **3**:179. doi:10.1038/s42003-020-0909-1

Dalal V, Arcario MJ, Petroff JT, Tan BK, Dietzen NM, Rau MJ, Fitzpatrick JAJ, Brannigan G, Cheng WWL.
2024. Lipid nanodisc scaffold and size alter the structure of a pentameric ligand-gated ion channel.
Nat Commun **15**:25. doi:10.1038/s41467-023-44366-w

Dämgen MA, Biggin PC. 2020. A Refined Open State of the Glycine Receptor Obtained via Molecular
Dynamics Simulations. *Structure* **28**:130-139.e2. doi:10.1016/j.str.2019.10.019

Darden T, York D, Pedersen L. 1993. Particle mesh Ewald: An N ·log(N) method for Ewald sums in large
systems. *J Chem Phys* **98**:10089–10092. doi:10.1063/1.464397

Dassa E, Bouige P. 2001. The ABC of ABCs: a phylogenetic and functional classification of ABC systems in
living organisms. *Res Microbiol* **152**:211–229. doi:10.1016/S0923-2508(01)01194-9

Decaestecker K, Decaestecker E, Castellani C, Jaspers M, Cuppens H, De Boeck K. 2004.
Genotype/phenotype correlation of the G85E mutation in a large cohort of cystic fibrosis patients.
European Respiratory Journal **23**:679–684. doi:10.1183/09031936.04.00014804

Deng Z, Maksaev G, Rau M, Xie Z, Hu H, Fitzpatrick JAJ, Yuan P. 2020. Gating of human TRPV3 in a lipid
bilayer. *Nat Struct Mol Biol* **27**:635–644. doi:10.1038/s41594-020-0428-2

Elborn JS. 2016. Cystic fibrosis. *The Lancet* **388**:2519–2531. doi:10.1016/S0140-6736(16)00576-6

Essmann U, Perera L, Berkowitz ML, Darden T, Lee H, Pedersen LG. 1995. A smooth particle mesh Ewald
method. *J Chem Phys* **103**:8577–8593. doi:10.1063/1.470117

Farkas B, Tordai H, Padányi R, Tordai A, Gera J, Paragi G, Hegedűs T. 2020. Discovering the chloride
pathway in the CFTR channel. *Cellular and Molecular Life Sciences* **77**:765–778.
doi:10.1007/s00018-019-03211-4

Fatehi M, Linsdell P. 2009. Novel Residues Lining the CFTR Chloride Channel Pore Identified by Functional
Modification of Introduced Cysteines. *Journal of Membrane Biology* **228**:151–164.
doi:10.1007/s00232-009-9167-3

Fiedorczuk K, Chen J. 2022a. Mechanism of CFTR correction by type I folding correctors. *Cell* **185**:158-
168.e11. doi:10.1016/j.cell.2021.12.009

Fiedorczuk K, Chen J. 2022b. Molecular structures reveal synergistic rescue of Δ508 CFTR by Trikafta
modulators. *Science (1979)* **378**:284–290. doi:10.1126/science.ade2216

Fiedorczuk K, Iordanov I, Mihályi C, Szollosi A, Csanády L, Chen J. 2024. The structures of protein kinase A
in complex with CFTR: Mechanisms of phosphorylation and noncatalytic activation. *Proceedings of*
the National Academy of Sciences **121**. doi:10.1073/pnas.2409049121

Fowler PW, Sansom MSP. 2013. The pore of voltage-gated potassium ion channels is strained when
closed. *Nat Commun* **4**:1872. doi:10.1038/ncomms2858

Gao X, Hwang T-C. 2015. Localizing a gate in CFTR. *Proceedings of the National Academy of Sciences*
112:2461–2466. doi:10.1073/pnas.1420676112

Ge N, Muise CN, Gong X, Linsdell P. 2004. Direct Comparison of the Functional Roles Played by Different
Transmembrane Regions in the Cystic Fibrosis Transmembrane Conductance Regulator Chloride
Channel Pore. *Journal of Biological Chemistry* **279**:55283–55289. doi:10.1074/jbc.M411935200

Gong X, Linsdell P. 2004. Maximization of the rate of chloride conduction in the CFTR channel pore by
ion–ion interactions. *Arch Biochem Biophys* **426**:78–82. doi:10.1016/j.abb.2004.03.033

Gonzalez-Gutierrez G, Wang Y, Cymes GD, Tajkhorshid E, Grosman C. 2017. Chasing the open-state
structure of pentameric ligand-gated ion channels. *Journal of General Physiology* **149**:1119–1138.
doi:10.1085/jgp.201711803

Gowers R, Linke M, Barnoud J, Reddy T, Melo M, Seyler S, Domański J, Dotson D, Buchoux S, Kenney I,
Beckstein O. 2016. MDAnalysis: A Python Package for the Rapid Analysis of Molecular Dynamics
Simulations. pp. 98–105. doi:10.25080/Majora-629e541a-00e

Gu R-X, de Groot BL. 2020. Lipid-protein interactions modulate the conformational equilibrium of a
potassium channel. *Nat Commun* **11**:2162. doi:10.1038/s41467-020-15741-8

Gumbart J, Khalili-Araghi F, Sotomayor M, Roux B. 2012. Constant electric field simulations of the
membrane potential illustrated with simple systems. *Biochimica et Biophysica Acta (BBA) -*
Biomembranes **1818**:294–302. doi:10.1016/j.bbamem.2011.09.030

Hess B. 2008. P-LINCS: A Parallel Linear Constraint Solver for Molecular Simulation. *J Chem Theory*
Comput **4**:116–122. doi:10.1021/ct700200b

Hoover WG. 1985. Canonical dynamics: Equilibrium phase-space distributions. *Phys Rev A (Coll Park)*
31:1695–1697. doi:10.1103/PhysRevA.31.1695

Huang J, MacKerell AD. 2013. CHARMM36 all-atom additive protein force field: Validation based on
comparison to NMR data. *J Comput Chem* **34**:2135–2145. doi:10.1002/jcc.23354

Humphrey W, Dalke A, Schulten K. 1996. VMD: Visual molecular dynamics. *J Mol Graph* **14**:33–38.
doi:10.1016/0263-7855(96)00018-5

Hwang T-C, Yeh J-T, Zhang J, Yu Y-C, Yeh H-I, Destefano S. 2018. Structural mechanisms of CFTR function
and dysfunction. *Journal of General Physiology* **150**:539–570. doi:10.1085/jgp.201711946

Ikuma M, Welsh MJ. 2000. Regulation of CFTR Cl- channel gating by ATP binding and hydrolysis.
Proceedings of the National Academy of Sciences **97**:8675–8680. doi:10.1073/pnas.140220597

Ishiguro H, Steward MC, Naruse S, Ko SBH, Goto H, Case RM, Kondo T, Yamamoto A. 2009. CFTR
Functions as a Bicarbonate Channel in Pancreatic Duct Cells. *Journal of General Physiology*
133:315–326. doi:10.1085/jgp.200810122

Jo S, Kim T, Iyer VG, Im W. 2008. CHARMM-GUI: A web-based graphical user interface for CHARMM. *J*
Comput Chem **29**:1859–1865. doi:10.1002/jcc.20945

Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. 1983. Comparison of simple potential
functions for simulating liquid water. *J Chem Phys* **79**:926–935. doi:10.1063/1.445869

Klauda JB, Venable RM, Freites JA, O'Connor JW, Tobias DJ, Mondragon-Ramirez C, Vorobyov I, MacKerell
AD, Pastor RW. 2010. Update of the CHARMM All-Atom Additive Force Field for Lipids: Validation on
Six Lipid Types. *J Phys Chem B* **114**:7830–7843. doi:10.1021/jp101759q

Lemkul JA. 2020. Pairwise-additive and polarizable atomistic force fields for molecular dynamics
simulations of proteins. pp. 1–71. doi:10.1016/bs.pmbts.2019.12.009

Levring J, Chen J. 2024. Structural identification of a selectivity filter in CFTR. *Proceedings of the National*
Academy of Sciences **121**. doi:10.1073/pnas.2316673121

Levring J, Terry DS, Kilic Z, Fitzgerald G, Blanchard SC, Chen J. 2023. CFTR function, pathology and
pharmacology at single-molecule resolution. *Nature* **616**:606–614. doi:10.1038/s41586-023-05854-
7

Linsdell P. 2018. Cystic fibrosis transmembrane conductance regulator (CFTR): Making an ion channel out
of an active transporter structure. *Channels* **12**:284–290. doi:10.1080/19336950.2018.1502585

Linsdell P. 2017. Architecture and functional properties of the CFTR channel pore. *Cellular and Molecular*
Life Sciences **74**:67–83. doi:10.1007/s00018-016-2389-5

Linsdell P. 2016. Anion conductance selectivity mechanism of the CFTR chloride channel. *Biochimica et*
Biophysica Acta (BBA) - Biomembranes **1858**:740–747. doi:10.1016/j.bbamem.2016.01.009

Linsdell P, Evagelidis A, Hanrahan JW. 2000. Molecular Determinants of Anion Selectivity in the Cystic
Fibrosis Transmembrane Conductance Regulator Chloride Channel Pore. *Biophys J* **78**:2973–2982.
doi:10.1016/S0006-3495(00)76836-6

Linsdell P, Hanrahan JW. 1998. Adenosine Triphosphate–dependent Asymmetry of Anion Permeation in
the Cystic Fibrosis Transmembrane Conductance Regulator Chloride Channel. *J Gen Physiol*
111:601–614. doi:10.1085/jgp.111.4.601

Linsdell P, Tabcharani JA, Rommens JM, Hou Y-X, Chang X-B, Tsui L-C, Riordan JR, Hanrahan JW. 1997.
Permeability of Wild-Type and Mutant Cystic Fibrosis Transmembrane Conductance Regulator
Chloride Channels to Polyatomic Anions. *J Gen Physiol* **110**:355–364. doi:10.1085/jgp.110.4.355

Linsdell P, Zheng S-X, Hanrahan JW. 1998. Non-pore lining amino acid side chains influence anion
selectivity of the human CFTR Cl – channel expressed in mammalian cell lines. *J Physiol* **512**:1–16.
doi:10.1111/j.1469-7793.1998.001bf.x

Liu F, Zhang Z, Csanády L, Gadsby DC, Chen J. 2017. Molecular Structure of the Human CFTR Ion Channel.
Cell **169**:85-95.e8. doi:10.1016/j.cell.2017.02.024

Liu F, Zhang Z, Levit A, Levring J, Touhara KK, Shoichet BK, Chen J. 2019. Structural identification of a
hotspot on CFTR for potentiation. *Science (1979)* **364**:1184–1188. doi:10.1126/science.aaw7611

Liu Y, Wang D-K, Chen L-M. 2012. The Physiology of Bicarbonate Transporters in Mammalian
Reproduction1. *Biol Reprod* **86**. doi:10.1095/biolreprod.111.096826

Lomize MA, Pogozheva ID, Joo H, Mosberg HI, Lomize AL. 2012. OPM database and PPM web server:
resources for positioning of proteins in membranes. *Nucleic Acids Res* **40**:D370–D376.
doi:10.1093/nar/gkr703

Lukacs GL, Verkman AS. 2012. CFTR: folding, misfolding and correcting the ΔF508 conformational defect.
Trends Mol Med **18**:81–91. doi:10.1016/j.molmed.2011.10.003

Mackerell AD. 2004. Empirical force fields for biological macromolecules: Overview and issues. *J Comput
Chem* **25**:1584–1604. doi:10.1002/jcc.20082

MacKerell AD, Bashford D, Bellott M, Dunbrack RL, Evanseck JD, Field MJ, Fischer S, Gao J, Guo H, Ha S,
Joseph-McCarthy D, Kuchnir L, Kuczera K, Lau FTK, Mattos C, Michnick S, Ngo T, Nguyen DT,
Prodhom B, Reiher WE, Roux B, Schlenkrich M, Smith JC, Stote R, Straub J, Watanabe M,
Wiórkiewicz-Kuczera J, Yin D, Karplus M. 1998. All-Atom Empirical Potential for Molecular Modeling
and Dynamics Studies of Proteins. *J Phys Chem B* **102**:3586–3616. doi:10.1021/jp973084f

Michaud-Agrawal N, Denning EJ, Woolf TB, Beckstein O. 2011. MDAnalysis: A toolkit for the analysis of
molecular dynamics simulations. *J Comput Chem* **32**:2319–2327. doi:10.1002/jcc.21787

Negoda A, Hogan MS, Cowley EA, Linsdell P. 2019. Contribution of the eighth transmembrane segment
to the function of the CFTR chloride channel pore. *Cellular and Molecular Life Sciences* **76**:2411–
2423. doi:10.1007/s00018-019-03043-2

Nosé S. 1984. A molecular dynamics method for simulations in the canonical ensemble. *Mol Phys*
52:255–268. doi:10.1080/00268978400101201

Nosé S, Klein ML. 1983. Constant pressure molecular dynamics for molecular systems. *Mol Phys*
50:1055–1076. doi:10.1080/00268978300102851

Páll S, Hess B. 2013. A flexible algorithm for calculating pair interactions on SIMD architectures. *Comput
Phys Commun* **184**:2641–2650. doi:10.1016/j.cpc.2013.06.003

Parrinello M, Rahman A. 1980. Crystal Structure and Pair Potentials: A Molecular-Dynamics Study. *Phys
Rev Lett* **45**:1196–1199. doi:10.1103/PhysRevLett.45.1196

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P, Weiss R,
Dubourg V, Vanderplas J, Passos A, Cournapeau D, Brucher M, Perrot M, Duchesnay É. 2011. Scikit-
learn: Machine Learning in Python. *Journal of Machine Learning Research* **12**:2825–2830.

Riordan, Rommens J, Kerem B, Alon N, Rozmahel R, Grzelczak Z, Zielenski J, Lok S, Plavsic N, Chou J, Et Al.
1989. Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA.
Science (1979) **245**:1066–1073. doi:10.1126/science.2475911

Roux B. 2008. The Membrane Potential and its Representation by a Constant Electric Field in Computer Simulations. *Biophys J* **95**:4205–4216. doi:10.1529/biophysj.108.136499

Saint-Criq V, Gray MA. 2017. Role of CFTR in epithelial physiology. *Cellular and Molecular Life Sciences* **74**:93–115. doi:10.1007/s00018-016-2391-y

Sakoe H, Chiba S. 1978. Dynamic programming algorithm optimization for spoken word recognition. *IEEE Trans Acoust* **26**:43–49. doi:10.1109/TASSP.1978.1163055

Sheppard DN, Rich DP, Ostedgaard LS, Gregory RJ, Smith AE, Welsh MJ. 1993. Mutations in CFTR associated with mild-disease-form Cl- channels with altered pore properties. *Nature* **362**:160–164. doi:10.1038/362160a0

Sheppard DN, Travis SM, Ishihara H, Welsh MJ. 1996. Contribution of Proline Residues in the Membrane-spanning Domains of Cystic Fibrosis Transmembrane Conductance Regulator to Chloride Channel Function. *Journal of Biological Chemistry* **271**:14995–15001. doi:10.1074/jbc.271.25.14995

Sheppard DN, Welsh MJ. 1999. Structure and Function of the CFTR Chloride Channel. *Physiol Rev* **79**:S23–S45. doi:10.1152/physrev.1999.79.1.S23

Shrivastava IH, Sansom MSP. 2000. Simulations of Ion Permeation Through a Potassium Channel: Molecular Dynamics of KcsA in a Phospholipid Bilayer. *Biophys J* **78**:557–570. doi:10.1016/S0006-3495(00)76616-1

Simon MA, Csanády L. 2021. Molecular pathology of the R117H cystic fibrosis mutation is explained by loss of a hydrogen bond. *Elife* **10**. doi:10.7554/eLife.74693

Smart OS, Neduvelil JG, Wang X, Wallace BA, Sansom MSP. 1996. HOLE: A program for the analysis of the pore dimensions of ion channel structural models. *J Mol Graph* **14**:354–360. doi:10.1016/S0263-7855(97)00009-X

Smith JJ, Travis SM, Greenberg EP, Welsh MJ. 1996. Cystic Fibrosis Airway Epithelia Fail to Kill Bacteria Because of Abnormal Airway Surface Fluid. *Cell* **85**:229–236. doi:10.1016/S0092-8674(00)81099-5

S.P. Sansom M, Weinstein H. 2000. Hinges, swivels and switches: the role of prolines in signalling via transmembrane α-helices. *Trends Pharmacol Sci* **21**:445–451. doi:10.1016/S0165-6147(00)01553-4

Tavenard R, Faouzi J, Vandewiele G, Divo F, Androz G, Holtz C, Payne M, Yurchak R, Rußwurm M, Kolar K, Woods E. 2020. Tslearn, A Machine Learning Toolkit for Time Series Data. *Journal of Machine Learning Research* **21**:1–6.

Thomas C, Aller SG, Beis K, Carpenter EP, Chang G, Chen L, Dassa E, Dean M, Duong Van Hoa F, Ekiert D, Ford R, Gaudet R, Gong X, Holland IB, Huang Y, Kahne DK, Kato H, Koronakis V, Koth CM, Lee Y, Lewinson O, Lill R, Martinoia E, Murakami S, Pinkett HW, Poolman B, Rosenbaum D, Sarkadi B, Schmitt L, Schneider E, Shi Y, Shyng S, Slotboom DJ, Tajkhorshid E, Tieleman DP, Ueda K, Váradi A, Wen P, Yan N, Zhang P, Zheng H, Zimmer J, Tampé R. 2020. Structural and functional diversity calls for a new classification of ABC transporters. *FEBS Lett* **594**:3767–3775. doi:10.1002/1873-3468.13935

Tordai H, Leveles I, Hegedűs T. 2017. Molecular dynamics of the cryo-EM CFTR structure. *Biochem*
Biophys Res Commun **491**:986–993. doi:10.1016/j.bbrc.2017.07.165

Verlet L. 1967. Computer "Experiments" on Classical Fluids. I. Thermodynamical Properties of Lennard-
Jones Molecules. *Physical Review* **159**:98–103. doi:10.1103/PhysRev.159.98

Visiers I, Braunheim BB, Weinstein H. 2000. Prokink: a protocol for numerical evaluation of helix
distortions by proline. *Protein Engineering, Design and Selection* **13**:603–606.
doi:10.1093/protein/13.9.603

Wagner K, Greil I, Schneditz P, Rosenkranz W. 1994. A New Missense Mutation G126D in Exon 4 of the
Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) Gene. *Hum Hered* **44**:56–57.
doi:10.1159/000154190

Wang W, Linsdell P. 2012. Relative Movements of Transmembrane Regions at the Outer Mouth of the
Cystic Fibrosis Transmembrane Conductance Regulator Channel Pore during Channel Gating.
Journal of Biological Chemistry **287**:32136–32146. doi:10.1074/jbc.M112.385096

Zeng ZW, Linsdell P, Pomès R. 2023. Molecular dynamics study of Cl– permeation through cystic fibrosis
transmembrane conductance regulator (CFTR). *Cellular and Molecular Life Sciences* **80**:51.
doi:10.1007/s00018-022-04621-7

Zhang Z, Chen J. 2016. Atomic Structure of the Cystic Fibrosis Transmembrane Conductance Regulator.
Cell **167**:1586-1597.e9. doi:10.1016/j.cell.2016.11.014

Zhang Z, Liu F, Chen J. 2018. Molecular structure of the ATP-bound, phosphorylated human CFTR.
Proceedings of the National Academy of Sciences **115**:12757–12762. doi:10.1073/pnas.1815287115

Zhang Z, Liu F, Chen J. 2017. Conformational Changes of CFTR upon Phosphorylation and ATP Binding.
Cell **170**:483-491.e8. doi:10.1016/j.cell.2017.06.041

Zhang Z-R, Song B, McCarty NA. 2005. State-dependent Chemical Reactivity of an Engineered Cysteine
Reveals Conformational Changes in the Outer Vestibule of the Cystic Fibrosis Transmembrane
Conductance Regulator. *Journal of Biological Chemistry* **280**:41997–42003.
doi:10.1074/jbc.M510242200

1103