Using FoxP2 to Distinguish Direct and Indirect Basal Ganglia Pathways for Vocal Learning in Songbirds

Aditi Jagannathan[1,2], Mira Nigudkar[1,3], Sarah W Bottjer[1]

[1]Section of Neurobiology, University of Southern California, Los Angeles, CA 90089 USA

Correspondence to: Sarah W. Bottjer, Section of Neurobiology, University of Southern California, 3641 Watt Way, Los Angeles, CA 90089 (bottjer@usc.edu)

Abbreviated title: FoxP2 as a marker for IND pallidal cells

Number of pages: 35
Number of figures: 7
Number of tables: 3

Number of words, abstract: 244
Number of words, introduction: 649
Number of words, discussion: 1499

Conflicts of Interest

The authors declare no competing financial interests.

Acknowledgements

This research was supported by NSF grant 1940957. We thank Dr. Jason Junge and Dr. Arkadi Shwartz of the Translational Imaging Center at the USC Michelson Center for Convergent Bioscience for their assistance in training. We thank Dr. Hugh C. Hemmings for the gift of antibody to DARPP-32. [2]Current affiliation: Netherlands Institute for Neuroscience, Amsterdam, The Netherlands 1105 BA; [3]Current affiliation: Loyola University Chicago, Stritch School of Medicine, Chicago, IL 60301 USA

ABSTRACT

The cortico-basal ganglia pathways that mediate vocal learning in zebra finches (*Taeniopygia guttata*) are localized in parallel circuits formed by CORE and SHELL subregions (Fig. 1). These circuits traverse a cortical region (LMAN), a specialized region of the basal ganglia essential for vocal learning (Area X), and a region of dorsomedial thalamus (DLM). Area X includes functional analogs of mammalian direct and indirect pathways that may have opposing effects and thereby increase or inhibit thalamic activity respectively. Direct pallidal neurons of Area X send projections to DLM, whereas indirect pallidal neurons form intrinsic connections onto DLM-projecting neurons. Expression of the transcription factor FoxP2 in the basal ganglia is necessary for normal vocal learning and production in both humans and songbirds. We used tract-tracing techniques to label direct pallidal Area X→DLM projection neurons and immunohistochemical techniques to label neurons expressing the transcription factor FoxP2 in adult and juvenile male zebra finches. Our results showed that DLM-projecting neurons in Area X did not express FoxP2 in either adults or juveniles. We also observed a population of large neurons that expressed FoxP2 but were not retrogradely-labeled from DLM. A putative marker of striatal neurons (DARPP-32) did not co-localize with FoxP2 in most of these large neurons, suggesting that they form a class of indirect pallidal neurons. These findings offer FoxP2 as a possible marker for indirect pallidal neurons and support the existence of different subpopulations of neurons that correspond to direct and indirect pathways within Area X.

Key words: vocal learning, Area X, thalamus, FoxP2, pallidal

Significance statement:
Area X is a localized region of the basal ganglia in songbirds that mediates vocal learning and behavior and includes CORE and SHELL subregions. Some pallidal neurons in Area X project to thalamus, forming a direct pathway; a separate class of Area X pallidal neurons form local contacts onto direct neurons, forming an indirect pathway. FoxP2 is a transcription factor that is necessary for normal vocal learning and production in both humans and songbirds. We present evidence in adult and juvenile zebra finches that direct thalamus-projecting pallidal neurons do not express FoxP2 while local indirect pallidal neurons do express FoxP2. Thus, large FoxP2-positive neurons appear to form a class of indirect pallidal neurons.

1. INTRODUCTION

The basal ganglia in songbirds includes a specialized region essential for vocal learning and behavior called Area X, which receives input from two sensorimotor cortical regions that also regulate song-related functions: HVC and LMAN (Nottebohm et al., 1976; Bottjer et al., 1989; Sohrabji et al., 1990; Scharff and Nottebohm, 1991; Nixdorf-Bergweiler et al., 1995; Vates and Nottebohm, 1995; Sanchez-Valpuesta et al., 2019). In addition, Area X receives cerebellar inputs via thalamic nuclei adjacent to DLM (Person et al., 2008; Nicholson et al., 2018; Pidoux et al., 2018). Area X sends inhibitory projections to a dorsal thalamic nucleus (DLM), which projects back to LMAN (Nixdorf-Bergweiler et al., 1995; Vates and Nottebohm, 1995; Luo and Perkel, 1999b; Luo et al., 2001). The Area X→DLM→LMAN pathway consists of CORE and SHELL subregions, forming parallel topographically organized recurrent circuits (Fig. 1a) (Bottjer et al., 1989; Johnson and Bottjer, 1992; Johnson et al., 1995; Iyengar et al., 1999; Luo et al., 2001; Gale et al., 2008; Person et al., 2008). Both CORE and SHELL loops are essential for learning (Bottjer and Altenau, 2010; Achiro and Bottjer, 2013; Achiro et al., 2017).

The output of the CORE pathway from LMAN to vocal motor cortex (RA) drives song production in juvenile birds (Bottjer et al., 1984; Scharff and Nottebohm, 1991; Aronov et al., 2008). In contrast, the output of LMAN-SHELL projects to a region of motor cortex (AId) adjacent to RA that lacks direct motor targets; AId neurons make a strong projection to a dorsal thalamic zone which forms a transcortical loop that converges with information from Area X in DLM (Fig. 1a) (Foster et al., 1997; Bottjer et al., 2000). Lesions of AId disrupt accurate imitation of the tutor song but do not impact vocal motor output in juvenile songbirds (Bottjer and Altenau, 2010). Therefore, SHELL circuitry does not drive vocal motor output, but may play a cognitive-limbic role in evaluating vocal output. The organization of CORE and SHELL loops is highly similar to sensorimotor and associative-limbic circuits of mammalian basal ganglia (Yin and Knowlton, 2006; Graybiel, 2008; Redgrave et al., 2010).

Area X contains analogs of both direct and indirect basal ganglia pathways (Farries et al., 2005; Goldberg et al., 2010). In the direct (DIR) pathway, pallidal neurons form calyceal inhibitory synapses on DLM neurons (Fig. 1b) (Luo and Perkel, 1999a; Carrillo and Doupe, 2004; Reiner et al., 2004; Gale et al., 2008; Person et al., 2008). A separate population of local pallidal neurons in Area X makes axonal contacts with DIR DLM-projecting neurons, forming an indirect (IND) pathway. These local neurons express GAD, and DLM-projecting neurons are surrounded by inhibitory GAD+ processes (Luo and Perkel, 1999b), suggesting that this non-

projecting pallidal population forms intrinsic inhibitory projections onto DIR neurons of Area X (Fig. 1b).

Because dopamine receptors do not provide a marker for direct versus indirect neurons in songbirds (Kubikova et al., 2010), the transcription factor FoxP2 may serve as one possible marker to distinguish between direct and indirect pallidal neurons in Area X. FoxP2 is essential for human vocal development, with mutations leading to speech disorders such as childhood apraxia (den Hoed and Fisher, 2020). Likewise, disruptions of the FoxP2 gene disrupt vocal learning and production in songbirds (Haesler et al., 2007; Shi et al., 2018; Xiao et al., 2021). Single nucleus RNA sequencing showed that a cluster of cells in Area X identified as pallidal formed two subgroups: one expressed the gene for enkephalin (ENK) and the other expressed the gene for FoxP2 (Xiao et al., 2021). Because ENK is expressed in DLM-projecting pallidal neurons in songbirds (Carrillo and Doupe, 2004), this pattern suggested that the direct neurons that project to DLM would not express FoxP2, while the indirect non-projecting pallidal neurons would be FoxP2-positive (Fig. 1b). To test this idea, we used tract-tracing techniques to label direct pallidal Area X→DLM projection neurons and immunohistochemical techniques to label neurons expressing FoxP2.

2. MATERIALS AND METHODS

2.1 Subjects

All animal procedures were performed in accordance with national regulatory guidelines and followed protocols approved by the Animal Care and Use Committee at the University of Southern California. Birds were bred in group aviaries in our colony and remained with their parents until at least 43-47 days post-hatch (dph), which is beyond the age of adequate tutor song exposure and social experience (Immelmann, 1969; Böhner, 1983; Eales, 1985; Clayton, 1987; Böhner, 1990; Catchpole and Slater, 1995; Mann and Slater, 1995; Roper and Zann, 2006). Birds were moved to sex-specific holding cages in the colony after ~45-50 dph. Surgeries involving intracranial injections of a fluorescent tracer were performed on adult male birds >90 dph (age range: 91-182, n = 4) and juvenile male birds 45 ± 2 dph (n = 3). Following surgery, birds were placed in individual holding cages located in the main colony for three days before being perfused (see below).

2.2 Alexa Fluor 488 Injections

Birds were anesthetized with 1.5-1.7% isoflurane (inhalation) and placed in a stereotaxic apparatus. Stereotaxic coordinates were used to target the location of the thalamic nucleus

DLM (dorsolateral nucleus of the medial thalamus; Fig. 1). The head angle was set as follows: we measured the depth of the skull at the bifurcation of the mid-sagittal sinus and at the midline 5 mm anterior to the bifurcation; the head angle was adjusted so that the difference in depth between these two locations was 2.2 ± 0.1. DLM injection coordinates were approximately 0.6 to 0.8 mm anterior and 0.9 to 1.1 mm lateral to the bifurcation of the mid-sagittal sinus at a depth of 4.5 mm. Birds received unilateral left iontophoretic injections of dextran labeled with Alexa Fluor 488 (MW 10,000, 10% in sterile-filtered PBS, ThermoFisher #D-22910) into DLM by applying pulses of current through a silver wire (3-5 µA, 6 sec on/6 sec off) for 20 minutes through a glass pipette (20-30 µm outer diameter). Injections sometimes extended beyond the borders of DLM, but we analyzed retrograde label only within Area X-CORE and SHELL regions. Birds were anesthetized with an injection of 0.04 - 0.06 mL 5% Euthasol 72 hours following surgery and perfused trans-cardially with 0.7% saline followed by 4% periodate-lysine-paraformaldehyde (PLP) fixative (4% para in 0.1 M PB with 0.01 M sodium periodate and 0.1 M L-lysine; pH 7.8). Brains were removed and post-fixed in 4% paraformaldehyde overnight before being cryo-protected in 25% buffered sucrose solution with 0.005% sodium azide. Brains were frozen sectioned at a thickness of 40 µm in the coronal plane; sections were serially collected in PBS into separate series as described below.

2.3 Immunohistochemistry

2.3.1 Double Label

Sections from four adult birds and one juvenile bird were stained for FoxP2 and labeled with retrograde Alexa Fluor 488 dye from DLM. Three separate series were processed as follows: Series 1 was stained using rabbit polyclonal anti-FoxP2 (Abcam Cat# ab16046, RRID:AB_2107107), Series 2 was stained using goat polyclonal anti-FoxP2 (Abcam Cat# ab1307, RRID:AB_1268914), and Series 3 was counterstained with thionin for Nissl and coverslipped with Permount (Fisher Scientific). The ab1307 antibody was raised against a synthetic peptide corresponding to amino acids 703-715 (C-terminal) of human FoxP2 (specificity has been described previously, Thompson et al., 2013). The ab16046 antibody was generated using a synthetic peptide corresponding to human FoxP2 aa 700 to the C-terminus conjugated to keyhole limpet haemocyanin (specificity has been described previously, Campbell et al., 2009).

Sections were stained using standard immunohistochemistry techniques. Following three washes in 0.02 M PBS, Series 1 and 2 were blocked in 5% Normal Goat Serum or 5% Normal Rabbit Serum, respectively, for 30-40 minutes and then incubated in the primary

antibody (1:1,000) overnight at room temperature. Sections were again triple-washed with PBS and incubated in the appropriate secondary antibody (anti-rabbit-Alexa 594, 1:200, ThermoFisher, #A-11012, made in goat for Series 1; anti-goat-Alexa 594, 1:200, ThermoFisher, #A-11080, made in rabbit for Series 2) for 1 hour. Sections of both series were given three final washes in PBS with the last wash containing 1 µL DAPI (1:10,000, gift from Dr. Jason Junge) before being mounted onto gelatin-coated glass slides. Sections were allowed to air dry before being coverslipped with Prolong Diamond Antifade (ThermoFisher, #P36970).

2.3.2 Triple Label

DARPP-32 is a phosphoprotein that is expressed in virtually all medium spiny neurons of mammals (Ouimet et al., 1998), although it has been reported to be immunohistochemically detectable in only 30-60% of spiny neurons in birds (Reiner et al., 1998; Reiner et al., 2004; Singh and Iyengar, 2019). In order to test the identity of cells labeled by FoxP2, we performed triple labeling for FoxP2, DARPP-32, and the retrograde label from DLM in two juvenile birds. During prior analysis of double-labeled tissue, we observed that the anti-FoxP2 used in Series 1 (ab16046) resulted in a higher density of labeled cells compared to the anti-FoxP2 used in Series 2 (ab1307). We therefore created a cocktail of rabbit polyclonal anti-FoxP2 (ab16046, 1:1,000) and mouse monoclonal anti-DARPP-32 (1:30,000, C24-6a, gift from Dr. Hugh C. Hemmings, specificity validated in Hemmings and Greengard (1986) and see Durstewitz et al. (1998) for pigeon data); alternate sections were stained for Nissl. The same blocking and primary staining procedures outlined for Series 1 above were followed before incubating sections in a secondary antibody cocktail containing anti-rabbit Alexa 594 (1:200, ThermoFisher, #A11012, made in goat) and anti-mouse Alexa 647 (1:200, ThermoFisher, #A-21235, made in goat). All markers used in this study are outlined in Table 1.

2.4 Data Analysis

2.4.1 Overlap Analysis of FoxP2 and Retrogradely-Labeled Neurons in Area X

Patterns of fluorescent label were analyzed in all adult (n = 4) and juvenile (n = 3) birds. Images of Medial Striatum, which includes Area X, were taken with a 40x objective using ZEN imaging software on a confocal microscope (Zeiss LSM 780 inverted microscope or Zeiss 880 LSM inverted microscope) at the USC Translational Imaging Center. The excitation and emission wavelength settings used to image each marker are given in Table 1. The tilescan feature of ZEN was used to acquire multiple individual image fields ("tiles") at adjacent XY positions. To collect depth information, the ZEN Z-stack feature was used to image multiple Z

positions ("Z-steps") for each tile. Each tile within a tissue section included 8-12 Z-steps that were taken approximately 3-4 µm apart. All tiles from a single section were stitched together using the Imaris Stitcher application (Oxford Instruments); the Z-steps were reconstructed into a 3D projection of the section in Imaris image analysis software. 3D projections of individual or multiple fluorescent channels were used in all figures except Figure 7, which was taken at a single Z-step (see below, 2.4.4). All 3D projections were visualized using the maximum intensity projection display mode in Imaris, which presents the voxel with maximum intensity from all Z-steps in the viewing plane.

3D projections were analyzed for overlap between neurons retrogradely-labeled from DLM and neurons expressing FoxP2. Because qualitative examination in the four adult birds and one juvenile bird revealed that anti-FoxP2 16046 produced a higher incidence of labeled cells than anti-FoxP2 1307, we used ab16046 to stain the remaining two juvenile birds. Both antibodies were used in all four adults for overlap analysis, whereas for juvenile overlap analysis, only one bird was labeled for both antibodies and the remaining two out of three birds were labeled with the 16046 antibody only (Table 2). Retrogradely-labeled cells (adults: n = 469; juveniles: n = 87) were classified as being within either Area X-CORE (outlined by DAPI staining), Area X-SHELL (defined conservatively as 500 µm extending from the medial, ventromedial, and ventrolateral edges of Area X-CORE) (Iyengar et al., 1999), or "Border" if the boundary between Area X-CORE and SHELL was not easily identifiable by DAPI staining. Each retrogradely-labeled cell was visually inspected at every Z-step for signs of double-label with FoxP2.

2.4.2 Size Analysis of Retrogradely-Labeled Neurons in Area X

Size analysis of Area X neurons retrogradely-labeled from DLM was performed on all four adult birds (n = 100 neurons chosen randomly for analysis out of a total of 469 retrogradely-labeled neurons) and all three juvenile birds (n = 87 neurons). Using the Surfaces auto-detection feature in Imaris, 3D models of labeled cells were created based on the Z-intervals described above in 2.4.1 (3-4 µm). Cells were detected using automatic absolute intensity thresholding on the Alexa Fluor 488 channel and smoothed with a surface detail of 0.50 µm to ensure close fitting to the staining. Once the Surfaces feature created 3D models of labeled cells, we manually inspected each neuron and adjusted each surface to include only the cell body, excluding the primary dendrites.

We measured the diameter of these 3D models of retrogradely-labeled neurons by using the average of two measures of Imaris Surfaces: the longest diameter of each soma (referred to

as Object-Oriented Bounding Box Length C) and the perpendicular diameter (Object-Oriented Bounding Box Length B); we refer to this measure as the average diameter. This measure is consistent with previous studies in songbirds that reported the average of the major and minor axis of somata as the diameter (Luo and Perkel, 1999b; Reiner et al., 2004). The average diameter values of each Alexa 488-labeled surface were exported to Microsoft Excel for further analysis.

2.4.3 Size Analysis of FoxP2-Positive Nuclei in Area X

Size analysis of FoxP2+ nuclei in Area X-CORE and SHELL was performed on three adult and two juvenile birds. Only sections stained for anti-FoxP2 16046 were examined in the FoxP2+ size analysis. Because the full depth profile of smaller FoxP2+ neurons was not represented in the Z-stack intervals used in the overlap analysis of DIR pallidal cells described above, two to five tissue sections per bird were reimaged with smaller Z-stack intervals (0.46 - 0.47 µm apart). This interval ensured that smaller cell types, such as medium spiny neurons, were included in the analysis. The boundaries of Area X-CORE in each section were determined using DAPI staining and divided into 212.55 µm x 212.55 µm fields. Images of two randomly selected fields per section were taken at a magnification of 40x with the updated Z-stack intervals. Two 212.55 µm x 212.55 µm fields per section were also taken for Area X-SHELL: one image was taken ventromedial and the other ventrolateral to the boundaries of Area X-CORE.

We determined the size of FoxP2+ nuclei using the Surfaces auto-detection feature of Imaris on the Alexa Fluor 594 channel; surfaces were smoothed with a surface detail of 0.50 µm (Fig. 2). Adjoining surfaces were split based on morphology with a seed point diameter of 3.20 µm to account for the smallest possible striatal medium spiny neuron diameter (Carrillo and Doupe, 2004). Cells were then filtered based on area (>25 µm^2) and distance from the XYZ border (>0.5 µm) to remove cells that were only partially included in the section. Each 3D projection generated by the Surfaces feature was visually inspected to see if the 3D model matched the FoxP2 staining; if the model's outline was not accurate, the surface was deleted. Additionally, all neurons that exhibited a fusiform appearance were excluded from analysis; these cells are likely migratory young neurons (Alvarez-Buylla and Nottebohm, 1988). In adults, 3,953 and 1,535 FoxP2+ nuclei were obtained from Area X-CORE and Area X-SHELL, respectively. In Area X-CORE and Area X-SHELL of juveniles, 1,813 and 835 FoxP2+ nuclei were analyzed. Measurements of average diameter were calculated as explained in section 2.4.2 and exported to Microsoft Excel for further analysis.

2.4.4 Overlap Analysis of Large FoxP2+ Neurons with DARPP-32 in Area X

As detailed below in Results, we observed a population of large FoxP2+ neurons that were not retrogradely-labeled from DLM, representing possible IND pallidal neurons. However, because FoxP2 was expressed in nuclei of different sizes along a continuous distribution (see below, Fig. 5), it was not obvious how to segregate non-projecting FoxP2+ pallidal neurons from other FoxP2+ cells. We decided to use nuclear diameter as a means of segregating putative IND neurons given that their nuclear diameter should be similar to that of DIR DLM-projecting neurons. Based on this idea, we first measured the nuclear diameter of 50 FoxP2- retrogradely-labeled DIR neurons from one adult bird in which DAPI staining was particularly good. The full depth profile of each DIR neuron was inspected using the original Z-steps of 3-4 µm (see above, 2.4.1) to select the Z-step that included the largest visible nucleus. The Measurement Points feature of Imaris was then used to determine the longest diameter and the perpendicular diameter of the nucleus at the selected Z-step. The mean of these two values was calculated in Microsoft Excel and reported as the average nuclear diameter for DIR neurons. It should be noted that the average diameter values may represent an underestimate since there was no way to confirm that the cell was imaged at the Z-plane where the nucleus was the largest. The mean diameter of these DIR pallidal nuclei was 7.95 ± 1.01 µm and did not differ between neurons in Area X-CORE and SHELL (p = 0.2941; unpaired t-test). We therefore established an initial cutoff of 8 µm to separate putative IND pallidal neurons from smaller medium spiny neurons based on nuclear size. While this measurement was established from adult data, we applied the nuclear size cutoff to juveniles because the Alexa 488+ DIR somal size did not differ between adult and juvenile songbirds (see below) and DAPI staining was clearer in adults.

Large (≥8 um) FoxP2+ neurons in two juvenile birds with staining for the striatal marker DARPP-32 were analyzed for overlap to assess whether these cells were pallidal in nature. We examined all FoxP2+ neurons with an average diameter ≥8 µm (n = 96; putative IND pallidal neurons) across Area X-CORE and SHELL. Each neuron was visually inspected at every Z position (at a step size of 0.46-0.47 µm) for signs of FoxP2/DARPP-32 co-expression. For purposes of illustration, the voxel-based Colocalization feature in Imaris was used to display voxels that co-expressed both FoxP2 and DARPP-32 channels. The threshold for FoxP2 and DARPP-32 channel colocalization was set at the single Z-step where the FoxP2+ nucleus appeared the largest and was then applied to the entire Z-stack to visualize double label (see Fig. 6).

2.4.5 Statistical Tests

Statistical analyses were performed with GraphPad Prism software. All continuous data were tested for normality using D'Agostino-Pearson tests. Two-tailed unpaired t-tests were used to compare means for normally distributed data, including Area X-CORE versus SHELL comparisons within juvenile and adult birds. Unpaired t-tests were used due to the differing numbers of neurons between Area X-CORE and SHELL. Nonparametric Mann-Whitney tests were used to compare medians for non-normally distributed data for Area X-CORE versus SHELL regions within age groups. Differences in the proportions of neurons were assessed for statistical significance using the Chi-square test. When there were only two possible outcomes, a binomial test was used to analyze differences in the expected and observed proportions. The expected proportion of DIR neurons that co-expressed FoxP2 was set to 0.5 to represent random chance. Measures of central tendency for normally and non-normally distributed data are reported as mean ± standard deviation and median with first and third quartiles (Table 3), respectively.

3. RESULTS

3.1 Direct DLM-projecting neurons in Area X do not express FoxP2 in adult or juvenile songbirds

We injected dextran-Alexa 488 into DLM in order to retrogradely label thalamus-projecting (DIR) neurons within CORE and SHELL regions of Area X (see Methods; Fig. 3a, 4a). DAPI staining was used to identify the outline of Area X-CORE (Fig. 3b, 4b); Area X-SHELL was defined conservatively as extending 500 µm from the medial, ventromedial, and ventrolateral edges of Area X-CORE (Iyengar et al., 1999). Our injections revealed a topographically organized pattern of retrograde label in Area X-CORE and Area X-SHELL that was consistent with previous work (Johnson et al., 1995; Iyengar et al., 1999; Luo and Perkel, 1999b; Person et al., 2008) (data not shown). Some injections extended beyond the borders of DLM, labeling neurons outside of Area X-CORE and SHELL; we did not include these neurons in our analysis. The precise borders of Area X-CORE were poorly defined in some sections, and some retrogradely-labeled neurons were located very close to the CORE-SHELL border. In these cases, we assigned labeled neurons to the border region between Area X-CORE and Area X-SHELL.

Sections were stained in two series by different antibodies for FoxP2 (anti-FoxP2 16046 or anti-FoxP2 1307, see Methods; Fig. 3c). Table 2 shows the number of DIR DLM-projecting

neurons in Area X-CORE, Area X-SHELL, and the border between them in each bird, as well as which antibody was used to analyze overlap with FoxP2. We analyzed a total of 469 retrogradely-labeled neurons across four adult male birds (Fig. 3d). Retrogradely-labeled neurons across Area X-CORE and Area X-SHELL were inspected for overlap with FoxP2 (Fig. 3e, f; see Methods). Of these 469 DLM-projecting neurons, only two neurons in Area X-CORE showed possible co-expression of Alexa 488 (DLM-projecting) and Alexa 594 (FoxP2) labels ($p < 0.0001$, binomial test) (Fig. 3h). The somata and primary dendrites of Area X neurons that projected to DLM were robustly labeled (Fig. 3f). These primary dendrites were aspiny in nature, characteristic of pallidal neurons as described by Reiner et al. (2004). To determine whether the somal size of DIR neurons differed between Area X-CORE versus Area X-SHELL, 100 DIR neurons from the 469 retrogradely-labeled neurons across four adult birds were randomly selected for size measurements. Of these, 54 were in Area X-CORE, 35 were in Area X-SHELL, and 11 were in the border region; the border region neurons were excluded from size analysis because they could not be definitively categorized into either group. The DIR somata in Area X-CORE had a mean diameter of 16.29 ± 2.17 µm ($n = 54$). In comparison, DIR somata in Area X-SHELL had a mean diameter of 14.39 ± 1.59 µm ($n = 35$), indicating a significant difference between the two regions ($p < 0.0001$; unpaired t-test) (Fig. 3g). This difference shows that DIR neurons in Area X-CORE are larger than those in Area X-SHELL.

We next examined whether the absence of FoxP2 expression in adult DLM-projecting DIR neurons was also true in juvenile zebra finches ($n = 3$; Fig. 4c, d). Eighty-seven retrogradely-labeled cells were analyzed for FoxP2 co-expression (Fig. 4e, f), including 38 in Area X-CORE, 26 in Area X-SHELL, and 23 in the border region between CORE and SHELL (Table 2). None of these juvenile DIR neurons showed expression of FoxP2 (Fig. 4h). The size of DLM-projecting somata in juvenile Area X-CORE (17.05 ± 2.28 µm, $n = 38$) was larger than those in Area X-SHELL (15.13 ± 2.28 µm, $n = 26$) (Fig. 4g). As in adults, this size difference in retrogradely-labeled neurons was significant ($p = 0.0016$; unpaired t-test). The DIR somal sizes in juvenile Area X-CORE did not differ compared to adult Area X-CORE ($p = 0.1085$; unpaired t-test), nor did juvenile Area X-SHELL DIR somal size differ from that in adult Area X-SHELL ($p = 0.1403$; unpaired t-test). Together, these data show that DIR Area X→DLM projection neurons do not express FoxP2 in either adult or juvenile zebra finches. In addition, the somata of DLM-projecting neurons in both adults and juveniles are larger in Area X-CORE compared to X-SHELL.

3.2 FoxP2+ nuclei vary in size and intensity

FoxP2+ nuclei in Area X exhibited a range of sizes in both adult (n = 3) and juvenile birds (n = 2) (Fig. 5a, c). Frequency histograms of size across all FoxP2+ nuclei revealed a continuous, positively skewed distribution for adults and juveniles within both Area X-CORE and Area X-SHELL (Fig. 5b). In Area X-CORE, the average diameter of FoxP2-labeled nuclei ranged from 3.02 µm to 9.97 µm in adults and 3.21 to 10.14 µm in juveniles. In Area X-SHELL, the average diameter of labeled nuclei ranged from 3.21 µm to 9.61 µm in adults and 3.46 µm to 10.02 µm in juveniles. Thus, FoxP2 was expressed across a similar, continuous distribution of nuclear sizes in CORE and SHELL regions of both juveniles and adults.

FoxP2 is highly expressed in medium spiny neurons of Area X (Haesler et al., 2004). However, snRNA-seq analysis by Xiao et. al (2021) showed that FoxP2 is also expressed in other cell types of adult birds, including pallidal-like neurons. Pallidal cells in the Medial Striatum of zebra finches, including Area X, are larger than striatal medium spiny neurons (Luo and Perkel, 1999b; Reiner et al., 2004), as is true in mammals. Reiner et al. (2004) previously reported that DLM-projecting pallidal neurons in zebra finches had average somal diameters of 12.4 ± 1.8 µm. In contrast, they found that the somata of medium spiny neurons, marked by the striatal cytoplasmic marker DARPP-32, had a diameter of 7.1 ± 1.1 µm. Interestingly, we did not observe a bimodal distribution in size suggested by this difference in diameter between medium spiny neurons and pallidal neurons (Fig. 5b). This may be due to the fact that FoxP2 is also expressed in some types of interneurons as well as oligodendrocytes in Area X of zebra finches (Xiao et al., 2021). We hypothesized that most of the smaller FoxP2+ neurons corresponded to medium spiny neurons (Fig. 5a, c; white arrows), while the sparsely distributed larger FoxP2+ neurons corresponded to IND pallidal neurons (Fig. 5a, c; white arrowheads).

To gauge whether large FoxP2+ cells belonged to a distinct subpopulation, we first estimated the nuclear size of DLM-projecting pallidal neurons (see Methods). This analysis indicated that nuclei of DIR DLM-projecting neurons had a mean diameter of 7.95 ± 1.01 µm. We therefore established 8 µm as an initial cutoff for the average nuclear diameter of putative FoxP2+ indirect pallidal neurons. In Area X-CORE of adults, 97.42% of FoxP2+ nuclei had an average diameter <8 µm (referred to as "small FoxP2+ cells"); the median diameter of these nuclei was 5.31 µm (Table 3). In Area X-SHELL, small FoxP2+ nuclei accounted for 97.39% of the total cell count in adults, with a median diameter of 5.41 µm. In juveniles, small FoxP2+ nuclei comprised 97.35% of Area X-CORE, with a median diameter of 5.85 µm (Table 3). In juvenile Area X-SHELL, 94.25% of the total cell count were small FoxP2+ cells, and the median

diameter was 5.70 µm. Although modest in terms of absolute difference, all comparisons of the size of small FoxP2+ nuclei between Area X-CORE and SHELL and between adults and juveniles were statistically significant. In adults, small FoxP2+ neurons in Area X-CORE had a smaller nuclear size than those in Area X-SHELL (p = 0.0018; Mann-Whitney test. In contrast, small FoxP2+ neurons in Area X-CORE of juveniles were larger than those in Area X-SHELL (p = 0.0017; Mann-Whitney test). Comparisons across age groups showed that small FoxP2+ nuclei in juveniles were larger than the small FoxP2+ nuclei in adults across both Area X-CORE (p < 0.0001; Mann-Whitney test) and Area X-SHELL (p < 0.0001; Mann-Whitney test).

We next analyzed the nuclear size of large FoxP2+ cells – a putative population of IND pallidal neurons. In adult zebra finches, 2.58% of FoxP2+ nuclei in Area X-CORE fell above the cutoff of ≥8 µm (referred to as "large FoxP2+ cells"), with a median diameter of 8.35 µm (Table 3). In adult Area X-SHELL, 2.61% of FoxP2+ nuclei were large and had a median diameter of 8.17 µm. This difference in size between large FoxP2+ nuclei of adult Area X-CORE and Area X-SHELL was significant (p = 0.0297; Mann-Whitney test). The proportion of large FoxP2+ nuclei did not differ between the CORE and SHELL subregions (p = 0.9573; Chi-square test). In juvenile songbirds, 2.65% and 5.75% of FoxP2+ nuclei were categorized as large (≥8 µm) in Area X-CORE and Area X-SHELL, respectively. This higher proportion of large FoxP2+ nuclei in juvenile Area X-SHELL compared to juvenile Area X-CORE was statistically significant (p < 0.0001; Chi-square test). Despite the proportional difference, the size of large FoxP2+ nuclei diameters in juveniles did not differ between Area X-CORE (median: 8.43 µm) and Area X-SHELL (median: 8.37 µm) (p = 0.6549; Mann-Whitney test). The proportion of large FoxP2+ nuclei did not differ across adults and juveniles in Area X-CORE (p = 0.8816; Chi-square test); however, Area X-SHELL of juveniles contained a greater proportion of large FoxP2+ nuclei compared to adult Area X-SHELL (p = 0.0001; Chi-square test), suggesting a developmental difference in the cellular makeup of this region. Nuclear size did not differ between large FoxP2+ nuclei in adult versus juvenile Area X-CORE (p = 0.4050; Mann-Whitney test), but was larger in Area X-SHELL of juveniles compared to adults (p = 0.0251; Mann-Whitney test), providing additional evidence of a region-specific developmental difference.

Interestingly, our qualitative observations revealed that these large nuclei were generally more intensely stained for FoxP2 than their smaller counterparts in both adults and juveniles (Fig. 5a, c). This observation aligns with data from Thompson et al. (2013) and Kosubek-Langer and Scharff (2020), which showed a bimodal distribution of relatively larger intensely-stained FoxP2+ neurons and relatively smaller weakly-stained FoxP2+ neurons in Area X.

3.3 Most large FoxP2+ neurons do not express DARPP-32

In order to investigate whether the larger FoxP2+ neurons were pallidal in nature, we examined the expression of DARPP-32, which marks ~50% of striatal spiny neurons in avian brain (Reiner et al., 1998; Reiner et al., 2004; Singh and Iyengar, 2019). DLM-projecting cells were never co-labeled with DARPP-32 in Area X-CORE or Area X-SHELL (Fig. 6a), consistent with a pallidal identity of DIR neurons. Therefore, if larger FoxP2+ neurons belong to an indirect pallidal class, we expected to observe a similar lack of overlap with DARPP-32 expression.

We analyzed 96 large FoxP2+ neurons with an average nuclear diameter ≥8 μm across both Area X-CORE and Area X-SHELL in two juvenile birds (Table 3; see Methods). FoxP2 staining (red) was inspected for co-expression with DARPP-32 staining (magenta); white voxels depict colocalization of the two channels (Fig. 6b, c). The majority of large FoxP2+ neurons did not express DARPP-32 in either Area-X CORE (65%, 31/48) or Area X-SHELL (69%, 33/48); two examples from Area X-CORE are marked by yellow arrowheads in Figure 6b. Surprisingly, a substantial number of larger neurons did co-express FoxP2 and DARPP-32 in both CORE and SHELL regions of Area X. That is, 35% (17/48) of large FoxP2+ neurons were double-labeled in Area X-CORE and 31% (15/48) were double-labeled in Area X-SHELL in these two birds (Fig. 6d). Figure 6c shows a large neuron in Area X-SHELL that expressed both FoxP2 and DARPP-32 (yellow arrowhead). This surprising result is consistent with the snRNA-seq findings of Xiao et al. (2021) in Area X that DARPP-32 is expressed in multiple cell types, including pallidal neurons.

However, our cutoff for large neurons represented a fairly rudimentary criterion, so we decided to assess the incidence of FoxP2 and DARPP-32 co-expression when the cutoff value of 8.0 μm was increased by one or two standard deviations. Interestingly, when the cutoff value was increased by one standard deviation to 9.01 (SD = 1.01 μm), only 2% (1/48) of large FoxP2+ nuclei in Area X-CORE expressed DARPP-32. When the cutoff was raised by two standard deviations to 10.02, 0% (0/48) of large FoxP2+ nuclei in Area X-CORE were also DARPP-32+. Similarly, when the cutoff value was raised by one or two standard deviations only 2% (1/48) and 0% (0/48) of large FoxP2+ nuclei in Area X-SHELL expressed DARPP-32, respectively.

We examined smaller neurons qualitatively, which revealed that a large majority expressed both FoxP2 and DARPP-32 as seen by the white voxels depicting colocalization (Fig. 6b, c; yellow arrows). Most of these likely represent medium spiny neurons, although it is possible that other small cell types such as some interneurons and glial cells also express

FoxP2 in Area X (Xiao et al., 2021). These results indicate that DARPP-32 is not a selective marker for medium spiny neurons in zebra finch brain, a finding that is similar to results in pigeons (Reiner et al., 1998; Reiner et al., 2004; Singh and Iyengar, 2019).

Our qualitative observations of smaller neurons in Area X-CORE and SHELL also revealed that FoxP2+ nuclei of double-labeled neurons took up a large percentage of the cytoplasm labeled by DARPP-32, examples of which can be seen in Figures 6b and 6c (yellow arrows). These observations conform to the report of Ouimet et al. (1998) that DARPP-32+ medium spiny neurons of rat striatum have scant cytoplasm. Qualitative inspection also suggested that the nuclei of large FoxP2+/DARPP-32+ cells (possible IND pallidal neurons) in Area X-CORE and SHELL follow a similar pattern by taking up a large percentage of the cytoplasm. In Figure 6c, the nuclear FoxP2 staining in a large FoxP2+ neuron (yellow arrowhead) takes up a large percentage of the cytoplasmic DARPP-32 staining, as seen by the white pixels indicating colocalization. Conversely, when we examined DIR retrogradely-labeled neurons, we observed that the DAPI+ nuclei encompassed a considerably smaller percentage of the soma. This observation was consistent across both Area X-CORE and SHELL regions. Figure 7 shows a DIR neuron in Area X-CORE in which the DAPI+ nucleus (blue) is surrounded by relatively more cytoplasm (green) compared to the larger neuron in Figure 6c (yellow arrowhead). An adjacent FoxP2+ nucleus may represent an interneuron or a medium spiny neuron. These observations suggest that DIR and IND pallidal cells have a similar nuclear size but vary in the percentage of the soma that their nuclei encompass. DIR neurons may have a larger amount of cytoplasm in order to support a longer axon and/or other features.

4. DISCUSSION

Vocal learning is mediated in part by two parallel topographically organized cortico-basal ganglia circuits, CORE and SHELL, that form reentrant loops. Area X sends inhibitory projections to the thalamic nucleus DLM: Area X-CORE projects to the dorsolateral subregion of DLM (DLMdl) and Area X-SHELL projects to the ventromedial subregion of DLM (DLMvm) (Bottjer et al., 1989; Johnson et al., 1995; Iyengar et al., 1999; Carrillo and Doupe, 2004; Person et al., 2008). DLMdl and DLMvm project to LMAN-CORE and LMAN-SHELL respectively, forming recurrent pathways (Johnson et al., 1995; Iyengar et al., 1999; Luo et al., 2001; Pinaud et al., 2007). Thus, DLM receives information from independent Area X-CORE and SHELL pathways as well as convergent input from AId (Foster et al., 1997; Bottjer et al., 2000). Given the complexity of these connections, it remains unclear how motor production and vocal evaluation information converge (Johnson and Bottjer, 1992; Johnson et al., 1995; Iyengar et

al., 1999; Paterson and Bottjer, 2017). Our goal in this study was to elucidate specific patterns of connectivity underlying how vocal information is conveyed from Area X to DLM. A prior study using snRNA-seq showed that pallidal neurons in Area X express either FoxP2 or PENK (the precursor to enkephalin), but not both (Xiao et al., 2021). Given that enkephalin is the neurotransmitter of DIR DLM-projecting neurons (Carrillo and Doupe, 2004), this distinction suggested the hypothesis that DLM-projecting pallidal cells are FoxP2-/PENK+, while non-projecting pallidal cells are FoxP2+/PENK-. In this study, we tested FoxP2 expression as a possible marker for direct versus indirect pallidal neurons in Area X as a first step to distinguish how information may be differently conveyed via these two pathways from Area X→DLM.

Consistent with this hypothesis, we found that DIR DLM-projecting neurons in CORE and SHELL regions of Area X of both adult and juvenile songbirds lacked expression of FoxP2. Neurons with nuclei that did express FoxP2 exhibited various sizes and intensities in both CORE and SHELL regions of Area X. Qualitative inspection revealed a subpopulation of FoxP2+ neurons that were larger and more intensely stained, consistent with findings from Thompson et al. (2013) and Kosubek-Langer and Scharff (2020). Coupled with the results from Xiao et al. (2021) regarding a lack of colocalization between FoxP2 and PENK in Area X, this pattern of results suggests that the larger, more intense FoxP2+ neurons represent a class of indirect pallidal neurons (Farries et al., 2005), while the smaller neurons represent spiny neurons, and other cell types.

As a preliminary test of whether larger FoxP2+ (putative IND) neurons had a pallidal identity, we examined expression of the striatal spiny neuron marker DARPP-32. Surprisingly, we discovered that 32/96 (33%) large (≥8 μm diameter) FoxP2+ neurons expressed DARPP-32 in juvenile birds. We found that increasing our average size cutoff value by one or two standard deviations greatly reduced or eliminated the incidence of cells that co-expressed FoxP2 and DARPP-32, suggesting perhaps that most IND pallidal neurons do not express DARPP-32. An alternative interpretation is that Area X contains diverse classes of FoxP2+ pallidal cells which have a larger range of nuclear sizes compared to DIR pallidal cells, including some that may be homologous to diverse classes of GPe pallidal neurons (Hegeman et al., 2016; Mastro et al., 2017). DARPP-32 is not expressed in pallidal neurons of rodents or pigeons (Anderson and Reiner, 1991; Reiner et al., 1998). FoxP2 does not colocalize with antibody markers for choline acetyltransferase (ChAT), nitric oxide (nNOS/SS/NPY), or parvalbumin (PV) interneurons in Area X (Haesler et al., 2004). DARPP-32 is also not detectable in cholinergic, PV+, or NPY+ striatal interneurons of rat or pigeon (Anderson and Reiner, 1991; Reiner et al., 1998).

However, recent studies of the molecular profile of songbird and chicken brains have shown that a distinct class of GABAergic neurons derived from the LGE (lateral ganglionic eminence) expresses FoxP2 and occurs throughout the telencephalon (Colquitt et al., 2021; Zaremba et al., 2025). In mammals, this population is expressed primarily in striatum (spiny neuron) and ventral pallium (olfactory bulb and pallial amygdala), but in songbird Area X this class includes both spiny and pallidal neurons. In addition, the snRNA-seq analysis of Xiao et al. (2021) in adult Area X indicated that FoxP2 is expressed in interneurons (type unknown), oligodendrocytes, and endothelial cells, and that DARPP-32 (PPP1R1B) is expressed at a low level in some pallidal neurons. This latter finding is consistent with our observation that DARPP-32 and FoxP2 co-localized to a subset of pallidal neurons in juvenile birds.

Indeed, the growing complexity of cell types and connections in basal ganglia generally (Hegeman et al., 2016; Oorschot, 2016; Soares-Cunha et al., 2016; Bostan et al., 2018; Spix et al., 2021; Arber and Costa, 2022), as well as the combined presence of spiny neurons, pallidal cells, interneurons, and possibly other neuronal cell types within Medial Striatum of avian brain, offer multiple possibilities for the identity of large FoxP2+/DARPP-32+ neurons in Area X. Mammalian GPe contains a heterogeneous assortment of neurons with diverse connections (Hegeman et al., 2016; Spix et al., 2021), suggesting that Area X may also contain multiple subclasses of GPe-like pallidal neurons. Xiao et al. (2021) reported that a cluster identified as pallidal neurons by snRNA-seq expressed high levels of markers for a type of GPe neuron known as arkypallidal (Hegeman et al., 2016), which projects back to striatum in mammals. In accordance with this idea, there are no pallidal projections from Area X to the subthalamic nucleus (STN), and pallidal neurons within Area X seem to make collateral projections onto spiny neurons (Farries et al., 2005; Gale and Perkel, 2010). Further characterization based on additional molecular markers combined with patterns of connectivity in both juvenile and adult zebra finches is necessary to distinguish whether the non-projecting pallidal cells of CORE and SHELL regions of Area X contain distinct subclasses.

FoxP2 and DARPP-32 may colocalize in a specific subclass of pallidal neurons only in juvenile birds and could represent a transient feature of vocal learning. Haesler et al. (2004) reported developmental increases in the expression of FoxP2 within Area X that were highest in juvenile zebra finches between 35 and 50 dph, which corresponds to the age range of our juvenile birds (~45 dph). This age range includes early stages of sensorimotor integration when birds are learning to imitate a memorized tutor song. Increased co-expression of FoxP2 and DARPP-32 may occur within IND pallidal neurons only during this period of vocal learning.

Because we did not examine the expression of DARPP-32 in adult birds, this remains an interesting future question. DARPP-32 is a phosphoprotein that is coupled to multiple receptors, including D1/D5 receptors, D2 and adenosine A2a receptors, and NMDA receptors (Blank, 1997; Langley et al., 1997; Yger and Girault, 2011). In striatal tissue, DARPP-32 is necessary for the cAMP-dependent enhancement of NMDA responses (Blank, 1997). Pallidal neurons in Area X are known to receive direct glutamatergic projections from both HVC and LMAN (Farries et al., 2005). Although it is not known whether these cortico-pallidal projections include an NMDAR-mediated component, NMDA responses are ubiquitous in projections from DLM to LMAN and from HVC and LMAN to RA (Mooney and Konishi, 1991; Bottjer et al., 1998; Stark and Perkel, 1999; Bottjer, 2005). It is also not known whether pallidal neurons in Area X receive direct dopaminergic inputs, although DA projections to pallidal cells occur in primates and rodents (Lindvall and Bjorklund, 1979; Gauthier et al., 1999; Eid and Parent, 2016). One intriguing possibility is that DARPP-32 may help to regulate interactions between glutamatergic and dopaminergic afferents by regulating the enhancement of NMDA responses (Blank, 1997).

Overall, our findings indicate that FoxP2 staining, along with size measurements of nuclei and somata, may be useful as a marker to distinguish the indirect pallidal neuron population in Area X that forms close contacts with direct DLM-projecting pallidal neurons (Farries et al., 2005). In terms of functional roles, the direct pathway of the basal ganglia increases thalamic activity, whereas the indirect pathway is thought to oppose the DIR pathway to decrease thalamic activity (Alexander and Crutcher, 1990). When self-generated vocal motor output matches the memorized tutor song, the direct pathway may enhance firing in DLM to generate a positive reinforcement signal for "correct" vocal behavior. In contrast, pallidal neurons that receive inputs from spiny neurons in a "traditional" indirect pathway would suppress firing in DLM. In that case, the indirect pathway may be involved in generating a negative reinforcement signal for "incorrect" behavior when vocalizations differ from memorized tutor sounds. However, because some pallidal neurons in Area X receive direct cortical inputs from HVC and LMAN, those excitatory inputs might increase firing rates in a subset of IND neurons, which would decrease firing rates in DIR neurons and thus increase thalamic activity (Farries et al., 2005). Understanding how these pathways connect and process motor skill information within the LMAN→Area X→DLM→LMAN feedback loop will provide insight into what roles the basal ganglia pathways play in vocal learning in songbirds as well as motor learning in humans.

5. REFERENCES

Achiro JM, Bottjer SW (2013) Neural representation of a target auditory memory in a cortico-basal ganglia pathway. J Neurosci 33:14475-14488.

Achiro JM, Shen J, Bottjer SW (2017) Neural activity in cortico-basal ganglia circuits of juvenile songbirds encodes performance during goal-directed learning. Elife 6.

Alexander GE, Crutcher MD (1990) Functional architecture of basal ganglia circuits: neural substrates of parallel processing. Trends Neurosci 13:266-271.

Alvarez-Buylla A, Nottebohm F (1988) Migration of young neurons in adult avian brain. Nature 335:353-354.

Anderson KD, Reiner A (1991) Immunohistochemical localization of DARPP-32 in striatal projection neurons and striatal interneurons: implications for the localization of D1-like dopamine receptors on different types of striatal neurons. Brain Res 568:235-243.

Arber S, Costa RM (2022) Networking brainstem and basal ganglia circuits for movement. Nat Rev Neurosci 23:342-360.

Aronov D, Andalman AS, Fee MS (2008) A specialized forebrain circuit for vocal babbling in the juvenile songbird. Science 320:630-634.

Blank RH (1997) Assisted reproduction and reproductive rights: the case of in vitro fertilization. Politics Life Sci 16:279-288.

Böhner J (1983) Song Learning in the Zebra Finch (Taeniopygia-Guttata) - Selectivity in the Choice of a Tutor and Accuracy of Song Copies. Anim Behav 31:231-237.

Böhner J (1990) Early acquisition of song in the zebra finch, *Taeniopygia guttata*. Anim Behav 39:369-374.

Bostan AC, Dum RP, Strick PL (2018) Functional Anatomy of Basal Ganglia Circuits with the Cerebral Cortex and the Cerebellum. Prog Neurol Surg 33:50-61.

Bottjer SW (2005) Silent synapses in a thalamo-cortical circuit necessary for song learning in zebra finches. J Neurophysiol 94:3698-3707.

Bottjer SW, Altenau B (2010) Parallel pathways for vocal learning in basal ganglia of songbirds. Nat Neurosci 13:153-155.

Bottjer SW, Miesner EA, Arnold AP (1984) Forebrain lesions disrupt development but not maintenance of song in passerine birds. Science 224:901-903.

Bottjer SW, Brady JD, Walsh JP (1998) Intrinsic and synaptic properties of neurons in the vocal-control nucleus IMAN from in vitro slice preparations of juvenile and adult zebra finches. J Neurobiol 37:642-658.

Bottjer SW, Brady JD, Cribbs B (2000) Connections of a motor cortical region in zebra finches: relation to pathways for vocal learning. J Comp Neurol 420:244-260.

Bottjer SW, Halsema KA, Brown SA, Miesner EA (1989) Axonal connections of a forebrain nucleus involved with vocal learning in zebra finches. J Comp Neurol 279:312-326.

Campbell P, Reep RL, Stoll ML, Ophir AG, Phelps SM (2009) Conservation and diversity of Foxp2 expression in muroid rodents: functional implications. J Comp Neurol 512:84-100.

Carrillo GD, Doupe AJ (2004) Is the songbird Area X striatal, pallidal, or both? An anatomical study. J Comp Neurol 473:415-437.

Catchpole C, Slater PJB (1995) Bird song : biological themes and variations. Cambridge England ;: New York, NY Cambridge University Press.

Clayton NS (1987) Song Learning in Cross-Fostered Zebra Finches - a Reexamination of the Sensitive Phase. Behaviour 102:67-81.

Colquitt BM, Merullo DP, Konopka G, Roberts TF, Brainard MS (2021) Cellular transcriptomics reveals evolutionary identities of songbird vocal circuits. Science 371.

den Hoed J, Fisher SE (2020) Genetic pathways involved in human speech disorders. Curr Opin Genet Dev 65:103-111.

Durstewitz D, Kroner S, Hemmings HC, Jr., Gunturkun O (1998) The dopaminergic innervation of the pigeon telencephalon: distribution of DARPP-32 and co-occurrence with glutamate decarboxylase and tyrosine hydroxylase. Neuroscience 83:763-779.

Eales LA (1985) Song learning in zebra finches - some effects of song model availability on what is learnt and when Anim Behav 33:1293-1300.

Eid L, Parent M (2016) Chemical anatomy of pallidal afferents in primates. Brain structure & function 221:4291-4317.

Farries MA, Ding L, Perkel DJ (2005) Evidence for "direct" and "indirect" pathways through the song system basal ganglia. J Comp Neurol 484:93-104.

Foster EF, Mehta RP, Bottjer SW (1997) Axonal connections of the medial magnocellular nucleus of the anterior neostriatum in zebra finches. J Comp Neurol 382:364-381.

Gale SD, Perkel DJ (2010) Anatomy of a songbird basal ganglia circuit essential for vocal learning and plasticity. J Chem Neuroanat 39:124-131.

Gale SD, Person AL, Perkel DJ (2008) A novel basal ganglia pathway forms a loop linking a vocal learning circuit with its dopaminergic input. J Comp Neurol 508:824-839.

Gauthier J, Parent M, Levesque M, Parent A (1999) The axonal arborization of single nigrostriatal neurons in rats. Brain Res 834:228-232.

Goldberg JH, Adler A, Bergman H, Fee MS (2010) Singing-related neural activity distinguishes two putative pallidal cell types in the songbird basal ganglia: comparison to the primate internal and external pallidal segments. J Neurosci 30:7088-7098.

Graybiel AM (2008) Habits, rituals, and the evaluative brain. Annu Rev Neurosci 31:359-387.

Haesler S, Rochefort C, Georgi B, Licznerski P, Osten P, Scharff C (2007) Incomplete and inaccurate vocal imitation after knockdown of FoxP2 in songbird basal ganglia nucleus Area X. PLoS Biol 5:e321.

Haesler S, Wada K, Nshdejan A, Morrisey EE, Lints T, Jarvis ED, Scharff C (2004) FoxP2 expression in avian vocal learners and non-learners. J Neurosci 24:3164-3175.

Hegeman DJ, Hong ES, Hernandez VM, Chan CS (2016) The external globus pallidus: progress and perspectives. Eur J Neurosci 43:1239-1265.

Hemmings HC, Jr., Greengard P (1986) DARPP-32, a dopamine- and adenosine 3':5'-monophosphate-regulated phosphoprotein: regional, tissue, and phylogenetic distribution. J Neurosci 6:1469-1481.

Immelmann K, ed (1969) Song development in the zebra finch and other estrildid finches.

Iyengar S, Viswanathan SS, Bottjer SW (1999) Development of topography within song control circuitry of zebra finches during the sensitive period for song learning. J Neurosci 19:6037-6057.

Johnson F, Bottjer SW (1992) Growth and regression of thalamic efferents in the song-control system of male zebra finches. J Comp Neurol 326:442-450.

Johnson F, Sablan MM, Bottjer SW (1995) Topographic organization of a forebrain pathway involved with vocal learning in zebra finches. J Comp Neurol 358:260-278.

Kosubek-Langer J, Scharff C (2020) Dynamic FoxP2 levels in male zebra finches are linked to morphology of adult-born Area X medium spiny neurons. Scientific reports 10:4787.

Kubikova L, Wada K, Jarvis ED (2010) Dopamine receptors in a songbird brain. J Comp Neurol 518:741-769.

Langley KC, Bergson C, Greengard P, Ouimet CC (1997) Co-localization of the D1 dopamine receptor in a subset of DARPP-32-containing neurons in rat caudate-putamen. Neuroscience 78:977-983.

Lindvall O, Bjorklund A (1979) Dopaminergic innervation of the globus pallidus by collaterals from the nigrostriatal pathway. Brain Res 172:169-173.

Luo M, Perkel DJ (1999a) A GABAergic, strongly inhibitory projection to a thalamic nucleus in the zebra finch song system. J Neurosci 19:6700-6711.

Luo M, Perkel DJ (1999b) Long-range GABAergic projection in a circuit essential for vocal learning. J Comp Neurol 403:68-84.

Luo M, Ding L, Perkel DJ (2001) An avian basal ganglia pathway essential for vocal learning forms a closed topographic loop. J Neurosci 21:6836-6845.

Mann NI, Slater PJB (1995) Song Tutor Choice by Zebra Finches in Aviaries. Anim Behav 49:811-820.

Mastro KJ, Zitelli KT, Willard AM, Leblanc KH, Kravitz AV, Gittis AH (2017) Cell-specific pallidal intervention induces long-lasting motor recovery in dopamine-depleted mice. Nat Neurosci 20:815-823.

Mooney R, Konishi M (1991) Two distinct inputs to an avian song nucleu activate different glutamate receptor subtypes on individual neurons Proc Natl Acad Sci U S A 88:4075-4079.

Nicholson DA, Roberts TF, Sober SJ (2018) Thalamostriatal and cerebellothalamic pathways in a songbird, the Bengalese finch. J Comp Neurol 526:1550-1570.

Nixdorf-Bergweiler BE, Lips MB, Heinemann U (1995) Electrophysiological and morphological evidence for a new projection of LMAN-neurones towards area X. Neuroreport 6:1729-1732.

Nottebohm F, Stokes TM, Leonard CM (1976) Central control of song in the canary, Serinus canarius. J Comp Neurol 165:457-486.

Oorschot DE (2016) Cell types in the different nuclei of the basal ganglia In: Handbook of Behavioral Neuroscience. , pp 99-117.: Elsevier.

Ouimet CC, Langley-Gullion KC, Greengard P (1998) Quantitative immunocytochemistry of DARPP-32-expressing neurons in the rat caudatoputamen. Brain Res 808:8-12.

Paterson AK, Bottjer SW (2017) Cortical inter-hemispheric circuits for multimodal vocal learning in songbirds. J Comp Neurol 525:3312-3340.

Person AL, Gale SD, Farries MA, Perkel DJ (2008) Organization of the songbird basal ganglia, including area X. J Comp Neurol 508:840-866.

Pidoux L, Le Blanc P, Levenes C, Leblois A (2018) A subcortical circuit linking the cerebellum to the basal ganglia engaged in vocal learning. Elife 7.

Pinaud R, Saldanha CJ, Wynne RD, Lovell PV, Mello CV (2007) The excitatory thalamo-"cortical" projection within the song control system of zebra finches is formed by calbindin-expressing neurons. J Comp Neurol 504:601-618.

Redgrave P, Rodriguez M, Smith Y, Rodriguez-Oroz MC, Lehericy S, Bergman H, Agid Y, DeLong MR, Obeso JA (2010) Goal-directed and habitual control in the basal ganglia: implications for Parkinson's disease. Nat Rev Neurosci 11:760-772.

Reiner A, Perera M, Paullus R, Medina L (1998) Immunohistochemical localization of DARPP32 in striatal projection neurons and striatal interneurons in pigeons. J Chem Neuroanat 16:17-33.

Reiner A, Laverghetta AV, Meade CA, Cuthbertson SL, Bottjer SW (2004) An immunohistochemical and pathway tracing study of the striatopallidal organization of area X in the male zebra finch. J Comp Neurol 469:239-261.

Roper A, Zann R (2006) The onset of song learning and song tutor selection in fledgling zebra finches. Ethology 112:458-470.

Sanchez-Valpuesta M, Suzuki Y, Shibata Y, Toji N, Ji Y, Afrin N, Asogwa CN, Kojima I, Mizuguchi D, Kojima S, Okanoya K, Okado H, Kobayashi K, Wada K (2019) Corticobasal ganglia projecting neurons are required for juvenile vocal learning but not for adult vocal plasticity in songbirds. Proc Natl Acad Sci U S A 116:22833-22843.

Scharff C, Nottebohm F (1991) A comparative study of the behavioral deficits following lesions of various parts of the zebra finch song system: implications for vocal learning. J Neurosci 11:2896-2913.

Shi Z, Piccus Z, Zhang X, Yang H, Jarrell H, Ding Y, Teng Z, Tchernichovski O, Li X (2018) miR-9 regulates basal ganglia-dependent developmental vocal learning and adult vocal performance in songbirds. Elife 7.

Singh UA, Iyengar S (2019) The expression of DARPP-32 in adult male zebra finches (Taenopygia guttata). Brain structure & function 224:2939-2972.

Soares-Cunha C, Coimbra B, Sousa N, Rodrigues AJ (2016) Reappraising striatal D1- and D2-neurons in reward and aversion. Neurosci Biobehav Rev 68:370-386.

Sohrabji F, Nordeen EJ, Nordeen KW (1990) Selective impairment of song learning following lesions of a forebrain nucleus in the juvenile zebra finch. Behav Neural Biol 53:51-63.

Spix TA, Nanivadekar S, Toong N, Kaplow IM, Isett BR, Goksen Y, Pfenning AR, Gittis AH (2021) Population-specific neuromodulation prolongs therapeutic benefits of deep brain stimulation. Science 374:201-206.

Stark LL, Perkel DJ (1999) Two-stage, input-specific synaptic maturation in a nucleus essential for vocal production in the zebra finch. J Neurosci 19:9107-9116.

Thompson CK, Schwabe F, Schoof A, Mendoza E, Gampe J, Rochefort C, Scharff C (2013) Young and intense: FoxP2 immunoreactivity in Area X varies with age, song stereotypy, and singing in male zebra finches. Frontiers in neural circuits 7:24.

Vates GE, Nottebohm F (1995) Feedback circuitry within a song-learning pathway. Proc Natl Acad Sci U S A 92:5139-5143.

Xiao L, Merullo DP, Koch TMI, Cao M, Co M, Kulkarni A, Konopka G, Roberts TF (2021) Expression of FoxP2 in the basal ganglia regulates vocal motor sequences in the adult songbird. Nature communications 12:2617.

Yger M, Girault JA (2011) DARPP-32, Jack of All Trades... Master of Which? Front Behav Neurosci 5:56.

Yin HH, Knowlton BJ (2006) The role of the basal ganglia in habit formation. Nat Rev Neurosci 7:464-476.

Zaremba B, Fallahshahroudi A, Schneider C, Schmidt J, Sarropoulos I, Leushkin E, Berki B, Van Poucke E, Jensen P, Senovilla-Ganzo R, Hervas-Sotomayor F, Trost N, Lamanna F, Sepp M, Garcia-Moreno F, Kaessmann H (2025) Developmental origins and evolution of pallial cell types and structures in birds. Science 387:eadp5182.

Label	Type of stain	Excitation/emission (nm)	Color in figures
DAPI	Nuclear	410/497	Blue
DLM-projecting Alexa Fluor 488	Cytoplasmic	500/580	Green
FoxP2 Alexa Fluor 594	Nuclear	589/651	Red
DARPP-32 Alexa Fluor 647	Cytoplasmic	653/755	Magenta

Table 1. Fluorescent label, type of cellular stain, confocal microscopy excitation/emission wavelengths, and color used in figures for each label in the experiment.

Bird ID	Age (days post-hatch)	# of DIR neurons in Area X-CORE	# of DIR neurons in Area X-SHELL	# of DIR neurons on border of Area X-CORE and Area X-SHELL	# neurons stained with ab16046/ab1307
Db51	167	98	30	10	67/71
R27	182	58	55	20	60/73
Db69	104	39	31	26	43/53
R54	91	47	52	3	46/56
Adult total = 469		**242**	**168**	**59**	**216/253**
Gy124	46	27	15	18	60/na
Gy48	44	9	8	4	21/na
O63	45	2	3	1	2/4
Juvenile total = 87		**38**	**26**	**23**	**83/4**

Table 2. Age and number of DLM-projecting DIR neurons in Area X-CORE and Area X-SHELL. The right-most column indicates which antibody against FoxP2 was used for overlap analysis. If DAPI staining was poor on a given tissue section and could not be used to define the precise borders of Area X-CORE, the DLM-projecting direct neurons were placed in the "Border" column. See text for further details.

	Large FoxP2+		Small FoxP2+	
	Area X-CORE	Area X-SHELL	Area X-CORE	Area X-SHELL
Adult	8.35 (8.15, 8.77) n = 102 2.58%	8.17 (8.09, 8.47) n = 40 2.61%	5.31 (4.66, 6.02) n = 3851 97.42%	5.41 (4.74, 6.15) n = 1495 97.39%
Juvenile	8.43 (8.23, 8.70) n = 48 2.65%	8.37 (8.15, 8.87) n = 48 5.75%	5.85 (5.15, 6.61) n = 1765 97.35%	5.70 (4.82, 6.60) n = 787 94.25%

Table 3. Median diameter, Q1, and Q3 values reported in μm, cell count, and relative percentage for large (≥8 μm) and small (<8 μm) FoxP2+ nuclei in Area X-CORE and Area X-SHELL of adults and juveniles.

Figure 1. Schematics of cortico-basal ganglia circuits in zebra finches. (a) Simplified schematic of cortico-basal ganglia circuits that mediate vocal learning in songbirds. The recurrent pathway from LMAN→Area X→DLM comprises CORE and SHELL subregions that form independent parallel loops. (b) Schematic of putative direct (DIR) and indirect (IND) pathways from Area X to the thalamus (DLM). Abbreviations: RA, robust nucleus of the arcopallium; AId, dorsal intermediate arcopallium; LMAN, lateral magnocellular nucleus of the anterior nidopallium; DLM, medial portion of the dorsolateral anterior thalamic nucleus, HVC, high vocal center; MSN, medium spiny neuron.

Figure 2. Surfaces auto-detection feature was used to measure size of FoxP2+ nuclei.
The size of FoxP2+ nuclei was measured using the Surfaces auto-detection feature of Imaris;
confocal image shows white outlines around FoxP2+ nuclei. See text for further details.

Figure 3. DIR neurons in Area X of adult zebra finches do not express FoxP2.

(a) Example of fluorescent tracer injection into DLM (outlined by dashed line). Injections produced retrograde label in Area X-CORE and Area X-SHELL of an adult bird (d-f). Injection site extended beyond the ventral border of DLM, but we did not analyze labeled neurons outside of CORE and SHELL regions of Area X (see Methods). (b) DAPI staining outlines Area X-CORE (inner dashed circle) within Medial Striatum of basal ganglia (gray line marks the axon tract PSL, pallial-subpallial lamina, which separates striatum from cortex). Area X-SHELL (outer

dashed circle) was conservatively defined as extending 500 µm from the medial, ventromedial, and ventrolateral edges of Area X-CORE. (c) Red FoxP2 staining (Alexa 594) using 16046 antibody and (d) green retrogradely-labeled cells from DLM (Alexa 488) in the same section as (b). (e) Overlay of DAPI, FoxP2, and DLM-projecting dyes. (f) Enlarged image of boxed region in (e) demonstrating that red FoxP2 label and green DLM-projecting retrograde label did not colocalize in a DIR neuron in Area X-CORE. (g) Box plots showing that the mean somal diameter of DIR DLM-projecting neurons in Area X-CORE (n = 54) was larger than those in Area X-SHELL (n = 35) (*p < 0.0001; unpaired t-test). (h) Bar graph showing that 467/469 DIR neurons in Area X-CORE and SHELL of 4 adult birds did not express FoxP2 (p < 0.0001, binomial test). Percentages represent number of DIR neurons that expressed FoxP2 over total number of DIR neurons in each bird.

Figure 4. DIR neurons in Area X of juvenile songbirds do not express FoxP2. (a) Injection of a fluorescent tracer into DLM (outlined by dashed line) produced retrograde label in Area X-CORE and SHELL of a juvenile bird (d-f). (b) DAPI staining outlines Area X-CORE (inner dashed circle) within Medial Striatum (the border between striatum and cortex is demarcated by gray line, which corresponds to the axon tract PSL, pallial-subpallial lamina). As in adult birds, Area X-SHELL (outer dashed circle) was conservatively defined as extending 500 μm from the medial, ventromedial, and ventrolateral edges of Area X-CORE. (c) Red FoxP2 staining (Alexa

594) using 16046 antibody and (d) green DLM-projecting neurons (Alexa 488) in the same section as (b). (e) Overlay of DAPI, FoxP2, and DLM-projecting dyes. (f) Enlarged image of boxed region in (e) demonstrating that red FoxP2 label and green retrograde label in an Area X-CORE DIR neuron did not colocalize. (g) Box plot showing that the mean somal diameter of DIR DLM-projecting neurons in Area X-CORE (n = 38) was larger than those in Area X-SHELL (n = 26) (*p = 0.0016; unpaired t-test). (h) Bar graph detailing that no DIR neurons in three juvenile birds (n = 87) showed FoxP2 expression (p < 0.0001, binomial test). Percentages represent number of DIR neurons that expressed FoxP2 over total number of DIR neurons in each bird.

Figure 5. Average diameters of FoxP2+ nuclei in Area X-CORE and SHELL revealed continuous, positively skewed size distributions. FoxP2 staining distinguished sparsely distributed large nuclei (white arrowheads) and numerous small nuclei (white arrows) in Area X-CORE of (a) adult and (c) juvenile songbirds. This pattern was also present in Area X-SHELL (not shown). Most small FoxP2+ neurons were less intensely stained, whereas larger nuclei were more intensely stained. (b) Histograms showing average diameter of FoxP2+ neurons in Area X-CORE (black, n = 3953) and SHELL (gray, n = 1535) of three adult birds (top) and two juvenile birds (bottom; CORE, black, n = 1813; SHELL, gray, n = 835).

Figure 6. All DIR and most putative IND neurons did not express the striatal marker DARPP-32 in juvenile songbirds. (a) Alexa 488-labeled DIR DLM-projecting neurons in juveniles (one shown from Area X-CORE) did not colocalize with magenta cytoplasmic DARPP-32 label. (b) Most large (≥8 μm) putative IND FoxP2+ nuclei across both Area X-CORE and SHELL did not express DARPP-32; two examples from Area-X CORE are indicated by yellow arrowheads. A majority of small (<8 μm) FoxP2+ nuclei expressed DARPP-32 (two examples indicated by yellow arrows). Colocalization of both proteins is depicted in white. (c) Some large (≥8 μm) FoxP2+ nuclei expressed DARPP-32 in both CORE and SHELL regions of Area X; yellow arrowhead indicates a large FoxP2+/DARPP-32+ neuron from Area-X SHELL. An adjacent small (<8 μm; yellow arrow) FoxP2+ nucleus also expressed DARPP-32. White voxels indicating colocalization show that the FoxP2+ nuclei of the putative IND and medium spiny neuron take up a large percentage of the DARPP-32+ cytoplasm. (d) Bar graph detailing the number of large FoxP2+ neurons that show DARPP-32 expression in Area X-CORE (17/48

neurons) and Area X-SHELL (15/48 neurons) from two juvenile songbirds when a nuclear diameter cutoff of ≥8 μm was applied. Percentages represent number of large FoxP2+/DARPP32+ nuclei over total number of large FoxP2+ nuclei across CORE and SHELL subregions for each bird.

Figure 7. DLM-projecting DIR pallidal cell and adjacent FoxP2+ neuron have similar DAPI+ nuclear sizes. The nucleus of a green Alexa 488+ retrogradely-labeled cell is defined by blue DAPI nuclear stain in Area X-CORE of an adult bird. DAPI+ nucleus of DIR neuron encompasses a smaller percentage of the soma in comparison to putative IND large FoxP2+/DARPP-32+ cells (see Fig. 6c); this pattern was also present in Area X-SHELL (not shown). The adjacent red FoxP2+ nucleus may represent an interneuron or a medium spiny neuron.