Enantioselective photoenzymatic C(sp^3)-H functionalization via single-electron oxidation of carbanions

Jin Zhu[1,2,6], Qiaoyu Zhang[3,6], Tao Gu[2,6], Binbin Chen[2], Mingzhe Ma[2], Xiaoyu Wang[3], Xiao Liu[2], Mingjie Ma[2], Binju Wang[3*], and Yajie Wang[2,4,5*]

[1]School of Materials Science and Engineering, Zhejiang University, Hangzhou, Zhejiang 310058, China.

[2]School of Engineering, Westlake University, Hangzhou 310030, China.

[3]Key Laboratory of Theoretical and Computational Chemistry, College of Chemistry and Chemical Engineering, Xiamen University, Xiamen 361005, China.

[4]School of Life Science, Westlake University, Hangzhou 310030, China.

[5]Westlake Center of Synthetic Biology and Integrated Bioengineering, Westlake University, Hangzhou 310030, China.

[6]These authors contributed equally: Jin Zhu, Qiaoyu Zhang, and Tao Gu.

*Corresponding author Emails: wangbinju2018@xmu.edu.cn;

wangyajie@westlake.edu.cn

Abstract

C-H functionalization provides unparalleled benefits for the late-stage modification of

complex molecules. In recent years, photocatalysis has progressed significantly due to

its mild conditions, sustainability and selectivities. Current mechanisms for

photocatalytic C-H functionalization primarily involve direct single-electron oxidation

and direct hydrogen atom transfer (d-HAT) by radicals, both of which are invalid for

the functionalization of highly electron-deficient C(sp^3)-H bonds. Here, we developed

a cooperative photoenzymatic system consisting of a flavin-dependent 'ene'-reductase

(ER) and an exogenous photocatalyst fluorescein (Fl) to achieve enantiodivergent

functionalization of electron-deficient C(sp^3)-H bonds. Mechanistic studies revealed a

novel pathway for radical intermediate formation via excited-state Fl*-induced single-

electron oxidation of carbanions under alkaline conditions. The overall catalytic

efficiency is enhanced by the electron transfer (ET) between FMN_{ox} and Fl$^{\bullet}$, while the

stereoselectivity is controlled by ERs through enantioselective hydrogen atom transfer

(HAT). This suggests that thermodynamically unfavorable single-electron oxidation of

highly electron-deficient species can occur through ion-to-radical conversion. We

anticipate that our novel mode of photoenzymatic catalysis will inspire new pathways

for generating free radical intermediates and foster innovative strategies for achieving

photoenzymatic new-to-nature reactions.

- Enantioselective functionalization of electron-deficient C(sp^3)-H bonds
- Radical formation by single-electron oxidation of carbanion
- The synergistic action of fluorescein and enzyme
- Enantiodivergent synthesis

Introduction

C-H functionalization, particularly $C(sp^3)$-H, is pivotal in synthetic chemistry for its efficiency in constructing complex molecules from simple precursors, high atom economy, and versatility in late-stage modification of diverse precursors in pharmaceuticals and materials science.[1,2] Visible-light photoredox catalysis has rapidly emerged as a robust strategy in this context due to its catalytic versatility, mild conditions, sustainability, and high selectivities.[3-6] Photocatalytic $C(sp^3)$-H functionalization primarily relies on two mechanisms to generate carbon-centered radicals: direct single-electron oxidation of C-H bonds[4] and direct hydrogen atom transfer (*d*-HAT) (Fig. 1a)[5].The former requires electron-rich, easily oxidizable C-H bonds[7,8], while the latter depends on the hydrogen abstraction by polarity-matched radical species[3,9-12]. Both mechanisms pose challenges for selective electron-deficient $C(sp^3)$-H bonds functionalization. In addition, asymmetric $C(sp^3)$-H functionalization via hydrogen atom transfer (HAT) with prochiral radicals presents challenges, partly due to the small size of the H atom and the early transition state associated with this process.[13,14] Therefore, photocatalysis can have great potential in the enantioselective functionalization of electron-deficient $C(sp^3)$-H bonds, but it also presents substantial challenges.

Recent advancements in photoenzymatic catalysis have shown remarkable enantioselectivity in the asymmetric HAT of prochiral radicals[15-30]. The radicals are produced either through the single-electron reduction of an electron-donor-acceptor (EDA) complex[15-26] or via the single-electron oxidation of electron-rich precursors[27-30].

Notably, these methods have been successfully applied in enantioselective radical

cyalization[15,16], hydroalkylation[17,23-27], hydroamination[18], hydrosulfonylation[21,27,28] of

olefins, and cross-electrophile coupling[20]. Most recently, Huang accomplished

enantioselective photoenzymatic C(sp^2)-H functionalization through the single-electron

oxidation of electron-rich arenes and alkenes[29,30] with low oxidation potentials (e.g., 3-

methoxythiophene at +1.42V vs. the saturated calomel electrode (SCE))[31] (Fig. 1b).

However, electron-deficient C(sp^3)-H bonds present significant challenges for direct

oxidation[32], and their high bond dissociation energy (BDE) further limits C-H bond

cleavage. For instance, the BDE of the C-H bond in malononitrile is as high as 87

kcal/mol[33]. Therefore, developing a novel strategy for the photoenzymatic

enantioselective functionalization of electron-deficient C(sp^3)-H bonds is of great

scientific importance.

We noticed that in the Bamford-Stevens reaction, diazo compounds can be

transformed into carbonium ions in protic solvents[34], which might provide a theoretical

basis for the generation of carbon-centered free radicals from diazo compounds via

proton-coupled electron transfer (PCET)[35-37]. In light of this, we propose that generating

ions as intermediates for radical formation could effectively facilitate reactions with

substrates that do not readily form radicals. Building on this concept, we developed a

cooperative photoenzymatic system comprising an ene-reductase (ER) and an

organophotoredox catalyst to achieve the asymmetric functionalization of electron-

deficient C(sp^3)-H bonds. In the catalytic process, carbanions are generated under

alkaline conditions and oxidized to radicals via an excited-state photocatalyst-induced

single-electron transfer (SET). Stereoselectivity is then accomplished via an

enantioselective HAT step of prochiral intermediates mediated by ER (Fig. 1c). This

photoenzymatic system not only addresses a synthetic challenge unmet by either

traditional organocatalysis[38-41] or photoenzymatic catalysis but also introduces an

emerging mechanism for radical formation, one that has been rarely reported in photo-

redox catalysis.

a. Photocatalytic C(sp^3)-H functionalization

Direct single-electron oxidation of electron-rich precursors

Photoredox Catalysis

FG: functional group

Direct hydrongen atom transfer (d-HAT) of polarity-matched radicals

b. Photoenzymatic enantioselective C(sp^2)-H functionalization

c. Our hypothesis: enantioselective functionalization of electron-deficient C(sp^3)-H bonds (this work)

87   **Fig. 1| Mode of photocatalytic C-H functionalization and our hypothesis.**

89   **Results and Discussion**

Reaction development and engineering

Carbanions are rarely used as precursors for radical formation via single-electron oxidation in photoredox or photoenzymatic catalysis.[42] The main challenges is their high reactivity, which complicates regulating competition between single-electron oxidation and inherent activities like nucleophilic addition to unsaturated bonds[43]. Highly electrophilic radicals can also quickly convert into carbanions through radical polar crossover (RPC).[44] To develop a photoenzymatic system for the enantioselective functionalization of electron-deficient $C(sp^3)$-H bonds, suitable enzymes must remain active in alkaline conditions necessary for carbanion formation. The excited ER-FMN_{ox}^* must effectively oxidize carbanions into free radicals while minimizing undesirable pathways, such as carbanion nucleophilic addition, reductive quenching of FMN_{ox}^* by the reaction buffer or nearby amino acids, and RPC (Supplementary Fig. 6).

We investigated the proposed reactions using malononitrile **1a** and α-methylstyrene **2a** as model substrates with a library of ERs at 1 mol% under 455 nm blue LED illumination. Most ERs showed no catalytic activity (Table 1, entry 1 and Supplementary Table 3), but MorB, OYE2, SYE1, SYE2, SYE3, and GluER achieved yields of 1.2% to 3%. Notably, SYE2 and GluER exhibited the best and opposite enantioselectivity for **3a**. GluER had (*S*)-selectivity with a 4:96 enantiomeric ratio (er) (Table 1, entry 2), while SYE2 showed (*R*)-selectivity with a 93.3:6.6 er (Table 1, entry 3). Thus, SYE2 and GluER were chosen for further investigation.

To enhance the yield of **3a**, we tested various reaction conditions, including buffer systems, pH levels, and co-solvents (Supplementary Tables 4-5). Although these

adjustments did not significantly improve the yield, we decided to proceed with Tris

buffer at pH 9 and ethanol as a co-solvent. Notably, the reaction only occurs at pH 9 or

above, indicating that strongly alkaline conditions are essential for initiating carbanion

formation in the catalytic cycle (Table 1, entries 4-6).

Unlike other photoenzymatic systems[28,29], adding extra FMN-Na did not enhance

the yield of **3a** (Table 1, entry 7). This suggests that $FMN_{ox}{}^*$-induced single-electron

oxidation of carbanions may be the rate-limiting step, indicating the need for

photocatalysts with higher oxidative potential. We screened 18 transition metal and

organic photocatalysts (Supplementary Tables 6-7) and identified fluorescein (FI-PC7),

PC9, and PC12 as effective, increasing the yield of **3a** to 27%, 21%, and 19%,

respectively, in both SYE2 and GluER-catalyzed reactions. The combination of 5 mol%

FI with 1 mol% SYE2 or GluER gave the highest yields of **3a**, at 47.8% and 45.5%,

respectively, without compromising enantioselectivity (Table 1, entries 8-9). Control

experiments without light or in the presence of air did not produce the target product

(Table 1, entries 10-11), supporting the visible-light-induced radical mechanism.

Although neither FMN nor FI alone could catalyze the transformation (Table 1, entries

12-13), combining 10 mol% of both resulted in a 3% racemate of **3a** (Table 1, entry 14).

These findings indicate that FI and FMN cooperate to catalyze the $C(sp^3)$-H

functionalization process, highlighting the essential role of photoenzymes in improving

efficiency and controlling the stereochemistry of radical processes.

Table 1 | Optimization of reaction conditions for photoenzymatic enantioselective

$C(sp^3)$-H functionalization.

ER 1mol%,
Fluorescein 5mol%
455nm LEDs

Tris buffer (50 mM, pH 9)
EtOH (7.5 v/v%)
N$_2$, room temperature, 12h
Standard conditions

NC CN + Ph → 3a

1a, 2 equiv   2a, 0.004 mmol

Fluorescein
FI

Entry	ER	Deviation from standard conditions[a]	Yield (%)[b]	er (%) (R:S)[c]
1	ERs[d]	w/o FI	n.r.	n.a.
2	GluER	w/o FI	1.3	4:96
3	SYE2	w/o FI	2.6	93.3:6.6
4	SYE2	Tris buffer pH 7.4, w/o FI	n.r.	n.a.
5	SYE2	Tris buffer pH 8, w/o FI	n.r.	n.a.
6	SYE2	Tris buffer pH 10, w/o FI	1.5	85:13
7	SYE2	w/o FI, 10 mol% FMN-Na	2.6	93.3:6.6
8	**SYE2**	**None**	**47.8**	**95.6:4.4**
9	**GluER**	**None**	**45.5**	**1.2:98.8**
10	SYE2	Dark	n.r.	n.a.
11	SYE2	Air	n.r.	n.a.
12	/	FMN-Na instead FI	n.r.	n.a.
13	/	None	n.r.	n.a.
14	/	10 mol% FMN-Na + FI	3	racemate

[a]Conditions: the total volume of standard reaction was 400 uL. [b]Yield was determined by gas chromatography. c er was determined by HPLC analysis on a chiral stationary phase, and the absolute configuration was established by theoretical calculations of electronic circular dichroism (ECD); see Supplementary Fig. 31 and Fig. 32. [d]LacER, KYE1, OYE1, OYE3, OPR1, SYE4, TOYE and YqjM were tested. n.r. No reaction; n.a. Not applicable.

Substrate scope investigations

We explored the substrate scope using SYE2 (ER1) and GluER to access both enantiomers of the products. While both enzymes showed excellent enantioselectivity for **3a**, this decreased significantly for most derivatives (Supplementary Fig. 33). To address this, we engineered GluER using site-saturated mutagenesis. Molecular

docking of GluER's crystal structure with **1a** and **2a** identified key residues in the

substrate channel and FMN binding region, including T36, W66, H172, N175, Y177,

T231, Q232, P239, R261, F269, and Y343 (see Supplementary Fig. 2). Site-saturated

mutagenesis of these 11 sites led to GluER-Y343A (ER2), which showed improved

enantioselectivity for both **3a** (up to R:S = 99.5:0.5) and other derivatives (Fig. 2).

The photoenzymatic system enabled hydroalkylation of various substituted methyl

styrenes with functional groups at the para-/meta-/ortho positions of the phenyl group

(**3a-j**), producing stereodivergent products with moderate to good yields and up

to >1:99 er. Although 1-bromo-2-(prop-1-en-2-yl)benzene (**3k**) had a relatively low

yield, it exhibited good to excellent enantioselectivity. Additionally, 2-naphthyl (**3l**),

benzothiophene (**3m**), and biphenyl (**3n**) demonstrated the compatibility of

photoenzymatic catalysis with aromatic rings and heterocycle. Variations at the α

position of styrene had minimal impact on outcomes, as seen with **3p**. However,

replacing aromatic olefins with aliphatic olefins significantly reduced the yield of **3o**.

Although the mutant GluER-W66Q-Y343A (ER3) improved the yield of **3o**, the

enantioselectivity remained unsatisfactory. Beyond malononitrile **2a**, dimethyl

malonate **2b** could also alkylate with **3q** with excellent enantioselectivity. Moreover,

this photoenzymatic catalysis can be successfully scaled up to 1 mmol with minimal

changes in yield and enantioselectivity.

Fig. 2 | Substrate scope of enantioselective functionalization of electron-deficient C(sp^3)-H bonds via hydroalkylation catalyzed by cooperative photoenzymatic system. Conditions: 1 (0.008 mmol), 2 (0.004 mmol), ER (1 mol%), FI (5 mol%), and EtOH (7.5 v/v%) in Tris buffer (50 mM, pH 9) were stirred at room temperature for 12 hours under N_2 with 455 nm LEDs. The total reaction volume was 400 μL. Yields were determined by gas chromatography, and the er by HPLC on a chiral stationary phase. [#] Isolated yield was obtained by scaling up the reaction.

Mechanistic investigations

Based on the experimental results, we proposed a synergistic photoredox and biocatalytic mechanism for new-to-nature photoenzymatic C(sp^3)-H functionalization (Fig. 3a and detailed in Supplementary Fig. 22). In this catalytic cycle, malononitrile **1a** is first deprotonated in alkaline solutions to form carbanion intermediate **Int. A**. The

light-excited photocatalyst FI* then oxidizes **Int. A** via SET, producing malononitrile

free radical **Int. B** and fluorescein radical anion FI$^{-\bullet}$. **Int. B** subsequently attacks α-

methylstyrene **2a**, forming a C-C bond and generating prochiral radical intermediate

Int. C, alongside FMN_{sq} from FMN_{ox} capturing an electron from FI$^{-\bullet}$. Finally, **Int. C**

undergoes ER-mediated enantioselective HAT, yielding enantioenriched **3a**, while

FMNH$^{\bullet}$ is oxidized back to FMN_{ox}, completing the catalytic cycle.

Several experiments were performed to support the proposed mechanism. The

photophysical properties of SYE2 and FI were measured to identify visible-light-

excited species. UV-vis absorption spectra show that SYE2 and FI absorb in the ranges

of 400-520 nm and 420-530 nm, with maximum absorptions at 460 nm and 490 nm,

respectively (Fig. 3b). This confirms that FI can be effectively excited by 455nm LED

to initiate the catalytic cycle. Stern-Volmer luminescence quenching experiments also

support the initial reductive quenching of excited FI* by **1a** (**Int. A** → **Int. B**), as the

fluorescence intensity of FI decreased upon adding **1a**, unlike with **2a** (Supplementary

Fig. 10).

Several experiments were also performed to verify the generated radical

intermediates. Firstly, the radical trapping experiment was performed by adding 2

equivalents of 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO) to the reaction mixture

under standard conditions. Only trace amounts of **3a** were detected, as the prochiral

radical intermediate **Int. C** was trapped by TEMPO, confirmed by high-resolution mass

spectrometry (HRMS) (Fig. 3c). Additionally, when α-cyclopropyl styrene **4** was

subjected to the photoenzymatic system, the ring-opening product **5** was observed as

the major product, supporting the involvement of **Int. C** (Fig. 3d). To further investigate

the photo-oxidation step and clarify the resulting radical intermediates, electron

paramagnetic resonance (EPR) experiments were conducted. With 5,5-dimethyl-1-

pyrroline *N*-oxide (DMPO) as a radical trapping agent, slight changes in EPR signals

were observed in the presence of **1a** compared to **2a** (Fig. 3b), indicating the formation

of a potential **1a**-derived radical (**Int. B**), consistent with quenching experiments. The

difference in EPR signals with and without **2a** suggests that the malononitrile radical

Int. B, generated from the single-electron oxidation of the unstable carbanion **Int. A**,

rapidly reacts with **2a** to form a more stable prochiral benzyl radical **Int. C** before being

captured.[45]

Since apo-SYE2 (SYE2 without FMN) and FI alone did not catalyze the product

formation, and adding extra FMN restored reactivity, we confirmed that the catalytic

cycle requires the cooperative action of ER_FMN and FI (Fig. 3e and detailed in

Supplementary Table 8 and Fig. 17). Finaly, deuterium-labelling experiments were

conducted to investigate the final termination step (Fig. 3f). Using a deuterated buffer,

the hydrogen at the stereogenic center was labelled at 76% D, suggesting that water

may be involved in HAT steps.

Fig. 3 | Mechanistic experiments and proposed catalytic cycle. a, Proposed catalytic cycle. **b**, UV-vis absorption (abs) spectra and room-temperature EPR experiments. **c**, TEMPO trapping experiment. **d**, Radical clock experiment. **e**, Photoenzymatic reactions catalysed by apo-flavoprotein. **f**, Deuterium-labelling experiment.

Computational studies

Extensive computational studies were performed to rationalize the mechanism proposed in Fig. 3a. Using hybrid cluster-continuum (HCC) model calculations, we found that **1a** undergoes facile deprotonation by OH^- ions in aqueous solution, forming the anionic species **Int. A.** with a low energy barrier of 0.3 kcal/mol (Supplementary Figure 20). To ascertain that **Int. A**, rather than **1a**, serves as the reactant for the generation of the carbanion intermediates **Int. B**, density functional theory (DFT)

calculations and time-dependent DFT (TD-DFT) calculations were conducted. The

Gibbs reaction energy (ΔG) for the excited FI*-mediated oxidation of **1a** and **Int. A**

were calculated to be +105.2 kcal/mol and -32.1 kcal/mol (Fig. 4a and Supplementary

Fig. 15), respectively, supporting that photoinduced single-electron oxidation process

occurs through carbanion intermediates **Int. A**.

Since EPR experiments with FI, **1a**, and **2a** revealed distinctive radical signals,

suggesting that the formation of **Int. C** might be independent on ER participation.

Quantum mechanics (QM) model calculations show that the radical addition between

Int. B and **2a** has an energy barrier of 3.0 kcal/mol and an exothermicity of 10.7

kcal/mol (Fig. 4b), suggesting that formation of **Int. C** is quite facile. In addition,

deuteration experiments and the pKa value of FMN_{sq} (~8.5) suggested that the

protonation of FMN_{sq} to form neutral FMNH$^\bullet$ is feasible at pH 9, with a Gibbs reaction

energy (ΔG) of ~0.7 kcal/mol. Starting from FMNH$^\bullet$, HAT from FMNH$^\bullet$ to **Int. C** is

expected to yield the final product **3a**.

To elucidate the opposing enantioselectivity of GluER and SYE2, **Int. C** and FI

were docked into the enzyme active sites. Molecular dynamics (MD) simulations

indicate that FI was stabilized by Arg261 in GluER (Supplementary Fig. 23). **Int. C**

adopts two distinct conformations (CH$_3$-left and CH$_3$-right) in GluER, corresponding

to the *R* and *S* configurations, respectively. Moreover, the quantum mechanics

/molecular mechanics (QM/MM)-calculated HAT barriers are 16.2 kcal/mol for the

CH$_3$-right and 23.2 kcal/mol for the CH$_3$-left, aligning with the observed dominant

product (*S*)-**3a** (Fig. 4c and Fig. 4d). This is likely due to steric repulsion between the

bulky-CN substituent of **Int. C** and FMNH$^•$ in the CH$_3$-left conformation, which tends to enlarge the distance between the C3 atom of **Int. C** and the N5-H of FMNH$^•$ (4.10 Å in CH$_3$-left vs. 3.44 Å in CH$_3$-right), thereby increasing the energy barrier for HAT in CH$_3$-left conformation.

Unlike GluER, MD simulations reveal that the bulky-CN substituents in the CH$_3$-left conformation of **Int. C** forms a stable hydrogen bond with Tyr180 in SYE2, which positions bulky-CN away from FMNH$^•$ and facilitates the subsequent HAT process, that is favor of generating (*R*)-**3a** (Supplementary Fig. 26). Indeed, our calculations suggest that the HAT process in SYE2 involves a barrier of 17.2 kcal/mol in the CH$_3$-left conformation and 22.0 kcal/mol in the CH$_3$-right conformation, leading to the dominant formation of (*R*)-**3a** that agrees with experiments (Supplementary Fig. 26 and Fig. 27).

Furthermore, the newly formed FI$^{-•}$ could reduce FMN$_{ox}$ to FMN$_{sq}$ and release the initial ground state FI. DFT calculations show that FI$^{-•}$ has the lowest redox potential among all functional photocatalysts (PC1, PC2, PC5, FI, PC10, and PC12), making it highly prone to transfer an electron to FMN$_{ox}$ to form FMN$_{sq}$ (Supplementary Fig. 16). This suggests that ET between FMN$_{ox}$ and FI$^{-•}$ enhances the whole catalytic process, explaining why FI is the most effective among all tested photocatalysts.

Fig. 4 | Mechanistic studies. a. SET from the excited FI* to **Int. A**. **b**, The radical addition reaction between **Int. B** and **2a**. **c**, QM/MM-calculated energy of the HAT process from FMNH• to the **Int. C** (CH$_3$-right) result in the production of S-(**3a**). **d**, QM/MM-calculated energy of the HAT process from FMNH• to the **Int. C** (CH$_3$-left) result in the production of R-(**3a**).

Conclusions

In summary, we have developed a synergistic photoenzymatic system for the enantioselective functionalization of electron-deficient C(sp^3)-H bonds through radical intermediates generated by photoinduced single-electron oxidation of carbanions. This

enantiodivergent strategy, demonstrated in 17 cases with enantiomeric ratios up

to >1:99 and yields of up to 83%, offers potential for the one-step construction of

dicyano and diester compounds with remote stereocenters. Detailed mechanistic

experiments confirmed that the excited-state photocatalyst FI^* performs single-electron

oxidation of carbanion rather than directly oxidizing the electron-deficient

malononitrile **1a** to generate free radical intermediates. Unlike previous

photoenzymatic systems, the ET between the additional photocatalyst FI and FMN_{ox}

ensures catalytic cycle's smooth operation. This novel mode of photoenzymatic

catalysis is expected to inspire new strategies for forming free radical intermediates and

enabling new-to-nature biotransformation.

Methods

General procedure for photoenzymatic reactions

Taking **1a** + **2a** → **3a** as an example, in a glovebox (O_2 concentration <5 ppm), ,

solutions of ene-reductase (1 mol%), FI (10 μL, 2 mM stock in EtOH, 5 mol%), and

substrates mix (20 μL, 400 mM **1a** and 200 mM **2a** in EtOH) were added to a 4 mL vial

containing a magnetic stir bar and Tris buffer (50 mM, pH 9). The total volume of the

reaction mixture was 400 μL and the final concentrations for EtOH was 7.5%. The vial

was sealed with a screw cap, removed from the glove box, illuminated with blue LEDs,

and stirred for 12 h with a cooling fan (see Supplementary Fig. 3 for reaction setup).

For the reaction where the yield was determined by gas chromatography (GC), EtOAc

(1.0 mL) and 10.0 μL of an internal standard stock (20 mg/mL of n-dodecane in EtOAc)

were added to the reaction mixture and mixed thoroughly. For the reaction where the

yield was determined by reverse phase HPLC, MeCN (1.0 mL) and 10.0 μL of an

internal standard stock (16.8 mg/mL of 1,3,5-trimethoxybenzene in MeCN) were added

to the reaction mixture and mixed thoroughly. The organic phase was separated and

then analyzed by GC (or reverse phase HPLC), GC-MS and chiral HPLC. Product

formation was confirmed by comparing retention times in GC/HPLC, as well as mass

spectra and [1]H-NMR spectra of the crude material with those of the reference. Yield

determined by GC was relative to the n-dodecane internal standard, while yield

determined by reverse phase HPLC was relative to the 1,3,5-trimethoxybenzene

internal standard. Enantioselectivity was determined by chiral HPLC.

System setup and MD simulations

The initial wild-type structure of GluER, was constructed based on the crystal structure

(PDB code: 6O08). The SYE2 enzyme structure was modeled using AlphaFold2, and

the position of FMN was confirmed by superimposing it with the GluER enzyme. Then,

the **Int. C** and FI were docked into the active sites using AutoDock Vina tool. 100 ns

MD simulations under the NPT ensemble. All MD simulations were performed with

GPU version of Amber 18 package.

QM/MM calculations for enzymatic reactions

A representative snapshot extracted from the clustering results in the MD trajectory was

used for subsequent QM/MM calculations. All QM/MM calculations were performed

using ChemShell combining Turbomole for the QM region and DL_POLY for the MM

region. The electronic embedding scheme was used to account for the polarizing effect

of the enzyme environment on the QM region.In all QM/MM geometry optimizations,

the QM region was treated with the B3LYP functional using the double-zeta basis set

def2-SVP for all atoms (labeled as B1). The energies were further corrected with the

larger basis set def2-TZVP for all atoms (labeled as B2). Grimme's D3BJ empirical

dispersion was included in all calculations.

DFT Calculations

All DFT calculations were performed with the Gaussian 16, Revision A.03 software.

The initial structures were first optimized in conjunction with the SMD continuum

solvation model at the B3LYP-D3/6-31G(d) level of theory. The solvation energies

were further calculated at the M052X/6-31G* for all atoms. In particular, the solvent of

water was used to simulate the reaction system.

TDDFT Calculations

The initial structures were first optimized at B3LYP/6-31G(d) level. Then, the single-

point TDDFT calculations at wB97XD/6-311+G(d,p) levels were performed to

compute the three low-lying singlet excited states. All TDDFT calculations were

performed with Gaussian 16, Revision A.03.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting

Summary linked to this article.

Data availability

Data relating to the materials and methods, experimental procedures, mechanistic

studies and computational calculations, HPLC spectra, and NMR spectra are available

in the Supplementary Information or from the authors on reasonable request. The

atomic coordinates of the optimized computational models are available in the

Supplementary Data.

References

1. Yamaguchi, J., Yamaguchi, A. D. & Itami, K. C-H bond functionalization: emerging

synthetic tools for natural products and pharmaceuticals. *Angew. Chem. Int. Ed.* **51**,

8960-9009 (2012).

2. Leibler, I. N. M., Tekle-Smith, M. A. & Doyle, A. G. A general strategy for C(sp^3)-H

functionalization with nucleophiles using methyl radical as a hydrogen atom abstractor.

Nat. Commun. **12**, 6950-6959 (2021).

3. Li, Y., Lei, M. & Gong, L. Photocatalytic regio- and stereoselective C(sp^3)-H

functionalization of benzylic and allylic hydrocarbons as well as unactivated alkanes.

Nat. Catal. **2**, 1016-1026 (2019).

4. Bellotti, P., Huang, H.-M., Faber, T. & Glorius, F. Photocatalytic late-stage C-H

functionalization. *Chem. Rev.* **123**, 4237-4352 (2023).

5. Capaldo, L., Ravelli, D. & Fagnoni, M. Direct photocatalyzed hydrogen atom transfer

(HAT) for aliphatic C-H bonds elaboration. *Chem. Rev.* **122**, 1875-1924 (2022).

6. Yi, H. et al. Recent advances in radical C-H activation/radical cross-coupling. *Chem.*

Rev. **117**, 9016-9085 (2017).

7. Miyake, Y., Nakajima, K. & Nishibayashi, Y. Visible-light-mediated utilization of α-

aminoalkyl radicals: addition to electron-deficient alkenes using photoredox catalysts.

J. Am. Chem.Soc., **134**, 3338-3341 (2012).

8. Loh, Y. Y. et al. Photoredox-catalyzed deuteration and tritiation of pharmaceutical

compounds. *Science* **358**, 1182-1187 (2017).

9. Cai, C.-Y., et al. Photoelectrochemical asymmetric catalysis enables site- and

enantioselective cyanation of benzylic C-H bonds. *Nat. Catal.* **5**, 943-951 (2022).

10. Shaw, M. H., Shurtleff, V. W., Terrett, J. A., Cuthbertson, J. D. & MacMillan, D. W.

C. Native functionality in triple catalytic cross-coupling: sp^3 C-H bonds as latent

nucleophiles. *Science* **352**, 1304-1308 (2016).

11. Xu, A., Guo, J.-J., Pan, H. & Zuo, Z. Selective functionalization of methane, ethane,

and higher alkanes by cerium photocatalysis. Science 361, 668-672 (2018).

12. Lei, G., Xu, M., Chang, R., Funes-Ardoiz, I. & Ye, J. Hydroalkylation of unactivated

olefins via visible-light-driven dual hydrogen atom transfer catalysis. *J. Am. Chem. Soc.*

143, 11251-11261 (2021).

13. Nagib, D. A. Asymmetric catalysis in radical chemistry. *Chem. Rev.* **122**, 15989-

15992 (2022).

14. Mondal, S. et al. Enantioselective radical reactions using chiral catalysts. *Chem. Rev.*

122, 5842-5976 (2022).

15. Biegasiewicz, K. F. et al. Photoexcitation of flavoenzymes enables a stereoselective radical cyclization. *Science* **364**, 1166-1169 (2019).

16. Black, M. J. et al. Asymmetric redox-neutral radical cyclization catalysed by flavin-dependent 'ene'-reductases. *Nat. Chem.* **12**, 71-75 (2020).

17. Fu, H. et al. Ground-state electron transfer as an initiation mechanism for biocatalytic C-C bond forming reactions. *J. Am. Chem. Soc.* **143**, 9622-9629 (2021).

18. Ye, Y. et al. Using enzymes to tame nitrogen-centred radicals for enantioselective hydroamination. *Nat. Chem.* **15**, 206-212 (2023).

19. Gao, X., Turek-Herman, J. R., Choi, Y. J., Cohen, R. D. & Hyster, T. K. Photoenzymatic synthesis of α-tertiary amines by engineered flavin-dependent 'ene'-reductases. *J. Am. Chem. Soc.* **143**, 19643-19647 (2021).

20. Fu, H., et al. An asymmetric sp^3-sp^3 cross-electrophile coupling using 'ene'-reductases. *Nature* **610**, 302-307 (2022).

21. Chen, X. et al. Photoenzymatic hydrosulfonylation for the stereoselective synthesis of chiral sulfones. *Angew. Chem. Int. Ed. Engl.* **62**, e202218140 (2023).

22. Chen, B. et al. Modular access to chiral amines via imine reductase-based photoenzymatic catalysis. *J. Am. Chem. Soc.* **20**, 14278-14286 (2024).

23. Huang, X. et al. Photoenzymatic enantioselective intermolecular radical hydroalkylation. *Nature* **584**, 69-74 (2020).

24. Li, M., Harrison, W., Zhang, Z., Yuan, Y. & Zhao, H. Remote stereocontrol with azaarenes via enzymatic hydrogen atom transfer. *Nat. Chem.* **16**, 277-284 (2024).

25. Huang, X. et al. Photoinduced chemomimetic biocatalysis for enantioselective

intermolecular radical conjugate addition. *Nat. Catal.* **5**, 586-593 (2022).

26. Li, M., Yuan, Y., Harrison, W., Zhang, Z. & Zhao, H. Asymmetric photoenzymatic

incorporation of fluorinated motifs into olefins. *Science* **385**, 416-421 (2024).

27. Shi, Q. et al. Single-electron oxidation-initiated enantioselective hydrosulfonylation

of olefins enabled by photoenzymatic catalysis. *J. Am. Chem. Soc.* **146**, 2748-2756

(2024).

28. Jiang, L. et al. Photoenzymatic redox-neutral radical hydrosulfonylation initiated

by FMN. *ACS Catal.* **14**, 6710-6716 (2024).

29. Zhao, B. et al. Direct visible-light-excited flavoproteins for redox-neutral

asymmetric radical hydroarylation. *Nat. Catal.* **6**, 996-1004 (2023).

30. Yu, J. et al. Repurposing visible-light-excited ene-reductases for diastereo-and

enantioselective lactones synthesis. *Angew. Chem. Int. Ed. Engl.* **63**, e202402673

(2024).

31. Chen, B.-H., Du, Y.-D. & Shu, W. Organophotocatalytic regioselective C-H

alkylation of electron-rich arenes using activated and unactivated alkenes. *Angew.*

Chem. Int. Ed. Engl. **61**, e202200773 (2022).

32. Poudel, D. P., Pokhrel, A., Tak, R. K., Shakar, M. & Giri, R. Photosensitized O_2

enables intermolecular alkene cyclopropanation by active methylene compounds.

Science **381**, 545-553 (2023).

33. Goebbert, D. J., Velarde, L., Khuseynov, D. & Sanov. A. C-H bond dissociation

energy of malononitrile. *J. Phys. Chem. Lett.* **1**, 792-795 (2010).

34. Bayless, J. H., Friedman, L., Cook, F. B. and Shechter, H. Effect of solvent on the

course of the Bamford-Stevens reaction. *J. Am. Chem. Soc.* **90**, 531-533 (1968).

35. Huang, X., Webster, R. D., Harms, K & Meggers, E. Asymmetric catalysis with

organic azides and diazo compounds initiated by photoinduced electron transfer. *J. Am.*

Chem. Soc. **138**, 12636-12642 (2016).

36. Duan, X. et al. A photoenzymatic strategy for radical-mediated stereoselective

hydroalkylation with diazo compounds. *Angew. Chem. Int. Ed. Engl.* **62**, e e202214135

(2023).

37. Gu, Y.-J., Luo, M.-P., Yuan, H., Liu, G.-K. & Wang, S.-G. Photocatalytic

enantioselective radical cascade multicomponent Minisci reaction of β-carbolines using

diazo compounds as radical precursors. *Adv. Sci.* **11**, 2402272 (2024).

38. Guo, H., Fan, Y., Sun, Z., Wu, Y. & Kwon, O. Phosphine organocatalysis. *Chem.*

Rev. **118**, 10049-10293 (2018).

39. Ni, H., Chan, W.-L. & Lu, Y. Phosphine catalyzed asymmetric organic reactions.

Chem. Rev. **118**, 9344-9411 (2018).

40. Zhu, Z., Hu, Q., Fu, Y., Tong, Y. & Zhou, Z. Design and evolution of an enzyme for

the asymmetric Michael addition of cyclic ketones to nitroolefins by enamine catalysis.

Angew. Chem. Int. Ed. Engl. **63**, e202404312 (2024).

41. Cruz, P. D. J, Cassels, W. R., Chen, C.-H. & Johnson, J. S. Doubly stereoconvergent

crystallization enabled by asymmetric catalysis. *Science* **376**, 1224-1230 (2022).

42. Yamashita, Y., Ogasawara, Y., Banik, T. & Kobayashi, S. Photoinduced efficient

catalytic α-alkylation reactions of active methylene and methine compounds with

nonactivated alkenes. *J. Am. Chem. Soc.* **145**, 23160-23166 (2023).

43. Northrup, A. B. & MacMillan, D. W. C. The first direct and enantioselective cross-

aldol reaction of aldehydes. *J. Am. Chem. Soc.* **124**, 6798-6799 (2002).

44. Tian, J. & Zhou, L. Photoredox radical/polar crossover enables C-H gem-

difunctionalization of 1,3-benzodioxoles for the synthesis of monofluorocyclohexenes.

Chem. Sci. **14**, 6045-6051 (2023).

45. Bowry, V. W. & Inlgold, K. U. Kinetics of nitroxide radical trapping. 2. Structural

effects. *J. Am. Chem. Soc.***114**, 4992-4996 (1992).

Acknowledgements

This work was supported by the National Key R&D Program of China

(2022YFA0912000), the Center of Synthetic Biology and Integrated Bioengineering

(WU2022A006, WU2022A007, WU2023A009), the Research Center for Industries of

the Future (WU2022C032). We thank Instrumentation and Service Center for

Molecular Sciences (ISCMS) at Westlake University for the assistance in measurement

and data interpretation. We thank Dr. Yinjuan Chen for the assistance with high

resolution mass spectrometry (HRMS), Dr. Yuan Chen for the assistance with circular

dichroism (CD), and Danyu Gu for the assistance with electron paramagnetic resonance

(EPR).

Author contributions

J.Z. and Y.W. conceived and designed the overall project. J.Z. performed all synthetic

experiments and wet mechanism experiments. T.G., X.L., and M.-Z.M. created

mutations. Q.Z., and X.W. carried out the computational studies and the calculation of

absolute configuration (ECD) under the supervision of B.W. B.C. completed the

calculation of redox potentials and Gibbs reaction energy (ΔG) for SET of Int. A. M.-

J.M. performed the purification of some GluER variants. J.Z., B.W., and Y.W. wrote the

paper with input of all authors.

Competing interests

The authors declare no competing interests.