Generation of a Transgenic Mouse Model for Investigating Mitochondria in Sperm

Hironmoy Sarkar[1, 2]*, Bhola Shankar Pradhan[1,3,4]*, Suryaprakash R Batta[1], Neerja Wadhwa[5] and Subeer S Majumdar[1,6,7]

1 Cellular Endocrinology Laboratory, National Institute of Immunology, Aruna Asaf Ali Marg, JNU complex, New Delhi 110067, India.

2 Cell Biology and Bacteriology Laboratory, Department of Microbiology, Raiganj University, Raiganj, West Bengal, India - 733734

3 Łukasiewicz Research Network—PORT Polish Center for Technology Development, 147

Stabłowicka Street, 54-066 Wrocław, Poland.

4 The International Institute of Molecular Mechanisms and Machines Polish Academy of Sciences, ul. Flisa 6, 02-247 Warsaw, Poland

5 Embryo Biotechnology Lab, National Institute of Immunology, New Delhi, India.

6 National Institute of Animal Biotechnology, Miyapur, Hyderabad-500049, Telengana, India

7 Gujarat Biotechnology University, Gandhinagar, GIFT City, Gandhinagar, 382355, Gujarat, India.

*Corresponding author:

Hironmoy Sarkar, Email : h.sarkar@raiganjuniversity.ac.in

Bhola Shankar Pradhan, Email : bholapnl@gmail.com

Abstract

Mitochondria play an important role in the development of sperm, yet the mechanisms regulating their function in sperm remain poorly understood. The development of a method to regulate the expression of a target gene in the mitochondria of the sperm is an essential step in this line of research. In this study, we attempted to generate a system to express the transgene in the mitochondria of the sperm. As a proof of concept, we generated the transgenic mice expressing green fluorescent protein (GFP) fused with a mitochondrial localization signal (MLS) driven by PGK2 promoter which expresses the transgene in the sperm. Although the PGK2 promoter has been previously demonstrated to drive gene expression in spermatocytes and spermatids, the novelty of our method is the combination of the PGK2-driven MLS-GFP expression to study the mitochondria in vivo. We established two founder lines of transgenic mice by pronuclear microinjection, and MLS-GFP expression was confirmed in the mitochondria of the sperm cells by fluorescence microscopy and flow cytometry. Therefore, we provide a novel platform for studying mitochondrial function in sperm where the GFP can be replaced with other genes of interest to study their effect on mitochondria. These results may also help in understanding mitochondrial inheritance, and mitochondrial dysfunction in vivo.

Introduction

Mitochondria play a crucial role in the metabolism of spermatozoa, and they are implicated in energy production, cell signaling, as well as apoptotic pathways, all of which are necessary for the proper function of sperm (Aitken et al., 2001; Aitken et al., 2012; Amaral et al., 2013; Boguenet et al., 2021). The mitochondria in spermatozoa regulates important functions relating to human fertility and infertility (Spinelli et al., 2018; Fu et al., 2021; Tesarik et al., 2023). The genetic modification of mitochondria and its role in sperm development and in spermatogenesis has gained significant attention in recent years. The understanding of the function of mitochondria in sperm cells may provide critical insights into the molecular basis of male infertility (Park et al., 2021; Zini et al., 1993; Freitas et al., 2017). Recent reports suggested that male infertility is associated with a loss of mitochondrial proteins in spermatozoa, which induces low sperm motility, reduces OXPHOS activity, and results in male infertility (Park et al., 2021).

56   The dynamics of mitochondria in sperm, including their distribution, fusion, and fission, are important for sperm function and viability. During the development of sperm, mitochondria undergo significant structural and functional changes. The role of mitochondrial fusion and fission in sperm motility is increasingly understood, as proper mitochondrial dynamics are necessary for the motility and structural integrity of the sperm tail (Green et al., 2022; Raad M et al., 2024). Therefore, any defect in these processes leads to sperm dysfunction and infertility. However, studying the mitochondria in sperm has been challenging due to the difficulty of tracking these dynamics in live sperm cells.

64   The PGK2 promoter has been used as a testis-specific promoter for studying spermatogenesis (Zhang et al., 1999). The PGK2 promoter drives expression of transgene in spermatocytes and spermatids in transgenic models to study gene function during sperm development (Kramer et al., 1981; Danshina et al., 2010; McCarrey et al., 1992; Yoshioka et al., 2007). Green fluorescent protein (GFP) is often used as a marker to visualize the expression and localization of specific proteins in live cells. Moreover, the combination of GFP with a mitochondrial localization signal (MLS) allows for study of mitochondria within sperm (Otera et al., 2013; Wakai et al., 2014; Sandoval et al., 2014; Bader et al., 2020). This strategy will help to visualize the mitochondria of the sperm in vivo.

74   Although PGK2 has been used in a previous study related to spermatogenesis, it isnot used for studying mitochondria of the sperm. In this study, we used the PGK2 promoter to regulate the expression of GFP-tagged to a mitochondrial localization signal (PGK2-MLS-GFP) in transgenic mice.

78   **Material and methods**

79   **Animals.** Mice (FVB/J), were obtained from the small animal facility of National Institute of Immunology, were used for the present study. All animals were kept at 24 ± 2°C under 14 h light and 10 h dark cycle and used as per the National Guidelines provided by the Committee for the Purpose of Control and Supervision of the Experiments on Animals (CPCSEA). Protocols for the experiments were approved by the Institutional Animal Ethics Committee. Animals were kept in a hygienic air condition with suitable humidity and handled by trained personnel. The animals were killed individually in a separate room by

86   carbon dioxide inhalation in an in-house built carbon-dioxide device and by cervical

87   dislocation (Russell and Burch;1959).

Transgene constructs.

A total of three constructs were used in this study, and the schematic diagram were

provided in **Fig 1A**. The EGFPN$_2$ vector was used, and in this vector, mitochondrial

localization signal of *COXVIII* gene

(TCCGTCCTGACGCCGCTGCTGCTGCGGGGCTTGACAGGCTCGGCCCGGCGGCTC

CCAGTGCCGCGCGCCAAGA) (MLS) was added just before the coding sequence of

EGFP transgene. The CMV promoter from this vector was later replaced by PGK2

promoter of mice (1kb upstream region from the start codon of PGK2). The PGK2

promoter, a testis-specific promoter, was used to drive the expression of a green

fluorescent protein (GFP) tagged to a mitochondrial localization signal (MLS). Cloning was

performed using the standard method as described previously (Pradhan et al., 2016). The

sequences of the construct were provided in the Supplementary data.

Validation of the mitochondrial localization signal

HEK293 cells were transfected with the CMV-GFP-MLS plasmid using Lipofectamine

2000 (Invitrogen) following the manufacturer's recommendation. After the transfection,

cells were incubated for 24 hours and then analyzed for GFP expression and

mitochondrial localization as previously described (Pradhan et al, 2016). The

mitochondrial-specific dye, MitoTracker Red, was used to stain the mitochondria, and the

colocalization was evaluated for the GFP signal to determine the mitochondrial targeting

of the MLS.

Pronuclear microinjection of transgene cassette and oviductal embryo transfer.

This method was performed as described previously by us (Sarkar et al, 2020). Transgenic

mice were generated via pronuclear microinjection (Gordon et al., 1980; Costantini et al.,

1981; Auerbach et al., 2004). For pronuclear microinjection, female mice were super-

ovulated using PMSG and hCG followed by cohabitation with male mice. Oviducts were

collected from donor mice (euthanized by cervical dislocation) and placed into Brinster's

Modified Oocyte Culture Media (BMOC) containing hyaluronidase (1 mg/mL). Embryos

were collected and transferred to the 60 mm dish contain 100 µL drops of BMOC overlaid
with mineral oil. Linearized transgene cassette (PGK2-MLS-EGFP) was prepared at a
final concentration of 4ng/µl for microinjection. DNA was microinjected into the male
pronucleus of fertilized eggs using Narishighe micromanipulator. Each manipulated
embryo was then transferred into the pre-incubated BMOC containing dish and
maintained at 37°C in a 5% CO_2 incubator. Embryos were incubated till 2-cell stage.
Pseudo-pregnant female mice were generated by mating with vasectomized male mice.
2-cell stage embryos were transferred into the ampulla of the oviduct of pseudo-pregnant
recipient mice. Approximately 20 microinjected embryos were transferred in the oviductal
ampulla. After the gestation period (21 ± 2 days), recipient pups were born. Pups born
were analyzed for transgene integration by PCR.

Establishment of Transgenic Lines

127 Two founder mice expressing PGK2-GFP-MLS were screened through PCR (Pradhan et
128 al., 2016; Usmani et al., 2013). The founder mice were mated with wild-type females to
129 establish the separate transgenic lines, and the transgene integration in the progeny was
130 evaluated by PCR.

Fluorescence Microscopy

132 Fluorescence microscopy was employed to confirm GFP expression in the mitochondria
133 of sperm cells. Sperm from transgenic mice were isolated by standard methods and
134 immediately imaged using a confocal microscope (Pradhan et al., 2020).

Flow Cytometry Analysis

136 The sperm were isolated from the cauda epididymis from the wild-type mice and
137 transgenic mice. Samples were analyzed by a flow cytometer (FACS caliber, BD
138 Biosciences). Data were collected and analyzed using FlowJo software (TreeStar, Inc.) to
139 assess the percentage of GFP-positive sperm (Pradhan et al., 2016).

Results and Discussion

Validation of MLS Functionality in HEK Cells

The MLS sequences for mitochondrial targeting were used previously (Wakai et al., 2014;

Bader et al., 2020*). The mitochondrial targeting ability of the MLS of *COXVIII* gene was

validated by expressing the CMV-MLS-GFP construct in HEK 293 cells. We observed a

colocalization of GFP with mitochondrial markers suggesting that the MLS was functional

in the mammalian system (**Fig 1B**). As a control, we used the CMV-GFP construct without

MLS.

Generation of Transgenic Mice

To determine the in vivo implications of MLS sequence, we generated two transgenic

mouse lines using pronuclear microinjection. For this we used the PGK2-MLS-GFP

construct. The tissue-specific activity of the PGK2 promoter was previously demonstrated

to drive post-meiotic gene expression in the male germline (Zhang et al., 1999). We are

providing a diagram of the generation of the transgenic line (**Fig 2**). The transgene

integration was confirmed by PCR (**Fig 3**).

Localization of GFP in the mitochondria of sperm

We performed the fluorescence microscopy to determine the GFP fluorescence in the

sperm. We observed that GFP signal was localized to the midpiece of sperm, where

mitochondria are concentrated (**Fig 4**). Previous reports suggested that GFP expression

in the mitochondria of the sperm was nontoxic (Wang *et al., 2023*). Our study was in

agreement with the previous findings demonstrating the functional importance of

mitochondrial localization sequences in directing proteins to mitochondria (Wakai et al.,

2014; Bader et al., 2020). Mitochondria in the midpieces of the sperm are critical for

energy production and the motility and previous studies suggest a link between

mitochondrial dysfunction and male infertility (Park et al., 2021).

Analysis of GFP Expression by Flow Cytometry (FACS)

We performed the FACS analysis of sperm samples to validate the microscopy data. We

observed that GFP-positive sperm in various transgenic mice, suggesting that the

efficiency of the PGK2 promoter in directing mitochondrial GFP expression in vivo (**Fig 5**).

Our data also validated the microscopy data suggesting that the method is reproducible.

This study may help in studies of mitochondrial health and dynamics in large populations

of sperm cells (Park et al., 2021). Our study provides a platform to investigate
mitochondrial biology during spermatogenesis. The specificity of the PGK2 promoter
ensures minimal ectopic expression, which is critical for accurate functional analyses
(Zhang et al., 1999). Such advancements address limitations of ubiquitous promoters in
transgenic models. This work is a significant advancement for studying mitochondrial
dynamics in sperm cells. Mitochondria are vital for sperm function, playing key roles in
ATP production and motility, and defects in mitochondrial function are a major contributor
to male infertility (Tesarik et al., 2023).

The integration of a tissue-specific promoter with an MLS sequence enables precise
visualization of mitochondrial architecture and bioenergetics in vivo. This approach
addresses a critical gap in male reproductive biology, providing insights into mitochondrial
dynamics in sperm development, motility, and fertilization (Matilainen et al., 2017; Tesarik
et al., 2023). This approach can be used to study the effect of various mutations caused
by environmental stress on mitochondrial function, mitochondrial dynamics using
advanced imaging techniques during spermatogenesis and the role of mitochondria in
assisted reproduction technologies. These findings may have broad implications for
mitochondrial research, particularly in male infertility and sperm development.

Data availability

All the data were provided in the manuscript, and the reagents used in the manuscript
will be made available upon considerable request.

References

Aitken RJ, Koppers AJ. Apoptosis and DNA damage in human spermatozoa. Asian J
Androl. 2011 Jan;13(1):36-42. doi: 10.1038/aja.2010.68. Epub 2010 Aug 30. PMID:
20802502; PMCID: PMC3739394.

Aitken RJ, Whiting S, De Iuliis GN, McClymont S, Mitchell LA, Baker MA. Electrophilic
aldehydes generated by sperm metabolism activate mitochondrial reactive oxygen
species generation and apoptosis by targeting succinate dehydrogenase. J Biol Chem.
2012 Sep 21;287(39):33048-60. doi: 10.1074/jbc.M112.366690. Epub 2012 Jul 31. PMID:
22851170; PMCID: PMC3463304.

Amaral A, Lourenço B, Marques M, Ramalho-Santos J. Mitochondria functionality and
sperm quality. Reproduction. 2013 Oct 1;146(5):R163-74. doi: 10.1530/REP-13-0178.
PMID: 23901129.

Auerbach AB. Production of functional transgenic mice by DNA pronuclear microinjection.
Acta Biochim Pol. 2004;51(1):9-31. PMID: 15094822.

Bader G, Enkler L, Araiso Y, et al. Assigning mitochondrial localization of dual localized
proteins using a yeast Bi-Genomic Mitochondrial-Split-GFP. Elife. 2020 Jul 13;9:e56649.
doi: 10.7554/eLife.56649. PMID: 32657755; PMCID: PMC7358010.

Boguenet M, Bouet PE, Spiers A, Reynier P, May-Panloup P. Mitochondria: their role in
spermatozoa and in male infertility. Hum Reprod Update. 2021 Jun 22;27(4):697-719. doi:
10.1093/humupd/dmab001. PMID: 33555313.

Costantini F, Lacy E. Introduction of a rabbit beta-globin gene into the mouse germ line.
Nature. 1981 Nov 5;294(5836):92-4. doi: 10.1038/294092a0. PMID: 6945481.

Danshina PV, Geyer CB, Dai Q, et al. Phosphoglycerate kinase 2 (PGK2) is essential for
sperm function and male fertility in mice. Biol Reprod. 2010 Jan;82(1):136-45. doi:
10.1095/biolreprod.109.079699. Epub 2009 Sep 16. PMID: 19759366; PMCID:
PMC2802118.

Green A, Hossain T, Eckmann DM. Mitochondrial dynamics involves molecular and
mechanical events in motility, fusion and fission. Front Cell Dev Biol. 2022 Oct
19;10:1010232. doi: 10.3389/fcell.2022.1010232. PMID: 36340034; PMCID:
PMC9626967.

Gordon JW, Scangos GA, Plotkin DJ, Barbosa JA, Ruddle FH. Genetic transformation of
mouse embryos by microinjection of purified DNA. Proc Natl Acad Sci U S A. 1980
Dec;77(12):7380-4. doi: 10.1073/pnas.77.12.7380. PMID: 6261253; PMCID:
PMC350507.

Fu L, Luo YX, Liu Y, Liu H, Li HZ, Yu Y. Potential of Mitochondrial Genome Editing for
Human Fertility Health. Front Genet. 2021 Jul 20;12:673951. doi:
10.3389/fgene.2021.673951. PMID: 34354734; PMCID: PMC8329452.

Freitas MJ, Vijayaraghavan S, Fardilha M. Signaling mechanisms in mammalian sperm
motility. Biol Reprod. 2017 Jan 1;96(1):2-12. doi: 10.1095/biolreprod.116.144337. PMID:
28395326.

Kramer JM. Immunofluorescent localization of PGK-1 and PGK-2 isozymes within specific
cells of the mouse testis. Dev Biol. 1981 Oct 15;87(1):30-6. doi: 10.1016/0012-
1606(81)90057-9. PMID: 7026322.

Matilainen O, Quirós PM, Auwerx J. Mitochondria and Epigenetics - Crosstalk in
Homeostasis and Stress. Trends Cell Biol. 2017 Jun;27(6):453-463. doi:
10.1016/j.tcb.2017.02.004. Epub 2017 Mar 6. PMID: 28274652.

McCarrey JR, Berg WM, Paragioudakis SJ, Zhang PL, Dilworth DD, Arnold BL, Rossi JJ.
Differential transcription of Pgk genes during spermatogenesis in the mouse. Dev Biol.
1992 Nov;154(1):160-8. doi: 10.1016/0012-1606(92)90056-m. PMID: 1426623.

Otera H, Ishihara N, Mihara K. New insights into the function and regulation of
mitochondrial fission. Biochim Biophys Acta. 2013 May;1833(5):1256-68. doi:
10.1016/j.bbamcr.2013.02.002. Epub 2013 Feb 20. PMID: 23434681.

Park YJ, Pang MG. Mitochondrial Functionality in Male Fertility: From Spermatogenesis
to Fertilization. Antioxidants (Basel). 2021 Jan 12;10(1):98. doi: 10.3390/antiox10010098.
PMID: 33445610; PMCID: PMC7826524.

Pradhan BS, Majumdar SS. An Efficient Method for Generation of Transgenic Rats
Avoiding Embryo Manipulation. Mol Ther Nucleic Acids. 2016 Mar 8;5(3):e293. doi:
10.1038/mtna.2016.9. PMID: 27111419; PMCID: PMC5014465.

Pradhan BS, Bhattacharya I, Sarkar R, Majumdar SS. Pubertal down-regulation of
Tetraspanin 8 in testicular Sertoli cells is crucial for male fertility. Mol Hum Reprod. 2020
Oct 1;26(10):760-772. doi: 10.1093/molehr/gaaa055. PMID: 32687199.

Russell, WMS and Burch, RL (1959) The principles of humane experimental technique.
Methuen: London.

Sandoval H, Yao CK, Chen K, Jaiswal M, Donti T, Lin YQ, Bayat V, Xiong B, Zhang K,
David G, Charng WL, Yamamoto S, Duraine L, Graham BH, Bellen HJ. Mitochondrial

fusion but not fission regulates larval growth and synaptic development through steroid
hormone production. Elife. 2014 Oct 14;3:e03558. doi: 10.7554/eLife.03558. PMID:
25313867; PMCID: PMC4215535.

Sarkar RK, Sen Sharma S, Mandal K, et al. Homeobox transcription factor Meis1 is crucial
to Sertoli cell mediated regulation of male fertility. Andrology. 2021 Mar;9(2):689-699. doi:
10.1111/andr.12941. Epub 2020 Nov 23. PMID: 33145986.

Spinelli JB, Haigis MC. The multifaceted contributions of mitochondria to cellular
metabolism. Nat Cell Biol. 2018 Jul;20(7):745-754. doi: 10.1038/s41556-018-0124-1.
Epub 2018 Jun 27. PMID: 29950572; PMCID: PMC6541229.

Tesarik J, Mendoza-Tesarik R. Mitochondria in Human Fertility and Infertility. Int J Mol Sci.
2023 May 18;24(10):8950. doi: 10.3390/ijms24108950. PMID: 37240296; PMCID:
PMC10218931.

Usmani A, Ganguli N, Sarkar H, Dhup S, Batta SR, Vimal M, Ganguli N, Basu S,
Nagarajan P, Majumdar SS. A non-surgical approach for male germ cell mediated gene
transmission through transgenesis. Sci Rep. 2013 Dec 5;3:3430. doi: 10.1038/srep03430.
PMID: 24305437; PMCID: PMC3852150.

Wang S, Duan Y, Chen B, et al. Generation of Transgenic Sperm Expressing GFP by
Lentivirus Transduction of Spermatogonial Stem Cells In Vivo in Cynomolgus Monkeys.
Vet Sci. 2023 Feb 1;10(2):104. doi: 10.3390/vetsci10020104. PMID: 36851408; PMCID:
PMC9966439.

Wakai T, Harada Y, Miyado K, Kono T. Mitochondrial dynamics controlled by mitofusins
define organelle positioning and movement during mouse oocyte maturation. Mol Hum
Reprod. 2014 Nov;20(11):1090-100. doi: 10.1093/molehr/gau064. Epub 2014 Aug 11.
PMID: 25113836.

Yoshioka H, Geyer CB, Hornecker JL, Patel KT, McCarrey JR. In vivo analysis of
developmentally and evolutionarily dynamic protein-DNA interactions regulating
transcription of the Pgk2 gene during mammalian spermatogenesis. Mol Cell Biol. 2007
Nov;27(22):7871-85. doi: 10.1128/MCB.00990-07. Epub 2007 Sep 17. PMID: 17875925;
PMCID: PMC2169153.

Zhang LP, Stroud J, Eddy CA, et al. Multiple elements influence transcriptional regulation
from the human testis-specific PGK2 promoter in transgenic mice. Biol Reprod. 1999
Jun;60(6):1329-37. doi: 10.1095/biolreprod60.6.1329. PMID: 10330089.

Zini A, de Lamirande E, Gagnon C. Reactive oxygen species in semen of infertile patients:
levels of superoxide dismutase- and catalase-like activities in seminal plasma and
spermatozoa. Int J Androl. 1993 Jun;16(3):183-8. doi: 10.1111/j.1365-
2605.1993.tb01177.x. PMID: 8359932.

Conflicts of Interest: The authors declare no conflict of interest.

Acknowledgments

We are thankful to all the staff of the Small Animal Facility. Thanks are due to Ram Singh,
Dharamvir Singh and Birendar Roy for the technical assistance. We are grateful to the
Director, NII for valuable support. We are grateful to Department of Biotechnology, Govt.
of India, for providing the financial assistance under grants
BT/PR11313/AAQ/01/376/2008, BT/HRD/35/01/01/2010 (TATA Innovation Award) and
BT/PR10805/AAQ/1/576/2013. BSP was supported the Polish National Science Centre
grant (2020/39/D/NZ5/02004). H.S. was supported by the Department of Biotechnology,
Govt. of India grant (BT/PR46488/AAQ/1/886/2022). The authors declare that they have
no competing interests.

Author Contributions

304 Experiments were conceived and designed by B.S.P., H.S., and S.S.M. Experiments
305 were performed by H.S., S.B., N.W. and B.SP. Data of the manuscript was analyzed by
306 B.S.P, H.S. Manuscript was written and reviewed by B.S.P. and H.S. All authors have
307 read and agreed to the published version of the manuscript.

Figure Legend

Figure 1. Validation of MLS sequence in Hek293 cells. (A) A schematic diagram of the
constructs used in this study. (B) Hek 293 cells were transfected with CMV-GFP control
and CMV-MLS-GFP; and costained with mitotracker red dye. The images were analyzed
24h after transfection. Scale bar: 10µm.

Figure 2. A schematic diagram of generation of PGK2-MLS-GFP transgenic mice after

embryo transfer.

Figure 3. Generation of PGK2-MLS-GFP transgenic mice. Detection of transgene by

PCR. PCR screening of progeny generated through pronuclear microinjection. A band

size of 258 bp represented the transgene positive mice. b = blank containing only PCR

mastermix, numerical indicated the progeny numbers in the gel. Ppia was used as a

loading control.

Figure 4. Detection of EGFP-positive sperm in the epididymis of the transgenic mice by

microscopy. The EGFP positive sperm was detected by microscopy in the live sperm of

the wild-type mice (A) and transgenic mice (B). Scale bar: 20µm.

Figure 5. Detection of EGFP-positive sperm in epididymis of the transgenic mice by flow

cytometry. (A) Scatter plot of one of the representative sperm samples and the

corresponding histogram of wild-type mice. (B) Scatter plot of one of the representative

sperm samples and the corresponding histogram of transgenic mice.

Figure 1

Figure 2

PCR positive

Figure 3

Figure 4

Control

PGK2-MLS-GFP mice

Figure 5
Control

PGK2-MLS-GFP mice