Hearing in Two Closely Related *Peromyscus* Species (*Peromyscus maniculatus* and *P.*
***leucopus*)**

Luberson Joseph[1], Emily Margaret New[1], Desi Marie Joseph[1], Tamara Chenell Woodley[1],

Vanessa Yamileth Franco[1], Ben-Zheng Li[2], Guinevere OU Wogan[1], and Elizabeth A.

McCullagh[1*]

[1]Oklahoma State University (OSU), Department of Integrative Biology, College of Arts and

Sciences, Stillwater, Oklahoma 74078, USA

[2]University of Colorado Anschutz Medical Campus, Department of Physiology and Biophysics,

Aurora, Colorado 80045, USA

*Corresponding author: elizabeth.mccullagh@okstate.edu

501 Life Sciences West, Stillwater, OK 74074 USA

ABSTRACT

The genus *Peromyscus* has been extensively used as a model for ecological, behavioral, and

evolutionary investigations. We used auditory brainstem responses (ABRs), craniofacial

morphology, and pinna measurements to compare characteristics that impact hearing in two

wild-caught species, *P. leucopus P. maniculatus*. We observed significant statistical differences

in craniofacial and pinna attributes between species with *P. leucopus* overall exhibiting larger

features than *P. maniculatus*. ABR recordings indicated that both species showed similar best

frequency thresholds between 8-24 kHz. We found significant effects of intensity on amplitude

ratio of wave I and IV for *P. maniculatus*, but not *P. leucopus* and effects of wave number on

slope of the latency-intensity function with higher wave IV and shorter wave I slope of latency

intensity function in *P. leucopus*. Finally, the data showed significant differences in latency shift

of the DN1 component of the BIC in relation to ITD between species, while no significant

differences were observed across relative DN1 amplitude. This study supports the used of *P.*

leucopus and *P. maniculatus* as future model species for auditory research.

Keywords: binaural hearing, *Peromyscus*, auditory brainstem recordings, hearing, pinna,

interaural timing difference (ITD)

INTRODUCTION

Hearing and sound localization are critical for the survival and fitness of all taxa. In small

mammals, sound localization facilitates predator avoidance, capturing prey, finding mates,

foraging, and conspecific communication (Colburn et al., 1987; Grothe et al., 2010; Kidd et al.,

1995). To perceive sound source location, mammals rely on interaural time differences (ITD)

(for low frequency sound in the horizontal plane) and interaural level differences (ILD) (for high

frequency sound) between the two pinnae. ITD and ILD cues are influenced by the size of the

head and the shape of the pinna (Blauert, 1997; Grothe et al., 2010). The auditory brainstem

consists of specialized regions that integrate ITD and ILD information from each ear. Despite

decades of research on hearing and sound localization in small mammals (Blauert, 1997; Grothe

et al., 2010; Heffner, 2001), our understanding of species-specific biological variation in sound

localization and their hearing ranges continues to need to be explored (Capshaw et al., 2023).

To understand the mechanism of hearing, animal models, including laboratory and wild

rodents, can serve as valuable tools. Most studies have used the laboratory house mouse (*Mus*

musculus) as a model in hearing research due to its sensitive hearing, ease to breed and

maintenance in laboratory settings, and genetic manipulability (Capshaw et al., 2022; Ehret and

Dreyer, 1984). Yet, the limited genomic diversity in inbred laboratory rodents pose challenges to

fully recapitulating the broad spectrum of human disorder phenotypes (Voelkl et al., 2020). The

house mouse has faced criticism as a model for auditory research, owing to its poor sensitivity to

low frequency sounds, increased vulnerability to noise, and minimal audiometric variation within

strains (Capshaw et al., 2022). Moreover, the house mouse has a short lifespan and may not

exhibit aging patterns similar to other mammals with long lifespan such as bats, African mole-

rats or humans (Dammann, 2017). In hearing studies, there has been relatively limited used of

wild rodent models to understand the mechanism of hearing with aging. However, the auditory

field has leveraged many alternative species including the Mongolian gerbil (*Meriones*

unguiculatus), which is valued for its similarity of hearing range to human (Heeringa et al., 2020;

Jüchter et al., 2022; Mills et al., 1990). Although these model taxa have shed insights in

understanding the fundamental mechanism of hearing, it is essential to continue to consider

comparative approaches that include a wider range of species, particularly those that reflect

natural diversity in their vocalization, have long life spans, and vary in habitat use (Capshaw *et*

al 2023).

The white-footed mouse (*Peromyscus leucopus*) and the North American deer mouse

(*Peromyscus maniculatus*) are two of the most abundant rodents in North America (Kirkland and

Layne, 1989). They both belong to the family Cricetidae and are more closely related to hamsters

and voles than to mice of the family Muridae. Both species have been extensively studied as

model systems in ecological, behavioral, biogeographical, and evolutionary investigations with

regard to their physiological adaptation to varying habitat types (arboreal habitats, grassland,

woodlands, brushlands, swamps, and desert), their social system (mainly promiscuous), and

behaviors (maternal, winter nesting, climbing, and agonistic behaviors) (Bedford and Hoekstra,

2015; Harney and Dueser, 1987; Lewarch and Hoekstra, 2018). Both species are also of human

health concern with regards to their carrying viruses and pathogens including hantavirus,

leptospirosis, and plague (Childs et al., 1994; Larson et al., 2018).

Recently, members of the genus *Peromyscus* have emerged as valuable model systems in

the field of neuroscience for studying age-related hearing loss due to their considerable lifespan

compared to mammals of similar size. *Peromyscus* species exhibit an average lifespan nearly

double that of *M. musculus,* when reared in laboratory settings, with the potential to live up to

eight years (Burger and Gochfeld, 1992; Guo et al., 1993). In addition, *Peromyscus* species hear

best between 8 to 16 kHz, as demonstrated by low ABR thresholds, with the ability to hear up to

65 kHz (Capshaw et al., 2022; Dice and Barto, 1952; Ralls, 1967). In comparison to *M.*

musculus, *Peromyscus* rodents display lower production of reactive oxygen species and

enhanced resistance to oxidative stress, resulting in delayed accumulation of oxidative damage to

deoxyribonucleic acid over the *Peromyscus* lifespan (Csiszar et al., 2007; Labinskyy et al., 2009;

Shi et al., 2013), among other preventive effects which may slow cochlear aging (Ohlemiller and

Frisina, 2008). *Peromyscus* species occupy a large range of habitats, which offer unique

opportunities to identify alleles underlying natural variation in biomedically relevant behaviors

(Dewey and Dawson, 2001; Vrana et al., 2014). Moreover, both species are phylogenetically

closely related (Bradley et al., 2007; Fiset et al., 2015; King, 1968), occur in sympatry, and share

diverse morphological similarities such as tail length and pelage color (Millien et al., 2017; Platt

et al., 2015) . However, the two species differ in their craniofacial and pinna sizes (Light et al.,

2021) which may contribute to differences in their hearing as we know that pinna size impact

hearing by enhancing sound collection and amplification, improving frequency discrimination,

and facilitating more accurate sound localization (Heffner et al., 2020; Heffner and Heffner,

1982). Therefore, members of the genus *Peromyscus* show promise as models that can be used to

complement auditory research across species and consequently can be reference taxa to explore

small mammals' hearing across longer lifespans.

The purpose of this investigation is to compare hearing-related anatomy, hearing range,

and binaural hearing of *P. leucopus* and *P. maniculatus* measured by craniofacial features, pinna

size, and auditory brainstem responses (ABRs). We expect that *P. maniculatus* will have shorter

binaural latency compared to *P. leucopus*, due to smaller overall size, pinna, and craniofacial

measurements. We also expect that there will be differences in other measures of hearing

including best frequency hearing thresholds and monaural ABRs (monaural amplitudes and

monaural latencies) due to differences in body size, habitat, and other variability between these

two closely related species.

MATERIALS AND METHODS

All procedures used for all experiments complied with the guidelines of the American

Society of Mammologists (Sikes et al., 2011), were approved by the Oklahoma State University

Institutional Animal Care and Use Committee (IACUC), and permission from an Oklahoma

Department of Wildlife Conservation scientific collecting permit.

Animals

113
Experiments were conducted on 26 individuals, including 15 wild *P. leucopus* (9 males, 6

females) and 11 wild *P. maniculatus* (9 males, 2 females). Animals were live trapped using

aluminum Sherman (H.B Sherman Traps, Inc. Tallahassee, FL) non-folding traps (3" x 3" x 10")

between June 2021 and July 2022 at three different locations across Oklahoma, USA

(Packsaddle Wildlife Management Area, James Collins Wildlife Management Area, and Payne

County) (Figure 1). The traps were baited with old fashioned oats and creamy peanut butter, left

overnight, and collected the next morning (~12 hours). Upon capture, animals were aged,

morphologically identified to species in the wild according to (Caire, 1989) and confirmed with

DNA barcoding polymerase chain reaction (PCR) of collected tail snips. Animals ages were

calculated for each species based on body mass and were divided into three age groups (Juvenile,

subadult, and adult) (see table 1). Animals were then transported to the laboratory for ABRs.

Figure 1: Map showing trapping site locations in Oklahoma. Packsaddle wildlife management area
(WMA) sites are represented by pink triangles, James Collin wildlife management area (WMA) sites are
represented by blue squares, and Payne County sites are represented by green circles.
130
131
132
DNA extraction, amplification, and sequencing

Deoxyribonucleic acid (DNA) was extracted from tail tissue samples by proteinase K

digestion using a Qiagen DNeasy blood and tissue kit (Hilden, Germany) and the protocol

outlined by (Nicolas et al., 2012). The DNA concentration and purity were first determined by

using a Thermo Scientific Nano-Drop Lite-Spectrophotometer (Fisher Scientific,

Spectrophotometer, Nanodrop Lite 6V 18W, Wilmington, DE). The CO1 gene was amplified

using the primer sequences (CCTACTCRGCCATTTTACCTATG) and

(ACTTCTGGGTGCCAAAGAATCA) (Ducroz et al., 2001; Robins et al., 2007). DNA samples

were assayed in a 50 µl reaction with 25 µl Phusion Master mix, 2.5 µl forward primer, and 2.5

µl reverse primer and mQ water. The PCR comprised of 35 cycles: 95º C for 300 seconds; 30

seconds at 94º C, 40 seconds at 55º C, 90 seconds at 72 º C, and a final 300 second extension at

72 º C. The double-stranded PCR products were purified and sequenced at the Center for

Genomics and Proteomics of Oklahoma State University (Stillwater, Oklahoma, USA). All

sequences were compared with other COI sequences using the NCBI GenBank (Sayers et al.,

2022, 2021) databases to confirm species identification (supplemental Figure 1, supplemental

Table 1).

Morphological measures

Craniofacial morphology features including pinna size, tail length, body length, and body

mass were recorded for each animal using a six-Inch Stainless Steel Electronic Vernier Caliper

(DIGI-Science Accumatic digital Caliper Gyros Precision Tools Monsey, New York, USA) and

a digital stainless Steel Electronic scale (Weighmax W-2809 90 LB X 0.1 OZ Durable Stainless

Steel Digital Postal scale, Chino, California, USA). Measurements of animal head and pinna

including inter-pinna distance (mm) (measurement between the two ear canals), nose to pinna

distance (mm) (measurement from the tip of the nose to the middle of the pinna), pinna length

(mm) (basal notch to tip, excluding hairs), and pinna width (mm) were measured (Figure 3A).

Pinna measurements (pinna width and pinna length) were used to calculate the effective pinna

diameter which is the square root of the pinna length multiplied by the pinna width (Anbuhl et

al., 2017). Tail length (sacrum to caudal tip, excluding hairs), body length (tip of nose to caudal

tip), and weight to the nearest gram were taken for each animal. To assess dependence of

morphological traits on body size, log values of traits (pinna width, length etc.) were plotted

against the log body length (supplemental Figure 2).

Auditory Brainstem Response (ABR) recordings

We recorded ABRs from wild *P. leucopus* and *P. maniculatus* using similar procedures

as previous publications(Chawla and McCullagh, 2022; McCullagh et al., 2020; New et al.,

2024). Rodents were sedated with an intraperitoneal injection of 60 mg/kg ketamine and 10

mg/kg xylazine for initial anesthesia followed by maintenance dosage of 25 mg/kg ketamine and

12 mg/kg xylazine. After being fully sedated, as indicated by lack of toe pinch reflex, the

animals were transported to a small sound-attenuating chamber (Noise Barriers, Lake Forest, IL,

USA), and positioned on a water pump heating pad to maintain a body temperature of 37^o C.

Subdermal needle electrodes were inserted under the skin at midline between the ears over the

brainstem (apex, active electrode), directly behind the apex on the nape (reference), and in the

back leg of the sedated animals (ground electrode) for differential recordings. To obtain and

amplify evoked potentials from electrodes positioned below the skin of the animal, we used a

Tucker-Davis Technologies (TDT, Alachua, FL, USA) RA4LI head stage, a RA16PA

preamplifier, and a Multi I/O processor RZ5 attached to a PC with custom Python software to

record the data. Data were processed using a second order 50-3000 Hz filter and averaged across

10-12 ms of recording time over 500-1000 repetitions. Acoustic stimuli for frequencies of 32 –

64 kHz were presented to animals using TDT Electrostatic Speakers (TDT EC-1) or TDT

Electrostatic Speaker-Coupler Model (TDT MF-1) for frequencies of 1 – 24 kHz and broadband

clicks attached through custom ear bars with Etymotic ER-7C probe microphones (Etymotic

Research Inc. Elk Grove, IL) for calibration of sounds in the ears. Ear bars had the ER-7C probe

microphones threaded through the tapered end of the tube with the speakers connected through

the provided tubing to the front of the tube creating a closed-field presentation of stimuli. Tone

stimuli were 4 ms in total duration with a 1 ms on-ramp and 1 ms off-ramp (2 ms ± 1 ms on/off

ramps). Click stimuli were 0.1 ms in duration with alternating polarity. Acoustic stimuli were

presented to the animal with a 30 ms interstimulus interval with a standard deviation of 5 ms

(Laumen et al., 2016a). Randomizing interstimulus interval between each presentation has been

shown to optimize the ABR waveform (Wang et al., 2020). Stimuli were produced at a sampling

rate of 97656.25 Hz through a TDT RP2.1 real-time processor controlled by a custom Python

program.

ABR response threshold

ABR response thresholds were determined by using the visual technique outlined by

(Brittan-Powell and Dooling, 2004). In short, threshold was defined to be between the intensity

at which the waveforms were no longer present and the previous intensity at which they were

visible in 5- and 10-dB increments (5 dB increments were used when near threshold). This

method was used for analyzing the audiogram for best hearing frequencies presented (1, 2, 4, 8,

16, 24, 32, 46, 64 kHz) and intensities (90 - 10 dB SPL) in addition to click threshold.

Monaural auditory brainstem response recordings

To generate monaural evoked potentials, broadband click stimuli were presented

independently to each ear of the sedated animal. Peak amplitude (voltage from peak to absolute

trough) and peak latency (time to peak amplitude) were measured across the four peaks of the

auditory brainstem recording waveforms (Figure 4A, 4B). To calculate the monaural latency and

amplitude for each species, we calculated the average of the monaural amplitude or latency of

waves I and IV from the ABRs data obtained for sound presentation in each ear across intensities

(60-90 dB SPL) (New et al., 2024; Zhou et al., 2006). We next calculated the slope of latency for

each individual to demonstrate the effects of click intensity on the peak latency of the ABR

waves I and IV following the methods of Zhou et al. (2006). In brief, the slope of each latency

intensity function was estimated by taking the change in peak latency and dividing it by the

intensity difference of each wave for each animal (Figure 2, Ba, b). The latency slope data was

used to make comparisons in monaural peak latency between species.

Figure 2: Auditory Brainstem response patterns of a female *P. leucopus* determined with clicks of
different intensities. Peak latency of monaural wave I and IV decrease with increasing click intensity
(dotted lines). B (a) represents latency intensity functions of wave I and B (b) shows latency intensity
functions of wave IV. The slope of the latency intensity function was calculated as the amount of change
in peak latency per decibel.

Binaural auditory brainstem response recordings

To produce the binaural ABR response, we simultaneously played broadband click

stimuli (same as above) at 90 dB SPL to both pinnae of the sedated animal. The binaural

interaction component (BIC) of the ABR was determined by subtracting the sum of the two

monaural auditory brainstem evoked responses from the binaural auditory brainstem response

recordings (Benichoux et al., 2018; Laumen et al., 2016a). Custom Python software was used to

measure the BIC amplitude and latency, with amplitude calculated to the baseline of the

recording (Chawla and McCullagh, 2022). BIC was defined as the negative peak wave (DN1) at

wave IV of the ABR after subtraction of the summed monaural and binaural responses. To

calculate how BIC varies with interaural time difference (ITD), both species were presented with

click stimuli that had shifting ITDs of -2.0 to 2.0 ms in 0.5 ms steps. We calculated the peak

latency and amplitude of DN1 for each ITD for each species. The ITD latency shift of the DN1

component of the BIC was determined in relation to the latency of DN1 at 0 ITD. The DN1

amplitude is highest at 0 ITD therefore amplitudes for ITD shifts were transformed to relative

amplitude with respect to 0 ms ITD to normalize recorded data (Laumen et al., 2016a). The

average latency shift and relative DN1 amplitude values were used to make comparison of

binaural auditory brainstem responses as function of ITD between species.

Statistical analyses

All analyses and figures were created in R Studio version 4.0.3 (R Core Team 2020),

using ggplot2 (Wickham, 2016) and lme4 (Bates et al., 2014) packages. Two-way analysis of

variances (ANOVAs) were used to statistically compare morphological characteristics between

species. Log-transformed morphological features (pinna width, length, etc.) were compared with

log body length and slope of the linear fit to describe potential allometry (slope > 1 indicating

positive allometry and < 1 indicating negative allometry). Linear mixed-effects models (LMMs)

were performed on multivariate data (hearing range and ABR amplitudes and latencies) with

species, frequency, percentage relative DN1 amplitude, and shifts in DN1 latency of BIC as

fixed effects and animal as a random effect. Estimated marginal means were used for pairwise

comparisons of frequency, relative amplitude and latency between both studied species (Russell,

2018). To estimate potential effects of body size or species on BIC relative DN1 amplitude and

latency of the DN1 component of the BIC relative to ITD at 0, LMMs were performed with fixed

effects of species, ITD, and body size and animal as a random effect compared to a null model

that excluded body size.

RESULTS

Based on the NCBI GenBank species identification systems for COI, of the tail samples

from the 26 animals collected, 15 individuals were identified as *P. leucopus*, and 11 as *P.*

maniculatus (supplemental table 1, Figure 1). *P. maniculatus* body length ranged from 55 to 78

millimeters with mean body mass of 12 grams, while *P. leucopus* body length ranged from 65 to

93 millimeters with mean body mass of 21 grams (Table 1).

Species	Age group	Body mass range (g)	Sample size (male, female)
P. maniculatus	Juvenile	< 14	8 (7, 1)
	Subadult	14 – 17	0
	Adult	> 17	3 (2, 1)
P. leucopus	Juvenile	< 13	1 (1,0)
	Subadult	13 – 18	4 (3, 1)
	Adult	> 18	10 (5, 5)

264
265   **Table 1:** Age was estimated based on body mass for each species based on published literature. Ages for
266   *P. maniculatus* was describe as follows: Juveniles < 14 grams, subadults, between 14-17 grams, and
267   adults, > 17 grams (Fairbairn, 1977). We inferred ages for *P. leucopus* as follow: Juveniles < 13 grams,
268   subadults, between 13 – 18 grams, and adults > 18 grams (Cummings and Vessey, 1994). We did not
269   make comparisons by ages due to limited sample sizes by age groups.
270

Morphological characteristics

Previous studies have shown that *P. leucopus* and *P. maniculatus* have significant

differences in pinna sizes and craniofacial features (Choate, 1973; Light et al., 2021; Millien et

al., 2017). We observed significant statistical differences for pinna attributes including pinna

length (Df = 1, 24; F = 11.79; p = 0.0021), and pinna width (Df = 1, 24; F = 8.47; p = 0.0076)

(Figure 3B and 3C respectively). In general, *P. leucopus* had longer and wider pinnae compared

to *P. maniculatus*, with mean pinna length and width estimated at 15.30 and 8.02 mm for *P.*

leucopus and those of *P. maniculatus* were 13.15 and 6.42 mm, respectively. Similarly, effective

pinna diameter (Df = 1, 24; F = 13.69; p = 0.0011) was significantly different between species,

with *P. leucopus* having a wider effective pinna diameter compared to *P. maniculatus* (Table 2,

Figure 3D). Craniofacial features including inter-pinna distance (Figure 2F; Df = 1, 24; F = 9.08;

p = 0.0060) and distance from the nose to the pinna (Figure 3E; Df = 1, 24; F = 5.82; p = 0.0239)

were significantly different between species with *P. leucopus* exhibiting a wider distance

between pinnae and a longer distance from the nose to the pinna. Like pinnae morphology and

craniofacial features, there were significant differences in body mass (Df = 1, 24; F = 24.2; p =

0.00005 , Figure 3I), tail length (Df = 1, 24; F = 25.76, p = 0.0245, Figure 2H), and body length

(Df = 1, 24; F = 18.32; p = 0.0002, Figure 3G) between both species, with *P. leucopus* weighing

significantly more including longer tails and longer body length than *P. maniculatus* (Table 2).

We tested if there were sex differences in craniofacial and pinna sizes in *P. leucopus*. There were

no significant differences in craniofacial and pinna sizes between male and female *P. leucopus*

(all p-value > 0.05). Sex differences were not explored for *P. maniculatus* due to limited number

of female subjects of this species (9 males, 2 females). When anatomical data were compared for

potential effects of body size (allometry), log features (pinna width, etc.) compared to log body

length did not show positive allometry except for tail length, which indicated positive allometry

(slope of 1.437 (*maniculatus*) and 1.218 (*leucopus*), supplemental Figure 1).

Figure 3: Morphological differences between *P. leucopus* and *P. maniculatus*. Pinnae, head, and body
features (**A**) were evaluated between species (pink boxplot = *leucopus*, blue boxplot = *maniculatus*).
Measurements JK show the inter pinnae distance, JN the nose to pinna distance, MK the pinna width, LM
the pinna height, OP the tail length, and PQ the body length. Effective pinna diameter was calculated by
taking the square root of pinna height multiplied by pinna width (MK/LM). Significant differences were
observed for all features including Pinna width (**B**), Pinna length (**C**), Effective diameter (**D**), Nose to
pinna distance (**E**), Inter pinna distance (**F**), Body length (**G**), Tail length (**H**), and Body mass (**I**).
Peromyscus pictured is a wild caught *P. leucopus* captured in Payne County, Stillwater, Oklahoma. Image
is presented only for demonstration of measurements.

307
308

Morphological characteristics	*Maniculatus* mean ± S. E	*Leucopus* mean ± S. E	degrees of freedom	F-statistic	p-value
Effective pinna diameter	9.16 ± 0.36	11.01 ± 0.34	1, 24	13.69	0.0011

Pinna length	13.15 ± 0.39	15.30 ± 0.45	1, 24	11.79	0.0021
Pinna width	6.42 ± 0.36	8.02 ± 0.39	1, 24	8.47	0.0076
Inter pinna distance	12.59 ± 0.47	14.72 ± 0.49	1, 24	9.08	0.0060
Nose to pinna distance	23.84 ± 0.96	26.82 ± 0.79	1, 24	5.82	0.0239
Body length	67.19 ± 2.28	80.87 ± 2.17	1, 24	18.32	0.0002
Tail length	48.01 ± 2.84	61.21 ± 4.21	1, 24	25.76	0.0245
Body mass	12 ± 1	22 ± 1	1, 24	24.2	0.00005

309
310 **Table 2:** Morphological characteristics features of *P. maniculatus* and *P. leucopus* of the Packsaddle
311 wildlife management area (WMA), James Collin wildlife management area (WMA) and Payne County.
312 Values presented represent the mean of different morphological features recorded ± standard error, the
313 degrees of freedom, F-statistic, and p-value of morphological differences between species.
314

Frequency thresholds between species

Both *P. maniculatus* and *P. leucopus* displayed the best sensitivity to tones between 8 to

24 kHz, as indicated by lower ABR thresholds (Figure 4C). We detected no significant statistical

difference in best frequency thresholds between species across the frequencies tested (LMM, p =

0.4692). Similarly, no significant difference in best frequency hearing threshold was observed

between male and female *P. leucopus* (F = 0.054, p = 0.82). We next investigated whether

craniofacial or pinna measurements features are correlated with or influence best frequency

thresholds in both species. We found that none of the morphological measurements had a

significant effect on best frequency hearing threshold between species (p-value > 0.05)

(supplemental materials, Figure 3).

Figure 4: Figure 4A and 4B show Auditory Brainstem response patterns of a female *P. leucopus* and a female *P. maniculatus* determined with clicks of different intensities, respectively. Hearing range was measured across frequency (1-64 kHz) for both *P. leucopus* and *P. maniculatus* (Figure 4C). No significant main effects of frequency between species were found. Unfilled blue circles represent *P. maniculatus* while filled pink squares represent *P. leucopus*.

ABR waveform amplitudes

We measured the responses of *P. leucopus* and *P. maniculatus* to monaural transient

click stimuli across intensities (60 -90 dB SPL) to ensure responses above threshold. We

observed that the amplitude of waves I and IV (Figure 5A and 5B, wave I filled circles and wave

IV unfilled circles), increased monotonically with increasing intensity (60 to 90 dB SPL). At 90

dB SPL, the average amplitude of wave I and wave IV were 2.75 and 1.79 µV for *P. leucopus*.

For *P. maniculatus*, the average amplitude of wave I and wave IV were 3.27 and 1.83 µV at 90

dB SPL. *P. leucopus* monaural wave I and IV amplitude were statistically different across

intensities (LMM: p-value < 0.0001). There were no significant statistical differences in

monaural wave I and IV amplitude between male and female *P. leucopus* across intensities

(LMM: p-value = 0.677). Similarly, we detected significant main effects of intensity on

monaural wave I and IV amplitude of *P. maniculatus* (LMM: p-value = 0.002). For both species,

wave IV generally had smaller amplitude than wave I (Figure 5A and 5B: unfilled vs filled

circles) and as a result, the wave IV/I amplitude ratio was generally lower than 1.0 at most

intensities tested in both species (Figure 5A, 5B: diamond with dotted line).

Monaural Amplitude Ratio

Monaural amplitude ratio was calculated by dividing the amplitude value of wave IV by

the amplitude value of wave I for left and right pinnae at each intensity. As displayed in figure

5C, the wave IV/I amplitude ratio typically decreased with increasing intensity from 60 to 90 dB

SPL for both species (Figure 5A, 5B). A linear mix-effect model revealed significant statistical

differences of intensity on the amplitude ratio for *P. maniculatus* (LMM: p-value = 0.014).

However, no statistically significant differences of either intensity or sexes were observed on the

amplitude ratio for *P. leucopus* (Intensity: LMM: p-value = 0.332; Sex: LMM: p-value = 0.84).

When combined, the results of the linear mix-effect model revealed no significant main effects

of either intensity (LMM: p-value = 0.332) or species (LMM: p-value = 0.474) on the amplitude

ratio of wave I and IV between species.

Figure 5: Average amplitude of wave I (filled circles) and wave IV (unfilled circles) of auditory brainstem responses determines with clicks of different intensities (Pink = *P. leucopus* (n = 15), Blue = *P. maniculatus* (n = 11)). The average wave IV/I amplitude ratio at each intensity (filled diamond with dotted line represents wave IV/I for *P. leucopus* and unfilled diamond with dotted line represents wave IV/I for *P. maniculatus*) is shown in each figure (right ordinate). The vertical bars represent the standard error at each point. Significant main effects of intensity on wave I and IV amplitude were detected for each species.

371
Absolute Latency

We next calculated the average peak latency of waves I and IV of both studied species

across click intensities (60 to 90 dB SPL). We detected no significant decrease in peak latency

for either wave I or wave IV with increasing intensity for *P. leucopus* (Wave I: LMM: p-value =

0.353; Wave IV: LMM: p-value = 0.122) (Figure 6A). Similarly, no significant statistical

differences were observed in peak latency for either wave I and wave IV between male and

female *P. leucopus* (Wave I: LMM: p-value = 0.841; Wave IV: p-value = 0.341). There were no

significant statistical differences in peak latency for either wave I or wave IV with increasing

intensity for *P. maniculatus* (Wave I: LMM: p-value = 0.353; wave IV: LMM: p-value = 0.392)

(Figure 6B). When data were combined, we detected no significant main effects of either

intensity or species in peak latency for either wave I or wave IV with increasing intensity

between both species (LMM: All p-value >0.05).

Inter-peak latency

Inter-peak latency was calculated as the difference in latency from the wave I peak to

the other designated peak (IV) for left and right pinnae at each intensity. We observed a

significant decreased in wave I-IV inter peak latency (Figure 6C) with increasing intensity for *P.*

leucopus (LMM: p-value = 0.043). However, no significant decrease in wave I-IV inter peak

latency was detected with increasing intensity between male and female *P. leucopus* (LMM: p-

value = 0.341). There was a significant decrease in wave I-IV inter peak latency (Figure 6C) with

increasing intensity for *P. maniculatus* (LMM: p-value = 0.010). When data were combined, we

observed a significant main effect of intensity on the inter-peak latency of waves I and IV

between both species (LMM: p-value = 0.002). However, no main effect of species was detected

on the inter-peak latency of waves I and IV between both species (LMM: p-value = 0.145).

398
Figure 6: Average peak latency of wave I (filled circles) and wave IV (unfilled circles) of auditory
brainstem responses determined with clicks of different intensities (Pink = *P. leucopus* (n = 15), Blue = *P.*
maniculatus (n = 11)). The average wave I-IV inter-peak latency at each intensity (filled diamond with
dotted line represents wave I-IV for *P. leucopus* and unfilled diamond with dotted line represents wave I-
IV for *P. maniculatus*) is shown in each figure (right ordinate). The vertical bars represent the standard
error at each point.

Slope latency intensity function between species

Peak latency is the time interval between the presentation of a sound stimulus and

the peak at maximum amplitude of the designated wave. For both waves I and IV, we calculated

the slope of each latency intensity function following the methods outlined by Zhou et al. 2006

(Figure 2 B(a), B(b), View methods section). As displayed in figure 3B, the slope of latency-

intensity function of the exemplar female *P. leucopus* rodent was 7.80 µs/dB for wave I and

11.33 µs/dB for wave IV. *P. leucopus* had an average slope latency-intensity function of 6.386

µs/dB for wave I and 14.088 µs/dB for wave IV, while the average slope of latency-intensity

function for wave I and wave IV of *P. maniculatus* was 7.291 µs/dB and 12.905 µs/dB,

respectively (Figure 7). We detected significant main effects of wave number on slope of the

latency-intensity function (LMM: p-value = 0.0003). However, no significant main effects of

species were detected on slope of the latency-intensity function of both waves I and IV (LMM:

p-value = 0.906). No pairwise comparisons were made for species since there was no main

effect. A linear mix-effect model revealed that the slope of the latency-intensity function of wave

IV was larger than wave I in both *P. leucopus* (t-value = -3.562, p-value = 0.001), and *P.*

maniculatus (t-value = -2.084, p-value = 0.048).

Figure 7: Average slope of latency-intensity function of waves I, and IV of ABRs (Pink = *P. leucopus* (n = 15), Blue = *P. maniculatus* (n = 11)).

Binaural hearing measures

We used the latency shift of the DN1 component of the BIC and relative DN1 amplitude to show the relationship of ITD on the latency and relative amplitude of the BIC in both studied species. The average DN1 amplitudes at 0 ITD were 2.72 µV and 1.74 µV for *P. maniculatus* and *P. leucopus*, respectively. The average latency for the DN1 component for 0 ITD was 5.01 ms for *P. maniculatus*, compared with 5.6 ms for *P. leucopus*. Linear mixed-effects models indicated no significant differences between *P. maniculatus* and *P. leucopus* across relative DN1 amplitude in relation to ITD normalized to the DN1 amplitude for 0 ITD (Figure 8B) (LMM: p-value = 0.82). Similarly, there were no significant statistical differences across relative amplitude in relation to ITD normalized to the DN1 amplitude for 0 ITD between male and female *P.*

leucopus (LMM: p-value > 0.05). There were statistically significant differences in latency shift

of the DN1 component of the BIC in relation to ITD normalized for 0 ITD between *P.*

maniculatus and *P. leucopus* (Figure 8A) (LMM: p = 0.016). Shift in DN1 latencies of the BIC

were significantly faster in *P. maniculatus* compared to *P. leucopus* at 1.0 ms (t-value = 2.101, p-

value = 0.037) and 2.0 ms (t-value = 2.316, p-value = 0.022) ITD (Figure 8A). When we added

body size to our LMM to test if body size contributes to BIC latencies, we saw a significant

effect of body size with a significant effect of species (all p-values < 0.05). We detected no

significant statistical differences in latency shift of the DN1 component of the BIC in relation to

ITD normalized for 0 ITD between male and female *P. leucopus* (LMM: p-value = 0.843).

446
447

Figure 8: Binaural ABRs in wild *P. leucopus* (pink filled square) and *P. maniculatus* (blue unfilled
circle). 8A, Shift in DN1 latency (ms) relative to ITD; reference latency at ITD = 0 is set to 0 ms. 8B,
percentage relative DN1 amplitude relative to ITD normalized to the DN1 amplitude for ITD = 0 ms.
Relative amplitude and latency of the DN1 BIC with varying ITD between - 2 to + 2 ms in 0.5 ms steps
were measured. Significant differences were detected in BIC shift in DN1 latencies between both species
at ITDs 1.0 and 2.0 ms. No significant differences were observed between both species for relative
amplitude of the BIC across ITDs.
456
457

DISCUSSION

In this study, we used craniofacial morphology, pinna features, and ABRs to compare

morphological features important for hearing with physiological measures of ABR amplitude

and latency of two species of the genus *Peromyscus*. Like previous findings (Choate, 1973; Light

et al., 2021; Millien et al., 2017), we detected significant morphological differences between

both species including pinna length, pinna width, effective pinna diameter, inter-pinna distance,

and other measures with *P. leucopus* displaying larger features. ABR-derived detection threshold

revealed that both species share similar ABR response threshold across frequencies with the best

frequency hearing between 8-24 kHz, which is in agreement with previous findings that showed

Peromyscus species have best hearing sensitivity between 8-16 kHz (Capshaw et al., 2022; Dice

and Barto, 1952; Ralls, 1967). Significant main effects of intensity were detected in monaural

amplitude of ABR wave I and IV between both studied species, which is in accordance with

similar findings using laboratory strains mice (Zhou et al., 2006). Measurements of the BIC,

indicated similar amplitude across ITDs with differences in latency of the BIC across ITDs

between the two species. Overall, our results revealed that both species have similar ABR best

frequency threshold with *P. maniculatus* slightly having shorter latency BIC and smaller

anatomical features compared to *P. leucopus*.

Morphological features including cranial size and shape, size of the pinnae, body and tail

length differ widely within species of the genus *Peromyscus* (Light et al., 2021; Ordóñez-Garza

et al., 2010). Our data showed differences in all measured anatomical traits between species

consistent with previous studies (Choate, 1973; Light et al., 2021; Millien et al., 2017). Previous

studies used two-dimensional (2D) geometric morphometrics (Light et al., 2021), and micro-CT

(Riede et al., 2022) as tools to morphologically differentiate rodents of the genus *Peromyscus*.

Light *et al* (2021) showed differences between *P. leucopus* and *P. maniculatus* based on head

size, pinnae features, hindfoot length, and other morphological features. Consistency in

morphological features (pinna length, pinna width, and body weight) documented in this study

provide additional evidence supporting the use of these morphology traits as reliable indicators

for distinguishing species within the genus *Peromyscus*.

Using ABRs, Capshaw *et al.* (2022) observed decreased hearing sensitivity to frequencies

below 2 kHz in two laboratory *Peromyscus* species (*P. leucopus* and *P. californicus*). Our

findings are consistent with Capshaw *et al.* (2022), with hearing thresholds around 85 dB SPL at

frequencies below 2 kHz in both studied species, suggesting relatively poor hearing sensitivity of

both studied species to frequencies 1-2 kHz. Small-headed mammals generally are not as

sensitive to low frequencies and therefore do not generate significant directional information

using low frequencies, where differences in timing and intensity between pinnae are minimal

(Lauer et al., 2018). Therefore, it is thought that small mammals rely on high frequencies for

directional hearing with exception of some subterranean mammals including the naked mole-rat

(*Heterocephalus glaber*), the plain pocket gopher (*Geomys bursarius*), and the blind mole rat

(*Spalax ehrenbergi*) that lack the capability to localize sound and lack high frequency hearing

(Heffner and Heffner, 1993, 1992, 1990), though see (Barker et al., 2021; Gessele et al., 2016;

McCullagh et al., 2022). The limited ability of small mammals (with exception of Mongolian

gerbils, chipmunks, groundhogs, hamsters, and others), like members of the genus *Peromyscus*,

to detect low frequency sounds has been attributed to selective pressure linked with the absence

of cues for localizing sounds in the horizontal plane (Heffner et al., 2001). Therefore, it is not

surprising that we did not observe low frequency sensitivity between the two studied species in

this current investigation.

P. leucopus and *P. maniculatus* are both highly territorial and produce both sonic and

ultrasonic vocalizations between 0.8 to 28 kHz (sustained vocalizations: frequency ranges

between 10-25 kHz, sweep vocalization: frequencies above 25 kHz, and barks: frequency ranges

between 0.8 and 6 kHz) (Miller and Engstrom, 2012; Pomerantz and Clemens, 1981; Riede et al.,

2022). The frequency ranges of ultrasonic vocalizations of both studied species correlate with

their best frequency threshold (Figure 4C, best frequency threshold ranging from 8-24 kHz).

Related species' (California mouse, *P. californicus*) defensive and distress vocalizations are

known to be associated with sounds ranging from 2 -30 kHz (Rieger and Marler, 2018). While

limited studies have described distress and defensive vocalizations across the genus *Peromyscus*,

previous investigations have reported that members of this genus produce agonistic calls such as

chits and barks at frequencies between 6 to 15 kHz (Houseknecht, 1968; Pasch et al., 2017).

These agonistic calls are likely associated with lower auditory thresholds at these frequencies for

the genus *Peromyscus*. These findings suggest that the good match of *Peromyscus's* vocalization

with their frequency threshold sensitivity (8 – 24 kHz) likely contributes to vocal air-borne

communication in the wild. In addition, *Peromyscus* species are relatively long-lived but due to

limited studies on the ability of *Peromyscus* to hear sound, it is hard to speculate the

physiological mechanisms that govern hearing sensitivity over their lifespan. It is possible that

the decreased emission of mitochondrial reactive oxygen species and improved activity of

antioxidant enzymes might play key roles in sustaining healthy auditory sensitivity across

Peromyscus species (Csiszar et al., 2007).

The white-footed mice (*P. leucopus*) and the deer mice (*P. maniculatus*) occur

throughout Oklahoma but generally occupy different habitats, with *P. maniculatus* being more

common in grasslands and *P. leucopus* primarily inhabiting shaded forests (Hackney and

Stancampiano, 2015; Stancampiano and Schnell, 2004). In our study, *P. leucopus* subjects were

mainly captured in shrubland and forested habitats, while *P. maniculatus* subjects were found in

open grassland habitats. Our findings revealed that *P. leucopus* has similar sensitivity to sound as

P. maniculatus across all frequencies tested, except at 1 kHz (t-value = 2.009, p-value = 0.046),

where *P. maniculatus* show slightly better hearing. One possibility is that slightly higher

frequency hearing sensitivity of *P. leucopus* may have coevolved with their vocal signal

characteristics to facilitate effective communication in forested and shrubland environments,

where acoustic information is often encoded at higher frequencies (Charlton et al., 2019). In

addition, weight distribution suggests that eight of the 11 *P. maniculatus* subjects in the current

study were juveniles, while 10 of the 15 *P. leucopus* were adults. Age differences could also

explain the shifted high frequency hearing in *P. leucopus* compared to *P. maniculatus,* as small

shifts in audiogram threshold has been observed in *P. leucopus* with aging (Capshaw et al.,

2022). A comparative study evaluating the vocalization content and sound attenuation of both

species in their respective habitats, across different age groups, would shed novel insights into

how habitat-related factors and age might influence the evolution of sound reception and

communication strategy both within and among closely related *Peromyscus* species.

Amplitude of wave I and IV tend to increase monotonically in most small mammals with

increasing intensity when measured by click stimuli (Zhou et al., 2006). Similar patterns have

been reported in other taxa commonly used in evoked potential studies (Backoff and Caspary,

1994; Neil J. Ingham, 1998). Observed differences in wave amplitudes between the two species

is likely a result of difference in craniofacial size relative to body mass. Previous studies indicate

that smaller craniofacial size with small body mass may bring the recording electrodes into

closer proximity to the generators, resulting in larger amplitudes compared to those with large

body mass (Merzenich et al. 1983). Prior publications indicate that other factors such as neural

synchronicity and the number of neural elements firing in the generators can also contribute to

the amplitude of ABR waves (Merzenich et al., 1983).

ABR wave amplitude can be affected by several factors including electrode position,

animal body temperature, external noise, recording protocol, and equipment characteristics,

therefore normalization between waves can help control for this variability. In humans, it has

been shown that auditory deficits related to retrocochlear pathology may lead to a decrease in

wave IV amplitude, and ultimately cause a decrease in the wave IV/I amplitude ratio (Arnold

2000). Our data revealed that the wave IV has a smaller amplitude than wave I in both species at

most intensities tested, resulting in a wave IV/I smaller than 1.0 (Figure 5C). Previous work

measuring ABR in inbred mouse strains, rats, gerbils, cats, guinea pigs, and humans indicated

that the ABR waves I and II are generally larger amplitude than ABR waves III and IV, which is

in agreement to this current results (Moore, 1983). However, wave II and III are relatively larger

in rats and guinea pigs, while shifted to wave IV in cats and wave IV-V complex in humans

(Merzenich et al., 1983). Accordingly, the species-specific differences in individual ABR wave

amplitude may result from complex factors including the evolution of the central nervous

system, neuronal response characteristics within the brainstem, and the neural conduction

velocity.

The slope of the latency-intensity function when combined with ABR threshold has been

shown to be a useful parameter to estimate hearing sensitivity (Zhou et al., 2006). Previous

studies have reported the slope of the latency-intensity function of wave I and IV of different

laboratory inbred strains of mice, gerbils, cats, and humans (Burkard et al., 1990; Burkard and

Voigt, 1989; Fullerton et al., 1987; Zhou et al., 2006). Zhou et al. described that the slope of the

latency-intensity functions of wave I and IV were 4.1 to 14.0 μs/dB in laboratory inbred stains of

mice (BALB/cJ, C3H/HeJ/ SJL/J, CBA/j, ect.) (Zhou et al., 2006). In gerbils and rats, wave I and

IV slope latency-intensity function have been reported to be ~ 8 to 9 and ~13 to 16 μs/dB,

respectively (Burkard et al., 1990; Burkard and Voigt, 1989). Other publications reported that the

slope of the latency-intensity function of wave I and IV were ~14 to 16 μs/dB in cats (Fullerton

et al., 1987). In addition, the slope of the latency-intensity function of wave V and other ABR

waves was ~ 40 μs/dB in humans and Dalmatian puppies but were ~ 28 μs/dB in Beagle puppies

(Burkard and Hecox, 1983; Poncelet et al., 2000). Accordingly, we conclude that the slope of the

latency-intensity function of wild *Peromyscus* rodents ABR waves is similar to that of laboratory

inbred strain of mice and gerbils, slightly less than those of rats and cats, but significantly less

than those of humans and dogs.

ITD and ILD are two cues that animals with external pinnae use for sound localization.

ITDs are generally processed by neurons in the medial superior olive (MSO <2kHz) while ILDs

are mainly processed by neurons in the lateral superior olive (LSO >2 kHz) (Grothe et al., 2010;

Suzuki and Horiuchi, 1981). Previous studies reported that the mean DN1 amplitude at 0 ITD

was 0.2 µV in humans, about 5 µV in guinea pig, 1.8 µV in gerbil, and 2.3 µV in cats (Goksoy et

al., 2005; Jones and Van der Poel, 1990; Laumen et al., 2016a; Riedel and Kollmeier, 2006;

Ungan et al., 1997). Comparatively, the DN1 amplitude of 0 ITD of wild *Peromyscus* species is

similar to that of gerbils and cats, higher than that of humans and significantly lower than that of

guinea pig. Differences in DN1 amplitude at 0 ITD observed could be a result of smaller distance

of the recorded electrodes to the subjects, as well as electrode configuration, or other procedural

differences.

Numerous publications have reported that latency of the DN1 component in humans

ranges from 5.6 to 6.8 ms, while those of other animal models (gerbil, cats, guinea pig) range

from 3.7 to 4.8 ms (Goksoy et al., 2005; Jones and Van der Poel, 1990; Riedel and Kollmeier,

2006; Ungan et al., 1997). Our data of the latency DN1 component is consistent with latencies

observed in human and is somewhat slower than what is seen in other animal models. Our results

are similar with others that show the latency of the DN1 component increases with longer ITDs

in cats, gerbil, guinea pig, and humans (Goksoy et al., 2005; Laumen et al., 2016b; Riedel and

Kollmeier, 2006; Ungan et al., 1997). Indeed, it has been suggested that the increase in DN1

latency with increasing ITD reflects the anatomy and interaction between excitatory and

inhibitory neurons in the superior olivary complex (Karino et al., 2011).

We observed faster latencies in DN1 in *P. maniculatus* compared to *P. leucopus*. It is hard

to speculate whether the difference in DN1 latency observed between both species is associated

with head size or the number of cells in the SOC nuclei. Studies characterizing the number of

excitatory and inhibitory cells in the SOC of both species would be beneficial to allow for

evaluation of the effects of head size or MSO and LSO size in shifts of the DN1 latency among

Peromyscus species. Further studies involving more *Peromyscus* species and other techniques,

such as head-related transfer functions, are needed to assess if larger external pinna sizes

contribute to additional features of *Peromyscus* hearing such as the use of spectral notches and

the contribution of the pinna to horizontal cues like ITD and ILD, particularly since our in-ear

presentation of ITD stimuli bypass the pinna. We calculated the functional interaural distance for

each species by summing the mean inter-pinna distance and pinna width divided by the speed of

sound in air to evaluate the availability of ITD cues for each species. While this technique is

limited due to our use of calipers and is not exactly the same as the time delay caused by sound

traveling around the head, we nonetheless used this is to roughly estimate the functional

interaural distance for each species. We found that *P. maniculatus* have a shorter functional

interaural distance ($\pm$ 55 μs) compared to *P. leucopus* ($\pm$ 67 μs) which is consistent with smaller

heads in *P. maniculatus*.

There are some limitations to the techniques employed in this study. Calipers are less

accurate as features get smaller due to their measurement sensitivity, therefore measures of pinna

and head morphology are likely to be less accurate than larger measurements such as body length

and tail length. We conducted analyses correcting for overall body length; however, they did not

show significant positive allometry (slope > 1) indicating that either these features were not

allometric or the loss of accuracy of measurements at smaller distances contributed significantly

to error. However, the one measurement that showed positive allometry was tail length, which is

one of the longer, or perhaps more accurate, measures suggesting that finer measurement tools

might be needed to make further arguments about effects of overall body size and morphological

features on hearing in these species. There are also limitations to using ABRs as measures of

hearing, including that interpretation of thresholds using visual observation, as performed in our

study, can be subjective (Suthakar and Liberman, 2019). However, others have shown minimal

differences between algorithms and observers to auditory threshold measurements (Capshaw et

al., 2022). Further validation of our observer method with more quantitative algorithms would be

useful to confirm threshold values reported here, though our thresholds coincide well with the

published literature in one of these species (Capshaw et al., 2022). Lastly, behavioral measures

of hearing can show differences compared to ABRs, and indeed anesthetics used, montage of

electrodes, calibration of sounds (in ear or other methods), sound presentation, and other factors

all may influence ABR results making cross-species and cross-publication results difficult to

interpret (Ramsier and Dominy, 2010; Wolski et al., 2003). However, the current study used the

same parameters across both species and showed results consistent with the literature and what

might be expected for species that are closely related but differ primarily in size giving us

confidence in the results presented here.

CONCLUSIONS

Our findings provide a deeper understanding of auditory similarities and differences

between two species of *Peromyscus* and validate that the highly abundant *Peromyscus* may serve

as a future model for auditory studies. Both species show differences in craniofacial and pinna features and exhibit best hearing thresholds at frequencies ranging from 8 to 24 kHz. *P. maniculatus* showed shorter relative latencies of the DN1 component of the BIC, while relative DN1 amplitude was not different between the species. Further physiological assessment exploring hearing between the sexes at different ages and across the lifespan are needed to further show whether there are differences in hearing in under these conditions. In addition, clarifying the role of the BIC between sexes across species of the genus *Peromyscus* is important to understand its relevance for sex differences.

ACKNOWLEDGMENTS

We would like to thank Game wardens, Benny Farrar and Marcus Thibodeau for housing and giving us opportunities to sample at James Collin and Packsaddle Wildlife Management Areas. Also, we would like to thank members of team wild rodent of the McCullagh lab which helped in trapping and performed ABRs. We would also like to thank Dr. Tim Lei and Benzheng Li for their creation of the ABR acquisition and analysis custom python software. Dr. Fabio Machado helped with interpreting analyses of allometry and body size measurements. NIH NICHD funding 1R15HD105231-01 and 3R15HD105231-01S1, NSF RaMP DEB 2216648, and Oklahoma State University College of Arts & Sciences (CAS) Research Program to EAM helped fund summer support, RA support for LJ and undergraduates involved in the project, and some materials (CAS research award). Support for VYF was provided by a Wentz and CAS AURCA program support, and EMN with Wentz fellowship support as well as additional support for LJ from the Payne County Audubon Society.

DATA AVAILABILITY

The data of the study will be made available upon request.

AUTHOR CONTRIBUTIONS

LJ, EMN, TCW, VYF, And DMJ captured the animals and LJ and EMN collected the ABR data for the manuscript. LJ, DMJ, and GOUW performed DNA analysis on tails snip samples while BL helped with data analysis and interpretation. LJ and EAM completed the statistical analysis and developed the idea of the paper. LJ wrote the manuscript and all other authors revised and edited the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

Anbuhl, K.L., Benichoux, V., Greene, N.T., Brown, A.D., Tollin, D.J., 2017. Development of the head, pinnae, and acoustical cues to sound location in a precocial species, the guinea pig (Cavia porcellus). Hearing Research 356, 35–50. https://doi.org/10.1016/j.heares.2017.10.015

Arnold: The auditory brainstem response - Google Scholar, n.d.

Backoff, P.M., Caspary, D.M., 1994. Age-related changes in auditory brainstem responses in fischer 344 rats: effects of rate and intensity. Hearing Research 73, 163–172. https://doi.org/10.1016/0378-5955(94)90231-3

Barker, A.J., Koch, U., Lewin, G.R., Pyott, S.J., 2021. Hearing and Vocalizations in the Naked Mole-Rat, in: Buffenstein, R., Park, T.J., Holmes, M.M. (Eds.), The Extraordinary Biology of the Naked Mole-Rat. Springer International Publishing, Cham, pp. 157–195. https://doi.org/10.1007/978-3-030-65943-1_6

Bates, D., Mächler, M., Bolker, B., Walker, S., 2014. Fitting Linear Mixed-Effects Models using lme4. https://doi.org/10.48550/ARXIV.1406.5823

Bedford, N.L., Hoekstra, H.E., 2015. Peromyscus mice as a model for studying natural variation. eLife 4, e06813. https://doi.org/10.7554/eLife.06813

Benichoux, V., Ferber, A., Hunt, S., Hughes, E., Tollin, D., 2018. Across Species "Natural Ablation" Reveals the Brainstem Source of a Noninvasive Biomarker of Binaural Hearing. J. Neurosci. 38, 8563–8573. https://doi.org/10.1523/JNEUROSCI.1211-18.2018

Blauert, J., 1997. Spatial Hearing: The Psychophysics of Human Sound Localization. MIT Press.

Bradley, R.D., Durish, N.D., Rogers, D.S., Miller, J.R., Engstrom, M.D., Kilpatrick, C.W., 2007. Toward a Molecular Phylogeny for Peromyscus: Evidence from Mitochondrial Cytochrome-b Sequences. Journal of Mammalogy 88, 1146–1159. https://doi.org/10.1644/06-MAMM-A-342R.1

Brittan-Powell, E.F., Dooling, R.J., 2004. Development of auditory sensitivity in budgerigars (*Melopsittacus undulatus*). The Journal of the Acoustical Society of America 115, 3092–3102. https://doi.org/10.1121/1.1739479

Burger, J., Gochfeld, M., 1992. Survival and reproduction in Peromyscus leucopus in the laboratory: viable model for aging studies. Growth Dev Aging 56, 17–22.

Burkard, R., Feldman, M., Voigt, H.F., 1990. Brainstem Auditory-Evoked Response in the Rat Normative Studies, with Observations Concerning the Effects of Ossicular Disruption. Audiology 29, 146–162. https://doi.org/10.3109/00206099009072847

Burkard, R., Hecox, K., 1983. The effect of broadband noise on the human brainstem auditory evoked response. I. Rate and intensity effects. The Journal of the Acoustical Society of America 74, 1204–1213. https://doi.org/10.1121/1.390024

Burkard, R., Voigt, H.F., 1989. Stimulus dependencies of the gerbil brain-stem auditory-evoked response (BAER). I: Effects of click level, rate, and polarity. The Journal of the Acoustical Society of America 85, 2514–2525. https://doi.org/10.1121/1.397746

Caire, W., 1989. Mammals of Oklahoma. University of Oklahoma Press.

Capshaw, G., Brown, A.D., Peña, J.L., Carr, C.E., Christensen-Dalsgaard, J., Tollin, D.J., Womack, M.C., McCullagh, E.A., 2023. The continued importance of comparative

auditory research to modern scientific discovery. Hearing Research 433, 108766. https://doi.org/10.1016/j.heares.2023.108766

Capshaw, G., Vicencio-Jimenez, S., Screven, L.A., Burke, K., Weinberg, M.M., Lauer, A.M., 2022. Physiological Evidence for Delayed Age-related Hearing Loss in Two Long-lived Rodent Species (Peromyscus leucopus and P. californicus). JARO 23, 617–631. https://doi.org/10.1007/s10162-022-00860-4

Charlton, B.D., Owen, M.A., Swaisgood, R.R., 2019. Coevolution of vocal signal characteristics and hearing sensitivity in forest mammals. Nat Commun 10, 2778. https://doi.org/10.1038/s41467-019-10768-y

Chawla, A., McCullagh, E.A., 2022. Auditory Brain Stem Responses in the C57BL/6J Fragile X Syndrome-Knockout Mouse Model. Frontiers in Integrative Neuroscience 15.

Childs, J.E., Ksiazek, T.G., Spiropoulou, C.F., Krebs, J.W., Morzunov, S., Maupin, G.O., Gage, K.L., Rollin, P.E., Sarisky, J., Enscore, R.E., Frey, J.K., Peters, C.J., Nichol, S.T., 1994. Serologic and Genetic Identification of Peromyscus maniculatus as the Primary Rodent Reservoir for a New Hantavirus in the Southwestern United States. Journal of Infectious Diseases 169, 1271–1280. https://doi.org/10.1093/infdis/169.6.1271

Choate, J.R., 1973. Identification and Recent Distribution of White-Footed Mice (Peromyscus) in New England. Journal of Mammalogy 54, 41–49. https://doi.org/10.2307/1378871

Colburn, H.S., Zurek, P.M., Durlach, N.I., 1987. Binaural Directional Hearing—Impairments and Aids, in: Yost, W.A., Gourevitch, G. (Eds.), Directional Hearing, Proceedings in Life Sciences. Springer, New York, NY, pp. 261–278. https://doi.org/10.1007/978-1-4612-4738-8_11

Csiszar, A., Labinskyy, N., Zhao, X., Hu, F., Serpillon, S., Huang, Z., Ballabh, P., Levy, R.J., Hintze, T.H., Wolin, M.S., Austad, S.N., Podlutsky, A., Ungvari, Z., 2007. Vascular superoxide and hydrogen peroxide production and oxidative stress resistance in two closely related rodent species with disparate longevity. Aging Cell 6, 783–797. https://doi.org/10.1111/j.1474-9726.2007.00339.x

Cummings, J.R., Vessey, S.H., 1994. Agricultural Influences on Movement Patterns of White-Footed Mice (Peromyscus leucopus). The American Midland Naturalist 132, 209–218. https://doi.org/10.2307/2426575

Dammann, P., 2017. Slow aging in mammals—Lessons from African mole-rats and bats. Seminars in Cell & Developmental Biology, Science communication in the field of fundamental biomedical research 70, 154–163. https://doi.org/10.1016/j.semcdb.2017.07.006

Dewey, M.J., Dawson, W.D., 2001. Deer mice: "The Drosophila of North American mammalogy." genesis 29, 105–109. https://doi.org/10.1002/gene.1011

Dice, L.R., Barto, E., 1952. Ability of Mice of the Genus Peromyscus to Hear Ultrasonic Sounds. Science 116, 110–111. https://doi.org/10.1126/science.116.3005.110

Ducroz, J.F., Volobouev, V., Granjon, L., 2001. An Assessment of the Systematics of Arvicanthine Rodents Using Mitochondrial DNA Sequences: Evolutionary and Biogeographical Implications. Journal of Mammalian Evolution 8, 173–206. https://doi.org/10.1023/A:1012277012303

Ehret, G., Dreyer, A., 1984. Localization of tones and noise in the horizontal plane by unrestrained house mice (Mus musculus). Journal of Experimental Biology 109, 163–174. https://doi.org/10.1242/jeb.109.1.163

Fairbairn, D.J., 1977. The spring decline in deer mice: death or dispersal? Can. J. Zool. 55, 84–92. https://doi.org/10.1139/z77-009

Fiset, J., Tessier, N., Millien, V., Lapointe, F.-J., 2015. Phylogeographic Structure of the White-Footed Mouse and the Deer Mouse, Two Lyme Disease Reservoir Hosts in Québec. PLoS ONE 10, e0144112. https://doi.org/10.1371/journal.pone.0144112

Fullerton, B.C., Levine, R.A., Hosford-Dunn, H.L., Kiang, N.Y.S., 1987. Comparison of cat and human brain-stem auditory evoked potentials. Electroencephalography and Clinical Neurophysiology 66, 547–570. https://doi.org/10.1016/0013-4694(87)90102-7

Furst, M., Eyal, S., Korczyn, A.D., 1990. Prediction of binaural click lateralization by brainstem auditory evoked potentials. Hear Res 49, 347–359. https://doi.org/10.1016/0378-5955(90)90113-4

Gessele, N., Garcia-Pino, E., Omerbašić, D., Park, T.J., Koch, U., 2016. Structural Changes and Lack of HCN1 Channels in the Binaural Auditory Brainstem of the Naked Mole-Rat (Heterocephalus glaber). PLOS ONE 11, e0146428. https://doi.org/10.1371/journal.pone.0146428

Goksoy, C., Demirtas, S., Yagcioglu, S., Ungan, P., 2005. Interaural delay-dependent changes in the binaural interaction component of the guinea pig brainstem responses. Brain Research 1054, 183–191. https://doi.org/10.1016/j.brainres.2005.06.083

Grothe, B., Pecka, M., McAlpine, D., 2010. Mechanisms of sound localization in mammals. Physiol Rev 90, 983–1012. https://doi.org/10.1152/physrev.00026.2009

Guo, Z., Wang, M., Tian, G., Burger, J., Gochfeld, M., Yang, C.S., 1993. Age- and gender-related variations in the activities of drug-metabolizing and antioxidant enzymes in the white-footed mouse (Peromyscus leucopus). Growth Dev Aging 57, 85–100.

Hackney, S., Stancampiano, A.J., 2015. Microhabitat Preferences of a Small Mammal Assemblage in Canadian County, Oklahoma. Proceedings of the Oklahoma Academy of Science 95.

Harney, B.A., Dueser, R.D., 1987. Vertical Stratification of Activity of Two Peromyscus Species: An Experimental Analysis. Ecology 68, 1084–1091. https://doi.org/10.2307/1938380

Heeringa, A.N., Zhang, L., Ashida, G., Beutelmann, R., Steenken, F., Köppl, C., 2020. Temporal Coding of Single Auditory Nerve Fibers Is Not Degraded in Aging Gerbils. J. Neurosci. 40, 343–354. https://doi.org/10.1523/JNEUROSCI.2784-18.2019

Heffner, H.E., Heffner, R.S., 1985. Hearing in two cricetid rodents: Wood rat (Neotoma floridana) and grasshopper mouse (Onychomys leucogaster). Journal of Comparative Psychology 99, 275–288. https://doi.org/10.1037/0735-7036.99.3.275

Heffner, H.E.H., Rickye S., 2001. Behavioral Assessment of Hearing in Mice, in: Handbook of Mouse Auditory Research. CRC Press.

Heffner, R.S., Heffner, H.E., 1993. Degenerate hearing and sound localization in naked mole rats (Heterocephalus glaber), with an overview of central auditory structures. J of Comparative Neurology 331, 418–433. https://doi.org/10.1002/cne.903310311

Heffner, R.S., Heffner, H.E., 1992. Hearing and sound localization in blind mole rats (Spalax ehrenbergi). Hearing Research 62, 206–216. https://doi.org/10.1016/0378-5955(92)90188-S

Heffner, R.S., Heffner, H.E., 1990. Vestigial hearing in a fossorial mammal, the pocket gopher (Geomys bursarius). Hearing Research 46, 239–252. https://doi.org/10.1016/0378-5955(90)90005-A

Heffner, R.S., Heffner, H.E., 1982. Hearing in the elephant (Elephas maximus): absolute
sensitivity, frequency discrimination, and sound localization. J Comp Physiol Psychol 96,
926–944.
Heffner, R.S., Koay, G., Heffner, H.E., 2020. Hearing and sound localization in Cottontail
rabbits, Sylvilagus floridanus. J Comp Physiol A 206, 543–552.
https://doi.org/10.1007/s00359-020-01424-8
Heffner, R.S., Koay, G., Heffner, H.E., 2001. Audiograms of five species of rodents:
implications for the evolution of hearing and the perception of pitch. Hearing Research
157, 138–152. https://doi.org/10.1016/S0378-5955(01)00298-2
Houseknecht, C.R., 1968. Sonographic Analysis of Vocalizations of Three Species of Mice.
Journal of Mammalogy 49, 555. https://doi.org/10.2307/1378232
Jones, S.J., Van der Poel, J.C., 1990. Binaural interaction in the brain-stem auditory evoked
potential: evidence for a delay line coincidence detection mechanism.
Electroencephalography and Clinical Neurophysiology/Evoked Potentials Section 77,
214–224. https://doi.org/10.1016/0168-5597(90)90040-K
Jüchter, C., Beutelmann, R., Klump, G.M., 2022. Speech sound discrimination by Mongolian
gerbils. Hearing Research 418, 108472. https://doi.org/10.1016/j.heares.2022.108472
Karino, S., Smith, P.H., Yin, T.C.T., Joris, P.X., 2011. Axonal Branching Patterns as Sources of
Delay in the Mammalian Auditory Brainstem: A Re-Examination. J. Neurosci. 31, 3016–
3031. https://doi.org/10.1523/JNEUROSCI.5175-10.2011
Kidd, G., Mason, C.R., Rohtla, T.L., 1995. Binaural advantage for sound pattern identification.
The Journal of the Acoustical Society of America 98, 1977–1986.
https://doi.org/10.1121/1.414459
King, J.A., 1968. Biology of Peromyscus (Rodentia). Biology of Peromyscus (Rodentia).
Kirkland, G.L., Layne, J.N., 1989. Advances in the Study of Peromyscus (Rodentia) Texas Tech
University Press. Lubbock, TX.
Labinskyy, N., Mukhopadhyay, P., Toth, J., Szalai, G., Veres, M., Losonczy, G., Pinto, J.T.,
Pacher, P., Ballabh, P., Podlutsky, A., Austad, S.N., Csiszar, A., Ungvari, Z., 2009.
Longevity is associated with increased vascular resistance to high glucose-induced
oxidative stress and inflammatory gene expression in Peromyscus leucopus. American
Journal of Physiology-Heart and Circulatory Physiology 296, H946–H956.
https://doi.org/10.1152/ajpheart.00693.2008
Larson, S.R., Lee, X., Paskewitz, S.M., 2018. Prevalence of Tick-Borne Pathogens in Two
Species of Peromyscus Mice Common in Northern Wisconsin. Journal of Medical
Entomology 55, 1002–1010. https://doi.org/10.1093/jme/tjy027
Lauer, A.M., Engel, J.H., Schrode, K., 2018. Rodent Sound Localization and Spatial Hearing, in:
Dent, M.L., Fay, R.R., Popper, A.N. (Eds.), Rodent Bioacoustics, Springer Handbook of
Auditory Research. Springer International Publishing, Cham, pp. 107–130.
https://doi.org/10.1007/978-3-319-92495-3_5
Laumen, G., Ferber, A.T., Klump, G.M., Tollin, D.J., 2016a. The Physiological Basis and
Clinical Use of the Binaural Interaction Component of the Auditory Brainstem Response.
Ear Hear 37, e276–e290. https://doi.org/10.1097/AUD.0000000000000301
Laumen, G., Tollin, D.J., Beutelmann, R., Klump, G.M., 2016b. Aging effects on the binaural
interaction component of the auditory brainstem response in the Mongolian gerbil:
Effects of interaural time and level differences. Hearing Research 337, 46–58.
https://doi.org/10.1016/j.heares.2016.04.009

Lewarch, C.L., Hoekstra, H.E., 2018. The evolution of nesting behaviour in Peromyscus mice.
Animal Behaviour 139, 103–115. https://doi.org/10.1016/j.anbehav.2018.03.008
Light, J.E., Siciliano-Martina, L., Dohlanik, E.G., Hafner, D.J., Lawing, A.M., Greenbaum, I.F.,
Light, J.E., Siciliano-Martina, L., Dohlanik, E.G., Hafner, D.J., Lawing, A.M.,
Greenbaum, I.F., 2021. Morphological differentiation of Peromyscus leucopus and P.
maniculatus in East Texas. Therya 12, 369–387. https://doi.org/10.12933/therya-21-1116
McCullagh, E.A., Peacock, J., Lucas, A., Poleg, S., Greene, N.T., Gaut, A., Lagestee, S., Zhang,
Y., Kaczmarek, L.K., Park, T.J., Tollin, D.J., Klug, A., 2022. Auditory brainstem
development of naked mole-rats (Heterocephalus glaber). Proceedings of the Royal
Society B: Biological Sciences 289, 20220878. https://doi.org/10.1098/rspb.2022.0878
McCullagh, E.A., Poleg, S., Greene, N.T., Huntsman, M.M., Tollin, D.J., Klug, A., 2020.
Characterization of Auditory and Binaural Spatial Hearing in a Fragile X Syndrome
Mouse Model. eNeuro 7. https://doi.org/10.1523/ENEURO.0300-19.2019
Miller, J.R., Engstrom, M.D., 2012. Vocal stereotypy in the rodent genera Peromyscus and
Onychomys (Neotominae): taxonomic signature and call design. Bioacoustics 21, 193–
213. https://doi.org/10.1080/09524622.2012.675176
Millien, V., Ledevin, R., Boué, C., Gonzalez, A., 2017. Rapid morphological divergence in two
closely related and co-occurring species over the last 50 years. Evol Ecol 31, 847–864.
https://doi.org/10.1007/s10682-017-9917-0
Mills, J.H., Schmiedt, R.A., Kulish, L.F., 1990. Age-related changes in auditory potentials of
mongolian gerbil. Hearing Research 46, 201–210. https://doi.org/10.1016/0378-
5955(90)90002-7
Moore, E.J., 1983. Bases of auditory brain-stem evoked responses. Grune & Stratton.
Neil J. Ingham, S.K.T., Spiro D. Comis, Deborah J. Withington, 1998. The Auditory Brainstem
Response of Aged Guinea Pigs. Acta Oto-Laryngologica 118, 673–680.
https://doi.org/10.1080/00016489850183160
New, E.M., Hurd, J.A., Alarcon, G.A., Miller, C.S., Williams, P.A., Greene, N.T., Sergott, C.E.,
Li, B.-Z., Lei, T.C., McCullagh, E.A., 2024. Hearing ability of prairie voles (Microtus
ochrogaster). J Acoust Soc Am 155, 555–567. https://doi.org/10.1121/10.0024357
Nicolas, V., Schaeffer, B., Missoup, A.D., Kennis, J., Colyn, M., Denys, C., Tatard, C., Cruaud,
C., Laredo, C., 2012. Assessment of Three Mitochondrial Genes (16S, Cytb, CO1) for
Identifying Species in the Praomyini Tribe (Rodentia: Muridae). PLOS ONE 7, e36586.
https://doi.org/10.1371/journal.pone.0036586
Ohlemiller, K.K., Frisina, R.D., 2008. Age-Related Hearing Loss and Its Cellular and Molecular
Bases, in: Schacht, J., Popper, A.N., Fay, R.R. (Eds.), Auditory Trauma, Protection, and
Repair, Springer Handbook of Auditory Research. Springer US, Boston, MA, pp. 145–
194. https://doi.org/10.1007/978-0-387-72561-1_6
Ordóñez-Garza, N., Matson, J.O., Strauss, R.E., Bradley, R.D., Salazar-Bravo, J., 2010. Patterns
of phenotypic and genetic variation in three species of endemic Mesoamerican
Peromyscus (Rodentia: Cricetidae). Journal of Mammalogy 91, 848–859.
https://doi.org/10.1644/09-MAMM-A-167.1
Pasch, B., Tokuda, I.T., Riede, T., 2017. Grasshopper mice employ distinct vocal production
mechanisms in different social contexts. Proc. R. Soc. B. 284, 20171158.
https://doi.org/10.1098/rspb.2017.1158
Platt, R.N., II, Amman, B.R., Keith, M.S., Thompson, C.W., Bradley, R.D., 2015. What Is
Peromyscus? Evidence from nuclear and mitochondrial DNA sequences suggests the

need for a new classification. Journal of Mammalogy 96, 708–719.
https://doi.org/10.1093/jmammal/gyv067

Pomerantz, S.M., Clemens, L.G., 1981. Ultrasonic vocalizations in male deer mice (Peromyscus
maniculatus bairdi): Their role in male sexual behavior. Physiology & Behavior 27, 869–
872. https://doi.org/10.1016/0031-9384(81)90055-X

Poncelet, L., Coppens, A., Deltenre, P., 2000. Brainstem Auditory Evoked Potential Wave V
Latency-Intensity Function in Normal Dalmatian and Beagle Puppies. Journal of
Veterinary Internal Medicine 14, 424–428. https://doi.org/10.1111/j.1939-
1676.2000.tb02251.x

Ralls, K., 1967. Auditory sensitivity in mice: Peromyscus and Mus musculus. Animal Behaviour
15, 123–128. https://doi.org/10.1016/S0003-3472(67)80022-8

Ramsier, M.A., Dominy, N.J., 2010. A comparison of auditory brainstem responses and
behavioral estimates of hearing sensitivity in Lemur catta and Nycticebus coucang.
American Journal of Primatology 72, 217–233. https://doi.org/10.1002/ajp.20780

Riede, T., Kobrina, A., Bone, L., Darwaiz, T., Pasch, B., 2022. Mechanisms of sound production
in deer mice (Peromyscus spp.). Journal of Experimental Biology 225, jeb243695.
https://doi.org/10.1242/jeb.243695

Riedel, H., Kollmeier, B., 2006. Interaural delay-dependent changes in the binaural difference
potential of the human auditory brain stem response. Hearing Research 218, 5–19.
https://doi.org/10.1016/j.heares.2006.03.018

Rieger, N.S., Marler, C.A., 2018. The function of ultrasonic vocalizations during territorial
defence by pair-bonded male and female California mice. Animal Behaviour 135, 97–
108. https://doi.org/10.1016/j.anbehav.2017.11.008

Robins, J.H., Hingston, M., Matisoo-Smith, E., Ross, H.A., 2007. Identifying *Rattus* species
using mitochondrial DNA. Molecular Ecology Notes 7, 717–729.
https://doi.org/10.1111/j.1471-8286.2007.01752.x

Russell, L., 2018. Emmeans: estimated marginal means, aka least-squares means. R package
version 1.

Sayers, E.W., Bolton, E.E., Brister, J.R., Canese, K., Chan, J., Comeau, D.C., Connor, R., Funk,
K., Kelly, C., Kim, S., Madej, T., Marchler-Bauer, A., Lanczycki, C., Lathrop, S., Lu, Z.,
Thibaud-Nissen, F., Murphy, T., Phan, L., Skripchenko, Y., Tse, T., Wang, J., Williams,
R., Trawick, B.W., Pruitt, K.D., Sherry, S.T., 2022. Database resources of the National
Center for Biotechnology Information. Nucleic Acids Research 50, D20.
https://doi.org/10.1093/nar/gkab1112

Sayers, E.W., Cavanaugh, M., Clark, K., Pruitt, K.D., Schoch, C.L., Sherry, S.T., Karsch-
Mizrachi, I., 2021. GenBank. Nucleic Acids Research 49, D92–D96.
https://doi.org/10.1093/nar/gkaa1023

Shi, Y., Pulliam, D.A., Liu, Y., Hamilton, R.T., Jernigan, A.L., Bhattacharya, A., Sloane, L.B.,
Qi, W., Chaudhuri, A., Buffenstein, R., Ungvari, Z., Austad, S.N., Van Remmen, H.,
2013. Reduced mitochondrial ROS, enhanced antioxidant defense, and distinct age-
related changes in oxidative damage in muscles of long-lived Peromyscus leucopus.
American Journal of Physiology-Regulatory, Integrative and Comparative Physiology
304, R343–R355. https://doi.org/10.1152/ajpregu.00139.2012

Sikes, R.S., Gannon, W.L., the Animal Care and Use Committee of the American Society of
Mammalogists, 2011. Guidelines of the American Society of Mammalogists for the use

of wild mammals in research. Journal of Mammalogy 92, 235–253.
https://doi.org/10.1644/10-MAMM-F-355.1

Stancampiano, A.J., Schnell, G.D., 2004. Microhabitat Affinities of Small Mammals in Southwestern Oklahoma. Journal of Mammalogy 85, 948–958.

Suthakar, K., Liberman, M.C., 2019. A simple algorithm for objective threshold determination of auditory brainstem responses. Hearing Research 381, 107782. https://doi.org/10.1016/j.heares.2019.107782

Suzuki, T., Horiuchi, K., 1981. Rise Time of Pure-Tone Stimuli in Brain Stem Response Audiometry. Audiology 20, 101–112. https://doi.org/10.3109/00206098109072688

Ungan, P., Yağcioğlu, S., Özmen, B., 1997. Interaural delay-dependent changes in the binaural difference potential in cat auditory brainstem response: implications about the origin of the binaural interaction component 1. Hearing Research 106, 66–82. https://doi.org/10.1016/S0378-5955(97)00003-8

Voelkl, B., Altman, N.S., Forsman, A., Forstmeier, W., Gurevitch, J., Jaric, I., Karp, N.A., Kas, M.J., Schielzeth, H., Van de Casteele, T., Würbel, H., 2020. Reproducibility of animal research in light of biological variation. Nat Rev Neurosci 21, 384–393. https://doi.org/10.1038/s41583-020-0313-3

Vrana, P.B., Shorter, K.R., Szalai, G., Felder, M.R., Crossland, J.P., Veres, M., Allen, J.E., Wiley, C.D., Duselis, A.R., Dewey, M.J., Dawson, W.D., 2014. Peromyscus (deer mice) as developmental models. WIREs Developmental Biology 3, 211–230. https://doi.org/10.1002/wdev.132

Wang, Xin, Zhu, M., Samuel, O.W., Wang, Xiaochen, Zhang, H., Yao, J., Lu, Y., Wang, M., Mukhopadhyay, S.C., Wu, W., Chen, S., Li, G., 2020. The Effects of Random Stimulation Rate on Measurements of Auditory Brainstem Response. Frontiers in Human Neuroscience 14.

Wickham, H., 2016. Programming with ggplot2, in: Wickham, H. (Ed.), Ggplot2: Elegant Graphics for Data Analysis, Use R! Springer International Publishing, Cham, pp. 241–253. https://doi.org/10.1007/978-3-319-24277-4_12

Wolski, L.F., Anderson, R.C., Bowles, A.E., Yochem, P.K., 2003. Measuring hearing in the harbor seal (Phoca vitulina): Comparison of behavioral and auditory brainstem response techniques. The Journal of the Acoustical Society of America 113, 629–637. https://doi.org/10.1121/1.1527961

Zhou, X., Jen, P.H.-S., Seburn, K.L., Frankel, W.N., Zheng, Q.Y., 2006. Auditory brainstem responses in 10 inbred strains of mice. Brain Research 1091, 16–26. https://doi.org/10.1016/j.brainres.2006.01.107

Figure legends:

Figure 1: Map showing trapping site locations in Oklahoma. Packsaddle wildlife management area (WMA) sites are presented by yellow triangle, James Collin wildlife management area (WMA) sites are presented by blue squares, and Payne County sites are presented by red circle.

Figure 2: Auditory Brainstem response patterns of a female *P. leucopus* determined with clicks of different intensities. Peak latency of monaural wave I and IV decrease with increasing click intensity (dotted lines). B (a) represents latency intensity functions of wave I and B (b) shows

1037   latency intensity functions of wave IV. The slope of the latency intensity function was calculated
1038   as the amount of change in peak latency per decibel.
1039
Figure 3: Morphological differences between *P. leucopus* and *P. maniculatus*. Pinnae, head, and
body features (**A**) were evaluated between species (pink boxplot = *leucopus*, blue boxplot =
maniculatus). Measurements JK show the inter pinnae distance, JN the nose to pinna distance,
MK the pinna width, LM the pinna height, OP the tail length, and PQ the body length. Effective
pinna diameter was calculated by taking the square root of pinna height multiplied by pinna
width (MK/LM). Significant differences were observed for all features: Pinna width (**B**), Pinna
length (**C**), Effective diameter (**D**), Nose to pinna distance (**E**), Inter pinna distance (**F**), Body
length (**G**), Tail length (**H**), and Body mass (**I**). Peromyscus head image (**A**) was obtained from
Rose Pest Solutions website and body/tail is from the OSU Collection of Vertebrates and is a
preserved sample, not an animal that was measured in this current study. Image is presented only
for demonstration of measurements.
1051
1052   **Figure 4:** Figure 4A and 4B show Auditory Brainstem response patterns of a female *P. leucopus* and a
1053   female *P. maniculatus* determined with clicks of different intensities, respectively. Hearing range was
1054   measured across frequency (1-64 kHz) for both *P. leucopus* and *P. maniculatus* (Figure 4C). No
1055   significant main effects of frequency between species were found. Unfilled blue circles represent *P.
1056   maniculatus* while filled pink squares represent *P. leucopus*.
1057
1058   **Figure 5:** Amplitudes of auditory brainstem responses wave I-IV. Data represent the response
1059   evoked by 90 dB SPL click stimuli between both species. No significant main effects of wave
1060   amplitude between species were found. Blue represents *P. maniculatus* while pink represents *P.
1061   leucopus*.
1062
1063   **Figure 6:** Latencies of auditory brainstem responses. Data represent latencies of ABR wave I-IV
1064   evoked responses by 90 dB SPL click stimulus between both species. No significant main effects
1065   of wave latency between species were found. Blue represents *P. maniculatus* while pink
1066   represents *P. leucopus*.
1067
1068   **Figure 7:** Average slope of latency-intensity function of waves I, and IV of ABRs (Pink = *P.
1069   leucopus* (n = 15), Blue = *P. maniculatus* (n = 11)).
1070
1071
1072   **Figure 8:** Binaural hearing in wild *P. leucopus* (pink) and *P. maniculatus* (blue). Binaural
1073   amplitude and latency for the BIC with varying ITD between - 2 to + 2 ms in 0.5 ms steps were
1074   measured. No significant differences were observed between both species at BIC amplitudes.
1075   Significant differences were detected in BIC latencies between both species across all ITDs.
1076
1077   **Table 1:** Age was estimated based on body mass for each species based on published literature.
1078   Ages for *P. maniculatus* was describe as follows: Juveniles < 14 grams, subadults, between 14-
1079   17 grams, and adults, > 17 grams (Fairbairn, 1977). We inferred ages for *P. leucopus* as follow:
1080   Juveniles < 13 grams, subadults, between 13 – 18 grams, and adults > 18 grams (Cummings and
1081   Vessey, 1994). We did not make comparisons by ages due to limited sample sizes by age groups.
1082
1083

1084

1085 **Table 2:** Morphological characteristics features of *P. maniculatus* and *P. leucopus* of the
1086 Packsaddle wildlife management area (WMA), James Collin wildlife management area (WMA)
1087 and Payne County. Values presented represent the mean of different morphological features
1088 recorded, the degrees of freedom, F-statistic and p-value of morphological differences between
1089 species.

1107